1   ***Smad4* and TGF-β1 dependent gene expression biomarkers in conditional intestinal ade-**
2   **noma, organoids and colorectal cancer.**
3

4   Mirvat Surakhy[1], Julia Matheson[1], David Barnes[1,3], Emma J. Carter[1], Jennifer Hughes[1], Clau-
5   dia Bühnemann[1], Sabina Sanegre[1], Hans Morreau[2], Paul Metz[1], Charlotte J. Imianowski[1] and
6   A. Bassim Hassan[1,¶].
7
8   [1]Oxford Molecular Pathology Institute, Sir William Dunn School of Pathology, University of
9   Oxford, South Parks Road, Oxford, OX1 3RE, United Kingdom.
10   [2]Department of Pathology, Leiden University Medical Centre, Leiden, The Netherlands.
11   [3]Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham B15 2TT

12
13
14   Short title;
15   *Smad4 and TGF-β1 dependent functional gene expression in intestinal adenoma*
16
17   Key words; Smad4, Apc, intestine, adenoma, TGF-β1, Id1, Spp1, Pak3, gene expression, co-
18   lorectal cancer, biomarkers
19

Corresponding author:
[¶]Prof A B Hassan: bass.hassan@path.ox.ac.uk
Oxford Molecular Pathology Institute, Sir William Dunn School of Pathology, University of
Oxford, South Parks Road, Oxford, OX1 3RE, UK
Tel: +44 (0)1865 275044
Fax: +44 (0)1865 275515

26
27

Abstract

TGF-β ligand activation suppresses cell growth yet can paradoxically and potently

promote cancer invasion and metastasis depending on downstream pathway mutational

context. Here, we evaluated the basis of this observation in conditional murine intestinal

adenoma models with and without loss of Mothers against decapentaplegic homolog 4

(*Smad4*), with the aim of identifying TGF-β-BMP-SMAD4 pathway dependent gene expres-

sion biomarkers for translational application. Conditional *Lgr5*-CreER^{T2} activation in

Apc$^{fl/fl}$Smad4$^{fl/fl}$ resulted in adenoma formation with recombined homozygote floxed alleles

(Apc$^{Δ/Δ}$Smad4$^{Δ/Δ}$). The adenoma phenotype was discordant, with a reduced small intestinal

adenoma burden yet development of large non-metastatic caecal adenoma with nuclear

localisation of phospho-Smad2/3. Derived *Apc$^{Δ/Δ}$Smad4$^{Δ/Δ}$* adenoma organoids resisted

TGF-β1 dose dependent growth arrest and cell death (IC$_{50}$ 534pM) compared to

Apc$^{Δ/Δ}$Smad4$^{+/+}$ (IC$_{50}$ 24pM). TGF-β1 (390pM) modified adenoma mRNA expression (bulk

RNA-Seq) most significantly for *Id1^{low}* and *Spp1^{high}* in *Apc$^{Δ/Δ}$Smad4$^{Δ/Δ}$*. Single cell RNAseq of

caecal adenoma identified expansion of *Lgr5^{low}, Pak3^{high}* and *Id1^{low}* progenitor populations

in *Apc$^{Δ/Δ}$Smad4$^{Δ/Δ}$*. Of the 76 *Smad4* and TGF-β1 dependent genes identified in adenoma

organoids, 7 human equivalent genes were also significantly differentially expressed in co-

lorectal cancer, including *ID1^{low}, SPP1^{high}* and *PAK3^{high}* that also correlated with poorer sur-

vival (TCGA cohorts). Murine conditional models identified *Smad4* loss of function mRNA

expression biomarkers that require further evaluation as functional classifiers of colorectal

cancer subtypes.

212

Introduction

Mutation of the downstream TGF-β-BMP ligand-dependent signalling pathway can

result in context-dependent and paradoxical gain of function (GOF) of the cancer pheno-

type, i.e. the ligand activation of the normally functioning signalling pathway exerts a tu-

mour suppressive effect, yet following downstream pathway mutation, ligand activation

can then result in 'oncogenic' GOF activation, epithelial-mesenchymal transition (EMT) and

potent promotion of the invasive and metastatic cancer phenotype [1]. It is well established

that human cancer associated variants (mutations) of the TGF-β-BMP-SMAD4 pathway are

correlated with poorer prognosis in enriched sub-types of intestinal cancers such colorectal

and pancreatic adenocarcinoma [2]. Despite the incorporation of whole genome sequencing,

gene level RNA sequencing and proteomic biomarkers into supervised machine learning

prognostic classifiers, for example in colorectal cancer, there remains a paucity of pathway-

specific gene expression biomarkers more closely linked to molecular mechanisms and in-

dicative of pathway activity [3,4]. This background provided the impetus for further evalua-

tion of the critically important TGF-β-BMP pathway that when disrupted, is associated with

the worse prognostic sub-types of intestinal cancer. Here, we use murine conditional *Apc*

dependent intestinal models to generate *in vivo* and *in vitro* adenoma with and without

TGF-β-BMP pathway modification to prospectively identify pathway and context specific

gene expression.

The TGF-β1 ligand acts as a dimer to bring together specific heterodimeric cell sur-

face type I and II serine/threonine kinase receptors to activate signals by phosphorylation

of R-Smad transcription factors. For example, TGF-β1 binds to TGFBR1 and TGFBR2 to sig-

nal through p-Smad2/3, but can also bind to activin receptor type 1A (ACVR1) to transduce

signal through p-Smad1/5 [5,6]. Following receptor activation, phospho-R-Smads form either

heterotrimeric or heterodimeric complexes with Smad4, the co-Smad, as it continuously

shuttles between the cytoplasm and nucleus [5,7]. Smad4-R-Smad complexes bind with co-

factors and DNA to co-ordinate gene expression [6]. Chromatin regions can alter quickly

(within 10 minutes) following TGF-β1 stimulation, with assembled Smad complexes binding

to *cis*-regulatory elements with AP-1 footprints [8,9]. Smad4 and R-Smad proteins consist of

two highly conserved globular domains, named Mad homology domains, both at the N-

terminus (MH1) and C-terminus (MH2) separated by a less well-conserved serine and

proline-rich linker region critical for protein-protein interactions and stability [6,9,10]. The

MH1 domain is involved in DNA binding through a conserved identical hairpin structure

present in all members except Smad2 [5,11,12]. The majority of SMAD4 loss of function (LOF)

cancer somatic mutations are missense, although nonsense, splice site, frameshift, and in-

frame insertion/deletion mutations exist. In terms of protein localisation, frequent hot

spot mutations occur within the conserved R-SMAD binding surface of the MH2 domain

and missense mutations reported in the MH1 domain along the DNA-binding interface [13].

The specialised intestinal epithelium is comprised of crypt-villus functional cell

units that provide an ideal experimental system for the investigation of specific compo-

nents of the TGF-β signalling pathway [14]. Differentiated cells in the crypt and villous com-

prise of four epithelial lineages: enterocytes, goblet, entero-endocrine and Paneth cells [14].

Leucine-rich repeat containing G-protein-coupled receptor (*Lgr5*) positive crypt base co-

lumnar cells (CBC) are stem cells that interact with Paneth cells to maintain the stem cell

niche through *Wnt* activation and β-catenin stability [15,16]. *Lgr5* positive CBC cells divide,

renew and create transit amplifying cells that undergo 4 to 5 cycles of cell division before

differentiating as they ascend the villus. By using a *Lgr5* driven Cre recombinase, genes

with associated loxP sites can be randomly disrupted in CBCs once the Cre is activated.

BMP signalling is active at the crypt-villous border towards the villous tip and is controlled

by a ligand gradient to promote differentiation towards secretory and enteroendocrine

cells [17-19]. BMP concentrations are regulated in the intestinal crypts by several different

BMP antagonists, such as Gremlin, Noggin and Chordin-like1 [20]. TGF-β type I/II receptors

and ligands are present and expressed in the muscularis, lamina propria and the differenti-

ated compartment of both the small intestine and colon [21]. TGF-β is also frequently pre-

sent in the intestinal tumour microenvironment and is released either by cancer cells, tu-

mour stromal cell or immune cells [22]. Intestinal adenoma can be conditionally generated

when *Apc* loss of function no longer regulates the degradation of β-catenin, resulting in

constitutive activation of transcription by β-catenin/ Tcf4. Subsequent mutational events

are selected for during the progression of adenoma to invasive carcinoma in humans, e.g.

RAS/MAPK GOF, *PTEN* and *TP53* LOF, and later acquired LOF mutation of the TGF-β-BMP

ligand activated receptor pathway, such a mutation of receptors and Smad2/3/4 [23]. The

conditional *Apc*$^{fl/fl}$ model using *Lgr5CreER*^{T2} is well characterised and so we incorporated

loxP dependent (LOF) of *Smad4* (*Smad4*$^{fl/fl}$ alleles) into this model.

Here, we first set out to evaluate loss of the TGF-β pathway signal transduction

regulator, Mothers against decapentaplegic homolog 4 (*Smad4*), *in vivo* and *in vitro*, with

respect to a murine intestinal adenoma phenotype and dependent gene expression. In

view of the complexity of the interacting pathways that can alter TGF-β1-*Smad4* induced

phenotypic effects, especially in immortalised human cancer cell lines, we sought to better

define the contextual impact of *Smad4* LOF to specifically evaluate the *Smad4* and TGF-β1

ligand dependency on gene expression biomarkers and to translate these findings with re-

spect to human intestinal cancer.

Results

Combined Apc and Smad4 conditional disruption results in discordant adenoma pheno-

types

To investigate the genetic interactions between *Apc* and *Smad4* in the murine in-

testine, we combined conditional alleles with a *Cre* driven by the promoter *Lgr5* and a

ROSA26-loxSTOP reporter allele (*Lgr5-EGFP-IRES-CreER*^{T2} or *Lgr5-CreER*^{T2})[24] (Fig. 1a). Het-

erozygote alleles were inter-crossed (*Apc*$^{fl/+}$*Smad4*$^{fl/+}$*Lgr5-CreER*^{T2}) on a C57Bl/6J back-

ground (backcrossed to C57Bl/6J for >10 generations) to generate homozygotes and het-

erozygote littermates that differ in Smad4 conditional alleles [25,26]. Conditional recombina-

tion followed tamoxifen injection (4OHT) in adult animals (Fig. 1a). The floxed allele of

Smad4 (*Smad4*$^{Δ/Δ}$) resulted in loss of the MH1 DNA binding domain and the production of a

truncated 43kDa protein which was not capable of translocating to the nucleus (see below,

Supplementary Fig. 1). The germline homozygote loxP alleles are denoted *Smad4*$^{fl/fl}$ and

Apc$^{fl/fl}$, and where somatic tissues have been directly tested for the presence of the result-

ing Cre-induced recombined alleles (floxed) using PCR, were denoted as *Smad4*$^{Δ/Δ}$ and

Apc$^{Δ/Δ}$. Recombination rates in intestinal epithelial cells were lower overall with the *Lgr5*

transgene (5-6%), but highly specific for *Lgr5* CBC stem cells [15,27]. Rapid adenoma devel-

opment occurred as expected (median 35-40 days post injection 4OHT) and animals were

culled if they exceeded predefined humane endpoints. Comparison of overall survival

(Kaplan-Meier) between *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$*Lgr5-CreER*^{T2} and *Apc*$^{fl/fl}$*Smad4*$^{fl/+}$*Lgr5-CreER*^{T2} with

Apc$^{fl/fl}$*Smad4*$^{+/+}$*Lgr5-CreER*^{T2} controls showed a modest but significant improvement in sur-

vival (p=0.011, Fig. 1a). These observations suggest that introducing *Smad4* LOF in *Apc* LOF

intestinal adenoma does not result in overwhelming tumour progression and metastasis.

However, analysis of intestinal adenoma distribution between the small intestine and colon

showed a marked discordance in growth phenotypes in *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$*Lgr5-CreER*^{T2}. A sig-

nificant reduction in small intestinal adenoma burden was observed with paradoxical

growth promotion of adenoma in the caecum (Fig. 1b). As the distribution of adenoma on

a C57BL/6 background is predominantly in the small intestine rather than the colon and

caecum, the multiple small intestinal adenoma would be expected to promote anaemia,

small intestinal obstruction and decreased survival [27,28]. The significantly reduced small in-

testinal adenoma burden in *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$*Lgr5-CreER*^{T2} compared to controls may have

improved overall survival if this was the only phenotypic effect (Fig. 1a, b). Examination of

the caecum and large intestine revealed markedly enlarged coalesced adenoma where

numerical analysis of single adenoma was impossible, hence we reported caecal weight as

a surrogate (Fig 1b).   The adenoma burden in heterozygote *Apc*$^{fl/fl}$*Smad4*$^{+/fl}$*Lgr5-CreER*^{T2}

compared to controls was statistically non-significant and does not support either *Smad4*

haplo-insufficiency or a dominant negative effect of *Smad4*$^{fl/+}$. Histological grading of small

intestinal adenoma showed low-grade dysplasia in all genotypes, whereas more aggressive

carcinoma *in situ* was observed in regions of *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$*Lgr5-CreER*^{T2} in the caecum

(Fig. 1c, author HM pathological review). Examination of H&E stained small intestinal ade-

noma tissue sections showed that *Apc*$^{fl/fl}$*Smad4*$^{+/+}$ mice had sessile adenomatosis appear-

ance compared to *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$ that had mainly tubular adenomas.   Crypt abscesses

were detected in both genotypes but no tissue invasion and metastasis were observed (Fig.

1c). The efficiency of Cre-mediated *Smad4* allele deletion was examined by immunohisto-

chemistry (IHC, Fig. 1c). Smad4 protein was clearly detected in *Apc*$^{Δ/Δ}$*Smad4*$^{+/+}$ adenoma

and adjacent tissue as expected.   Adenoma from *Apc*$^{Δ/Δ}$ *Smad4*$^{Δ/Δ}$ did not label strongly

with anti-Smad4 antibodies, consistent with adenoma-specific disruption in both the small

and large intestine (Fig.1c). The presence of active TGF-β (BMP) signaling was supported

by p-Smad2 and p-Smad1/5/8 nuclear immunolabeling, respectively, as was confirmed by

immunoblots (Fig. 1d). Both p-Smad2 and p-Smad 1/5/8 labelling were detected in adeno-

ma from *Apc*$^{Δ/Δ}$*Smad4*$^{Δ/Δ}$ compared to normal tissue and littermate control adenoma, con-

sistent with either TGF-β alone or with BMP ligand activation in combination (p<0.001, Fig.

1d).

Combined Apc and Smad4 disruption modifies growth of adenoma organoids

In order to generate an organoid culture, we isolated *Apc*$^{Δ/Δ}$*Smad4*$^{Δ/Δ}$ (*Smad4*$^{Δ/Δ}$) and

Apc$^{Δ/Δ}$*Smad4*$^{+/+}$ (*Smad4*$^{+/+}$) caecal adenoma and generated *in vitro* cultures in Matrigel us-

ing   published   conditions   [24,29].   Using   a   ROSA26-LSL-YFP   reporter   line   bred   with

Apc$^{fl/fl}$*Smad4*$^{fl/fl}$*Lgr5-CreER*^{T2}, we confirmed that 40HT exposure resulted in uniform YFP ex-

pression in adenoma organoid culture that was also TGF-β1 independent (Fig. 2a). In cul-

ture, *Smad4*$^{Δ/Δ}$ adenomas appeared smaller and grew slower compared to *Smad4*$^{+/+}$ ade-

nomas, as measured by both area and the alamarBlue (AB) growth assay (Supplementary

Fig. 2). Noggin is a BMP pathway inhibitor that normally prevents enterocyte differentia-

tion and is required for crypt culture maintenance [23,24,29]. Noggin addition had no effect on

adenoma growth (Supplementary Fig. 2). SB-505124, a TGF-βR1 small molecule inhibitor

also had no significant inhibitory effect on Smad4$^{+/+}$ and *Smad4*$^{Δ/Δ}$ adenoma growth (1 μM)

without the exogenous addition of TGF-β1 (Supplementary Fig.2). These data suggest that

the adenoma culture conditions appeared to be independent of endogenous TGF-β supply.

6   To exclude whether there may have been a genotype dependent difference in the

floxing efficiency between *Apc* and *Smad4* that may have accounted for the two-fold dif-

ferences in growth in culture, we also generated adenoma with an alternative conditional

Cre (*Villin-CreER*^{T2}). Here, cultured small intestinal organoids from healthy animals bred

with non-floxed alleles, *Apc*$^{fl/fl}$*Smad4*$^{+/+}$*Villin-CreER*^{T2}*Rosa26-YFP* and *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$*Villin-*

CreER^{T2}*Rosa26-YFP* genotypes, were cultured in organoid media containing R-spondin-1,

Noggin and EGF. Intestinal organoids that formed a normal crypt-villous morphology were

exposed to 4OHT followed by withdrawal of R-spondin-1. This resulted in adenoma organ-

oid formation in both *Apc*$^{Δ/Δ}$*Smad4*$^{+/+}$ and *Apc*$^{Δ/Δ}$*Smad4*$^{Δ/Δ}$ (Supplementary Fig. 3). Allele

recombination was confirmed by PCR product of 259bp and 512bp for *Apc* and *Smad4* al-

leles, respectively. Both the *Apc*$^{fl/fl}$ and *Smad4*$^{fl/fl}$ alleles recombined at the same time and

rate following 4OHT treatment, and were detected as early as day 2 post treatment (Sup-

plementary Fig. 3). Thus, floxing efficiency appeared similar between alleles.

Deletion of MH1 domain of Smad4 following Cre-mediated recombination.

To evaluate the expression of *Smad4* mRNA in *Smad4*$^{Δ/Δ}$ and *Smad4*$^{+/+}$ *in vitro*, we

used bulk RNA-seq of adenoma organoids and discovered two different *Smad4* mRNA tran-

scripts. *Smad4* gene tracking with Cuffdiff revealed transcripts from the intact *Smad4*

gene, and a second transcript from the *Smad4* gene with exon 2 excised following loxP me-

diated recombination (Supplementary Fig. 1). Immunoblots confirmed expression of

Smad4 full-length protein (61kDa) in both *Smad4*$^{+/+}$ and *Smad4*$^{+/Δ}$ adenomas, with a smaller

band of (43kDa) detected in protein extracts from *Smad4*$^{+/Δ}$ and *Smad4*$^{Δ/Δ}$ adenomas (Sup-

plementary Fig. 1). *In silico* analysis revealed an open reading frame in exon 5 of the *Smad4*

sequence and mapping with ExPASy revealed that exon 2 deletion results in a truncated

protein in which the *Smad4* MH1 DNA binding domain is selectively deleted. The predicted

size of the truncated protein was 43kDa, comprising the linker and Smad4 MH2 domain

(Supplementary Fig. 1). The full-length Smad4 protein in *Smad4*$^{+/+}$ adenoma showed nucle-

ar localisation following exposure of 390pM TGF-β1 for 2 hours. Under the same condi-

tions, no nuclear localisation of Smad4 (43kDa) was detected in *Smad4*$^{Δ/Δ}$ adenomas, con-

sistent with loss of DNA binding function (Supplementary Fig. 1).   Taken together, Cre-

mediated recombination of the Smad4 gene resulted in selective MH1 deletion and a trun-

cated protein that did not undergo nuclear localisation following TGF-β1 stimulation. The-

se data, and the apparent reduction of detection using the same anti-Smad4 antibody in

IHC of *Smad4*$^{\Delta/\Delta}$ adenoma, are consistent with non-DNA binding and presumably soluble

truncated SMAD4 protein that depletes from of cells during tissue processing resulting in

reduced detection.

TGF-β1 dose dependency in Smad4$^{\Delta/\Delta}$ adenoma organoids

We next tested the dose and timing effects of exogenous addition of TGF-β1 to ade-

noma organoid cultures and the subsequent localisation of R-Smads.   TGF-β1 induced

dose-dependent growth inhibition of both the *Smad4*$^{+/+}$ and *Smad4*$^{\Delta/\Delta}$ adenomas, but with

a 20-fold differential sensitivity (Fig. 2b).   *Smad4*$^{+/+}$ adenomas were more sensitive to TGF-

β1, with an IC$_{50}$ of 24pM and threshold of 0.39pM.   For *Smad4*$^{\Delta/\Delta}$, the TGF-β1 IC$_{50}$ was

534pM with a threshold between 39-390pM. The decrease in adenoma viability with TGF-

β1 dosing was prevented when the adenoma cultures were pre-incubated with SB-505124

(Fig. 2b). For subsequent experiments, a TGF-β1 concentration of 390pM was used to dif-

ferentiate the TGF-β1 context-dependent sensitivity effects between *Smad4*$^{+/+}$ (100 fold

higher than the threshold) and *Smad4*$^{\Delta/\Delta}$ (at threshold but below IC$_{50}$) adenomas. TGF-β1

addition (390pM) resulted in the detection of phosphorylated p-Smad2 and p-Smad3 in

both the *Smad4*$^{+/+}$ and *Smad4*$^{\Delta/\Delta}$ adenomas as expected (Fig. 2c). Despite the differential

sensitivity, the kinetics of response to TGF-β1 stimulation were similar between genotypes,

with p-Smad2 maximally detected after 1-2 hours and associated with nuclear localisation

(Fig. 2c). Within 24 hours of TGF-β1 administration, >90% of the *Smad4*$^{+/+}$ adenoma cells

either had either morphological changes of apoptosis with organoid retraction, or changes

in shape toward a spindle or mesenchymal phenotype (Supplementary Movies 1-4). These

features appeared delayed to 72 hours in *Smad4*$^{\Delta/\Delta}$ adenomas but were still detectable (Fig.

2e). The reduced viability and altered morphology were accompanied by a significant de-

creased in proliferation and increased apoptosis in *Smad4*$^{+/+}$ adenoma when quantified by

EdU and cleaved caspase 3 (CC3) labelling, respectively (Fig. 2f, g).

TGF-β1 and Smad4-dependent Id1 and Spp1 gene expression in adenoma organoids

To assess the transcriptome response to TGF-β1 that might inform differences

between *Smad4*$^{+/+}$ and *Smad4*$^{\Delta/\Delta}$ adenoma organoids, bulk paired-end RNA-seq detected

16,143 gene transcripts that we utilised for the analysis of differentially expressed genes

(DEG). Three separate adenoma biological replicates per genotype and time were har-

vested at t=0, t=1 and t=12 hours after exposure to a 390pM TGF-β1 (Fig. 3a). Principle

component analysis (PCA) of gene expression showed a clear separation of the samples

based on genotype and duration of TGF-β1 exposure (Fig. 3b). Comparison of DEG be-

tween *Smad4*$^{\Delta/\Delta}$ and *Smad4*$^{+/+}$ at t=0 showed 111 upregulated genes (log$_2$FC >1.5) and 51

downregulated genes (log$_2$FC < -1.5) (Fig. 3c, Supplementary Table 1). The most important

observation was the magnitude of the genotype dependent expression of a group of genes

that included *Id1* (Inhibitor of DNA binding 1)*, Ddit4, Serpine1, Skil, Mucins, Myc, JunB* and

E2f target genes (Fig. 3c, d, Supplementary Fig. 4). Gene ontology (GO) analysis of biologi-

cal processes showed enrichment for epithelial cell proliferation and response to TGF-β1

(Supplementary Fig. 4, Supplementary Table 2). Gene set enrichment analysis (GSEA) using

hallmark gene sets with "tmod" in *Smad4*$^{\Delta/\Delta}$ showed downregulation of *Myc* target genes,

differential regulation of TGF-β signalling genes, mainly by upregulation, and enrichment of

other pathways including IL2-Stat5 and TNF signalling (AUC >0.5, padj<0.0001, Fig. 3e, Sup-

plementary Table 2). Importantly, mucins in *Smad4*$^{\Delta/\Delta}$ adenomas were among the DEGs

with *Muc4, Muc20, Muc2* being upregulated and *Muc6* being downregulated (padj p<0.05,

Supplementary Table 1, supplementary Fig. 5). Subsequent Mucin 2 immunolabeling con-

firmed protein overexpression in *Smad4*$^{\Delta/\Delta}$ adenoma tissue sections (Supplementary Fig. 5).

For early timepoints, DEG were mainly observed in *Smad4*$^{+/+}$ adenoma organoids

and included *Skil, Junb, Serpine 1, Ddit4* and *Id1* (Fig. 3d, Supplementary Fig. 4, Supplemen-

tary Table 1). The differential suppression of *Id1* in *Smad4*$^{\Delta/\Delta}$ compared to *Smad4*$^{+/+}$ after 1

hour achieved a 2.83 log$_2$FC difference.   By 12 hours, *Id1* maintained suppression (1.2

log2FC, Fig. 3d, Supplementary Table 1). Despite the repression of expression, Id1 protein

abundance and immunolocalisation did not appear to be significantly altered between

adenoma genotypes (Supplementary Fig. 6). We also observed that *Spp1* (Secreted phos-

phoprotein 1), an extracellular matrix gene, was the only gene significantly upregulated

gene following TGF-β1 exposure (12 hours) in *Smad4*$^{\Delta/\Delta}$ (log$_2$FC >1.5 threshold, Fig. 3c, d.,

Supplementary Table 3).

Later time points displayed further significant time dependent differences in gene

expression. For example, 35 DEGs were observed in *Smad4*$^{\Delta/\Delta}$ compared to 689 in *Smad4*$^{+/+}$

adenomas at t=12 hours vs t=0 hours (Fig.3d, Supplementary Table 1). Pathway enrich-

ment analysis using tmod also showed that *E2f, G2m, Myc* and mTORC targets were more

significantly downregulated in *Smad4*$^{+/+}$ adenomas compared to *Smad4*$^{\Delta/\Delta}$ adenomas (Fig.

3e, Supplementary Table 2). Time series analysis using the likelihood ratios showed differ-

ences in expression of genes including *Fos*, *Lama3, Aldh1a3, Col7a1* and *Doks* associated

with increased induction by t=12 hours, whereas *E2f8, Exo1, Clspn, Kif14* and *Kif12* all were

decreased in *Smad4$^{+/+}$* adenomas. Overall, the suppression of cell cycle genes was evident

in *Smad4$^{+/+}$* compared to *Smad4$^{Δ/Δ}$* adenomas, as shown by Metascape overrepresentation

(Fig.3f, g, Supplementary Table 3). Finally, TRRUST (Transcriptional Regulatory Relationship

Unravelled by Sentence based Text mining) enrichment analysis implemented in Me-

tascape also suggested potential regulation by transcription factors, with *Smad4*-

dependent downregulated genes being co-regulated by Sp1, Smad3 and Jun, and down-

regulated genes by Rbl2, Tp53 and E2f1 (Supplementary Fig.4). Several gene signatures

were therefore determined that were *Smad4* and TGF-β1 dependent, with the two genes

with the greatest magnitude of differential expression being *Id1* (low) and *Spp1* (high).

TGF-β1 dependent gene expression and protein correlations

Smad4 LOF did not fundamentally alter the range of TGF-β1 phenotypic responses,

but did change the overall magnitude and relative gene expression. Whilst TGF-β1 admini-

stration in *Smad4$^{+/+}$* adenomas resulted in a >4.5 Log$_2$FC relative increase in expression *of*

Fn1, we did not observe a similar change in Fn1 protein by Western blot (fibronectin 1,

Supplementary Fig. 7a). In addition, TGF-β1 exposure resulted in a 0.25 log$_2$FC reduction in

Survivin (Birc5) expression after 12 hours in *Smad4$^{Δ/Δ}$* compared to a 2.55 log$_2$FC reduction

in *Smad4$^{+/+}$* adenoma. Anti-survivin antibody western blots and nuclear localisation sug-

gested detectable reduction in protein levels in *Smad4$^{+/+}$* adenoma by 18 hours post- TGF-

β1 (Supplementary Fig. 8). The *Wnt* dependent intestinal stem cell marker *Lgr5* was down-

regulated in *Smad4$^{+/+}$* following TGF-β1 (12 hours) compared to *Smad4$^{Δ/Δ}$* adenoma (2

log$_2$FC, Supplementary Table 1), suggesting modification in Wnt pathway activity. E-

cadherin and β-catenin proteins remained immunolocalised at the adherens junctions and

in nuclei in both adenoma genotypes as expected (Supplementary Fig. 9). Immunoblots of

adenomas from both genotypes showed multiple molecular weight bands for β-catenin

and E-cadherin in the context of disruption of the β-catenin destruction complex by LOF of

Apc$^{Δ/Δ}$ (Supplementary Fig. 9b, c). Gene expression for *Myc* was not markedly different

between genotypes, yet c-Myc protein levels appeared increased in *Smad4$^{Δ/Δ}$* (12-18 hours)

compared to *Smad4$^{+/+}$* adenomas (Supplementary Fig. 9c, Supplementary Table 1). Overall,

there did not appear to be significant correlation between changes in gene expression and

protein localisation and abundance between genotypes. This implies that protein abun-

dance, presumably regulated by variable processes independent of gene expression, is not

a universal and reliable surrogate for gene expression in this context.

Single cell RNA-seq of Smad4^{fl/fl} caecal adenoma are enriched in Pak3^{Hi} epithelial progeni-

tor cells.

6   In view of the differential changes in bulk mRNA expression, we next sought to

quantify mRNA expression at the single cell level in the large adenoma directly harvested

from the caecum. Epithelial cells (Epcam cluster) consisted of the *Lgr5* stem cell (*Lgr5* ISC)

and seven stem cell-like (SC-L) clusters based on differential expression of *Lgr5* and other

stem-cell markers such as *Lrig1, Smoc2, Hopx* as previously defined [30-33] (Fig. 4a, b. Supple-

mentary Fig. 10, Supplementary Table 4). Differential enrichment in stem cell clusters was

observed between the two genotypes, consisting of 61.8% of the total cell population in

Smad4$^{fl/fl}$ adenoma compared to 47.3% in *Smad4*$^{+/+}$ (Figure 4c, d, Supplementary Table 3). A

Pak3^{High} cluster (*Pak3*; P21 Rac1/CDC42 activated kinase 3) was the most enriched in

Smad4$^{fl/fl}$ (10.5%) compared to *Smad4*$^{+/+}$ (2.5%) control adenoma. Pathway enrichment

analysis for DEGs between *Smad4*$^{fl/fl}$ and *Smad4*$^{+/+}$ for the *Pak3*^{High} cluster showed an en-

richment for the Wnt pathway and pluripotency markers (Supplementary Fig. 11). The

same comparison in adenoma organoids between genotypes and all timepoints post TGF-

β1 showed a non-significant difference in expression of *Pak3*, suggesting this was an *in vivo*

adaptation that was not reproduced in organoids (Supplementary Table 1).   The

Slc14a1^{High}cluster was also enriched in *Smad4*$^{fl/fl}$ (7%) compared to *Smad4*$^{+/+}$ (3.5%), where

expression of the stem cell marker *Ly6a* was enriched (*Sac1*). Conversely, the *Lgr5* ISC clus-

ter was significantly enriched in *Smad4*$^{+/+}$ (5.8%) compared to *Smad4*$^{fl/fl}$ adenoma (2.7%)

(Figure 4d). *Smad4*$^{fl/fl}$ showed significantly lower proportions of enterocytes (EC) as a con-

sequence (7.8% *Smad4*$^{fl/fl}$ vs 23.4% *Smad4*$^{+/+}$) yet *Fmnl2*^{Hgh}, *Cachd1*^{High} and *Slc12a2*^{High}

showed similar proportions between genotypes. Note that the SC-L and the Lgr5 ISC clus-

ters originate from the same node when viewed with cluster tree (0.2 resolution) (Fig. 4a).

The *Tacstd2*^{Hgh} cluster (murine foetal intestinal progenitor gene [34]) and the *Cl*^{Hgh} cluster

(*Clusterin* marker of the revival stem cell [35]) are SC-L clusters with a similar proportions and

originate from the same node (0 at 0.2 resolution). Markers for non-stem cells that lacked

stem cell expression markers, included ribosomal gene expression [33] in transient amplifying

cells(TA) and the Paneth cells marker *Pnliprp1* (pancreatic lipase-related protein 1) [30],

enteroendocrine precursor *Gadd45g* [33], *Mt3* and *Pnliprp2* in secretory precursor cells (se-

cretory pr).

Using both the 1.2 and 2.2 resolution cluster tree, we identified two groups of

cells; the enterocyte early precursor (ECepr) expressing *Apoa1 Dmbt1, Reg4,* and *Reg3g,*

and the enterocyte late precursor (EClpr) expressing *Reg3b, Mt1,* and *Mt2* [33]. In *Smad4*[fl/fl],

EClpr (0.26% vs 4.1%) and ECepr (1.3% vs 6.5%) were depleted compared to *Smad4*[+/+] ade-

noma. The mature enterocyte clusters, EC-1 and EC-2, showed expression of *Krt19, Krt8,*

Lgals3, Lypd8, Emp1, and *Krt20* with differences in *Mxd1, Cdkn1a* and *Fam3*b expression.

Significantly, 2.2% of *Smad4*[fl/fl] were EC-1 cells compared to 5.4% for *Smad*[+/+] (Fig. 4a, b, c,

d). Thus, scRNA-seq of *in vivo* adenomas identified *Pak3*[High] and *Slc14a1*[High] as genes with

increased expression and relatively differentially enriched in progenitor epithelial cells in

Smad4[fl/fl] caecal adenoma. Moreover, comparison of combined scRNAseq using a pseudo-

bulk approach confirmed significant downregulation of *Id1* and upregulation of *Pak3* (Fig.

5f). In addition, *Id1* and *Sema5a* were the uniquely deregulated genes that were common

between organoid bulk RNA-seq and the primary pseudo-bulk caecal adenoma scRNA-seq

data (Supplementary Fig. 12). For *Spp1*, the high overall mRNA expression in adenoma de-

rived organoids was not matched in the caecal derived single cell mRNA in epithelial cells,

as expression was confined mainly to the monocyte-macrophage lineage (Fig. 6e).

Smad4 dependent ID1, SPP1 and PAK3 expression in human colorectal cancer.

Having identified Smad4- TGF-β1 pathway dependency of *Id1*[low] and *Spp1*[high] gene

expression and *Pak3*[high] expressing cell enrichment in primary caecal intestinal adenoma,

we next considered whether these functional gene expression biomarkers have prognostic

relevance in human colorectal cancer (Fig. 5, Fig. 6, Supplementary Fig. 1, Supplementary

Table 5). Data was derived from 382 individual cases from the human Cancer Genome At-

las project (TCGA; COAD, Colon Adenocarcinoma, READ, Rectal adenocarcinoma) that had

both mutational profiles in c-BioPortal and RNA-seq data. We first tested whether the co-

existing mutational context in colorectal cancer altered biomarker expression and key

regulatory gene expression pathways by assembling mutational subgroups of colorectal

cancer based on WNT-APC-β-catenin, RAS-MAPK, PI3K, and TP53 pathway variants (Fig.

5a). This test matrix derived four variant based sub-groups named Group 1-4 (Fig. 5a).

Firstly, colorectal cancers with confirmed WNT-APC-β-catenin variants were selected and

subsequently sub-divided. Group 1 and Group 2 cases were *SMAD4* wild-type and Group 3

and 4 cases had *SMAD4* pathogenic variants. Groups 1&3, and 2&4 differed depending on

the presence or absence of second pathway variants in either the RAS-MAPK, PI3K or TP53

pathway, respectively. We next utilised the 76 DEGs identified in murine adenoma

organoids and applied the equivalent genes to the human colorectal RNA-seq data. Volca-

no plots revealed only 7 genes that were significantly differentially expressed based on the

presence (n=44) or absence (n=191) of a SMAD4 pathogenic variant (Fig. 5b).   These genes

were *BCAS1, CACNB2, CREB3L1, ID1, ID3, RASGRF2* and *SLC30A2.*   Further comparison of

the subgroups based on these genes with second pathway variants revealed 26 gene-

context pairs that were significantly upregulated and 15 gene-context pairs that were sig-

nificantly downregulated (Fig. 5c, 5d).   Despite the correlation established for *Spp1* and

Pak3 in adenoma organoids, these two genes were not significantly differentially expressed

in human colorectal cancer (Fig. 5c).   Importantly, the upregulated expression of *CREB3L1*

and BCAS1 and downregulated expression of *ID1* and *ID3* appeared least impacted by the

co-existing 2^{nd} pathway variants based on visualisation of the heat maps.   However, analy-

sis based on our three predefine criteria (see Methods) of the comparison of the increased

expression of *CREB3L1* and *BCAS1*, and repression of *ID1* and *ID3*, only *ID1^{low}* met the sig-

nificance criteria.   Analysis of the TCGA cohort with respect to either the presence or ab-

sence of any additional *TGF-β/BMP* pathway variants also showed that *ID1* expression was

also significantly downregulated with co-existing *SMAD4* and other *TGF-β* pathway down-

stream mutations compared to normal control tissues (Supplementary Table 5).   Compared

to non-SMAD4 associated TGF-β/BMP pathway pathogenic variants (e.g. in Type 1 recep-

tors, other SMADs or in the BMP pathway), *SMAD4* mutations specifically appeared to have

the greatest impact on overall *ID1* repression (Fig. 6a, p<0.0001).   In the context of all

combined TGF-β/BMP pathway mutations, however, *SPP1* mRNA did appear to be ex-

pressed significantly higher compared to cases without mutations in the TGF-β and BMP

pathway (p=0.044, respectively, Dunn test, Fig. 6a).   For *PAK3*, no significant overall expres-

sion differences were observed between *SMAD4* mutational groups (Fig. 6b).   We next

evaluated whether any of the expression biomarkers impacted on the survival outcome of

colorectal cancer using maximum selected rank statistics to define a single cut-point (Fig.

6b).   For example, *Smad4^{low}* expression (p=0.0019), *ID1^{low}* expression (p=0.0062), *SPP1^{High}*

(p= 0.0072) and *PAK3^{High}* (p=0.012) appeared to be associated with poorer overall survival

in the TCGA cohort (Kaplan-Meier, Fig. 6b).   The less significant impact of *PAK3^{High}* may re-

flect the relative low proportion of *PAK3^{High}* expressing sub-populations of cells within tu-

mours.

Discussion

The identification of cancer cells that have activation of the TGF-β-BMP-SMAD4

pathway in the context of pathway gene disruption requires validated models in order to

discover mechanism and pathway specific gene expression biomarkers [2]. Moreover, RNA

based molecular prognostic classifiers of colorectal cancer have developed in line with the

now clinically applied breast cancer equivalents. Such machine learning classifiers have

identified several clusters (e.g. CRISB) that include a stromal based signature that corre-

lates with poor prognosis in colorectal cancer [3,36,37]. A supervised machine learning CMS1-4

classifier (Consensus Molecular Classifier) has correlated gene level expression signatures

with four prognostic groups, where the poor prognostic group CSM4 was associated with

EMT and TGF-β dependency, yet the classifier does not utilise RNA expression biomarkers

that are specific readouts of the TGF-β-BMP-SMAD4 pathway. With respect to the latter,

human colorectal polyp derived organoids have been recently used to discover TGF-β de-

pendent gene sets in human serrated adenomas combined with a *BRAF^{V600E}* mutations,

where there was an induced mesenchymal phenotype, yet this model did not address spe-

cific *Smad4* loss of function [38]. Further analysis of human expression data has recently

identified three classes of pathway derived subtypes (PDS1-3) in colorectal cancer with and

without *KRAS* mutations, where CSM4 and CSM1 combine in the PDS2 subtype where TGF-

β-BMP-SMAD4 pathway expression profiles appear enriched [4]. These subtypes appeared

independent of *KRAS* driver gene variant classes and phenotypic appearance and for PDS1

and 2, correlated with RNA expression data from mouse genetically modified models. Thus,

gene-pathway specific gene expression biomarkers their dependencies are overall becom-

ing better understood and applied in intestinal cancer classifiers.

In a murine system, we discovered that the generation of homozygous *Smad4* with

Apc LOF alleles was associated with a discordant intestinal phenotype, with adenoma sup-

pression in the small intestine and adenoma progression in the caecum. By deriving or-

ganoid cultures from the adenoma, we compared control wild-type *Smad4* (*Smad$^{+/+}$*) with

Smad4 LOF (*Smad4$^{Δ/Δ}$*). We discovered that the high sensitivity to TGF-β1 induces similar

EMT and cell death phenotypes in a dose dependent manner, with a relative 10-fold dose

resistance to TGF-β1 in *Smad4$^{Δ/Δ}$* adenoma. Following TGF-β1 exposure, we identified

genes with extremes of differential expression by genotype, including reduced *Id1* and in-

creased *Spp1* gene expression as consistent features in *Smad4$^{Δ/Δ}$* murine adenoma organ-

oids. The gene expression changes as consistent with *Smad4* independent activity of the R-

Smads, in this case this is likely to be mainly Smad3 [39-41]. Single cell RNA sequencing of the

primary caecal adenomas detected enrichment of the *Pak3^{High}* expressing progenitor

epithelial cell population in *Smad4$^{Δ/Δ}$*. We assume that that the differentially expressed

genes are the main underlying difference that may account for functional discordant pro-

gression of adenoma, for example $Id1^{Low}$.

3   Mouse models of *Smad4* LOF in the intestine have been validated and have shown

that $Smad4^{+/-}$ mice spontaneously developed juvenile polyposis (JP) at 1.5 years, and inva-

sive adenocarcinoma when crossed with $Apc^{\Delta716/+}$ [42-45]. In addition, an

$Apc^{1638N/+}Smad4^{fl/fl}K19\text{-}CreERT2$ model showed an increased adenoma burden[46], and

CRISPR/Cas9-mediated genome editing of human intestinal epithelial organoid cultures

introduced mutations in *APC*, *TP53*, *KRAS* and *SMAD4* which, when combined and selected,

developed aggressive invasive carcinomas[23]. Moreover, *SMAD4* LOF blocked differentiation

during tumour progression in an orthotopic organoid transplantation model mimicking an

intestinal adenoma-carcinoma transition [47]. These results may also help to explain the se-

lection of deletions of chromosome arms 18p and 18q (including *SMAD4*) detected in 66%

of invasive colorectal cancers [48].

The generation of homozygous alleles in our conditional model results in rapid

adenoma formation within days, with animals reaching humane endpoints because of

anaemia and intestinal obstruction before the detection of invasive and metastatic sites.

The discordant pattern of adenoma development in $Apc^{fl/fl}Smad4^{fl/fl}$ suggests that the colon

may be more prone to adenoma initiation and progression when both genes are disrupted.

One possibility is that this may be due to the caecal inflammatory microenvironment and

the local TGF-β-BMP ligand supply. Our RNA-Seq analysis showed downregulation of

Pla2g2a (2.5log$_2$FC) in $Smad4^{\Delta/\Delta}$ adenoma. *Pla2g2a* is considered to be a tumour suppres-

sor independent of the Wnt/β-catenin signalling pathway, and intraluminal secretion of

Pla2g2a is a host defence mechanism during the active phase of ulcerative colitis [49]. More-

over, Reg3γ and *Reg3β* were also downregulated (1.2log$_2$FC) in $Smad4^{\Delta/\Delta}$ adenomas and as

both genes are important for epithelial defence against bacterial invasion [50,51]. Another

confounder may be modifier alleles co-segregating with LoxP alleles, although this was

mitigated by backcrossing to >10 generations of all conditional alleles to C57Bl/6J [27]. Ex-

cess TGF-β-BMP may also promote adenoma formation independent of *Apc* LOF, as tar-

geted deletion of *Bmpr1a* can lead to crypt expansion, fused villi and adenoma formation

due to an increase in β-catenin [52]. Overall, adenoma *Smad4* LOF may create the conditions

for local TGF-β1 ligand supply and further stimulation of the pathway, contributing to dif-

ferentiation arrest, chronic inflammation and subsequent adenoma progression.

The dose dependency of TGF-β1 ligand supply with respect to cell growth phenotype

has not been previously described to our knowledge, but may contribute functionally to

the manifestation of differential phenotypes with *Smad4* LOF; at a given dose, pathway gene expression positively modifies differentiation and EMT function, but at the same time there is resistance to cell death and cell cycle arrest. The time-dependent TGF-β1 on gene expression, using a fixed concentration of 390pM which exceeds the IC_{50} of $Smad4^{+/+}$ adenoma, resulted in differential expression of genes that were *Smad4* and TGF-β1 dependent, with the most significant being decreased expression of *Id1*. *Id1* codes for a non-DNA binding inhibitor of basic-helix-loop-helix (bHLH) transcription factors (e.g. GATA4) to inhibit multiple cell processes including differentiation. ChiP-Seq assays that have shown that Smad3 and Smad4 can bind the *Id1* promoter to directly stimulate early induction following TGF-β1 exposure [40]. Feedback *Id1* repression is then thought to be also pSmad3 dependent but through feedback activation of *ATF3*, that then directly represses *Id1* expression [53]. *Id1* over-expression in pancreatic ductal adenocarcinoma in mouse and human is also associated with more invasive and undifferentiated cells, a mechanism proposed to be dependent on EIF2 signalling [54] and independent of the TGF-β pathway[55]. Id1 proteins detected by IHC show increased expression with in colorectal cancer and mouse models, and one report (although the paper was subsequently retracted) showed that a genetic knock-out of *Id1* results in adenoma suppression in the small intestine of $Apc^{Min/+}$ but not when colitis is chemically induced, in support of an inflammatory basis of the caecal adenoma [56]. Id1 protein localisation has not been shown to correlate with gene expression, as we have observed, and so is unlikely to be a reliable functional biomarker of the pathway activity [56]. Recently, a Nur77 based molecular switch has also been proposed to regulate *Id1*, where stabilisation of TGF-β induced Smad3 degradation leads to an increase in *Id1* expression, whereas in the absence of TGF-β, Nur77 enhances Smad3 degradation and results in reduces *Id1* expression[57].

Spp1 (osteopontin) expression normally occurs in osteoclasts and macrophages, but has also been specifically detected in tumour associated M2 macrophages, with senescence associated secretary phenotype (SASP) epithelial tumour cells and at the invasive front of human colorectal tumours [58-62]. As well as extensive circumstantial data associating high *SPP1* expression in human colorectal cancer, *Spp1* was also previously reported to be a key *Smad4* dependent expressed gene functionally associated with murine prostatic cancer invasion in a *Pten* prostate deletion model [63]. High *SPP1* expression may have therapeutic implications, as immune exclusion correlates in the tumour microenvironment with cancer associated fibroblasts resulting in resistance to immune checkpoint therapy [64,65].

Pak3 is a well-known downstream effector of Cdc42 and Rac1 GTPases that mediates

motility and MAPK activation associated with cancer invasion and metastasis [66-68]. In a re-

cent adenovirus Cre-induced *Kras*^{G12D}*, p53*$^{fl/fl,}$ *Smad4*$^{fl/fl}$ model of lung cancer, increased me-

tastasis was associated with *Smad4* LOF and *Pak3* activation, the latter identified as a

downstream effector of *Smad4* via the PAK3-JNK-Jun pathway [68]. Interestingly, the pro-

posed mechanism appeared to be due to attenuation of *Smad4* dependent transcription of

miR-495 and miR-543 that can directly bind to the PAK3 3'UTR. Overexpression of PAK

family members are also associated with CRC outcome, transition of adenoma to carci-

noma and stabilisation of β-catenin [69-75].

Finally, having determined 76 differentially expressed TGF-β-BMP-SMAD4 gene-

pathway dependent candidate gene expression biomarkers in mouse, we evaluated

whether these same genes in humans were also differentially expressed with respect to

the presence or absence of SMAD4 pathogenic variants. At least 7 such DEGs were identi-

fied, several of which (*CREB3L1, BCAS1, ID1* and *ID3*) also appeared independent of com-

mon co-existing pathogenic variants in the RTK-RAS, TP53 and PI3K pathways. *SMAD4,*

ID1, SPP1 and *PAK3* mRNA expression also have prognostic implications for colorectal can-

cer using the TCGA cohorts, and are potential functional readouts of the TGF-β-BMP-

SMAD4 pathway. We provide evidence these genes, and potentially their associated

pathway gene sets, should be incorporated into future prospective evaluation of pathway-

based prognostic-predictive classifiers for the TGF-β-BMP-SMAD4 pathway in intestinal

cancer subtypes[4].

Materials and methods

Conditional Cre intestinal adenoma model

Work was performed under a UK Home Office licence, and approved by University of

Oxford Animal ethics committee. The mouse lines were backcrossed to C57Bl/6J as re-

ported including *Smad4[fl/fl]* (*Smad4[tm1Rob]*) (Supplementary Material and Methods). Genotyp-

ing   PCRs   were   briefly   (5'-3',   forward-F   and   reverse-R):   *Apc*   F-

GTTCTGTATCATGGAAAGATAGGTGGTC and R- CACTCAAAACGCTTTTGAGGGTTG (GAG-

TACGGGGTCTCTGTCTCAGTGAA   for   recombined   product),   *Smad4*   F-

CTTTTATTTTCAGATTCAGGGGTTC (F-AAAATGGGAA- AACCAACGAG for recombined product)

and R- TACAAGTGTATGTCTTCAGCG. Genotype combinations were generated by experi-

mental breeding. *Smad4[fl/fl]RosaYFP[fl/fl]* and *Apc[fl/fl]Smad4[fl/fl]RosaYFP[fl/fl]* were crossed with ei-

ther *Lgr5-CreER[T2]* or *Vil-CreER[T2]* mice. *Apc[fl/fl]Smad4[+/+]RosaYFP[fl/fl] Lgr5-CreER[T2]/Vil-CreER[T2]*

animals were used as controls. Mice bred with *Vil-CreER[T2]* mice were used for an *in vitro*

experiment only. For *Lgr5-CreER[T2]* activation, adult mice age ≥ 8 weeks had an intra-

peritoneal injection of 5mg (200 mg.kg[-1]) tamoxifen dissolved in corn oil. Mice were moni-

tored daily, small intestinal and colonic adenomas were scored for number, diameter, dis-

tribution and weight as described[27].

Intestinal adenoma organoids

Adenoma formed *in vivo* from *Smad[fl/fl]* or *Smad4[+/+]* Lgr5 mice were cultured as de-

scribed[27,76] . For crypt culture, the small intestines were collected from *Smad4[fl/fl]RosaYFP[fl/fl]*

and *Apc[fl/fl]Smad4[fl/fl]RosaYFP[fl/] Vil-CreER[T2]* mice and processed as described [76]. Briefly,

crypts were cultured in Matrigel and supplemented with ADF2 and crypt growth factors,

ENR (8.1 nM EGF (E), 500ng ml-1 R-spondin1 (R), and 2.16nM Noggin (N)) and incubated at

37°C, 5% as described. At D3, crypts generated from transgenic mice were incubated with

500nM 4-hydroxy-tamoxifen (4-OHT) (Millipore, 579002) dissolved in 100% EtOH (test),

vehicle (negative control) with 5μl EtOH for 12h at 37°C. The test and the vehicle treated

organoids were supplemented with EN only (no R) and the normal crypts with ENR. Pro-

teins were extracted from organoids as described [77].

Cell labelling and imaging

Neutral buffered formalin fixed tissues were paraffin embedded and 5μm sections

were processed either by immunohistochemistry (IHC) or immunofluorescence (IF). The

antibodies that used in this study are listed (Supplementary Materials and Methods). Or-

ganoids growing on coverslips in 24 well plates were used for IF. Organoid proliferation

used the thymidine analogue 5-ethynyl-2'-deoxyuridine (EdU) for 2 hrs using Click-iT EdU

Alexa Fluor-555 Imaging Kit (Invitrogen, C10338). H&E and IHC slides were viewed by

brightfield- fluorescent microscopy (Olympus BX60). Images were acquired by CCD camera

and Nuance 2.10 software (CRI, Massachusetts). Confocal images were captured with an

Olympus FluoView FV1000 using 40x 1.3NA lens and associated software. Labelled slides

(EdU, IHC or IF) were analysed using ImageJ 1.48 software. AlamarBlue (AB) (AbD Serotec,

BUF012B), was used to quantify adenoma growth. Absorbance after 24hr was read at

570nm and 600nm using a Spectramax M5 plate reader (MTX Lab Systems). Live cell imag-

ing was conducted with a Zeiss Axiovert 200 inverted microscope with a BSI Prime CMOS

camera with a Neofluar 10x 0.3NA lensfor adenoma organoids treated with TGF-β for

24hrs. MetaMorph software (v 7.8.12.0) multidimensional Z-stack acquisition with 100 mil-

liseconds exposure acquired at regular intervals of 15 minutes over an extended duration

of 6-18 hours culture at 37.5 °C, and 5% CO_2 in air.

TGF-β1 induced gene expression bulk RNASeq.

Early passage adenomas seeded at 9,000 cells per 25µl well were used for RNA-Seq

and miRNA-Seq. Adenomas were grown in ADF2 medium supplemented with 8.1nM EGF,

10µM Y-27632 and 1µM SB-505142. The medium was changed on D2, the cells were

washed, EGF and 390pM TGF-β1 added and samples extracted from three different time

points (t=0, t=1, t=12 hours post TGF-β1 treatment). Three biological replicates from each

genotype and RNA was pooled from 8 wells per treatment. RNA extraction was performed

using Quick-RNA MiniPrep (Zymo Research, R1054), assessed using a Nanodrop-1000 spec-

trophotometer and RNA Clean & concentrator[TM]-5 (Zymo Research, R1015). RNA-Seq and

miRNA-Seq were performed at the Wellcome Trust Center for Human Genetics, University

of Oxford. Briefly, directional multiplexed mRNA and miRNA Libraries were prepared. All

mRNA was ribo-depleted and converted to cDNA. Paired-end sequencing (100nt) were run

on a HiSeq 2500 (Illumina) for mRNA-Seq.

Transcript abundances from RNA-Seq data were quantified using the Tuxedo pipe-

line[78]. Fastq reads were aligned to the mouse genome (mm10/GRCm38) with the TopHat-

Bowtie2 aligner, versions 2.1.0 and 2.3.4, respectively and expression of transcripts was

quantified with Cufflinks 2.2.1, as fragments per kilobase per million mapped reads (FPKM).

Counts were also generated using the feature 'Counts' from the Subread package. Pairwise

differential expression analysis between groups was performed in R using DESeq2 version

1.28[79]. Genes that have less than 5 reads in at least 3 samples were removed. Genes with

adjusted p-values (FDR) < 0.05 were considered significant. Shrunk log2fold change

(log2FC) was calculated using as a shrinkage estimator. Genes considered as differentially

regulated between 2 conditions have ±1.5log2FC. Time-series analysis was performed us-

ing the likelihood ratio test implemented in DESeq2. Gene ontology term enrichment

analysis for the significant DEG genes performed using "clusterProfiler" where we evalu-

ated biological processes and molecular functions[80]. For easier visualisation, we used

"tmod"package for enrichment analysis (https://peerj.com/preprints/2420v1). In tmod, we

applied CERNOtest, Coincident Extreme Ranks in Numerical Observations (CERNO) after

converting mouse genes to human orthologs. We performed the enrichment analysis using

Hallmark gene sets from MSigDB v7.2 (Broad Institute) and the blood signature from LI

modules in tmod, gene sets with the area under the curve (AUC) $\geq$ 0.5 and padj<0.05 were

considered positive. Metascape website was utilised for pathway analysis of certain gene

lists, and to identify the transcription regulatory network based on TRRUST ontology[81].

Early TGF-β target genes were classified based on log2FC ±1.5 expression and ad-

justed p <0.05 after 1h compared to t=0 in *Smad4$^{+/+}$* adenomas. Transient early upregu-

lated/ downregulated genes were the genes with opposite directions to their initial regula-

tion when compared to late time points (t=12 vs t=0 and/ or t=12 vs t=1 hour) in *Smad4$^{+/+}$*

adenomas. Late TGF-β upregulated/downregulated genes were classified according to

their upregulation/downregulation at 12 hours in *Smad4$^{+/+}$* adenomas compared to t=0

and/or t=1 hour with log2FC ±1.5 and adjusted p <0.05. Genes were considered *Smad4*-

independent if they were significantly expressed in both *Smad4$^{+/+}$* and *Smad4$^{\Delta/\Delta}$* adenomas

with TGF-β1 treatment compared to their corresponding untreated baseline. However, if

genes were differentially expressed in an opposite direction to that of TGF-β target genes

in *Smad4$^{\Delta/\Delta}$* vs *Smad4$^{+/+}$* at t=0, they were classified as Smad4-dependent genes and ex-

cluded from the *Smad4$^{\Delta/\Delta}$* vs *Smad4$^{+/+}$* t=0 list. TGF-β target genes that were only

upregulated in *Smad4$^{+/+}$* adenomas were classified as *Smad4*-dependent genes. Genes that

were expressed in *Smad4$^{\Delta/\Delta}$* vs *Smad4$^{+/+}$* at t=0 hour and were not among TGF-β target

genes and considered for baseline comparisons.

Caecal adenoma single cell mRNA sequencing

Pooled single-cell suspensions from were obtained from caecal tumours from three

Apc$^{fl/fl}$,Smad4$^{+/+}$, Lgr5Cre^{ERT2} and five *Apc$^{fl/fl}$, Smad4$^{fl/fl}$, Lgr5Cre^{ERT2}* mice. Cell suspensions

underwent library preparation using the Chromium Single Cell 3' kit (10X Genomics) and

sequencing using Illumina HiSeq 2x150bp (Azenta, USA), with approximately 25,000 reads

per cell. Sample demultiplexing, barcode processing and single-cell 5☐ unique molecular

identifier (UMI) counting (Cell Ranger Software Suite v7.0.1) with FASTQ alignment to the

mouse reference transcriptome mm10-3.0.0. The following criteria were then applied to

each cell: gene number between 250 and 7,000, UMI count >500 and mitochondrial gene percentage <0.15, log10 Genes Per UMI >0.80. Doublets were removed using the scDblFinder library. Genes with ten read counts or less were removed. A total of 20886 cells were analysed (11212 for *Smad4^fl/fl^* and 9674 for *Smad4^+/+^*). Dimensionality reduction and clustering used the filtered gene-barcode matrix normalised using the SCTransform with Seurat (v4.3.0) [82]. Variables including mitochondria Ratio, nUMI, nGene and the cell cycle gene difference were regressed and the top 3000 variable genes were used. The filtered gene-barcode matrix of all samples were then integrated, k nearest neighbour (k-NN) generated using FindNeighbors (Seurat) using the top 40 principal components and clusters identified using FindCluster function (Louvain algorithm). We used resolution 1.2 to classify the different cell types and resolution 2.2 to determine the enterocyte precursors based on clustree (v0.5.1) [83]. Gene markers that identified each cluster relative to all other clusters was determined with FindAllMarkers function in Seurat using the Wilcox test (log-fold change threshold 0.25). Manual cell labelling of the top marker genes for each cluster utilised previous scRNA-seq studies [30,32,84]. Significant differences (p values, and confidence intervals via bootstrapping) in cell type proportion between each genotype was determined with the permutation test implemented in scProportionTest package in R (V4.2.3). Data was visualised using Uniform Manifold Approximation and Projection (UMAP) dimensional reduction and plots were generated using Seurat and the scCustomize library. Each cluster was determined using FindAllMarkers function in Seurat (Wilcox test). The log fold change threshold was set to 0.25. Manual cell labelling utilised gene clusters in previous studies [30-33], the Lgr5 intestinal stem cell signature [32] and the CellMarker 2.0 database [84]. The significant differences in cell type proportion used the permutation test implemented in scProportionTest package in R which calculate a p-value for each cluster, and a confidence interval for the magnitude difference via bootstrapping [85]. All analysis was performed in R V4.2.3. Differential gene expression analysis for each cluster between *Smad4^fl/fl^* and *Smad4+/+* were performed using MAST [86] with FindAllMarkers function in Seurat. Differentially expressed genes were selected based on log fold change threshold and expressed in at least in 25% in one of the groups. Pathway analysis was performed using Metascape [81]. Pseudo-bulk expression analysis was performed using MAST cell implemented in Seurat's FindMarkers function with the first and second identities using all *Smad4+/+* and *Smad4^fl/fl^* cells, respectively.

TCGA colorectal cancer analysis

Normalised gene expression from RNA-seq (illumine hiseq Level 3 RSEM normalized
genes) retrieved from http://firebrowse.org/?cohort=COADREAD accessed October 2019.
Genes and pathways that are frequently mutated in colorectal cancer, retrieved from Can-
cer Genome Atlas project (TCGA)[48], include the Wnt/β-catenin pathway (*APC, CTNNB1,*
DKK1, DKK2, DKK3, DKK4, LRP5, FZD10, FAM123B:AMER1, AXIN2, TCF7L2, FBXW7, and
ARID1A), TP53 pathway (*ATM, TP53*). PI3K (*IGF2, IRS2, PTEN, PIK3R1, and PIK3CA*), RTK-
RAS pathway (*NRAS, KRAS, BRAF, ERBB2, and ERBB3*). TGF-β pathway (*ACVR1B, ACVR2A,*
SMAD2, SMAD3, SMAD4, TGFBR1 and *TGFBR2*) and BMP pathway (*SMAD1, SMAD9,*
SMAD5, BMPR1A, BMPR1B and *ACVR1*). Genomic variants (missense mutation, homozy-
gote deletion, amplification or fusion gene) in these genes for COAD and READ TCGA co-
horts were retrieved from cBioPortal (https://www.cbioportal.org), accessed December
2019. In COAD and READ TCGA cohorts the variant status was available for 56 patients
with *SMAD4* variants. 98.1% of these patients had alterations in the Wnt/β-catenin path-
way, 59.3% had alteration in the TP53 pathway, 46.3% in the PI3K pathway and 64.8% had
variants in the RTK-RAS pathway. Normalized RNA-seq was available for 51 normal tissues
and 326 patient tumours with variant data. In TCGA COAD-READ cohorts, 44 patients with
probable pathogenic and possibly pathogenic SMAD4 variants, but with no other variants
in the TGF-β pathway, had RNA-seq data. One patient did not have a variant in the Wnt
pathway and was excluded from the analysis. Patients with known variants in the TGF-β
pathway were excluded from the control group. Using this patient group, we validated the
top 76 DEG in the mouse adenomas, genes with ±2log2FC and padj<0.05 from Smad4$^{\Delta/\Delta}$ vs
Smad4$^{+/+}$ (t=0 hr), either in the presence or absence of variants in the PI3K, TP53, PTK-RAS
pathways in the context of *SMAD4* variant. Statistical analysis was performed using
Kruskal-Wallis test followed by FDR adjustment. Genes with p<0.05 following FDR adjust-
ment were considered significant. Multiple comparisons were performed using Dunn test
followed by Holm adjustment to detect *SMAD4*-dependent genes. The expression be-
tween 2 groups were considered positive if adjusted p<0.05. We considered genes to be
SMAD4 dependent genes based on three criteria. First, there needs to be a significant dif-
ference in gene expression between the groups (Group 1-Group 3, and Group 1-Group 4,
Group 2-Group 3 and Group 2-Group 4) in the context of the three second pathways with
and without variants. Second, there should be no significant difference in gene expression
between Group 1-Group 2, and between Group 3 -Group 4, respectively. Third, the any
change in gene expression should follow the same direction as observed in the mouse ade-
nomas. Note that 84 genes differentially regulated at t=0 with absolute log2FC >=2 and

adjusted p<= 0.05. Of those, 76 genes had human orthologs, and 73 genes had RNA data in

the TCGA cohorts. Survival data for the COAD-READ TCGA cohort were retrieved (Table S1,

Tab TCGA-CDR, accessed May 2019[87]).

Statistical analysis

Statistical and graphical data analyses were performed using either Graph-Pad Prism

version 6.0. software or R version 4.0.2. Images were analysed using ImageJ 1.48 software.

Final Figures were generated using Photoshop (CS5). Immunoblot analysis was performed

as follow; loading control and test from scanned immunoblots was aligned horizontally on

Photoshop (CS5) and analyzed using ImageJ 1.48 software. For TCGA COAD-READ expres-

sion data, the cut-off point between high and low mRNA expression levels for each gene

was computed based on the maximally selected rank statistics (maxstat) using the

surv_cutpoint function from the 'survminer' package in R.

Acknowledgements

We thank Elisabeth Robertson for the conditional Smad4 model and Caroline Hill for dis-

cussion. Linda Randall and Nicky Hamp for mouse technical assistance.

Disclosure of potential conflicts of interest

None.

Author Contributions

Experiment conceived by ABH. Mouse breeding and phenotypic analysis by MS and JM,

imaging by MS, SB, and CB, bulk gene expression MS and DB, genotyping by MS, EC, and JH,

histopathology HM, evaluation of cancer cell model systems, MS, PM and CJI, single cell

sequencing analysis MS, TCGA analysis, MS, manuscript was written by ABH, MS and DB.

All authors approved the manuscript.

Grant Support

We thank Cancer Research UK (A429- ABH), Clarendon-Arab Bank scholarship programme

(MS), Rhodes Trust (JM) and Dunn School (ABH) for funding.

Figure Legends

Figure 1

Adenoma growth following conditional disruption of *Apc* and *Smad4* loxP alleles

using *Lgr5-CreER^{T2}*.

a. Breeding strategy to combine loxP conditional alleles with *Lgr5-CreER^{T2}* prior to

4OHT injection in adult mice. Kaplan-Meier survival of *Apc^{fl/fl}Smad4^{+/+}*,

Apc^{fl/fl}Smad4^{fl/+} and *Apc^{fl/fl}Smad4^{fl/fl}* mice post-Cre induction. Comparison between

Apc^{fl/fl}Smad4^{+/+} and *Apc^{fl/fl}Smad4^{fl/fl}* genotypes, *p=0.011, Log-rank test. Caecal

enlargement (arrow) from *Apc^{fl/fl}Smad4^{fl/fl}* mice. Bar 1cm. **b**. Caecum (adenoma)

weight, small intestine and colon total adenoma from *Apc^{fl/fl}Smad4^{+/+}* (n=9),

Apc^{fl/fl}Smad4^{fl/+} (n=17) and *Apc^{fl/fl}Smad4^{fl/fl}* (n=13). Horizontal bars; mean. One-way

ANOVA with Bonferroni's multiple comparison test. ***p<0.001. **c**. Haematoxylin

and Eosin (H&E) sections and anti-Smad4 antibody immunohistochemistry localisa-

tion in *Apc^{fl/fl}Smad4^{+/+}* and *Apc^{fl/fl}Smad4^{fl/fl}* caecal adenoma. Arrows show crypt

abscesses, upper panel. Smad4 localisation is reduced in adenoma and confined to

adjacent normal colonic crypt cells in *Apc^{fl/fl}Smad4^{fl/fl}* mice. Scale bar 100µm. **d.**

Western blot of adenoma (T) and normal tissue (N) from *Apc^{fl/fl}Smad4^{+/+}* and

Apc^{fl/fl}Smad4^{f/fl} with anti-phospho-Smad2 (p-Smad2) and anti-phospho-Smad1/5/8

antibodies. Densitometric analysis for p-Smad2 represented as ratio to β-actin (for

Apc^{fl/fl}Smad4^{+/+} n=3 and for *Apc^{fl/fl}Smad4^{fl/fl}* n=4 experimental replicates). Signifi-

cant increase in p-Smad2 in adenoma (T) compared to normal tissues (N) in

Apc^{fl/fl}Smad4^{fl/fl} mice. ***p<0.001. Non-parametric one-way ANOVA with Dunn's

multiple comparison test. Mean ± S.D.

Figure 2

TGF-β1 and Smad4 dependent signalling in intestinal adenoma organoids.

a. Expression of YPF reporter, E-cadherin localisation and DNA (DAPI) in

Apc^{Δ/Δ}Smad4^{+/+} and *Apc^{Δ/Δ}Smad4^{Δ/Δ}* adenoma organoids with and without exposure

to TGF-β1 (390pM). **b**. TGF-β1 (range: 0.039- 3900pM) induced growth inhibition of

adenoma organoids quantified by alamarBlue (measured at 24h. Data normalised

to untreated control (100%) (n=4 experimental replicates). Growth inhibition of

390pM TGF-β1 rescued by pre-treatment with SB-505124 (ALK4,5,7 inhibitor) for

2h (1mM). Error bars ± S.D. **c**. Representative immunoblots of Smad4, total R-

Smads and p-R-Smads from $Apc^{\Delta/\Delta}Smad4^{+/+}$ and $Apc^{\Delta/\Delta}Smad4^{\Delta/\Delta}$ adenoma organ-

oids with and without SB-505124. Graphs of p-Smad2/ Smad2 and p-Smad3/

Smad3 densitometry relative to β-actin (n=3 experimental replicates). Non-

parametric one way-ANOVA with Dunn's multiple comparison. **d**. Confocal im-

munofluorescence images of Smad2 and Smad3 nuclear localisation (DAPI) follow-

ing TGF-β1 (390pM) exposure and SB-505124 inhibition. Scale bar 50μm. Associ-

ated Immunoblots of p-Smad2 at different time points (h) following TGF-β1

(390pM), densitometry relative to β-actin. **e**. Bright field confocal images of organ-

oids 24-72 hr following TGF-β1 exposure. Red arrows intact spherical (circular) or-

ganoid morphology, yellow arrows show shape change, red dotted line indicates

organoids with marked disruption, cell spreading and cellular transformation.

Quantification shows relative preservation of spheroid (circular) organoids in

$Apc^{\Delta/\Delta}Smad4^{\Delta/\Delta}$. One-way ANOVA with Sidak's multiple comparison test

****p<0.0001. **f**. Suppression of organoid proliferation following TGF-β1 treatment

assessed with EdU incubation for 2h. Confocal images of EdU incorporation (green)

in nuclei (DAPI- blue) from $Apc^{\Delta/\Delta}Smad4^{+/+}$ and $Apc^{\Delta/\Delta}Smad4^{\Delta/\Delta}$ and quantification

using ImageJ (n=10 organoids). One-way ANOVA with Sidak's multiple comparison

test *** p<0.001, ****, p<0.0001, respectively. **g**. Cleaved Caspase 3 (CC3) activa-

tion following 24-72 hrs of TGF-β1 (390pM). CC3 (arrowheads) labelling in confocal

images supported by immunoblots. Arrows indicate margin of organoids. Scale bar

50μm.

Figure 3

TGF-β1 and Smad4 dependent bulk RNA expression signatures in adenoma or-

ganoids.

a. Adenoma organoids derived from 3 mice per genotype of $Apc^{\Delta/\Delta}Smad4^{+/+}$ and

$Apc^{\Delta/\Delta}Smad4^{\Delta/\Delta}$ were exposed to TGF-β1 (390pM) (0, 1 and 12h, eight wells per

time point), were then pooled for paired-end bulk RNA-Seq (total of 18 RNA-Seq

samples). **b**. Principal components of transcriptional clusters according to genotype

and time (vst-transformed). **c**. Venn diagram classifies TGF-β1 and Smad4 depend-

ent down-regulated and upregulated genes. Threshold utilised 1.5 log2FC and ad-

justed p-value <0.05. **d**. Volcano plots of differentially expressed genes comparing

time and genotype ($^{+/+}$=*Apc*$^{Δ/Δ}$*Smad4*$^{+/+}$, $^{Δ/Δ}$ = *Apc*$^{Δ/Δ}$*Smad4*$^{Δ/Δ}$). In all volcano plots,

genes with -log(p-adj)=30 are shown. Cut-off values for log2FC (1.5) and adjusted-p-

value (1e-3) are marked with dashed lines. *Apc*$^{Δ/Δ}$*Smad4*$^{Δ/Δ}$ DEGs are much reduced

in number following TGF-β1 treatment. **e**. Enriched gene modules of DEG using the

hallmark gene set enrichment (GSEA with tmod package). Comparisons shown in

each column. Red and blue indicate the proportion of genes in a module that are

either upregulated or downregulated, respectively. Width of each box relates to its

effect size, where lighter, less saturated colours indicate lower significance with p-

values. **f.** Heatmap showing time-dependent changes in expression (log2FC) for the

50 genes with the lowest adjusted p-values (Likelihood ration test) and **g**. pathway

enrichment for these genes are shown generated using Metascape. **h**. Summary of

gene specific expression comparing *Apc*$^{Δ/Δ}$*Smad4*$^{+/+}$ and *Apc*$^{Δ/Δ}$*Smad4*$^{Δ/Δ}$ with time

post TGF-β1 exposure. Statistical comparison performed using the Wald test in De-

seq2, followed by Benjamini and Hochberg multiple hypothesis correction.

Figure 4

Smad4 **dependent adenoma cell populations using single cell RNA-seq from** *in*

vivo **caecal adenoma.**

a. Pooled caecum adenoma cells derived from different mice, *Apc*$^{fl/fl}$*Smad4*$^{+/+}$ n=3

and *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$ n=5, were subject to scRNA-Seq. Cluster trees show the resolu-

tion from 0 to 1.2 with the EClpr and ECepr clusters identified at resolution 2.2. **b**.

Heatmap of the scaled expression of the top 5 genes identified in each cluster

(genes selected based on the biggest difference between the cluster compared to

all clusters). Lgr5 ICS = Lgr5 Intestinal stem cell; EC =enterocyte; EClpr =enterocyte

late precursor; ECepr =enterocyte early precursor; TA= transit amplifying; CAF

=cancer associated fibroblast; Secretory Pr =secretory precursor; hi =high referring

to specific genes. **c**. UMAP plot of the cell-type clusters from *Apc*$^{fl/fl}$*Smad4*$^{+/+}$ (9674,

n=3) and *Apc*$^{fl/fl}$*Smad4*$^{fl/fl}$ (11212, n=5) based on the expression of known marker

genes. **d**. Bar graph; frequencies of clusters of the single cells in *Apc*$^{fl/fl}$*Smad4*$^{+/+}$ and

Apc$^{fl/fl}$*Smad4*$^{fl/fl}$ adenoma. Point range plot; the confidence interval for the absolute

log2FC (obs_log2FC) for the different identified cell types. Significant results are

labelled in red. Note enrichment of *Pak3*^high^, *Slc14a1*^high^, CAF and neutrophil popu-

lations and the reduction in ECepr and EClpr populations in $Apc^{fl/fl}Smad4^{fl/fl}$ ade-

noma. Permutation test with FDR adjustment for p-value. **e.** UMAP plots compar-

ing genotypes for single cell expression of *ID1* (suppression in all cell types in

$Apc^{fl/fl}Smad4^{fl/fl}$), *Spp1* (monocytes and macrophage expression) and *Pak3* (in-

creased expression in progenitor enterocyte population of $Apc^{fl/fl}Smad4^{fl/fl}$). **f.** Violin

plots of pseudobulk RNA expression of *Id1*, *Spp1* and *Pak3*. Statistical analysis using

MAST with Bonferroni adjustment for p-value.

Figure 5

***SMAD4* pathogenic variant dependent expression biomarkers for the Smad4-TGF-**

β gene-pathway in human colorectal cancer (TCGA)

a. Flow diagram of TCGA bulk somatic DNA and RNA NGS sequencing data (COAD

=colon adenocarcinoma, READ =rectal adenocarcinoma) with known WNT pathway

driver variants (n=326). Pathogenic variants for TGF-β, activin or BMP receptors and

Smad 2/3/4/5/7/9 were identified (n=135) from wild-type cases (n=191). Sub-

groups (G1-4) were then assembled with or without co-existing common pathway

pathogenic variants in either RTK-RAS, TP53 or PI3K pathways. **b.** The human

equivalents of the 76 significantly differentially expressed genes obtained from the

comparison of bulk mRNA expression in the mouse adenoma were then compared

in the human TCGA mRNA data set. Volcano plot of median $\log_2$ fold-change

against adjusted p-value (-Log10) using Mann-Whitney U test (non-normal distribu-

tion using Shapiro-Wilk test) identified 7 human genes (in red) significantly differ-

ent in expression between Smad4 wild-type (n=191) and Smad4 pathogenic vari-

ants (n=44)[BCAS1, CACNB2, CREB3L1, ID1, ID3, RASGRF2 and SLC30A2]. **c., d.**

Comparison between sub-groups with and without context dependent variants G1-

G4 are shown as heat maps only for the significantly differentially expressed genes

(median RSEM expression $\log_{10}$). Non-parametric Kruskal-Wallis test (FDR adjust-

ment p<0.05) was followed by Dunn's test and Holm adjustment for multiple com-

parisons. 26 gene expression-context pairs were identified to be upregulated in the

TCGA cohort (c) and 15 gene expression-context pairs were significantly down regu-

lated (d). *Spp1* and *Pak3* median expression are shown separately but were non-

significant between groups (right heatmap). Of the comparison of the increased

expression of *CREB3L1 and BCAS1* and repression of *ID1* and *ID3,* only *ID1* met the

significance criteria.

Figure 6

***SMAD4, ID1, SPP1 and PAK3* expression biomarkers for the Smad4-TGF-β gene-**

pathway and human colorectal cancer (TCGA) survival.

a. Comparison of *ID1, SPP1 and PAK3 mRNA* expression in cases with somatic path-

ogenic variants in either *SMAD4* alone, *SMAD4* with other genes in the TGF-β/BMP

pathway (TGF-β *SMAD4* mutated), single gene variants in either the TGF-β or BMP

pathway (TGF-β or BMP mutated), more than one variant in TGF-β /BMP (TGF-β

and BMP mutated) or variants of BMP pathway genes alone (BMP mutated). Statis-

tical analysis was performed using Kruskal-Wallis test ($p < 0.0001$) followed by Dunn

test multiple comparison test adjusted by Holm test (shown in the plot). Note the

significant repression of *ID1* and increase in *SPP1* in *SMAD4* mutated cases. **b**. Kap-

lan-Meier survival curves showing survival differences between high and low gene

expression of *SMAD4*, *ID1*, *SPP1* and *PAK3* (left). Cut points were determined using

the maximally selected rank statistics (maxstat) method and are shown for each

gene (right). Statistical significance between the High vs low expression groups de-

termined with the Log-rank test (p values shown).

References

David, C. J. & Massague, J. Contextual determinants of TGFbeta action in development, immunity and cancer. *Nat Rev Mol Cell Biol* **19**, 419-435, doi:10.1038/s41580-018-0007-0 (2018).

Massague, J. & Sheppard, D. TGF-beta signaling in health and disease. *Cell* **186**, 4007-4037, doi:10.1016/j.cell.2023.07.036 (2023).

Dunne, P. D. & Arends, M. J. Molecular pathological classification of colorectal cancer-an update. *Virchows Arch*, doi:10.1007/s00428-024-03746-3 (2024).

Malla, S. B. *et al.* Pathway level subtyping identifies a slow-cycling biological phenotype associated with poor clinical outcomes in colorectal cancer. *Nat Genet*, doi:10.1038/s41588-024-01654-5 (2024).

Heldin, C.-H. & Moustakas, A. Role of Smads in TGFβ signaling. *Cell and tissue research* **347**, 21–36 (2012).

Macias, M. J., Martin-Malpartida, P. & Massague, J. Structural determinants of Smad function in TGF-beta signaling. *Trends Biochem Sci* **40**, 296-308, doi:10.1016/j.tibs.2015.03.012 (2015).

Inman, G. J., Nicolas, F. J. & Hill, C. S. Nucleocytoplasmic shuttling of Smads 2, 3, and 4 permits sensing of TGF-beta receptor activity. *Mol Cell* **10**, 283-294, doi:10.1016/s1097-2765(02)00585-3 (2002).

Guerrero-Martinez, J. A., Ceballos-Chavez, M., Koehler, F., Peiro, S. & Reyes, J. C. TGFbeta promotes widespread enhancer chromatin opening and operates on genomic regulatory domains. *Nat Commun* **11**, 6196, doi:10.1038/s41467-020-19877-5 (2020).

Inman, G. J. & Hill, C. S. Stoichiometry of active smad-transcription factor complexes on DNA. *J Biol Chem* **277**, 51008-51016, doi:10.1074/jbc.M208532200 (2002).

Massague, J. TGFbeta signalling in context. *Nat Rev Mol Cell Biol* **13**, 616-630, doi:10.1038/nrm3434 (2012).

Gaarenstroom, T. & Hill, C. S.   107–118 (Elsevier).

Massagué, J., Seoane, J. & Wotton, D. Smad transcription factors. *Genes & development* **19**, 2783–2810 (2005).

Fleming, N. I. *et al.* SMAD2, SMAD3 and SMAD4 mutations in colorectal cancer. *Cancer Res* **73**, 725-735, doi:10.1158/0008-5472.CAN-12-2706 (2013).

Gehart, H. & Clevers, H. Tales from the crypt: new insights into intestinal stem cells. *Nat Rev Gastroenterol Hepatol* **16**, 19-34, doi:10.1038/s41575-018-0081-y (2019).

Sato, T. *et al.* Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature* **469**, 415-418, doi:10.1038/nature09637 (2011).

Tan, S. H. *et al.* A constant pool of Lgr5(+) intestinal stem cells is required for intestinal homeostasis. *Cell Rep* **34**, 108633, doi:10.1016/j.celrep.2020.108633 (2021).

Beumer, J. *et al.* Enteroendocrine cells switch hormone expression along the crypt-to-villus BMP signalling gradient. *Nat Cell Biol* **20**, 909-916, doi:10.1038/s41556-018-0143-y (2018).

Beumer, J. & Clevers, H. Cell fate specification and differentiation in the adult mammalian intestine. *Nat Rev Mol Cell Biol* **22**, 39-53, doi:10.1038/s41580-020-0278-0 (2021).

Haramis, A.-P. G. *et al.* De novo crypt formation and juvenile polyposis on BMP inhibition in mouse intestine. *Science* **303**, 1684–1686 (2004).

Zhang, Y. & Que, J. BMP Signaling in Development, Stem Cells, and Diseases of the Gastrointestinal Tract. *Annu Rev Physiol* **82**, 251-273, doi:10.1146/annurev-physiol-021119-034500 (2020).

Winesett, M. P., Ramsey, G. W. & Barnard, J. A. Type II TGF(beta) receptor expression in intestinal cell lines and in the intestinal tract. *Carcinogenesis* **17**, 989-995, doi:10.1093/carcin/17.5.989 (1996).

Hahm, K. B. *et al.* Loss of transforming growth factor beta signalling in the intestine contributes to tissue injury in inflammatory bowel disease. *Gut* **49**, 190-198, doi:10.1136/gut.49.2.190 (2001).

Drost, J. *et al.* Sequential cancer mutations in cultured human intestinal stem cells. *Nature* **521**, 43-47, doi:10.1038/nature14415 (2015).

Barker, N. *et al.* Crypt stem cells as the cells-of-origin of intestinal cancer. *Nature* **457**, 608-611, doi:10.1038/nature07602 (2009).

Chu, G. C., Dunn, N. R., Anderson, D. C., Oxburgh, L. & Robertson, E. J. Differential requirements for Smad4 in TGFbeta-dependent patterning of the early mouse embryo. *Development* **131**, 3501-3512, doi:10.1242/dev.01248 (2004).

Costello, I., Biondi, C. A., Taylor, J. M., Bikoff, E. K. & Robertson, E. J. Smad4-dependent pathways control basement membrane deposition and endodermal cell migration at early stages of mouse development. *BMC Dev Biol* **9**, 54, doi:10.1186/1471-213X-9-54 (2009).

Matheson, J. *et al.* Epithelial-mesenchymal transition and nuclear beta-catenin induced by conditional intestinal disruption of Cdh1 with Apc is E-cadherin EC1 domain dependent. *Oncotarget* **7**, 69883-69902, doi:10.18632/oncotarget.11513 (2016).

Hassan, A. B. & Howell, J. A. Insulin-like growth factor II supply modifies growth of intestinal adenoma in Apc(Min/+) mice. *Cancer Res* **60**, 1070-1076 (2000).

Sato, T. *et al.* Long-term expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and Barrett's epithelium. *Gastroenterology* **141**, 1762-1772, doi:10.1053/j.gastro.2011.07.050 (2011).

Haber, A. L. *et al.* A single-cell survey of the small intestinal epithelium. *Nature* **551**, 333-339, doi:10.1038/nature24489 (2017).

Yan, K. S. *et al.* Non-equivalence of Wnt and R-spondin ligands during Lgr5(+) intestinal stem-cell self-renewal. *Nature* **545**, 238-242, doi:10.1038/nature22313 (2017).

Munoz, J. *et al.* The Lgr5 intestinal stem cell signature: robust expression of proposed quiescent '+4' cell markers. *EMBO J* **31**, 3079-3091, doi:10.1038/emboj.2012.166 (2012).

Grun, D. *et al.* Single-cell messenger RNA sequencing reveals rare intestinal cell types. *Nature* **525**, 251-255, doi:10.1038/nature14966 (2015).

Mustata, R. C. *et al.* Identification of Lgr5-independent spheroid-generating progenitors of the mouse fetal intestinal epithelium. *Cell Rep* **5**, 421-432, doi:10.1016/j.celrep.2013.09.005 (2013).

Ayyaz, A. *et al.* Single-cell transcriptomes of the regenerating intestine reveal a revival stem cell. *Nature* **569**, 121-125, doi:10.1038/s41586-019-1154-y (2019).

Bramsen, J. B. *et al.* Molecular-Subtype-Specific Biomarkers Improve Prediction of Prognosis in Colorectal Cancer. *Cell Rep* **19**, 1268-1280, doi:10.1016/j.celrep.2017.04.045 (2017).

Isella, C. *et al.* Stromal contribution to the colorectal cancer transcriptome. *Nat Genet* **47**, 312-319, doi:10.1038/ng.3224 (2015).

Fessler, E. *et al.* TGFbeta signaling directs serrated adenomas to the mesenchymal colorectal cancer subtype. *EMBO Mol Med* **8**, 745-760, doi:10.15252/emmm.201606184 (2016).

Fink, S. P., Mikkola, D., Willson, J. K. & Markowitz, S. TGF-beta-induced nuclear localization of Smad2 and Smad3 in Smad4 null cancer cell lines. *Oncogene* **22**, 1317-1323, doi:10.1038/sj.onc.1206128 (2003).

Liang, Y. Y., Brunicardi, F. C. & Lin, X. Smad3 mediates immediate early induction of Id1 by TGF-beta. *Cell Res* **19**, 140-148, doi:10.1038/cr.2008.321 (2009).

Zhang, Y. *et al.* High throughput determination of TGFbeta1/SMAD3 targets in A549 lung epithelial cells. *PLoS One* **6**, e20319, doi:10.1371/journal.pone.0020319 (2011).

Kitamura, T. *et al.* SMAD4-deficient intestinal tumors recruit CCR1+ myeloid cells that promote invasion. *Nat Genet* **39**, 467-475, doi:10.1038/ng1997 (2007).

Takaku, K. *et al.* Gastric and duodenal polyps in Smad4 (Dpc4) knockout mice. *Cancer Res* **59**, 6113-6117 (1999).

Takaku, K. *et al.* Intestinal tumorigenesis in compound mutant mice of both Dpc4 (Smad4) and Apc genes. *Cell* **92**, 645-656, doi:10.1016/s0092-8674(00)81132-0 (1998).

Takaku, K., Wrana, J. L., Robertson, E. J. & Taketo, M. M. No effects of Smad2 (madh2) null mutation on malignant progression of intestinal polyps in Apc(delta716) knockout mice. *Cancer Res* **62**, 4558-4561 (2002).

Freeman, T. J. *et al.* Smad4-mediated signaling inhibits intestinal neoplasia by inhibiting expression of beta-catenin. *Gastroenterology* **142**, 562-571 e562, doi:10.1053/j.gastro.2011.11.026 (2012).

Fumagalli, A. *et al.* Genetic dissection of colorectal cancer progression by orthotopic transplantation of engineered cancer organoids. *Proc Natl Acad Sci U S A* **114**, E2357-E2364, doi:10.1073/pnas.1701219114 (2017).

Cancer Genome Atlas, N. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **487**, 330-337, doi:10.1038/nature11252 (2012).

Fijneman, R. J. *et al.* Expression of Pla2g2a prevents carcinogenesis in Muc2-deficient mice. *Cancer Sci* **99**, 2113-2119, doi:10.1111/j.1349-7006.2008.00924.x (2008).

Dessein, R. *et al.* Toll-like receptor 2 is critical for induction of Reg3 beta expression and intestinal clearance of Yersinia pseudotuberculosis. *Gut* **58**, 771-776, doi:10.1136/gut.2008.168443 (2009).

Haapamaki, M. M. *et al.* Gene expression of group II phospholipase A2 in intestine in ulcerative colitis. *Gut* **40**, 95-101, doi:10.1136/gut.40.1.95 (1997).

He, X. C. *et al.* BMP signaling inhibits intestinal stem cell self-renewal through suppression of Wnt-beta-catenin signaling. *Nat Genet* **36**, 1117-1121, doi:10.1038/ng1430 (2004).

Kang, Y., Chen, C. R. & Massague, J. A self-enabling TGFbeta response coupled to stress signaling: Smad engages stress response factor ATF3 for Id1 repression in epithelial cells. *Mol Cell* **11**, 915-926, doi:10.1016/s1097-2765(03)00109-6 (2003).

Tang, Y., Zhang, S., Li, J., Wu, C. & Fan, Q. ID1 marks the tumorigenesis of pancreatic ductal adenocarcinoma in mouse and human. *Sci Rep* **12**, 13555, doi:10.1038/s41598-022-17827-3 (2022).

Huang, Y. H. *et al.* ID1 Mediates Escape from TGFbeta Tumor Suppression in Pancreatic Cancer. *Cancer Discov* **10**, 142-157, doi:10.1158/2159-8290.CD-19-0529 (2020).

Zhang, N. *et al.* Id1 Deficiency Protects against Tumor Formation in Apc(Min/+) Mice but Not in a Mouse Model of Colitis-Associated Colon Cancer. *Cancer Prev Res (Phila)* **8**, 303-311, doi:10.1158/1940-6207.CAPR-14-0411 (2015).

Niu, B. *et al.* Interplay between transforming growth factor-beta and Nur77 in dual regulations of inhibitor of differentiation 1 for colonic tumorigenesis. *Nat Commun* **12**, 2809, doi:10.1038/s41467-021-23048-5 (2021).

Li, N. *et al.* Mapping and modeling human colorectal carcinoma interactions with the tumor microenvironment. *Nat Commun* **14**, 7915, doi:10.1038/s41467-023-43746-6 (2023).

Lin, D. *et al.* Identification of a novel macrophage-related prognostic signature in colorectal cancer. *Sci Rep* **14**, 2767, doi:10.1038/s41598-024-53207-9 (2024).

Lv, H. *et al.* Comparative analysis of single-cell transcriptome reveals heterogeneity and commonality in the immune microenvironment of colorectal cancer and inflammatory bowel disease. *Front Immunol* **15**, 1356075, doi:10.3389/fimmu.2024.1356075 (2024).

Nakajima, T. *et al.* Osteopontin expression in the invasive front stroma of colorectal adenocarcinoma is associated with tumor budding and prognosis. *Pathol Res Pract* **240**, 154190, doi:10.1016/j.prp.2022.154190 (2022).

Ozato, Y. *et al.* Spatial and single-cell transcriptomics decipher the cellular environment containing HLA-G+ cancer cells and SPP1+ macrophages in colorectal cancer. *Cell Rep* **42**, 111929, doi:10.1016/j.celrep.2022.111929 (2023).

Ding, Z. *et al.* SMAD4-dependent barrier constrains prostate cancer growth and metastatic progression. *Nature* **470**, 269-273, doi:10.1038/nature09677 (2011).

Luo, H. *et al.* Pan-cancer single-cell analysis reveals the heterogeneity and plasticity of cancer-associated fibroblasts in the tumor microenvironment. *Nat Commun* **13**, 6619, doi:10.1038/s41467-022-34395-2 (2022).

Qi, J. *et al.* Single-cell and spatial analysis reveal interaction of FAP(+) fibroblasts and SPP1(+) macrophages in colorectal cancer. *Nat Commun* **13**, 1742, doi:10.1038/s41467-022-29366-6 (2022).

Gao, Z. *et al.* PAK3 promotes the metastasis of hepatocellular carcinoma by regulating EMT process. *J Cancer* **13**, 153-161, doi:10.7150/jca.61918 (2022).

Ma, T., Li, H., Yang, W., Liu, Q. & Yan, H. Over-expression of miR-193a-3p regulates the apoptosis of colorectal cancer cells by targeting PAK3. *Am J Transl Res* **14**, 1361-1375 (2022).

Tan, X. *et al.* Loss of Smad4 promotes aggressive lung cancer metastasis by de-repression of PAK3 via miRNA regulation. *Nat Commun* **12**, 4853, doi:10.1038/s41467-021-24898-9 (2021).

Carter, J. H. *et al.* Pak-1 expression increases with progression of colorectal carcinomas to metastasis. *Clin Cancer Res* **10**, 3448-3456, doi:10.1158/1078-0432.CCR-03-0210 (2004).

Gong, W. *et al.* P21-activated kinase 5 is overexpressed during colorectal cancer progression and regulates colorectal carcinoma cell adhesion and migration. *Int J Cancer* **125**, 548-555, doi:10.1002/ijc.24428 (2009).

Huynh, N. *et al.* Depletion of p21-activated kinase 1 up-regulates the immune system of APC(Δ14/+) mice and inhibits intestinal tumorigenesis. *BMC Cancer* **17**, 431, doi:10.1186/s12885-017-3432-0 (2017).

King, H., Nicholas, N. S. & Wells, C. M. Role of p-21-activated kinases in cancer progression. *Int Rev Cell Mol Biol* **309**, 347-387, doi:10.1016/B978-0-12-800255-1.00007-7 (2014).

Li, C. *et al.* High-level expression of P21-Cdc/Rac-activated kinase 7 is closely related to metastatic potential and poor prognosis of colon carcinoma. *Oncotarget* **7**, 46042-46055, doi:10.18632/oncotarget.10017 (2016).

Li, X. & Li, F. p21-Activated Kinase: Role in Gastrointestinal Cancer and Beyond. *Cancers (Basel)* **14**, doi:10.3390/cancers14194736 (2022).

Song, B., Wang, W., Zheng, Y., Yang, J. & Xu, Z. P21-activated kinase 1 and 4 were associated with colorectal cancer metastasis and infiltration. *J Surg Res* **196**, 130-135, doi:10.1016/j.jss.2015.02.035 (2015).

Sato, T. *et al.* Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* **459**, 262-265, doi:10.1038/nature07935 (2009).

Lee, G. Y., Kenny, P. A., Lee, E. H. & Bissell, M. J. Three-dimensional culture models of normal and malignant breast epithelial cells. *Nat Methods* **4**, 359-365, doi:10.1038/nmeth1015 (2007).

Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562-578, doi:10.1038/nprot.2012.016 (2012).

Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).

Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284-287, doi:10.1089/omi.2011.0118 (2012).

Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* **10**, 1523, doi:10.1038/s41467-019-09234-6 (2019).

Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821, doi:10.1016/j.cell.2019.05.031 (2019).

Zappia, L. & Oshlack, A. Clustering trees: a visualization for evaluating clusterings at multiple resolutions. *Gigascience* **7**, doi:10.1093/gigascience/giy083 (2018).

Hu, C. *et al.* CellMarker 2.0: an updated database of manually curated cell markers in human/mouse and web tools based on scRNA-seq data. *Nucleic Acids Res* **51**, D870-D876, doi:10.1093/nar/gkac947 (2023).

Miller, S. A. *et al.* LSD1 and Aberrant DNA Methylation Mediate Persistence of Enteroendocrine Progenitors That Support BRAF-Mutant Colorectal Cancer. *Cancer Res* **81**, 3791-3805, doi:10.1158/0008-5472.CAN-20-3562 (2021).

Finak, G. *et al.* MAST: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biol* **16**, 278, doi:10.1186/s13059-015-0844-5 (2015).

Liu, J. *et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. *Cell* **173**, 400-416 e411, doi:10.1016/j.cell.2018.02.052 (2018).

Smad4 and TGF-β1 dependent gene expression biomarkers in conditional intestinal adenoma, organoids and colorectal cancer.

Mirvat Surakhy[1], Julia Matheson[1], David Barnes[1,3], Emma J. Carter[1], Jennifer Hughes[1], Claudia Bühnemann[1], Sabina Sanegre[1], Hans Morreau[2], Paul Metz[1], Charlotte J. Imianowski[1] and A. Bassim Hassan[1,¶].

Figure 1

Smad4 and TGF-β1 dependent gene expression biomarkers in conditional intestinal adenoma, organoids and colorectal cancer.
Mirvat Surakhy[1], Julia Matheson[1], David Barnes[1,3], Emma J. Carter[1], Jennifer Hughes[1], Claudia Bühnemann[1], Sabina Sanegre[1], Hans Morreau[2], Paul Metz[1], Charlotte J. Imianowski[1] and A. Bassim Hassan[1,¶].

Figure 2

Smad4 and TGF-β1 dependent gene expression biomarkers in conditional intestinal adenoma, organoids and colorectal cancer.
Mirvat Surakhy[1], Julia Matheson[1], David Barnes[1,3], Emma J. Carter[1], Jennifer Hughes[1], Claudia Bühnemann[1], Sabina Sanegre[1], Hans Morreau[2], Paul Metz[1], Charlotte J. Imianowski[1] and A. Bassim Hassan[1,¶].

Figure 3

Smad4 and TGF-β1 dependent gene expression biomarkers in conditional intestinal adenoma, organoids and colorectal cancer.

Mirvat Surakhy[1], Julia Matheson[1], David Barnes[1,3], Emma J. Carter[1], Jennifer Hughes[1], Claudia Bühnemann[1], Sabina Sanegre[1], Hans Morreau[2], Paul Metz[1], Charlotte J. Imianowski[1] and A. Bassim Hassan[1,¶].

Figure 4

Smad4 and TGF-β1 dependent gene expression biomarkers in conditional intestinal adenoma, organoids and colorectal cancer.

Mirvat Surakhy[1], Julia Matheson[1], David Barnes[1,3], Emma J. Carter[1], Jennifer Hughes[1], Claudia Bühnemann[1], Sabina Sanegre[1], Hans Morreau[2], Paul Metz[1], Charlotte J. Imianowski[1] and A. Bassim Hassan[1,¶].

Figure 5

***Smad4* and TGF-β1 dependent gene expression biomarkers in conditional intestinal adenoma, organoids and colorectal cancer.**

Mirvat Surakhy[1], Julia Matheson[1], David Barnes[1,3], Emma J. Carter[1], Jennifer Hughes[1], Claudia Bühnemann[1], Sabina Sanegre[1], Hans Morreau[2], Paul Metz[1], Charlotte J. Imianowski[1] and A. Bassim Hassan[1,¶].

Figure 6