Dendritic Cell – Fibroblast Crosstalk via TLR9 and AHR Signaling Drives Lung Fibrogenesis

Hannah Carter [1], Rita Medina Costa [1], Taylor S. Adams [2], Talon Gilchrist [1], Claire E. Emch [1], Monica Bame [1], Justin M. Oldham [3], Angela L. Linderholm [4], Imre Noth [5], Naftali Kaminski [2], Bethany B. Moore [1,3], Stephen J. Gurczynski* [1].

1) Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI
2) Section of Pulmonary, Critical Care and Sleep Medicine, Yale University School of Medicine, New Haven, Connecticut, USA.
3) Division of Pulmonary and Critical Care Medicine, University of Michigan, Ann Arbor, MI
4) Division of Pulmonary and Critical Care Medicine, University of California, Davis, California.
5) Division of Pulmonary and Critical Care Medicine, University of Virginia, Charlottesville, Virginia.

* Corresponding author:

Stephen J. Gurczynski, PhD

5608C Medical Science II

1137 Catherine St

Ann Arbor, MI 48109-2200

sgurczyn@umich.edu

(734) 647-9968 phone

(734)615-2331 fax

The Authors declare no conflicts of interest.

Funding sources. This work was funded by NIH R35 HL144481 awarded to BBM, T32GM007863 awarded to HC, and a Boehringer Ingelheim discovery award in ILD. This work was made possible by the Pulmonary Fibrosis Foundation, by an independent grant from Boehringer Ingelheim Pharmaceuticals, Inc. who provided the financial support. The authors meet criteria for authorship as recommended by the International Committee of Medical Journal Editors (ICMJE) and were fully responsible for all aspects of the trial and publication development.

Author contributions. HC performed experiments, analyzed data, wrote, and edited the manuscript. TSA analyzed and interpreted data. RMC, TG, CE, and MB performed experiments and analyzed data. JMO, ALL, and IN provided clinical data. BBM analyzed and interpreted data and edited the manuscript. SJG wrote and edited the manuscript as well as analyzing and interpreting data.

Word count: 5387

Abstract

Idiopathic pulmonary fibrosis (IPF) is characterized by progressive scarring and loss of lung function. With limited treatment options, patients succumb to the disease within 2-5 years. The molecular pathogenesis of IPF regarding the immunologic changes that occur is poorly understood. We characterize a role for non-canonical aryl-hydrocarbon receptor signaling (ncAHR) in dendritic cells (DCs) that leads to production of IL-6 and IL-17, promoting fibrosis. TLR9 signaling in myofibroblasts is shown to regulate production of TDO2 which converts tryptophan into the endogenous AHR ligand kynurenine. Mice with augmented ncAHR signaling were created by crossing floxed AHR exon-2 deletion mice ($AHR_{\Delta ex2}$) with mice harboring a CD11c-Cre. Bleomycin was used to study fibrotic pathogenesis. Isolated CD11c+ cells and primary fibroblasts were treated ex-vivo with relevant TLR agonists and AHR modulating compounds to study how AHR signaling influenced inflammatory cytokine production. Human datasets were also interrogated. Inhibition of all AHR signaling rescued fibrosis, however, $AHR_{\Delta ex2}$ mice treated with bleomycin developed more fibrosis and DCs from these mice were hyperinflammatory and profibrotic upon adoptive transfer. Treatment of fibrotic fibroblasts with TLR9 agonist increased expression of TDO2. Study of human samples corroborate the relevance of these findings in IPF patients. We also, for the first time, identify that AHR exon-2 floxed mice retain capacity for ncAHR signaling.

Abstract word count 211

Introduction

Idiopathic Pulmonary Fibrosis (IPF) is a progressive, fibrotic disease with estimated survival of 2-5 years after diagnosis. Risk factors include male sex, cigarette smoking, repeated viral infection, with median age at diagnosis of 66 years.(1) Treatments slow progressive fibrosis, but only lung transplant is curative.(2) Fibroblast activation and excess production of collagens are critical factors that underlie disease progression, however, fibrogenesis is a complex and poorly understood process requiring cooperation / crosstalk between multiple cell subsets. Myriad inflammatory cells accumulate in fibrotic lungs including dendritic cells (DCs), macrophages, and lymphocytes, however, the ultimate role of the immune system in initiation and progression of fibrosis is not understood. Immune responses may promote progressive scarring but treating patients with steroids is counterintuitively harmful highlighting the need for more research into the interactions between immune cell populations and fibrogenesis.(3) We have previously defined an important role for tissue resident DCs, and more specifically DCs expressing the aryl-hydrocarbon receptor (AHR) in viral-mediated pulmonary fibrosis following bone marrow transplantation.(4, 5) Tissue resident DCs have been shown to accumulate in the lungs of both IPF patients and bleomycin treated mice, however their exact role in the initiation or progression of pulmonary fibrosis is unclear. (6-8)

The canonical Aryl-Hydrocarbon Receptor (AHR) pathway is an intracellular signaling cascade originally described as the major sensor of a poly-cyclic aromatic hydrocarbon (PAH) toxicants, e.g., dioxin.(9) More recently, endogenous AHR ligands such as the kynurenine family of tryptophan metabolites have been described.(10, 11) Kynurenine can be made by indoleamine-2,3-dioxygenase (IDO) or tryptophan 2,3-dioxygenase (TDO), and kynurenine / tryptophan ratios are increased in patients with fibrotic lung disease.(12) Increased kynurenine / tryptophan ratios in the blood are also prognostic indicators for many inflammatory lung diseases, including cancers and SARS-CoV-2.(13-18)

AHR is constitutively expressed in an inactive form sequestered in the cytoplasm bound to various chaperone proteins.(19) Binding of ligand exposes a nuclear localization signal which facilitates translocation of AHR to the nucleus wherein the chaperone coat is recycled and AHR dimerizes with its major canonical partner ARNT.(20, 21) This AHR / ARNT dimer binds to xenobiotic response elements in the genome and activates transcription of a variety of gene products including the CYP family of cytochrome p450 oxidases.(22)

AHR is present and involved in inflammation in several lung cell types including immune

cells such as DCs, macrophages, and lymphocytes as well as structural cells such as epithelial

cells and fibroblasts. Recently AHR has been studied for its interactions with other binding

partners outside of ARNT, i.e., NFkB subunits RelA / RelB and KLF6, in what is termed "non-

canonical AHR signaling" (ncAHR) which is especially important in regulating expression of pro-

inflammatory cytokines in immune cells like dendritic cells (DCs) and macrophages.(4, 23-25)

Production of pro-inflammatory cytokines by DCs is augmented in the presence of AHR ligands

suggesting a role for AHR in concert with NFkB when triggered by pro-inflammatory signals.(26,

27) AHR and TLR signaling, especially TLR9, have also been studied separately for their ability

to promote fibrogenesis by directly augmenting fibroblast differentiation into pro-fibrotic

myofibroblasts.(28, 29) Pro-inflammatory TLR9 / AHR responsiveness has been linked to

worsened IPF progression(30), however, the functional binding partners of AHR in inflammatory

vs normal conditions, the triggers which divert AHR from its canonical pathway, and the exact

role of AHR in complex pathologies such as fibrosis are not well understood.

Here, we present data that ncAHR signaling is an important modulator of DC immune

responses during the progression of pulmonary fibrosis. Using the bleomycin mouse model of

pulmonary fibrosis, we demonstrate that AHR+ DCs accumulate in the lungs and drive

production of IL-6 in an AHR dependent fashion. AHR plays a role in differentiation of several

immune cells, including Treg, Th-17 and Th-22 cells(31-33), which makes whole-body knock

outs non-ideal for studying immune responses. Using a DC-specific deletion of the DNA binding

region of AHR (CD11c-Cre / AHR$_{\Delta ex2}$), we further demonstrate deletion of canonical AHR

function in DCs leads to significant upregulation of ncAHR signaling, pro-inflammatory cytokines

and worsened fibrosis in the bleomycin model. This work also demonstrated that adoptive

transfer of ncAHR-activated DCs can promote fibrogenesis in a CD4 T cell-independent

manner, highlighting an unappreciated role for DCs in fibrogenesis. We also provide evidence

that myofibroblasts in the lungs of both IPF patients and fibrotic mice express the enzymes that

convert tryptophan into endogenous AHR ligands highlighting an important role for TDO2

specifically in the fibrogenic circuit that is established between structural cells and immune cells

in the fibrotic lung.

Materials and Methods

Mouse line, cell lines, and reagents. All experiments were conducted under an IACUC

approved animal protocol. WT C57Bl/6J mice were ordered from Jackson labs and used

between 6-8 weeks of age. AHR$_{\Delta ex2}$ mice were generated by crossing the AHR^{fx} line

(Ahrtm3.1Bra/J, strain 6203, Jackson labs) with a CD11c-Cre expressing line (B6.Cg-Tg(Itgax-

cre)1-1Reiz/J, strain 8068, Jackson labs). Resulting pups were verified to contain both Cre and

the AHR floxed alleles via PCR using Jackson labs recommended primers. CD11c+ cells were

further verified to contain the AHR exon2 excised allele via PCR using Jackson labs

recommended primers. AHR$_{\Delta ex2}$ mice were bred at the University of Michigan mouse breeding

facility and used between 6-8 weeks of age. For some experiments AHR^{fx} mice which did not

express CD11c-Cre were used as littermate controls. Bleomycin was purchased as Blenoxane

from the pharmacy at the University of Michigan hospital, 1U/mL stocks were made using sterile

saline and mice were administered 0.75U/kg via the oral-pharyngeal route. The AHR and TDO2

inhibitors CH223191 and 680C91 (Tocris) were used at 5 mg/kg and 15mg/kg respectively for *in*

vivo experiments, CH223191 was used between 10-30 µM for *in vitro* treatment of cell culture.

The TiPARP (PARP7) inhibitor RBN2397 (Selleckchem) was used at a concentration of 10 µM.

The TLR9 agonist ODN2395 (Invivogen) was used between 0.1 – 1.0 µM.

Generation of inducible iCD103+ DCs. Inducible CD103 (iCD103) DCs were generated from

mouse bone marrow as previously described.(4, 34) In short, bone marrow cells were

aseptically isolated from the femurs of indicated mouse lines and incubated for 16-days in RPMI

media containing 10% heat inactivated FBS, 1% Pen/Strep cocktail, 50µM β-mercapto-ethanol,

3 ng/mL GMCSF (R&D systems) and 200 ng/mL Flt3-L (R&D systems) before any additional

treatment. iCD103 cells were characterized to be CD11c+, MHCII+, and CD103+ by flow

cytometry. For adoptive transfer experiments, Groups of WT-B6 mice were first administered

BLM before receiving 1x10^6 iCD103+ cells from AHR$_{\Delta ex2}$ mice via tail vein injection. CD4 T-cell

depletion was carried out via i.p. injection of 100 µg anti-CD4 depleting antibody (clone GK1.5

Bioxcell) at 0- and 7-days post BLM instillation.(35)

Fibroblast isolation. Groups of mice were administered BLM or saline as control. 21-days post

BLM, lungs were harvested and minced with scissors. Minced lungs were incubated in complete

DMEM (DMEM + 10% FBS and 1% pen/strep cocktail) for 7 days to let fibroblasts adhere.

Media was changed at 7 days and every 2 days afterwards. Cells were trypsonized and plated

on day 14-post isolation for subsequent assays.

Single Cell RNA-seq analysis. Processed gene expression data from Habermann et al.(36)

was downloaded from GSE135893. Raw sequencing data from Adams et al.(37) was

downloaded from GSE136831 for reprocessing. Read 2 poly(A) and template-switch oligo

contaminants were trimmed using cutadapt (version 3.1). Reads were aligned to genome

GRCh38 with STAR (version 2.7.8a) using Gencode annotation release 38.

Control and IPF cells previously classified as stromal were filtered from each dataset and

combined in R (version 4.0.3) using the library Seurat (version 4.3.0.1). For normalization, raw

expression counts were scaled to 10,000 transcripts per cell then natural log transformed with a

pseudocount of 1. Cells were split by dataset and integrated using Seurat's reciprocal principal

component analysis (RPCA) implementation, followed by dimension reduction, graph

embedding, high granularity clustering and cell type re-annotation. These steps were performed

iteratively until all multiplets were identified and removed.

Of the filtered data presented: Habermann et al. consists of 2,801 cells from 10 controls and 12

IPF lungs; Adams et al. consists of 6,850 cells from 26 control and 32 IPF lungs. Integrated

gene expression values were only used for PCA-based graph embedding; cluster interrogation

and subsequent visualizations were performed with normalized, empirical values. For the

heatmap, normalized gene expression values of each cell type were averaged for each subject

with 3 or more respective cells. To avoid batch effect between datasets when visualizing gene

expression, normalized expression values were further min-max normalized across cells within

each dataset.

Generation of single cell suspensions and flow cytometry. Single cells suspensions were

generated via collagenase digestion of lung tissue as previously described.(38) In short, mouse

lungs were first perfused via injection of 3-5mL of saline through the right ventricle of the heart.

Lungs were resected and minced with scissors before incubation in a collagenase buffer

containing 1 mg/mL Collagenase-A (Roche), 2000U DNaseI (Sigma), in complete DMEM media

supplemented with 10% heat inactivated FBS and 1% Pen/Strep cocktail for 45 minutes at 37°C

in a shaking incubator. Cell suspensions were disrupted by drawing through a 10 mL syringe

before filtration through a 100 μm pore sized cell strainer. Resulting single cell suspensions

were assayed for cell viability via trypan blue exclusion, staining for flow cytometry was

subsequently carried out via incubation with appropriately diluted fluorophore conjugated

antibodies. Antibodies used for flow cytometry were as follows: Myeloid flow cytometry staining

panel, BV650-CD11b (clone M1/70), BV510-CD45 (clone 30-F11), BV421-I-Ab (MHCII clone

AF6-120.1), APC-Cy7-SiglecF (clone E50-2440), APC-Ly6G (clone 1A8) purchased form BD

Horizon. BV605-CD64 (clone X54-5/7.1), PerCP-Cy5.5-CD24 (clone M1/69), PE-Cy7-Ly6C

(clone HK1.4), PE-CD193 (CCR3 clone J073E5) purchased from Biolegend. PE-eFluor610-

CD11c (clone N418), PE-AHR (clone 4MEJJ) purchased from eBioscience. Lymphocyte flow

cytometry staining panel. PE-Cy7-IFNγ (clone XMG1.2), FITC-CD3 (clone 17A2) purchased

from BD bioscience. APC-Cy7-IL17A (clone TC11-18H10.1), BV570-CD8a (clone 53-6.7),

AF700-CD90.2 (Thy1.2 clone 30-H12), BV510-CD4 (clone GK1.5) purchased from Biolegend.

PE-eFluor 610-FoxP3 (clone FJK-16s) purchased from eBioscience. Note: Some markers were

stained for but were not included in this study. Flow analysis was carried out in FlowJo v. 10.5.

Quantification of collagen via hydroxyproline assay. Collagen was quantified as described

in(39). In short, lungs from BLM or WT mice were resected at the indicated timepoints and

homogenized in a buffer of PBS and complete protease inhibitor (Roche). Lung homogenates

were diluted 1:1 with 12N hydrochloric acid and incubated overnight at 120^{o}C in well-sealed

glass tubes. The following day, aliquots of acid-hydrolyzed lung homogenate were incubated

with a chloramine-T solution and then reacted with 4-(Dimethylamino)benzaldehyde.

Hydroxyproline was detected via absorbance at 560 nM on a synergy H1 spectrophotometer

(Biotek).

Isolation of RNA for qRT-PCR. RNA was purified from lung tissue or cell pellets via incubation

with Trizol reagent following manufacturer directions. RNA quantity and purity was assessed on

a nanodrop spectrophotometer (Thermo). Transcript expression was analyzed via reverse

transcriptase quantitative real-time PCR using Lunaprobe reagent (New England Biolabs) on a

Quantstudio3 thermocycler (ABI). Expression analysis was conducted using a ΔΔct calculation.

Transcripts were normalized to expression level of a housekeeping gene, RPL38, using a pre-

validated primer probe set (Integrated DNA technologies, Hs.PT.58.40595235.gs). Other primer

and probe sequences are detailed in Table 1.

Analysis of plasma concentrations of IL-6 and IL-17 isoforms. Circulating plasma

concentration for IL6 and IL17 (a, c, d and f isoforms) was determined used the Olink (Uppsala,

Sweden) Explore platform, which uses proximity extension assays to generate semi-quantitative

protein data, which are Log2 transformed for modeling. The association between plasma

biomarker concentration and three-year transplant-free survival was assessed in a combined

cohort of patients with IPF (n=366) from the University of Virginia, University of Chicago and

University of California at Davis using univariable Cox proportional hazards regression [IRB

approval and subject characteristics were previously reported(40)]. Survival was plotted for each

quartile of IL6 concentration using the Kaplan-Meier estimator. The proportional hazards

assumption was checked and satisfied for each biomarker tested.

Statistical analysis. Statistical analysis was conducted in Graphpad Prism v.9.1.0. Statistical

significance was determined using students T-test when two groups were compared and

ANOVA using Holm-Sidak's post-test when three or more groups were compared. Relevant

statistical information, i.e., P-values and n-values, are given in individual figure legends.

Results

CD103+ DCs accumulate in the fibrotic lung, promote fibrogenesis and show evidence of

pro-inflammatory AHR signaling. We previously published that AHR+ CD103+ DCs were

recruited to lungs, and drove lung injury and fibrosis following bone marrow transplantation and

herpesvirus infection(4); however, a pathogenic role for AHR and DCs in non-infectious forms of

pulmonary fibrosis is not known. To address this, we characterized lung DCs using a bleomycin

model of pulmonary fibrosis. Lung leukocytes were analyzed via flow cytometry at 7- and 14-

days post-treatment and a significant increase in CD103+ DCs (CD45+, CD11c+, MHCII +,

CD103+, CD11b-) was noted at both time points (Fig. 1A). Differentiation of CD103+ DCs

requires the transcription factor BATF3 and CD103+ DCs are completely absent in BATF3[-/-]

mice.(41) To clarify the role of lung CD103+ DCs in mediating fibrogenesis we treated BATF3[-/-]

mice with bleomycin for 14-21 days and verified the absence of CD103+ DCs in the lung by flow

cytometry (Fig. 1B). BATF3[-/-] mice exhibited durable protection from bleomycin induced

pulmonary fibrosis exhibiting less collagen production at 14 days as assessed by hydroxyproline

assay and at 21-days as assessed by qRT-PCR for collagen I transcript (Fig. 1C and 1D).

Interestingly, BATF3[-/-] mice also displayed less IL-6 transcript indicating that CD103+ DCs are

important regulators of pro-inflammatory / fibrotic cytokine production following bleomycin

challenge (Fig. 1E). We next analyzed expression of AHR in various lung immune cells via flow

cytometry at 7-days post bleomycin treatment (markers for cell subsets are detailed in Sup.

Table 1). The highest percentage of AHR+ cells was found in CD103+ DCs (cDC1s ~60%)

while most other immune cells had AHR+ rates between 3-5% (Fig. 1F). To further verify that

expression of AHR was augmented in lung DCs we isolated CD11c+ cells via magnetic bead

isolation from the lungs of bleomycin treated or control mice. CD11c+ cells from the lungs of

bleomycin treated mice had increased expression of AHR in comparison to control mice but

interestingly had decreased expression of the canonical AHR gene product Cyp1b1 (Fig. 1G

and 1H). In agreement with our previous data using BATF3[-/-] mice, CD11c+ cells isolated from

bleomycin treated mice had increased expression of IL-6 further indicating that lung DCs are an

important source of pro-inflammatory cytokines (Fig. 1I).

Production of AHR ligands is augmented during BLM-induced fibrosis and complete AHR

inhibition ameliorates disease. Several endogenous ligands for AHR have been described,

most notably, kynurenine which is a metabolite of tryptophan processed by the heme-dependent

oxygenases IDO1/2 and TDO2.(42) Whole lung homogenates were prepared from saline-

treated or BLM-instilled mice. Concentrations of lung-associated kynurenine were assayed via

ELISA. A significant increase in lung-associated kynurenine was observed at both 7- and 14-

days post-BLM (Fig. 2A). These data suggest activation of AHR signaling in the fibrotic lung

may promote development of BLM-induced fibrosis. To test this hypothesis, we administered

the complete AHR inhibitor, CH223191, following BLM instillation. Complete inhibition of AHR

signaling ameliorated development of bleomycin-induced pulmonary fibrosis. Mice treated with

CH223191 exhibited less collagen deposition than non-treated mice (Fig. 2B) and had

decreased evidence of fibrotic pathology on trichrome stained lung sections (Fig. 2C).

Deletion of the AHR DNA binding domain reduces canonical AHR signaling in DCs but

AHR expression is retained and augments inflammatory non-canonical AHR signaling. To

further address AHR signaling in lung DCs we bred mice that harbored floxed copies of AHR

exon-2 (43) crossed to a strain that harbored Cre recombinase under control of the CD11c

promoter (CD11c-AHR$_{\Delta ex2}$ mice referred to as AHR $_{\Delta ex2}$, Fig. 3A). Previous reports using these

AHR exon-2 floxed mice described them as AHR-null mutants (43); however, we identified an

alternative start codon (Supplemental Fig. 1A red text) that is introduced as part of the fusion of

AHR exon-1 and exon-3 resulting in a 14 amino acid N-terminal leader sequence (Supplemental

Fig. 1A green text) fused in frame to AHR exons 3 – 10 (Supplemental Fig. 1A blue text). DCs

generated from these mice are indeed null for production of canonical AHR response genes

following stimulation with kynurenine (Fig. 3B), however, primers specific for AHR exon3-4

showed no difference in the amounts of AHR transcript while primers specific for the junction of

exon1-2 showed a significant reduction in AHR$_{\Delta ex2}$ mice (Fig. 3C). Furthermore, AHR was still

detectable in CD103+ DCs produced from AHR$_{\Delta ex2}$ mice as assessed by intracellular flow

cytometry (Fig. 3D and 3E).

We next generated cDNA from BMDCs isolated from either heterozygous B6 / AHR $_{\Delta ex2}$ or

homozygous AHR$_{\Delta ex2}$ mice. PCR of the AHR ORF produced two bands in the heterozygous

mice of approximately 2400 and 2200 bp (Supplemental Fig. 1B) corresponding to the predicted

sizes of the full length AHR or AHR$_{\Delta ex2}$ ORF, respectively, and a single band of approximately

2200bp in the AHR$_{\Delta ex2}$ mice. We cloned these products into pl-Cherry-NEO vector which co-

expressed each construct with mCherry and verified sequences of both full length and AHR$_{\Delta ex2}$

PCR products via sanger sequencing (data not shown) . To verify the specificity of the AHR flow

cytometry antibody, We conducted flow cytometry on 3T3 cells transfected with either AHR full

length or AHR$_{\Delta ex2}$ constructs and noted ~30% of cells expressed mCherry after 48 hours

(Supplemental Fig. 1C). We also examined expression of AHR in these transfected cells and

noted a ~2-fold increase in AHR MFI in both full length and AHR$_{\Delta ex2}$ transfections indicating both

constructs produced an in frame AHR protein. Thus, we now report that although canonical

AHR signaling is ablated following excision of exon-2, a truncated AHR protein is produced that

likely mediates effects through the ncAHR pathway.

To assess impact of DC AHR signaling on development of fibrosis we administered BLM to

groups of WT B6 or AHR$_{\Delta ex2}$ mice. AHR$_{\Delta ex2}$ mice exhibited increased fibrosis and ~1.5-fold

increased collagen deposition 21-days post BLM versus WT mice (Fig. 3F and 3G). We were

surprised by this result as complete AHR inhibition ameliorated fibrosis (Fig. 1C). We next

administered CH223191 to groups of BLM-treated WT or AHR$_{\Delta ex2}$ mice. Interestingly, complete

inhibition of AHR was still ameliorative (Fig. 3F) indicating that, while deletion of AHR exon-2

effectively ablated canonical AHR signaling in CD11c+ cells (Fig 3B), ncAHR signaling was

likely still functional and could mediate pro-fibrotic effects.

Loss of canonical AHR signaling in CD11c+ cells exacerbates fibrosis via a ncAHR-

dependent mechanism. To further test the role AHR in CD103+ DCs we generated pure

populations of induced CD103+ DCs (iCD103) from bone marrow stem cells.(34) We previously

published iCD103 DCs express many cDC1 markers and express AHR.(4) We addressed the

role of CD103+ DCs in mediating fibrosis by adoptively transferring iCD103+ DCs from AHR$_{\Delta ex2}$

mice into groups of BLM treated WT mice. Lungs were harvested at 21-days post BLM and

collagen content was assessed via hydroxyproline assay. B6 BLM treated mice that had

received an adoptive transfer of AHR$_{\Delta ex2}$ iCD103+ cells developed increased fibrosis exhibiting

significantly increased lung collagen over mice receiving a sham transfer (Fig. 4A). We also

administered neutralizing αCD4 antibody to BLM-treated mice with AHR$_{\Delta ex2}$ iCD103+ adoptive

transfer, however, no change in fibrosis was observed indicating the pro-fibrotic effects of

AHR$_{\Delta ex2}$ CD11c+ cells are independent of CD4+ T-cells(Fig. 4A).

The role of AHR in modulating cytokine responses is complex. Genes such as TiPARP

induced through the canonical AHR pathway downregulate cytokine and interferon responses

through post-translational modification of critical kinases such as TBK1 thus shutting down

NFκB and IRF responses.(44-47) In contrast, AHR itself can directly bind NFκB and drive pro-

inflammatory cytokine responses.(27, 48-50) Multiple TLRs, including TLR9, have been

implicated in the progression of pulmonary fibrosis.(29, 51, 52) To address if DC IL-6 expression

was dependent on cooperation between AHR and TLR signaling, we stimulated iCD103 DCs

with the TLR9 agonist ODN2395. Cells generated from AHR$_{\Delta ex2}$ mice expressed over 2-fold

more IL-6 in comparison to ODN2395-treated iCD103 cells from WT mice (Fig. 4B).

Interestingly, while ODN treatment could stimulate expression of TiPARP in WT iCD103 DCs,

this effect was abrogated in AHR$_{\Delta ex2}$ iCD103 DCs (Fig 4C). Together, these data indicate

expression of TiPARP is dependent on canonical AHR signaling while expression of IL-6 is

augmented by ncAHR signaling. We next treated iCD103 cells with TiPARP inhibitor RBN2397

and observed expression of IL-6 was increased by ~2.5-fold when TiPARP was inhibited

indicating production of TiPARP by canonical AHR signaling significantly inhibits non-canonical

signaling and decreases production of inflammatory IL-6 (Fig. 4D). We next examined levels of

IL-6 in BLM treated WT or AHR$_{\Delta ex2}$ mice that had been treated with the complete AHR inhibitor

CH223191. In agreement with the increased IL-6 we observed in iCD103 DCs, AHR$_{\Delta ex2}$ mice

exhibited increased expression of IL-6 both in whole lung and in CD11c+ cells isolated by

magnetic bead purification from single cell lung digestions (Fig. 4E and 4F). Expression of IL-6

could be diminished by AHR inhibition (CCH223191) both at the whole lung and CD11c+ cell

level indicating a direct effect of AHR in CD11c+ cells in potentiating IL-6 production in the

fibrotic lung (Fig. 4E and 4F). Furthermore, we verified that expression of TiPARP was similarly

down-regulated in lung CD11c+ cells from AHR$_{\Delta ex2}$ mice and thus conclude that the increased

IL-6 observed is a product of the loss of anti-inflammatory canonical AHR signaling and a

concomitant increase in pro-inflammatory non-canonical AHR signaling.

DC ncAHR signaling potentiates IL-17 production from lymphocytes. Given that AHR$_{\Delta ex2}$

mice developed increased fibrosis and that ex-vivo generated DCs produced increased IL-6 in

response to TLR9 stimulus, we next examined if AHR$_{\Delta ex2}$ mice would over-produce IL-17 in

response to BLM. We generated lung single cell suspensions at 7-days post-BLM and analyzed

expression of IL-17 in lymphocyte populations via flow cytometry. Flow cytometric analysis

revealed AHR$_{\Delta ex2}$ mice displayed increases in multiple IL-17 producing lymphocyte populations

including conventional Th17 cells, IL-17A producing gamma-delta T-cells, and putative IL-17A+

iNKT cells and ILC3s (Fig 5A quantified in 5B). Although we determined CD4+ lymphocytes

were not responsible for increased fibrosis (Fig. 4A), we noted the largest increase in IL-17A

expressing lymphocytes was in CD4- γδ TCR+ populations which would not be affected by CD4

depletion. AHR$_{\Delta ex2}$ mice expressed ~5-fold more IL-17 at 21-days post-BLM (Fig. 5C).

Production of both IL-6 and IL-17 was susceptible to complete inhibition of AHR via

administration of CH223191 (Fig. 4E, 4F and Supplemental Fig. 2) indicating DCs and

macrophages in CD11c-Cre / AHR$_{\Delta ex2}$ mice still had functional ncAHR signaling. We further

verified the involvement of CD103+ DCs by examining expression of IL-17 in the lungs of BLM

treated BATF3$^{-/-}$ mice and found that loss of CD103+ DCs almost completely abrogated lung IL-

17 responses (Fig. 5D).

Fibroblasts from fibrotic lungs express kynurenine producing enzymes in response to TLR9 stimulus. Endogenous AHR ligands are largely produced as a result of the metabolism of tryptophan along the kynurenine pathway by enzymes such as IDO1 and TDO2. To determine what cells produce these enzymes in fibrotic lungs, we investigated data from two previously published single-cell RNAseq studies of IPF.(36, 37) Over 9000 stromal cells from 36 controls and 44 lungs were collectively analyzed and reclassified based on the markers and nomenclature described in Adams et al.(37) Though IDO1 was undetected, expression of TDO2 was observed upregulated specifically in IPF alveolar fibroblasts across both datasets (Fig. 6A and Supplemental Fig. 3). In addition to myeloid cells, TLR9 activation in fibroblasts has also been shown to be important in driving fibrosis.(29, 53) Thus, we next isolated fibroblasts from mouse lungs on d-21 post-Saline or BLM and subsequently treated these cells with ODN2395 to simulate an inflammatory / fibrotic environment *ex vivo*. Expression of both IDO1 and TDO2 increased in ODN2395-treated fibroblasts from BLM-treated mice but not in ODN2395-treated control fibroblasts (Fig. 6B and 6C). AHR was also upregulated in the same conditions (Fig 6D); however, TLR9 was upregulated with ODN2395 regardless of BLM treatment (Fig 6E). We further assayed for expression of the main enzymes involved in the production of kynurenine, i.e., IDO1 and TDO2, in whole lung following BLM treatment. Interestingly, expression of IDO1 was decreased at both 7- and 14-days post BLM (Fig. 7A); however, TDO2 expression was significantly increased at both time points (Fig. 7B) thus it appears that TDO2 is preferentially induced over IDO1 in pulmonary fibrosis *in vivo*. To address the importance of TDO2 during fibrosis we next treated groups of mice with the specific TDO2 inhibitor 680C91. Strikingly, inhibition of TDO2 significantly decreased both collagen deposition and expression of the myofibroblast marker alpha smooth muscle actin (αSMA) during BLM induced pulmonary fibrosis as well fibrotic pathology as assessed by trichrome staining of lung sections (Fig. 7C, 7D, and 7E).

Discussion

Role of DCs in regulating pulmonary fibrosis. The role of myeloid cells, especially DCs, in altering immune responses during development / exacerbation of pulmonary fibrosis is currently not well understood. Increased numbers of DC subsets infiltrate lungs of IPF patients and mice with pulmonary fibrosis.(8, 54-56) However, the study of DCs in pulmonary fibrosis has been overlooked due to publication of seminal work demonstrating that CD4+ and CD8+

lymphocytes, and by extension the DCs that prime those responses, are not important in

mediating fibrotic effects.(57) Recently, it is appreciated that DCs can modulate immune

responses outside of antigen presentation through production of pro-inflammatory / fibrotic

cytokines. We, and others, have demonstrated that lung DCs can prime and activate pathogenic

IL-17 responses through production of inflammatory cytokines such as IL-6 and TGF-β in innate

lymphocytes as well as CD4+ T cells.(5, 58)

Here we demonstrate CD103+ DCs infiltrate the fibrotic lung and mediate production of

pro-inflammatory / fibrotic cytokines through cooperation of AHR and pro-inflammatory TLR

signaling. We noted significant accumulation of CD103+ DCs in lungs of BLM-treated mice

beginning in the early inflammatory phase (7-days post-treatment) and continuing into the

fibrotic phase (21-days post treatment, Fig. 1F). Loss of CD103+ DCs in BATF3$^{-/-}$ mice was

protective indicating a pathogenic role for lung resident DCs (Fig. 1B – 1E). In agreement with

our previously published study(4), CD103 cells that infiltrate the fibrotic lung express high levels

of AHR (Fig. 1F). Canonical AHR signaling is immuno-suppressive and AHR target genes such

as TiPARP can constrain production of interferons and pro-inflammatory cytokines via ADP-

ribosylation of activators of NF-κB signaling such as TBK1.(44-47) Alternatively, ncAHR

signaling is well characterized to bind NF-κB subunits like RelA and RelB and enhance

production of pro-inflammatory cytokines.(27, 48-50) We noted increased expression of IL-6

following BLM both in the whole lung and from isolated CD11c+ cells which correlated with

increased expression of AHR (Fig. 1); however, expression of the AHR inducible gene product,

Cyp1b1 was significantly decreased (Fig. 1H). Thus, even though CD11c+ cells express AHR

and there is abundant AHR ligand produced in fibrotic lungs (Fig. 2A), these cells do not

activate canonical AHR signaling and instead activate pro-inflammatory ncAHR signaling in the

BLM treated lungs.

Role of ncAHR / NF-Kb and TIPARP in modulating cellular cytokine responses. The role of

tryptophan metabolism and AHR signaling in mediating pulmonary fibrosis is unclear.

Kynurenine / tryptophan ratios increase in serum and lungs of patients with interstitial

pneumonia and treatment with anti-fibrotic Pirfenidone significantly decreased levels of

kynurenine suggesting a positive correlation between tryptophan metabolism, AHR activation,

and development of fibrosis.(12, 59) More recently, activation of AHR through exogenous

administration of AHR ligand FICZ ameliorated pulmonary fibrosis presumably by augmenting

numbers of FoxP3+ regulatory T-cells, and decreasing IFNγ+ CD4+ T-cells and IL-17A+ γδ+ T-

cells.(60) However, this study did not look specifically at AHR signaling in pulmonary immune

cells nor did it demonstrate if FICZ is produced during fibrosis. Furthermore, mice were treated

directly after administration of BLM on days 0, 1, and 2 making it difficult to determine direct

effects of pulmonary AHR signaling in inhibiting fibrosis versus modulating early lung injury post-

BLM. We detected significantly elevated kynurenine from 7 – 14 days post-BLM corresponding

to the early fibrotic phase of BLM induced fibrosis (Fig. 2A). In contrast to previous work,

complete inhibition of AHR signaling via administration of CH223191 reduced fibrosis when

started late in the progression of disease (10-days post-BLM) indicating a role for pulmonary

AHR signaling in promoting fibrogenesis (Fig. 2B – 2C).

We generated mice that lacked canonical AHR signaling in CD11c+ cells including

pulmonary DCs and macrophage (Fig. 3A and Supplemental Fig. 1). Interestingly, while DCs

produced from these mice lacked canonical AHR signaling (Fig. 3B), we still detected AHR at

the transcript and protein level (Fig. 3C – 4E). Furthermore, sequencing cDNA generated from

AHR$_{\Delta ex2}$ DCs indicated an alternative start codon is formed when exon 2 is deleted

(Supplemental Fig. 1A). Together, these data indicated that a truncated AHR product still

mediated ncAHR interactions. When we administered BLM to AHR$_{\Delta ex2}$ mice we noted more

fibrosis that was still AHR-dependent (Fig. 3F and 3G) indicating loss of canonical AHR

signaling in CD11c+ cells resulted in increased ncAHR signaling that exacerbated pulmonary

fibrosis.

To determine if increased fibrosis seen in AHR$_{\Delta ex2}$ was intrinsic to DCs we adoptively

transferred ex-vivo generated iCD103 DCs from AHR$_{\Delta ex2}$ mice into BLM treated B6 mice and

noted more fibrosis when DCs from CD11c-Cre+ / AHR$_{\Delta ex2}$ were transferred in comparison to

either Cre- cells (data not shown) or non-transferred mice (Fig. 4A). We further show AHR can

synergize with TLR9 activation and promote expression of IL-6 in DCs; however, expression of

canonical AHR genes that limit cytokine production, e.g., TiPARP is lost in AHR$_{\Delta ex2}$ DCs (Fig 4B

– 4D). Thus, BLM-induced fibrosis results in increased expression of AHR; however, the

signaling is shunted towards non-canonical signaling resulting in increased expression of pro-

inflammatory cytokines. It is currently unclear why ncAHR signaling is promoted over canonical

signaling in fibrotic lungs, however, several transcription factors such as HIF1α which are up-

regulated during fibrogenesis bind to the canonical AHR partner ARNT (HIF1β).(61) Thus, it is

plausible that dimerization of HIF1α-ARNT may reduce the available pool of ARNT which

promotes AHR-NFκB dimerization shifting the balance of AHR signaling from canonical, anti-

inflammatory, to non-canonical, inflammatory, signaling which drives excessive fibrogenesis.

Role of IL-6 in promoting profibrotic IL-17 responses. The role of the IL-6 – IL-17 axis is appreciated in the progression of pulmonary fibrosis. Depletion of either IL-6 or IL-17 is protective for BLM-induced fibrosis and we have previously published that IL-17 can act directly on fibroblasts to promote collagen production and fibroproliferation.(5, 62-64) Even in IPF patients, expression of IL-6 and IL-17 isoforms are associated with more rapid time to death or lung transplantation (Supplemental Fig. 3).

In agreement with this we isolated fibroblasts from control or BLM-treated mice and treated with the TLR9 agonist, ODN2395. TLR9 activation upregulated expression of both IDO1 and TDO2 in BLM fibroblasts but not control (WT) fibroblasts, we corroborated expression of TDO2 in IPF patient myofibroblasts, however, we did not detect appreciable expression of IDO1 in human myofibroblasts and overall expression of IDO1 was shown to decrease in the lungs of fibrotic mice while expression of TDO2 increased (Fig. 6A – 6C, 7A and7B). Treatment of mice with the TDO2 inhibitor 680C91 ameliorated fibrosis, reducing collagen production, and decreasing expression of the myofibroblast marker αSMA (Fig. 7C, 7D, and 7E). Furthermore, we noted increased expression of AHR and TLR9 itself following TLR9 stimulation in BLM fibroblasts indicating that AHR plays a fundamentally pro-fibrotic role (Fig. 6D – 6E). However, In direct contrast to this, a previous study found that treatment of fibroblasts with the AHR activating compound ITE inhibited TGFβ induced myofibroblast differentiation and collagen production.(28) The exact cause for the discrepancy between these two studies will require further investigation, however, it is plausible that induction of AHR signaling in the context of inflammatory stimulus, e.g., TLR stimulation as would happen in the fibrotic lung behaves fundamentally different than in an isolated *ex vivo* condition. Expression of TLR9 is associated with rapid progression of IPF in patients and loss of TLR9 signaling is protective from BLM-induced fibrosis.(29, 51, 52) Furthermore, fibroblasts isolated from IPF patients that had more robust activation profiles when exposed to TLR9 agonist correlated with increased AHR activation in matched myeloid cells.(30)

Together, these studies indicate that a fibrogenic circuit is established between TLR9+ fibroblasts and AHR+ myeloid cells that is important in progression of pulmonary fibrosis through modulation of pro-fibrotic cytokines like IL-6 and IL-17 (Working model in Fig. 8). More work is needed to fully understand the switch to ncAHR signaling during fibrogenesis and to understand the spatial crosstalk between fibroblasts and DCs. However, given the need for new treatments for IPF patients, drugs that modulate kynurenine pathway enzymes, i.e., IDO1 and TDO2, or AHR directly represent an exciting possibility with untapped therapeutic potential.

References

1. Raghu G, Collard HR, Egan JJ, Martinez FJ, Behr J, Brown KK, et al. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med.* 2011;183(6):788-824.

2. Chambers DC, Cherikh WS, Harhay MO, Hayes D, Hsich E, Khush KK, et al. The International Thoracic Organ Transplant Registry of the International Society for Heart and Lung Transplantation: Thirty-sixth adult lung and heart-lung transplantation Report-2019; Focus theme: Donor and recipient size match. *J Heart Lung Transplant.* 2019;38(10):1042-55.

3. Douglas WW, Ryu JH, and Schroeder DR. Idiopathic pulmonary fibrosis: Impact of oxygen and colchicine, prednisone, or no therapy on survival. *Am J Respir Crit Care Med.* 2000;161(4 Pt 1):1172-8.

4. Gurczynski SJ, Pereira NL, Hrycaj SM, Wilke C, Zemans RL, and Moore BB. Stem cell transplantation uncovers TDO-AHR regulation of lung dendritic cells in herpesvirus-induced pathology. *JCI Insight.* 2021;6(2).

5. Zhou X, Loomis-King H, Gurczynski SJ, Wilke CA, Konopka KE, Ptaschinski C, et al. Bone marrow transplantation alters lung antigen-presenting cells to promote TH17 response and the development of pneumonitis and fibrosis following gammaherpesvirus infection. *Mucosal Immunol.* 2016;9(3):610-20.

6. Misharin AV, Morales-Nebreda L, Mutlu GM, Budinger GR, and Perlman H. Flow cytometric analysis of macrophages and dendritic cell subsets in the mouse lung. *Am J Respir Cell Mol Biol.* 2013;49(4):503-10.

7. Marchal-Somme J, Uzunhan Y, Marchand-Adam S, Kambouchner M, Valeyre D, Crestani B, et al. Dendritic cells accumulate in human fibrotic interstitial lung disease. *Am J Respir Crit Care Med.* 2007;176(10):1007-14.

8. Tsoumakidou M, Karagiannis KP, Bouloukaki I, Zakynthinos S, Tzanakis N, and Siafakas NM. Increased bronchoalveolar lavage fluid CD1c expressing dendritic cells in idiopathic pulmonary fibrosis. *Respiration; international review of thoracic diseases.* 2009;78(4):446-52.

9. Piwarski SA, and Salisbury TB. The effects of environmental aryl hydrocarbon receptor ligands on signaling and cell metabolism in cancer. *Biochem Pharmacol.* 2023;216:115771.

10. Poland A, Glover E, and Kende AS. Stereospecific, high affinity binding of 2,3,7,8-tetrachlorodibenzo-p-dioxin by hepatic cytosol. Evidence that the binding species is receptor for induction of aryl hydrocarbon hydroxylase. *J Biol Chem.* 1976;251(16):4936-46.

11. Adams S, Braidy N, Bessede A, Brew BJ, Grant R, Teo C, et al. The kynurenine pathway in brain tumor pathogenesis. *Cancer Res.* 2012;72(22):5649-57.

12. Meyer KC, Arend RA, Kalayoglu MV, Rosenthal NS, Byrne GI, and Brown RR. Tryptophan metabolism in chronic inflammatory lung disease. *J Lab Clin Med.* 1995;126(6):530-40.

13. Mandarano M, Orecchini E, Bellezza G, Vannucci J, Ludovini V, Baglivo S, et al. Kynurenine/Tryptophan Ratio as a Potential Blood-Based Biomarker in Non-Small Cell Lung Cancer. *Int J Mol Sci.* 2021;22(9).

14. Wang M, Dong X, Huang Y, Su J, Dai X, Guo Y, et al. Activation of the kynurenine pathway is associated with poor outcome in Pneumocystis pneumonia patients infected with HIV: results of 2 months cohort study. *BMC Infect Dis.* 2019;19(1):223.

15. Zinellu A, Fois AG, Zinellu E, Sotgiu E, Sotgia S, Arru D, et al. Increased kynurenine plasma concentrations and kynurenine-tryptophan ratio in mild-to-moderate chronic obstructive pulmonary disease patients. *Biomark Med.* 2018;12(3):229-37.

16. Almulla AF, Supasitthumrong T, Tunvirachaisakul C, Algon AAA, Al-Hakeim HK, and Maes M. The
tryptophan catabolite or kynurenine pathway in COVID-19 and critical COVID-19: a systematic
review and meta-analysis. *BMC Infect Dis.* 2022;22(1):615.
17. Cihan M, Doğan Ö, Ceran Serdar C, Altunçekiç Yıldırım A, Kurt C, and Serdar MA. Kynurenine
pathway in Coronavirus disease (COVID-19): Potential role in prognosis. *J Clin Lab Anal.*
2022;36(3):e24257.
18. D'Amora P, Silva IDCG, Budib MA, Ayache R, Silva RMS, Silva FC, et al. Towards risk stratification
and prediction of disease severity and mortality in COVID-19: Next generation metabolomics for
the measurement of host response to COVID-19 infection. *PLoS One.* 2021;16(12):e0259909.
19. Henry EC, and Gasiewicz TA. Transformation of the aryl hydrocarbon receptor to a DNA-binding
form is accompanied by release of the 90 kDa heat-shock protein and increased affinity for
2,3,7,8-tetrachlorodibenzo-p-dioxin. *Biochem J.* 1993;294 (Pt 1)(Pt 1):95-101.
20. Whitelaw M, Pongratz I, Wilhelmsson A, Gustafsson JA, and Poellinger L. Ligand-dependent
recruitment of the Arnt coregulator determines DNA recognition by the dioxin receptor. *Mol Cell
Biol.* 1993;13(4):2504-14.
21. Wang GL, Jiang BH, Rue EA, and Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-
helix-PAS heterodimer regulated by cellular O2 tension. *Proc Natl Acad Sci U S A.*
1995;92(12):5510-4.
22. Okino ST, Pendurthi UR, and Tukey RH. Phorbol esters inhibit the dioxin receptor-mediated
transcriptional activation of the mouse Cyp1a-1 and Cyp1a-2 genes by 2,3,7,8-
tetrachlorodibenzo-p-dioxin. *J Biol Chem.* 1992;267(10):6991-8.
23. Wright EJ, De Castro KP, Joshi AD, and Elferink CJ. Canonical and non-canonical aryl hydrocarbon
receptor signaling pathways. *Curr Opin Toxicol.* 2017;2:87-92.
24. Wilson SR, Joshi AD, and Elferink CJ. The tumor suppressor Kruppel-like factor 6 is a novel aryl
hydrocarbon receptor DNA binding partner. *J Pharmacol Exp Ther.* 2013;345(3):419-29.
25. Ho CC, Lee HL, Chen CY, Luo YH, Tsai MH, Tsai HT, et al. Involvement of the cytokine-IDO1-AhR
loop in zinc oxide nanoparticle-induced acute pulmonary inflammation. *Nanotoxicology.*
2017;11(3):360-70.
26. Vogel CF, Khan EM, Leung PS, Gershwin ME, Chang WL, Wu D, et al. Cross-talk between aryl
hydrocarbon receptor and the inflammatory response: a role for nuclear factor-κB. *J Biol Chem.*
2014;289(3):1866-75.
27. Kado S, Chang WLW, Chi AN, Wolny M, Shepherd DM, and Vogel CFA. Aryl hydrocarbon receptor
signaling modifies Toll-like receptor-regulated responses in human dendritic cells. *Arch Toxicol.*
2017;91(5):2209-21.
28. Lehmann GM, Xi X, Kulkarni AA, Olsen KC, Pollock SJ, Baglole CJ, et al. The aryl hydrocarbon
receptor ligand ITE inhibits TGFbeta1-induced human myofibroblast differentiation. *Am J Pathol.*
2011;178(4):1556-67.
29. Trujillo G, Meneghin A, Flaherty KR, Sholl LM, Myers JL, Kazerooni EA, et al. TLR9 differentiates
rapidly from slowly progressing forms of idiopathic pulmonary fibrosis. *Science translational
medicine.* 2010;2(57):57ra82.
30. Huang Y, Ma SF, Espindola MS, Vij R, Oldham JM, Huffnagle GB, et al. Microbes Are Associated
with Host Innate Immune Response in Idiopathic Pulmonary Fibrosis. *Am J Respir Crit Care Med.*
2017;196(2):208-19.
31. Wang F, Liu M, Ma D, Cai Z, Liu L, Wang J, et al. Dendritic cell-expressed IDO alleviates
atherosclerosis by expanding CD4. *Int Immunopharmacol.* 2023;116:109758.
32. Rankin LC, Kaiser KA, de Los Santos-Alexis K, Park H, Uhlemann AC, Gray DHD, et al. Dietary
tryptophan deficiency promotes gut RORγt. *Cell Rep.* 2023;42(3):112135.

33. Holloman BL, Cannon A, Wilson K, Nagarkatti P, and Nagarkatti M. Aryl Hydrocarbon Receptor
Activation Ameliorates Acute Respiratory Distress Syndrome through Regulation of Th17 and
Th22 Cells in the Lungs. *mBio.* 2023:e0313722.
34. Mayer CT, Ghorbani P, Nandan A, Dudek M, Arnold-Schrauf C, Hesse C, et al. Selective and
efficient generation of functional Batf3-dependent CD103+ dendritic cells from mouse bone
marrow. *Blood.* 2014;124(20):3081-91.
35. Woroniecka KI, Rhodin KE, Dechant C, Cui X, Chongsathidkiet P, Wilkinson D, et al. 4-1BB
Agonism Averts TIL Exhaustion and Licenses PD-1 Blockade in Glioblastoma and Other
Intracranial Cancers. *Clin Cancer Res.* 2020;26(6):1349-58.
36. Habermann AC, Gutierrez AJ, Bui LT, Yahn SL, Winters NI, Calvi CL, et al. Single-cell RNA
sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in
pulmonary fibrosis. *Sci Adv.* 2020;6(28):eaba1972.
37. Adams TS, Schupp JC, Poli S, Ayaub EA, Neumark N, Ahangari F, et al. Single-cell RNA-seq reveals
ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. *Sci Adv.*
2020;6(28):eaba1983.
38. Gurczynski SJ, Procario MC, O'Dwyer DN, Wilke CA, and Moore BB. Loss of CCR2 signaling alters
leukocyte recruitment and exacerbates gamma-herpesvirus-induced pneumonitis and fibrosis
following bone marrow transplantation. *Am J Physiol Lung Cell Mol Physiol.* 2016;311(3):L611-
27.
39. Gurczynski SJ, Zhou X, Flaherty M, Wilke CA, and Moore BB. Bone marrow transplant-induced
alterations in Notch signaling promote pathologic Th17 responses to gamma-herpesvirus
infection. *Mucosal Immunol.* 2018;11(3):881-93.
40. Bowman WS, Newton CA, Linderholm AL, Neely ML, Pugashetti JV, Kaul B, et al. Proteomic
biomarkers of progressive fibrosing interstitial lung disease: a multicentre cohort analysis.
Lancet Respir Med. 2022;10(6):593-602.
41. Hildner K, Edelson BT, Purtha WE, Diamond M, Matsushita H, Kohyama M, et al. Batf3 deficiency
reveals a critical role for CD8alpha+ dendritic cells in cytotoxic T cell immunity. *Science.*
2008;322(5904):1097-100.
42. Badawy AA. Kynurenine Pathway of Tryptophan Metabolism: Regulatory and Functional
Aspects. *Int J Tryptophan Res.* 2017;10:1178646917691938.
43. Walisser JA, Glover E, Pande K, Liss AL, and Bradfield CA. Aryl hydrocarbon receptor-dependent
liver development and hepatotoxicity are mediated by different cell types. *Proc Natl Acad Sci U S*
A. 2005;102(49):17858-63.
44. Matthews J. AHR toxicity and signaling: Role of TIPARP and ADP-ribosylation. *Current Opinion in*
Toxicology. 2017;2:50-7.
45. Yamada T, Horimoto H, Kameyama T, Hayakawa S, Yamato H, Dazai M, et al. Constitutive aryl
hydrocarbon receptor signaling constrains type I interferon-mediated antiviral innate defense.
Nat Immunol. 2016;17(6):687-94.
46. Kozaki T, Komano J, Kanbayashi D, Takahama M, Misawa T, Satoh T, et al. Mitochondrial damage
elicits a TCDD-inducible poly(ADP-ribose) polymerase-mediated antiviral response. *Proc Natl*
Acad Sci U S A. 2017;114(10):2681-6.
47. Ahmed S, Bott D, Gomez A, Tamblyn L, Rasheed A, Cho T, et al. Loss of the Mono-ADP-
ribosyltransferase, Tiparp, Increases Sensitivity to Dioxin-induced Steatohepatitis and Lethality. *J*
Biol Chem. 2015;290(27):16824-40.
48. Chen PH, Chang H, Chang JT, and Lin P. Aryl hydrocarbon receptor in association with RelA
modulates IL-6 expression in non-smoking lung cancer. *Oncogene.* 2012;31(20):2555-65.
49. Vogel CF, Sciullo E, Li W, Wong P, Lazennec G, and Matsumura F. RelB, a new partner of aryl
hydrocarbon receptor-mediated transcription. *Mol Endocrinol.* 2007;21(12):2941-55.

50.   Vogel CF, Wu D, Goth SR, Baek J, Lollies A, Domhardt R, et al. Aryl hydrocarbon receptor
636   signaling regulates NF-kappaB RelB activation during dendritic-cell differentiation. *Immunol Cell*
637   *Biol.* 2013;91(9):568-75.
638   51.   Hogaboam CM, Trujillo G, and Martinez FJ. Aberrant innate immune sensing leads to the rapid
639   progression of idiopathic pulmonary fibrosis. *Fibrogenesis Tissue Repair.* 2012;5(Suppl 1):S3.
640   52.   Alzahrani B, Gaballa MMS, Tantawy AA, Moussa MA, Shoulah SA, and Elshafae SM. Blocking Toll-
641   like receptor 9 attenuates bleomycin-induced pulmonary injury. *J Pathol Transl Med.*
642   2022;56(2):81-91.
643   53.   Fang F, Marangoni RG, Zhou X, Yang Y, Ye B, Shangguang A, et al. Toll-like Receptor 9 Signaling Is
644   Augmented in Systemic Sclerosis and Elicits Transforming Growth Factor beta-Dependent
645   Fibroblast Activation. *Arthritis Rheumatol.* 2016;68(8):1989-2002.
646   54.   Chakraborty K, Chatterjee S, and Bhattacharyya A. Modulation of CD11c+ lung dendritic cells in
647   respect to TGF-beta in experimental pulmonary fibrosis. *Cell biology international.*
648   2017;41(9):991-1000.
649   55.   Serezani APM, Pascoalino BD, Bazzano JMR, Vowell KN, Tanjore H, Taylor CJ, et al. Multiplatform
650   Single-Cell Analysis Identifies Immune Cell Types Enhanced in Pulmonary Fibrosis. *Am J Respir*
651   *Cell Mol Biol.* 2022;67(1):50-60.
652   56.   Bantsimba-Malanda C, Marchal-Somme J, Goven D, Freynet O, Michel L, Crestani B, et al. A role
653   for dendritic cells in bleomycin-induced pulmonary fibrosis in mice? *Am J Respir Crit Care Med.*
654   2010;182(3):385-95.
655   57.   Helene M, Lake-Bullock V, Zhu J, Hao H, Cohen DA, and Kaplan AM. T cell independence of
656   bleomycin-induced pulmonary fibrosis. *J Leukoc Biol.* 1999;65(2):187-95.
657   58.   Wilke CA, Chadwick MM, Chan PR, Moore BB, and Zhou X. Stem cell transplantation impairs
658   dendritic cell trafficking and herpesvirus immunity. *JCI Insight.* 2019;4(18).
659   59.   Fois AG, Sotgiu E, Scano V, Negri S, Mellino S, Zinellu E, et al. Effects of Pirfenidone and
660   Nintedanib on Markers of Systemic Oxidative Stress and Inflammation in Patients with
661   Idiopathic Pulmonary Fibrosis: A Preliminary Report. *Antioxidants (Basel).* 2020;9(11).
662   60.   Takei H, Yasuoka H, Yoshimoto K, and Takeuchi T. Aryl hydrocarbon receptor signals attenuate
663   lung fibrosis in the bleomycin-induced mouse model for pulmonary fibrosis through increase of
664   regulatory T cells. *Arthritis Res Ther.* 2020;22(1):20.
665   61.   Epstein Shochet G, Bardenstein-Wald B, McElroy M, Kukuy A, Surber M, Edelstein E, et al.
666   Hypoxia Inducible Factor 1A Supports a Pro-Fibrotic Phenotype Loop in Idiopathic Pulmonary
667   Fibrosis. *Int J Mol Sci.* 2021;22(7).
668   62.   Wilson MS, Madala SK, Ramalingam TR, Gochuico BR, Rosas IO, Cheever AW, et al. Bleomycin
669   and IL-1beta-mediated pulmonary fibrosis is IL-17A dependent. *J Exp Med.* 2010;207(3):535-52.
670   63.   Cipolla E, Fisher AJ, Gu H, Mickler EA, Agarwal M, Wilke CA, et al. IL-17A deficiency mitigates
671   bleomycin-induced complement activation during lung fibrosis. *FASEB journal : official*
672   *publication of the Federation of American Societies for Experimental Biology.* 2017;31(12):5543-
673   56.
674   64.   Saito F, Tasaka S, Inoue K, Miyamoto K, Nakano Y, Ogawa Y, et al. Role of interleukin-6 in
675   bleomycin-induced lung inflammatory changes in mice. *Am J Respir Cell Mol Biol.*
676   2008;38(5):566-71.

679   **Figure Legends**

Figure 1. CD103+ DCs accumulate in the fibrotic lung and show evidence of pro-inflammatory AHR signaling. A and B) Mice (n = 6 – 10 mice per group (panel A), 5 per group (panel B) were treated with 0 75 U/kg bleomycin for 0, 7 or 14 days (panel A) or 14 days (panel B). Lungs were harvested and single cell suspensions were analyzed by flow cytometry for the presence of CD103+ DCs (CD45+, CD11c+, MHCII+, CD24+, CD64-, CD103+). **(C, D, and E)** groups of mice (n = 5-8 mice per group (panel C) or n = 6-8 per group (panel D and E)) were treated with 0.75 U/kg BLM for 14 – 21 days. Lungs were harvested and collagen was quantified by hydroxyproline assay or qRT-PCR. **F)** Mice (n = 4 per group) were treated with 0.75 U/kg BLM for 7 days after which lungs were harvested and single cell suspensions were analyzed via flow cytometry for AHR expression in the indicated cell subsets. **G through I)** Mice (n = 5 per group) were treated with 0.75 U/kg bleomycin or saline control (WT). Single cell suspensions were prepared from collagenase digested lung tissue at 7-days post bleomycin and CD11c+ cells were purified via magnetic bead isolation. Expression of the indicated transcripts was analyzed via qRT-PCR. All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA in panel A and C-E, or student's t-test in panels G-I (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, **** = $p < 0.0001$).

Figure 2. Production of AHR ligands is augmented during BLM induced fibrosis and complete AHR inhibition ameliorates disease. A) Mice (n = 4-5 mice per group) were treated with 0.75 U/kg BLM. Lungs were harvested at the indicated times and whole lung homogenates were assayed for KYN concentration by ELISA. **B)** Mice, (n = 3 – 5 per group) were treated with 0.75 U/kg bleomycin. Beginning at day 10 mice were treated daily with 5 mg/kg CH223191 dissolved in a vehicle of 2% Tween-20, 0.5% Carboxy methylcellulose, in PBS or vehicle alone by oral gavage until day 18. Lungs were harvested at Day 21 and lung collagen was quantified by hydroxyproline assay. **C)** Representative histology of lungs from B). Sections were stained with trichrome which highlights collagen fibers as blue staining (scale bar = 50 µM). All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA (* = $p < 0.05$, ** = $p < 0.01$, **** = $p < 0.0001$).

Figure 3. Deletion of AHR exon-2 reduces canonical AHR signaling in DCs but AHR expression is retained and augments inflammatory non-canonical AHR signaling. A) Cartoon schematic of murine AHR depicting the deletion of AHR exon-2 containing the DNA binding bHLH domain. Numbered boxes correlate to individual exons. **(B and C)** iCD103+ DCs were generated from WT B6 mice or AHR$_{\Delta ex2}$ mice and treated with 200 µM KYN or 10 µM CH223191 (CH223). 18 hours after treatment, RNA was harvested and expression of the canonical AHR gene Cyp1b1 (A) or AHR (B) was analyzed via qRT-PCR. **D)** iCD103+ DCs were analyzed via flow cytometry for expression of CD11c and CD103 as well as intracellular expression of AHR. **E)** Representative histogram showing AHR expression in WT or AHR$_{\Delta ex2}$ iCD103 DCs analyzed via flow cytometry. **F and G)** Mice (n = 5 per group) were treated with 0.75 U/kg bleomycin. At 10-d post bleomycin groups of mice were administered CH223191 (5 mg/kg) or vehicle alone daily until 18-d post bleomycin. Lungs were harvested at 21-d post bleomycin collagen content was quantified via hydroxyproline assay (F) and lung fibrosis was examined via histopathologic examination of fixed lung sections (A: Scale bar = 100 µM panel G). All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA in panel A or student's t-test in panel B (* = $p < 0.05$, ** = $p < 0.01$, **** = $p < 0.0001$).

Figure 4. Loss of canonical AHR signaling in CD11c+ cells exacerbates fibrosis via a ncAHR dependent mechanism. A) Mice (n = 7 mice no transfer group and 11 mice Cre+ transfer groups) were treated with 0.75 U/kg bleomycin. At 1-d post-BLM 1×10^6 iCD103 cells

generated from naïve CD11c-Cre+ AHR$_{\Delta ex2}$ mice were adoptively transferred via tail vein
injection. Additionally, one group was administered 100 µg αCD4 neutralizing antibody at 1-
and 7-d post bleomycin. Lungs were harvested 21-d post bleomycin and collagen content was
quantified via hydroxyproline assay. **B through D)** iCD103 cells were generated as described.
Cells were stimulated with 1 µM ODN2395 (TLR9 agonist), 10 µM RBN2397 (TiPARP inhibitor),
or a vehicle control for 18 hours. Cells were harvested and RNA was analyzed for expression of
IL-6 or TiPARP transcript via qRT-PCR. **E)** WT-B6 or AHR$_{\Delta ex2}$ mice were treated with 0.75 U/kg
BLM, CH223191 (CH223) was administered from day 10- to 18-post BLM. Lungs were
harvested at 21-day post BLM and expression of IL-6 transcript were quantified via qRT-PCR. **F**
and G) Mice (n = 5 - 7 per group) were treated with 0.75 U/kg bleomycin or saline control (WT).
Single cell suspensions were prepared from collagenase digested lung tissue at 7-days post
bleomycin and CD11c+ cells were purified via magnetic bead isolation after which cells were
pooled and treated overnight *ex-vivo* (n = 4 per group*)* with either 30µM CH223191 (CH223) or
DMSO vehicle control. Expression of the indicated transcripts was analyzed via qRT-PCR. All
experiments shown are representative of at least three independent experiments, statistical
significance was determined via ANOVA (* = p < 0.05, ** = p < 0.01, *** = p < 0.001, **** = p <
0.0001).
749
Figure 5. ncAHR signaling in CD11c+ cells drives IL-17 production in the lung. A) WT B6
or AHR$_{\Delta ex2}$ mice (n = 5 per group) were treated o.p. with 0.75 U/kg BLM. 7-days post-BLM lungs
were harvested and lung leukocytes were analyzed by flow cytometry for the following
populations Th17 (CD45+, Thy1.2+, CD3+, CD4+, IL-17A+), IL-17+ γδ t-cell (CD45+, Thy1.2+,
CD3+, CD4-, γδ-TCR+, IL-17A+), IL-17+ ILC3 (CD45+, Thy1.2+, CD3-, CD4-, IL-17+), IL-17A+
iNKT (CD45+, Thy1.2+, CD4-, CD3+, γδ-TCR+, IL-17A+). **B)** Quantification of cell populations
identified in A. **C)** WT-B6 or AHR$_{\Delta ex2}$ mice were treated with 0.75 U/kg BLM, CH223191
(CH223) was administered from day 10- to 18-post BLM. Lungs were harvested at 21-day post
BLM and expression of IL-17 transcript was quantified via qRT-PCR. **D)** Mice (n = 5-8 per
group) were treated with 0.75 U/kg BLM. 21-days post BLM treatment, lungs were harvested
and expression of IL-17 transcript from whole lung was analyzed via qRT-PCR. All experiments
shown are representative of at least three independent experiments, statistical significance was
determined via ANOVA (* = p < 0.05, ** = p < 0.01, **** = p < 0.0001).
763
Figure 6. Fibroblasts isolated from fibrotic individuals express the kynurenine pathway
enzymes IDO and TDO in response to TLR9 stimulation. A) Uniform manifold approximation
and projections (UMAPs) of human lung stromal cells labeled by normalized expression of
TDO2, cell type, and disease. **B through E**) Lung fibroblasts were cultured from groups of mice
treated with BLM for 21-d or untreated as control. Fibroblasts were subsequently treated with
the TLR9 agonist ODN2395 (ODN) or vehicle control (veh) for 18-h. Expression of the indicated
transcript was analyzed via qRT-PCR. All experiments shown are representative of at least two
independent experiments, statistical significance was determined via ANOVA (* = p < 0.05, ** =
p < 0.01, *** = p < 0.001).
773
Figure 7. Inhibition of TDO2 rescues BLM induced fibrosis. (A and B) Groups of mice (n = 5
per group) were treated with 0.75 U/kg BLM or saline control for 7-14 days after which
expression of the indicated transcript was analyzed in whole lung via qRT-PCR. **(C and D)**
Group of mice (n = 8 – 9 per group) were treated with 0.75 U/kg BLM or saline control.
Beginning on day 10 one group was treated with 15 mg/kg TDO2 inhibitor (680C91) daily by
oral gavage until day 18 after which lungs were harvested and expression of the indicated
transcript was assayed via qRT-PCR. **(E)** Representative histology (n =2 mice per group)
showing assessment of collagen deposition by trichrome staining (blue staining) scale bar
represents 100µm. All experiments shown are representative of at least two independent

experiments, statistical significance was determined via ANOVA (* = $p < 0.05$, ** = $p < 0.01$, ***
= $p < 0.001$).
785
Figure 8. Working model for how fibroblast DC crosstalk contributes to fibrosis.
Myofibroblasts express the kynurenine pathway enzyme TDO2 in response to TLR9 stimulation.
Kynurenine pathway metabolites active ncAHR signaling in DCs which, in cooperation with
inflammatory TLR singling, augments production of IL-6 resulting in production of IL-17 from
lymphocytes. Fibroblasts can be directly activated by IL-17 resulting in collagen production and
fibroproliferation.
792
793
794
795

Figure 1

Figure 1. CD103+ DCs accumulate in the fibrotic lung and show evidence of pro-inflammatory AHR signaling. A and B) Mice (n = 6 – 10 mice per group (panel A), 5 per group (panel B) were treated with 0.75 U/kg bleomycin for 0, 7 or 14 days (panel A) or 14 days (panel B). Lungs were harvested and single cell suspensions were analyzed by flow cytometry for the presence of CD103+ DCs (CD45+, CD11c+, MHCII+, CD24+, CD64-, CD103+). **(C, D, and E)** groups of mice (n = 5-8 mice per group (panel C) or n = 6-8 per group (panel D and E)) were treated with 0.75 U/kg BLM for 14 – 21 days. Lungs were harvested and collagen was quantified by hydroxyproline assay or qRT-PCR. **F)** Mice (n = 4 per group) were treated with 0.75 U/kg BLM for 7 days after which lungs were harvested and single cell suspensions were analyzed via flow cytometry for AHR expression in the indicated cell subsets. **G through I)** Mice (n = 5 per group) were treated with 0.75 U/kg bleomycin or saline control (WT). Single cell suspensions were prepared from collagenase digested lung tissue at 7-days post bleomycin and CD11c+ cells were purified via magnetic bead isolation. Expression of the indicated transcripts was analyzed via qRT-PCR. All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA in panel A and C-E, or student's T-test in panels G-I (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$, **** = $p < 0.0001$).

Figure 2

Figure 2. Production of AHR ligands is augmented during BLM induced fibrosis and complete AHR inhibition ameliorates disease. A) Mice (n = 4-5 mice per group) were treated with 0.75 U/kg BLM. Lungs were harvested at the indicated times and whole lung homogenates were assayed for KYN concentration by ELISA. **B)** Mice, (n = 3 – 5 per group) were treated with 0.75 U/kg bleomycin. Beginning at day 10 mice were treated daily with 5 mg/kg CH223191 dissolved in a vehicle of 2% Tween-20, 0.5% Carboxy methylcellulose, in PBS or vehicle alone by oral gavage until day 18. Lungs were harvested at Day 21 and lung collagen was quantified by hydroxyproline assay. **C)** Representative histology of lungs from B). Sections were stained with trichrome which highlights collagen fibers as blue staining (scale bar = 50 μM). All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA (* = p < 0.05, ** = p < 0.01, **** = p < 0.0001).

Figure 3

Figure 3. Deletion of AHR exon-2 reduces canonical AHR signaling in DCs but AHR expression is retained and augments inflammatory non-canonical AHR signaling. A) Cartoon schematic of murine AHR depicting the deletion of AHR exon-2 containing the DNA binding bHLH domain. Numbered boxes correlate to individual exons. **(B and C)** iCD103+ DCs were generated from WT B6 mice or AHR$_{\Delta ex2}$ mice and treated with 200 µM KYN or 10 µM CH223191 (CH223). 18 hours after treatment, RNA was harvested and expression of the canonical AHR gene Cyp1b1 (A) or AHR (B) was analyzed via qRT-PCR. **D)** iCD103+ DCs were analyzed via flow cytometry for expression of CD11c and CD103 as well as intracellular expression of AHR. **E)** Representative histogram showing AHR expression in WT or AHR$_{\Delta ex2}$ iCD103 DCs analyzed via flow cytometry. **F and G)** Mice (n = 5 per group) were treated with 0.75 U/kg bleomycin. At 10-d post bleomycin groups of mice were administered CH223191 (5 mg/kg) or vehicle alone daily until 18-d post bleomycin. Lungs were harvested at 21-d post bleomycin collagen content was quantified via hydroxyproline assay (F) and lung fibrosis was examined via histopathologic examination of fixed lung sections (A: Scale bar = 100 µM panel G). All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA in panel A or student's t-test in panel B (* = p < 0.05, ** = p < 0.01, **** = p < 0.0001).

Figure 4

Figure 4. Loss of canonical AHR signaling in CD11c+ cells exacerbates fibrosis via a ncAHR dependent mechanism. A) Mice (n = 7 mice no transfer group and 11 mice Cre+ transfer groups) were treated with 0.75 U/kg bleomycin. At 1-d post-BLM 1x10⁶ iCD103 cells generated from naïve CD11c-Cre+ AHR$_{\Delta ex2}$ mice were adoptively transferred via tail vein injection. Additionally, one group was administered 100 µg αCD4 neutralizing antibody at 1- and 7-d post bleomycin. Lungs were harvested 21-d post bleomycin and collagen content was quantified via hydroxyproline assay. **B through D)** iCD103 cells were generated as described. Cells were stimulated with 1 µM ODN2395 (TLR9 agonist), 10 µM RBN2397 (TiPARP inhibitor), or a vehicle control for 18 hours. Cells were harvested and RNA was analyzed for expression of IL-6 or TiPARP transcript via qRT-PCR. **E)** WT-B6 or AHR$_{\Delta ex2}$ mice were treated with 0.75 U/kg BLM, CH223191 (CH223) was administered from day 10- to 18-post BLM. Lungs were harvested at 21-day post BLM and expression of IL-6 transcript were quantified via qRT-PCR. **F and G)** Mice (n = 5 - 7 per group) were treated with 0.75 U/kg bleomycin or saline control (WT). Single cell suspensions were prepared from collagenase digested lung tissue at 7-days post bleomycin and CD11c+ cells were purified via magnetic bead isolation after which cells were pooled and treated overnight *ex-vivo* (n = 4 per group) with either 30µM CH223191 (CH223) or DMSO vehicle control. Expression of the indicated transcripts was analyzed via qRT-PCR. All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA (* = p < 0.05, ** = p < 0.01, *** = p < 0.001, **** = p < 0.0001).

Figure 5

Figure 5. ncAHR signaling in CD11c+ cells drives IL-17 production in the lung. A) WT B6 or AHR$_{\Delta ex2}$ mice (n = 5 per group) were treated o.p. with 0.75 U/kg BLM. 7-days post-BLM lungs were harvested and lung leukocytes were analyzed by flow cytometry for the following populations Th17 (CD45+, Thy1.2+, CD3+, CD4+, IL-17A+), IL-17+ γδ t-cell (CD45+, Thy1.2+, CD3+, CD4-, γδ-TCR+, IL-17A+), IL-17+ ILC3 (CD45+, Thy1.2+, CD3-, CD4-, IL-17+), IL-17A+ iNKT (CD45+, Thy1.2+, CD4-, CD3+, γδ-TCR+, IL-17A+). **B)** Quantification of cell populations identified in A. **C)** WT-B6 or AHR$_{\Delta ex2}$ mice were treated with 0.75 U/kg BLM, CH223191 (CH223) was administered from day 10- to 18-post BLM. Lungs were harvested at 21-day post BLM and expression of IL-17 transcript was quantified via qRT-PCR. **D)** Mice (n = 5-8 per group) were treated with 0.75 U/kg BLM. 21-days post BLM treatment, lungs were harvested and expression of IL-17 transcript from whole lung was analyzed via qRT-PCR. All experiments shown are representative of at least three independent experiments, statistical significance was determined via ANOVA (* = $p < 0.05$, ** = $p < 0.01$, **** = $p < 0.0001$).

Figure 6

Figure 6. Fibroblasts isolated from fibrotic individuals express the kynurenine pathway enzymes IDO and TDO in response to TLR9 stimulation. A) Uniform manifold approximation and projections (UMAPs) of human lung stromal cells labeled by normalized expression of TDO2, cell type, and disease. **B through E)** Lung fibroblasts were cultured from groups of mice treated with BLM for 21-d or untreated as control. Fibroblasts were subsequently treated with the TLR9 agonist ODN2395 (ODN) or vehicle control (veh) for 18-h. Expression of the indicated transcript was analyzed via qRT-PCR. All experiments shown are representative of at least two independent experiments, statistical significance was determined via ANOVA (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$).

Figure 7

Figure 7. Inhibition of TDO2 rescues BLM induced fibrosis. (A and B) Groups of mice (n = 5 per group) were treated with 0.75 U/kg BLM or saline control for 7-14 days after which expression of the indicated transcript was analyzed in whole lung via qRT-PCR. **(C and D)** Group of mice (n = 8 – 9 per group) were treated with 0.75 U/kg BLM or saline control. Beginning on day 10 one group was treated with 15 mg/kg TDO2 inhibitor (680C91) daily by oral gavage until day 18 after which lungs were harvested and expression of the indicated transcript was assayed via qRT-PCR. **(E)** Representative histology (n =2 mice per group) showing assessment of collagen deposition by trichrome staining (blue staining) scale bar represents 100µm. All experiments shown are representative of at least two independent experiments, statistical significance was determined via ANOVA (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$).

Figure 8

Figure 8. Working model for how fibroblast DC crosstalk contributes to fibrosis.
Myofibroblasts express the kynurenine pathway enzyme TDO2 in response to TLR9 stimulation. kynurenine pathway metabolites active ncAHR signaling in DCs which, in cooperation with inflammatory TLR singling, augments production of IL-6 resulting in production of IL-17 from lymphocytes. Fibroblasts can be directly activated by IL-17 resulting in collagen production and fibroproliferation.