Role of ADMA-histones in dual-strand piRNA source loci recognition by Rhino

Raku Saito[1,3]*, **Hirotsugu Ishizu**[2], **Ritsuko Harigai**[1], **Kensaku Murano**[2], **Yurika Namba**[1], **and Mikiko C. Siomi**[1]*

[1]Department of Biological Sciences, Faculty of Science, The University of Tokyo, Tokyo 113-0032, Japan

[2]Department of Molecular Biology, Keio University School of Medicine, Tokyo 162-8582, Japan

[3]Present address: Laboratory of Transcription Dynamics, Institute for Quantitative BioSciences, The University of Tokyo, Tokyo 113-0032, Japan

*Corresponding authors: R.S., rakusaito@g.ecc.u-tokyo.ac.jp and M.C.S., siomim@bs.s.u-tokyo.ac.jp

Keywords: piRNA cluster, Rhino, histone arginine methylation, ADMA, piRNA

Abstract

In *Drosophila* germ cells, piRNAs arise from dual-strand piRNA clusters, which are marked by a repressive histone mark H3K9me3, but are transcribed from internal sites in a manner dependent on the HP1 homolog Rhino. Rhino binds to H3K9me3 on these clusters, yet the mechanism controlling its binding to clusters remains unclear. Here, we used cultured ovarian somatic cells (OSCs), which lack endogenous Rhino and its stabilizer Kipferl, whose absence renders the dual-strand clusters inert, and found that exogenous Rhino tends to bind to the ends of dual-strand clusters with asymmetric dimethylarginine histones (ADMA-histones). Depletion of the arginine methyltransferases responsible for ADMA modifications, such as DART4, affected the genomic localization of Rhino in both OSCs and ovaries. We also identified primitive, cluster-like genomic regions, termed DART4 piSL, where Rhino propagates in an DART4-dependent but unstable manner. Our study proposes that ADMA-histones play a crucial role in the initial genome loading of Rhino and may establish the potential sites of its propagation, which are subsequently stabilized to support piRNA production.

Introduction

PIWI-interacting RNAs (piRNAs) are germline-enriched small non-coding RNAs that repress transposons in collaboration with PIWI proteins to maintain fertility[1,2,3]. In *Drosophila* germ cells, piRNAs are primarily derived from dual-strand piRNA clusters, intergenic elements rich in transposon fragments[4,5]. These clusters are occupied with the repressive histone mark H3K9me3 and lack typical promoters, but are transcribed in both directions from the internal sites[6,7]. This unconventional transcription requires special factors, including heterochromatin protein 1 (HP1) homolog HP1d/Rhino (Rhino), and provides piRNA precursors with a variety of sequences, which are then processed to mature piRNAs in the cytoplasm[5,6,7,8,9,10,11,12,13,14]. The mechanism dedicated to this piRNA processing is known as the ping-pong cycle, where Aubergine and Argonaute 3 cleave the precursors in a manner dependent on the sequences of piRNAs already bound to them[1,2,3]. This pathway also consumes transposon RNA transcripts as the sources of mature piRNAs, resulting in transposon repression at the post-transcriptional level[2,3].

Like other HP1 family members, Rhino has a chromodomain through which it binds to H3K9me3 scattered on dual-strand piRNA clusters[6,8,9,12,13,14,15,16]. Deadlock and Cutoff then join Rhino, assembling the Rhino–Deadlock–Cutoff (RDC) complex, which further recruits the transcription factor IIA-L (TFIIA-L) paralogue Moonshiner, the transcription factor IIA-S (TFIIA-S), and the animal TATA-box-binding protein (TBP) paralog TRF2, initiating transcription of the clusters[6,7,10]. Upon transcription, Cutoff binds to piRNA precursors and protects them from splicing, contributing to the production of an abundance of piRNAs[17,18,19,20].

A recent study identified the zinc-finger protein Kipferl as a factor required for

Rhino to distinguish between H3K9me3 located on transposons and dual-strand piRNA

clusters[12]. Kipferl stabilizes Rhino on guanine-rich sequence motifs present in dual-

strand clusters in an H3K9me3-independent manner[12]. Maternally inherited piRNAs also

play a role in the localization of Rhino to dual-strand clusters during early embryogenesis

via Piwi-dependent H3K9me3 epigenetic marks [9,21].

Rhino has a propensity to spread across the genome. For example, depletion

of the histone demethylase Kdm3 in the female germline causes Rhino to spread to

adjacent chromatin[13]. Deletion of the region proximal to the transcription start site of the

phospholipase D gene, which is in the region flanking piRNA cluster 42AB, also resulted

in Rhino spreading toward the deleted genomic region[10]. In addition, Rhino tethering

experiments with artificial constructs showed that Rhino spreads as the constructs are

transcribed[7]. In each case, production of piRNAs from regions after Rhino spreading was

detected. These findings suggest that piRNA clusters are defined by Rhino spreading

over the genome and imply that the process of piRNA cluster formation is composed of

three steps: Rhino spreading, Kipferl-dependent stabilization of Rhino–genome

association, and epigenetic inheritance of the Rhino positional memory by maternally

inherited piRNAs. Since the genomic localization of both Rhino and Kipferl is

interdependent, it is hypothesized that the initial loading of Rhino on the genome

precedes its spread or stabilization, and that this process is independent of Kipferl. To

investigate this, it is necessary to observe the behavior of Rhino in a cell environment

where piRNAs are functional, but where both Kipferl and Rhino are not expressed.

However, the mechanism of such Rhino behavior remains unclear.

In this study, we describe the important role of asymmetric dimethylarginine

histones (ADMA-histones) in the initial loading of Rhino onto the genome and their

stabilization. We first established cultured ovarian somatic cells (OSCs)[22] expressing

FLAG-Rhino by induction with doxycycline (Dox). Unlike ovarian germ cells, OSCs lack

Rhino and other factors required for the activation of dual-strand piRNA clusters, such

as Deadlock and Cutoff[23]; therefore, dual-strand clusters are not in an active state for

piRNA biogenesis[24]. We found that Rhino in OSCs assembles nuclear foci similar to

Rhino foci in the ovary. We then performed chromatin immunoprecipitation sequencing

(ChIP-seq) and showed that Rhino tends to bind to the ends of dual-strand clusters with

ADMA-histones, such as H3R17me2a. Depletion of the arginine methyltransferases

responsible for ADMA modifications, such as *Drosophila* arginine methyltransferase 4

(DART4), affected the genomic localization of Rhino in OSCs and in the ovary. We also

identified genomic regions that have ADMA-histones at their ends and exhibit broad

Rhino spreading across their internal regions in a DART4-dependent manner. In contrast

to authentic piRNA clusters, Kipferl was lost together with Rhino upon DART4 depletion

in these regions, suggesting that Kipferl by itself is not sufficient to stabilize Rhino

binding; rather, their localization depends on DART4. piRNAs were mapped to these

regions, albeit at low levels. The levels of these piRNAs decreased also in a DART4-

dependent manner. Therefore, these regions, referred to as DART4-dependent piRNA

source loci (DART4 piSL), might be nascent dual-strand piRNA source loci driven by the

propagation of Rhino initially loaded onto the genome via ADMA-histones.

Results

Rhino localizes to the ends of dual-strand clusters in the OSC genome

To observe how Rhino in OSCs behaves intracellularly, we first performed

immunofluorescence in FLAG-Rhino inducible OSCs to examine the subcellular

localization of FLAG-Rhino upon Dox induction. This showed that Rhino assembled

nuclear speckles in OSCs similar to Rhino-positive foci in the ovary[5,10] when the cells

were treated with Dox at a concentration of 5 ng/mL (Fig. 1A). At lower Dox

concentrations and lower levels of Rhino expression, no clear foci appeared

(Supplementary Fig. 1A, B), suggesting that Rhino-positive foci form in a manner

dependent on the level of Rhino expression.

We next conducted ChIP-seq of FLAG-Rhino under conditions where Rhino

assembles nuclear foci as in Fig. 1A. Genome mapping of the ChIP-seq reads showed

that Rhino can be found, albeit weak, on dual-strand piRNA clusters, such as 42AB and

38C on chromosome 2, 80F on chromosome 3, and 102F on chromosome 4, all of which

are in an active state in piRNA production in the ovary, but the signals seem to be biased

toward the ends of the clusters (Fig. 1B and Supplementary Fig. 1C). ChIP-seq of Rhino

was also performed in the ovary: Rhino was strongly enriched at highly transcribed

piRNA clusters, 42AB, 38C, 80F, and 102F, as previously reported[5,6,7,8,9,10,11,12,13,14] (Fig.

1B and Supplementary Fig. 1C), which suggests that our ChIP-seq procedure is reliable.

We then compared the mode of Rhino genome binding in OSCs with that in the

ovary in a more comprehensive manner. The results show that Rhino expressed in OSCs

bound predominantly to genomic sites exhibiting sharp and interspersed Rhino localization

patterns in the ovary, while showing little localization within broad Rhino domains, including

major piRNA cluster (Fig. 1C). The metaplot indicates a tendency for Rhino in OSCs to

be toward the terminal regions of broad domains, such as the cluster ends (Fig. 1D).

Rhino also binds to the cluster bodies in OSCs, but the signals were weaker compared

to those at the ends. Based on these results, we interpret that OSCs contain a factor(s)

that promotes Rhino binding toward the ends of the clusters, but unlike the ovary, they

lack a factor(s) that promotes and stabilizes Rhino spreading throughout the clusters.

The terminal localization of Rhino in OSCs does not appear to be due to

H3K9me2 (Fig. 1B, E, F): Of the 823 Rhino signals, only 14 signals overlapped with the

H3K9me2 signals (Fig. 1E). The heatmap also showed that H3K9me2 barely

accumulated at the Rhino location (Fig. 1F). The positional correlation between Rhino

and H3K9me3 is stronger than between Rhino and H3K9me2 (i.e., 77%; 636 in 823

signals) (Fig. 1E). Nonetheless, the overlap between Rhino and H3K9me3 was not very

strong according to the heatmap (Fig. 1F, G). Immunofluorescence also showed weak

overlap between the prominent H3K9me3 speckles and FLAG-Rhino foci (Fig. 1A).

Since immunofluorescence primarily visualizes H3K9me3 foci that correspond to broad

heterochromatic domains in the genome, such as those at centromeres, pericentromeres,

or telomeres (named chromocenters), the sharp and interspersed H3K9me3 signals

along chromosome arms, where Rhino partially colocalizes (Fig. 1F), are difficult to

visualize, resulting in little apparent overlap with Rhino (Fig. 1A). HP1a, another HP1

family member required for constitutive heterochromatin and known to be a general

H3K9me3 marker, co-localizes with H3K9me2/3 and is distributed along the entire length

of piRNA clusters similarly in both OSCs and ovary (Fig. 1B and Supplementary Fig. 1C).

HP1a is present in the broad H3K9me3 peak (Fig. 1G). Therefore, although H3K9me3

and HP1a might be involved in some aspect of the mechanism underlying Rhino

localization, they do not contribute to the localization specificity within that mechanism

at least.

FLAG-Rhino ChIP-seq was also performed at low Rhino expression levels. The

trend toward terminal localization of Rhino at 5 ng/mL Dox was observed at 0.05 ng/mL

and 0.5 ng/mL Dox as well (Supplementary Fig. 1D, E). This indicates that Rhino binds

to the OSC genome even at low expression levels. However, Rhino foci were not clearly

observed with the lower Dox concentrations (Supplementary Fig. 1A). It is interpreted

that genome binding occurs first, followed by Rhino foci formation, which is dependent

on Rhino expression levels.

The use of anti-FLAG antibodies in Rhino ChIP in OSCs was justified by the

high similarity between the signals of anti-FLAG and anti-Rhino antibodies (Fig. 1B and

Supplementary Fig. 1C, D).

Rhino co-localizes with asymmetric dimethylarginine histones on the OSC genome

To identify genomic scaffolds that are potentially involved in determining the

genomic localization of Rhino in OSCs, we performed ChIP-seq of key histone

modifications and compared the data with Rhino ChIP-seq data. Heatmap revealed that

specific ADMA-histones, including H3R2me2a, H3R8me2a, H3R17me2a, and

H4R3me2a, tend to co-localize with Rhino across the OSC genome (Fig. 2A). Consistent

with this, these ADMA-histones showed a tendency to localize to the ends of Rhino-

bound dual-strand piRNA clusters (Fig. 2B, C, and Supplementary Fig. 2A). H3K4me1

and H3K27ac have been reported to co-localize with H3R8me2a in mammals[25], which

was not observed in OSCs (Fig. 2A, B, and Supplementary Fig. 2A). Co-localization of

H3K4me1 and H3K27ac with Rhino was also not observed.

Examination of the correlation coefficients revealed that signal peaks of each

ADMA-histone were significantly correlated with those of FLAG-Rhino (Fig. 2D). The

ChIP-seq signals of H3R2me2s and H3R8me2s correlate more strongly with

FLAG(Dox-) than with FLAG(Dox+), suggesting that these histone modifications may

represent background noise (see "background noise" in Fig. 2D). In contrast, ADMA-histones show a higher correlation coefficient with FLAG(Dox+) than with FLAG(Dox-): it is also worth noting that in hierarchical clustering, ADMA-histones form a monophyletic group with FLAG(Dox+) (see "Rhino localization" in Fig. 2D). Histone modifications related to transcription, such as H3K4 and H3K27, as well as constitutive heterochromatin markers like H3K9me2 are not associated with the localization of FLAG-Rhino (see "transcriptional regulation" in Fig. 2D). Correlation of ADMA-histones in the genomic localization of Rhino was implicated.

The overlaps of ChIP-seq peaks were examined. In the diagram shown in Fig. 1E, approximately 23% of Rhino ChIP peaks did not overlap with H3K9me3 (187 in 823 sites), but these Rhino peaks were almost encapsulated by the read sets of ADMA-histones (Fig. 2E). This suggests that ADMA-histones are involved in the localization of Rhino. However, the degree of overlap of those sites is not very high in each case (Supplementary Fig. 2B). It appears that each ADMA-histone contributes differently to Rhino localization. HP1a was found at the dual-strand clusters in OSCs (Fig. 1B). However, unlike Rhino, genomic localization of HP1a does not appear to be determined by ADMA-histones and H3K9me3 because approximately 30% of HP1a ChIP peaks did not overlap with the read sets of ADMA-histones and H3K9me3 (492 in 1,672 sites) (Supplementary Fig. 2C). These results raise the possibility that ADMA-histones, together with H3K9me3, may contribute specifically to the recruitment of Rhino to the ends of dual-strand clusters in OSCs.

Arginine methyltransferases DART1 and DART4 differently regulate Rhino genomic binding

Arginine methylation of histones is catalyzed by protein arginine

methyltransferases (PRMTs), and the *Drosophila* genome encodes 11 of these PRMTs,

also known as *Drosophila* arginine methyltransferases (DARTs)[26,27]. Of those, DART1, a

homolog of mammalian PRMT1, and DART4/CARMER (DART4), a homolog of

mammalian PRMT4/CARM1 (CARM1), were reported to be type I PRMTs that catalyze

the methylation of monomethylarginine to asymmetric dimethylarginine[28,29,30,31,32,33,34,35].

DART8 is involved in asymmetric dimethylation of H3R2[36]. CG17726 is considered to be

a DART member because of its sequence similarity and is expressed in the ovary

(Flybase 2024). However, its function remains unknown. If ADMA-histones play an

important role in the local recruitment of Rhino to the OSC genome, it is highly likely that

the loss of DARTs responsible for ADMA modification would influence the subcellular

localization of Rhino.

A previous study showed that HP1a germline knockdown (GLKD) causes Rhino

foci to be more prominent in the ovary and reduces Rhino occupancy at the clusters[37].

Because a similar phenotype was observed for Rhino foci in HP1a-knockdown (KD)

OSCs (Fig. 3A, B), we used this phenotype to examine the loss-of-function effects of

DARTs. Similar phenotypes were observed in DART1- and DART4-KD OSCs as in

HP1a-KD OSCs, suggesting that DART1/4-dependent histone ADMA modification is

involved in Rhino recruitment (Fig. 3A, B). Moreover, the localization of Rhino in

DART1/4 KD cells closely matches prominent H3K9me3 foci, suggesting that the

arginine methylation activity of DART1/4 contributes to the genomic localization of Rhino

in OSCs as a genome scaffold beyond H3K9me3 (Supplementary Fig. 3A). In contrast,

this phenotype was not observed for HP1a, suggesting that the effects of DART1/4 are

specific to Rhino (Supplementary Fig. 3B).

Fluctuation in the localization of Rhino in the DART1- and DART4-KD OSCs

was also observed in ChIP experiments. In ChIP-seq, Rhino tended to be decreased in

DART1 KD and DART4 KD (Fig. 3C, D), which was confirmed by ChIP-qPCR (Fig. 3E).

Among the ADMA-histones that co-localized with Rhino, H3R17me2a decreased in a

DART4-dependent manner and H4R3me2a decreased in a DART1-dependent manner

(Fig. 3C), consistent with previous findings for human PRMT1 and CARM1[38,39,40,41].

Detailed analysis revealed that the genomic localizations of H4R3me2a lost in DART1

KD (575 sites) and H3R17me2a lost in DART4 KD (653 sites) overlapped at 48 sites

(Fig. 3F). Correspondingly, the genomic localizations of Rhino that decreased in DART1

KD (115 sites) and those that decreased in DART4 KD (207 sites) overlapped at 25 sites

(Fig. 3G). These results suggest that DART1 and DART4 could contribute to Rhino

recruitment at distinct genomic sites through the decreases in ADMA-histones in each of

their KD conditions (H4R3me2a and H3R17me2a, respectively).

Among ADMA-histones, only H3R8me2a tended to decrease in both DART1

and DART4 KD (Fig. 3C). To examine the specificity of the regulation of H3R8me2a by

DART1 and DART4, the degree of overlap was visualized in Venn diagrams for

H3R8me2a, as well as H3R17me2a and H4R3me2a. The genomic localizations of

H3R8me2a decreased by DART1 KD (235 sites) and DART4 KD (58 sites) weakly

overlapped (8 sites) (Supplementary Fig. 3C). The genomic localizations of H3R8me2a

(58 sites) and H3R17me2a (652 sites) that decreased in DART4 KD overlapped only at

7 sites (Supplementary Fig. 3D), as was the case for H3R8me2a (235 sites) and

H4R3me2a (575 sites), which were decreased in DART1 KD (Supplementary Fig. 3E).

These results suggest that DART1 and DART4 regulate Rhino localization through

H3R8me2a at different sites, as well as through H3R17me2a and H4R3me2a.

The level of Rhino expression was unchanged in DART1/4-KD cells

(Supplementary Fig. 3F). Rhino and HP1a even in normal OSCs were under detection

levels with anti-ADMA antibodies (Supplementary Fig. 3G). Thus, the loss of DART1/4

does not appear to affect Rhino or HP1a function directly through methylation. Although

DART4 has been reported to be highly expressed in the ovary[28], public RNA-seq data

showed that its reads were almost undetectable in both OSCs and ovary (Supplementary

Fig. 3H). Consistent with this, qPCR analysis revealed that DART4 mRNA expression

was only about 3% of that of DART1 (Supplementary Fig. 3I). Nevertheless, this low-

level mRNA expression appears sufficient to support its functional role, such as the

modification of H3R17me2a (Fig. 3C and Supplementary Fig. 3F). The knockdown of

each gene in OSCs was confirmed to be effective (Supplementary Fig. 3J).

ADMA-histones co-localize with Rhino in a manner dependent of DART4 in the

ovary

We examined the localization of ADMA-histones in the Oregon-R ovary. In the

wild-type (WT) ovary, we detected Rhino peaks in FLAG-Rhino localization sites in OSCs

and ADMA-histones, H3R8me2a, H3R17me2a, and H4R3me2a, co-localized with them

(Fig. 4A and Supplementary Fig. 4). Heatmap showed that these ADMA-histones are

detected in the narrow Rhino-positive regions as in the case of Rhino in OSCs (Fig. 4B).

As well, the metaplot showed that ADMA-histones tend to localize to the ends of piRNA

clusters (Fig. 4C). The trend of Rhino genome binding observed in OSCs was also

recapitulated in ovaries at corresponding genomic locations.

To determine whether the destabilization of Rhino genome binding observed in

OSCs under DART4-KD conditions is also observed in the ovary, we performed GLKD

of DART4 using the GAL4-UAS system. Under the DART4-GLKD conditions (Supplementary Fig. 5A), Rhino foci were more prominent in the absence of DART4 (Fig. 5A, B). ChIP-seq experiments showed that Rhino localizes almost the same genomic regions in the ovary as in OSCs, although these localizations are weaker compared to piRNA clusters such as 42AB (Fig. 5C, D). Notably, Rhino levels significantly decreased with the loss of H3R17me2a under DART4-GLKD conditions (Fig. 5C–F, and Supplementary Fig. 5C, D). These results indicate that the relationship between ADMA-histones and Rhino found in OSCs is conserved in the ovary.

Interestingly, Rhino spreading across clusters annotated in the piRNA cluster database[42], such as 42AB, 38C, 80F, and 102F, was little affected by DART4 GLKD (Supplementary Fig. 5D). These results suggest that the weak Rhino localization dependent on ADMA-histones (such as at the terminal sites of piRNA clusters) and the strong Rhino localization within the core regions of piRNA clusters are fundamentally governed by distinct regulatory mechanisms.

Loss of Kdm3 in the ovary (i.e., Kdm3-GLKD ovary) causes an increase in H3K9me2 and regional expansion, accompanied by Rhino spreading and the appearance of ectopic piRNA clusters[13]. To determine whether ADMA-histones are involved in this process, we examined the localization of ADMA-histones in the ectopic piRNA clusters. ADMA-histones were found near the 5′ and 3′ ends of the Kdm3 GLKD-dependent piRNA clusters, as well as in the authentic piRNA clusters (Supplementary Fig. 5E). Relatively few ADMA-histones were observed inside the clusters, even in the Kdm3 GLKD-dependent piRNA clusters (Supplementary Fig. 5F). This raises the possibility that ADMA-histones may be involved in the Rhino propagation by defining the spreading regions.

Genomic regions where Rhino binds broadly in a DART4-dependent manner, but

not stably anchored, produce some piRNAs

Counting the genomic sites where H3R17me2a was no longer localized in

DART4-GLKD ovaries, 3,082 sites were identified. The genomic sites where Rhino was

decreased under the same circumstance were also counted (2,078 sites). Of these, 497

sites overlapped in the two groups (Fig. 6A). This indicates that 76.1% of the Rhino

genomic localization reduced by DART4 GLKD appears to be H3R17me2a-independent.

Forty-three such regions were found in the *Drosophila* genome (Fig. 6B, C).

Representative examples are shown in Fig. 6D and Supplementary Fig. 6A: In control

KD, these regions have Rhino in their internal regions and H3R17me2a at their ends

(and other ADMA-histones at each end in WT Oregon-R strain, albeit at low levels). In

DART4 GLKD, H3R17me2a disappeared from the terminal regions as expected. The

internal Rhino also disappeared, although the areas did not contain H3R17me2a (Fig.

6D, E). These regions are distinctly different in nature from authentic piRNA clusters,

where Rhino localization remained largely unaffected by DART4 GLKD (Fig. 6B).

Not only H3R17me2a but also other ADMA-histones were found at the ends of

these regions, and H3K9me2 accumulated in the internal areas (Supplementary Fig. 6B),

consistent with previous studies suggesting H3K9me2 involvement in the Rhino

spreading[13]. FLAG-Rhino in OSCs was also localized at the ends of the regions (Fig.

6E). These results raise the possibility that a subset of Rhino localized to genomic

regions correlating with ADMA-histones may serve as origins of spreading.

We next investigated the potential for piRNA production from the genomic

regions where Rhino propagates in a DART4-dependent manner but does not stably

anchor. Under control conditions, piRNAs were detected mapping to these regions, albeit

at low levels (Fig. 6D, F, and Supplementary Fig. 6A). Under DART4-GLKD conditions,

piRNA levels were significantly reduced, suggesting that piRNA production from these

regions is dependent on DART4. Although piRNA production was also influenced by

Rhino, the reduction observed upon Rhino loss was much less pronounced compared

to annotated piRNA clusters (Fig. 6G).

The expression levels of transposons and genes presumed to be targeted by

piRNAs originating from genomic regions showed minimal changes under DART4-GLKD

conditions (Supplementary Fig. 6C). Similarly, egg hatching rates remained largely

unaffected (Supplementary Fig. 6D). These observations could be attributed to the

relatively low levels of piRNAs arising from these regions compared to authentic piRNA

clusters (Supplementary Fig. 6E, F), as well as their shorter genomic lengths

(Supplementary Fig. 6G). This can also be explained by the fact that piRNA production

from authentic clusters remained largely unaffected. Notably, even the functional loss of

authentic piRNA clusters in germ cell piRNA production does not usually result in

infertility[43]. Based on these findings, which highlight that these genomic regions differ

from authentic piRNA clusters in many aspects yet possess the ability to produce piRNAs,

we designate them as DART4-dependent piRNA source loci (DART4 piSL).

Kipferl stabilizes the Rhino–genome binding[12]. ChIP-seq using anti-Kipferl

antibodies showed that Kipferl localizes to DART4 piSL and decreases in DART4 GLKD

like Rhino (Supplementary Fig. 7A, B). These results suggest that, although Kipferl is

present at DART4 piSL, the apparent stabilization of Rhino may depend more on DART4

than on Kipferl, unlike at other authentic piRNA clusters.

In Kipferl mutants, Rhino localization is biased toward satellite repeats[12]. In

DART4 GLKD, there appears to be no significant difference in Rhino localization (Supplementary Fig. 7C), nor does Rhino become enriched at satellite repeats (Supplementary Fig. 7D).

A recent comprehensive annotation of active piRNA clusters across eight highly inbred *Drosophila* strains classified the clusters into eight levels of evolutionary novelty[44]. We subsequently analyzed the enrichment of ADMA-histones, which significantly correlate with Rhino localization in OSCs (Fig. 2A–E) and ovaries (Fig. 4 A–C) (e.g., H3R2me2a, H3R8me2a, H3R17me2a, and H4R3me2a), around these piRNA clusters. Notably, a trend of greater ADMA-histone enrichment was observed at the ends of the newer clusters (Fig. 7A). This trend was also observed in DART4 piSL (Supplementary Fig. 8A). In contrast, such bias was not observed in older clusters or other genomic loci (Fig. 7A, B, Supplementary Fig. 8A). H3R2me2a, H3R8me2a, H3R17me2a, and H4R3me2a showed a loss of enrichment as the clusters aged (Fig. 7A). The DART4-dependent H3R17me2a differences observed between newer and older clusters were less pronounced in DART4 GLKD (Fig. 7B). We next extracted piRNA clusters where piRNAs were detected in the Oregon-R strain and examined them for ADMA-histone enrichment. The results remained consistent (Supplementary Fig. 8B). A similar analysis using GLKD strains yielded identical findings (Supplementary Fig. 8C). These results suggest that DART4 piSL represent relatively new piRNA clusters.

Based on all these results, we propose that ADMA-histones play a crucial role in determining Rhino's initial genomic binding (Fig. 7C, left). Furthermore, ADMA-histone-dependent Rhino binding to the genome may also contribute to the propagation of Rhino within the piRNA cluster regions (Fig. 7C, right). Our evolutionary analysis suggests that DART4 piSL appear to share features with young piRNA clusters and may

eventually develop into *bona fide* piRNA clusters like 42AB; however, it does not provide direct evidence for this process. Further studies will be necessary to explore this possibility.

Discussion

In this study, we took advantage of cultured OSCs for our analysis and found that chromatin marks (i.e., bivalent nucleosomes containing H3K9me3 and ADMA-histones) appear to contribute to the initial loading of Rhino onto the genome. Comparative analysis of OSC and ovarian ChIP-seq data also allowed us to identify "primitive" piRNA source loci, DART4 piSL. Based on these results, DART4 piSL are piRNA source loci where Rhino propagates in a DART4-dependent manner from the ADMA marks at the ends, but this spread is not stable. In DART4 piSL, Rhino spreads from the initial loading sites but does not stabilize, indicating that these regions may be nascent piRNA clusters. The evolutionary analysis showing a stronger tendency for ADMA accumulation at the ends of younger piRNA clusters (Fig. 7A) provides indirect evidence that DART4 piSL may represent evolutionarily young, nascent piRNA clusters. While DART4 piSL may represent a "later" stage compared to annotated piRNA clusters, the essential role of piRNAs in reproduction and species conservation makes it unlikely that a large number of clusters exist in a degenerate state. Thus, we speculate that these piRNA source loci represent an "earlier" stage in piRNA cluster formation. As note, both ctrl GLKD and DART4 GLKD share the nos-GAL4 background, but we cannot fully exclude the possibility that the identified DART4 piSL reflect strain-specific genomic variation between ctrl UAS-RNAi strain and DART4 UAS-RNAi strain rather than DART4-dependent regulation.

Whether DART4 piSL evolve into large piRNA clusters such as 42AB and 38C

during the course of evolution remains an open question. Although these large clusters

were not affected by DART4 GLKD, ADMA-histones were still detectable, albeit at low

levels, at the ends of their Rhino peaks (Fig. 2B and 4A). This observation raises the

possibility that even large piRNA clusters like 42AB and 38C may have initially formed

through a process involving ADMA-histone–mediated Rhino loading followed by

propagation, as seen in DART4 piSL. While it remains possible that these clusters

formed through a different mechanism without passing through a DART4 piSL -like phase,

such a scenario would leave the presence of residual ADMA-histones at their ends

unexplained. Nonetheless, evolutionary analyses provide only indirect evidence, and

experimental validation will be necessary to determine whether DART4 piSL can indeed

mature into larger piRNA clusters.

In general, ADMA-histones are known to function as transcriptional

coactivators[45]. In mammalian cells, H3R8me2a does not co-localize with H3K9me3, but

with enhancer modifications such as H3K4me1 and H3K27ac[25]. This may indicate that

the function of ADMA-histones has evolved in a *Drosophila*-specific manner, which is not

unlikely given that Rhino itself is *Drosophila*-specific. However, there are regions where

Rhino does not localize in the read sets of ADMA-histones and H3K9me3, suggesting

that an unknown factor may contribute to the specificity of Rhino targeting (Fig. 2E).

This study showed that multiple ADMA-histones, such as H3R17me2a and

H4R3me2a, play a role in the initial Rhino loading. This suggests that a co-factor(s) with

an affinity for ADMAs might be involved. Tudor domains are known to bind partner

proteins through methylation of arginine and lysine residues[46], and Tudor domain-

containing protein 3 (TDRD3) was identified as a factor that specifically binds to

H3R17me2a and H4R3me2a in humans[47]. Therefore, TDRD protein family members

may be involved in Rhino loading via ADMAs. It should also be noted that not all ADMA-

enriched regions recruit Rhino; rather, Rhino association is observed only where ADMA-

histones overlap with H3K9me3 (Fig. 2E). Even the combination of H3K9me3 and

ADMA-histones does not fully explain the specificity of Rhino recruitment, implying the

requirement for additional co-factors. Such factors may include other ADMA marks (e.g.,

H3R42me2a) or chromatin-interacting proteins that are arginine-methylated by DART1

or DART4. DART1 and DART4 might contribute to Rhino recruitment not only through

direct histone methylation but also via arginine methylation of non-histone chromatin

factors. However, at least under our conditions, HP1a methylation did not appear to be

involved in Rhino behavior (Fig. S3G). Given the potential functional redundancy among

multiple DARTs, future studies may require simultaneous knockdown or knockout of

several DART proteins to fully assess their contribution to Rhino recruitment and piRNA

cluster regulation.

The mechanism by which ADMA-histones is involved in Rhino propagation

remains unclear. Because Rhino propagation has been shown to be correlated with

transcription[7,10], one interpretation is that ADMA-histones may affect Rhino spreading via

transcriptional activity. Insertion of insulators into piRNA clusters inhibits piRNA

production[48], indicating that the three-dimensional (3D) structure of the genome within

the piRNA cluster also affects the extent of Rhino spreading. ADMA-histones were found

at the ends of piRNA clusters, reminiscent of the boundaries of topologically associated

domains. An unknown stabilizing factor(s) suggested by the absence of Kipferl in some

DART4 piSL (Supplementary Fig. 7A, B) may be a relevant factor(s) determining such

genomic 3D structures. Functional validation of ADMA-histones in piRNA cluster

specification will be required in future studies.

Maternally inherited siRNAs initiate piRNA biogenesis from a transgene inserted

into the uni-strand piRNA cluster 20A[14]. In this previous report, evidence was presented

that siRNAs induce "*de novo*" Rhino loading onto this transgene (20A-X) via H3K9me3.

This led us to hypothesize that ADMA-histones should be present in the vicinity of 20A-

X. Indeed, we detected ADMA-histones in the region and Rhino, albeit weakly

(Supplementary Fig. 4). Thus, this study provided experimental evidence in support of

the "*de novo* model" by reinforcing the findings of siRNA induced piRNA biogenesis. This

model has been becoming supported by evolutionary analyses[44,48,49], but experimental

results had been limited. The evolutionarily young piRNA clusters, which appear in a

limited number of strains, may have formed through the spread of Rhino into newly

inserted transposons, facilitated by the presence of ADMA-histones in their vicinity (Fig.

7A).

In *Drosophila*, major piRNA clusters such as 42AB and 38C are located at the

boundary between heterochromatin and euchromatin[4,51]. It was unclear why these highly

transcribed clusters, which are capable of producing substantial piRNAs, tend to appear

in regions where two different types of chromatins collide. If DART4 piSL represent the

initial stage in the formation of piRNA clusters, our model may help explain this question

by showing that cluster formation begins with initial Rhino loading by ADMA-histones

onto euchromatic regions and then spreads to adjacent regions of H3K9me2-rich

heterochromatin (Fig. 7C). Rhino shows a tendency to spread from an DART4 piSL on

chromosome 2 to heterochromatin/euchromatin boundary regions where 42AB and 38C

are present (Fig. 6C), supporting our hypothesis.

Methods

Establishment and cloning of FLAG-Rhino inducible OSCs

To construct Tet-ON 3G inducible FLAG-Rhino plasmid, cDNA encoding FLAG-Rhino was cloned into pPB-2× Ty1-Tjen-EGFP-P2A-BlastR[52] using In-Fusion cloning (TAKARA), bearing pPB-FLAG-Rhino-Tjen-EGFP-P2A-BlastR. The Actin promoter and *EGFP* cDNA were replaced respectively with the TRE3G promoter and *rtTA* cDNA from inducible Caspex expression plasmid (Addgene). PCR was performed using PrimeSTAR MAX (TAKARA) with primers (Supplementary Table 1). FLAG-Rhino inducible OSCs were established using the piggyBac system as described previously[52]. In brief, 2×10^6 OSCs were transfected with 4 μg of pPB-TRE3G-FLAG-Rhino-Tjen-trTA-P2A-BlastR and 1 μg of pHsp70-Myc-hyPBase using Nucleofector Kit V (Lonza). After incubation in media containing blasticidin (50 μg/mL), colonies were isolated. One of the colonies contained a single FLAG-Rhino copy in the intron of the *jing interacting gene regulatory 1* gene, and this was used in this study.

Cell culture and RNAi

FLAG-Rhino inducible OSCs were cultured in Shields and Sang M3 Insect Medium (Sigma) supplemented with 10% fly extract, 10% fetal bovine serum, 0.6 mg/mL glutathione, and 10 mU/mL insulin[26,53]. To perform RNAi, 400 pmol of siRNA duplex was introduced into 3.0×10^6 cells using the Cell Line 96-well Nucleofector Kit SF (Lonza) with program DG150 of the 96-well Shuttle Device (Lonza). After 48 h incubation, cells were transfected with another 400 pmol of siRNA and incubated further for 48 h. For HP1a KD, RNAi was performed once to avoid cell damage. All siRNAs used are shown in Supplementary Table 2.

Fly stock

Flies were raised in standard Bloomington medium at 26 °C. Oregon-R was employed as a WT strain. For DART4 GLKD, BDSC 4937 (nosGAL4, w[1118]; P{w[+mC]=GAL4::VP16-nanos.UTR}CG6325[MVD1]) and BDSC 36833 (y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GL01073}attP2) were used as the maternal GAL4 driver and DART4 UAS-RNAi lines, respectively. BDSC 36303 (y[1] v[1]; P{y[+t7.7]=CaryP}attP2) was used as a control (ctrl).

Western blotting

Western blotting was performed as described previously[54]. Antibodies are summarized in Supplementary Table 3. Nuclear extraction and immunoprecipitation were performed as described previously[55].

Immunofluorescence

Immunofluorescence of OSCs was performed essentially as described previously[56]. OSCs were fixed using 4% formaldehyde (Wako) and permeabilized in PBS containing 0.1% Triton X-100 for 24 h at RT. For DAPI staining, VECTASHIELD Mounting Medium with DAPI (Vector Laboratories) was used. Immunofluorescence of ovaries was performed as described previously[10]. All images were obtained using a confocal laser scanning microscope (Olympus FV3000 or Carl Zeiss LSM 710). For image processing, ZEN software (Carl Zeiss) was used. Antibodies used are listed in Supplementary Table 3.

    Generation of mouse anti-Rhino monoclonal antibody

Anti-Rhino monoclonal antibody for immunofluorescence in the ovary were raised

against N-terminal 200 residues of Rhino using a protocol published previously[57]. The

N-terminal 200 residues of Rhino were fused to glutathione S-transferase (GST) and the

purified fusion protein was used as antigen to immunize mice. The expression vector

was produced by inserting the cDNA encoding the Rhino fragment into pGEX-5X-3

(Cytiva) by Infusion (Takara). The primers used are listed in Supplementary Table 1. The

specificity of the antibody was validated using OSCs (Supplementary Fig. 9A).

RNA extraction and RT-qPCR

Total RNAs were extracted from OSCs or ovaries with the ISOGEN II reagent (Nippon

Gene) and treated with DNase (Life Technologies) to eliminate DNA contaminations. The

cDNA libraries were prepared using the Transcriptor First Strand cDNA Synthesis Kit

(Roche). For RT-qPCR, cDNAs were amplified using the StepOnePlus Real-Time PCR

System (Applied Biosystems) with PowerUp SYBR Green Master Mix (Thermo Fisher

Scientific). The expression level of target genes and transposons in OSCs and ovary

were calculated by $\Delta\Delta$Ct method using *ribosomal protein 32* (*RP49/RpL32*) as an internal

control. Data are presented as mean values ± SE. The primers used are listed in

Supplementary Table 1.

OSC ChIP

OSCs were fixed in PBS containing 1% formaldehyde for 10 min at RT and glycine was

added to 150 mM to stop the fixation. Cells were collected and suspended in swelling

buffer [25 mM HEPES-KOH (pH 7.3), 1.5 mM $MgCl_2$, 10 mM KCl, 1 mM DTT, 0.1% NP-

40, supplemented with cOmplete ULTRA Tablets, EDTA-free (Roche)]. The nuclei were

collected by centrifugation at 1,000 g for 5 min at 4°C, suspended in sonication buffer

[50 mM HEPES-KOH (pH 7.4), 140 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1%

sodium deoxycholate, 0.1% SDS, supplemented with cOmplete ULTRA Tablets, EDTA-

free], and sonicated using a Covaris S220 Focused ultrasonicator for 10 min at 4°C with

setting of peak power of 200, duty factor of 10.0, and 200 cycles/burst. The chromatin

fraction was obtained by centrifugation at 20,000 g for 15 min at 4°C and diluted with

sonication buffer to make the DNA concentration to 0.2 mg/mL. Immunoprecipitation was

performed O/N at 4°C using 1mL of chromatin fraction per ChIP. Antibodies used are

listed in Supplementary Table 3. The reaction mixture was mixed with 25 μL of

Dynabeads ProteinG (Invitrogen) and incubated for 1 h at 4°C. Beads were washed one

time each with low-salt buffer [20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 2 mM EDTA,

0.1% Triton X-100, 0.1% SDS, and 1 mM phenylmethylsulfonyl fluoride (PMSF)], high-

salt buffer [20 mM Tris-HCl (pH 8.0), 500 mM NaCl, 2 mM EDTA, 0.1% Triton X-100,

0.1% SDS, and 1 mM PMSF], and LiCl buffer [20 mM Tris-HCl (pH 8.0), 250 mM LiCl, 2

mM EDTA, 1% NP-40, 1% sodium deoxycholate, and 1 mM PMSF], and then twice with

TE buffer [20 mM Tris-HCl (pH 8.0), 1 mM EDTA, and 1 mM PMSF]. The beads were

suspended in elution buffer [50 mM Tris-HCl (pH 8.0), 10 mM EDTA, and 1% SDS] and

incubated at 65°C for 30 min. For de-crosslinking, the supernatant was incubated O/N

at 65°C after supplement of NaCl to 200mM. RNAs and proteins were eliminated from

the sample by incubating at 37°C for 30 min with 10 mg/mL RNase A and 20 mg/mL

proteinase K at 55°C for 1 h. The DNA in the supernatant was purified by EconoSpin

column IIa (Funakoshi). 50 μL of nuclease-free water was added to it and the sample

was centrifuged at 13,000 rpm for 1 min to obtain ChIP product. The following

experiments were conducted to evaluate the specificity of the antibodies involved in the

main claims of this paper in ChIP-seq: H3R8me2a: ChIP-seq using two different anti-

H3R8me2a antibodies showed that signal was obtained at the same genomic locations

(Supplementary Fig. 9B). H3R17me2a: A reduction in DART4-KD cells was

demonstrated in separate experiments using western blotting (Supplementary Fig. 9C).

H4R3me2a: A histone peptide array kit (Active motif) confirmed that the antibody

recognizes the H4R3me2a peptide with high specificity (Supplementary Fig. 9D). The

histone peptide array was performed according to the manufacturer's protocol.

Ovary ChIP

150 ovary pairs were crosslinked in PBS containing 1% formaldehyde for 10 min at RT,

quenched with 180 mM glycine, and washed twice in PBS. Ovaries were flash frozen

and suspended with swelling buffer (see "OSC ChIP"). Ovaries were disrupted using a

Dounce homogenizer and incubated at 4°C for 10 min. The nuclei were collected by

centrifugation at 1,000 g for 5 min at 4°C, suspended in sonication buffer, and sonicated

(see "OSC ChIP"). The process after this is as described above (see "OSC ChIP").

Antibodies used are listed in Supplementary Table 3.

ChIP-seq library preparation

ChIP-seq libraries were constructed using NEBNext Ultra II FS DNA Library Prep Kit for

Illumina (New England BioLabs) using ChIP products from "OSC ChIP" and "Ovary ChIP".

In the adaptor ligation step, the original adaptor reagent was diluted 10-fold. Adapter

ligated DNAs were PCR-amplified (10−12 cycles) with Illumina primers. The PCR

products of 100−1,000 bp (including adaptor and primer sequences) were purified using

AMPure XP beads (Beckman Coulter) and captured on an Illumina flow cell. The libraries

were sequenced using the Illumina MiSeq.

Computational analysis of ChIP-seq

Prior to mapping, low-quality reads were removed and adapters were trimmed using

Fastp[58]. The filtered reads were mapped to the genome of *Drosophila melanogaster*

(dm6) using Bowtie 2[59]. Only uniquely mapped reads were used in the analysis using

samtools view command option -q 42[60]. For genome browser view, we mapped these

unique reads and the scores of the *Drosophila* genome mappability calculated using

GenMap software with the k-mer size set to 150[61]. We performed ChIP-seq experiment

using at least two biological replicates, except for ChIP with anti-FLAG antibody in OSCs

under Dox = 0.05 and 0.5ng/mL conditions. The replicates were highly correlated, so we

merged them together and used the merged libraries for analyses. Because of the low

mapping score, four replicates were prepared for Rhino and three for Kipferl in the GLKD

condition. Peakcall was completed with MACS2 with q = 0.01 settings using input ChIP-

seq data as control unless otherwise noted in figure legends[62]. For FLAG-Rhino peakcall

in OSCs, anti-FLAG ChIP-seq data in Dox = 0 ng/mL condition was used as control. For

Rhino peakcall in Oregon-R ovary in Fig. 1C, D and Fig. 4B, and H3K9me3 peakcall in

OSCs in Fig. 1F, options --broad --broad-cutoff 0.01 were used. For Fig. 1D, called peak

of more than 1.5 kb were extracted to select piRNA clusters where Rhino was localized

broadly. For heatmaps and profile plots, ChIP scores were calculated by deeptools[63].

Bigwig files of ChIP were produced by the bamCoverage / bamCompare program of

deeptools and matrix data were calculated by the computeMatrix program of deeptools

using bigwig files and plotted. For bamCompare program, anti-FLAG ChIP data with Dox

= 0 (ng/mL) was subtracted as control for FLAG-Rhino in OSCs and input for the other

data. Mapping file correlation and hierarchical clustering were computed using

multibamSummary programs of deeptools. To draw the Venn diagram, the narrowPeak

file obtained with Macs2 callpeak was stored in the findOverlapsOfPeaks function in R's

ChIPpeakanno library and visualized with makeVennDiagram program[64]. The manorm

was used for quantitative comparison of called peaks[65]. As ChIP-seq quality metrics,

aligned rates are provided in Supplementary Table 4, while library complexity values and

the SSP (strand cross-correlation peak)[66], a signal-to-noise ratio that can be calculated

independently of read number or peak shape, are listed in Supplementary Table 5.

ChIP-qPCR

Real-time qPCR was performed using PowerUp SYBR Green Master Mix and

StepOnePlus real-time PCR system. Data are presented as mean values ± SE. The

primers used are listed in Supplementary Table 2.

Definition of DART4-dependent piRNA source loci

DART4-dependent piRNA source loci were defined using manorm results based on the

macs2 callpeak with option p = 0.01 and --broad --broad cutoff 0.1 output files that met

the following seven criteria. First, M value > 0.5 is required for Rhino to be predominantly

enriched against DART4 GLKD. Second, the sharp peaks of the mononucleosome units

were excluded because of their non-spreading initial Rhino localization. Third, peaks

overlapped with DART4-dependent H3R17me2a were excluded because these peaks

cannot distinguish spread Rhino and initial Rhino loading. Fourth, each Rhino peak was

checked in the genome browser, and if there were consecutive Rhino peaks around it

that met the first, second and third criteria, they were grouped together as one peak.

Fifth, peaks that overlap with major piRNA clusters such as 42AB, 38C, 80F, and 102F

were excluded because they are highly transcribed piRNA clusters. Sixth, peaks on

subtelomeres and pericentromeres were excluded because ChIP-seq reads cannot

distinguish piRNA clusters or repeats. Finally, peaks whose localization spanned the

entire length of a single transposon were excluded because they may reflect genomic

differences between ctrl GLKD and DART4 GLKD, such as copy number variation of

transposons. The combined peaks that met these seven criteria were defined as single

DART4 piSL together with the surrounding contiguous ctrl GLKD-specific Rhino

localizations if they were more than 5 kb. The genomic location of DART4 piSL in Fig.

6C was visualized using Phenogram software[67]. The regions of DART4 piSL are listed in

Supplementary Table 6. The regions where Rhino localization increased upon DART4

GLKD were identified as regions with an M value of less than -2 and are listed in

Supplementary Table 7. Genes or transposons matched by piRNAs produced from

DART4 piSL were analyzed using bedtools and are listed in Supplementary Table 8.

Ovary piRNA extraction

100 ovary pairs were disrupted using a Dounce homogenizer and incubated in lysis

buffer [20 mM HEPES-KOH (pH7.3), 150 mM NaCl, 1 mM EDTA, 1 mM DTT, 0.1% NP-

40, 40 U/ml RNasin (Promega), 2 µg/ml pepstatin, 2 µg/ml leupeptin, 0.5% aprotinin] at

4°C for 10 min. The homogenized ovaries were centrifuged at 13,500rpm for 15 min at

4°C. The supernatant was then collected and cooled on ice. 10µg of anti-Piwi antibody

was immobilized on Dynabeads protein G (Invitrogen) in lysis buffer at 4°C for 5 min.

Beads were incubated with ovarian lysates at 4°C for 2 h, and then washed three times

with low-salt wash buffer [20 mM HEPES-KOH (pH7.3), 300 mM NaCl, 1 mM DTT, 0.05% NP-40, 2 µg/ml pepstatin, 2 µg/ml leupeptin, 0.5% aprotinin], three times with high-salt wash buffer [20 mM HEPES-KOH (pH7.4), 500 mM NaCl, 1 mM DTT, 0.05% NP-40, 2 µg/ml pepstatin, 2 µg/ml leupeptin, 0.5% aprotinin]. After wash, Piwi-bound RNAs were extracted from the beads with phenol–chloroform (Ambion) and then precipitated with ethanol. The precipitated RNAs were denatured with Gel Loading Buffer II (Ambion) for 5 min at 95°C, then electrophoresed on 12% urea-PAGE gel. 20-30 nt nucleic acids were gel purified as piRNAs. piRNA-seq library preparation followed a previous study[68].

Computational analysis of piRNA-seq

Raw reads were adapter trimmed and the random 4 bases of each side were removed using cutadapt[69]. Trimmed reads were mapped to the dm6 genome using bowtie with the option -m 1[70]. For read count in Supplementary Fig. 7D, this option was excluded. Read counts for each piRNA cluster were performed using bedtools[71]. For the analysis of piRNA in Rhino GLKD, GSM1580790 and GSM1580793 datasets[72] were used as piRNA-seq data in the ovaries of ctrl GLKD and Rhino GLKD, respectively.

Egg hatching assay

50 males and 50 virgin females were collected and mated in a cage with apple juice agar plate and yeast paste for 2 days at 25°C. After exchanging plates twice in the morning, the eggs on the plates were collected within 3 hours. Two hundred eggs were placed on new apple juice agar plates and incubated at 25°C for 2 days, followed by counting the number of hatched larvae.

Evolutionary analysis of piRNA clusters

The 500 bp around the ends of a novel dataset comprising piRNA clusters categorized

into eight levels of evolutionary novelty[44] ("master-list" of Supplementary Table S4 in

reference 44) was used for analysis. The ADMA-histones and input ChIP-seq reads

mapped to the piRNA cluster regions were counted by bedtools and reads per piRNA

cluster were calculated.Genomic regions with RPKM values above 30 in the input were

excluded as they were considered ChIP-seq background noise. For Supplemntary Fig.

8B and 8C, only piRNA clusters with non-zero piRNA RPKM were used for analysis in

the Oregon-R (Supplemntary Fig. 8B) and ctrl GLKD (Supplemntary Fig. 8C) strains,

respectively. SRR4473614 was used as the piRNA of Oregon-R. Extraction of piRNA

from SRR4473614 followed the methods of the original paper[50,73].

Computational analysis of RNA-seq

Prior to mapping, low-quality reads were removed and adapters were trimmed using

Fastp[58]. The filtered reads were mapped to the genome of *Drosophila melanogaster*

(dm6) using STAR[74] with default settings. Reads of each gene were calculated using

featureCount[75] with default settings.

Data availability

ChIP-seq and piRNA-seq data generated in this study have been deposited in the

Sequence Read Archive (SRA) under accession number PRJNA1085006 and the Gene

Expression Omnibus (GSE) under accession number GSE261403. All other data

supporting the findings of this study are available from the corresponding authors upon

request.

Acknowledgments

We are grateful to all members of the Siomi laboratory for discussions and comments on this work, especially Kaoru Sato. We also thank Soichiro Yamanaka, Tetsuji Kakutani, and Soichi Inagaki for critical reading of the manuscript, Takashi Fukaya and Hiroshi Kimura for discussion, Yuica Koga for technical assistance with *Drosophila* genetics, and Akashi Taguchi for support with the Illumina sequencing. We are also grateful to Julius Brennecke and Laszlo Tirian for antibodies and Harvard TRiP and Bloomington stock centers for *Drosophila* flies. This study was supported by MEXT KAKENHI Grant Number JP19H05466 and JSPS Grant-in Aid for Transformative Research Areas [A] 25H01303 (to M.C.S.), JSPS KAKENHI Grant Number 21K20629 and 23K14232 (to R.S.), JSPS KAKENHI Grant Numbers 17K08644 and 20H03439 (to K.M.), and JSPS KAKENHI Grant Numbers 24K18093 (to Y.N.)

Author contributions

R.S. and R.H. conducted biochemical experiments. R.S. and Y.N. performed bioinformatic analysis. H.I. produced FLAG-Rhino inducible OSCs. K.M. produced anti-Rhino antibody. R.S. and M.C.S. conceived the project and designed the experiments. R.S. and M.C.S. wrote the manuscript with input from all authors.

Disclosure and competing interest statement

The authors declare that they have no conflicts of interest.

References

1. Ghildiyal M, Zamore PD. Small silencing RNAs: an expanding universe. *Nat Rev Genet* **10,** 94-108 (2009)

2. Iwasaki YW, Siomi MC, Siomi H. PIWI-Interacting RNA: Its Biogenesis and Functions. *Annu Rev Biochem* **84,** 405-433 (2015).

3. Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD. PIWI-interacting RNAs: small RNAs with big functions. *Nat Rev Genet* **20,** 89-108 (2019).

4. Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell* **128,** 1089-1103 (2007).

5. Klattenhoff C, Xi H, Li C, Lee S, Xu J, Khurana JS, Zhang F, Schultz N, Koppetsch BS, Nowosielska A, Seitz H, Zamore PD, Weng Z, Theurkauf WE. The *Drosophila* HP1 homolog Rhino is required for transposon silencing and piRNA production by dual-strand clusters. *Cell* **138,** 1137-1149 (2009).

6. Mohn F, Sienski G, Handler D, Brennecke J. The rhino-deadlock-cutoff complex licenses noncanonical transcription of dual-strand piRNA clusters in *Drosophila*. *Cell* **157,** 1364-1379 (2014).

7. Zhang Z, Wang J, Schultz N, Zhang F, Parhad SS, Tu S, Vreven T, Zamore PD, Weng Z, Theurkauf WE. The HP1 homolog rhino anchors a nuclear complex that suppresses piRNA precursor splicing. *Cell* **157,** 1353-1363 (2014).

8. Yu B, Cassani M, Wang M, Liu M, Ma J, Li G, Zhang Z, Huang Y. Structural insights into Rhino-mediated germline piRNA cluster formation. *Cell Res* **25,** 525-528 (2015).

9. Akkouche A, Mugat B, Barckmann B, Varela-Chavez C, Li B, Raffel R, Pélisson A, Chambeyron S. Piwi Is Required during *Drosophila* Embryogenesis to License Dual-Strand piRNA Clusters for Transposon Repression in Adult Ovaries. *Mol Cell* **66,** 411-419 (2017).

10. Andersen PR, Tirian L, Vunjak M, Brennecke J. A heterochromatin-dependent transcription machinery drives piRNA expression. *Nature* **549,**54-59 (2017).

11. Yu B, Lin YA, Parhad SS, Jin Z, Ma J, Theurkauf WE, Zhang ZZ, Huang Y. Structural insights into Rhino-Deadlock complex for germline piRNA cluster specification. *EMBO Rep* **19,** e45418 (2018).

12. Baumgartner L, Handler D, Platzer SW, Yu C, Duchek P, Brennecke J. The *Drosophila* ZAD zinc finger protein Kipferl guides Rhino to piRNA clusters. *Elife* **11,** e80067 (2022).

13. Casier K, Autaa J, Gueguen N, Delmarre V, Marie PP, Ronsseray S, Carré C,

Brasset E, Teysset L, Boivin A. The histone demethylase Kdm3 prevents auto-immune piRNAs production in *Drosophila*. *Sci Adv* **9,** eade3872 (2023).

14. Luo Y, He P, Kanrar N, Fejes Toth K, Aravin AA. Maternally inherited siRNAs initiate piRNA cluster formation. *Mol Cell* **83,** 3835-3851 (2023).

15. Volpe AM, Horowitz H, Grafer CM, Jackson SM, Berg CA. *Drosophila* rhino encodes a female-specific chromo-domain protein that affects chromosome structure and egg polarity. *Genetics* **159,** 1117-1134 (2001).

16. Vermaak D, Henikoff S, Malik HS. Positive selection drives the evolution of rhino, a member of the heterochromatin protein 1 family in *Drosophila*. *PLoS Genet* **1,** 96-108 (2005).

17. Pane A, Jiang P, Zhao DY, Singh M, Schüpbach T. The Cutoff protein regulates piRNA cluster expression and piRNA production in the *Drosophila* germline. *EMBO J* **30,** 4601-4615 (2011).

18. Czech B, Preall JB, McGinn J, Hannon GJ. A transcriptome-wide RNAi screen in the *Drosophila* ovary reveals factors of the germline piRNA pathway. *Mol Cell* **50,** 749-761 (2013).

19. Chen YA, Stuwe E, Luo Y, Ninova M, Le Thomas A, Rozhavskaya E, Li S, Vempati S, Laver JD, Patel DJ, Smibert CA, Lipshitz HD, Toth KF, Aravin AA. Cutoff Suppresses RNA Polymerase II Termination to Ensure Expression of piRNA Precursors. *Mol Cell* **63,** 97-109 (2016).

20. Pritykin Y, Brito T, Schupbach T, Singh M, Pane A. Integrative analysis unveils new functions for the *Drosophila* Cutoff protein in noncoding RNA biogenesis and gene regulation. *RNA* **23,** 1097-1109 (2017).

21. Le Thomas A, Stuwe E, Li S, Du J, Marinov G, Rozhkov N, Chen YC, Luo Y, Sachidanandam R, Toth KF, Patel D, Aravin AA. Transgenerationally inherited piRNAs trigger piRNA biogenesis by changing the chromatin of piRNA clusters and inducing precursor processing. *Genes Dev* **28,** 1667-1680 (2014).

22. Saito K, Inagaki S, Mituyama T, Kawamura Y, Ono Y, Sakota E, Kotani H, Asai K, Siomi H, Siomi MC. A regulatory circuit for piwi by the large Maf gene traffic jam in *Drosophila*. *Nature* **461,** 1296-1299 (2009).

23. Sumiyoshi T, Sato K, Yamamoto H, Iwasaki YW, Siomi H, Siomi MC. Loss of l(3)mbt leads to acquisition of the ping-pong cycle in *Drosophila* ovarian somatic cells. *Genes Dev* **30,** 1617-1622 (2016).

24. Sato K, Siomi MC. The piRNA pathway in *Drosophila* ovarian germ and somatic cells. *Proc Jpn Acad Ser B Phys Biol Sci* **96,** 32-42 (2020)

25. Dong F, Li Q, Yang C, Huo D, Wang X, Ai C, Kong Y, Sun X, Wang W, Zhou Y, Liu

X, Li W, Gao W, Liu W, Kang C, Wu X. PRMT2 links histone H3R8 asymmetric dimethylation to oncogenic activation and tumorigenesis of glioblastoma. *Nat Commun* **9,** 4552 (2018).

26. Wang YC, Li C. Evolutionarily conserved protein arginine methyltransferases in non-mammalian animal systems. *FEBS J* **279,** 932-945 (2012).

27. Wang YC, Wang JD, Chen CH, Chen YW, Li C. A novel BLAST-Based Relative Distance (BBRD) method can effectively group members of protein arginine methyltransferases and suggest their evolutionary relationship. *Mol Phylogenet Evol* **84,** 101-111 (2015).

28. Boulanger MC, Miranda TB, Clarke S, Di Fruscio M, Suter B, Lasko P, Richard S. Characterization of the *Drosophila* protein arginine methyltransferases DART1 and DART4. *Biochem J* **379,** 283-289 (2004).

29. Scaramuzzino C, Monaghan J, Milioto C, Lanson NA Jr, Maltare A, Aggarwal T, Casci I, Fackelmayer FO, Pennuto M, Pandey UB. Protein arginine methyltransferase 1 and 8 interact with FUS to modify its sub-cellular distribution and toxicity *in vitro* and *in vivo*. *PLoS One* **8,** e61576 (2013).

30. Xu J, Wang AH, Oses-Prieto J, Makhijani K, Katsuno Y, Pei M, Yan L, Zheng YG, Burlingame A, Brückner K, Derynck R. Arginine Methylation Initiates BMP-Induced Smad Signaling. *Mol Cell* **51,** 5-19 (2013).

31. Jäckel S, Summerer AK, Thömmes CM, Pan X, Voigt A, Schulz JB, Rasse TM, Dormann D, Haass C, Kahle PJ. Nuclear import factor transportin and arginine methyltransferase 1 modify FUS neurotoxicity in *Drosophila*. *Neurobiol Dis* **74,** 76-88 (2015).

32. Yuan Q, Tian R, Zhao H, Li L, Bi X. Multiple Arginine Residues Are Methylated in *Drosophila* Mre11 and Required for Survival Following Ionizing Radiation. *G3* **8,** 2099-2106 (2018).

33. Chen D, Ma H, Hong H, Koh SS, Huang SM, Schurter BT, Aswad DW, Stallcup MR. Regulation of transcription by a protein methyltransferase. *Science* **284,** 2174-2177 (1999).

34. Cakouros D, Daish TJ, Mills K, Kumar S. An arginine-histone methyltransferase, CARMER, coordinates ecdysone-mediated apoptosis in *Drosophila* cells. *J Biol Chem* **279,** 18467-18471 (2004).

35. Urwyler O, Zhang L, Li X, Imboden H, Suter B. Tissue-dependent subcellular localization of *Drosophila* arginine methyl-transferase 4 (DART4), a coactivator whose overexpression affects neither viability nor differentiation. *Differentiation* **75,** 757-765 (2007).

36. Villar-Garea A, Forne I, Vetter I, Kremmer E, Thomae A, Imhof A. Developmental

regulation of N-terminal H2B methylation in *Drosophila* melanogaster. *Nucleic*

Acids Res **40,** 1536-1549 (2012).

37. Teo RYW, Anand A, Sridhar V, Okamura K, Kai T. Heterochromatin protein 1a

functions for piRNA biogenesis predominantly from pericentric and telomeric

regions in *Drosophila. Nat Commun* **9,** 1735 (2018).

38. Schurter BT, Koh SS, Chen D, Bunick GJ, Harp JM, Hanson BL, Henschen-

Edman A, Mackay DR, Stallcup MR, Aswad DW. Methylation of histone H3 by

coactivator-associated arginine methyltransferase 1. *Biochemistry* **40,** 5747-5756

(2001).

39. Miao F, Li S, Chavez V, Lanting L, Natarajan R. Coactivator-associated arginine

methyltransferase-1 enhances nuclear factor-kappaB-mediated gene transcription

through methylation of histone H3 at arginine 17. *Mol Endocrinol* **20,** 1562-1573

(2006).

40. Wang H, Huang ZQ, Xia L, Feng Q, Erdjument-Bromage H, Strahl BD, Briggs SD,

Allis CD, Wong J, Tempst P, Zhang Y. Methylation of histone H4 at arginine 3

facilitating transcriptional activation by nuclear hormone receptor. *Science* **293,**

853-857 (2001).

41. Huang S, Litt M, Felsenfeld G. Methylation of histone H4 by arginine

methyltransferase PRMT1 is essential *in vivo* for many subsequent histone

modifications. *Genes Dev* **19,** 1885-1893 (2005).

42. Rosenkranz D, Zischler H, Gebert D. piRNAclusterDB 2.0: update and expansion

of the piRNA cluster database. *Nucleic Acids Res* **50**,D259-D264 (2022).

43. Gebert D, Neubert LK, Lloyd C, Gui J, Lehmann R, Teixeira FK. Large Drosophila

germline piRNA clusters are evolutionarily labile and dispensable for transposon

regulation. *Mol Cell* **81,** 3965-3978 (2021).

44. Srivastav S, Feschotte C, Clark AG. Rapid evolution of piRNA clusters in the

Drosophila melanogaster ovary. *Genome Res* 34,711-724 (2024).

45. Blanc RS, Richard S. Arginine Methylation: The Coming of Age. *Mol Cell* **65,** 8-24

(2017).

46. Šimčíková D, Gelles-Watnick S, Neugebauer KM. Tudor-dimethylarginine

interactions: the condensed version. *Trends Biochem Sci* **48,** 689-698 (2023).

47. Yang Y, Lu Y, Espejo A, Wu J, Xu W, Liang S, Bedford MT. TDRD3 is an effector

molecule for arginine-methylated histone marks. *Mol Cell* **40,** 1016-1023 (2010).

48. Radion E, Sokolova O, Ryazansky S, Komarov PA, Abramov Y, Kalmykova A. The

Integrity of piRNA Clusters is Abolished by Insulators in the *Drosophila* Germline.

Genes (Basel) **10,** 209 (2019).

49. Luo S, Zhang H, Duan Y, Yao X, Clark AG, Lu J. The evolutionary arms race

between transposable elements and piRNAs in *Drosophila* melanogaster. *BMC*

Evol Biol **20,** 14 (2020).

50. Wierzbicki F, Kofler R. The composition of piRNA clusters in *Drosophila*

melanogaster deviates from expectations under the trap model. *BMC Biol* **21,** 224

(2023).

51. Yamanaka S, Siomi MC, Siomi H. piRNA clusters and open chromatin structure.

Mob DNA **5,** 22 (2014).

52. Takeuchi C., Murano K., Ishikawa M., Okano H., Iwasaki W.Y. Generation of stable

Drosophila ovarian somatic cell lines using piggyBac system. *Methods Mol Biol*

2509, 143–153 (2022).

53. Niki Y, Yamaguchi T, Mahowald AP. Establishment of stable cell lines of *Drosophila*

germ-line stem cells. *Proc Natl Acad Sci U S A* **103,** 16325-16330 (2006).

54. Saito K, Nishida KM, Mori T, Kawamura Y, Miyoshi K, Nagami T, Siomi H, Siomi

MC. Specific association of Piwi with rasiRNAs derived from retrotransposon and

heterochromatic regions in the *Drosophila* genome. *Genes Dev* **20,** 2214-2222

(2006).

55. Yamamoto-Matsuda H, Miyoshi K, Moritoh M, Yoshitane H, Fukada Y, Saito K,

Yamanaka S, Siomi MC. Lint-O cooperates with L(3)mbt in target gene

suppression to maintain homeostasis in fly ovary and brain. EMBO Rep. **23,**

e53813 (2022).

56. Saito K, Ishizu H, Komai M, Kotani H, Kawamura Y, Nishida KM, Siomi H, Siomi

MC. Roles for the Yb body components Armitage and Yb in primary piRNA

biogenesis in *Drosophila*. *Genes Dev* **24,** 2493-2498 (2010).

57. Murano K, Iwasaki YW, Ishizu H, Mashiko A, Shibuya A, Kondo S, Adachi S,

Suzuki S, Saito K, Natsume T, Siomi MC, Siomi H. Nuclear RNA export factor

variant initiates Piwi-piRNA-guided co-transcriptional silencing. *EMBO J* **38,**

e102870 (2019).

58. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34,** i884-i890 (2018).

59. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat*

Methods **9,** 357-359 (2012).

60. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

Durbin R. 1000 Genome Project Data Processing Subgroup. The Sequence

Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078-2079 (2009).

61. Pockrandt C, Alzamel M, Iliopoulos CS, Reinert K. GenMap: ultra-fast computation of genome mappability. Bioinformatics. **36,** 3687-3692. (2020).

62. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, Liu XS. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9,** R137 (2008).

63. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44,** W160-5 (2016).

64. Zhu L, Gazin C, Lawson N, Pagès H, Lin S, Lapointe D, Green M. ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data. *BMC Bioinformatics*, **11**, 237 (2010).

65. Shao Z, Zhang Y, Yuan GC, Orkin SH, Waxman DJ. MAnorm: a robust model for quantitative comparison of ChIP-Seq data sets. *Genome Biol* **13,** R16 (2012).

66. Nakato R, Shirahige K. Sensitive and robust assessment of ChIP-seq read distribution using a strand-shift profile. *Bioinformatics* **34,** 2356-2363 (2018).

67. Wolfe D, Dudek S, Ritchie MD, Pendergrass SA. Visualizing genomic information across chromosomes with PhenoGram. BioData Min **6,** 18 (2013).

68. Iwasaki YW, Ishino K, Siomi H. Deep sequencing and high-throughput analysis of PIWI-associated small RNAs. *Methods* **126,** 66-75. (2017).

69. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal* **17,** 10–12 (2011).

70. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10,** R25 (2009).

71. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26,** 841-842 (2010)

72. Mohn F, Handler D, Brennecke J. piRNA-guided slicing specifies transcripts for Zucchini-dependent, phased piRNA biogenesis. *Science* **348,** 812-817 (2015).

73. Fast I, Hewel C, Wester L, Schumacher J, Gebert D, Zischler H, Berger C, Rosenkranz D. Temperature-responsive miRNAs in *Drosophila* orchestrate adaptation to different ambient temperatures. RNA. **23,** 1352-1364 (2017).

74. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner. Bioinformatics **29**, 15-21 (2013).

75. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. **30**, 923-30 (2014).

Figure 1. Rhino forms nuclear foci and binds the genome in OSCs

(A) Confocal immunofluorescent images of FLAG-Rhino (red) in FLAG-Rhino inducible

OSCs upon Dox induction (5 ng/mL). H3K9me3 and OSC nucleus (DAPI staining) are

shown in green and blue. Scale bar represents 10 μm. (B) Genome snapshot showing

ChIP-seq in major dual-strand piRNA clusters (42AB and 38C). The gray-shaded areas

indicate the sites at which FLAG-Rhino is localized in each piRNA cluster in OSCs.

piRNA cluster 38C has two piRNA-producing regions, labeled 38C1 and 38C2. (C)

Heatmap showing FLAG-Rhino localization of OSCs in Rhino peaks of *Drosophila* wild-

type Oregon-R ovary. The regions of 6 kb around the Rhino localization site were plotted

based on macs2 callpeak results. (D) Metaplot showing the localization of FLAG-Rhino

in OSCs at piRNA clusters. For piRNA clusters, broad peaks of Rhino localization in the

ovary greater than 1.5 kb were used. The overlap of each Rhino peak with annotated

piRNA clusters (e.g., 42AB, 38C, and 80F) was identified, as listed in Supplementary

Table 9. (E) Venn diagram showing the overlap of Rhino, H3K9me3, and H3K9me2

peaks in OSCs. (F) Heatmap showing the signals of ChIP-seq using anti-FLAG, anti-

H3K9me3, and anti-H3K9me2 in FLAG-Rhino peaks of OSCs. The regions of 2 kb

around the FLAG-Rhino localization sites were plotted based on macs2 callpeak results.

(G) Heatmap showing the signals of ChIP-seq using anti-H3K9me3, anti-FLAG, anti-

HP1a antibodies at H3K9me3 broad peaks of OSCs. The regions of 10 kb around the

FLAG-Rhino localization sites were plotted based on macs2 callpeak results.

Figure 2. ADMA-histones tend to co-localize with FLAG-Rhino in OSCs

(A) Heatmap showing key histone modifications and HP1a ChIP-seq results at FLAG-

Rhino localization sites (828 regions) in OSCs. The regions of 2 kb around the FLAG-

Rhino localization sites were plotted based on macs2 callpeak results. (B) Genome

snapshot showing all histone modifications' ChIP-seq data in major dual-strand piRNA

clusters (38C). The gray-shaded areas indicate the localization sites of FLAG-Rhino in

each piRNA cluster in OSCs. (C) Metaplot showing the localization of ADMA-histones

and FLAG-Rhino in OSCs at piRNA clusters. For piRNA clusters, broad peaks of Rhino

localization in the ovary greater than 1.5 kb were used. (D) Pearson correlation

coefficient comparing the coverage per 100 bp bin size of all ChIP-seq mapping results

to the dm6 genome and its hierarchical clustering. (E) Venn diagram showing sites of

overlapping localization of FLAG-Rhino, H3K9me3, H3R2me2a, H3R8me2a,

H3R17me2a, H3R26me2a, H4R3me2a, and all ADMA-histones in OSCs.

Figure 3. Arginine methyltransferases DART1 and DART4 play a role in Rhino

genomic localization and foci formation in OSCs

(A) Confocal immunofluorescent images of FLAG-Rhino (red) and nuclei (blue; DAPI

stained) in the EGFP-, HP1a-, DART1-, DART4-, DART8-, and CG17726-KD OSCs.

Scale bar represents 10 μm. (B) Comparison of FLAG-Rhino foci sizes among the

EGFP-, HP1a-, DART1-, DART4-, DART8-, and CG17726-KD OSCs in (A). Thirty OSCs

from three biological replicates were analyzed. Statistical significance in differences of

Rhino foci between the genotypes was examined using Kolmogorov–Smirnov test. $*P <$

0.05, $**P < 0.01$, and $***P < 0.001$. HP1a KD: $***P = 8.49 \times 10^{-13}$, DART1 KD: $***P =$

1.83×10^{-7}, DART4 KD: $***P = 2.83 \times 10^{-4}$, DART8 KD: n.s. $P = 0.655$, CG17726 KD:

n.s. $P = 0.454$. The box indicates the lower (25%) and upper (75%) quartiles, and the

open circle indicates the median. Whiskers extend to the most extreme data points. (C)

ChIP-seq profile showing the localization of FLAG-Rhino, H3R2me2a, H3R8me2a,

H3R17me2a, H3R26me2a, and H4R3me2a in EGFP-, DART1-, and DART4-KD OSCs

at FLAG-Rhino localization sites. The genomic regions used to draw the profile are the same as in Fig. 1G. (D) Heatmap showing the localization of FLAG-Rhino, H3R17me2a, and H3R8me2a in EGFP- and DART4-KD OSCs at the sites of decreased FLAG-Rhino localization in DART4 KD. The regions with decreased FLAG-Rhino were defined as those with an M value greater than 0.5 by manorm. macs2 callpeak p = 0.05 was used for manorm with input as a control. (E) FLAG-Rhino ChIP-qPCR results of one site at which Rhino initially localized. DART1 KD: †P = 5.88×10^{-2}, DART4 KD: * P = 3.17×10^{-2}. (F) Venn diagram showing the overlap of H3R17me2a localization decreased in DART4 KD and H4R3me2a localization decreased in DART1 KD. The regions with decreased ADMA-histones were defined as those with an M value greater than 1 in manorm. macs2 callpeak p = 0.01 was used for manorm. (G) Venn diagram showing the overlap of Rhino localizations that were decreased by DART1 KD and DART4 KD. The decreased genomic sites are identified as in (F).

Figure 4. ADMA-histones co-localize with Rhino prior to Rhino spread in the ovary

(A) Genome browser snapshot showing the localization of H3R2me2a, H3R8me2a, H3R17me2a, H3R26me2a, and H4R3me2a in piRNA clusters in OSCs and Oregon-R ovary. The gray-shaded areas indicate the localization sites of FLAG-Rhino in each piRNA cluster in OSCs. (B) Heatmap showing the localization of H3R2me2a, H3R8me2a, H3R17me2a, H3R26me2a, and H4R3me2a at Rhino peaks in Oregon-R ovary. The Rhino peak regions shown in Fig. 1C were used. (C) Metaplot showing the localization of ADMA-histones in the ovary at piRNA clusters. For piRNA clusters, broad peaks of Rhino localization in the ovary greater than 1.5 kb were used.

Figure 5. DART4 GLKD has impacts on Rhino foci and Rhino genomic localization

(A) Confocal immunostaining images of Rhino (red) in ctrl- and DART4-GLKD ovaries.

DAPI shows nuclei (blue). Scale bar represents 10 μm. (B) Comparison of sizes of Rhino

foci between the ctrl- and DART4-GLKD ovaries in (A). Ten ovaries from two biological

replicates were analyzed. Statistical significance in differences of Rhino foci between the

two genotypes was examined as in Fig. 3B. DART4 GLKD: ***$P = 1.26 \times 10^{-10}$. (C) ChIP-

seq profile showing the localization of Rhino and H3R17me2a in ctrl- and DART4-GLKD

ovaries at the FLAG-Rhino localization sites in OSCs. The FLAG-Rhino peak regions,

as in Fig. 1G, are used. (D) Rhino ChIP-qPCR results of one site at which Rhino initially

localized and piRNA cluster 42AB. t-test was used for statistical hypothesis testing.

Rhino initial localization point: * $P = 4.03 \times 10^{-2}$, piRNA cluster 42AB: n.s. $P = 0.11$. (E)

Genome snapshot of ChIP-seq data in euchromatic regions. The gray-shaded areas

indicate genomic locations where Rhino localization is decreased in a H3R17me2a-

dependent manner. (F) Scatter plot comparing Rhino enrichment in ChIP-seq between

ctrl GLKD and DART4 GLKD. Each dot represents the square root of the read count per

million mapped reads (RPM) within the annotated piRNA clusters in piRNAcluster

database[42] (colored in black, 184 clusters, without clusters on chromosome Y and uni-

strand piRNA clusters flamenco/20A) and FLAG-Rhino loaded points in OSCs as in Fig.

5C (colored in red, 828 regions). The dotted lines represent the least-squares linear

regression fit to the data points: the black line for 184 piRNA clusters and the red line for

828 FLAG-Rhino–loaded points in OSCs.

Figure 6. DART4-dependent piRNA source loci (DART4 piSL**) are genomic loci**

where Rhino spreads from H3R17me2a

(A) Venn diagram showing the overlap between sites with reduced Rhino localization and sites where H3R17me2a was lost in DART4 GLKD. The regions where Rhino was decreased were defined as those with an M value greater than 0.5 by manorm, and the regions where H3R17me2a was lost were defined as those with an M value greater than 0. For manorm, the output results of macs2 callpeak p = 0.01 were used with input as a control. (B) Scatter plot comparing Rhino enrichment in ChIP-seq between ctrl GLKD and DART4 GLKD. Each dot represents the square root of the read count per million mapped reads (RPM) within the annotated piRNA clusters in piRNAcluster database[42] (colored in black, 184 clusters, without clusters on chromosome Y and uni-strand piRNA clusters flamenco/20A) and Rhino reduced regions in DART4 GLKD (colored in green, 43 clusters). The dotted lines represent an approximation by a linear regression model. (C) Chromosomal locations of Rhino reduced regions in DART4 GLKD and the estimated direction of Rhino spread based on initial localization in OSCs. Depressed areas indicate pericentromeres. Arrows show the spread direction based on ADMA-histones. Bidirectional arrows indicate that ADMA-histones are present at both ends and the direction of spread cannot be determined. Horizontal bars indicate clusters without ADMA-histones at both ends. (D) Genome snapshot showing Rhino reduced regions in DART4 GLKD around the *bx* gene. Black lines mark Rhino reduced regions boundaries, and gray-shaded areas highlight potential ADMA-histone localization at cluster ends. (E) ChIP-seq profiles of Rhino and H3R17me2a in ctrl GLKD and DART4 GLKD across all 43 Rhino reduced regions in DART4 GLKD, and Rhino in OSCs. (F) Scatter plot comparing piRNA enrichment between ctrl GLKD and DART4 GLKD. Each dot represents the logged RPM value within piRNA clusters (black: 184 annotated clusters; green: 43 Rhino reduced regions in DART4 GLKD). Dotted lines indicate linear

 regression. (G) Box plot showing RPKM of piRNAs uniquely mapped to Rhino sites in

 OSCs, Rhino reduced regions in DART4 GLKD, all piRNAcluster database[42]-annotated

 clusters, and highly transcribed clusters (42AB, 38C, 80F, 102F). Clusters with RPKM

 below 400 are shown.

Figure 7. ADMA-histones are enriched at the ends of evolutionarily young piRNA

clusters

(A) RPKM of each ChIP-seq dataset (Oregon-R) in the piRNA cluster dataset with eight

levels of evolutionary novelty. RPKM was calculated for the 500 bp region surrounding

the ends of the piRNA clusters. t-test was used for statistical hypothesis testing between

clusters found only in 1 strain (cluster 1, newly emerged) and clusters found in all 8

strains (cluster 8, evolutionarily conserved). All ADMA-histones: ** $P = 1.11 \times 10^{-3}$,

H3R2me2a * $P = 1.45 \times 10^{-2}$, H3R8me2a: *** $P = 1.27 \times 10^{-4}$, H3R17me2a: ** $P = 7.28$

$\times 10^{-3}$, H3R26me2a: n.s. $P = 0.23$, H4R3me2a: ** $P = 1.91 \times 10^{-3}$, input: * $P = 0.53$. (B)

Same analysis of Fig. 7A for each ChIP-seq dataset (ctrl GLKD and DART4 GLKD).

H3R17me2a (ctrl GLKD): ** $P = 2.77 \times 10^{-3}$, H3R17me2a (DART4 GLKD): n.s. $P = 0.24$,

input (ctrl GLKD): n.s. $P = 0.42$, input (DART4 GLKD): n.s. $P = 0.47$. (C) Model for the

piRNA source loci formation. In the cell conditions lacking Kipferl such as in OSCs, Rhino

initially localizes to bivalent nucleosomes containing H3K9me3 and ADMA-histones (left).

In the second step, Rhino spreading occurs, leading to the formation of DART4 piSL.

Transcription of adjacent regions and/or H3K9me2 enrichment may play a role in this

process (right). Rhino in DART4 piSL is lost if ADMA-histones are lost. It remains unclear

whether DART4 piSL can eventually develop into highly transcribed piRNA clusters, such

as 42AB, over the course of evolutionary time.

Supplementary Figure 1. Expression and localization of FLAG-Rhino at different

Dox concentrations

(A) Confocal immunofluorescent images of FLAG-Rhino (green) in FLAG-Rhino

inducible OSCs upon Dox induction (from 0.01 to 1.0 ng/mL). DAPI shows nuclei (blue).

Scale bar represents 10 μm. (B) Western blotting of FLAG-Rhino under each Dox

concentration. HP1a was detected as an internal control. (C) Genome snapshot showing

ChIP-seq in major dual-strand piRNA clusters (80F and 102F). The gray-shaded areas

indicate the sites at which FLAG-Rhino is localized in each piRNA cluster in OSCs. (D)

Heatmap showing the localization of FLAG-Rhino in OSCs under various Dox

concentrations and Rhino in ovary at FLAG-Rhino peaks of OSCs. The regions of 2 kb

around the FLAG-Rhino localization sites were plotted based on macs2 callpeak results.

(E) Genome snapshot showing ChIP-seq in major dual-strand piRNA clusters (42AB and

38C). The gray-shaded areas indicate the sites at which FLAG-Rhino is localized in each

piRNA cluster in OSCs. piRNA cluster 38C has two piRNA-producing regions, labeled

38C1 and 38C2.

Supplementary Figure 2. Behavior of ADMA-histones in FLAG-Rhino inducible

OSCs

(A) Genome snapshot showing all histone modifications ChIP-seq data in major dual-

strand piRNA clusters (42AB, 80F and 102F). The gray-shaded areas indicate the

localization sites of FLAG-Rhino in each piRNA cluster in OSCs. (B) Venn diagram

showing the sites of overlapping localization of each ADMA-histone. (C) Venn diagram

showing the sites of overlapping localization of ADMA-histones, HP1a, and H3K9me3.

Supplementary Figure 3. Behavior of FLAG-Rhino, HP1a, and ADMA-histones

 under DART1-KD and DART4-KD conditions

 (A) Confocal immunofluorescent images of FLAG-Rhino (red) and H3K9me3 (green) in

 FLAG-Rhino inducible OSCs. Dox was added at a final concentration of 5 ng/mL. DAPI

 shows nuclei (blue). Scale bar represents 10 μm. (B) Confocal immunofluorescent

 images of HP1a (red) and H3K9me3 (green) in FLAG-Rhino inducible OSCs. Dox was

 added in final concentration of 5 ng/mL. DAPI shows nuclei (blue). Scale bar represents

 10 μm. (C) Venn diagram showing the overlap of the decreased H3R8me2a regions in

 DART1 KD and DART4 KD. The decreased regions were defined as those with an M

 value greater than 0.5 in manorm. macs2 callpeak p = 0.01 was used for manorm with

 input as a control. (D) Venn diagram showing the overlap of the regions where

 H3R8me2a and H3R17me2a decreased in DART4 KD. The decreased regions were

 identified using the same conditions as in (c). (E) Venn diagram showing the overlap of

 the regions where H3R8me2a and H4R3me2a were decreased in DART1 KD. The

 decreased regions were identified using the same conditions as in (C) and Fig. 3F, G.

 (F) Western blotting using anti-FLAG, anti-βTub anti-H3R17me2a antibodies on total

 lysates of FLAG-Rhino inducible OSCs. Dox was added at a final concentration of 5

 ng/mL. (G) Western blotting using anti-ADMA, anti-HP1a, anti-FLAG antibodies on

 immunoprecipitated nuclear lysates of FLAG-Rhino inducible OSCs. Dox was added at

 a final concentration of 5 ng/mL. Anti-H3R17me2a antibody was used as a positive

 control for the reactivity of the anti-ADMA antibody. (H) Scatter plot comparing gene

 expression levels between OSCs and ovaries. Each dot represents a single gene plotted

 by its TPM values (log10). RNA-seq data for OSCs and ovaries were obtained from

 DRR05098823 and SRR11879526, respectively. (I) Comparison of qRT–PCR results

 for DART1, DART4, DART8, and CG17726 in OSCs under EGFP knockdown conditions

(n = 3). (J) qRT–PCR analysis of DART1, DART4, DART8, and CG17726 in OSCs under each gene knockdown condition (n = 3).

Supplementary Figure 4. ADMA-histones localize at the ends of Kdm3 GLKD specific ectopic piRNA clusters

Genome browser snapshot showing the localization of H3R2me2a, H3R8me2a, H3R17me2a, H3R26me2a, and H4R3me2a in piRNA clusters in OSCs and Oregon-R ovary. The gray-shaded areas indicate the localization sites of ADMA-histones in Oregon-R ovary.

Supplementary Figure 5. Behavior of Rhino and ADMA-histones under DART4-GLKD conditions

(A) qRT-PCR for DART4 in ovaries under ctrl-GLKD and DART4-GLKD conditions (n = 3). (B) ChIP-seq heatmap showing the localization of Rhino and H3R17me2a in ctrl-GLKD and DART4-GLKD conditions at the FLAG-Rhino localization sites in OSCs. (C) ChIP-seq heatmap showing the localization of Rhino and H3R17me2a in ctrl-GLKD and DART4-GLKD conditions at Rhino localized regions in WT Oregon-R ovary. (D) Genome snapshot of the indicated ChIP-seq data in piRNA clusters, 42AB, 80F, 38C, and 102F. (E) Genome browser snapshot of the indicated ChIP-seq data in representative ectopic piRNA clusters appearing in Kdm3 GLKD (R2_bi region, R4_jing region and R7_opa/laf region). The gray-shaded areas indicate the localization sites of potential FLAG-Rhino considered to be the ends of the ectopic piRNA cluster in OSCs. (F) Metaplot showing ADMA-histones at the major ectopic piRNA clusters R1−R12 (R5 was excluded because of low Rhino localization in Kdm3 GLKD) that appeared in Kdm3 GLKD. Kdm3 GLKD-dependent ectopic piRNA cluster ends were extracted using the following method.

Among Rhino peaks listed in the MACS2 callpeak files

"GSE203279_MACS2_narrowpeak_Rhino_Ctrl.txt" and "GSE203279_MACS2_

narrowpeak_Rhino_Kdm3GLKD.txt" in the GSE203279 dataset, we first extracted the

Rhino peaks that appeared only in the latter[13]. The peaks closest to the 5′ and 3′ ends of

the R1−R12 regions in Table S4 of the Casier et al. (2023) were designated as the ends

of ectopic piRNA clusters[13].

Supplementary Figure 6. piRNA source loci with H3R17me2a at the ends lose

Rhino localization in DART4 GLKD

(A) Genome snapshot of ChIP-seq results of Rhino reduced regions in DART4 GLKD,

as shown in Fig. 6D. Rhino reduced region in DART4 GLKD with remarkable loss of

Rhino are shown. Black lines indicate boundaries of Rhino reduced regions in DART4

GLKD. The gray-shaded areas indicate the localization sites of potential ADMA-histones

considered to be the ends of Rhino reduced regions in DART4 GLKD. (B) ChIP-seq

profile showing enrichment of the indicated histone modifications in Rhino reduced

regions in DART4 GLKD. ChIP-seq data from WT Oregon-R ovaries were used for each

histone modification. SDMA: symmetric dimethylarginine. (C) qRT-PCR showing the

expression of transposons and genes within Rhino reduced regions in DART4 GLKD (bx,

CG31612, CG14629, and Nf-Yb). Primers are listed in Supplementary Table 1. (D) Egg

hatching rate showing fertility in DART4 GLKD. (E) Proportion of transposon/gene

sequences contained within Rhino reduced regions in DART4 GLKD and annotated all

piRNA clusters. (F) Ratio of piRNA reads mapped to transposons in ctrl-GLKD ovaries[72]

within annotated all piRNA clusters and Rhino reduced regions in DART4 GLKD. The

number of piRNA reads in each category is as follows: Transposons in all clusters:

1,020,456; intragenic regions in all clusters: 156,244; intergenic regions in all clusters:

123,370; transposons in Rhino reduced regions in DART4 GLKD: 2,568; intragenic

regions in Rhino reduced regions in DART4 GLKD: 71,318; intergenic regions in Rhino

reduced regions in DART4 GLKD: 9,907. (G) Box plot showing the lengths of Rhino

localization sites in OSCs, Rhino reduced regions in DART4 GLKD, all piRNA clusters

annotated in proTRAC (as in Fig. 6B), and highly transcribed piRNA clusters (42AB, 38C,

80F, and 102F).

Supplementary Figure 7. Kipferl and Rhino behavior in DART4 GLKD

(A) ChIP-seq profile of Kipferl enrichment in bx, CG14629, and CG31612 DART4 piSL.

(B) Genome snapshot of DART4 piSL in which a significant decrease in Kipferl was

observed in DART4 GLKD. (C) Pie chart showing the number of Rhino peaks in piRNA

clusters, transposons, and DART4 piSL in ctrl GLKD and DART4 GLKD. The numbers

for each are as follows. piRNA clusters in ctrl GLKD: 282; transposons in ctrl GLKD: 114;

DART4 piSL in ctrl GLKD: 60; piRNA clusters in DART4 GLKD: 402; transposons in

DART4 GLKD: 173; DART4 piSL in DART4 GLKD: 30. (D) Boxplot showing Rhino

enrichment in satellite repeats.

Supplementary Figure 8. Control for the evolutionary analysis in Fig. 7

(A) RPKM of each ChIP-seq dataset in the piRNA cluster found in all 8 strains (cluster 8,

evolutionarily conserved), random genomic regions, the whole genome, and DART4

piSL. Genomic regions with RPKM values above 30 in the input were excluded as they

were considered ChIP-seq background noise. For the random genomic regions, 500 bp

regions were randomly selected using Python's random.choice to match the number of

piRNA clusters. The whole genome represents the full length of chromosomes 2L, 2R,

3L, 3R, 4, and X. t-test was used for statistical hypothesis testing between clusters found in all 8 strains (cluster 8, evolutionarily conserved) and other indicated genomic regions. Random genomic regions are shown in Supplementary Table 10, and the p-values from the t-test are shown in Supplementary Table 11. (B) Same analysis of Fig. 7A and Supplementary Fig. 8A for each ChIP-seq dataset (Oregon-R) in the piRNA cluster dataset with eight levels of evolutionary novelty in which piRNA reads were detected in Oregon-R strain. (C) Same analysis of Fig. 7A for each ChIP-seq dataset (ctrl GLKD and DART4 GLKD) in the piRNA cluster dataset with eight levels of evolutionary novelty in which piRNA reads were detected in ctrl GLKD strain.

Supplementary Figure 9. Validation of antibody specificity

(A) Western blotting on total lysates of FLAG-Rhino-inducible OSCs. The anti-Rhino antibody used in immunofluorescence (Fig. 5A) was used. (B) Heatmap showing the localization of H3R8me2a (antibodies produced by different industries) in OSCs at ChIP-seq peaks using the H3R8me2a antibody (Active Motif). The regions of 2 kb around the H3R8me2a localization sites were plotted based on macs2 callpeak results. (C) Western blotting using anti-H3R17me2a antibody on total lysates of FLAG-Rhino inducible OSCs. (D) Histone peptide array using H4R3me2a antibody. The table shows the top 10 single modification peptides with the highest signal intensity.

Figure 1

Saito et al

A

FLAG (Rhino) | H3K9me3 | DAPI | FLAG / H3K9me3 | Merge

B

Genome coordinates — chr2R:6,186,563-6,590,984 | chr2L:20,065,753-20,263,204

FLAG-Rhino inducible OSCs:
- input [0 - 50]
- FLAG (Dox-) [0 - 50]
- FLAG (Dox+) [0 - 50]
- Rhino (Dox+) [0 - 50]
- H3K9me3 [0 - 20]
- H3K9me2 [0 - 30]
- HP1a [0 - 15]

OregonR ovary:
- input [0 - 50]
- Rhino [0 - 50]
- H3K9me3 [0 - 20]
- H3K9me2 [0 - 30]
- HP1a [0 - 15]

Mappability [0 - 1.00]
piRNA clusters
Transposons
Genes

38C1 | 38C2

piRNA cluster, 42AB (Chr 2R) | piRNA cluster, 38C (Chr 2L)

C Rhino peaks in ovary (2629 regions) — Ovary | OSCs

D piRNA clusters in ovary (Rhino peaks larger than 1.5kb) — Ovary / OSCs

E FLAG-Rhino, H3K9me3, H3K9me2 Venn: 187, 622, 3097, 14, 1090, 5718

F FLAG-Rhino peaks (828 regions) — FLAG (OSCs) | H3K9me3 | H3K9me2

G H3K9me3 peaks (4696 regions) — H3K9me3 | FLAG-Rhino | HP1a

A FLAG (Rhino) | DAPI | Merge

EGFP KD
HP1a KD
DART1 KD
DART4 KD
DART8 KD
CG17726 KD

B
n.s.
n.s.
***
***
***

Area FLAG-Rhino foci (micron²)

EGFP KD | HP1a KD | DART1 KD | DART4 KD | DART8 KD | CG17726 KD

C
FLAG-Rhino | H3R17me2a
EGFPKD
DART1KD
DART4KD

H3R2me2a | H3R26me2a

H3R8me2a | H4R3me2a

-1.0 | 1.0Kb

D FLAG-Rhino reduced peaks in DART4KD (207 regions)
FLAG-Rhino | H3R17me2a | H3R8me2a
KD: EGFP DART4 | EGFP DART4 | EGFP DART4

Normalized reads (CPM)

-1Kb +1Kb -1Kb +1Kb -1Kb +1Kb -1Kb +1Kb -1Kb +1Kb -1Kb +1Kb

E
*
†

Rhino ChIP-qPCR (% of input)

EGFPKD | DART1KD | DART4KD

OSC FLAG-Rhino point
Chr4:1051240

F
527 | 48 | 605

H4R3me2a loss
in DART1 KD

H3R17me2a loss
in DART4 KD

G
90 | 25 | 182

FLAG-Rhino decrease
in DART1 KD

FLAG-Rhino decrease
in DART4 KD

A

B

Rhino domains in ovary (2629 regions)

C

piRNA clusters in ovary
(Rhino peaks larger than 1.5kb)

A

B

C

transposon enrichment?

transcription of nearby genes?

accumulation of H3K9me2?

D

E

A

A Kipferl — ctrl GLKD, DART 4GLKD; ChIP enrichment (CPM); -7.5, 5', 3', 7.5Kb

C ctrl GLKD, DART4 GLKD; piRNA clusters, transposons, DART4 piSL

D satellite repeats; Rhino ChIP enrichment (RPM); ctrl GLKD, DART4 GLKD

B (GLKD); coordinates; input ctrl/DART4; Rhino ctrl/DART4; Kipferl ctrl/DART4; piRNA ctrl/DART4; Mappability; DART4 piSL; annotated clusters; Transposons; Genes

DART4 piSL bx (ChrX)

DART4 piSL CG14629 (ChrX)

DART4 piSL CG31612 (Chr2L)

A