# Mitochondrial dysfunction drives a neuronal exhaustion phenotype in methylmalonic aciduria

**Matthew C.S. Denley[1], Monique S. Straub[1], Giulio Marcionelli[1], Miriam A. Güra[1], David Penton Ribas[2], Igor Delvendahl[3], Martin Poms[4], Beata Vekeriotaite[6], Federica Conte[5], Ferdinand von Meyenn[6], D. Sean Froese[1 ✉], Matthias R. Baumgartner[1 ✉]**

[1]Division of Metabolism and Children's Research Center, University Children's Hospital Zurich, University of Zurich, Zurich, Switzerland CH-8032.
[2]Electrophysiology Core Facility, University of Zurich, Zurich, Switzerland CH-8057.
[3] Department of Molecular Life Sciences, University of Zurich, Zurich, Switzerland CH-8057.
[4] Clinical Chemistry and Biochemistry and Children's Research Center, University Children's Hospital Zurich, University of Zurich, Zurich, Switzerland CH-8032.
[5] Department of Neurology, Donders Institute for Brain, Cognition and Behavior, Radboud University Medical Center, 6525 GA Nijmegen, Netherlands.
[6] Laboratory of Nutrition and Metabolic Epigenetics, Institute for Food, Nutrition and Health, Department of Health Sciences and Technology, ETH Zurich, Zurich, Switzerland CH-8603.

✉e-mail: Sean.Froese@kispi.uzh.ch; Matthias.Baumgartner@kispi.uzh.ch

**Methylmalonic aciduria (MMA) is an inborn error of metabolism resulting in loss of function of the enzyme methylmalonyl-CoA mutase (MMUT). Despite acute and persistent neurological symptoms, the pathogenesis of MMA in the central nervous system is poorly understood, which has contributed to a dearth of effective brain specific treatments. Here we utilised patient-derived induced pluripotent stem cells and *in vitro* differentiation to generate a novel human neuronal model of MMA. We reveal strong evidence of mitochondrial dysfunction caused by deficiency of MMUT in patient neurons. By employing patch-clamp electrophysiology, targeted metabolomics, and bulk transcriptomics, we further expose an altered state of excitability, which we suggest may be connected to metabolic rewiring which is exacerbated by application of 2-dimethyloxoglutrate. Our work provides first evidence of mitochondrial driven neuronal dysfunction in MMA, which through our comprehensive characterisation of this paradigmatic model, enables first steps to identifying effective therapies.**

Mitochondrial diseases are a diverse group of genetic disorders that lead to dysfunctional mitochondria and energetic depletion. Pathogenic variants in approximately 1136 genes which encode mitochondrial-resident proteins[1], found on both nuclear and mitochondrial DNA, may cause mitochondrial diseases. Together, 23 per 100,000 people harbour pathogenic variants that have or will convey a mitochondrial disease[2]. However, the inherent clinical and genetic heterogeneity of mitochondrial diseases make them difficult to diagnose, hard to treat, and complex to study[3,4]. Nevertheless, mitochondrial diseases affecting components of oxidative phosphorylation (OXPHOS) or proteins proximal to the tricarboxylic acid cycle (TCA cycle) often report overlapping symptoms[5,6], particularly within the central nervous system (CNS)[7,8].

Isolated methylmalonic aciduria (MMA) is an archetypical mitochondrial disease caused by pathogenic variants leading to the absence or deficiency of the nuclear-encoded mitochondrial matrix-residing enzyme methylmalonyl-CoA mutase (MMUT; MIM #251000)[9]. It is typified by accumulation of methylmalonic acid and related compounds in blood and urine; however, increased levels of these disease-related compounds are likely not sufficient to explain overall clinical presentation and disease progression[10,11]. Rather, reduced mitochondrial function and energy production may also be important contributors to disease progression[10,12–15]. Mitochondrial structural changes and dysfunction have been widely

described in MMA[12,16,17]. Previous liver and kidney tissue models have demonstrated MMUT-
deficiency causes enlarged mitochondria and altered OXPHOS activity[16,17]. The consequences
of this appear to be impaired mitophagy[12] and  shifted anaplerosis into the TCA cycle[10,18].
MMUT-KO cells and MMA patient fibroblasts have demonstrated a preference for reductive
TCA cycle pathway, which was rescued by supplying cells with 2-dimethyloxoglutarate, an
analogue of oxoglutarate[10]. However, neither mitochondrial nor metabolic response to MMUT-
deficiency in the CNS have been investigated due to the lack of an adequate model.
In the CNS, glutamine and glutamate have tissue-specific functions as stimulatory
neuroactive molecules[19]. Additionally, mitochondria preferentially utilise OXPHOS for
synaptic transmission and maintaining membrane potential through Na/K-ATPases[20–23].
Despite the clear role of neurological dysfunction in MMA and the long-term symptoms (i.e.
metabolic stroke, epilepsy, optic neuropathy) invoked, no appropriate models have been
employed to investigate this aspect of the disease. Such a shortfall is important because current
treatments of MMA are symptomatic and do not protect against long-term CNS sequelae.
Recent advances of patient-derived induced pluripotent stem cell (iPSC)-models offer an
invaluable route to model neuronal dysfunction in MMA, and provide valuable insights into
CNS dysfunction.
In this study, we generated iPSC-derived neuronal models of MMA from variant-
matched patient fibroblasts. We demonstrate that MMUT-deficient neurons have perturbed
mitochondrial networks and reduced mitochondrial function. These appear to drive action
potential attenuation in patient-derived neurons, potentially through reduced sodium currents.
To tie metabolism to cellular function, metabolomic and transcriptomic exploration suggest
altered glutamine processing, which is exacerbated by application of 2-dimethyloxoglutarate.
Altogether, our results demonstrate that mitochondrial dysfunction may lead to a neuronal
exhaustion-like phenotype, which may be an important component of neurological dysfunction
in MMA.
77
**Generation of MMUT-deficient iPSC-derived neuronal lines**
Deficiency of MMUT causes MMA leading to biochemical alterations which drive disease
pathology (Fig. 1a). Using Sendai virus based OKSM (OCT4, KLF4, SOX2, MYC)
reprogramming, we generated iPSC clones from two unrelated individuals, each homozygous
for the pathogenic variant MMUT-p.(Asn219Tyr) (AV & KK, n = 2), and a wildtype control
individual (BJ, n = 1) (Fig. 1b and Extended Data Fig. 1a). Isolated clones from all three iPSC
lines did not retain Sendai virus markers used in ectopic pluripotency induction (Extended Data
Fig. 1b). These cell lines were characterised alongside a commercially available control (ND2,
n = 1). All four iPSC lines expressed the endogenous pluripotency markers: KLF-4, SOX2,
OCT4 and NANOG, which were not found in the progenitor fibroblasts (Extended Data Fig.
1c). We demonstrated expression of the pluripotency markers NANOG and SSEA-4, and pan-
stem cell marker SOX2, as well as the proliferative marker Ki67 (Fig. 1b), and could generate
the three germ layers in all four cell lines (Fig. 1c). Comparative genomic hybridization
indicated that induction of pluripotency did not generate *de novo* copy number variations
between fibroblasts and their daughter iPSC clones (Supplementary Data Fig. 1).
To evaluate whether the generated iPSC clones demonstrated MMUT deficiency, we
examined MMUT protein expression by immunocytochemistry (Fig. 1d) and Western blotting
analysis (Fig. 1e). Using a MMUT specific antibody (Extended data Fig. 1d and 1e), we found
reduced MMUT protein in lysates of patient iPSCs (Fig. 1e), consistent with reduced enzymatic
activity (2.58% of control), as demonstrated by an indirect activity assay which monitors
propionate incorporation via MMUT into the TCA cycle[24] (Fig. 1f).

**Figure 1 | Generation of iPSC-derived MMUT-deficient neurons from individuals with methylmalonic aciduria**. **a**, Schematic of the affected metabolic pathway in MMUT deficiency. **b,** Brightfield and epifluorescent acquisitions of pluripotency (NANOG, SSEA4, SOX2) and proliferation (Ki67) markers in representative iPSC cultures from control and patient lines. Scale, 50 μm. **c**, Ectoderm lineage indicated by beta-tubulin III (TUJ1) and nestin (NES) positive cells after 12 DIV. Endoderm lineage indicated by anti-α-Fetoprotein (AFP) positive cells after 4 days *in vitro* (DIV). Mesoderm lineage indicated by alpha-smooth muscle actin (SMA) and brachyury positive cells after 4 DIV. Scale, 10 μm. **d**, Comparison of the anti-MMUT (green) staining pattern from representative patient fibroblast-derived iPSCs compared to controls. For reference, anti-TOMM20, a mitochondrial protein, is also shown (red). Scale, 50 μm. **e**, Western blotting analysis of derived iPSCs using anti-MMUT. MMUT is anticipated at 83 kDa. The loading control ACTB is anticipated at 42 kDa. Uncropped membranes are available in Supplementary Data Fig. 2a. **f**, Propionate incorporation of two BJ wildtype iPSC sub-clones and two separate patient cell lines. Data points represent independent measurements taken from 3 separate cultures per cell line. Mean control iPSC incorporation values were 22.71 and 24.36 pmol/mg.protein/16h. Patient iPSC incorporation values were 0.77 pmol/mg.protein/16h in AV and 0.44 pmol/mg.protein/16h in KK. P-value was tested using Mann-Whitney U-test (**, <0.01).

**Figure 2 | Derivation of cortical neurons from iPSCs affected by methylmalonic aciduria**. **a**, Schematic of the 2D neuron differentiation protocol used in this article. Created using BioRender **b**, Staining for SSEA4 and SOX2 at day 0 for pluripotent stem cells (iPSC). Staining for Nestin (NES) and SOX2 at day 13 for neuroepithelium. Staining for PAX6 and SOX2 at day 21 for neural stem cells (NSCs). Staining for TUBB3 and EOMES at day 40 for cortical progenitors. Staining for TUBB3/TBR1 and MAP2/NeuN at day-50 signal postmitotic deep layer cortical neurons and pan-neuronal molecular markers in cultures, respectively. Scale: iPSCs and neurons have 10 µm bars. Neuroepithelium, NSC, and NPCs have 50 µm bars. **c**, Real-time RT-qPCR of SOX2, PAX6, and EOMES in iPSCs at d0, neuroepithelium d13, and NSCs d21. Datapoints are representative of at least 3 independent experiments and error is reported as standard deviation. Expression values are relative to measurements at d0 for each cell line. **d,** Immunocytochemistry of patient and control NSC and neurons. Scale, 10 µm. **e**, Western blotting analysis of neurons using anti-MMUT. MMUT is anticipated at 83 kDa. The loading control ACTB is anticipated at 42 kDa. Uncropped membranes are available in Supplementary Data Fig. 2b.

To assess if MMUT deficiency affects neuronal development and function, we employed a 2D
differentiation protocol that utilises dual-SMAD inhibition to generate dorsal forebrain-
enriched neuronal progenitors[25,26] (Fig. 2a). For all control and patient-derived iPSCs, we
identified  successful  generation  of  neuroectodermal  Nestin-  and  SOX2-positive
neuroepithelium which progressively differentiated from a population including SOX2- and
PAX6-positive neural stem cells (NSC) to TUBB3 and EOMES-positive neural progenitor
cells and finally TUBB3, TBR1, MAP2 and NeuN-positive neurons (Fig. 2b). RT-qPCR
revealed that patient cells express *SOX2*, *PAX6*, and *EOMES* at days 13 and 21 *in vitro*
comparably to control cells when expression is normalized to day 0 (Fig. 2c). This suggests
they adopt cortical NSC identity at the same rate as control cell lines. Immunocytochemistry
(Fig. 2d) and Western blotting analysis (Fig. 2e) indicate MMUT protein expression is retained
in NSC and neurons (Fig. 2d), however patient-derived neurons show reduced MMUT protein
levels (Fig. 2e).
114
**Patient neurons have mitochondrial dysfunction and disrupted networks**
Translocase of the outer mitochondrial membrane complex subunit 20 (TOMM20) is an
essential mitochondrial protein often used as a proxy for mitochondrial abundance and shape[27].
Immunocytochemistry demonstrated altered TOMM20 staining patterns in patient-derived
neurons (Extended Data Fig. 2a), suggestive of disrupted mitochondrial networks. This was
consistently observed in neurons derived from both patients, but not controls and was related
to the extent to which MMUT could be detected within individual neurons (Fig. 3a and
Extended Data Fig. 2b). Investigation of spatial mitochondrial characteristics of patient
neurons using TOMM20 staining revealed reduced mitochondrial volume and diameter but not
sphericity (Extended Data Fig. 2c,d). Further, disrupted mitochondrial networks in patient
neurons appeared to impair mitochondrial response to OXPHOS insult, as demonstrated by
comparable MitoSOX fluorescence at baseline but significantly reduced intensity after
treatment with complex I inhibitor rotenone (Extended Data Fig. 2e). In patient derived cells,
we identified two apparent populations, those expressing detectable MMUT (MMUT+) which
overlapped with TOMM20, and those without (MMUT-) (Fig. 3a,b). To investigate the
functional capacity of patient MMUT+ and MMUT- mitochondria, we used detectable
fluorescence of tetramethylrhodamine-methyl-ester (TMRM) as a measure of membrane
potential. We found a reduced TMRM signal in patient derived neurons, a difference that was
exacerbated in those cells with no detectable MMUT (Fig. 3c,d). We hypothesized that the
lower TMRM signal denoted lower electron transport chain function, which we tested by
inhibiting complex I with rotenone. We found reduced TMRM fluorescence in rotenone treated
MMUT+ cells, less so in patient MMUT-, but not in control (Fig. 3d), suggesting that MMUT
deficient mitochondria have a reduced proton motive force leaving them less able to react to
events that would dissipate this potential. We hypothesized that reduced membrane potential
could trigger programmed cell death. This was supported by patient-derived neurons having
increased proportion of pro-apoptotic cleaved Caspase 3 (Fig. 3e) as well as DAPI nuclei
staining patterns consistent with apoptotic cells (Fig. 3f and Extended Data Fig. 2f).

**Figure 3 | Patient neurons show mitochondrial dysfunction that overlaps with loss of MMUT protein a,** Left and middle: MMUT+ signal (green) and TOMM20+ mitochondrial signal (red). Right: TOMM20+ signal (magenta) and colocalization with MMUT (white). Orthogonal representative slices are also shown. Scale is 10 μm. **b,** Pearson correlation coefficient of TOMM20+ regions of interest (ROIs). One data point represents a Pearson's coefficient from one ROI. 3-5 ROIs per image, total images used from control n = 6 (18 total ROIs) and from patient n = 11 (50 total ROIs). P-value determined by one-way ANOVA with multiple testing (ns, >0.05; ****, <0.0001). **c,** Neuronal mitochondria stained with MMUT and TMRM (both grey). Scale is 10 μm. **d,** TMRM+ ROIs selected from non-overlapping confocal images. Cumulative data is representative of N = 6 in control without rotenone, n = 4 in control with rotenone, n = 8 in patient without rotenone, n = 9 in patient with rotenone. One datapoint represents one background/size-corrected ROI. P-value determined by Kruskal-Wallis with multiple testing (ns, >0.05; *, <0.05; ****, <0.0001). **e,** immunoblot of MMUT, caspase-3 (CASP3), its cleaved product, and beta-actin (ACTB) in neurons. Each column represents an independent lysate. Uncropped membranes are available in Supplementary Data Fig. 2d. **f,** Quantification of DAPI+ apoptotic cells determined from representative, non-overlapping, epifluorescent images acquired using a 63x objective. Cumulative data is representative of n = 43 (776 cells) in control and n = 87 (1317 cells) in patient images. Significance is assessed by Mann-Whitney test (****, <0.0001). Dunn's multiple comparisons test used in b, d.

**Electrophysiological assessment reveals an exhaustion phenotype in patient-derived neurons**

In order to assess generated neurons' functional properties, we performed whole-cell analysis on patient (n = 132, AV & KK) and control (n = 94, BJ & ND2) neurons (Supplementary Data Table 1). Similar to previous reports[28], principle component analysis (PCA) of all 31 extracted features from step current injection protocols (Supplementary Data Table 2) revealed three distinct clusters (Extended Data Fig. 3a), which were further discriminated by utilizing 21 primary variables (Fig. 4a and Extended Data Fig. 3b). These clusters could be best represented as non-action potential (AP) firing (Type 1, T1), single-AP firing (T2), and multiple AP-firing (T3) neurons (Fig. 4a,b and Extended Data Fig. 3a). They were not separated by genotype (patient vs control) (Extended Data Fig. 3b), but rather by variation within inherent electrophysiological properties, such as resting membrane potential, input resistance, capacitance, number of APs, and AP threshold (Extended Data Fig. 3c). Unguided hierarchical clustering of 21 primary variables separated assessed neurons by T1, T2 or T3, but not by genotype (Extended data Fig. 3d). However, we observed an apparent decrease in T3 and corresponding increase in T2 in patient-derived neurons (Fig. 4c), suggesting that generating successive APs may be corrupted in patient neurons.

To explore this further, we performed correlation analysis on 21 primary variables in T3 neurons derived from patients (n = 50) and controls (n = 52) (Fig. 4d). For all cells, we identified anticipated correlations between variables, such as AP threshold and AP full width at half maximum (FWHM, i.e. time between slope rise and fall at half-max amplitude) with AP amplitude (negative correlations), as well as AP amplitude with AP number and resting membrane potential with AP FWHM (positive correlations) (Fig. 4d and Extended Data Fig. 4a,b). Striking, however, was that variables expected to correlate with AP attenuation, and found to do so in control neurons (e.g. AP amplitude, AP threshold and FWHM), did not correlate in patient neurons (Fig. 4d and Extended Data Fig. 4c). Closer inspection of the reduction of AP amplitude following multiple APs (measured as the attenuation ratio), revealed a significantly stronger loss of AP amplitude and hence enhanced attenuation in patient neurons (Fig. 4e). This is indicative of a mitochondrial exhaustion-like phenotype [29,30].

To link the loss of AP amplitude over multiple firings and mitochondrial dysfunction, we further assessed attenuation in selected cells whose mitochondrial morphology was visualized using MitoTracker (Extended Data Fig. 4d). Consistent with previous data (Extended Data Fig. 2a-d), all control cells appeared to have intact mitochondrial networks, whereas patient derived cells had either intact or disrupted mitochondrial networks (Extended Data Fig. 4d). Patient cells with intact mitochondrial networks had an attenuation ratio similar to control neurons, whereas neurons with disrupted mitochondrial networks had increased attenuation (Extended Data Fig. 4d), suggesting an association between AP exhaustion and mitochondrial dysfunction.

Depolarisations generating higher AP amplitudes are achieved by influxes of cations; the velocity of these influxes can be calculated through the first derivative of the membrane potential slope over time (dVdt, Fig. 4f). dVdt ratios in patient-derived neurons were decreased compared to controls (Extended Data Fig. 4e), which was driven by the depolarisation (reported as maximum) and not repolarisation (potassium efflux, reported as minimum) phase (Fig. 4g). In control neurons, the maximum dVdt positively correlated with the attenuation ratio, a relationship which was lost in patient-derived neurons (Extended Data Fig. 4f). Similarly, the positive correlation of dVdt with AP number and with AP amplitude found in control neurons was lost or reduced in patient-derived cells (Fig. 4d and Extended Data Fig. 4g). These findings suggest that patient-derived cells cannot sustain sufficient current densities to maintain action potential firing, which may lead to the observed exhaustion-like phenotype.

192   To test the hypothesis that reduced sodium currents drive these observations, we
193 performed additional voltage-clamp recordings on patient (n = 92) and control (n = 89)
194 neurons. The selected neurons contained the same patient T3 proportional decrease (Extended
195 Data Fig. 5a), increased attenuation and decreased dVdt ratio as the overall population
196 (Extended Data Fig. 5b). We found substantially reduced sodium current densities in patient-
197 derived neurons when compared to control, with peak densities reduced 5-fold in patient-
198 derived neurons (Fig. 4i,j; Extended Data Fig. 5c-e).

**Figure 4 | Patient neurons show action potential exhaustion driven by reduced sodium currents. a**, First two dimensions (44% variance explained) of a principal component analysis of 21 measured features from patient (n = 132) and control (n = 94) neurons. Greyscale represents 3 identified neuronal types, type 1 (light grey, T1, n = 60), type 2 (dark grey, T2, n = 64), and type 3 (black, T3, n = 102). **b**, Current-clamp, representative traces from T1, T2, or T3 neurons in response to depolarizing current injection. **c**, Proportional generation of classified neurons of control (ND2, BJ) and patient (AV, KK). **d**, Correlation matrix of 21 features in patient (bottom-left, orange) and control (top-right, blue) neurons. Positive correlations indicated by red and negative by blue. Features with significance P = <0.01 are displayed on the matrix, values above are blank. **e**, Top, overlayed traces from T3 control and patient neurons. Bottom, attenuation ratio from 102 measured T3 neurons. **f**, Phase plots of 2 overlayed single action potentials from representative T3 control and patient neurons. **g**, Depolarisation (left, maximum dV/dt) and repolarisation (right, minimum dV/dt) velocities. **h**, Voltage-clamp, representative overlay of currents evoked by step to -30 or -20 mV in control and patient neurons, respectively. Inset is an enlarged section of the initial segment. **i**, Peak control (n = 67) and patient (n = 67) sodium current densities. **j**, Peak and steady control (n = 67) and patient (n = 67) potassium current densities. In i-j, current densities are normalised to capacitance, and variation is SD and present as shaded area. In e and g, datapoints represent one neuron, data are pooled into control (2 independent lines) and patient (2 independent lines), significance is by t-test and reported with p-value adjustment using Holm.

**Rewired glutamine-glutamate metabolism alters glutamatergic identity**

We next utilised targeted metabolomics to investigate the connection between mitochondrial dysfunction and altered neuronal excitability. MMA related metabolites, including hydroxypropionic acid and propionyl-carnitine were elevated as expected in patient-derived neurons, while lactate tended to be increased with methylmalonic acid and 2-methylcitrate appearing unchanged (Fig. 5a and Extended Data Fig. 6a). Despite our finding of disturbed mitochondrial networks in patient derived neurons, we found no difference in the pools of the anaplerotic intermediates pyruvate, alanine, glutamine and glutamate (Extended Data Fig. 6b) as well as TCA cycle intermediates (Extended Data Fig. 6c) between patient and control-derived neurons. This prompted us to investigate the anaplerotic contribution to the TCA cycle from labelled glutamine (Fig. 5b). Glutamine flows into the TCA cycle via glutamate and 2-oxoglutarate (Fig. 5b). Following labelled glutamine supplementation, we found a high proportion of labelled glutamine, which was reduced in glutamate and 2-oxoglutarate, but found no difference between fractional labelling (Extended Data Fig. 6d) nor pool levels (Extended Data Fig. 6e) of all three intermediates between patient and control-derived neurons. We also found no difference in the fractional incorporation nor pool levels of measured TCA intermediates (Extended Data Fig. 6f,g). However, we did notice reduced fractional incorporation, but not pool levels, of aspartate (Fig. 5c). Glutamine-derived aspartate can be produced via oxaloacetate by cytosolic or mitochondrial aminotransferases, whereby the direction of the TCA cycle (oxidative or reductive) influences the number of labelled carbons (Fig. 5b). We found the ratio of M+5 to M+4 citrate, and M+3 to M+4 malate (M+$x$ refers to the number of carbon isotopes that are present), to be reduced in patient-derived neurons (Fig. 5d), suggestive of anaplerotic rewiring into the TCA cycle in these cells.

We also investigated the pool sizes of glycolytic intermediates, which follows report of increased hexoses in primary patient fibroblasts[10]. Using a targeted approach to measure levels of specific detectable sugar phosphates[31], we identified elevated galactose-1-phosphate, mannose-6-phosphate, and glucosamine-6-phosphate in patient-derived neurons compared to controls (Extended Data Fig. 7a). We found reduced *N*-acetyl-glucosamine-6-phosphate and *N*-acetyl-mannosamine-6-phosphate (Extended Data Fig. 7b) and elevate nucleotide sugars GDP-fucose and GDP-mannose (Extended Data Fig. 7c), but no change in other glucose or fructose phosphate derivatives nor pentose phosphate derivates (Extended Data Fig. 7a,d and Supplementary Data Table 3). Our finding of elevated galactose-1-phosphate and mannose-6-phosphate suggest glycolytic flux is increased. Further investigation is required to determine a link between increased glycolytic metabolites and rewired TCA cycle metabolism, as this phenomena has been reported previously[10].

Based on the above changes, we attempted to rescue the phenotype in patient-derived neurons using dimethyl-2-oxoglutarate (DM-2OG) a membrane permeable analogue the TCA cycle intermediate 2-oxoglutarate (Fig. 5e). Application of DM-2OG increased some but not all pools of TCA cycle intermediates measured, including glutamate and glutamine (Extended Data Fig. 8a-b), and resulted in amelioration of the difference between pool levels of hydroxypropionic acid, propionyl-carnitine, and lactate (Fig. 5f, Extended Data Fig. 8c) as well as fractional contribution into aspartate of control and patient-derived neurons and normalisation of TCA cycle flux (Fig. 5g-h). DM-2OG supplementation increased selected hexose phosphates but patient galactose-1-phosphate, mannose-6-phosphate, and glucosamine-6-phosphate remained significantly increased compared to treated control neurons (Extended Data Fig. 8d). However, DM-2OG treatment reduced the difference between patient and control *N*-acetyl-glucosamine-6-phosphate and *N*-acetyl-mannosamine-6-phosphate (Extended Data Fig. 8e).

**Figure 5 | Metabolomics reveal dysregulated glutamate and glutamine neuronal metabolism. a,** Box plots of absolute ion abundance of disease-related metabolites. P value is reported as one-way ANOVA for hydroxypropionic acid, and unpaired t-test for propionyl-carnitine, and lactate. **b**, Schematic of labelled [13]C-glutamine metabolism into the TCA cycle. Labelled carbons = filled black circles. **c** Stacked bar chart of contribution of labelled glutamine carbons to aspartate (left) and box plot of absolute ion abundance of aspartate pool (right). P value calculated via unpaired t-test. **d**, Box plot of ratio between M+5 to M+4 labelled citrate fractions (left) and box plot of ratio between M+4 to M+3 labelled malate or aspartate fractions (right). P value is reported as unpaired t test. **e**, Representative schematic for the entry of dimethyl-2-oxoglutarate (DM-2OG) into the TCA cycle and selected anaplerotic entry points. **f**, Box plot of absolute ion abundance of propionyl-carnitine and lactate in DM-2OG treated neurons. P value is reported as unpaired t test. **g**, Box plot of total fractional contribution of carbons to untreated and DM-2OG treated neuronal abundance of aspartate. P value is reported as Mann-Whitney (left). **h,** Stacked bar chart of contribution of labelled carbons to untreated and DM-2OG treated neuronal abundance of malate (left) and citrate (right). In **a, c, d, f, g, h,** error = min/max (box plot) or SD (fractional bar plot). Each datapoint represents 3 technical replicates from one independent sample. Genotype conditions are pooled into control (blue) and patient (no colour). In significance tests: ns = >0.05, * = <0.05, *** = <0.001, **** = <0.0001.

**TCA cycle rewiring impacts neuronal excitability at the synaptic level.**

DM-2OG altered the neuronal levels of neurotransmitter amines, such as glutamate, leading us to investigate the effect of DM-2OG supplementation and subsequent metabolic rewiring on electrophysiological properties. We performed current-clamp recordings on 36 neurons with supplementation of DM-2OG between 0.1 – 6.0 mM (Extended Data Fig. 9a). We found that higher doses abolished AP firing capacity (Extended Data Fig. 9a), which may be linked to increasing hyperpolarisation and decreasing capacitance (Extended Data Fig. 9b-c). Voltage-clamp recordings on control neurons demonstrated that 1 mM and 6 mM DM-2OG reduced sodium current densities to levels that were similar to untreated patient-derived neurons (Fig. 6a,b). At lower doses, control but not patient-derived neurons were still able to produce APs and maintain sodium current densities (Fig. 6a-b and Extended Data Fig. 9a). We hypothesized that patient metabolic rewiring altered neuronal function at the synaptic level. We investigated the packaging of glutamate in the synapse through *SLC17A7*, which encodes the vesicular glutamate transporter 1 (VGLUT1). We found a greater number of VGLUT1+ foci in patient neurons (Extended Data Fig. 9d,e), suggesting a predisposition towards glutamate excitotoxicity that could contribute to the exhaustion phenotype, and, more generally, to the neurological phenotype. Consequently, we recorded spontaneous excitatory postsynaptic currents of untreated patient (n = 46) and control (n = 48) as well as 0.1mM DM-2OG treated patient (n = 7) and control (n = 8) neuronal networks. We observed an increased charge of synaptic events in patient-derived neurons (Fig. 6c,d). However, we did not observe difference in event amplitude, decay kinetics, or frequency (Extended Data Fig. 9f-h). The event duration (charge) may be related to glutamate receptor subunits, which combine to form functional neuronal glutamate receptors[32].

Given the electrophysiological and metabolic changes in patient-derived neurons, we performed bulk transcriptomics on control and patient neurons. Analysis of the neuronal transcriptome via PCA revealed successful separation of controls and patients based on inherent variance (Extended Data Fig. 10a). Using over-representation analysis, we found gene networks involved in "glutamatergic synapse", "ion channel activity", and "chemical synaptic transmission" to be differentially expressed (Fig. 6e and Extended Data Fig. 10b-c). Further analysis revealed glutamate packaging machinery (*SLC17A7*), and glutamate receptors, such as *GRM7*, *GRIN3A*, *GRIA3*, *GRIA4*, to have altered expression in patient-derived neurons (Fig. 6f and Extended Data Fig. 10d). We also observed several ion channel genes, including sodium (*SCN9A*), calcium (*CACNA1H*), and potassium (*KCNH3*, *KCNN1*, *KCNIP3*) channels, to be differentially expressed (Fig. 6f). Moreover, GAD1/2, which catalyses the production of 4-aminobutanoate (GABA) from glutamate, depending on the supply pool size of glutamate in the cytosol, appeared to be downregulated in patient-derived neurons (Fig. 6f).

To test if subunit composition is altered, we treated control and patient neurons with 0.1 mM DM-2OG and analysed the transcriptome. Comparison of untreated control and DM-2OG supplemented patient-derived neurons revealed the absence of many previously identified significantly differentially regulated synaptic related GO terms (Fig. 6g). Of previously identified glutamate synapse differentially expressed genes, only GRM7, GRIN3A, SCN9A, KCNN1, and GRIA3 were still significant (Fig. 6h). Further investigation on GRIA2 and GRIN1 demonstrated that patient-derived neurons were sensitive to DM-2OG treatment when compared to gene count in control derived neurons (Extended Data Fig. 10e,f). Disease-state metabolic rewiring changes the TCA cycle, an impact that we can replicate with DM-2OG treatment. We demonstrate DM-2OG increases metabolite pools, such as glutamate, and these treatments recapitulate aspects of the exhaustion phenotype.

**Figure 6 | Metabolic rewiring affects glutamatergic synaptic transmission a,** Line graph indicating maximum current density at evoked voltages in control neurons with various DM-2OG treatments. Datapoint represents at least 5 independent samples, error = SD represented by shading. **b,** Line graph indicating maximum current density at evoked voltages in patient neurons with various DM-2OG treatments. Datapoint represents at least 6 independent samples, error = SD represented by shading. **c,** Representative averaged traces of spontaneous synaptic currents from single control (n events = 114) or patient (n events = 36) neurons in voltage-clamp. **d,** Box plot of spontaneous synaptic event charge of individual neurons in voltage-clamp. P value is calculated via mann-whitney **e,** Over-representation analysis of top 10 up and down-regulated GO terms from untreated control (n = 5) and patient (n = 6) derived neurons. Count refers to number of contributing genes. **f,** Volcano plot of differential regulation between untreated control (n = 5) and patient (n = 6) neurons. **g,** Over-representation analysis of top 10 up- and down-regulated GO terms. Count refers to number of contributing genes from untreated control (n = 5) and 0.1 mM DM-2OG treated patient (n = 4) derived neurons. **h,** Volcano plot of differential regulation between untreated control (n = 5) and 0.1mM DM-2OG treated patient (n = 4) neurons. In **c, d,** each datapoint represents one independent sample. Genotypes are control (blue) and patient (orange). In **a, b** center is the mean and error are SD. In **c, e,** colour represents *P* value magnitude. In **f, h,** values are FDR, genes highlighted contribute to a top differentially regulated GO term. Dotted lines indicate Log2 fold change of 1 and -log10 P-value of 0.01. In **d,** box plot is median and min/max.

# Discussion

In this study, we have gained insight into the conserved role of mitochondrial disruption in MMA and uncovered an exhaustion-like phenotype in neurons affected by mitochondrial disease. We described how mitochondrial disruption and reduced sodium current density underlie AP attenuation. MMUT deficiency rewires neuronal metabolism and conferred a greater demand of glutamine and aspartate to the TCA cycle, we also documented reduced current density through modulation of synaptic ion channel dynamics.

The occurrence of neurological symptoms despite treatment highlights the need to explore neuronal pathomechanisms [33–39]. Our findings reinforce previous comprehensive evidence that mitochondrial dysfunction drives pathophysiology in MMA[12,16,17]. These evidence fit well within the current literature that describe reduced mitochondrial respiration and/or mitochondrial ultrastructure alterations in primary patient kidney cells [12]; liver, kidney, heart, and skeletal muscle of patients [13]; Mut$^{-/-}$ mice kidney [17] and liver [16]. Here we demonstrate that mitochondrial dysfunction is also an important component of neuronal pathology. Although electrophysiology has never been assessed in MMA, two separate iPSC-derived models of the primary mitochondrial disease Leigh Syndrome demonstrate that pathogenic variants in the electron transport chain reduced oxygen consumption, decreased current densities, confer strong attenuation, and reduced AP firing [29,30].

Neurons have been shown to have a greater reliance on OXPHOS during AP firing[40,41]. One route to meeting such a demand in the brain is through the oxidation of glutamine[42–45], which is coupled to incremental glucose oxidation[45]. Derangement of OXPHOS puts a greater reliance on glutamine and aspartate metabolism[46]. Our finding of reduced glutamine contribution to aspartate suggests rewiring of the TCA flux[47], consistent with reduced reliance on reductive TCA cycle flux in patient derived neurons. This pattern is consistent with a greater importance on anaplerotic metabolites. Elevated levels of lactate and hexose phosphate sugars suggest glucose flowing into glycolysis may also be rewired to side pathways[48]. Altered dependence of patient neurons on glutamine and aspartate may feasibly lead to differential expression of glutamatergic receptor subunits. Hippocampal neurons rapidly redistribute their glutamate receptors after glutamate and glycine treatment[49,50], indicating that metabolic shift may be strong enough to cause the differential expression of ionotropic glutamate receptors observed in patient neurons. Concurrently this may result in patient neurons' prolonged synaptic events or reduced ability to sustain successive action potentials. *GRIA4*, which is downregulated in patient cells, is the fastest glutamate receptor and its expression is linked to receptor kinetics[51]. GRIA4-containing AMPA receptors also appear responsive to changes in spontaneous synaptic events[52]. Manipulation of disease-state metabolic rewiring with DM-2OG demonstrated a rescue of disease-related metabolites but exacerbated components of the exhaustion phenotype in control-derived neurons. This suggests an association between metabolic rewiring and gene expression. Decreased measurements of *N*-acetyl-glucosamine-6-phosphate and *N*-acetyl-mannosamine-6-phosphate may link changes in the hexosamine biosynthesis pathway to altered transcription, as this pathway contributes posttranslational modifications to intracellular proteins, including transcription factors. Furthermore, the use of DM-2OG and its effect on glutamate, a neurotransmitter, evidenced these effects by abolition of AP firing capacity.

Here, we describe a novel mechanism of mitochondrial disruption generating reduced AP firing in patient-derived cells affected by MMA. Underlying this AP exhaustion was a reduced sodium current density, rewired glutamine metabolism, and differential glutamate receptor transcript expression. Further manipulation of glutamine metabolism reverses some of the metabolic and transcriptomic observations. More work will be required to elucidate the exact mechanism that reduces excitability in patient neurons.

# Online Methods

**Ethics statement, patients and fibroblast lines**

Approval to use patient fibroblasts was granted by the Cantonal Ethics Commission of Zurich (KEK-ZH-Nr. 2014-0211, amendment PB_2020-00053). Both patient-derived fibroblast lines carried the *MMUT* homozygous pathogenic variant c.655A>T (p.Asn219Tyr), confirmed by Sanger sequencing at the DNA level, and corresponding to the $mut^0$ phenotype. In addition to fibroblasts from two affected individuals (AV, female; KK, male), fibroblasts from BJ (ATCC: CRL-2522, male) were used for iPSC generation. An additional control (CI, male; NHLBli003-A, RRID: CVCL_1E78) was acquired from Rutgers University (RUCDR, USA) [53].

**Fibroblast and HEK293 culture, and maintenance**

Cells were cultured at 37°C and 5% CO2 in a humidified incubator. Cells were cultured in Dulbecco's modified eagle medium (DMEM, Gibco, Cat. 31966-047) supplemented with 10% Fetal bovine serum (FBS, Gibco, Cat. 102070-106) and 1% Anti-anti (100x, Antibiotic-antimycotic, Gibco, Cat. 15240-062). Passaging was performed once cells reached 80% confluency, using 0.05% Trypsin (1x, Trypsin-EDTA, Gibco, Cat. 25300-054) and washed with Dulbecco's Phosphate Buffered Saline (DPBS, Gibco, Cat. 14190-250). Cells were maintained frozen in liquid nitrogen in cryopreservation medium (90% FBS (as before) and 10% DMSO (Sigma, Cat. D2438-50ML)).

**iPSC culture, differentiation, and maintenance**

All cells were cultured at 37°C and 5% $CO_2$ in a humidified incubator. The generation of iPSCs from fibroblasts was performed as previously described [54]. Briefly, the transduction was performed using the CytoTune®-iPS 2.0 Sendai Reprogramming Kit (Invitrogen, Cat. A16518) according to the manufacturer's instructions. Transduction was performed on fibroblasts with passage numbers lower than five. Cells were plated at a density of $1x10^5$ cells/well in a 12-well plate. Non-integrating Sendai virus was added to the fibroblast medium at a MOI (ratio virus to cells) of 5 for KOS (Klf4, Oct4, Sox2) and c-Myc, and MOI 3 for Klf4. 8000 transduced fibroblasts were plated on inactivated mouse embryonic fibroblasts (MEFs). Derived iPSCs colonies were transferred to feeder-free conditions using 1mg/ml Collagenase type IV solution (Gibco, Cat. 17104019).

Feeder-free iPSC colonies were maintained as previously described [55] with minor modifications. Between culturing, cells were maintained frozen in liquid nitrogen in cryopreservation medium (90% KnockOut™ Serum Replacement, Gibco, Cat. 10828028; 10% DMSO, Sigma, Cat. D2438). Cells were cultured on 0.17 mg/well (6-well) or 0.083 mg/well (12-well) BME matrix (Cultrex® Reduced Growth Factor Basement Membrane Extract, PathClear®, Cat. 3433-010-01) in Essential 8 medium (Gibco, Cat. A1517001) supplemented with 100 U/mL Penicillin-Streptomycin (10'000 U/mL, Gibco, Cat. 15140122). Passaging was performed once colonies reached 80% confluency using EZ-LiFT Stem Cell Passaging Reagent per manufactures instructions (Sigma-Aldrich, SCM139) after which 10 µL/mL RevitaCell™ Supplement (100X, Gibco, Cat. A2644501) was applied to passaged iPSC colonies. For neural induction, single-cell passaging of iPSCs was performed using 0.5 mL or 1 mL ACCUTASE™ Cell detachment solution (1X, StemCellTechnologies, Cat. 07920) for 12-well and 6-well plates, respectively, and 2 µL/mL from 5 mM ROCK-inhibitor Y-27632 (Dihydrochloride; Sigma-Aldrich, Cat. SCM075) was applied for single-cell passaging. Cells were inspected daily using the Leica DMIL microscope and bright-field culturing images were taken using the Leica Application Suite (LAS) Version 2.8.1 software.

398
399
400

**Array comparative genomic hybridization**

To test the genetic integrity of fibroblast-derived iPSCs pre- and postinduction, DNA extracts from fibroblasts and iPSCs were assessed using array comparative genomic hybrid analysis (Cell Guidance Systems). Briefly, DNA was isolated from cell pellets using DNeasy Blood & Tissue Kit (Qiagen, Cat. 69504) per manufacturer's instructions. All extracted DNA had a concentration >50 ng/µl and contained more than 1 µg. Purity measured on using an ND-1000 Spectrophotometer (NanoDrop Technologies, Inc.) fell within A260/A230 = 1.8 – 2.2 and A260/A280 > 1.8. The array contained 754,027 distinct probes and analysis was conducted on the Infinium Global Screening Array v3.0 platform. The genome-wide resolution was ~200 kb. Copy number variations and loss of heterozygosity was reported in samples when present in at least ~15 – 20% of cells.

**Germ-layer differentiation assay**

iPSCs were differentiated to the three germ layers as previously described [56,57] with minor modifications. Cells were plated onto a 12-well plate in chemically defined medium with polyvinyl alcohol (CDM-PVA: 250 mL IMDM, LifeTechnologies Cat. 21980032; 250 mL DMEM/F-12 + GlutaMAX, Gibco, Cat. 31765027; 1 mL Concentrated Lipids, Life Technologies, Cat. 11905031; 20 µL 1-Thioglycerol, Sigma, Cat. M6145-25ML; 350 µL Insulin, Sigma, Cat. I9278-5ML; 250 µL Transferrin, Sigma, Cat. 10652202001; 10 mL PVA (10 g [stock: 0.05 g/mL], Sigma, Cat. P8136-250G in 200 mL embryo-grade water, Sigma, Cat. W1503)). For one day CDM-PVA was supplemented with 10 ng/mL Activin (Gibco, PHC9564) and 12 ng/mL fibroblast growth factor-2 (FGF2) (Gibco, Cat. PHG0024). Thereafter, lineage specific protocols were followed, as reported below.

For endoderm, 100 ng/mL Activin, 80 ng/mL FGF2, 10 ng/mL BMP-4 (Gibco, PHC9534), 10 µM LY294002 (Gibco, Cat. PHZ1144), and 3 µM CHIR99021 (Sigma, Cat. SML1046-5MG) were supplemented in CDM-PVA medium for one day. The next day, the medium was refreshed at the same concentrations without CHIR99021. On the third day, the medium was changed to RPMI medium (500 mL RPMI1640 + GlutaMAX, LifeTechnologies, Cat. 61870010; 10 mL B27, Gibco, Cat. 17504044; 5 mL NEAA, Gibco, Cat. 11140035; 5 mL Pen/Strep, Gibco, Cat. 15140122) containing 100 ng/mL Activin and 80 ng/mL FGF2. On day four cells were fixed with 4% formaldehyde (Sigma, Cat. 158127) in DPBS and were tested for markers of endoderm as described below.

For mesoderm, 100 ng/mL Activin, 20 ng/mL FGF2, 10 ng/mL BMP4, 10 µM LY294002, and 3 µM CHIR99021 was supplemented to CDM-PVA for two days, with media change each day. On day three cells were fixed with 4% formaldehyde in DPBS and were tested for markers of mesoderm as described below.

For ectoderm, 10 nM SB431542 (Sigma, Cat. S4317-5MG), 12 ng/mL FGF2, and 150 ng/mL Noggin (Sigma, Cat. H6416-10UG) was supplemented to CDM-PVA medium for ten days, with media change each day. On day eleven cells were fixed with 4% Formaldehyde in DPBS and were tested for markers of ectoderm as described below.

To assess differentiation capacity, samples were analysed using the 3-Germ Layer Immunocytochemistry Kit (LifeTechnologies, Cat. A25538) per manufacturer's instructions. TUJ1, AFRP and SMA were included in this kit but anti-FoxA2/hHNF-Abs (R&D Systems, Cat. AF2400), diluted 1:100, and anti-Brachyury-Abs (R&D Systems, Cat. AF2085), diluted 1:100; and anti-Nestin-Abs (Abcam, Cat. ab22035), diluted 1:100 were used.

The next day, samples were incubated with the corresponding secondary Abs, which were diluted 1:250 in 1% BSA in PBS. AFP, SMA, and TUJ1 were counter-stained with AlexaFluor488 goat anti-mouse IgG1 (Invitrogen, Cat. A25536), AlexaFluor555 goat anti-

mouse IgG2a (Cat. A25533), and AlexaFluor647 donkey anti-rabbit (Invitrogen, Cat. A25537),
respectively from the 3-Germ Layer Immunocytochemistry Kit. FoxA2/hHNF and Brachyury
were counter-stained with mouse-anti-goat Texas Red (Santa Cruz Biotechnology, Cat. sc-
3916), and Nestin counter-stained with AlexaFluor 488 goat-anti mouse (Invitrogen, Cat.
A21121). After the secondary Ab incubation, the samples were mounted using ProLong™
Diamond Antifade Mountant with DAPI (Invitrogen, Cat. P36971). Images were taken with
the CLSM Leica SP5 resonant APD at the Center for Microscopy and Image Analysis,
University of Zurich.
456
**Mature neuronal culture**
To generate neural progenitors, iPSCs were disassociated into single cells using
ACCUTASE™ Cell detachment solution as described above, and seeded at high density ($1\times10^6$
cells per well of a 12-well plate) in SMAD inhibition medium as previously described [25]. Cell
counting was performed using the Countess 3 Automated Cell Counter (ThermoFisher
Scientific) with trypan blue (Gibco, Cat. 15250061) staining to identify dead cells, as per
manufactures instructions. Neural induction of iPSC was performed using the STEMdiff™
SMADi Neural Induction Kit (StemCellTechnologies, Cat. 08582) and protocol per
manufacturer was followed [26]. Briefly, iPSCs were cultured for 8-12 days on BME matrix (as
before) and passaged using 1mg/ml Collagenase type IV solution (Gibco, Cat. 17104019) upon
the appearance of neuroepithelial cells onto BME matrix coated 6-well plates in STEMdiff™
SMADi Neural Induction medium. The next day, medium was switched to neural maintenance
medium (NMM: 48.4 mL KnockOut DMEM/F12, Gibco, Cat. 12660-102; 48.5 mL
Neurobasal, Gibco, Cat. 12348-017; 1 mL B-27 Supplement, Gibco, Cat. 17504-044; 500 µL
N-2 Supplement, Gibco, Cat. 17502-048; 500 µL Non-essential amino acids, Gibco, Cat.
11140-050; 500 µL GlutaMAX, Gibco, Cat. 35050-061; 500 µL Penicillin-Streptomycin,
Gibco, Cat. 15140-122; 90 µL 2-Mercaptoethanol, Gibco, Cat. 31350010; 25 µL Insulin
solution, Sigma, Cat. I9278-5ML; as previously described [25]) and henceforth medium was
refreshed daily. Upon formation of rosettes (as described in [25,58]), cultures were treated with 5
µL/mL Recombinant Human Basic Fibroblast Growth Factor (bFGF, 4 µg/mL, Gibco, Cat.
PHG0024). After four consecutive days of bFGF treatment, cells were passaged with 1mg/ml
Collagenase type IV solution and transferred to BME matrix (Cultrex®) coated 6-plates. Once
morphologically apparent neuronal cells (polarised with leading axon and trailing dendrite)
could be seen migrating from the apical border of the rosette, cultures were passaged with
ACCUTASE™ Cell detachment solution and plated onto 10 µg/mL laminin (Sigma-Aldrich,
Cat. L2020) and poly-L-Lysine (Sigma-Aldrich, Cat. P6282-5MG). Cells were passaged
between day 21 and 26 and either continued as neural cultures or frozen for storage. For
freezing, cells were detached using ACCUTASE™ Cell detachment solution, transferred to
neural freezing medium (90% NMM, 10% DMSO (Sigma, Cat. D2438), 20 ng/mL bFGF), and
placed in a Mr. Frosty (Thermo Scientific, Cat. 5100-0001) at -80°C and stored long-term in
liquid nitrogen, as described [25].
488            Neural cultures were fed daily with NMM until day 45, after which 48-hour cycles were
489    implemented whereby every second day 2/3 of the medium was removed and 2 mL NMM plus
490    maturation supplements GDNF and BDNF (both at 20 ng/mL, StemCellTechnologies, Cat.
491    78058 and 78005) were supplied to neuronal cultures. In this way, neuronal cultures were
492    maintained up to a maximum of day 120.
493
**Mitochondrial fluorometric stains and dyes**
Mitochondrial oxidative phosphorylation complex I was inhibited using 1 µM rotenone
(Sigma, Cat. R8875-1G) in DMSO for 30 minutes added directly to the medium. Mitochondrial
membrane potential was assessed using Tetramethylrhodamine Methyl Ester Perchlorate

(Sigma, Cat. T668) reconstituted in DMSO and supplemented at 15 nM incubated for 60
minutes. Mitochondrial superoxide production was measured using MitoSOX™ Mitochondrial
Superoxide Indicators (Invitrogen, Cat. M36008) reconstituted in DMSO was supplemented at
5 µM for 15 minutes.

**Electrophysiology**

Cells were used for electrophysiology studies after at least 50 days *in vitro* (range: 50 – 113,
mean: 73) and at least 20 days following addition of the maturation supplements GDNF and
BDNF in 10 µg/mL laminin (Sigma-Aldrich, Cat. L2020) and transfer to poly-L-Lysine
(Sigma-Aldrich, Cat. P6282-5MG) coated Nunc™ Cell Culture/Petri Dishes (Thermo
Scientific, Cat. 150318). For treatment studies, 0.1 mM, 6 mM or 12 mM dimethyl 2-
oxoglutarate (Sigma, Cat. 349631) (DM-2OG) was added for 24 hours before cell collection.
For recording, iPSC-derived neurons were placed in a recording chamber and whole-cell patch-
clamp recordings were performed using an EPC10 USB amplifier (HEKA). iPSC-derived
neurons were bathed with extracellular solution containing (in mM) 140 NaCl, 2 $MgCl_2$, 2
$CaCl_2$, 10 HEPES, 3 KCl, 10 D-Glucose, pH 7.4 (all from Sigma). Pipette solution contained
(in mM) 4 NaCl, 120 K-gluconate, 10 HEPES, 10 EGTA, 3 Mg-ATP, 0.5 $CaCl_2$, 1 $MgCl_2$ pH
7.2 (all from Sigma). Patch pipettes ~9 MOhm were pulled from borosilicate glass capillaries
(Harvard Apparatus, Cat. 30-0038). Voltages were corrected for a liquid junction potential of
+12 mV. Series resistance was not compensated.

We used current-clamp recordings to examine AP firing and membrane properties. Step
current injection protocols (duration, 2 s; step, 5 pA; −20 to +60 pA) were applied from the
neuron's resting membrane potential. Current-clamp data were filtered at 2.7 kHz and digitized
with 20–100 kHz. To study whole-cell currents, voltage steps (duration, 1 s) were applied from
a holding potential of −80 mV to voltages between −80 and + 60 mV. Peak inward and outward
currents were quantified and normalized to whole-cell capacitance to obtain current densities.
Spontaneous excitatory postsynaptic currents were measured for 120 s at a holding potential of
−80 mV. Voltage-clamp data were filtered at 2.9 kHz and digitized with 50 kHz.

Where described, MitoTracker™ Deep Red FM (Invitrogen, Cat. M22426) was used at
0.1 µM in NMM and provided directly to cells for 15 minutes. Thereafter, medium was
aspirated, and cells washed with extracellular solution. Tested doses of MitoTracker™ Deep
Red FM $\geq$ 1 µM were not compatible with electrophysiological recording.

Electrophysiological data were analysed with custom-written routines in IgorPro
software (WaveMetrics, version 6.37) for current-clamp and voltage-clamp step protocols.
Spontaneous synaptic events were analysed using miniML[59] in Python (version 3.9). Extracted
data were integrated using R Studio (version 2022.12.0+353). Analysis scripts are available
upon request.

**Transcriptomics**

Bulk RNA-seq was performed by Novogene from RNA extracted from liquid-nitrogen snap
frozen cell pellets (DIV range: 65 – 78, mean: 75) that were lysed using QIAshredder (Qiagen,
79656) with RNA extracted using the RNeasy® Mini Kit (Qiagen, Cat. 74106). Prior to
sequencing, RNA quality was assessed using an ND-1000 Spectrophotometer (NanoDrop
Technologies, Inc.) to ensure sample purity was between A260/280 = 1.8-2.2 and A260/230 $\geq$
1.8. Peak illustration was conducted using an Agilent 5400 and the low marker was used to
calibrate the size of peaks. Between 960 to 8800 ng of total RNA was provided and $\geq$ 20 million
read pairs per sample were acquired.

Raw sequencing reads were preprocessed using the SUSHI framework [60,61], which was
developed by the Functional Genomics Center Zurich (FGCZ). Adapter sequences and low-
quality reads were trimmed off using fastp v0.02 [61]. Filtered reads were pseudo-aligned against

the reference human genome assembly GRCh38.p13 and quantified using Kallisto v0.46.1 [62].
Differential gene expression analysis was performed between different conditions using the R
package edgeR v3.42 [63]. 21057 features were identified and 15530 features had counts above
the threshold. $P < 0.05$ (reported as false discovery rate unless otherwise stated, correction is
done via Benjamini-Hochberg method) and one-fold change were considered to be
differentially expressed. R package clusterProfiler was used for functional enrichment analysis
based on Go terms and KEGG pathways, which utilised Over-Representation Analysis [64].

**Metabolomics – Sugar phosphates**
Polar metabolites were extracted from neuronal cells at day 60 cultured in 6-well plates, either
untreated or after 24 hours supplementation with 12 mM DM-2OG, according to the described
protocol [65]. After extraction, the metabolic extracts were concentrated and dried via an Alpha
3-4 LSCbasic (Martin Christ, 102394, 102395) in vacuo centrifuge with an external oil-pump
overnight at ambient temperature and stored at -80°C until analysis.
The metabolite extracts were dissolved in 100 μL of ultrapure water upon analysis, and
a volume of 8 μL per sample was injected into an Agilent 1290 ultra-high performance liquid
chromatography (UHPLC) module connect to an Agilent 6490A tandem quadruple (QqQ)
mass spectrometer, as previously described [31]. Chromatography was performed using a 0.25
μL/min flow rate and a gradient from 0 to 100% mobile phase B over a 25-minute total run
time. Mobile phase A (12 mM acetic acid, 10 mM tetrabutylammonium, 2 mM acetylacetone,
3% methanol in mQ) and B (12 mM acetic acid, 10 mM tetrabutylammonium, 2 mM
acetylacetone, 3% methanol, 80% acetonitrile in mQ) were used for phosphate sugar
separation. The gradient method was as follows (time: % B): 11 min: 0%; 14 min: 15%; 19
min; 40%; 20 min; 100%; 21.5 minutes 0%. The QqQ MS was operated in dynamic multiple-
reaction monitoring (MRM) mode. The list of transitions used to detect targeted compounds is
reported in Supplementary data 4. Skyline Software (v20.2, MacCoss Lab Software) was used
for peak integration and resulting peak areas were normalized on total peak area [31]. Data
processing and statistical data analysis were performed in PRISM GraphPad (v.5.03).
Technical triplicates were measured for each sample. Shapiro-Wilk test was used to assess
normality. Significance was evaluated for multiple comparison via Kruskal-Wallis test with
significance set at p<0.05 (without post hoc correction).

**Metabolomics – TCA cycle metabolites and stable isotope tracing**
Treated cells were supplemented with 12 mM DM-2OG for 24 hours. Four hours before cell
collection, medium was changed to NMM without GlutaMAX™ Supplement (Gibco, Cat.
35050061) and with 4 mM [U-$^{13}$C] glutamine (Sigma-Aldrich, Cat. 605166). At collection,
medium was removed, coverslips quickly dipped into sterile double-distilled water at 37 °C
and quenched in 80% methanol at −20 °C. Cells were scrapped in methanol and centrifuged at
15,000g for 15 minutes at 4 °C. Supernatants were collected, snap-frozen in liquid nitrogen and
stored at −80 °C before LC–MS analysis. Mass spectrometry preparation and processing was
performed as previously described [10].

**Immunohistochemistry**
To detect native protein markers, immunochemical assay was performed on cells grown on
glass coverslips treated with poly-L-Lysine. Depending on cell state, coverslips were also
treated with BME matrix (iPSCs, NECs, NSCs) or laminin (NPCs, Neurons). Briefly, cells
were washed with DPBS (Gibco, Cat. 14190144), and fixed with 4% paraformaldehyde
(Sigma-Aldrich, Cat. 158127) in DPBS for 15 minutes at ambient temperature. Cells were then
permeabilised for 30 minutes in 0.1% triton (diluted in DPBS), PFA was quenched using 100
mM glycine solution for 15 minutes, and coverslips blocked in 1% BSA for 60 minutes. Both

primary and secondary antibodies were diluted in 1% BSA. All solutions were prepared in
DPBS. A full list of antibodies, their sources, dilution ratios, incubation times and
manufacturers can be found in Supplementary Data Table 5.
601
**Imaging and microscopy**
Immunofluorescence images, unless otherwise stated, were taken with a DMi8 S Inverted
Microscope fitted with a DFC480 camera (Leica) and RGB filter boxes. Image acquisition was
adjusted for each fluorophore-antibody combination. All cells, unless otherwise stated, were
plated on glass coverslips (Epredia, Cat. 630-2124) treated with poly-L-Lysine (Sigma-
Aldrich, Cat. P6282-5MG) and mounted on Adhesive Microscopic Slides 25 x 75mm Menzel
(ThermoFisher, Cat. J1800AMNZ) with either ProLong™ Gold Antifade Mountant with DNA
Stain DAPI (Invitrogen, Cat. P36935) or ProLong™ Gold Antifade Mountant without DAPI
(Invitrogen, Cat. P10144).
Images were acquired using a Leica Confocal SP8 Inverse Falcon with Power HyD R
photon detector in counting mode, and fluorophores excited using a white light laser or
photomultiplier tube with a 405 diode to excite fluorophores in the UV spectrum. Images were
acquired using an HC PL APO 63x/1,40 OIL CS2 objective (Leica) with oil immersion liquid
("Type-F" immersion liquid, Leica). Acquisition parameters for high resolution images and
deconvolution: Z-step at 130 nm (Image slices range 18-25), Image size at 42.91 um. Image
deconvolution was performed using Bitplane Imaris.
618
**Correlation colocalization image analysis**
The analysis of colocalization of the MMUT and TOMM20 channels was performed using the
Coloc 2 plugin in ImageJ. This plugin performed a spatial pixel intensity correlation, creating
a scatterplot with a linear regression fit, which indicated the degree of colocalization between
the two channels. Pearson's correlation analysis and the calculated Pearson's R coefficient
were         extracted         and         reported,         (https://imagej.net/Coloc_2         and
https://imagej.net/Colocalization_Analysis).
626
**Immunoblotting**
Cells were collected with Accutase or scraping, washed with DPBS, and centrifuged to retrieve
pellets. The supernatant was removed, and the remaining pellet was either used immediately
for the lysate production or stored at -80°C for a later use. The cell pellets were resuspended in
100 µl of lysis buffer containing 150 mM NaCl (Sigma, Cat.#S7653), 50 mM Trizma-Base
(pH 8, Sigma, Cat.#T1503), 1% NP-40 (IGEPAL CA-630, Sigma, Cat.#I8896), 10% sodium
deoxycholate (Sigma, Cat.#D6750) and 1% HaltTM Protease & Phosphatase Inhibitor Cocktail
(100X, Thermo Fischer Scientific, Cat.#78440) and centrifuged max speed at 4°C for 5
minutes. The supernatant was collected in a pre-cooled microfuge tube. The protein
concentration of each sample was assessed by performing a Bradford assay (Quick Start
Bradford 1X Dye Reagent, BioRad, Cat.#5000205) and measuring the absorbance at 595 nm.
The samples were prepared by adding the protein extract to a mixture of lysis buffer and 4X
Laemmli buffer (BioRad, Cat.#161-0747) with 5% β-Mercaptoethanol (Sigma, Cat.#M6250)
to obtain a total protein concentration of 1 µg/µL. The samples were incubated at 96°C for 5
minutes, before 15 µg was loaded, alongside a molecular weight standard (PageRulerTM Plus
Prestained Protein Ladder, Thermo Fischer Scientific, Cat.#26619) on Novex™ Tris-Glycine
Mini Protein Gels, 4–20%, 1.0 mm, WedgeWell™ format (Invitrogen, XP04200BOX) or self-
cast 12% running gel (in µL: 2500 $H_2O$, 3000 Bis-Acrylamide (Bio-Rad, Cat.#1610158)
30:0.8, 1875 1.5M Tris-HCl (pH 8.8), 75 10% SDS, 50 10% APS, TEMED 5). The 10x running
running buffer comprised 1.92 M glycine (PanReac AppliChem, Cat.#A1067,1000), 250 mM
Trizma-Base and 35 mM Sodium dodecyl sulfate (SDS, Sigma, Cat.#L3771), which were

dissolved in double-distilled water. The gels were run at 140V until the sample front reached
the end of the gel. Proteins were transferred onto a nitrocellulose (pore size 0.45 μm; Cytiva,
Cat.#10600007) or polyvinylidene fluoride (pore size 0.22 μm, activated in MeOH; Bio-Rad,
Cat.#1620177) blotting membrane using the Trans-Blot® TurboTM Transfer Pack in running
buffer + 20% methanol with Bio-Rad standard program (30 mins). Ponceau staining (Sigma,
Cat.#P7170-1L) was performed to visualise protein transfer. Ponceau was washed off with
distilled water and blocked in blocking buffer at ambient temperature (10% 10x TBS-tween
(24 g Tris base, 88 g NaCl (Sigma, Cat.#71380-1kG), in 1L distilled water neutralised to pH
7.6 with HCl) 0.2% Tween (Sigma, Cat.#P1379-100ML), 5% milk powder (Millipore,
Cat.#70166-500G)) for 1 hour. Antibody staining was performed with agitation for 2 hours in
ambient conditions or overnight at 4°C. Membranes were washed in 1xTBS-tween in between
each step. Stained protein signal was developed using Cytiva GE ECL Solution (Amersham,
RPN2109) or Clarity Max ECL solution (Bio-Rad, 1705062) on the ChemiDoc™ Touch
imaging system (Bio-Rad). A full list of antibodies, their sources, dilution ratios, incubation
times and manufacturers can be found in Source data 5.
663
664 **Propionate Incorporation Assay**
665 The assay to quantify propionate pathway catabolism of iPSCs was assessed according to a
666 protocol described previously [66] and included modifications as described [24].
667
**Quantitative PCR**
To test for incorporation of remnants of the Sendai virus, RNA was extracted from fibroblasts,
Sendai virus-positive fibroblasts and iPSCs using either the QIAamp RNA Blood Mini Kit
(QIAGEN) or Qiagen RNeasy micro kit (QIAGEN) following manufacturer's instructions. The
isolated RNA was converted into cDNA, utilizing PrimeScript™ 1st strand cDNA Synthesis
Kit (Takara, Cat.#6210A). cDNA was amplified by PCR, using Taq Polymerase (New England
Biolabs) using the primers listed in Supplementary Data 7. The PCR products were stained
with EZ-Vision® One DNA Dye as Loading Buffer, 6X (Amresco), analyzed by gel
electrophoresis on a 2% agarose gel and visualized by an UV transilluminator (Alpha Imager
HP).
To test for expression of SOX2, PAX6 and EOMES, RNA was extracted from iPSC,
neuroepithelium and NSCs as described above. cDNA was synthesised from 500-3000 ng RNA
using the PrimeScriptTM II 1st Strand cDNA Synthesis Kit using a Professional TRIO
Thermocycler (Analytik Jena) following manufacturer's instructions. Real-time quantitative
PCR consisted of 10 μL of GoTaq® qPCR Master Mix (2X, Promega, Cat.#A600A), 5 μL
diethylpyrocarbonate (DEPC BioChemica, PanReac AppliChem, Cat.#A0881,0020) treated
water and 1 μL of a 10 mM solution containing forward and reverse primers for the genes of
interest, to which 4 μL of 300-3000 ng cDNA were added. The RT-qPCR was performed using
either a LightCycler® 480 II (Roche), a 7900HT Fast Real-Time PCR System (Applied
Biosystems), or a QuantStudio™ 7 Pro Flex Real-Time PCR System (Applied Biosystems,
4485701). In all cases the following conditions were used: 96°C for 5 minutes, followed by 45
cycles composed of 10 seconds at 96°C and 1 minute at 60°C. Technical triplicates were
performed for each sample. Data were collected using the LightCycler® 480 Software release
1.5.0 SP4 or exported from the QuantStudio as CSV. The relative gene expressions were
normalized using the housekeeping gene β-Actin and analysed using the $2^{-\Delta\Delta Ct}$ method [67].
Primer sequences can be found in Source data 6.
694
695 **Data availability**
696 All main data supporting the findings of this study are available within the article, Extended
697 Data, and Supplementary Information. Source data are provided with this paper.

# References

1. Rath, S. *et al.* MitoCarta3.0: an updated mitochondrial proteome now with sub-organelle localization and pathway annotations. *Nucleic Acids Res.* **49**, D1541–D1547 (2021).

2. Schaefer, A., Lim, A. & Gorman, G. Epidemiology of Mitochondrial Disease. in *Diagnosis and Management of Mitochondrial Disorders* (eds. Mancuso, M. & Klopstock, T.) 63–79 (Springer International Publishing, Cham, 2019). doi:10.1007/978-3-030-05517-2_4.

3. Rahman, S. Mitochondrial disease in children. *J. Intern. Med.* **287**, 609–633 (2020).

4. Rahman, J. & Rahman, S. Mitochondrial medicine in the omics era. *Lancet Lond. Engl.* **391**, 2560–2574 (2018).

5. Wajner, M. Neurological manifestations of organic acidurias. *Nat. Rev. Neurol.* **15**, 253–271 (2019).

6. DiMauro, S. & Schon, E. A. Mitochondrial disorders in the nervous system. *Annu. Rev. Neurosci.* **31**, 91–123 (2008).

7. Chinnery, P. F. Primary Mitochondrial Disorders Overview. in *GeneReviews®* (eds. Adam, M. P. et al.) (University of Washington, Seattle, Seattle (WA), 1993).

8. Forny, P. *et al.* Guidelines for the diagnosis and management of methylmalonic acidaemia and propionic acidaemia: First revision. *J. Inherit. Metab. Dis.* **44**, 566–592 (2021).

9. Froese, D. S. & Gravel, R. A. Genetic disorders of vitamin B12 metabolism: eight complementation groups – eight genes. *Expert Rev. Mol. Med.* **12**, e37 (2010).

10. Forny, P. *et al.* Integrated multi-omics reveals anaplerotic insufficiency in methylmalonyl-CoA mutase deficiency. (2023) doi:https://doi.org/10.1038/s42255-022-00720-8.

11. Kölker, S. *et al.* no Methylmalonic acid, a biochemical hallmark of methylmalonic

acidurias but no inhibitor of mitochondrial respiratory chain. *J. Biol. Chem.* **278**, 47388–

47393 (2003).

12. Luciani, A. *et al.* Impaired mitophagy links mitochondrial disease to epithelial stress in

methylmalonyl-CoA mutase deficiency. *Nat. Commun.* **11**, 970 (2020).

13. de Keyzer, Y. *et al.* Multiple OXPHOS deficiency in the liver, kidney, heart, and skeletal

muscle of patients with methylmalonic aciduria and propionic aciduria. *Pediatr. Res.* **66**,

91–95 (2009).

14. Collado, M. S. *et al.* Biochemical and anaplerotic applications of in vitro models of

propionic acidemia and methylmalonic acidemia using patient-derived primary

hepatocytes. *Mol. Genet. Metab.* **130**, 183–196 (2020).

15. Wongkittichote, P. *et al.* Tricarboxylic acid cycle enzyme activities in a mouse model of

methylmalonic aciduria. *Mol. Genet. Metab.* **128**, 444–451 (2019).

16. Chandler, R. J. *et al.* Mitochondrial dysfunction in mut methylmalonic acidemia. *FASEB*

*J.* **23**, 1252–1261 (2009).

17. Manoli, I. *et al.* Targeting proximal tubule mitochondrial dysfunction attenuates the renal

disease of methylmalonic acidemia. *Proc. Natl. Acad. Sci.* **110**, 13552–13557 (2013).

18. Ramon, C., Traversi, F., Bürer, C., Froese, D. S. & Stelling, J. No Cellular and

computational models reveal environmental and metabolic interactions in MMUT-type

No methylmalonic aciduria. *J. Inherit. Metab. Dis.* **n/a**,.

19. Kandel, E. R., Koester, J. D., Mack, S. H. & Siegelbaum, S. A. Synaptic Transmission. in

*Principles of Neural Science, 6e* (McGraw Hill, New York, NY, 2021).

20. Zheng, X. *et al.* Metabolic reprogramming during neuronal differentiation from aerobic

glycolysis to neuronal oxidative phosphorylation. *eLife* **5**, e13374 (2016).

21. Yellen, G. Fueling thought: Management of glycolysis and oxidative phosphorylation in

neuronal metabolism. *J. Cell Biol.* **217**, 2235–2246 (2018).

22. Bouzier-Sore, A.-K. *et al.* Competition between glucose and lactate as oxidative energy

substrates in both neurons and astrocytes: a comparative NMR study. *Eur. J. Neurosci.*

**24**, 1687–1694 (2006).

23. Hall, C. N., Klein-Flügge, M. C., Howarth, C. & Attwell, D. Oxidative Phosphorylation,

Not Glycolysis, Powers Presynaptic and Postsynaptic Mechanisms Underlying Brain

Information Processing. *J. Neurosci.* **32**, 8940–8951 (2012).

24. Froese, S. & Baumgartner, M. Lysosomal Vitamin B12 Trafficking. in *Ion and Molecule*

*Transport in Lysosomes* (CRC Press, 2020).

25. Shi, Y., Kirwan, P. & Livesey, F. J. Directed differentiation of human pluripotent stem

cells to cerebral cortex neurons and neural networks. *Nat. Protoc.* **7**, 1836–1846 (2012).

26. Chambers, S. M. *et al.* Highly efficient neural conversion of human ES and iPS cells by

dual inhibition of SMAD signaling. *Nat. Biotechnol.* **27**, 275–280 (2009).

27. Wurm, C. A. *et al.* Nanoscale distribution of mitochondrial import receptor Tom20 is

adjusted to cellular conditions and exhibits an inner-cellular gradient. *Proc. Natl. Acad.*

*Sci.* **108**, 13546–13551 (2011).

28. Bardy, C. *et al.* Predicting the functional states of human iPSC-derived neurons with

single-cell RNA-seq and electrophysiology. *Mol. Psychiatry* **21**, 1573–1588 (2016).

29. Inak, G. *et al.* Defective metabolic programming impairs early neuronal morphogenesis

in neural cultures and an organoid model of Leigh syndrome. *Nat. Commun.* **12**, 1929

(2021).

30. Galera-Monge, T. *et al.* Mitochondrial Dysfunction and Calcium Dysregulation in Leigh

Syndrome Induced Pluripotent Stem Cell Derived Neurons. *Int. J. Mol. Sci.* **21**, 3191

(2020).

31. van Scherpenzeel, M. *et al.* Dynamic tracing of sugar metabolism reveals the

mechanisms of action of synthetic sugar analogs. *Glycobiology* **32**, 239–250 (2022).

32. Hansen, K. B. *et al.* Structure, Function, and Pharmacology of Glutamate Receptor Ion

Channels. *Pharmacol. Rev.* **73**, 298–487 (2021).

33. Martinelli, D. *et al.* Neurologic outcome following liver transplantation for

methylmalonic aciduria. *J. Inherit. Metab. Dis.* **n/a**,.

34. Kaplan, P., Ficicioglu, C., Mazur, A. T., Palmieri, M. J. & Berry, G. T. Liver

transplantation is not curative for methylmalonic acidopathy caused by methylmalonyl-

CoA mutase deficiency. *Mol. Genet. Metab.* **88**, 322–326 (2006).

35. Molema, F. *et al.* Neurotoxicity including posterior reversible encephalopathy syndrome

after initiation of calcineurin inhibitors in transplanted methylmalonic acidemia patients:

Two case reports and review of the literature. *JIMD Rep.* **51**, 89–104 (2020).

36. Chakrapani, A., Sivakumar, P., McKiernan, P. J. & Leonard, J. V. Metabolic stroke in

methylmalonic acidemia five years after liver transplantation. *J. Pediatr.* **140**, 261–263

(2002).

37. Kasahara, M. *et al.* Current role of liver transplantation for methylmalonic acidemia: A

review of the literature. *Pediatr. Transplant.* **10**, 943–947 (2006).

38. Nyhan, W. L., Gargus, J. J., Boyle, K., Selby, R. & Koch, R. Progressive neurologic

disability in methylmalonic acidemia despite transplantation of the liver. *Eur. J. Pediatr.*

**161**, 377–379 (2002).

39. Mc Guire, P. J. *et al.* Combined liver–kidney transplant for the management of

methylmalonic aciduria: A case report and review of the literature. *Mol. Genet. Metab.*

**93**, 22–29 (2008).

40. Groten, C. J. & MacVicar, B. A. Mitochondrial Ca2+ uptake by the MCU facilitates

pyramidal neuron excitability and metabolism during action potential firing. *Commun.*

*Biol.* **5**, 1–15 (2022).

41. Chang, D. T. W., Honick, A. S. & Reynolds, I. J. Mitochondrial trafficking to synapses in

cultured primary cortical neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **26**, 7035–7045

(2006).

42. McKenna, M. C. The glutamate-glutamine cycle is not stoichiometric: Fates of glutamate

in brain. *J. Neurosci. Res.* **85**, 3347–3358 (2007).

43. Aldana, B. I. *et al.* Glutamate-glutamine homeostasis is perturbed in neurons and

astrocytes derived from patient iPSC models of frontotemporal dementia. *Mol. Brain* **13**,

125 (2020).

44. Andersen, J. V. *et al.* Deficient astrocyte metabolism impairs glutamine synthesis and

neurotransmitter homeostasis in a mouse model of Alzheimer's disease. *Neurobiol. Dis.*

**148**, 105198 (2021).

45. Shen, J. *et al.* Determination of the rate of the glutamate/glutamine cycle in the human

brain by in vivo 13C NMR. *Proc. Natl. Acad. Sci.* **96**, 8235–8240 (1999).

46. Chen, Q. *et al.* Rewiring of Glutamine Metabolism Is a Bioenergetic Adaptation of

Human Cells with Mitochondrial DNA Mutations. *Cell Metab.* **27**, 1007-1025.e5 (2018).

47. Yoo, H. C., Yu, Y. C., Sung, Y. & Han, J. M. Glutamine reliance in cell metabolism.

*Exp. Mol. Med.* **52**, 1496–1516 (2020).

48. Conte, F., van Buuringen, N., Voermans, N. C. & Lefeber, D. J. Galactose in human

metabolism, glycosylation and congenital metabolic diseases: Time for a closer look.

*Biochim. Biophys. Acta BBA - Gen. Subj.* **1865**, 129898 (2021).

49. Lissin, D. V., Carroll, R. C., Nicoll, R. A., Malenka, R. C. & Zastrow, M. von. Rapid,

Activation-Induced Redistribution of Ionotropic Glutamate Receptors in Cultured

Hippocampal Neurons. *J. Neurosci.* **19**, 1263–1272 (1999).

50. Lu, W.-Y. *et al.* Activation of Synaptic NMDA Receptors Induces Membrane Insertion

of New AMPA Receptors and LTP in Cultured Hippocampal Neurons. *Neuron* **29**, 243–

254 (2001).

51. Mosbacher, J. *et al.* A Molecular Determinant for Submillisecond Desensitization in

Glutamate Receptors. *Science* **266**, 1059–1062 (1994).

52. Zhu, J. J., Esteban, J. A., Hayashi, Y. & Malinow, R. Postnatal synaptic potentiation:

Delivery of GluR4-containing AMPA receptors by spontaneous activity. *Nat. Neurosci.*

**3**, 1098–1106 (2000).

53. Patterson, K., Linask, K. L., Beers, J. & Zou, J. Generation of two tdTomato reporter

induced pluripotent stem cell lines (NHLBIi003-A-1 and NHLBIi003-A-2) by AAVS1

safe harbor gene-editing. *Stem Cell Res.* **42**, 101673 (2020).

54. Chichagova, V., Sanchez-Vera, I., Armstrong, L., Steel, D. & Lako, M. Generation of

Human Induced Pluripotent Stem Cells Using RNA-Based Sendai Virus System and

Pluripotency Validation of the Resulting Cell Population. *Methods Mol. Biol. Clifton NJ*

**1353**, 285–307 (2016).

55. Ghaedi, M. & Niklason, L. E. Human Pluripotent Stem Cells (iPSC) Generation, Culture,

and Differentiation to Lung Progenitor Cells. *Methods Mol. Biol. Clifton NJ* **1576**, 55–92

(2019).

56. Vallier, L. & Pedersen, R. Differentiation of human embryonic stem cells in adherent and

in chemically defined culture conditions. *Curr. Protoc. Stem Cell Biol.* **Chapter 1**, Unit

1D.4.1-1D.4.7 (2008).

57. Vallier, L. *et al.* Early Cell Fate Decisions of Human Embryonic Stem Cells and Mouse

Epiblast Stem Cells Are Controlled by the Same Signalling Pathways. *PLoS ONE* **4**,

e6082 (2009).

58. Shi, Y., Kirwan, P., Smith, J., Robinson, H. P. C. & Livesey, F. J. No Human cerebral

cortex development from pluripotent stem cells to functional excitatory synapses. *Nat.*

*Neurosci.* **15**, 10.1038/nn.3041 (2012).

59. O'Neill, P. S. *et al.* Deep learning-based synaptic event detection. 2023.11.02.565316

Preprint at https://doi.org/10.1101/2023.11.02.565316 (2023).

60. Hatakeyama, M. *et al.* SUSHI: an exquisite recipe for fully documented, reproducible and

reusable NGS data analysis. *BMC Bioinformatics* **17**, 228 (2016).

61. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

*Bioinforma. Oxf. Engl.* **34**, i884–i890 (2018).

62. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

63. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinforma. Oxf. Engl.*

**26**, 139–140 (2010).

64. Yaari, G., Bolen, C. R., Thakar, J. & Kleinstein, S. H. Quantitative set analysis for gene

expression: a method to quantify gene set differential expression including gene-gene

correlations. *Nucleic Acids Res.* **41**, e170 (2013).

65. Conte, F. *et al.* In Vitro Skeletal Muscle Model of PGM1 Deficiency Reveals Altered

Energy Homeostasis. *Int. J. Mol. Sci.* **24**, 8247 (2023).

66. Willard, H. F., Ambani, L. M., Hart, A. C., Mahoney, M. J. & Rosenberg, L. E. Rapid

prenatal and postnatal detection of inborn errors of propionate, methylmalonate, and

cobalamin metabolism: A sensitive assay using cultured cells. *Hum. Genet.* **34**, 277–283

(1976).

67. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-

time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods San Diego Calif*

**25**, 402–408 (2001).

# Acknowledgements

The authors thank Adhideb Ghosh for his assistance with transcriptomic data and Caroline Frei for her assistance with analysis R script development. MCSD was supported by the FZK Fellowship of the University Children's Hospital Zurich. DSF is supported by the Swiss National Science Foundation [310030_192505] and the University Research Priority Program of the University of Zurich ITINERARE – Innovative Therapies in Rare Diseases. MRB is supported by the Swiss National Science Foundation [310030_175779 and 310030_212505] and the University Research Priority Program of the University of Zurich ITINERARE – Innovative Therapies in Rare Diseases.

# Author contributions

MCSD, DSF, and MRB contributed to the conception and design of the study. MCSD carried out the experimental work contributing to the majority of the paper. MCSD and DSF prepared the manuscript with input from the other authors. MS generated patient-derived iPSCs. GM performed western blots and ICC. MAG performed western blots. DPR performed electrophysiology and ID helped with analysis and wrote some of the analysis scripts. MP performed TCA cycle metabolomics. FC performed HexP metabolomics. BV and FvM facilitated bulk transcriptomics and data integration. All authors contributed to the manuscript and approved the submitted version.

# Competing interests

All authors declare no competing interests.

**Correspondence and requests for materials** should be addressed to D. Sean Froese, or Matthias R. Baumgartner

# Extended Figures

Extended Figure 1. **Fibroblasts generate iPSCs without incorporating viral factors. a.** Brightfield
microscope images of iPSC generation process. **b.** Sendai virus clearance from fibroblasts to iPSCs in
control (BJ), and patient (AV, KK) cell lines. **c.** Endogenous mRNA expression of pluripotency markers
in generated control, and patient cell lines, with purchased ND2 control line. **d.** Immunochemistry of
selected antibody against MMUT in WT and KO HEK293 cells. Scale bar = 10 μm. **e.** Western blot
analysis of MMUT in WT and KO HEK293 cells. Uncropped membranes are available in Supplementary
Data Fig. 2c.

Extended Figure 2. **Patient-derived neurons have disrupted mitochondria. a.** Confocal microscope images from control and patient demonstrating TOMM20 staining pattern. Images are representative. Scale bar = 5 µm. **b.** Confocal images of localisation of MMUT (green) to TOMM20 (grey) in control (ND2, BJ) and patient (KK, AV) cell lines. Scale = 20 µm. **c.** Volume and sphericity measurements for Z-stack confocal images of TOMM20 in control and patient cells. Patient images were separated into MMUT localised (MMUT+) or MMUT mislocalised (MMUT-). Datapoints are representative of ROIs from one non-overlapping image. **d.** Maximum projection of confocal TOMM20 immunofluorescent images in control and patient (left), and ferret diameter measurements of those mitochondrial networks identified through TOMM20 staining. Scale =1 µm. Datapoints are representative of one ROI. Statistical test = Mann-Whitney, **** = <0.0001 (Right). **e.** MitoSOX fluorescence measurements from confocal microscopy images in control and patient, without (left, baseline) and with (right, rotenone) 1 µM rotenone treatment. Datapoints are representative of one ROI. Statistical test = Kruskal-Wallis, ns = not significant, * = <0.05, ** = <0.01,*** = <0.001. **f.** Representative epifluorescence images of DAPI+ nuclei in control and patient during interphase and apoptosis. Scale = 10 µm.

Extended Figure 3. **Electrophysiological characterisation of 226 neurons. a.** PCA of 31 features in 226 independent cells highlighting the 3 neuron types, measured during current-clamp recordings. List of features in Supplementary Data Table 1. **b.** Refined PCA of 21 features in 226 independent control (BJ, ND2) and patient (AV, KK) cells. **c.** Box plots indicate median and IQR 1.5 from Type 1-3 neurons for 3 selected features, each datapoint represents one independent cell (left). PCA loadings for the 21 features on 226 neurons in Fig. 4a, colour = percentage contribution to explained variance (right). **d.** Unguided hierarchical heatmap clustering on measured feature in 226 neurons. One row indicates one cell, one column represent one feature.

Extended Figure 4. **Loss of excitability in patient-derived neurons is explained by mitochondrial disruption. a.** Negative correlation between AP amplitude and threshold or full-width half-max in control and patient type 3 neurons. **b.** Positive correlation between AP amplitude and number or resting membrane potential and AP full-width half-max in control and patient type 3 neurons. **c.** Loss of correlation in patient-derived type 3 neurons between Attenuation ratio and AP amplitude, threshold or full-width half-max. **d.** Immunofluorescent images of MitoTracker in control and patient-derived neurons, showing intact (yellow arrowhead) or disrupted (magenta arrowhead) mitochondria, images are representative (left). Box plot indicates median and IQR 1.5 of attenuation ratio from 12 neurons measured in tandem with MitoTracker visualisation, each datapoint represents one independent cell (right). **e.** Box plot indicates median and IQR 1.5 of dVdt in type 3 control and patient neurons, each datapoint represents one independent cell. **f.** Correlation between attenuation ratio and dVdt measurements in type 3 neurons, each datapoint represents one independent cell. **g.** Loss of correlation in patient-derived type 3 neurons between dVdt ratio and AP number or amplitude.

Extended Figure 5. **Attenuation phenotype in patient-derived neurons coherent with reduced sodium and potassium current densities a,** Stacked bar graph of proportional generation of type 1, 2, or 3 control (ND2, BJ) and patient (AV, KK) neurons in each cell line of cells measured in both current clamp and voltage clamp (n = 181). **b,** Box plot indicates median and IQR 1.5 of attenuation (left) and dVdt (right) ratios in control and patient neurons. **c,** Sodium and Potassium current density measurements from voltage-clamp recordings in 39 type 1 neurons. **d,** Sodium and Potassium current density measurements from voltage-clamp recordings in 42 type 2 neurons. **e,** Sodium and Potassium current density measurements from voltage-clamp recordings in 53 type 3 neurons.

957
Extended Figure 6. **Absolute abundance of TCA cycle metabolites is unchanged in patient-**
**derived neurons. a.** Box plot of ion abundance of disease-related metabolites. **b.** Box plot of ion
abundance of anaplerotic metabolites. **c.** Box plot of ion abundance of TCA cycle metabolites. **d.**
Relative fractional incorporation of fully labelled glutamine into glutamine, glutamate and 2-oxoglutarate.
**e.** Bar plots of fractional contribution of labelled carbons to glutamine, glutamate and 2-oxoglutarate. **f.**
Relative fractional incorporation of fully labelled glutamine into TCA cycle metabolites. **g.** Bar plots of
fractional contribution of labelled carbons to TCA cycle metabolites. In **a, b, c, e, g,** each datapoint is
an averaged result from technical triplicates of an independent sample. In **d, f** data is presented as a
fraction of complete labelled and unlabelled measured metabolite. In **a-c**, significance = unpaired t-test.
967

968
969 Extended Figure 7. **Changes in metabolite pools of selected sugar phosphates. a.** Box plot of
970 normalised area of the peak of targeted hexose phosphates. **b.** Box plot of normalised area of the peak
971 of amine-hexoses. **c.** Box plot of normalised area of the peak of targeted nucleotide sugars. **d.** Box plot
972 of normalised area of the peak of targeted pentose phosphates. In **a-d,** significance testing = paired t-
973 test.
974
975
976

977
Extended Figure 8. **DM-2OG treatment rescues metabolic phenotype in patient neurons. a.** Box
plot of response of absolute ion abundance of TCA cycle metabolites to treatment with dimethyl-2-
oxoglutarate (DM). **b.** Box plot of glutamate and glutamine response to DM-2OG treatment. **c.** Box plot
of response of absolute ion abundance of disease-related metabolites to treatment with dimethyl-2-
oxoglutarate (DM). **d.** Box plot of normalised area of the peak of targeted hexoses and response to DM-
2OG treatment. **e.** Box plot of normalised area of the peak of *N*-acetylhexosamine phosphates and
response to DM-2OG treatment. In **a-e,** datapoints are representative of independent samples
measured in triplicate and averaged. In **a-e** multiple testing = Mann-Whitney.
986
987

988
Extended Figure 9. **Patient excitatory perturbation can be mimicked in controls using DM-2OG.**
**a.** Box plot of action potential count in current-clamp at various DM-2OG concentrations. **b.** Box plot
indicates median and IQR at 1.5 from the measurement of resting membrane potential at different
treatments of DM-2OG. **c.** Box plot indicates median and IQR at 1.5 from the measurement of
capacitance at different treatments of DM-2OG. In **a, b, c,** datapoints are representative of an
independently measured sample. **d.** Confocal immunofluorescence of an antibody against SLC17A7
(VGLUT1), images are representative. Scale = 10 µm. **e.** Box plot indicates median and min/max from
quantification of synaptic VGLUT1 events in control (n=4) and patient (n = 9) neurons at DIV 50.
Datapoints are representative of averages from non-overlapping images. **f.** Box plot of spontaneous
event amplitude in response to DM-2OG treatment at 0.1mM. **g.** Box plot of spontaneous event decay
in response to DM-2OG treatment at 0.1mM. **h.** Box plot of spontaneous event frequency in response
to DM-2OG treatment at 0.1mM. In **f,g,h,** box plot indicates median and min/max. Where state
numerically, *p* value is Bonferroni significance adjusted.

Extended Figure 10. **DM-2OG influences glutamate-sensitive synaptic apparatus. a.** PCA of control
(n = 5) and patient (n = 6) neuronal cultures' transcriptome. **b.** Differentially regulated genes
corresponding to GO terms associated to biological processes by overrepresentation analysis. **c.**
Differentially regulated genes corresponding to GO terms associated to molecular functions by
overrepresentation analysis. **d.** Box plot indicates median and min/max of count of the neuronal
glutamate solute carrier extracted from transcriptomic data. Box plot indicates median and min/max of
count of the selected genes extracted from transcriptomic data. In **a, d, f, g,** datapoints are
representative of an independently measured sample. In **b,c,** the top 10 (in either direction) differentially
expressed terms were selected.
1011
1012