Population structure and genetic diversity of the Critically Endangered

bowmouth guitarfish (*Rhina ancylostomus*) in the Northwest Indian Ocean

Marja J. Kipperman[1]*, Rima W. Jabado[2,3], Alifa Bintha Haque[4,5], Daniel Fernando[6], P.A.D.L

Anjani[6], Julia L.Y. Spaet[7], Emily Humble*[1]

[1]Royal (Dick) School of Veterinary Studies and the Roslin Institute, University of Edinburgh,

EH25 9RG, Edinburgh, UK

[2]College of Science and Engineering, Centre for Sustainable Tropical Fisheries and

Aquaculture, James Cook University, Townsville, Australia

[3]Elasmo Project, Dubai, United Arab Emirates

[4]Department of Zoology, University of Dhaka, Dhaka, Bangladesh

[5]Nature-based Solutions Initiative, Department of Biology, University of Oxford, Oxford, UK

[6]Blue Resources Trust, Colombo, Sri Lanka

[7]Evolutionary Ecology Group, Department of Zoology, University of Cambridge, Cambridge CB2

3EJ, UK

*Corresponding authors

Correspondence: m.j.kipperman@gmail.com; emily.humble@ed.ac.uk

Abstract

The bowmouth guitarfish (*Rhina ancylostomus*) is a unique and relatively understudied species

of wedgefish with a distribution spanning the Indo-Pacific Oceans. Due to targeted and bycatch

fisheries, this species is experiencing serious declines across its range. It is now considered

among the most threatened species of elasmobranch. Despite this, species-specific

management is limited, particularly around primary fishing hotspots. This is in part due to

knowing very little about fundamental population processes. Here, we combine mitochondrial

and single nucleotide polymorphism (SNP) data to carry out the first genetic assessment of *R.*

ancylostomus across the Northwest Indian Ocean. We find evidence for population structure

across the species' northwest range that is best explained by dispersal constraints and

therefore emphasises the importance of habitat connectivity. We also uncover a cline in genetic

variation from east to west of the sampling distribution and identify regions in potential need of

heightened protection. In doing so, our study provides baseline information to support

conservation management of *R. ancylostomus* and highlights the value of genomic data in

elasmobranch conservation.

Keywords

Conservation genomics, elasmobranchs, SNPs, mitochondrial DNA, fisheries management

Introduction

Characterising genetic diversity and population structure across extant species' ranges is crucial for both *in situ* and *ex situ* conservation management (Hohenlohe et al., 2021). However, for many threatened species, such baseline information is severely lacking (Borgelt et al., 2022; Hochkirch et al., 2021; Tella et al., 2013). This compromises both the development and effectiveness of conservation actions. For example, the appropriate geographic scale of management largely depends on the extent to which populations are connected across space (Lowe & Allendorf, 2010; Mills & Allendorf, 1996; Mönkkönen & Reunanen, 1999). Discrete populations exchanging limited genetic information will benefit from local or regional scale management and may warrant delineation as conservation units (Moritz, 1999; Palsbøll et al., 2007). In contrast, populations connected by dispersal will require a more coordinated approach across larger spatial scales.

Large marine organisms exhibit contrasting patterns of genetic connectivity (Bailleul et al., 2018; Bernard et al., 2021, 2025; Pirog et al., 2019) Pelagic species with high dispersal capabilities frequently display high levels of gene flow across continuous distributions (Palumbi, 2003; Waples, 1998). This presents a challenge for conservation since it is not always clear where to delineate management units (Turbek et al., 2023). For example, individuals at opposite ends of a distribution may display distinct genetic and phenotypic variation yet remain connected by populations continuously distributed across space (Burri et al., 2016; Devitt et al., 2011; Irwin et al., 2005). Conversely, species exhibiting high site-fidelity and restricted movement are more likely to develop genetic differentiation between populations. If certain populations become small and isolated, they face increased vulnerability to exploitation, primarily due to heightened risks of inbreeding and loss of adaptive potential resulting from reduced genetic variation (Kimura et al., 1963; Lande, 1993; Lande & Shannon, 1996). From observational monitoring alone, it can be virtually impossible to identify such populations, especially in a marine environment.

Genetic and genomic tools provide a fantastic opportunity to explore the landscape of genetic diversity and differentiation (Bertola et al., 2024; Funk et al., 2012; Stapley et al., 2010) and are increasingly being applied to elusive marine megafauna (Kelley et al., 2016). For example, recent work has uncovered contrasting patterns of genetic diversity in two recently diverged manta ray (Mobulidae) species (Humble et al., 2024), signals of inbreeding in great hammerhead sharks (*Sphyrna mokarran*) (Stanhope et al., 2023) and subtle stock structure in

blue shark (*Prionace glauca*) – a species previously thought to comprise a single global

population (Nikolic et al., 2023). These insights all have the potential to influence management

through highlighting populations in need of prioritisation, contributing to non-detriment findings

under the Convention on International Trade of Endangered Species of Wild Flora and Fauna

(CITES) and informing fisheries management (Casey et al., 2016; Domingues et al., 2018).

Guitarfish and wedgefish are among the most vulnerable and understudied groups of

elasmobranchs (Dulvy et al., 2014; Kyne et al., 2020; Moore, 2017; Pytka et al., 2023). Their

mostly coastal and in-shore habitat overlaps with much of the world's most intensive fishing

activities making them highly susceptible to bycatch (Jabado, 2018; White et al., 2013; Pytka et

al. 2023), while international demand for their sought-after fins and thorns drive targeted

exploitation (Clarke et al., 2006; Pytka et al., 2023). Consequently, 'shark-like rays' have been

listed on Appendix II of CITES which seeks to regulate commercial trade, including of their

derivative products. Among the most valuable fins on the market are those of the bowmouth

guitarfish (*Rhina ancylostomus*), an evolutionarily distinct wedgefish (Gumbs et al., 2024) with

the most widespread and continuous distribution of all shark-like rays (Figure 1). This species is

listed as Critically Endangered on the IUCN Red List of Threatened Species and is particularly

susceptible to population decline due to its slow growth, late maturity, and low fecundity (Kyne

et al., 2019). Despite this, *R. ancylostomus* receives little to no species-specific management

across its entire range, putting the species at serious risk of extinction.

Here, we carry out the first genetic assessment of *R. ancylostomus* across the Northwest Indian

Ocean. We use a combination of mitochondrial and single nucleotide polymorphism (SNP)

markers to build an atlas of genetic variation in order to address the following questions: (i) to

what extent are populations genetically connected across the distribution; (ii) what are the levels

of genetic variation at both mitochondrial DNA and SNPs and how is this partitioned among

populations; and (iii) what are the implications of our findings for conservation management.

Materials and Methods

Sampling and DNA extraction

Fin clips were collected from 70 *R. ancylostomus* individuals from five geographic locations

across the Northwest Indian Ocean: Saudi Arabia (*n* = 4), United Arab Emirates, Persian Gulf (*n*

= 40), Oman (*n* = 12), Sri Lanka (*n* = 9) and Bangladesh (*n* = 5) (Figure 1). In all cases, samples

were obtained from specimens captured in fisheries and stored in 99% ethanol at -22°C. Total

genomic DNA was extracted using the Qiagen DNeasy® Blood and Tissue Kit following

manufacturer protocols and quantified using both a Qubit dsDNA Broad Range Assay and

running on a 1% agarose gel.

Figure 1. Geographic distribution of *R. ancylostomus* (dark blue) visualised together with the locations of

samples used in this study. Sampling location points are distinguished by colour and scaled by the

number of samples. The sample from Taiwan represents the *R. ancylostomus* mitochondrial genome

assembly downloaded from NCBI and included in our mitochondrial DNA analysis. Further details are

provided in Table S1.

Mitochondrial sequencing

Two regions of the mitochondrial genome were targeted for sequencing: cytochrome oxidase I

(COI) and the control region (CR). A 650 bp region of the COI was amplified using universal

primers FishF2 (5' TCGACTAATCATAAAGATATCGGCAC 3') and FishR2 (5'

ACTTCAGGGTGACCGAAGAATCAGAA 3') (Ward et al., 2005). PCRs were carried out using

10 μl DreamTaq Green PCR Master Mix (2X) (ThermoFisher Scientific), 6 μl nuclease free

water, 1 μl of each primer and 2 μl of template DNA. The following PCR profile was used for

amplification: initial denaturation of 3 min at 95°C; 35 cycles of 30 sec at 95°C, 30 sec at 55°C,

and 30 sec at 72°C, followed by a final extension of 5 min at 72°C. To amplify the CR, we

designed a new set of primers (RACR_F: 5' TGGGCTGGCGAGAAATAACC 3', RACR_R: 5'

TTTTCGTTTCGACCCAGGGG 3') in Geneious Prime v11.0.14.1 using the *R. ancylostomus*

mitogenome as a reference (GenBank accession number KU721837.1). PCRs were carried out

using the same volumes as above, together with the following amplification profile: initial

denaturation of 1 min at 95°C; 35 cycles of 30 sec at 95°C, 30 sec at 58°C, and 30 sec at 70°C,

followed by a final extension of 5 min at 70°C. The target product size for the CR fragment was

560 bp. PCR products were run on a 1.5% agarose gel to determine if amplification was

successful. Successful amplicons were purified using the Exo-CIP Rapid PCR Cleanup Kit (New

England BioLabs®) and sent for bi-directional Sanger sequencing with Eurofins Genomics.

Sequences were manually checked and edited using Geneious v1.2.3. Forward and reverse

reads for each individual and each mitochondrial DNA (mtDNA) target region were *de novo*

assembled into consensus sequences. Prior to further processing, each COI and CR consensus

sequence was blasted to the NCBI non-redundant reference database using megablast and an

E-value threshold of 0.05 for species identification purposes. Any sequence whose top BLAST

hit based on sequence similarity and bit score was not *R. ancylostomus* was removed from the

dataset. Remaining COI and CR consensus sequences were then multiple aligned using the

Clustal Omega algorithm in Geneious and trimmed to the same length for downstream analysis.

To increase the geographic scope of our analysis, we added target regions from the *R.*

ancylostomus mitochondrial reference genome to each alignment. This reference sequence was

generated using a sample originating from the Taiwan Strait (Si et al., 2016).

SNP genotyping and filtering

SNP genotyping was carried out by Diversity Arrays Technology (DArT) who implemented a

reduced representation sequencing protocol (DArTseq) followed by SNP calling with the

DARTsoft14 pipeline (Cruz et al., 2013; Kilian et al., 2012). Briefly, DNA from 65 of the samples

identified as *R. ancylostomus* was digested using PstI and SphI enzymes followed by adaptor

ligation for each individual. Enzymes were chosen by DArT for their ability to isolate highly

informative, low copy fragments of the genome. Following competitive PCR amplification, the

resulting library was 250 bp paired-end sequenced on an Illumina HiSeq2500. To obtain a high

quality dataset for downstream analysis, we filtered the resulting SNP genotypes using the R

package *dartR* (Gruber et al., 2018; Mijangos et al., 2022). For this, we removed SNPs with low

reproducibility scores based on technical replicates, retained one SNP per locus to account for

potential linkage in the dataset and removed SNPs with a read depth less than 5 or greater than

50, and with a genotyping rate less than 80%. We then removed individuals with a call rate

below 96%, that displayed excess heterozygosity or that displayed high relatedness with any

other individual in the dataset. To determine relatedness, we estimated KING, R0 and R1

coefficients (Waples et al., 2019) using NgsRelate v2 (Korneliussen & Moltke, 2015). These

statistics are based on genome-wide patterns of identity by state sharing between two

individuals. This analysis was carried out using a highly informative dataset whereby SNPs that

deviated significantly from HWE with a p-value threshold of 0.001, with a minor allele frequency

< 0.1 or a genotyping rate <0.9 had been removed using PLINK v1.90. The individual with the

lowest genotyping rate from any pairing exhibiting high relatedness was removed from the final

dataset (Figure S1). At this point, we applied a final filter to the data to remove any SNPs with a

minor allele frequency (MAF) <0.03, equivalent to a minor allele count of 3.

Population structure

To investigate population structure using mtDNA, we generated median joining haplotype

networks with PopART v1.7 (Leigh & Bryant, 2015) based on both the COI and CR alignments.

Haplotype accumulation curves (HACs) were then calculated to determine how well within-

species diversity was captured with our sampling effort. For this, we used the R package

HACSim to simulate the number of samples required to observe the full range of haplotype

variation that exists for a species (Phillips et al., 2020). The simulation was run for 10,000

permutations, a confidence level of 0.95 and at different percentage recovery rates until no

additional individuals were necessary for recovery of haplotypes (0.95). We further explored

population differentiation by calculating pairwise F_{ST} between sampling locations using Arlequin

v3.5.22 (Excoffier & Lischer, 2010). This was run separately for each mtDNA gene with 1000

permutations to estimate 95% confidence intervals. Finally, to measure population genetic

structure both within and between groups, we ran an Analysis of Molecular Variance (AMOVA)

using Arlequin and tested significance using 1000 permutations.

To investigate population structure using the SNP data, we implemented three approaches.

First, we performed a principal components analysis using the R package *adegenet* (Jombart,

2008). Second, we used the Bayesian clustering algorithm in STRUCTURE to identify the

number of genetic clusters (K) in the dataset. STRUCTURE was run for values of K ranging

from $K = 1 – 8$, with ten simulations for each K and a burn-in of 50,000 iterations followed by

100,000 MCMC iterations. We used the admixture and correlated allele frequency models

without sampling location information. We then used the R package pophelper (Francis 2017) to

analyse the results, parse the output to CLUMPP for averaging across simulations and to

visualise assignment probabilities. The optimal K was identified based on the maximum value of

the mean estimated *In* probability of the data (Ln Pr($X \mid K$)) (Pritchard et al 2000) and the ΔK

method (Evanno et al 2005). Finally, we estimated pairwise genetic differentiation between

populations using the Weir and Cockerham F_{ST} value (Weir & Cockerham, 1984) in the R

package dartR. Confidence intervals and *p*-values were estimated based on bootstrap

resampling of loci within each sampling location 1000 times (Weir & Cockerham, 1984). To

account for the effects of unbalanced sampling on our inference of population structure

(Puechmaille, 2016), we randomly downsampled the UAE population to eight individuals,

resulting in a more uniform distribution of samples across locations. We recalculated MAF and

removed SNPs with a MAF <0.03 before re-running PCA, STRUCTURE and F_{ST} as described

above. Results displayed similar patterns for the subsampled dataset as the full dataset

(Figures S7–9) and therefore those for the full dataset are presented below.

Isolation by distance

To investigate patterns of isolation by distance in *R. ancylostomus*, we explored the relationship

between genetic and geographic distance between all pairs of populations. Geographic location

was assigned to each individual by determining the coordinates of the coastline directly adjacent

to the fisheries landing site in which the sample was collected. Although this is not the true

location of where the individual was caught, it represents a fair approximation given the coastal

nature of the species. Genetic distances were based on COI-based pairwise F_{ST} estimates as

calculated above. We determined geographic distances based on a least-cost path analysis

using the R package *marmap* (Pante & Simon-Bouhet, 2013). A minimum depth constraint of -

10 metres was applied to ensure no paths were overland. Significance between genetic and

geographic distance matrices was then calculated using distance-based Moran's eigenvector

maps (dbMEM) by redundancy analysis (RDA, Legendre et al., 2015). Geographical distances

were transformed into dbMEMs using the R package *adespatial*, and genetic distances were

decomposed into principal components using the R function *prcomp*. RDA was then performed

using the R package *vegan*, with significance tested using 1000 permutations.

Genetic diversity and heterozygosity

Genetic diversity estimates were calculated for the mtDNA data using the software DnaSNP

v6.12.03 (Rozas et al., 2017) at both a species and sampling location level. For the SNP data,

we assessed levels of genetic variation by estimating multi-locus heterozygosity for each

individual using the R package *inbreedR* (Stoffel et al., 2016).

Results

Mitochondrial sequencing

Out of a total of 70 samples, 66 were confirmed in a BLAST search using COI and / or CR

sequences to originate from *R. ancylostomus*. The remaining four samples were identified as

tiger shark (*Galeocerdo cuvier)*, painted sweetlips (*Diagramma pictum)*, starry puffer (*Arothron*

stellatus) and *Pseudomonas campi* and were removed from further analysis. These results are

likely due to mislabelling of samples. Of the 66 confirmed *R. ancylostomus* samples, a total of

63 successfully amplified at COI, and 65 successfully amplified at the CR. Final filtered

alignment lengths were 674 bp for COI and 512 bp for the CR. Analysis of mitochondrial

haplotypes revealed a total of eleven unique sequences for COI and three for the CR, including

the sequence originating from the mitochondrial reference genome. Of these, ten and two

haplotypes were newly discovered, respectively (Table 1). All measures of genetic diversity

were consistently higher for COI compared to the CR (Table 1).

Table 1. Genetic diversity measures for *R. ancylostomus* calculated from COI and CR sequences

generated as part of this study.

Indices	COI	CR
Number of sequences	63	65
Number of segregating sites (*S*)	18	4
Number of haplotypes (*H*)	10	3
Haplotype diversity (*Hd*)	0.65028	0.49135
Nucleotide diversity (*pi*)	0.00377	0.00181

SNP genotyping

A total of 19,436 SNPs were genotyped in 65 individuals using the DARTsoft14 pipeline. After

quality filtering, the final dataset contained 3,533 SNPs in 49 individuals from across all

sampling locations (Saudi Arabia = 2, Oman = 8, UAE = 33, Sri Lanka = 5, Bangladesh = 1). For

a full breakdown of the number of SNPs and individuals removed at each filtering step, see

Table S2.

Population structure

Haplotype networks for both gene regions revealed a noticeable geographic signal from west to

east of the sampling range (Fig 2A–B). This pattern was more distinct for COI, where a greater

number of haplotypes were observed overall. In particular, we observed no haplotype sharing

between locations at the extremes of the sampling range while Sri Lanka, which was

intermediate, shared haplotypes with every location except for Taiwan. We also observed

geographic signal among samples collected in Sri Lanka, where individuals originating from the

west coast shared no haplotypes with those originating from the east coast.

Figure 2. Mitochondrial-based population differentiation and isolation by distance in *R. ancylostomus*. Median-joining mitochondrial haplotype network for (A) CR and (B) COI sequences. Colours reflect geographic origin, node size is proportional to the number of individuals displaying each haplotype and hashed lines indicate the number of nucleotide substitutions between haplotypes. (C) Pairwise F_{ST} estimates between sampling locations based on COI. Population abbreviations: BAN, Bangladesh; SL, Sri Lanka; UAE, United Arab Emirates; OMAN, Oman; SA, Saudi Arabia. (D) Relationship between genetic (COI based F_{ST}) and geographic distance as calculated by least-cost path analysis for all pairwise population comparisons. Solid lines and shaded areas reflect the regression slopes and standard errors, respectively, based on a linear model.

Based on haplotype accumulation curves (Figure S2), our COI dataset represents around 85% of the haplotype diversity in the species. An additional 63 and 132 individuals would be required to recover 95% and 99% of the haplotype diversity respectively (Figure S2A–B). For the CR, our sample size of 65 individuals was sufficient to recover both 95% and 99% of the haplotype diversity in the species (Figure S2C–D).

Population differentiation

To further explore levels of differentiation, we carried out an AMOVA and calculated pairwise

F_{ST} using both mitochondrial gene regions. AMOVA revealed that most of the observed

variation is partitioned within populations for both COI and the CR (Table 2). However, there

was also significant evidence of variation at a population level, with 29.07% and 25.58% of the

variation occurring between populations for COI and CR respectively. Pairwise F_{ST} estimates

ranged from 0.04–0.75 for COI and from -0.18–0.76 for the CR (Figure 2C and Table S3).

Significant differences were observed between Bangladesh and all other populations for both

COI and CR (Table S3). This was also the case for Sri Lanka at COI. In contrast, no

comparisons between populations in the Arabian Peninsula were found to be significant

suggesting an effect of geographic proximity. In line with this, although no significant

relationship was observed between pairwise F_{ST} and geographic distance (adjusted $R2 = 0.79$,

$P = 0.075$) there was a clear tendency for populations separated by greater distances to display

higher genetic differentiation indicating a signal of isolation by distance (Figure 2D).

Table 2. Summary of analysis of molecular variance (AMOVA) for COI and CR sequences in *R.*

ancylostomus. Variation is partitioned among and within populations and a p-value is significant if <0.05.

Source of variation	Degrees of freedom	Sum of squares	Variance components	Percentage of variation
COI				
Among populations	4	5.170	0.10593	29.07
Within populations	58	14.989	0.25843	70.93
Total	62	20.159	0.36435	
Fixation index (F_{ST})			0.29072	
p-value			<0.0001	
CR				
Among populations	4	3.607	0.06939	25.58
Within populations	60	12.116	0.20193	74.42
Total	64	15.723	0.27133	
Fixation index (F_{ST})			0.25576	
p-value			0.00098	

SNP results

Analysis of SNP data revealed a similar yet more pronounced pattern of population structure

across the sampling range. In the PCA analysis, samples from most sites clustered apart along

PC1 (Figure 3A). In particular, individuals from Oman and Saudi Arabia clustered separately

from those originating from the UAE (Persian Gulf), with a deeper split observed between

samples originating from the Red Sea, Arabian Sea and the Persian Gulf, and those originating

from Sri Lanka and Bangladesh. No clustering was observed between sampling locations along

PC2 (Figure 3A) or PC3 (Figure S3). These patterns were in part mirrored by the STRUCTURE

analysis in which an optimal value of K = 2 was inferred based on ΔK and log likelihood values

(Figure 3B, Figure S4–5). Clusters correspond to the east and west proximities of the sampling

range with all individuals from Oman and Saudi Arabia being assigned to Cluster I with q > 0.85,

and individuals from Sri Lanka and Bangladesh being assigned to Cluster II with q > 0.81.

Individuals from intermediate locations in the UAE displayed admixture between inferred

clusters (Cluster I mean q = 0.53; Cluster II mean q = 0.47) suggesting some degree of gene

flow across the range. Pairwise F_{ST} estimates ranged from 0.01–0.27 and tended to be higher

between populations separated by greater distances, in line with our mitochondrial analysis

(Figure S6). All estimates including Bangladesh had larger confidence intervals than those

without, most likely due to the small sample size of this population.

Figure 3. SNP-based population structure and diversity in *R. ancylostomus*. (A) Scatterplot showing

individual variation in principal components (PC) one and two derived from principal components

analysis. The amount of variance explained by each PC is shown in parentheses. Blue circles correspond

to individuals from Oman, pink from Saudi Arabia, green from the UAE, purple from Sri Lanka and yellow

from Bangladesh (B) Individual assignment to genetic clusters based on STRUCTURE analysis for *K* = 2

using 3,565 SNPs and 49 individuals. Each horizontal bar represents a different individual and the relative

proportion of the different colours indicate the probabilities of belonging to each cluster. Individuals are

separated by sampling locations as indicated in Figure 1. (C) Variation in multi-locus heterozygosity

among populations. Centre lines of boxplots reflect the median, bounds of the boxes extend from the first

to the third quartiles, and upper and lower whiskers reflect the largest and smallest values but no further

than 1.5 x the interquartile range from the hinge. Population abbreviations: BAN, Bangladesh; SL, Sri

Lanka; UAE, United Arab Emirates; OMAN, Oman; SA, Saudi Arabia

Genetic diversity

To explore the landscape of genetic variation, we calculated genetic diversity estimates across

sampling locations. Haplotype and nucleotide diversity at COI were highest for individuals

originating from Sri Lanka and Bangladesh (Table 4). Individuals from the UAE and Oman had

intermediate levels of diversity, while the lowest levels of diversity were observed in Saudi

Arabia where only one haplotype was detected. Similar patterns were observed at the CR

except that Bangladesh displayed the lowest variation and Saudi Arabia was intermediate.

Table 4. Genetic diversity measures for each *R. ancylostomus* sampling location calculated from COI and

CR sequences.

Indices	COI					CR				
Location	BAN	SL	UAE	OMAN	SA	BAN	SL	UAE	OMAN	SA
Number of sequences	4	9	36	10	4	4	9	37	11	4
Number of segregating sites (*S*)	2	16	3	1	0	0	3	1	1	1
Number of haplotypes (*H*)	2	6	4	2	1	1	2	2	2	2
Haplotype diversity (*Hd*)	0.50	0.83	0.50	0.47	0	0	0.50	0.43	0.33	0.50
Nucleotide diversity (*pi*)	0.0015	0.0092	0.0008	0.0007	0	0	0.0029	0.0008	0.0006	0.0010

When comparing individual heterozygosity calculated from SNP data, we detected a noticeable

decrease in variation from east to west of the sampling range (Figure 3C). In line with findings

from the mtDNA analysis, individuals originating from Saudi Arabia and Oman displayed the

lowest levels of variation, with little overlap in heterozygosity values between any of the other

populations. Individuals from Sri Lanka and Bangladesh displayed the highest levels of

variation. Unsurprisingly, the greatest variance in heterozygosity was observed in individuals

originating from the UAE, likely driven by its high sample number.

Discussion

Rhina ancylostomus are among the most threatened elasmobranch species. While some

international measures are in place to reduce pressure on this species, national and regional

management is hindered in part due to gaps in our understanding of population connectivity and

genetic variation. We used a combination of mitochondrial and SNP data to shed light on
population processes in and around the Northwest Indian Ocean. Our findings have implications
for both *in situ* and *ex situ* management and conservation planning and provide a solid baseline
with which to build our knowledge of this threatened and understudied species.

Our results first highlighted an unexpected difference in haplotype variation between COI and
the CR. Across many vertebrates, the CR is the most polymorphic region of the mitochondrial
genome (McMillan & Palumbi, 1997; Saccone et al., 1991; Zhang et al., 1995) making it a
common marker for investigating intraspecific variation (Avise, 1995). It was therefore
somewhat surprising to observe lower levels of variation at the CR than at COI across multiple
R. ancylostomus populations. However, in many teleost fish it is well established that the CR
evolves at a similar rate to the whole mitochondrial genome (Bernatchez & Danzmann, 1993;
Shedlock et al., 1992; Zhu et al., 1994), and similar patterns have more recently been observed
in other elasmobranch species (Boomer et al., 2012; Castillo-Páez et al., 2014; Dudgeon et al.,
2009). While it is possible we inadvertently targeted a conserved sequence domain, our findings
indicate this pattern could extend into numerous shark and ray families. A possible driver of low
CR variation includes functional constraints for a particular base composition and therefore
selection against transition mutations (Apostolidis et al., 1997; Bernatchez & Danzmann, 1993).
Further work comparing complete gene regions and whole mitogenomes across species and
populations will shed light on the extent and mechanisms of mitochondrial evolutionary rate
variation in elasmobranchs. Nevertheless, those interested in intraspecific variation in related
species will benefit from targeting highly variable regions of the mitochondrial genome such as
COI and NADH2 (Naylor et al., 2012).

To explore patterns of intraspecific variation, we supplemented our mtDNA with SNP genotypes.
Both marker sets highlighted some degree of genetic variation across the Northwest Indian
Ocean. This was largely driven by differences between individuals from opposite ends of the
sampling distribution, with those from adjacent sites displaying evidence for gene flow, and
those from the UAE (Persian Gulf) forming an intermediate group. Interestingly, SNP-based
structure and PCA analysis supported the presence of at least two genetically differentiated
clusters associated with sampling location. It is important to acknowledge that sample
acquisition from this endangered species is challenging, resulting in limitations to our sampling
design. In addition to the locations sampled as part of this study, *R. ancylostomus* occurs along
the coastlines of all countries in between, including eastern Saudi Arabia, Iran, India, and

Pakistan (Kyne et al., 2019, Figure 1). Uneven sampling in a continuously distributed species

can lead to the presence of artificial clusters due to spatial autocorrelation in allele frequencies

(Chambers & Hillis, 2020; Perez et al., 2018). The population structure observed in our dataset

is therefore most likely driven by gaps in our sampling distribution as opposed to true population

boundaries. We therefore cautiously anticipate that the species displays a pattern of isolation by

distance across much of the range assessed here with clinal variation in allele frequencies

being driven by intrinsic dispersal constraints.

The distribution of *R. ancylostomus* extends beyond the Northwest Indian Ocean, down the

coast of East Africa in one direction (including offshore islands such as the Seychelles and

Reunion), and across to South East Asia, Australia and the East China Sea in the other (Kyne

et al., 2019). While our study provides a significant advance in our understanding of the species,

we acknowledge that sampling constraints prevented assessment across its full range, limiting

our ability to determine the complete landscape of genetic variation. On the one hand, a pattern

of decreasing genetic similarity with increasing genetic distance may persist across the entire

range, driven by the continuous nature of the species distribution. There are a handful of

examples demonstrating isolation by distance in similarly coastal shark species, including the

lemon shark (*Negaprion brevirostris*), sandbar shark (*Carcharhinus plumbeus)* and Caribbean

reef shark (*Carcharhinus perezi,* Ashe et al., 2015; Bernard et al., 2017; Pember et al., 2023),

however, it is rare for complete distributions to have been investigated. Yet, even under such

scenarios, adaptive loci may show signatures of differentiation despite clinal variation at neutral

loci (Turbek et al., 2023). On the other hand, barriers to gene flow can lead to breakpoints in

patterns of isolation by distance. For example, deep water trenches and strong ocean currents

have been shown to restrict movements and limit gene flow in many coastal elasmobranchs

(Dudgeon et al., 2009; Hirschfeld et al., 2021; Humble et al., 2024; Schultz et al., 2008). Given

its propensity for shallow inshore environments, we anticipate barriers to gene flow beyond the

Northwest Indian Ocean will likely give rise to population differentiation in *R. ancylostomus*.

Nevertheless, one record exists of an individual caught by a purse seiner in pelagic waters

between the African continent and the Seychelles (Forget & Muir, 2021) suggesting the species

may be capable of longer distance movements. Further sampling and investigation across the

entire species range, while challenging, will be crucial to explore this in more detail.

In addition to investigating population structure, we also explored the landscape of genetic

variation among populations. Findings from SNP data, and to some extent mitochondrial

haplotypes, highlighted a decline in variation from east to west of the sampling range. Such a

pattern may have emerged as a result of species range expansion, a process which inherently

impacts diversity as repeated founder events drive the loss of low frequency alleles (Le Corre &

Kremer, 1998; Peter & Slatkin, 2013). While we cannot infer the true geographic origin of the

species, we hypothesise this may have occurred somewhere in the Indo-Malayan archipelago.

This region has extraordinary levels of biodiversity and is thought to act as a centre of origin and

survival for many marine species (Evans et al., 2016), including some coast elasmobranchs

(Maisano Delser et al., 2019; Walsh et al., 2022). With this in mind, we cautiously expect

populations further east of the sampled range to harbour higher levels of genetic diversity and

those along the east African coast to harbour lower levels. In line with this, Saudi Arabia and

Oman displayed the lowest levels of individual variation in our study. Further sampling in these

regions, particularly at range edges, will be crucial to validate our results and to explore this

hypothesis in more detail. Genetic variation is fundamental not only for buffering the intrinsic

impacts of population decline but for enabling populations to adapt to extrinsic pressures

(Bonnet et al., 2022; Lai et al., 2019). Given the intensity of fishing pressure across the Arabian

Sea and its adjacent waters (Jabado et al., 2018; Jabado & Spaet, 2017; Spaet & Berumen,

2015) our results paint a worrying picture for the future of the species in this region.

Conservation implications and future directions

Guitarfish and wedgefish are notoriously neglected in conservation management (D'Alberto,

2022; Kyne et al., 2019; Pytka et al., 2023). While some local and national fisheries

management measures can provide indirect benefits, there are few species-specific measures

in place. By advancing our understanding of *R. ancylostomus* populations, our results provide

fundamental information for guiding targeted management and conservation planning for the

species. We uncover a pattern of genetic differentiation across the Northwest Indian Ocean that

is likely explained by dispersal constraints as opposed to population isolation. Such patterns are

common in marine taxa yet make the designation of discrete management units somewhat

inappropriate (Turbek et al., 2023). Instead, we highlight the importance of maintaining habitat

connectivity across as much of the northern range of *R. ancylostomus* as possible, combined

with local and national fisheries management. The former is crucial for maximising genetic

diversity and buffering the impacts of local exploitation while the latter recognizes how fisheries

management ultimately works in practice. We also uncover regions with markedly lower genetic

variation that should warrant heightened protective measures in these areas. Recently

designated Important Shark and Ray Areas (ISRAs) in the Western Indian Ocean (Jabado et al.,

2024) not only identify these locations but partly capture the continuous coastal distribution of *R.*

ancylostomus and should therefore help kickstart the development of appropriate area-based

conservation management.

Our findings also have implications for *ex situ* management. *Rhina ancylostomus* is one of few

wedgefish species routinely held in aquarium collections across the globe (Smith et al., 2017).

Through building an atlas of genetic variation for a key part of the species range, we provide a

valuable reference for determining the breadth of variation present in the *ex situ* collection.

Due to the risks associated with small population size in captivity, such knowledge is crucial for

breeding, movement and supplementation decisions (Frankham et al., 2002; Lacy, 1987). While

there are excellent examples of well managed terrestrial collections (Farquharson et al., 2022;

Humble et al., 2023; Witzenberger & Hochkirch, 2011), the use of genetic information in *ex situ*

management of marine species significantly lags behind. *Rhina ancylostomus* therefore

presents an excellent opportunity to integrate *ex situ* and *in situ* genetic management in an

aquatic species under a One Plan approach to conservation (Pritchard et al., 2012; Redford et

al., 2012; Schwartz et al., 2017). Nevertheless, we acknowledge the potential existence of sub-

populations in areas of the species range not assessed here. Furthermore, under an isolation by

distance model, individuals at opposite ends of the distribution may display phenotypic and even

ecological differences (Burri et al., 2016; Devitt et al., 2011; Irwin et al., 2005). If *R.*

ancylostomus becomes a candidate for restoration or reintroduction, this information will be

fundamental for establishing and managing an insurance population, identifying release sites

and selecting individuals for release. We therefore strongly advocate for further sampling and

investigation – particularly around East Africa and South East Asia– in order to support holistic

and coordinated conservation management of this unique and threatened species.

Acknowledgements

We thank all the volunteers that generously provided their time to support with data collection in the United Arab Emirates and Bangladesh. We are grateful to the Department of Wildlife Conservation and the Department of Fisheries and Aquatic Resources in Sri Lanka for their advice and support, and to all the BRT researchers and volunteers for their long hours at landing sites.

Data, scripts, code and supplementary information availability

Data analysis code is available at https://github.com/mkp131/Bowmouth_analysis and https://github.com/elhumble/rhina_popgen_2024. Mitochondrial haplotypes have been deposited on NCBI (PV468213– PV468222; PV467736– PV467738). SNP genotypes and metadata are available at https://github.com/elhumble/rhina_popgen_2024.

Conflict of interest disclosure

The authors declare no conflict of interest

Funding

This work was generously supported by the Save our Seas Foundation and the Shark Conservation Fund. Sample collection in Oman and the UAE was supported by the United Arab Emirates University. Sample collection in Sri Lanka was supported by the Save Our Seas Foundation, the Shark Conservation Fund, the Marine Conservation and Action Fund (MCAF) of the New England Aquarium, the Ocean Park Conservation Foundation, Hong Kong [FH02_1920], and the Tokyo Cement Group.

References

Apostolidis, A. P., Triantaphyllidis, C., Kouvatsi, A., & Economidis, P. S. (1997). Mitochondrial DNA sequence variation and phylogeography among *Salmo trutta L.* (Greek brown trout) populations. *Molecular Ecology*, *6*(6), 531–542. https://doi.org/10.1046/j.1365-294X.1997.d01-176.x

Ashe, J. L., Feldheim, K. A., Fields, A. T., Reyier, E. A., Brooks, E. J., O'Connell, M. T., Skomal, G., Gruber, S. H., & Chapman, D. D. (2015). Local population structure and context-

dependent isolation by distance in a large coastal shark. *Marine Ecology Progress*

Series, *520*, 203–216. https://doi.org/10.3354/meps11069

Avise, J. C. (1995). Mitochondrial DNA polymorphism and a connection between genetics and

demography of relevance to conservation. *Conservation Biology*, *9*(3), 686–690.

Bailleul, D., Mackenzie, A., Sacchi, O., Poisson, F., Bierne, N., & Arnaud-Haond, S. (2018).

Large-scale genetic panmixia in the blue shark (*Prionace glauca*): A single worldwide

population, or a genetic lag-time effect of the "grey zone" of differentiation? *Evolutionary*

Applications, *11*(5), 614–630. https://doi.org/10.1111/eva.12591

Bernard, A. M., Finnegan, K. A., Pavinski Bitar, P., Stanhope, M. J., & Shivji, M. S. (2021).

Genomic assessment of global population structure in a highly migratory and habitat

versatile apex predator, the tiger shark (*Galeocerdo cuvier*). *Journal of Heredity*, *112*(6),

497–507. https://doi.org/10.1093/jhered/esab046

Bernard, A. M., Horn, R. L., Chapman, D. D., Feldheim, K. A., Garla, R. C., Brooks, E. J., Gore,

M. A., & Shivji, M. S. (2017). Genetic connectivity of a coral reef ecosystem predator:

The population genetic structure and evolutionary history of the Caribbean reef shark

(*Carcharhinus perezi*). *Journal of Biogeography*, *44*(11), 2488–2500.

https://doi.org/10.1111/jbi.13062

Bernard, A. M., Mehlrose, M. R., Finnegan, K. A., Wetherbee, B. M., & Shivji, M. S. (2025).

Connections across open water: A bi-organelle, genomics-scale assessment of atlantic-

wide population dynamics in a pelagic, endangered apex predator shark (*Isurus*

oxyrinchus). *Evolutionary Applications*, *18*(1), e70071. https://doi.org/10.1111/eva.70071

Bernatchez, L., & Danzmann, R. (1993). Congruence in control-region sequence and restriction-

site variation in mitochondrial DNA of Brook Charr (Salvelinus Fontinalis Mitchill).

Molecular Biology and Evolution, *10*(5), 1002.

https://doi.org/10.1093/oxfordjournals.molbev.a040062

Bertola, L. D., Brüniche-Olsen, A., Kershaw, F., Russo, I.-R. M., MacDonald, A. J., Sunnucks,

P., Bruford, M. W., Cadena, C. D., Ewart, K. M., de Bruyn, M., Eldridge, M. D. B.,

Frankham, R., Guayasamin, J. M., Grueber, C. E., Hoareau, T. B., Hoban, S.,

Hohenlohe, P. A., Hunter, M. E., Kotze, A., … Segelbacher, G. (2024). A pragmatic

approach for integrating molecular tools into biodiversity conservation. *Conservation*

Science and Practice, *6*(1), e13053. https://doi.org/10.1111/csp2.13053

Bonnet, T., Morrissey, M. B., de Villemereuil, P., Alberts, S. C., Arcese, P., Bailey, L. D., Boutin,

S., Brekke, P., Brent, L. J. N., Camenisch, G., Charmantier, A., Clutton-Brock, T. H.,

Cockburn, A., Coltman, D. W., Courtiol, A., Davidian, E., Evans, S. R., Ewen, J. G.,

Festa-Bianchet, M., … Kruuk, L. E. B. (2022). Genetic variance in fitness indicates rapid

contemporary adaptive evolution in wild animals. *Science*, *376*(6596), 1012–1016.

https://doi.org/10.1126/science.abk0853

Boomer, J. J., Harcourt, R. G., Francis, M. P., & Stow, A. J. (2012). Genetic divergence,

speciation and biogeography of *Mustelus* (sharks) in the central Indo-Pacific and

Australasia. *Molecular Phylogenetics and Evolution*, *64*(3), 697–703.

https://doi.org/10.1016/j.ympev.2012.05.024

Borgelt, J., Dorber, M., Høiberg, M. A., & Verones, F. (2022). More than half of data deficient

species predicted to be threatened by extinction. *Communications Biology*, *5*(1), Article

1. https://doi.org/10.1038/s42003-022-03638-9

Burri, R., Antoniazza, S., Gaigher, A., Ducrest, A.-L., Simon, C., The European Barn Owl

Network, Fumagalli, L., Goudet, J., & Roulin, A. (2016). The genetic basis of color-

related local adaptation in a ring-like colonization around the Mediterranean. *Evolution*,

70(1), 140–153. https://doi.org/10.1111/evo.12824

Casey, J., Jardim, E., & Martinsohn, J. Th. (2016). The role of genetics in fisheries management

under the E.U. common fisheries policy. *Journal of Fish Biology*, *89*(6), 2755–2767.

https://doi.org/10.1111/jfb.13151

Castillo-Páez, A., Sosa-Nishizaki, O., Sandoval-Castillo, J., Galván-Magaña, F., Blanco-Parra,

M.-P., & Rocha-Olivares, A. (2014). Strong population structure and shallow

mitochondrial phylogeny in the banded guitarfish, *Zapteryx exasperata* (Jordan Y Gilbert,

1880), from the Northern Mexican Pacific. *Journal of Heredity*, *105*(1), 91–100.

https://doi.org/10.1093/jhered/est067

Chambers, E. A., & Hillis, D. M. (2020). The multispecies coalescent over-splits species in the

case of geographically widespread taxa. *Systematic Biology*, *69*(1), 184–193.

https://doi.org/10.1093/sysbio/syz042

Clarke, S. C., Magnussen, J. E., Abercrombie, D. L., Mcallister, M. K., & Shivji, M. S. (2006).

Identification of shark species composition and proportion in the Hong Kong shark fin

market based on molecular genetics and trade records. *Conservation Biology*, *20*(1),

201–211. https://doi.org/10.1111/j.1523-1739.2005.00247.x

Cruz, V. M. V., Kilian, A., & Dierig, D. A. (2013). Development of DArT marker platforms and

genetic diversity assessment of the U.S. collection of the new oilseed crop Lesquerella

and related species. *PLOS ONE*, *8*(5), e64062.

https://doi.org/10.1371/journal.pone.0064062

D'Alberto, B. M. (2022). *Conservation biology of wedgefishes (Family Rhinidae) and giant*

guitarfishes (Family Glaucostegidae) [PhD, James Cook University].

https://doi.org/10.25903/67rg-e763

Devitt, T. J., Baird, S. J., & Moritz, C. (2011). Asymmetric reproductive isolation between

terminal forms of the salamander ring species *Ensatina eschscholtzii* revealed by fine-

scale genetic analysis of a hybrid zone. *BMC Evolutionary Biology*, *11*(1), 245.

https://doi.org/10.1186/1471-2148-11-245

Domingues, R. R., Hilsdorf, A. W. S., & Gadig, O. B. F. (2018). The importance of considering

genetic diversity in shark and ray conservation policies. *Conservation Genetics*, *19*(3),

501–525. https://doi.org/10.1007/s10592-017-1038-3

Dudgeon, C. L., Broderick, D., & Ovenden, J. R. (2009). IUCN classification zones concord with,

but underestimate, the population genetic structure of the zebra shark Stegostoma

fasciatum in the Indo-West Pacific. *Molecular Ecology*, *18*(2), 248–261.

https://doi.org/10.1111/j.1365-294X.2008.04025.x

Dulvy, N. K., Fowler, S. L., Musick, J. A., Cavanagh, R. D., Kyne, P. M., Harrison, L. R.,

Carlson, J. K., Davidson, L. N., Fordham, S. V., Francis, M. P., Pollock, C. M.,

Simpfendorfer, C. A., Burgess, G. H., Carpenter, K. E., Compagno, L. J., Ebert, D. A.,

Gibson, C., Heupel, M. R., Livingstone, S. R., … White, W. T. (2014). Extinction risk and

conservation of the world's sharks and rays. *eLife*, *3*, e00590.

https://doi.org/10.7554/eLife.00590

Evans, S. M., McKenna, C., Simpson, S. D., Tournois, J., & Genner, M. J. (2016). Patterns of

species range evolution in Indo-Pacific reef assemblages reveal the Coral Triangle as a

net source of transoceanic diversity. *Biology Letters*, *12*(6), 20160090.

https://doi.org/10.1098/rsbl.2016.0090

Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5: A new series of programs to

perform population genetics analyses under Linux and Windows. *Molecular Ecology*

Resources, *10*(3), 564–567. https://doi.org/10.1111/j.1755-0998.2010.02847.x

Farquharson, K. A., McLennan, E. A., Cheng, Y., Alexander, L., Fox, S., Lee, A. V., Belov, K., &

Hogg, C. J. (2022). Restoring faith in conservation action: Maintaining wild genetic

diversity through the Tasmanian devil insurance program. *iScience*, *25*(7), 104474.

https://doi.org/10.1016/j.isci.2022.104474

Forget, F., & Muir, J. (2021). The critically endangered bowmouth guitarfish (*Rhina*

ancylostoma) in the open ocean with an associated tuna school. *Marine Biodiversity*,

51(4), 69. https://doi.org/10.1007/s12526-021-01195-8

Frankham, R., Ballou, J., & Briscoe, D. A. (2002). *Introduction to conservation genetics*.

Cambridge University Press.

572 Funk, W. C., McKay, J. K., Hohenlohe, P. A., & Allendorf, F. W. (2012). Harnessing genomics

573   for delineating conservation units. *Trends in Ecology & Evolution*, *27*(9), 489–496.

574   https://doi.org/10.1016/j.tree.2012.05.012

575 Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). dartr: An r package to facilitate

576   analysis of SNP data generated from reduced representation genome sequencing.

577   *Molecular Ecology Resources*, *18*(3), 691–699. https://doi.org/10.1111/1755-0998.12745

578 Gumbs, R., Oenone Scott, Bates, R., Böhm, M., Forest, F., Gray, C. L., Hoffmann, M., Kane, D.,

579   Low, C., Pearse, W. D., Pipins, S., Tapley, B., Turvey, S. T., Jetz, W., Owen, N. R., &

580   Rosindell, J. (2024). Global conservation status of the jawed vertebrate Tree of Life.

581   *Nature Communications*, *15*(1), 1101. https://doi.org/10.1038/s41467-024-45119-z

582 Hirschfeld, M., Dudgeon, C., Sheaves, M., & Barnett, A. (2021). Barriers in a sea of

583   elasmobranchs: From fishing for populations to testing hypotheses in population

584   genetics. *Global Ecology and Biogeography*, *30*(11), 2147–2163.

585   https://doi.org/10.1111/geb.13379

586 Hochkirch, A., Samways, M. J., Gerlach, J., Böhm, M., Williams, P., Cardoso, P., Cumberlidge,

587   N., Stephenson, P. J., Seddon, M. B., Clausnitzer, V., Borges, P. A. V., Mueller, G. M.,

588   Pearce-Kelly, P., Raimondo, D. C., Danielczak, A., & Dijkstra, K.-D. B. (2021). A strategy

589   for the next decade to address data deficiency in neglected biodiversity. *Conservation*

590   *Biology*, *35*(2), 502–509. https://doi.org/10.1111/cobi.13589

591 Hohenlohe, P. A., Funk, W. C., & Rajora, O. P. (2021). Population genomics for wildlife

592   conservation and management. *Molecular Ecology*, *30*(1), 62–82.

593   https://doi.org/10.1111/mec.15720

594 Humble, E., Hosegood, J., Carvalho, G., de Bruyn, M., Creer, S., Stevens, G. M. W., Armstrong,

595   A., Bonfil, R., Deakos, M., Fernando, D., Froman, N., Peel, L. R., Pollett, S., Ponzo, A.,

596   Stewart, J. D., Wintner, S., & Ogden, R. (n.d.). Comparative population genomics of

manta rays has global implications for management. *Molecular Ecology*, *n/a*(n/a). https://doi.org/10.1111/mec.17220

Humble, E., Stoffel, M. A., Dicks, K., Ball, A. D., Gooley, R. M., Chuven, J., Pusey, R., Remeithi, M. A., Koepfli, K.-P., Pukazhenthi, B., Senn, H., & Ogden, R. (2023). Conservation management strategy impacts inbreeding and mutation load in scimitar-horned oryx. *Proceedings of the National Academy of Sciences*, *120*(18), e2210756120. https://doi.org/10.1073/pnas.2210756120

Irwin, D. E., Bensch, S., Irwin, J. H., & Price, T. D. (2005). Speciation by distance in a ring species. *Science*, *307*(5708), 414–416. https://doi.org/10.1126/science.1105201

Jabado, R. W. (2018). The fate of the most threatened order of elasmobranchs: Shark-like batoids (Rhinopristiformes) in the Arabian Sea and adjacent waters. *Fisheries Research*, *204*, 448–457. https://doi.org/10.1016/j.fishres.2018.03.022

Jabado, R. W., Kyne, P. M., García-Rodríguez, E., Charles, R., Armstrong, A. O., Mouton, T. L., Gonzalez-Pestana, A., Battle-Morera, A., & Rohner, C. A. (2024). *Western Indian Ocean: A regional compendium of Important Shark and Ray Areas*. IUCN SSC Shark Specialist Group and IUCN Ocean Team. https://doi.org/10.59216/ssg.isra.2023.r7

Jabado, R. W., Kyne, P. M., Pollom, R. A., Ebert, D. A., Simpfendorfer, C. A., Ralph, G. M., Al Dhaheri, S. S., Akhilesh, K. V., Ali, K., Ali, M. H., Al Mamari, T. M. S., Bineesh, K. K., El Hassan, I. S., Fernando, D., Grandcourt, E. M., Khan, M. M., Moore, A. B. M., Owfi, F., Robinson, D. P., … Dulvy, N. K. (2018). Troubled waters: Threats and extinction risk of the sharks, rays and chimaeras of the Arabian Sea and adjacent waters. *Fish and Fisheries*, *19*(6), 1043–1062.

Jabado, R. W., & Spaet, J. L. Y. (2017). Elasmobranch fisheries in the Arabian Seas Region: Characteristics, trade and management. *Fish and Fisheries*, *18*(6), 1096–1118.

Jombart, T. (2008). adegenet: A R package for the multivariate analysis of genetic markers. *Bioinformatics*, *24*(11), 1403–1405.

Kelley, J. L., Brown, A. P., Therkildsen, N. O., & Foote, A. D. (2016). The life aquatic: Advances

in marine vertebrate genomics. *Nature Reviews Genetics*, *17*(9), Article 9.

https://doi.org/10.1038/nrg.2016.66

Kilian, A., Wenzl, P., Huttner, E., Carling, J., Xia, L., Blois, H., Caig, V., Heller-Uszynska, K.,

Jaccoud, D., Hopper, C., Aschenbrenner-Kilian, M., Evers, M., Peng, K., Cayla, C., Hok,

P., & Uszynski, G. (2012). Diversity arrays technology: A generic genome profiling

technology on open platforms. *Methods in Molecular Biology (Clifton, N.J.)*, *888*, 67–89.

https://doi.org/10.1007/978-1-61779-870-2_5

Kimura, M., Maruyama, T., & Crow, J. F. (1963). The mutation load in small populations.

Genetics, *48*(10), 1303–1312.

Korneliussen, T., & Moltke, I. (2015, December 15). *NgsRelate: A software tool for estimating*

pairwise relatedness from next-generation sequencing data. Bioinformatics (Oxford,

England); Bioinformatics. https://doi.org/10.1093/bioinformatics/btv509

Kyne, P. M., Jabado, R. W., Rigby, C. L., Dharmadi, Gore, M. A., Pollock, C. M., Herman, K. B.,

Cheok, J., Ebert, D. A., Simpfendorfer, C. A., & Dulvy, N. K. (2020). The thin edge of the

wedge: Extremely high extinction risk in wedgefishes and giant guitarfishes. *Aquatic*

Conservation: Marine and Freshwater Ecosystems, *30*(7), 1337–1361.

https://doi.org/10.1002/aqc.3331

Kyne, P. M., Rigby, C. L., Dharmadi, & Jabado, R. W. (2019). *Rhina ancylostoma. IUCN Red*

List of Threatened Species, *e.T41848A124421912*.

Lacy, R. C. (1987). Loss of genetic diversity from managed populations: Interacting effects of

drift, mutation, immigration, selection, and population subdivision. *Conservation Biology*,

1(2), 143–158.

Lai, Y.-T., Yeung, C. K. L., Omland, K. E., Pang, E.-L., Hao, Y., Liao, B.-Y., Cao, H.-F., Zhang,

B.-W., Yeh, C.-F., Hung, C.-M., Hung, H.-Y., Yang, M.-Y., Liang, W., Hsu, Y.-C., Yao,

C.-T., Dong, L., Lin, K., & Li, S.-H. (2019). Standing genetic variation as the predominant

source for adaptation of a songbird. *Proceedings of the National Academy of Sciences*,

116(6), 2152–2157. https://doi.org/10.1073/pnas.1813597116

Lande, R. (1993). Risks of population extinction from demographic and environmental

stochasticity and random catastrophes. *The American Naturalist*, *142*(6), 911–927.

https://doi.org/10.1086/285580

Lande, R., & Shannon, S. (1996). The role of genetic variation in adaptation and population

persistence in a changing environment. *Evolution*, *50*(1), 434–437.

https://doi.org/10.2307/2410812

Le Corre, V., & Kremer, A. (1998). Cumulative effects of founding events during colonisation on

genetic diversity and differentiation in an island and stepping-stone model. *Journal of*

Evolutionary Biology, *11*(4), 495–512. https://doi.org/10.1046/j.1420-

9101.1998.11040495.x

Legendre, P., Fortin, M.-J., & Borcard, D. (2015). Should the Mantel test be used in spatial

analysis? *Methods in Ecology and Evolution*, *6*(11), 1239–1247.

https://doi.org/10.1111/2041-210X.12425

Leigh, J. W., & Bryant, D. (2015). POPART: full-feature software for haplotype network

construction. *Methods in Ecology and Evolution*, *6*(9), 1110–1116.

Lowe, W. H., & Allendorf, F. W. (2010). What can genetics tell us about population connectivity?

Molecular Ecology, *19*(15), 3038–3051. https://doi.org/10.1111/j.1365-

294X.2010.04688.x

Maisano Delser, P., Corrigan, S., Duckett, D., Suwalski, A., Veuille, M., Planes, S., Naylor, G. J.

P., & Mona, S. (2019). Demographic inferences after a range expansion can be biased:

The test case of the blacktip reef shark (Carcharhinus melanopterus). *Heredity*, *122*(6),

759–769. https://doi.org/10.1038/s41437-018-0164-0

McMillan, W. O., & Palumbi, S. R. (1997). Rapid rate of control-region evolution in Pacific

butterflyfishes (Chaetodontidae). *Journal of Molecular Evolution*, *45*(5), 473–484.

https://doi.org/10.1007/pl00006252

Mijangos, J. L., Gruber, B., Berry, O., Pacioni, C., & Georges, A. (2022). dartR v2: An

accessible genetic analysis platform for conservation, ecology and agriculture. *Methods*

in Ecology and Evolution, *13*(10), 2150–2158. https://doi.org/10.1111/2041-210X.13918

Mills, L. S., & Allendorf, F. W. (1996). The one-migrant-per-generation rule in conservation and

management. *Conservation Biology*, *10*(6), 1509–1518.

Mönkkönen, M., & Reunanen, P. (1999). On critical thresholds in landscape connectivity: A

management perspective. *Oikos*, *84*(2), 302–305. https://doi.org/10.2307/3546725

Moore, A. B. M. (2017). Are guitarfishes the next sawfishes? Extinction risk and an urgent call

for conservation action. *Endangered Species Research*, *34*, 75–88.

Moritz, C. (1999). Conservation units and translocations: Strategies for conserving evolutionary

processes. *Hereditas*, *130*(3), 217–228. https://doi.org/10.1111/j.1601-

5223.1999.00217.x

Naylor, G. J. P., Caira, J. N., Jensen, K., Rosana, K. a. M., White, W. T., & Last, P. R. (2012). A

DNA sequence–based approach to the identification of shark and ray species and its

implications for global elasmobranch diversity and parasitology. *Bulletin of the American*

Museum of Natural History, *2012*(367), 1–262. https://doi.org/10.1206/754.1

Nikolic, N., Devloo-Delva, F., Bailleul, D., Noskova, E., Rougeux, C., Delord, C., Borsa, P.,

Liautard-Haag, C., Hassan, M., Marie, A. D., Feutry, P., Grewe, P., Davies, C., Farley,

J., Fernando, D., Biton-Porsmoguer, S., Poisson, F., Parker, D., Leone, A., … Arnaud-

Haond, S. (2023). Stepping up to genome scan allows stock differentiation in the

worldwide distributed blue shark *Prionace glauca*. *Molecular Ecology*, *32*(5), 1000–1019.

https://doi.org/10.1111/mec.16822

Palsbøll, P. J., Bérubé, M., & Allendorf, F. W. (2007). Identification of management units using

population genetic data. *Trends in Ecology & Evolution*, *22*(1), 11–16.

https://doi.org/10.1016/j.tree.2006.09.003

Palumbi, S. R. (2003). Population genetics, demographic connectivity, and the design of marine

reserves. *Ecological Applications*, *13*(sp1), 146–158. https://doi.org/10.1890/1051-

0761(2003)013[0146:PGDCAT]2.0.CO;2

Pante, E., & Simon-Bouhet, B. (2013). marmap: A package for importing, plotting and analyzing

bathymetric and topographic data in R. *PLOS ONE*, *8*(9), e73051.

https://doi.org/10.1371/journal.pone.0073051

Pember, B. M., Chaplin, J. A., Braccini, M., & Loneragan, N. R. (2023). Population genomic and

size distribution data suggest high genetic connectivity in the sandbar shark

(*Carcharhinus plumbeus*) along a 2700 km coastline. *Fisheries Research*, *266*, 106779.

https://doi.org/10.1016/j.fishres.2023.106779

Perez, M. F., Franco, F. F., Bombonato, J. R., Bonatelli, I. A. S., Khan, G., Romeiro-Brito, M.,

Fegies, A. C., Ribeiro, P. M., Silva, G. A. R., & Moraes, E. M. (2018). Assessing

population structure in the face of isolation by distance: Are we neglecting the problem?

Diversity and Distributions, *24*(12), 1883–1889. https://doi.org/10.1111/ddi.12816

Peter, B. M., & Slatkin, M. (2013). Detecting range expansions from genetic data. *Evolution;*

International Journal of Organic Evolution, *67*(11), 3274–3289.

https://doi.org/10.1111/evo.12202

Phillips, J. D., French, S. H., Hanner, R. H., & Gillis, D. J. (2020). HACSim: An R package to

estimate intraspecific sample sizes for genetic diversity assessment using haplotype

accumulation curves. *PeerJ. Computer Science*, *6*, e243. https://doi.org/10.7717/peerj-

cs.243

Pirog, A., Ravigné, V., Fontaine, M. C., Rieux, A., Gilabert, A., Cliff, G., Clua, E., Daly, R.,

Heithaus, M. R., Kiszka, J. J., Matich, P., Nevill, J. E. G., Smoothey, A. F., Temple, A. J.,

Berggren, P., Jaquemet, S., & Magalon, H. (2019). Population structure, connectivity,

and demographic history of an apex marine predator, the bull shark *Carcharhinus*

leucas. *Ecology and Evolution*, *9*(23), 12980–13000. https://doi.org/10.1002/ece3.5597

Pritchard, D. J., Fa, J. E., Oldfield, S., & Harrop, S. R. (2012). Bring the captive closer to the

wild: Redefining the role of *ex situ* conservation. *Oryx*, *46*(1), 18–23.

https://doi.org/10.1017/S0030605310001766

Puechmaille, S. J. (2016). The program structure does not reliably recover the correct

population structure when sampling is uneven: Subsampling and new estimators

alleviate the problem. *Molecular Ecology Resources*, *16*(3), Article 3.

Pytka, J. M., Moore, A. B. M., & Heenan, A. (2023). Internet trade of a previously unknown

wildlife product from a critically endangered marine fish. *Conservation Science and*

Practice, *5*(3), e12896. https://doi.org/10.1111/csp2.12896

Redford, K. H., Jensen, D. B., & Breheny, J. J. (2012). Integrating the captive and the wild.

Science, *338*(6111), 1157–1158. https://doi.org/10.1126/science.1228899

Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J. C., Guirao-Rico, S., Librado, P., Ramos-

Onsins, S. E., & Sánchez-Gracia, A. (2017). DnaSP 6: DNA sequence polymorphism

analysis of large data sets. *Molecular Biology and Evolution*, *34*(12), 3299–3302.

https://doi.org/10.1093/molbev/msx248

Saccone, C., Pesole, G., & Sbisá, E. (1991). The main regulatory region of mammalian

mitochondrial DNA: Structure-function model and evolutionary pattern. *Journal of*

Molecular Evolution, *33*(1), 83–91. https://doi.org/10.1007/BF02100199

Schultz, J. K., Feldheim, K. A., Gruber, S. H., Ashley, M. V., Mcgovern, T. M., & Bowen, B. W.

(2008). Global phylogeography and seascape genetics of the lemon sharks (genus

Negaprion). *Molecular Ecology*, *17*(24), 5336–5348. https://doi.org/10.1111/j.1365-

294X.2008.04000.x

Schwartz, K. R., Parsons, E. C. M., Rockwood, L., & Wood, T. C. (2017). Integrating in-situ and

ex-situ data management processes for biodiversity conservation. *Frontiers in Ecology*

and Evolution, *5*. https://www.frontiersin.org/article/10.3389/fevo.2017.00120

Shedlock, A. M., Parker, J. D., Crispin, D. A., Pietsch, T. W., & Burmer, G. C. (1992). Evolution

of the salmonid mitochondrial control region. *Molecular Phylogenetics and Evolution*,

1(3), 179–192. https://doi.org/10.1016/1055-7903(92)90014-8

Smith, M., warmolts, doug, thoney, dennis, Hueter, R., Murray, M., & Ezcurra, J. (2017). *The*

Elasmobranch Husbandry Manual II - Recent Advances in the Care of Sharks, Rays and

their Relatives.

Spaet, J. L. Y., & Berumen, M. L. (2015). Fish market surveys indicate unsustainable

elasmobranch fisheries in the Saudi Arabian Red Sea. *Fisheries Research*, *161*, 356–

364. https://doi.org/10.1016/j.fishres.2014.08.022

Stanhope, M. J., Ceres, K. M., Sun, Q., Wang, M., Zehr, J. D., Marra, N. J., Wilder, A. P., Zou,

C., Bernard, A. M., Pavinski-Bitar, P., Lokey, M. G., & Shivji, M. S. (2023). Genomes of

endangered great hammerhead and shortfin mako sharks reveal historic population

declines and high levels of inbreeding in great hammerhead. *iScience*, *26*(1), 105815.

https://doi.org/10.1016/j.isci.2022.105815

Stapley, J., Reger, J., Feulner, P. G. D., Smadja, C., Galindo, J., Ekblom, R., Bennison, C., Ball,

A. D., Beckerman, A. P., & Slate, J. (2010). Adaptation genomics: The next generation.

Trends in Ecology & Evolution, *25*(12), 705–712.

Stoffel, M. A., Esser, M., Kardos, M., Humble, E., Nichols, H., David, P., & Hoffman, J. I. (2016).

inbreedR: an R package for the analysis of inbreeding based on genetic markers.

Methods in Ecology and Evolution, *7*(11), 1331–1339.

Tella, J. L., Rojas, A., Carrete, M., & Hiraldo, F. (2013). Simple assessments of age and spatial

population structure can aid conservation of poorly known species. *Biological*

Conservation, *167*, 425–434. https://doi.org/10.1016/j.biocon.2013.08.035

Turbek, S. P., Funk, W. C., & Ruegg, K. C. (2023). Where to draw the line? Expanding the

delineation of conservation units to highly mobile taxa. *Journal of Heredity*, *114*(4), 300–

311. https://doi.org/10.1093/jhered/esad011

Walsh, C. A. J., Momigliano, P., Boussarie, G., Robbins, W. D., Bonnin, L., Fauvelot, C., Kiszka,

J. J., Mouillot, D., Vigliola, L., & Manel, S. (2022). Genomic insights into the historical

and contemporary demographics of the grey reef shark. *Heredity*, *128*(4), 225–235.

https://doi.org/10.1038/s41437-022-00514-4

Waples, R. (1998). Separating the wheat from the chaff: Patterns of genetic differentiation in

high gene flow species. *Journal of Heredity*, *89*(5), 438–450.

https://doi.org/10.1093/jhered/89.5.438

Waples, R., Albrechtsen, A., & Moltke, I. (2019). Allele frequency-free inference of close familial

relationships from genotypes or low-depth sequencing data. *Molecular Ecology*, *28*(1),

35–48. https://doi.org/10.1111/mec.14954

Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R., & Hebert, P. D. N. (2005). DNA barcoding

Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *360*(1462), 1847–1857. https://doi.org/10.1098/rstb.2005.1716

Weir, B. S., & Cockerham, C. C. (1984). Estimating f-statistics for the analysis of population

structure. *Evolution*, *38*(6), 1358–1370. https://doi.org/10.2307/2408641

White, J., Heupel, M. R., Simpfendorfer, C. A., & Tobin, A. J. (2013). Shark-like batoids in

Pacific fisheries: Prevalence and conservation concerns. *Endangered Species*

Research, *19*, 277–284. https://doi.org/10.3354/esr00473

Witzenberger, K. A., & Hochkirch, A. (2011). Ex situ conservation genetics: A review of

molecular studies on the genetic consequences of captive breeding programmes for

endangered animal species. *Biodiversity and Conservation*, *20*(9), 1843–1861.

https://doi.org/10.1007/s10531-011-0074-4

Zhang, D. X., Szymura, J. M., & Hewitt, G. M. (1995). Evolution and structural conservation of

801   the control region of insect mitochondrial DNA. *Journal of Molecular Evolution*, *40*(4),

802   382–391. https://doi.org/10.1007/BF00164024

803   Zhu, D., Jamieson, B. G., Hugall, A., & Moritz, C. (1994). Sequence evolution and phylogenetic

804   signal in control-region and cytochrome b sequences of rainbow fishes

805   (Melanotaeniidae). *Molecular Biology and Evolution*, *11*(4), 672–683.

806   https://doi.org/10.1093/oxfordjournals.molbev.a040146