The Origin of Movement Biases During Reaching

Tianhe Wang[1,2,*], Ryan J. Morehead[3], Jonathan S. Tsay[4,†,*], Richard B. Ivry[1,2,†],

[1]Department of Psychology, University of California, Berkeley

[2]Helen Wills Neuroscience Institute, University of California, Berkeley

[3]School of Psychology, University of Leeds

[4]Department of Psychology, Carnegie Mellon University

[†]Equal contribution as senior authors.

*Corresponding Authors: tianhewang@berkeley.edu, xiaotsay2015@gmail.com

Abstract

Goal-directed movements can fail due to errors in our perceptual and motor systems. While these errors

may arise from random noise within these sources, they also reflect systematic motor biases that vary

with the location of the target. The origin of these systematic biases remains controversial. Drawing on

data from an extensive array of reaching tasks conducted over the past 30 years, we evaluated the merits

of various computational models regarding the origin of motor biases. Contrary to previous theories, we

show that motor biases do not arise from systematic errors associated with the sensed hand position

during motor planning or from the biomechanical constraints imposed during motor execution. Rather,

motor biases are primarily caused by a misalignment between eye-centric and the body-centric

representations of position. This model can account for motor biases across a wide range of contexts,

encompassing movements with the right versus left hand, proximal and distal effectors, visible and

occluded starting positions, as well as before and after sensorimotor adaptation.

Main

Accurate movements are crucial for everyday activities, affecting whether a glass is filled or spilled, or whether a dart hits or misses the target. Some of movement errors arise from sensorimotor noise, including visual noise regarding the location of targets and effectors[1–3], planning noise introduced when issuing a motor command, and neuromuscular noise when executing a movement[4–6].

In addition, some of these errors arise from systematic biases that vary across the work space[7–10]. The origin of these biases remains controversial (Fig 1a): Whereas some studies postulate that motor biases stem from systematic distortions in perception[7,10–13], others posit that biases originate from inaccurate motor planning and/or biomechanical constraints associated with motor execution[14–16]. In the following section, we provide an overview of current models of systematic motor biases as well as outline a novel hypothesis, setting the stage for a re-analysis of published data and presentation of new experimental results.

Starting at the input side, motor biases may arise from systematic distortions in the representation of the perceived target position (Fig 1b). A prominent finding in the visual cognition literature is that the remembered location of a visual stimulus is biased towards diagonal axes[12,13,17]. That is, the reported visual location of a stimulus is shifted towards the centroid of each quadrant. This perceptual bias does not depend on the method of response, as this phenomenon can be observed when participants point to a cued location or press a key to indicate the remembered location of a briefly presented visual target[12,13]. While this literature has emphasized that this form of perceptual bias arises from processing within visual working memory, it is an open question whether it contributes to goal-directed reaching when the visual target remains visible.

Another perceptual cause of motor biases stems from proprioception (Fig 1c). Systematic distortions in

the perceived position of the hand[18–20] and/or joint position[21,22] can influence motor planning. For example,

if the perceived starting position of the hand is leftwards of its true location, a reaching movement to a

forward visual target would exhibit a rightward bias and a reaching movement to a rightward visual target

would fall short[7]. A proprioceptive perceptual bias at the starting position has been reported to be the

major source of bias in many previous studies.[18–22]

Whereas the preceding models have considered how distortions of visual or proprioceptive space might,

on their own, lead to reaching biases, motor biases could also originate from a misalignment in the

mapping between perceptual and motor reference frames. Based on classic theories of motor planning[23],

the start position and the target position are initially encoded in an eye-centric visual coordinate frame,

and then transformed to representations in a body-centric proprioceptive coordinate frame within which

the movement is planned (Fig 1d). Motor biases could arise from systematic distortions that occur during

this visuo-proprioceptive transformation process[24–26]. Indeed, when participants are required to match

the position of their unseen hand with a visual target, systematic transformation biases are observed

across the workspace (Fig 1d; also see Methods)[27,28]. In the current study, we develop a novel

computational model to capture how these transformation biases should result in systematic biases

during reaching.

On the output side, it has been posited that reaching biases could arise from biomechanical factors that

impact movement execution[8]. Specifically, movements may be biased toward trajectories that minimize

inertial resistance and/or energetic costs[15,16,29,30]. For example, minimizing energy expenditure would

result in biases towards trajectories that minimize resistive forces or changes in joint angles[14,31]. Moreover,

inaccuracies in the internal model of limb dynamics could produce systematic execution biases. For

example, underestimating the weight of the limb would result in reaches that overshoot the target[14,32].

To determine the origin of motor biases, we formalized four computational models to capture the

potential sources described above. As detailed in the Results section, the models predict distinct motor

bias patterns in a center-out reaching task (Fig 1f-h). While prior research has focused on the impact of

individual sources (e.g., vision or proprioception) on the pattern of motor errors, these studies often entail

a limited set of contexts (e.g., reaching behavior only when the start position is visible or only with the

right hand). However, looking across studies, the task context can result in dramatically different motor

bias patterns; indeed, when plotted in polar coordinates across the workspace, the bias functions range

from having single peak[7,10,21] to quadruple peaks[12,13,17]. This diversity underscores the absence of a

comprehensive explanation for motor bias phenomena across different experimental designs and setups.

Additionally, a notable limitation of earlier work is the reliance on small participant cohorts (n<10) and a

restricted number of targets (typically 8). The sensitivity of such experiments is limited in terms of their

capacity to discriminate between models.

To better evaluate sources of motor bias during reaching, we report a series of experiments involving a

range of contexts, designed to test predictions of the different models. We compared movements

performed with the right or left hand, proximal or distal effectors, under conditions in which the start

position was either visible or not visible, and before and after implicit sensorimotor adaptation. To

increase the power of model comparisons, we measured the motor bias function at a higher resolution

(24 targets) and in a bigger sample size (n >50 per experiment).

Figure 1. Different Causes of Motor Biases. (a) Motor biases may originate from biases in perceiving the initial hand position (proprioceptive bias), perceiving the location of the visual target (visual bias), transforming positional information from visual to proprioceptive space (transformation bias), and/or biomechanical constraints during motor execution. Previous models attribute motor biases to errors originating from the distinct contributions of visual (b) and proprioceptive biases (c). (d) Our model attributes motor biases to a transformation error between visual and proprioceptive coordinate systems. (e) A visuo-proprioceptive map showing the matching error between proprioceptive and visual space (Wang et al (2020)). Participants matched the position their hand (tip of the arrow) from a random starting location to the position of a visual target (end of the arrow). The blue dot depicts an example of a visual target in the workspace, and the red arrow indicates the corresponding matched hand position. Participants were asked to maximize spatial accuracy rather than focus on speed. (f-h) Model bias functions

predicted by four models. Top: Illustration of how each model is applied to a center-out reaching task. As an example,

the predicted motor plan and the corresponding real movement are provided for the 100° target in f and 135° target

in g and h. Bottom: The predicted motor bias functions qualitatively differ in terms of the number of peaks and

toughs. Note that the middle panel depicts two variants of a proprioception model.

Results

<u>Motor biases across the workspace</u>

To examine the pattern of motor biases during goal-directed movements, participants performed a

center-out reaching task with their right hand (Fig 2a). We ran two versions of the study in Experiment 1.

In Exp 1a, we used an 8-target version similar to that used in most previous studies[7–10,21]. To obtain better

resolution of the motor bias pattern, we also conducted a 24-target version in Exp 1b. Within each

experiment, participants first performed the task without visual feedback to establish their baseline bias

and then a block with veridical continuous feedback to examine how feedback influences their biases.

Motor biases were calculated as the angular difference between the target and hand when the movement

amplitude reached the target distance (Fig 2b), with a positive error indicating a counterclockwise bias

and a negative error indicating a clockwise error.

Across the workspace, the pattern of motor biases exhibited a two-peaked function (Fig 2c) characterized

by two peaks and two troughs. From the 8-target experiment, larger biases were observed for the diagonal

targets (45°, 135°, 225°, 335°) compared to the cardinal targets (0°, 90°, 180°, 270°)[33,34]. In terms of

direction, reaches to diagonal targets were biased toward the vertical axis, and reaches for cardinal targets

were biased in the counterclockwise direction. This pattern is similar to what has been observed in

previous studies for right-handed movements[8,9]. With the higher resolution in the 24-target experiment,

we see that the peaks of the motor bias function are not strictly aligned with the diagonal targets but are

shifted towards the horizonal axis. Moreover, the upward shift of the motor bias function relative to the

horizontal line suggests that clockwise biases are more prevalent compared to counterclockwise biases

across the workspace.

Figure 2. Motor biases across different experimental contexts. (a) Lab-based experimental apparatus for Exps 1-2.

(b) Vectors linking the start position to the average endpoint position when reach amplitude equaled the target

radius (pink lines; Exp 1a). (c) Motor biases as a function of target location. The dots indicate the mean angular error

across participants during the no-feedback block (pink) and veridical feedback block (grey). The pattern of motor

bias was similar in Exp 1a (8-targets; left panel) and Exp 1b (24-targets; right panel), characterized by two peaks and

two troughs. Error bars denote standard error. (d) Motor biases generated during left hand reaches (left), left-hand

results when the data are mirror reversed across the vertical meridian (middle), and right-hand reaches (right). (e)

Left: The motor bias generated by right-hand reaches was similar to that of mirror-reversed left-hand reaches. Right:

Difference in RMSE when the right-hand map is compared to the original left-hand map relative to when the right-

hand map is compared to the mirror reversed left-hand map. Positive values indicate better data alignment when

the left hand data are mirror-reversed. (f) Correlation of the motor bias function between the no-feedback and

feedback blocks is higher in the within-participant condition compared to the between-participant condition. Gray

bars indicate the 25% and 75% quartiles. White dots indicate the median and horizontal lines indicate the mean. (g)

Experimental setup for Exp 3. Participants were asked to make center-out reaching movements using a trackpad or

mouse. These movements predominantly involve finger and wrist movements. (h) The workspace is presumed to be

closer to the reference point (e.g., left shoulder) for finger/wrist movements (Exp 3) compared to that of arm

movements (Exp 1-2). (i) The pattern of motor biases in finger/wrist movements for 8-targets (left) and 24-targets

(right).

<u>Motor biases primarily emerge from a misalignment between visual and hand reference frames</u>

We developed a series of models to capture how systematic distortions at different sensorimotor

processing stages would cause systematic motor biases (Fig 3a). Here we consider processing associated

with the perceived position of the target, the perceived position of the arm/hand, and planning processes

required to transform a target defined in visual space to a movement defined in arm/proprioceptive space.

Biases could also arise from biomechanical constraints. Given that biomechanical biases are not easily

simulated, we will evaluate this hypothesis experimentally (see below).

We implemented four single-source models to simulate the pattern of the motor bias that would be

predicted in a center-out reaching task (Fig 1f-h; see Methods). For the Visual Bias model, we assumed

that the representation of the visual targets were biased towards the diagonals[13]. This model predicts a

four-peaked motor bias function (Fig 1f). For the Proprioceptive Bias model, we considered two variants

building on the core idea is that the perceived starting position of the hand is distorted: A Vector-Based

model in which the motor plan is a vector pointing from the perceived hand position to the target[7,10] and

a Joint-Based model in which the movement is encoded as changes in the shoulder and elbow joint angles

to move the limb from a start position to a desired target location[21,22] (See Fig S1). Importantly, both

models predict a motor bias function with a single peak (Fig 1g). Taken together, models that focus on

systematic distortions of perceptual information do not qualitatively capture the observed two-peaked

motor bias function (Fig 2c).

The fourth model, the Transformation model is based on the idea that the start and target positions are

initially encoded in visual space and transformed into proprioceptive space for motor planning[23]. Motor

biases may arise from a transformation error between coordinate systems. Two prominent features stand

out when this transformation error is empirically measured[18–20,27]. First, the direction of the

transformation error is similar across the workspace (e.g., a leftward and downward matching error for

right-handers). Second, the magnitude of the error increases with distance from the body (Fig 1d). As such,

we simulated a visuo-proprioceptive error map by using a leftward and downward error vector with the

magnitude scaled across the workspace based on the distance of each location to a reference point (Fig

1h Top). This model predicts a two-peaked motor bias function (Fig 1h Bottom), a shape strikingly similar

to that observed in Exps 1a and 1b.

To quantitively compare the models, we fit each model with the data in Exp 1b. The Transformation Bias

model provided a good fit of the two-peaked motor bias function (R^2=0.84, Fig 3a left, see Table S1 for

parameters). Fig 1h (top) shows the recovered visual-proprioceptive bias map based on the parameters

of the Transformation Bias model when fit to the reaching data in Exp 1b. The simulated results are very

similar to the map measured empirically in the previous study[20] (Fig 1e). In contrast, the Visual Bias and

Proprioceptive Bias models provide poor fits (all R^2<0.18, Fig 3a right). Thus, the simulations suggest that

motor biases observed in reaching primarily originate from a transformation between visual and

proprioceptive space.

A second way to evaluate the models is to compare the motor bias functions for the left and right hands.

The Proprioceptive and Transformation Bias models predict that the bias function will be mirror-reversed

for the two hands whereas the Visual Bias model predicts that the functions will be superimposed. We

compared the functions for right and left hand reaches in Exp 2 using the 8-target layout. We found that

the dissimilarity (RMSE) between the pattern of motor biases across two hands significantly decreased

when the left-hand data are mirror-reversed compared to when the bias patterns are compared without

mirror reversal (t(78) = 2.7, p = 0.008, Fig 2d-e). These results are consistent with the Transformation Bias

model and provide further confirmation that the Visual Bias model, at least as conceptualized here, fails

to provide a comprehensive account of reaching biases.

While the overall pattern of biases in the visuo-proprioceptive map are similar across individuals, there

are subtle individual differences[18,20]. As such, we would anticipate that the motor bias function would also

exhibit stable individual differences due to idiosyncrasies in the visuo-proprioceptive map. To examine

this, we correlated the bias functions obtained from blocks in which we either provided no visual feedback

or veridical endpoint feedback. The magnitude of the biases was attenuated when endpoint feedback was

provided, likely because the feedback reduced the visuo-proprioceptive mismatch. Nonetheless, the

overall pattern of motor bias was largely preserved, with the within-participant correlations (Exp 1a:

$r_{norm} = 0.999$, Exp 1b: r_{norm} = 0.974) significantly higher than the averaged between-participant

correlation in both Exp 1a and Exp 1b (Fig 2f).

<u>Visual bias also contributes to the motor bias.</u>

In the preceding section, we considered each model in isolation, testing the idea that motor biases arise

from a single source. However, the bias might originate from multiple sources. For example, there could

be a distortion in both vision and proprioception, or a visuo-proprioceptive transformation that operates

on distorted inputs. To address this, we evaluated hybrid models by combining the Visual Bias model with

the Proprioceptive or Transformation Bias models. Although theoretically plausible, we did not consider

a hybrid of the Proprioceptive and Transformation Bias models since they conflict in terms of whether the

start position is perceived visually or proprioceptively.

The hybrid model that combines the Transformation and Visual Bias models (T+V model) provided an

excellent fit of the motor bias pattern in Exp 1b (R^2=0.973, Fig 3a). Based on a comparison of BIC values,

this model not only outperforms the other hybrid models, but also significantly improved the fit compared

to the Transformation Bias model alone. These results are especially interesting in that the assumed visual

bias towards the diagonal axes has only been shown in studies in which perception was tested after the

target had been extinguished. The current results suggest that this bias is also operative when the target

remains visible, suggesting that the visual bias may reflect a general distortion in how space is represented,

rather than a distortion that arises as information is processed in visual working memory.

Figure 3. The pattern of motor biases is best explained by assuming systematic distortions in the perceived location of the target and the transformation between visual-proprioceptive coordinate frames. (a) For single-source models, the pattern of motor biases in the no feedback block of Exp 1a (pink dots) is best fit by the Transformation Bias model (left) compared to the other models (right). (b) A mixed model with transformation and visual biases (T+V) provides the best fit to the motor bias function in both Exp 1b (top) and Exp 3b (bottom). (c) Model comparison using BIC. ΔBIC values are provided by subtracting the BIC from the best performing model (i.e., the T+V model). A smaller ΔBIC signifies better model performance.

To further evaluate the T+V model, we examined its performance in explaining the motor bias function

obtained in an on-line study (Exp 3) in which participants performed the center-out task by moving a finger

across a trackpad. One major difference between the in-person and on-line setups is that the workspace

is much smaller and closer to the body when participants use a trackpad (Fig 2g). As such, the magnitude

of the motor biases generated by transformation errors should be smaller in the online compared to the

in-person setup (Fig 2h).

Consistent with the prediction of the T+V model, we found markedly smaller motor biases with the online

setup (Exp 3) compared to the in-person setup (Exp 1) (Fig 2c). While the motor bias functions were similar

across experiments, we observed two small peaks between 20° and 200° in Exp 3 that were not apparent

in Exp 1. When we fit this function to single source models, the Visual Bias model outperformed the

Transformation Bias model. This suggests that, when the movements are close to the body, visual biases

make a relatively stronger contribution to the motor biases compared to transformation biases.

Nonetheless, the T+V model again provides the best fit to the motor bias function (R^2=0.857, Fig 3e, see

Table S1 for parameters), significantly outperforming the other alternatives including the Visual Bias

model (Fig 3f).

<u>Transformation model accounts for qualitative changes in the motor bias function</u>

The Transformation Bias model assumes that, for normal reaching, both the start and the target positions

are encoded in visual space before being transformed into proprioceptive space for motor planning.

However, if the start position is not visible, then the sensed start position would be directly encoded in

proprioceptive space (i.e., where the hand is positioned), bypassing the need for a transformation

between coordinate frames. As such, biases arising from the transformation process would only arise

when the input is limited to the perceived position of the visual target. When we simulated the scenario

in which the start position is not visible, the Transformation Bias model predicts a single-peaked function

(Fig 4a right), a qualitative change from the two-peaked function predicted when both the start position

and target position are visible (Fig 4a left).

Figure 4. Motor bias patten changes when the start position is not visible. (a) Schematic showing the planned

movement under the Transformation Bias model when the start position is either visible (left) or not visible (right).

In the latter case, only the target position has to be transformed from visual to proprioceptive coordinates with the

start position directly encoded in proprioceptive space. The T+V model now predicts a single-peaked motor bias

function (lower row). (b) Consistent with this prediction, a two-peaked function is predicted when the start position

is visible (as in Exp 1) and a single-peaked function is predicted when start position is not displayed. Data (pink dots)

are from Vindras et al (2005).

To test this idea, we re-examined data from previous studies in which the participant's hand was passively

moved to a start position with no visual information given about the start location or hand position[7,10,21].

Strikingly, the motor bias function under this condition has only have one peak. Thus, the transformation

Bias model provides a novel account of the difference in motor biases observed when the start position

is visible (Exp 1-3) compared to when it is not visible.

We note that the one-peaked motor bias function has previously been interpreted as evidence in support

of a Proprioceptive Bias model (Fig 1f)[7,10,21]. We performed a model comparison on the data from one of

these studies[10] and the T+V outperformed the Proprioceptive Bias, as well as the P+V models (ΔBIC=10.9).

In addition, only the T+P model is able to account for the asymmetry between clockwise and

counterclockwise biases. In summary, these results suggest that motor biases when reaching from an

unseen start position arise when the target position is transformed from visual to proprioceptive

coordinates rather than from a proprioceptive bias impacting the sensed start position. Moreover, the

T+V model provides a parsimonious account of the bias functions, independent of the visibility of the start

position.

Another way to evaluate the Transformation model is to perturb the position of the visual start position

relative to the real hand position. Under this manipulation, a single peaked motor bias function is

observed (Fig S2) [21,22]. Interestingly, the functions exhibit opposing phase shifts when the starting position

is perturbed to the left versus to the right (Fig S2). This qualitative change in the motor bias function can

again be successfully captured by the Transformation model (Fig S2c). Taken together, these data provide

strong evidence favoring the notion that motor biases originate primarily from a misalignment between

visuo-proprioceptive reference frames.

<u>Biomechanical models fail to account for motor biases</u>

An alternative account of motor biases is that they arise from biomechanical constraints associated with

upper limb movements. For example, movement kinematics have been explained in terms of cost

functions to minimize energy and/or minimize jerk,[35] constraints that may result in an increase in endpoint

error[15,16,29]. One argument against a biomechanical model comes from evidence provided in the previous

section: Biomechanical constraints associated with movement execution would not predict qualitative

changes in the motor bias function in response to a visual manipulation of the start position.

As a second comparison of the T+V and Biomechanical Bias models, we examined how motor biases

change after the sensorimotor map is recalibrated following a form of motor learning, implicit

sensorimotor adaptation. Here we re-analyzed the data from previous experiments that had used a

perturbation technique in which the visual feedback was always rotated by 15° from the target,

independent of the hand position (Fig 5a, non-contingent clamped feedback[36]). Participants adapt to this

perturbation, with subsequent reaches to the same target shifted in the opposite direction (Fig 5b),

reaching an asymptote of around 20° and showing a robust aftereffect when the perturbation is removed.

Participants are unaware of their change in hand angle in response to clamped feedback, reporting their

perceived hand position to be close to the target[37].

Figure 5. The pattern of motor bias is preserved after implicit sensorimotor adaptation, consistent with the Transformation + Visual Bias model. (a) Illustration of the clamped perturbation. Feedback cursor is offset by a fixed angle from the target, independent of the participant's heading direction. (b) Time course of hand angle in response to clockwise or counterclockwise clamped feedback. Vertical lines demarcate the perturbation block which was preceded and followed by no-feedback baseline and washout phases, respectively (gray areas). Shaded area indicates standard error. (c) Predictions for the bias functions after adaptation for the T+V (top) and Biomechanical models (bottom). See text for details. The right column shows the predicted motor bias functions following adaptation in response to a clockwise (CW) or counterclockwise (CCW) clamp. (d) Motor bias functions before and after training in a CW (left) and a CCW (right) clamp. Data taken from Morehead et al. (2017) and Kim et al. (2018);

the height of the colored bars indicates the standard error for each data point. The best-fit lines for the T+V model

are shown. (e) Parameter values to capture vertical and horizontal shifts in motor bias functions before and after

training. The CW and CCW conditions both showed a significant vertical shift but no horizontal shift.

For the T+V model, the transformation between visual and proprioceptive space depends on the

perceived positions of the start and target locations in a visual-based reference space, one that remains

unchanged before and after adaptation. We assume that adaptation has changed a sensorimotor map

that is referenced after the transformation from visual to proprioceptive space. As such, the heading

angle after adaptation for each target location is obtained by summing the motor biases for that target

location and the extent of implicit adaptation. This would result in a vertical shift of the motor bias

function (Fig 5c top).

In contrast, the biomechanical model predicts that motor biases will be dependent on the actual

movement direction rather than the target location (e.g., a bias towards a movement that is energetically

efficient). Since the mapping between a target location and its corresponding reach direction is rotated

after adaptation, the motor bias pattern would also be rotated (Fig 5c bottom). As such, the biomechanical

model predicts that the motor bias function will be shifted along both the horizontal and vertical axes.

To arbitrate between these models, we analyzed the data from two previous studies, looking at the bias

function from no-feedback trials performed before (baseline) and after adaptation (washout) [36,38].

Consistent with the prediction of the T+V model, the motor bias function shifted vertically after adaptation

(Fig 5d) but did not shift horizontally.

To quantitatively evaluate these results, we first fit the motor bias function during the baseline phase with

the T+V model and fixed the parameters. We then examined the heading angles during the aftereffect

phase by fitting two additional parameters, one that allowed the function to shift vertically (v) and the

other to allow the function to shift horizontally (h). The T+V model predicts that only v will be different

than zero; in contrast, the Biomechanical model predicts that h and v will both be different than zero and

should be of similar magnitude. The results clearly favored the T+V model (Fig 5e). The vertical shift in the

bias functions was of a similar magnitude as the aftereffect, with the shift direction depending on the

direction of the clamped feedback (v: CW: 12.5°; CCW: -12.2°, p<0.001). In contrast, the best fitting value

for h was not significantly different from zero in both condition (Fig 5g). These results are consistent with

the hypothesis that visual representations are first transformed into proprioceptive space for motor

planning, with the recalibrated sensorimotor map altering the trajectory selected to achieve the desired

movement outcome.

Discussion

While motor biases are ubiquitous in goal-directed reaching movements, the origin of these biases has

been a subject of considerable debate. We addressed this issue by characterizing these biases across a

range of experimental conditions, and evaluated a set of computational models derived to capture

different possible sources of bias. Contrary to previous theories, our results indicate that motor biases do

not stem from a distortion in the sensed position of the hand[7,10,21,22] or from biomechanical constraints

during movement execution[15,16,29]. Instead, motor biases appear to arise from systematic distortions in

perceiving the location of the visual target and the transformation required to translate a perceived visual

target into a movement described in proprioceptive coordinates.[24–26] Strikingly, our model successfully

accounts for sensorimotor biases across a wide range of contexts, encompassing movements performed

with either hand as well as with proximal and distal effectors. Our model also accounts for the qualitative

changes in the motor bias function that are observed when vision of the starting position of the hand is

occluded, and when the sensorimotor map is perturbed following implicit adaptation.

While motor biases have been hypothesized to reflect a mismatch across perceptual and motor

coordinate systems,[25,26] it is unclear what information is transformed and what reference frame is

employed for motor planning. Interestingly, many previous studies posit that movement is planned in an

eye-centric visual reference frame[39–41]. While the target can be directly perceived in this reference space,

the start position of the hand would need to be transformed from a proprioceptive reference frame to a

visual one. Systematic error in this transformation would mean that the start position of the hand is

inaccurately represented in visual space, resulting in motor biases[7,25]. This idea underlies the

Proprioceptive Bias models described in this paper.

In contrast to these models, our Transformation Bias model posits that movement is planned in a hand-

centric proprioceptive reference frame. By this view, when both the target and start position are provided

in visual coordinates, the sensorimotor system transforms these positions from visual space to

proprioceptive space. Systematic error in this transformation process will result in motor biases. When

vision of the start position is available, the Transformation Bias model successfully accounts for the two-

peaked motor bias function (Exp 1), whereas the Proprioceptive Bias model in which motor planning is

eye-centric predicts only a single peak in the function.

Perhaps most compelling, the Transformation Bias model accounts for how the pattern of motor biases

change when the visibility of the start position is manipulated. When visible, the motor bias function is

two-peaked. In contrast, when the start position is occluded, the transformation from visual to

proprioceptive space is only relevant for the target position since the start position of the hand is already

represented in proprioceptive space. Here the model predicts a motor bias function with a single peak, a

function that has been observed in previous studies[7,10].

We note that there is a third scenario, one which both the start position and target position are provided

in proprioceptive space. We predict that under this condition, motor biases originating from the visuo-

proprioceptive transformation would completely disappear. Indeed, when the hand is passively moved

first to the target location and then to the start position, subsequent reaches to the target do not show

the signature of bias from a visuo-proprioceptive transformation.[12] Instead, the reaches exhibited a bias

towards the diagonal axes, consistent with the predicted pattern if the sole source of bias is visual.

Why would a sensorimotor system exhibit inherent biases during the transformation process? We

propose that these biases arise from two interrelated factors. First, these systems are optimally tuned for

distinct purposes: A body-centric system predominantly uses proprioceptive and vestibular inputs to

determine the orientation and position of the body in space, while an eye-centric system relies on visual

inputs to interpret the layout of objects in the external world, representations that should remain stable

even as the agent moves about in this environment.[42,43] Second, these sensory systems consistently

receive information with very different statistical distributions[2,44], perhaps because of these distinct

functions. For example, visual inputs tend to cluster around the principal axes (horizontal and vertical)

[45,46], whereas proprioceptive information during reaching is clustered around diagonal axes[47]. This is

because these movements are often the least effortful and are the most frequently enacted directions of

movement.[48]. The differences in computational goals and input distributions might have led to natural

divergences in how each system represents space[49], and consequently, result in a misalignment between

the reference frames.

The Transformation Bias model addresses how biases arise when the information is passed along from a

visual to a proprioceptive reference frame. However, the results indicate that another source of bias

originates from a distortion within the visual reference frame itself, manifesting as an attractive bias

towards the diagonal axes. Thus, the best fitting model posits two sources of bias, one related to the

representation of the visual target and a second associated with the transformation process. This hybrid

Transformation + Visual Bias model outperformed all single-source and hybrid models, providing an

excellent fit of the behavioral data across a wide variety of contexts.

There are several hypotheses concerning the origin of this bias. One account has focused on the idea that

these biases arise from distortions introduced in visual working memory. The biases are observed when

participants report the location of a remembered visual target[13,17,50], independent of the reporting

method (pointing or keypresses)[17]. However, as mentioned above, similar biases are observed even when

participants report the location of a proprioceptive target (i.e., participants match the position of their

unseen hand).[12] Moreover, as shown in our study, diagonal target biases make a sizable contribution even

when the visual target is always present, imposing no demands on working memory. As such, we postulate

that the visual biases may reflect a more domain-general distortion of spatial representations.

Our data suggest that biomechanical factors do not significantly impact motor biases. While we did not

formalize a biomechanical model, we provided several lines of evidence suggesting the biomechanical

factors have minimal influence on the pattern of motor biases. For example, it is hard to envision a

biomechanical model that would account for the qualitative change in the bias function when the start

position was visible (two-peak function) to when it was hidden (one-peak function).

Empirically, we evaluated biomechanical contributions to motor biases by examining the bias pattern

observed before and after implicit sensorimotor adaptation. We assume that adaptation mainly modifies

a sensorimotor map[51] but has a relatively smaller influence on a visuo-proprioceptive map[28,37,52]. That is,

adaptation may change the mapping between a target represented in the proprioceptive space and the

motor commands required to reach that location. Given that a biomechanical model assumes that motor

biases are associated with the direction of a movement, this model would predict that the pattern of

motor biases would be distorted by implicit motor adaptation. At odds with this prediction, the pattern

of motor biases remained unchanged after adaptation, a result consistent with the Transformation Bias

model.

Nonetheless, the current study does not rule out the possibility that biomechanical factors may cause

motor biases in different contexts. In our study, biomechanical constraints may not be influential since

the extent of the required movements was relatively modest and involved minimal interaction torques.

Moreover, we focused on examining biases that manifest at the movement endpoint rather than in the

movement trajectory where biomechanics are known to play a greater role.[15,16] Future studies are

necessary to explore whether biomechanical biases are more pronounced in contexts in which the limb

dynamics are crucial and/or the movements are energetically costly.

Methods

<u>Participants</u>

For the lab-based study (Exp 1, 2), 206 undergraduate students (age: 18-24) were recruited from

University of California, Berkeley. For the online study (Exp 3), 183 young adult participants (age: 18-30)

were recruited via Prolific, a website designed to recruit participants for online behavioral testing. All

participants were right-handed as assessed by the Edinburgh handedness test[53] with normal or corrected-

to-normal vision. Each participant was paid $15/h. The protocol was approved by the institutional review

board at the University of California Berkeley.

<u>Procedure</u>

Experiments 1a, 1b, and 2:

Experiments 1a, 1b, and 2 were conducted in the lab. Participants performed a center-out reaching task,

holding a digitizing pen in the right or left hand to make horizontal movements on a digitizing tablet

(49.3cm x 32.7cm, sampling rate= 100 Hz; Wacom, Vancouver, WA). The stimuli were displayed on a 120

Hz, 17-in. monitor (Planar Systems, Hillsboro, OR), which was mounted horizontally above the tablet (25

cm), to preclude vision of the limb. The experiment was controlled by custom software coded in MATLAB

(The MathWorks, Natick, MA), using Psychtoolbox extensions, and run on a Dell OptiPlex 7040 computer

(Dell, Round Rock, TX) with Windows 7 operating system (Microsoft Co., Redmond, WA).

Participants made reaches from the center of the workspace to targets positioned at a radial distance of

8 cm. The start position and target location were indicated by a white annulus (1.2 cm diameter) and a

filled blue circle (1.6 cm), respectively. Vision of the hand was occluded by the monitor, and the lights

were extinguished in the room to minimize peripheral vision of the arm. Feedback, when provided, was

in the form of a 4 mm white cursor that appeared on the computer monitor, aligned with the position of

the digitizing pen.

To start each trial, the participant moved the cursor to the start circle (5mm diameter). After maintaining

the cursor within the start circle for 500 ms, a target appeared at one of the target locations. The

participant was instructed to make a rapid slicing movement through the target. We did not impose any

reaction time guidelines, allowing the participant to set their own pace to initiate the movement. On no-

feedback trials, the cursor was blanked when the hand left the start circle, and the target was extinguished

once the radial distance of the movement reached the target distance (8 cm). On feedback trials, the

cursor was visible throughout the movement until the movement amplitude reached 8 cm; at that point,

its position was frozen for 1 s, providing feedback of the accuracy of the movement (angular position with

respect to the target). After this interval, the target and cursor were extinguished.

At the end of both the no-feedback and feedback trials, a white ring appeared denoting the participant's

radial distance from the start position. This ring was displayed to guide the participant back to the start

position without providing angular information about hand position. Once the participant moved within

2 cm of the start position, the ring was extinguished, and a veridical cursor appeared to allow the

participant to move their hand to the start position. Participants had to remain within the start position

for 200 ms before the next target appeared. If the amplitude of the hand movement did not reach the

target (<8 cm radial distance) within 300 ms, the message "too slow" would be displayed for 500 ms before

the white ring appeared.

For Exp 1a and Exp 2, there were 8 target locations, evenly spaced in 45° increments around the workspace

(primary axes and main diagonals). For Exp 1b, there were 24 target locations, evenly spaced in 15°

increments. Each experiment consisted of a no-feedback block followed by a feedback block. There were

5 trials per target (40 trials total) for each block in the Exps 1a. There were 4 trials per target (96 trials

total) in Exp 1b.

Experiments 3a and 3b

Exps 3a and 3b were conducted using our web-based experimental platform (Tsay et al., 2021).

Participants made center-out movements by controlling a cursor with the trackpad on their personal

computers. It was not possible to occlude vision of the hand. However, since the visual stimulus was

presented on a vertical monitor and the hand movement was in the horizontal plane, we assume vision

of the hand was limited to the periphery (based on observations that the eyes remain directed to the

screen during the trial). The size and position of visual stimuli were scaled based on each participant's

screen size (height = 239.6 ± 37.7 mm, width = 403.9 ± 69.5 mm). The experiment was controlled by

custom software written with JavaScript and presented on Google Chrome. Data were collected and

stored using Google Firebase.

The procedure was designed to mimic the lab-based experiments. On each trial, the participant made a

center-out planar movement from the start position to a visual target. A white annulus (1% of screen

height in diameter, 0.4 cm on average) indicated the start position, and a blue circle (1% of screen height

in diameter) indicated the target location. The radial distance of the target from the start position was 40%

of the screen height (5 cm on average). At the beginning of each trial, participants moved the cursor (0.6%

of the screen height in diameter) to the start position, located at the center of their screen. The cursor

was only visible when its distance from the start position was within 20% of the screen height. After

maintaining the cursor at the start position for 500 ms, the target appeared. The participant made a rapid

slicing movement through the blue target. As in the online experiments, there were feedback and no-

feedback trials. For feedback trials, the cursor was visible until it reached the target distance, and then

froze for 1 s at the target distance. On no-feedback trials, the cursor was extinguished after the hand

exited the start position and the target disappeared once the radial distance of the movement reached

the target distance. 500 ms after the end of the trial, the cursor became visible, repositioned at a random

location within 10% of the screen height from the start position. The participant then moved the cursor

to the start position to trigger the next trial.

There were 8 target locations in Exp 3a and 24 target locations in Exp 3b. As with the lab-based

experiments, each experiment included a no-feedback block followed by a feedback block. We obtained

larger data sets in the online studies: For each block, there were 20 trials/target (160 total trials for Exp

3a and 480 total trials for Exp 3b).

Reanalysis of prior data sets

Vindras et al (2005). This study used a design in which the participant did not see the start position of the

movement. This was achieved by not included start position information in the visual display and passively

moving the participant's hand to a start position prior to each reach. Once positioned, a visual target

would appear and the participant reached to that location. Across trials, there were two start positions,

12 target positions (spaced evenly by 30° around the workspace), and two target distances (6 and 12 cm).

In modeling these data, we used the movement endpoint averaged across start positions and target

distances.

Morehead et al (2017) & Kim et al (2018). We re-analyzed the data from the 15° conditions of Exp 4 in

Morehead et al (2017) and Exps 1 and 2 in Kim et al (2018). These three experiments examined visuomotor

adaptation using non-contingent clamped feedback. On perturbation trials, the feedback cursor was

presented at the radial position of the hand but with a fixed 15° angular offset relative to the target.

Participants were informed that the angular position was not contingent on their hand position and

instructed to move directly to the target, ignoring the feedback. This method results in robust implicit

adaptation, with the heading direction of the movement gradually shifting away from the target in the

opposite direction of the cursor. Participants are unaware of this change in behavior (Tsay et al, 2020). In

each experiment, there were three blocks: A no-feedback baseline block (10 trials/target), a clamped

feedback block (60 trials/target), and a no-feedback washout block (10 trials/target).

<u>Data analyses</u>

Motor bias refers to the angular difference between the position of the hand and target when the hand

reaches the endpoint target distance. Angular errors were plotted as a function of the target position with

0° corresponding to the rightward target (3 o'clock location) and 90° corresponding to the forward target.

Positive bias values indicate a counterclockwise error, and negative values indicate a clockwise error.

To assess the similarity of the motor bias functions across different conditions, we calculated the

normalized correlation coefficient as $r_{norm} = \frac{r_{data}}{r_{max}}$. r_{data} is the Pearson correlation coefficient between

the two motor bias functions. r_{max} is the correlation coefficient between the recorded motor bias

function and the true (but unknown) underlying motor bias function from that condition. To calculate

r_{max}, we used a method developed to measure the noise ceiling for EEG/fMRI data [54]:

579
$$r_{max} = \sqrt{\frac{2}{1 + \sqrt{\frac{1}{r_{half}^2}}}}$$

where r_{half} is determined by splitting the data set (based on participants) into random halves and

calculating the correlation coefficient between the first half and the second half of the data. We

bootstrapped r_{half} by resampling the data 2000 times and used the average value. r_{max} is calculated

separately for a pair of conditions and the smaller one is applied as the normalizer for r_{norm}.

<u>Models</u>

To examine the source of motor bias, we considered five single-source models and three multiple-source

models.

Visual Bias model

The Visual Bias model postulates that movement biases arise because the perceived position of the visual

target is systematically distorted (Fig 1b). Here we draw on the work of Huttenlocher et al[13]. In their study,

a visual target was picked from an invisible circle, presented for 1 s and then blanked. The participant then

indicated the remembered position of the target by pointing to a position on a circular digitizing pad. The

results showed a bias towards the four diagonal directions (45°, 135°, 225°, 315°), with the magnitude of

this bias increasing linearly as a function of the distance from the diagonals. As such, the maximum bias

was observed for targets close to four cardinal target locations (0°, 90°, 180°, 270°), and the sign of bias

flipped at the four cardinal target locations.

We used the shape of this function to model bias associated with the perception of the location of the

visual targets. To obtain a continuous function, we assumed a transition zone around the cardinal targets,

each with a half-width represented by the parameter a (Fig 1a), and the peak motor bias is represented

by the parameter b. As such, the angular bias (y) at a target located at $x°$ can be formalized as:

$$[1]\ x' = x\ mod(90)$$

$$[2]\ y = \frac{b * x'}{a}, if\ x' < a;$$

$$y = b - \frac{(x' - a)b}{45° - a}, if\ 90 - a > x' > a;$$

$$y = -\frac{b * (90 - x')}{a}, if\ x' > 90 - a;$$

This model has two free parameters (a and b). If participants directly reach to the perceived target location,

their motor biases will directly reflect their visual biases.

Vector-based Proprioceptive Bias model

Vindras et al. [7,10] proposed a model in which movement biases result from a misperception in estimating

the initial position of the hand (Fig 1c). Specifically, it has been shown that the perceived position of the

hand when placed near the center of the workspace is biased towards the ipsilateral side and away from

the body [18,19,27]). Assuming that the planned movement is formed by a vector pointing from the sensed

hand position to the visual target position, this proprioceptive distortion will result in systematic motor

biases around the workspace. For example, for the target at 90°, misperceiving the initial position of the

right hand to the right of the start position will result in a movement that is biased in the counterclockwise

(leftward) direction.

To simulate this Proprioceptive Bias model, we assumed the participants perceived the start position (0,

0) as a rightward bias away from the midline position, defining a proprioceptive error vector (x_e, y_e). For

a target i at $[x_i, y_i]$, the motor plan is a vector $[x_i - x_e, y_i - y_e]$. From this, we calculated the angular

difference between the motor plan vector and the target position to generate the motor bias for each

target. The two free parameters in this model are $[x_e, y_e]$.

Joint-based Proprioceptive Bias model

Reaching movements may also be planned in joint coordinates rather than the hand (endpoint) position

[21,22]. Based on this hypothesis, motor biases could come about if there is a misperception of the initial

elbow and shoulder joint angles. To implement a Joint-Based Proprioceptive Bias model, we represent the

length of the forearm and upper arm as $l1$ and $l1$, respectively. We denote the initial angles of the

shoulder and elbow joints as θ_0 and φ_0, respectively, and their associated perceived error as θ_e, and φ_e

(See Fig S1).

By setting the origin of the coordinate system for the right shoulder at P_0 (0, 0), the hand can be

represented as:

$$[3]\ P(\theta, \varphi) = [l1\cos\theta_0 + l1\cos\varphi_0, l1\sin\theta_0 + l2\sin\varphi_0].$$

For a fixed position in the workspace, there will be a unique solution pair for θ and φ ($\pi > \varphi > \theta > 0$),

should a solution exist. To calculate the required change in joint angle to reach a visual target, we assumed

that the system plans a movement based on the perceived hand position:

$$[4]\ P_p = P(\theta_0 + \theta_e, \varphi_0 + \varphi_e)$$

Then we solve the following equation to decide the proper $\sigma\theta_i$ and $\sigma\varphi_i$ that transfer the hand from the

start position to a target i at $[x_i, y_i]$:

$$[5]\ P_p - [x_i, y_i] = P(\theta_0 + \theta_e + \sigma\theta_i, \varphi_0 + \varphi_e + \sigma\varphi_i) - P_p$$

We calculated the real movement direction based on the real hand position:

$$[6]\ \sigma h_i = P_p - P(\theta_0, \varphi_0)$$

We compare the direction of σh_i and the target direction to calculate the motor bias. For simplicity, we

assume $l1 = l2 = 24$ cm.[55] The four free parameters in this model are θ_0, φ_0, θ_e, and φ_e

Transformation Bias model

The Transformation Bias model proposes attributes motor biases to systematic errors that arise during

the transformation from a visual to proprioceptive-based reference frame. To implement this model, we

refer to an empirically derived visuo-proprioceptive error map from a data set that sampled most of

reachable space (Fig 1d, [20]). Specifically, in that study, participants were asked to move their unseen hand

from a random start position to a visual target. Rather than require a discrete reaching movement, they

were told to continuously adjust their hand position, focusing on accuracy in aligning the hand with the

target. The direction of the error was relatively consistent across targets, with the final hand position

shifted to the right and undershooting the target. The magnitude of these biases increased as the radial

extent of the limb increased. This basic pattern has been observed across studies using different visuo-

proprioceptive matching methods[18,19,27,28,52,56].

The matching errors provide an empirical measure of the transformation from a visual reference frame to

a proprioceptive reference frame. To model these data, we defined a transformation error vector, $[x_e,$

$y_e]$, whose direction is fixed across space. We then defined a "reference position" with a coordinate of

$[x_r, y_r]$. For upper-limb movements, this reference position is often considered to be positioned around

the shoulder.[57] The transformation error vector at position i is scaled by its Euclidean distance (d) to the

referent position:

$$[7] T_i = d_i [x_e, y_e],$$

$$[8]\ d_i = \sqrt{(x_i - x_r)^2 + (y_i - y_r)^2}$$

669   Movements towards a target i is planned via the vector connecting the start position to the target in

670   proprioceptive space, denoted as:

$$[9]\ \sigma h_i = T_0 - (T_i + [x_i, y_i]),$$

where T_0 is the transformation vector at the start position, which is set as [0,0]. Motor bias is calculated

as the angular difference between the motor plan and the target. The four free parameters in the

Transformation Bias model are x_e, y_e, x_r, y_r.

Hybrid models

The four models described above each attribute motor biases to a single source. However, the bias might

originate from multiple processes. To formalize this hypothesis, we considered three hybrid models,

combining the Visual Bias model with the two versions of the Proprioceptive Bias model and with the

Transformation Bias model. We did not create a hybrid of the Proprioceptive and Transformation Bias

models since they make different assumptions about the information used to derive the motor plan.

Proprioceptive Bias + Visual Bias (P+V) model

We also created two hybrid models, combining the Visual Bias model with the Vector-Based and Joint-Based Proprioceptive Bias models. The Visual Bias model is used to estimate systematic error in the perceived location of the target and the proprioceptive Bias models are used to estimate systematic error in the perceived position of the hand at the start position. For these models, we calculated the biases from two models separately and then added them together:

$$[10]\ b_i(P + V) =\ b_i(\text{Proprioceptive Bias}) +\ b_i(\text{Visual Bias}),$$

where b_i refers to the bias at target i.

Transformation Bias + Visual Bias (T+V) model

The Transformation Bias model attributes motor biases to error that arises during the transformation of spatial information from visual space to proprioceptive space. However, the perceived location of the visual target may be biased. While we recognize that the misperception of the visual target may influence the transformation bias, the visual bias usually very small in our experiment, with a peak <2°. As such, we simplify by calculating the biases from the Transformation Bias and Visual Bias models separately, and then added the values together:

$$[11]\ b_i(T + V) =\ b_i(\text{Transformation Bias}) +\ b_i(\text{Visual Bias})$$

Model comparison

To compare the models, we fit each model with the data from Experiments 1b and 3b in which reaches were made to 24 targets. We evaluated our models on the average data across participants. We used the fminsearchbnd function in MATLAB to minimize the sum square errors (SSE) and used BIC for model comparison:

$$[12] \text{ BIC} = -2LL + k * \ln(n)$$

where k is the number of parameters of the models, and LL is the sum of loglikelihood across all trials of

all participants. Smaller BIC values correspond to better fits.

Modeling motor bias after implicit sensorimotor adaptation

To examine how the motor bias function changes after visuomotor adaptation, we first used the T+V

model to fit the motor bias function from a no-feedback baseline block tested prior to the introduction of

the perturbation. We then used the best-fitted baseline model (TVb) to estimate the shift in the motor

bias function from data obtained in a no-feedback aftereffect block following adaptation:

$$[13] b(i) = TVb(i + v) + h$$

where $b(i)$ is the motor bias at target i in the aftereffect; v and h indicate the vertical and horizontal shift

respectively. To estimate distribution of v and h, we bootstrapped the subjects with repetition for 200

times and fitted the v, h based on the group average of each bootstrapped sample.

Funding

RBI is funded by the NIH (grants NS116883 and NS105839). JST is funded by the NIH (F31NS120448).

Acknowledgments

We thank Zixuan Wang and Anisha Chandy for helpful discussions. We thank Anisha Chandy for data

collection.

Competing interests
RI is a co-founder with equity in Magnetic Tides, Inc.

Data availability statement: Data and code is available at https://github.com/shion707/Motor-Bias

Reference

1. Burge, J., Ernst, M. O. & Banks, M. S. The statistical determinants of adaptation rate in human reaching. *J. Vis.* **8**, 20.1–19 (2008).

2. Tassinari, H., Hudson, T. E. & Landy, M. S. Combining priors and noisy visual cues in a rapid pointing task. *J. Neurosci.* **26**, 10154–10163 (2006).

3. Osborne, L. C., Lisberger, S. G. & Bialek, W. A sensory source for motor variation. *Nature* **437**, 412–416 (2005).

4. Dideriksen, J. L., Negro, F., Enoka, R. M. & Farina, D. Motor unit recruitment strategies and muscle properties determine the influence of synaptic noise on force steadiness. *J. Neurophysiol.* **107**, 3357–3369 (2012).

5. Hamilton, A. F. de C., Jones, K. E. & Wolpert, D. M. The scaling of motor noise with muscle strength and motor unit number in humans. *Exp. Brain Res.* **157**, 417–430 (2004).

6. Dhawale, A. K., Smith, M. A. & Ölveczky, B. P. The Role of Variability in Motor Learning. *Annu. Rev. Neurosci.* **40**, 479–498 (2017).

7. Vindras, P., Desmurget, M., Prablanc, C. & Viviani, P. Pointing Errors Reflect Biases in the Perception of the InitialHand Position. *J. Neurophysiol.* **79**, 3290–3294 (1998).

8. Gordon, J., Ghilardi, M. F., Cooper, S. E. & Ghez, C. Accuracy of planar reaching movements. II. Systematic extent errors resulting from inertial anisotropy. *Exp. Brain Res.* **99**, 112–130 (1994).

9. Ghilardi, M. F., Gordon, J. & Ghez, C. Learning a visuomotor transformation in a local area of work space produces directional biases in other areas. *J. Neurophysiol.* **73**, 2535–2539 (06/1995).

10. Vindras, P., Desmurget, M. & Viviani, P. Error parsing in visuomotor pointing reveals independent

processing of amplitude and direction. *J. Neurophysiol.* **94**, 1212–1224 (2005).

11. Holden, M. P., Newcombe, N. S. & Shipley, T. F. Categorical biases in spatial memory: the role of

certainty. *J. Exp. Psychol. Learn. Mem. Cogn.* **41**, 473–481 (2015).

12. Yousif, S. R., Forrence, A. D. & McDougle, S. D. A common format for representing spatial location in

visual and motor working memory. *Psychon. Bull. Rev.* (2023) doi:10.3758/s13423-023-02366-3.

13. Huttenlocher, J., Hedges, L. V., Corrigan, B. & Crawford, L. E. Spatial categories and the estimation of

location. *Cognition* **93**, 75–97 (2004).

14. Goble, J. A., Zhang, Y., Shimansky, Y., Sharma, S. & Dounskaia, N. V. Directional biases reveal

utilization of arm's biomechanical properties for optimization of motor behavior. *J. Neurophysiol.* **98**,

1240–1252 (2007).

15. Alexander, R. M. A minimum energy cost hypothesis for human arm trajectories. *Biol. Cybern.* **76**,

97–105 (1997).

16. Nishii, J. & Taniai, Y. Evaluation of trajectory planning models for arm-reaching movements based

on energy cost. *Neural Comput.* **21**, 2634–2647 (2009).

17. Kosovicheva, A. & Whitney, D. Stable individual signatures in object localization. *Curr. Biol.* **27**, R700–

R701 (2017).

18. Rincon-Gonzalez, L., Buneo, C. A. & Helms Tillery, S. I. The Proprioceptive Map of the Arm Is

Systematic and Stable, but Idiosyncratic. *PLoS One* **6**, e25214- (2011).

19. van Beers, R. J., Sittig, A. C. & Denier van der Gon, J. J. The precision of proprioceptive position sense.

Exp. Brain Res. **122**, 367–377 (1998).

20. Wang, T. *et al.* Accuracy of hand localization is subject-specific and improved without performance

feedback. *Sci. Rep.* **10**, 19188 (12/2020).

21. Sober, S. J. & Sabes, P. N. Flexible strategies for sensory integration during motor planning. *Nat.*
Neurosci. **8**, 490–497 (2005).

22. Sober, S. J. & Sabes, P. N. Multisensory integration during motor planning. *J. Neurosci.* **23**, 6982–
6992 (2003).

23. Buneo, C. A., Jarvis, M. R., Batista, A. P. & Andersen, R. A. Direct visuomotor transformations for
reaching. *Nature* **416**, 632–636 (2002).

24. Soechting, J. F. & Flanders, M. Errors in pointing are due to approximations in sensorimotor
transformations. *J. Neurophysiol.* **62**, 595–608 (1989).

25. Tillery, S. I., Flanders, M. & Soechting, J. F. A coordinate system for the synthesis of visual and
kinesthetic information. *J. Neurosci.* **11**, 770–778 (1991).

26. Flanders, M. & Soechting, J. F. Frames of reference for hand orientation. *J. Cogn. Neurosci.* **7**, 182–
195 (1995).

27. Jones, S. A. H., Cressman, E. K. & Henriques, D. Y. P. Proprioceptive localization of the left and right
hands. *Exp. Brain Res.* **204**, 373–383 (2009).

28. Cressman, E. K. & Henriques, D. Y. P. Reach adaptation and proprioceptive recalibration following
exposure to misaligned sensory input. *J. Neurophysiol.* **103**, 1888–1895 (2010).

29. Balasubramanian, R., Howe, R. D. & Matsuoka, Y. Task performance is prioritized over energy
reduction. *IEEE Trans. Biomed. Eng.* **56**, 1310–1317 (2009).

30. Summerside, E. M., Courter, R. J., Shadmehr, R. & Ahmed, A. A. Slowing of movements in healthy
aging as a rational economic response to an elevated effort landscape. *J. Neurosci.* (2024)
doi:10.1523/JNEUROSCI.1596-23.2024.

31. Soechting, J. F., Buneo, C. A., Herrmann, U. & Flanders, M. Moving effortlessly in three dimensions:
does Donders' law apply to arm movement? *J. Neurosci.* **15**, 6271–6280 (1995).

32.   Gordon, J., Ghilardi, M. F. & Ghez, C. Impairments of reaching movements in patients without

proprioception. I. Spatial errors. *J. Neurophysiol.* **73**, 347–360 (1995).

33.   Slijper, H., Richter, J., Over, E., Smeets, J. & Frens, M. Statistics predict kinematics of hand

movements during everyday activity. *J. Mot. Behav.* **41**, 3–9 (2009).

34.   van der Kooij, K., Brenner, E., van Beers, R. J., Schot, W. D. & Smeets, J. B. J. Alignment to natural and

imposed mismatches between the senses. *J. Neurophysiol.* **109**, 1890–1899 (2013).

35.   Flash, T. & Hogan, N. The coordination of arm movements: an experimentally confirmed

mathematical model. *J. Neurosci.* **5**, 1688–1703 (1985).

36.   Morehead, J. R., Taylor, J. A., Parvin, D. E. & Ivry, R. B. Characteristics of Implicit Sensorimotor

Adaptation Revealed by Task-irrelevant Clamped Feedback. *J. Cogn. Neurosci.* **29**, 1061–1074 (2017).

37.   Tsay, J. S., Parvin, D. E. & Ivry, R. B. Continuous reports of sensed hand position during sensorimotor

adaptation. *J. Neurophysiol.* **124**, 1122–1130 (2020).

38.   Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R. & Ivry, R. B. Invariant errors reveal limitations

in motor correction rather than constraints on error sensitivity. *Commun Biol* **1**, 19 (12/2018).

39.   Batista, A. P., Buneo, C. A., Snyder, L. H. & Andersen, R. A. Reach plans in eye-centered coordinates.

Science **285**, 257–260 (1999).

40.   Blohm, G., Khan, A. Z. & Crawford, J. D. Spatial Transformations for Eye–Hand Coordination. in

Encyclopedia of Neuroscience 203–211 (Elsevier, 2009).

41.   Henriques, D. Y., Klier, E. M., Smith, M. A., Lowy, D. & Crawford, J. D. Gaze-centered remapping of

remembered visual space in an open-loop pointing task. *J. Neurosci.* **18**, 1583–1594 (1998).

42.   Proske, U. & Gandevia, S. C. The proprioceptive senses: their roles in signaling body shape, body

position and movement, and muscle force. *Physiol. Rev.* **92**, 1651–1697 (2012).

43.   Héroux, M. E., Butler, A. A., Robertson, L. S., Fisher, G. & Gandevia, S. C. Proprioception: a new look

at an old concept. *J. Appl. Physiol.* **132**, 811–814 (2022).

44.   Zhang, H., Daw, N. D. & Maloney, L. T. Human representation of visuo-motor uncertainty as mixtures

of orthogonal basis distributions. *Nat. Neurosci.* **18**, 1152–1158 (2015).

45.   van den Berg, R., Shin, H., Chou, W.-C., George, R. & Ma, W. J. Variability in encoding precision

accounts for visual short-term memory limitations. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 8780–8785

(2012).

46.   Hahn, M. & Wei, X.-X. A unifying theory explains seemingly contradictory biases in perceptual

estimation. *Nat. Neurosci.* (2024) doi:10.1038/s41593-024-01574-x.

47.   Mawase, F., Lopez, D., Celnik, P. A. & Haith, A. M. Movement Repetition Facilitates Response

Preparation. *Cell Rep.* **24**, 801–808 (2018).

48.   Shadmehr, R., Huang, H. J. & Ahmed, A. A. A representation of effort in decision-making and motor

control. *Curr. Biol.* **26**, 1929–1934 (2016).

49.   van Beers, R. J., Wolpert, D. M. & Haggard, P. When Feeling Is More Important Than Seeing in

Sensorimotor Adaptation. *Curr. Biol.* **12**, 834–837 (2002).

50.   Sheehan, T. C. & Serences, J. T. Distinguishing response from stimulus driven history biases. *bioRxiv*

(2023) doi:10.1101/2023.01.11.523637.

51.   Tsay, J. S., Kim, H., Haith, A. M. & Ivry, R. B. Understanding implicit sensorimotor adaptation as a

process of proprioceptive re-alignment. *Elife* **11**, e76639 (2022).

52.   Cressman, E. K. & Henriques, D. Y. P. Sensory recalibration of hand position following visuomotor

adaptation. *J. Neurophysiol.* **102**, 3505–3518 (2009).

53.   Oldfield, R. C. The assessment and analysis of handedness: The Edinburgh inventory.

Neuropsychologia **9**, 97–113 (1971).

54.   Schoppe, O., Harper, N. S., Willmore, B. D. B., King, A. J. & Schnupp, J. W. H. Measuring the

Performance of Neural Models. doi:10.3389/fncom.2016.00010.

55.   Fryar, C. D., Gu, Q. & Ogden, C. L. Anthropometric reference data for children and adults: United

848   States, 2007-2010. *Vital Health Stat. 11* 1–48 (2012).

849   56.   Johnson, E. O., Babis, G. C., Soultanis, K. C. & Soucacos, P. N. Functional neuroanatomy of

850   proprioception. *J. Surg. Orthop. Adv.* **17**, 159–164 (2008).

851   57.   Haggard, P., Newman, C., Blundell, J. & Andrew, H. The perceived position of the hand in space.

852   *Percept. Psychophys.* **62**, 363–377 (2000).

Figure S1. Schematic of a vector based and joint-based Proprioceptive Bias model. Previous studies

have considered two variants of the Proprioceptive Bias model. (a) A vector-based model in which the

motor plan is a vector pointing from the perceived hand position to the target[7,10]. (b) A joint-Based

model in which the movement is encoded as changes in the shoulder and elbow joint angles to move

the limb from a start position to a desired location[21,22]. See Method, Models for details.

Figure S2. The Transformation Bias model can explain the motor bias functions when the visual

information is shifted. (a) In Sober and Sabes (2003)[22] participants performed center-out reaches to a

visual target. To perturb the visual information, the start position was presented 6 cm to the left or right

of the actual start position of the hand. (b) Participants showed a one-peaked motor bias functions with

the shift-left and shift-right functions shifted in an antiphase relationship to one another. (c) These bias

functions are quantitively captured by the Transformation Bias model.

	x_r(cm)	y_r(cm)	x_e	y_e	a(°)	b(°)
Exp 1b	75.7	-17.2	-0.27	-0.05	19.6	3.8
Exp 3b	75.1[a]	-0.14[a,b]	-0.24	-0.00	20.0	1.5

Table S1. Parameter estimates from best fits of the T+V model for the data from Exps 1b and 3b.

See Methods for description of each parameter.

a. Participant moved on the trackpad in Exp 3b. We assumed the movement distance was 1 cm and

scaled the parameters accordingly.

b. The estimate of y_r is much smaller in Exp 3b compared to Exp 1b, suggesting the workspace in Exp 3b

is closer to the body. This attenuates the average magnitude of the bias.