Imprinting and DNA methylation in water lily endosperm: implications for seed evolution

Rebecca A. Povilus[1], Caroline A. Martin[1,2], Lindsey L. Bechen[1], and Mary Gehring[1,2,3,4*]

[1] Whitehead Institute for Biomedical Research, 455 Main Street, Cambridge MA 02142

[2] Department of Biology, Massachusetts Institute of Technology, Cambridge MA 02139

[3] Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge MA

02139

[4] Howard Hughes Medical Institute

*Corresponding author: Mary Gehring, mgehring@wi.mit.edu

Author Contributions: R.A.P. and M.G. conceived of the original premise of the study. R.A.P.

conducted experiments and wrote the paper with input from M.G. and C.A.M. C.A.M. assisted in

processing genomic data. L.L.B performed methylation profiling on leaf tissue.

Competing Interest Statement: The authors have no competing interests to declare.

Keywords: Endosperm, Imprinting, Early Angiosperms, DNA methylation

This PDF file includes:

Main Text

Figures 1-4

Legends/Titles for Supplementary Figures 1-5 and Supplementary Tables 1-5

Abstract

Endosperm is a key evolutionary innovation associated with the origin of angiosperms (flowering plants). This altruistic seed tissue supports the growth and development of the embryo by mediating the relationship of the mother plant as a nutrient source to the compatriot embryo as a nutrient sink. The endosperm is the primary site of gene imprinting in plants (where expression of an allele in offspring depends on which parent it was inherited from) and of parent-specific epigenetic modifications like DNA methylation, which are differentially patterned during male and female gamete development. Experimental results from a phylogenetically-wide array of monocot and eudicot plants suggest these parent-of-origin effects are a common feature across angiosperms. However, information about genetic imprinting and epigenetic modifications in seeds of angiosperm lineages whose origins predate the monocot-eudicot divergence (such as Nymphaeales, water lilies) is extremely limited. Additionally, Nymphaeales are an intriguing lineage in which to investigate seed genetic and epigenetic phenomena, as they are characterized by diploid endosperm and a maternal storage tissue (perisperm), both of which are unusual across angiosperm diversity. Here, we examined DNA methylation and genetic imprinting using two reproductively compatible water lily sister-species, *Nymphaea thermarum* and *N. dimorpha*. Our results suggest that maternally-expressed imprinted genes and differential DNA methylation of maternally and paternally inherited endosperm genomes are an ancestral condition for endosperm, while other seed characters like seed provisioning strategies, endosperm ploidy, and paternally-expressed imprinted genes might have evolved as coinciding, opposing strategies in the evolutionary dialogue over parental control of offspring development.

Main Text

Introduction

The evolutionary origin of endosperm, a second fertilization product in the seeds of flowering plants, fundamentally altered the relationship between an embryo and its mother during seed development. In non-flowering seed plants, the embryo is directly connected to tissue that only contains genome complements from its mother. However, in angiosperm seeds, endosperm largely separates the embryo and its mother and is the product of a fertilization event and thus biparental, with both maternal and paternal genome contributions. Endosperm is widely recognized as a key mediator of developmental and nutritional relationships between an embryo and its mother (Povilus and Gehring 2022). The balance of maternal and paternal genomes is important for endosperm (and seed) viability, as evidenced by the phenomena of parental genome dosage sensitivity ((Haig and Westoby 1991) and references therein). When extra paternal genome complements are added to the endosperm, it over-proliferates – often leading to initially larger but ultimately collapsed, inviable seeds. Conversely, when extra maternal genome complements are added to the endosperm, reduced endosperm proliferation is observed, resulting in smaller seeds with fewer invested maternal resources (Haig and Westoby 1991; Birchler 1993; Scott et al. 1998). The endosperm is subject to other parent-of-origin effects such as imprinted gene expression (where expression of an allele depends on which parent it was inherited from) and parent-of-origin specific epigenetic modifications like DNA and histone methylation, which are differentially patterned during male and female gamete development (Gehring et al. 2006; Park et al. 106; Moreno-Romero et al. 2019; Borg et al. 2020). Our knowledge of endosperm gene imprinting and its underlying mechanisms is built from experiments performed in a phylogenetically wide array of monocot and eudicot plants (Povilus and Gehring 2022; Picard and Gehring 2020). However, information from lineages whose origins predate the monocot-eudicot divergence is extremely limited.

The order Nymphaeales (water lilies) is sister to all other angiosperms except for

Amborella tricopoda. Endosperm parental genome dosage sensitivity has been documented in

a species of water lily (Povilus et al. 2018), but nothing is known about patterns of genetic

imprinting or endosperm epigenetic patterning in this or any other ANA-grade lineage

(Amborella, Nymphaeales, Austrobaileyales), magnollids, or Chloranthales. In addition to their

relationship to other angiosperms, Nymphaeales are a particularly intriguing system in which to

investigate genetic imprinting and associated epigenetic patterning given the unique

combination of seed characters found in this lineage. First, endosperm of the Nymphaeales is

ab initio-cellular (the first nuclear division of the endosperm is accompanied by cellular division)

and diploid with a 1:1 maternal:paternal genome ratio (Orban and Bouharmont 1998; Williams

and Friedman 2002; Friedman 2006; Friedman 2008; Rudall et al. 2008; Povilus et al. 2015),

whereas triploid endosperm (2:1 ratio) characterizes the majority of angiosperms and all taxa in

which endosperm epigenetic patterning and genetic imprinting have been studied (Haig and

Westoby 1991). Diploidy has been suggested to represent the ancestral ploidy of endosperm

(Williams and Friedman 2002) and thus Nymphaeales are an opportunity to test how these

processes operate in the context of different base maternal-paternal genome/gene dosage

ratios. Second, in seeds of Nymphaeales nutrients are primarily stored in a perisperm (which is

derived from maternal tissue and contains no paternal genome contribution) instead of in

offspring tissues, in contrast to the vast majority of flowering plants (Lersten 2004; Patten et al.

2014). Therefore, Nymphaeales is an excellent clade in which to investigate the suggested

connection between genetic imprinting in endosperm and control of nutrient storage (Haig and

Westoby 1991; Patten 2014). Nutrient storage in perisperm is only initiated after fertilization

(Povilus et al. 2015), suggesting influence of offspring tissues on this process.

Characterizing genetic imprinting and epigenetic modifications in water lilies therefore

offers a unique perspective on the evolution of key endosperm traits and processes that are

associated with the origin of angiosperms. Here, we sought to determine whether gene

imprinting and parent-of-origin effects on DNA methylation, which has been mechanistically

linked to gene imprinting, preceded the origin of parental dosage-imbalanced (triploid)

endosperm.

Results

Nymphaea thermarum has been developed as an experimental system for the

Nymphaeales (Povilus et al. 2015; Povilus et al. 2018; Povilus et al. 2020). Assessing parent-

of-origin effects at the molecular level requires sequence polymorphisms, of which there are few

within the highly-inbred extant populations of *N. thermarum* in cultivation (Povilus et al. 2015).

We therefore assessed parent-of-origin effects in Nymphaeales by examining F1 tissue from

crosses between *N. thermarum* and *Nymphaea dimorpha* (which was formerly known as *N.*

minuta). These two species are estimated to have diverged roughly 20 million years ago

(Borsch et al. 2011). We confirmed the internal structure of young seeds of *N. thermarum*, *N.*

dimorpha, and of F1 reciprocal crosses and determined that we could ensure consistency in

developmental stage among crosses (Figure 1A). The hybrid F1 seeds are fully viable and

germinate to give rise to viable F1 plants (Supplementary Figure 1), suggesting no large-scale

divergence in endosperm developmental programs that would lead to failure in seed

development. We performed long-read based, *de novo* genome assembly and annotation for *N.*

dimorpha (248 contigs with an NG50 of 13,941,033 bp, representing 83% of the estimated

genome size, with the set of 40,850 annotated genes having a BUSCO score of 85% for the

Embryophyta gene set) and an improved genome assembly and annotation for *N. thermarum*

(1,553 contigs with and NG50 of 4,352,861, representing 86% of the estimated genome size,

with the set of 42,431 annotated genes having a BUSCO score of 83% for the Embryophyta

gene set) (Supplementary Table 1, Supplementary Figure 2). To allow direct comparisons of

genomic regions, the *N. thermarum* and *N. dimorpha* genomes were aligned and re-annotated

to create "reorganized" genomes for each species (each 358,929,111 bp in length and with a

resolved annotation having 39,608 genes and a BUSCO score of 72% for the Embryophyta

gene set).

To examine parent-of-origin specific gene expression, we made use of reciprocal

crosses between *N. thermarum* (*Nt*) and *N. dimorpha* (*Nd*) (2 samples each from *N. thermarum*

x N. dimorpha and *N. dimorpha x N. thermarum* crosses) as well as self-fertilized seeds (3

samples each of *N. thermarum* and *N. dimorpha*), and isolated RNA from young endosperm at

9-10 days after pollination/anthesis (Supplementary Table 1). By performing mRNA-seq, we

detected expression of a total of 22,984 genes with a TPM >= 1 (expression averaged across all

samples). A principal component analysis (PCA) of total gene expression revealed that

biological replicates clustered together according to cross type, with hybrid endosperm samples

midway along PC1 (54.85%) between endosperm from *N. thermarum* self-fertilized seeds and

N. dimorpha self-fertilized seeds (Figure 2A). Differential gene expression analysis between

sets of hybrid and non-hybrid seeds revealed that 879 genes were consistently significantly

differentially expressed in all comparisons; this set of genes was not significantly enriched for

any KEGG pathways, but was enriched for the GO biological process term "RNA-dependent

DNA biosynthetic processes" (FDR = 1.9e-2, n=8, fold enrichment = 7.1). Importantly, while

expression differences between hybrid and non-hybrid endosperm existed, endosperm of the

two hybrid cross directions were more similar to each other than they were to endosperm of the

parental lines (Figure 2A). We implemented a previously developed pipeline to evaluate

imprinted gene expression (Gehring et al, 2011) (see Methods). In each possible comparison of

a *Nt* x *Nd* and *Nd* x *Nt* cross, we identified transcripts that showed a significant bias in the

number of reads mapping uniquely to either the maternally- or paternally-inherited alleles, in

both cross directions (Figure 2B, Supplementary Table 2). For these imprinting tests, 26,465

genes had at least one read that could be assigned to a parent-of-origin and 16,647 genes

passed our minimum allele-specific read count cut-off of 50 reads and were assessed for

imprinting. Our analysis revealed the presence of imprinted genes in *Nymphaea*. We identified

small numbers of paternally-biased genes in individual cross comparisons, but only 1 paternally

expressed imprinted gene (PEG) was consistent in at least 75% of comparisons (3 of 4 total

possible cross comparisons) (Figure 2C); this PEG is a homolog of *CELLULOSE SYNTHASE*

LIKE G2 (*ATCLSG2*) in *Arabidopsis thaliana*. A handful of PEGs have previously been identified

as conserved between monocots and dicots (Pignatta and Gehring 2012). We examined the

expression of homologs of these specific genes in *Nymphaea* endosperm. Although there was

some evidence for paternally-biased expression, there were also large *cis* or species effects on

transcription, and these genes did not meet all of our criteria for imprinting (Supplementary

Table 2). In contrast to PEGs, 157 MEGs were consistently identified in at least 75% of

comparisons, with 147 being identified as MEGs in all comparisons (Figure 2C) (Supplementary

Table 2). Previous studies have shown that imprinting can be altered or obscured in

interspecies hybrids (Josefsson et al. 2006; Burkart-Waco et al. 2015; Florez-Rueda et al.

2016), and we evaluated whether the identified MEGs were differentially expressed in the F1

hybrid endosperm. When endosperm gene expression profiles were compared between each

hybrid type and each parental species type (Figure 2D, Supplementary Figure 3, Supplementary

Table 3), only 10 MEGs were consistently (in 75% or more of comparisons) significantly

differentially expressed, including only 1 MEG that was significantly differentially expressed in all

comparisons. This indicates low overlap between identified MEGs and genes that are mis-

regulated in hybrid crosses. We conclude that imprinted expression exists in *Nymphaea*

endosperm, but is largely restricted to MEGs and mostly does not include genes whose

expression is altered in hybrid endosperm.

The identification of MEGs can be influenced by contamination with maternal tissue or

transcripts in endosperm samples and thus we evaluated the extent to which this might be

affecting our results. Notably, PCA results indicate that hybrid endosperm from both cross

directions was more like each other transcriptionally than endosperm of their mother species,

suggesting no significant, wide-spread maternal contamination (Figure 2A). We additionally

performed RNA *in situ* hybridizations on seeds of self-fertilized *N. thermarum* for a set of

identified MEGs to test whether they showed substantial expression in maternal tissue or lacked

expression in the endosperm (both of which would indicate potential for maternal tissue

contamination). Target genes were selected for high maternal expression bias and high

expression (Figure 2E), including a gibberellic acid oxidase homolog (*GA2OX2*) (Figure 2F) and

a sucrose synthase 3 homolog (*SUS3*) (Figure 2G). We furthermore confirmed that the high

starch and carbohydrate content of the perisperm was not interfering with the *in situ*

hybridization experimental protocol, as we were able to detect expression of a subfamily of

terpene synthases in the perisperm, as well as in the endosperm (Supplementary Figure 4). For

both *GA2OX2* and *SUS3*, we detected expression in the endosperm and not in the perisperm,

indicating that for these and likely other MEGs, the identified maternal expression bias is

unlikely to be caused by contamination with maternal tissue. For *SUS3*, we also performed *in*

situ hybridizations on seeds from reciprocal crosses and from *N. dimorpha* self-fertilizations, and

found similar endosperm expression patterns (Supplementary Figure 4).

Additionally, we took advantage of whole-seed expression datasets from *N. thermarum*

(Povilus and Friedman 2022) to explore the potential impact of maternal tissue transcript

contamination on our imprinting results. For each transcript expressed in the endosperm, we

calculated corrected endosperm maternal read counts based on assuming that 50% or 25% of

the isolated endosperm transcript pool was comprised of transcripts from a whole-seed pool

(Tonosaki et al. 2024). We found that correcting for an assumed 50% sample contamination

resulted in identification of 112 MEGs and correcting for an assumed 25% contamination

resulted in identification of 134 MEGs, compared to the originally identified 157 MEGs

(Supplementary Table 4). Both sets of 112 MEGs and 134 MEGs were subsets of the originally

identified 157 MEGs. The "corrected" MEGs were enriched for KEGG or GO biological process

terms related to response and development, carbohydrate processing, and secondary

metabolism (Supplementary Table 4), similar to the enrichments for the original, uncorrected set

of MEGs (Figure 2G). Furthermore, in the corrected datasets *SUS3* was not identified as a

MEG, although our *in situ* hybridizations demonstrated that *SUS3* was not highly expressed in

maternal seed tissues but was expressed in endosperm (Figure 2F, Supplementary Figure 4).

We concluded that the correction for extensive, assumed whole-seed transcript contamination

was likely inappropriately removing true MEGs, and therefore proceeded to use the uncorrected

set of MEGs for further analysis.

Overall, *Nymphaea* MEGs were significantly enriched for GO annotations associated

with response and development, carbohydrate processing, and secondary metabolism (Figure

2H, Supplementary Table 4). The enrichment for processes integral to development and nutrient

dynamics in seeds is similar to what has been found in other species (Xin et a. 2013; Picard et

al. 2021). Notably, *SUS3* is a maternally expressed imprinted gene in *Nymphaea* and is a key

part of nutrient processing in the endosperm in other species (Angeles-Núñez and Tiessen

2010). These results are congruent with genetic imprinting being associated with nutrient

dynamics during seed development.

Having found evidence for parent-of-origin effects on gene expression, we next

investigated parent-of-origin effects on DNA methylation by performing endosperm whole-

genome enzymatic methyl-sequencing (EM-seq). We again made use of reciprocal crosses

between *N. thermarum* and *N. dimorpha* to permit allele-specific characterization of DNA

methylation patterning in young F1 hybrid endosperm; biological replicates from two *N.*

thermarum x N. dimorpha crosses and from three *N. dimorpha x N. thermarum* crosses were

analyzed (Supplementary Table 1). We also obtained methylation profiles from single samples

of *N. thermarum* and *N. dimorpha* leaves and the leaves of an F1 *N. dimorpha x N. thermarum*

hybrid. In both endosperm and leaves, the average methylation profiles of *N. thermarum* and *N.*

dimorpha alleles of genes and repeats (transposable elements) (Figure 3A) was similar to other

angiosperms (Niederhuth et al. 2016), with CG methylation occurring in gene bodies and CG

and non-CG methylation in repeats. We identified differentially methylated regions (DMRs)

between *N. thermarum* and *N. dimporpha* genomes in F1 hybrid leaves and between *N.*

thermarum and *N. dimorpha* leaves. In hybrid leaves, the majority of DMRs occurred in the CG

context, and similar numbers of regions were more methylated in one species or one genome

versus the other (Figure 3B, Supplementary Figure 5). We then identified DMRs between

maternal and paternal alleles in F1 endosperm. The majority of CG and CHH DMRs were less

methylated on maternal alleles than on paternal alleles, regardless of which species was the

maternal parent in the F1 hybrid endosperm (Figure 3B). Additionally, maternal alleles of both

species were consistently hypermethylated in the CHG context and were generally more

methylated in endosperm than in hybrid or parental species' leaf tissue, for both genic flanking

regions and within repeats (Figure 3A). The higher methylation of maternal alleles in the CHG

context was also borne out by analyzing the number of DMRs between maternal and paternal

alleles (Figure 3B, Supplementary Table 5): the CHG context had the highest difference

between the portion of maternally- or paternally-hypermethylated DMR windows, with both

species having more maternally hypermethylated windows. These methylation patterns were

not found in leaf tissue (Figure 3B; Supplementary Figure 5) and may therefore be unique to

endosperm. Together, these findings parallel observations of methylation patterning in

endosperm of monocots and dicots, in which endosperm is CG hypomethylated on maternal

alleles due to active DNA demethylation that occurs in the central cell (the female gamete that is

the progenitor of the endosperm) before fertilization. Maternal allele CG hypomethylation has

been noted in rice, Arabidopsis, and other species (Gehring et al. 2009; Hsieh et al. 2009;

Waters et al. 2011; Rodrigues et al. 2013; Park et al. 2016; Zhang et al. 2016; Zhang et al.

2021). Endosperm maternal allele CHG hypermethylation has been observed in *Arabidopsis*

lyrata (Klosinska et al. 2016) and to a lesser extent in *A. thaliana* (Moreno-Romero et al. 2019),

although this occurs primarily in gene bodies, unlike the observations here where it occurs in

gene flanking regions and in repeats.

We further examined the set of genes that overlapped a maternal- or- paternal

hypermethylated DMR consistently across both species (had a DMR that overlapped the gene

region +/- 2kb in at least 75% of pairwise comparisons, while not overlapping a DMR of the

opposite type in more than 25% of comparisons) (Figure 3C, Supplementary Table 5). Similar to

looking at numbers of DMRs associated with genes separately in each species, more genes

were associated with paternally hypermethylated DMRs in the CG and CHH contexts, while in

the CHG context more genes overlapped with maternally hypermethylated DMRs. Few

imprinted loci were consistently associated with allele-specific DMRs (Supplementary Table 5).

Four MEGs were consistently associated with maternal allele hypomethylated DMRs in the CG

context, including homologs of *TARGET OF MONOPTEROS 6* (*TMO6*) and *GA2OX2* (Figure

3D)*,* and 2 MEGs were associated with maternally hypomethylated DMRs in the CHH context.

These regions were not differentially methylated in F1 hybrid leaves, indicating a parent-of-

origin effect on methylation that is specific to the endosperm (Figure 3D). The one identified

PEG was not associated with any significant methylation differences between parental alleles.

Thus, we conclude that there are parent-of-origin specific differences in DNA methylation in

Nymphaea endosperm, of a similar nature (maternal CG hypomethylation) to those observed in

monocots and eudicots. The majority of *Nymphaea* MEGs are not associated with differential

DNA methylation. For comparison, in maize and Arabidopsis approximately 50% and 40% of

MEGs, respectively, are associated with differential DNA methylation (Pignatta et al. 2014; Gent

et al. 2022).

Discussion

Our results illuminate the evolution of imprinting and potential mechanisms facilitating the

emergence of gene imprinting. In summary, we found that genetic imprinting and parent-of-

origin effects on DNA methylation patterning occur in endosperm of Nymphaea seeds. Both

DNA methylation and genetic imprinting have been suggested to be strategies that can alter the

effective maternal or paternal genome/gene dosage in endosperm. Changes to absolute

parental genome dosage (and ploidy) of endosperm have also occurred repeatedly during

angiosperm evolution. Our discovery of endosperm genetic imprinting and parent-of-origin

effects on DNA methylation in *Nymphaea* suggests that these characters/processes predate the

evolution of triploid endosperm and are likely to be have been either co-opted from preexisting

ancestral molecular programs or are novelties associated with the origin of endosperm itself

(Figure 4). In either case, these findings demonstrate that a 2:1 maternal:paternal genome

dosage ratio is not a requirement for either endosperm maternal allele CG hypomethylation and

CHG hypermethylation, or for genetic imprinting. Furthermore, our results suggest that parent-

of-origin effects on endosperm development in *Nymphaea* (Povilus et al. 2018) could be linked

to parent-of-origin-specific DNA methylation patterning or maternally-expressed imprinted

genes, but not extensively to paternally-expressed imprinted genes. This is perhaps surprising

given that Povilus et al. (2018) observed paternal effects when diploids were pollinated by

tetraploids: in mature seeds, endosperm of both 4n x 2n (maternal excess) and 2n x 4n

(paternal excess) is larger than endosperm of 2n x 2n crosses. However, the developmental

timing by which larger endosperm is achieved differs between maternal and paternal excess

crosses. During later development (7-32 DAA), the endosperm of 2n x 4n crosses grows

significantly faster than that of 2n x 2n crosses. By contrast, in 4n x 2n crosses, the endosperm

grows faster earlier (1-7 DAA), and then decelerates at later stages. The observed maternal and

paternal effects in *N. thermarum* endosperm are therefore distinct from those typically observed

in maize or Arabidopsis, where maternal excess seeds undergo early endosperm cellularization

and are typically smaller than 2n x 2n seed at maturity and paternal excess seeds undergo

extended endosperm proliferation and are larger and dead (Arabidopsis) or smaller and dead

(maize) at maturity (Scott et al. 1998; Pennington et al. 2008). Although it has been proposed

that an increased dosage of PEGs is the cause of interploidy paternal excess phenotypes, direct

evidence is limited. Indeed, in Arabidopsis it has been shown that PEG expression is increased

in both viable and non-viable seeds from Arabidopsis paternal excess crosses (Satyaki and

Gehring 2019), suggesting that PEG expression is not the determining factor, or sole

determining factor, of paternal excess interploidy phenotypes. Finally, a single asymmetry

between parental genomes, such as a MEG, has the potential to cause both maternal and

paternal parental effects as the endosperm dosage of the gene would differ between interploidy

crosses.

It remains possible that we failed to detect PEGs because of our use of inter-species

comparisons or because of the developmental stage at which we analyzed imprinting. Inter-

specific and inter-ploidy crosses can result in misregulation of imprinted gene expression. In

hybrid crosses between *A. thaliana* and *A. arenosa*, several PEGs gain expression from the

maternally-inherited allele in whole seed data, rendering them biallelic or even maternally

biased (Josefsson et al. 2006; Burkart-Waco et al. 2015). Other genes become newly paternally

biased. Similarly, endosperm of wild tomato hybrids exhibit a genome-wide increase in maternal

transcriptome contribution and reduced paternal bias of PEGs (Florez-Rueda et al. 2016).

However, in both of these examples, endosperm development is defective and the hybrid seeds

are inviable. This is not the case in crosses between *N. thermarum* and *N. dimorpha* – the

seeds and F1 plants are fully viable (Figure 1, Supplementary Figure 1), suggesting no defects

in endosperm development of the kind that is often correlated with misregulated genomic

dosage or misregulated imprinting. Thus, although we cannot exclude it, we think it unlikely that

PEGs were not detected in *Nymphaea* endosperm because of the genetic material used in our

crosses. We also cannot exclude the possibility that PEGs might be active at earlier stages of

Nymphaea seed development than we sampled, before significant development/functioning of

the perisperm, as has been suggested by a study on endosperm of a different Nymphaea

species (Florez-Rueda et al, 2024). Yet, in other species PEGs have been detected across all

assayed stages of endosperm development, including in seeds with similar stages of embryo

development as sampled in our study.

The observed differences in CG DNA methylation between endosperm maternal and

paternal alleles (Figure 3B,D) are consistent with the activity of a 5-methylcytosine DNA

glycosylase in the central cell before fertilization, which would be predicted to cause maternal

allele CG hypomethylation in endosperm after fertilization. Homologues of DME are present in

all angiosperms (Pei et al. 2019), including *Nymphaea* (Povilus and Friedman 2022). The lack of

high congruence between CG maternal allele hypomethylated DMRs and *Nymphaea* MEGs is

not inconsistent with data from other species. In Arabidopsis and maize, there are many more

endosperm CG hypomethylated regions than there are imprinted genes (Gehring et al. 2009;

Pignatta et al. 2014; Gent et al. 2022). Although the imprinting of individual Arabidopsis MEGs

like *FWA* and *SDC* is correlated with differential methylation between maternal and paternal

alleles (Kinoshita et al. 2004), as a group MEGs are not enriched for differential methylation

compared to non-imprinted genes (Pignatta et al. 2014). This is in contrast to PEGs, where

differential CG methylation of upstream or downstream regions is enriched compared to non-

imprinted genes (Pignatta et al. 2014; Moreno-Romero et al. 2019). We also observed

endosperm maternal allele CHG hypermethylation in *Nymphaea* endosperm in repeats and

gene-flanking regions (Figure 3A-C). In other angiosperm species, CHG methylation is

frequently associated with H3K9me2 and transcriptional silencing. The maternal alleles of PEGs

in Arabidopsis species are significantly associated with CG hypomethylation and CHG

hypermethylation in endosperm (Klosinska et al. 2016; Moreno-Romero et al. 2019). Although

we observed maternal allele CG hypomethylation and CHG hypermethylation in *Nymphaea*

endosperm, they were not associated with PEGs, suggesting that paternal expression bias is

not an inevitable outcome of these types of epigenetic patterns. Other epigenetic mechanisms

could regulate gene imprinting in *Nymphaea*. In mammals, imprinting of a small number of

genes is regulated by H3K27me3, without an apparent role for DNA methylation (Lewis et al.

2004; Inoue et al. 2017; Santini et al. 2021; Raas et al. 2022), and has been proposed as an

ancestral mechanism of imprinting in the placenta. H3K27me3 also plays an important role in

plant gene imprinting and is often coincident with differential DNA methylation (Gehring et al.

2006; Zhang et al. 2014; Moreno-Romero et al. 2019). Histone modification patterns in

Nymphaea endosperm is a potential area of future investigation.

In the context of parental conflict over the investment of maternal resources in offspring

and maternal-offspring coadaptation, the evolution of maternal storage tissues and the notable

paucity of PEGs in Nymphaea endosperm give rise to two distinct hypotheses about the early

evolution of angiosperm seed development and endosperm molecular/genetic processes

(Figure 4). For the first hypothesis, if MEGs and PEGs were both present in the ancestrally

diploid endosperm of the earliest flowering plants, then PEGs were largely lost in association

with the evolution of perisperm in the Nymphaeales (Figure 4, left). If genetic imprinting is a tool

to establish maternal or paternal control over resource investment in offspring (Haig and

Westoby 1991), perhaps PEGs are no longer an effective paternal strategy when the mother

establishes primary control by storing invested resources in a maternally-derived tissue that

does not have a paternal genome contribution. A second hypothesis is that genetic imprinting

has evolved in stepwise fashion along with endosperm ploidy changes (Figure 4, right). In this

case, MEGs may have evolved as a maternal strategy to balance the addition of a paternal

genome – and potential for paternal influence on seed development – that resulted in the origin

of endosperm. Subsequent addition of a second maternal genome complement with the

evolution of triploid endosperm may then be similarly associated with the evolution of PEGs.

Studying other members of ANA-grade lineages with diploid endosperm that lack perisperm

(such as in Austrobaileyales (Losada et al. 2017)) or taxa with triploid endosperm and perisperm

(as can be found in Amaranthaceae (Coimbra and Salema 1999; López-Fernández and

Maldonado 2013), within the eudicots) would help distinguish between these hypotheses by

specifically testing the relationship between maternally-derived storage tissues and the absence

of PEGs. Thus, while characterizing parent-of-origin effects on gene expression and epigenetic

modifications in Nymphaea endosperm is an important step in understanding the evolution of

molecular processes in endosperm, the genetic and developmental diversity across angiosperm

seeds deserves further attention.

Materials and Methods

Data availability

Data generated as part of this study are available as part of NCBI BioProjects PRJNA1085993,

PRJNA1086866, PRJNA1086863, PRJNA1085992, and PRJNA1087317, including raw

sequence data deposited in SRA, and genome assemblies deposited in NCBI WGS.

Plant growth and sample collection

Seeds of *N. thermarum* and *N. dimorpha* were sourced from the Arnold Arboretum of Harvard

University (Boston, MA, USA) and grown at the Whitehead Institute for Biomedical Research

(Cambridge, MA, USA) (Supplementary Materials and Methods).

Controlled pollinations and self-fertilizations were performed as previously described

(Povilus et al. 2015; Povilus et al. 2018). For collection of seeds from crosses and self-

fertilizations, first day of anthesis (time of female receptivity and fertilization (Povilus et al.

2015)) was defined by presence of stigmatic fluid. Fruits were collected at 10 days after

anthesis (DAA) and seeds were immediately removed and dissected with fine forceps in

dissection buffer. Endosperm tissue was washed with dissection buffer multiple times and

frozen in liquid nitrogen (Supplementary Materials and Methods).

Whole genome sequencing, assembly, and annotation

For long-read and short-read DNA sequencing, high molecular weight DNA was

extracted from > 1 g young leaf samples from a single individual plant using a modified CTAB-

based protocol (Supplementary Materials and Methods). Samples were prepared for PacBio

sequencing (PacBio Sequel SMRTcell, 20h, v3 chemistry) and were sequenced at the MIT

BioMicroCenter. *Nymphaea thermarum* was sequenced in both LR (long-read) and HiFi (high-

fidelity) modes; *N. dimorpha* was sequenced in LR mode. *N. dimorpha* short-read data was

obtained from the same sample, using 1 lane of an Illumina HiSeq2000 flow-cell (40 bp, paired-

end reads) at the MIT BioMicroCenter. Short-read genomic data for *N. thermarum* was

downloaded from BioProject PRJNA508901.

Genome assembly for *N. thermarum* and *N. dimorpha* was performed separately using

long-reads as input for Canu (version 2.1.1) (Koren et al. 2017); short-reads were used to polish

the resulting assemblies using POLCA (from MaSuRCA version 3.4.2) (Zimin et al. 2013).

Genome assemblies were visualized with Bandage (Wick et al. 2015). The resulting original

genome assemblies were separately annotated with MAKER (version 2.31.10)(Campbell 2014)

for both species, using an iterative approach to train AUGUSTUS (version 3.3.3)(Stanke 2006)

and SNAP (version 2006.07.28-1)(Korf 2004) gene-model predictors; initial input for all

annotation pipelines included the set of transcript and protein sequences from the published *N.*

thermarum genome assembly/annotation(Povilus 2020), the set of all protein sequences from

Nymphaeaceae available on NCBI, protein sequences from the *N. colorata* genome assembly

and annotation (Zhang 2020), all basal Magnoliophyta protein sequences on Uniprot, Amborella

protein sequences, and TAIR10 protein sequences. Three rounds of annotation and gene model

predictor training were performed for annotation of both species. Repeat identification and

masking was performed with RepeatMasker (version 4.0.5)(Chen 2004) using Spermatophyta

as the specified query clade and the Embryophyta repeat database.

To create genome assemblies of *N. thermarum* and *N. dimorpha* that shared positional

homology, the *N. thermarum* contigs were mapped to *N. dimorpha* contigs using minimap2 (Li

2018) (the *N. dimorpha* assembly was used as the reference as it had the fewest contigs;

genome alignment visualized using D-Genies (Cabanettes and Klopp 2018). The resulting

reorganized genomes for both species were separately re-annotated as described above, and

the final annotations were resolved using MAKER to the reorganized *N. thermarum* genome,

with positional homology used to apply the annotation to the *N. dimorpha* genome assembly.

For each species, assembled transcripts were then generated using the resolved annotation

and genome assembly of each species, resulting in a set of homologous *N. thermarum*

transcripts and a set of *N. dimorpha* transcripts.

For *N. thermarum* and *N. dimorpha* transcripts, homology to *Arabidopsis thaliana* was

determined by blastx searching the *N. thermarum* transcripts against the TAIR11 protein set (e-

value cut-off set at 1e-4). The top *A. thaliana* blastx hit for each *N. thermarum* transcript was

selected as the putative homolog. For each *N. dimorpha* transcript, the putative *N. thermarum*

homolog was similarly identified with a blastx search (e-value cut-off set at 1e-4), and the

corresponding *A. thaliana* homologs was assigned to the *N. dimorpha* transcript.

DNA methylation-sensitive sequencing and analysis

DNA was extracted from endosperm using the QIAamp micro kit (Qiagen, cat# 56304).

For enzymatic methyl conversion sequencing and library preparation, an NEBNext Enzymatic

Methyl-seq kit was used; one additional AMPure bead clean-up was performed on libraries to

remove primer dimer. Sequencing was performed at the Whitehead Institute Genome

Technology Core. Libraries were pooled and sequenced across 2 lanes of a NovaSeq SP

flowcell (50 bp, paired-end reads) (endosperm samples) or 2 lanes of a NovaSeq S4 flowcell

(150 bp, paired-end) (leaf samples) to give ~14x genome coverage. Enzymatic-methyl

sequencing conversion rate was assessed prior to sequencing (Supplementary Materials and

Methods). Conversion rates were calculated using CyMATE (Hetzl et al. 2007). Sample

conversion rate averaged 99.85%.

Reads from enzymatic-converted samples were first mapped to a concatenation of the

originally produced *N. thermarum* and *N. dimorpha* assemblies and annotations, using Bismark

(version 0.22.3) (Krueger and Andrews 2011). 150 bp reads of leaf samples were broken into 40

bp segments and all reads were treated as single-end during mapping to ensure consistency in

data processing. The reads that uniquely mapped to either species' genome were sorted into

separate sets of *N. thermarum* or *N. dimorpha* reads and mapped to their respective species'

reorganized genome annotation with Bismark and methylation data was extracted. Analysis of

average DNA methylation 5', 3' and interior of features was performed using previously

developed custom pipelines (Pignatta et al. 2014). Differentially methylated regions (DMRs)

between samples were identified in the CG, CHG, and CHH contexts using a previously

developed pipeline (Pignatta et al. 2014). DMRs were defined as 300-bp windows for which 3 or

more cytosines with a coverage of 5 or more reads had a methylation difference of 35% or

greater between samples for CG and CHG contexts and 10% or greater for the CHH context,

with a Fisher's exact test with Benjamini-Hochberg correction p-value cutoff of 0.01 to determine

significance. DMRs were called between all combinations of biological replicates. For total

number of DMRs between endosperm maternal and paternal alleles, the number of DMRs was

averaged across all replicate comparisons. Genes and repeat regions were identified as

associated with a DMR if the gene or repeat region had a DMR within the annotated region or

+/-2 kb.

GO enrichment analysis was performed using ShinyGO 0.77 (Ge et al. 202). Putative

Arabidopsis homologs of all transcripts were used, and the set of putative Arabidopsis homologs

of all transcripts expressed during seed development (TPM > 1) (Povilus and Friedman 2022)

was used as the background set.

RNA sequencing and data analysis

For mRNA sequencing, RNA was extracted from frozen endosperm samples via

RNAqueous Total RNA Isolation Kit (Invitrogen) according to the kit protocol. Libraries were

prepared and sequenced at the MIT BioMicroCenter via NEBNext Ultra II Directional RNA

Library Prep Kit for Illumina (polyA-based isolation). Samples were pooled and sequenced on

one NovaSeq S4 flowcell (50bp, single end reads).

For full analysis methods, see Supplementary Materials and Methods. Briefly, for initial

analysis of gene expression, reads from all hybrid samples were mapped to the concatenated

genomes of the originally produced *N. thermarum* and *N. dimorpha* assemblies and

annotations; reads from non-hybrid samples were mapped to the reorganized genome of their

respective species. For identification of imprinted genes in hybrid samples, the reads that

uniquely mapped to either species' genome were sorted into separate sets of *N. thermarum* or

N. dimorpha reads and were used for subsequent analysis. *N. thermarum* reads were mapped

to the reorganized *N. thermarum* genome annotation, *N. dimorpha* reads were mapped to the

reorganized *N. dimorpha* genome annotation. Resulting allele-specific count tables for each

transcript were used for calling genetic imprinting. Genetically imprinted genes were called as

previously described (Gehring et al. 2011; Pignatta et al. 2014), using a pairwise comparison of

all possible combinations of each hybrid cross sample. For each gene we tested whether there

was a significant difference (Benjamini-Hochberg adjusted p-value < 0.01) between p_1 and p_2,

where p_1 is the portion of Nt reads for a gene in a Nt x Nd cross and p_2 is the portion of Nt reads

for the same gene in a Nd x Nt cross. While mapping reads, a slight maternal expression bias

was noted for both cross directions (Supplemental Table 1). Therefore, when calling imprinted

genes, the expected maternal : paternal expression ratio was adjusted from 1 (the anticipated

null ratio for diploid endosperm) to the average maximum observed maternal expression bias of

1.32 (null hypothesis: p_1=1.32, p_2=0.57). To increase stringency, minimum allelic-specific read

count was set to 50, a minimum imprinting factor was set to 2, and a maximum cis-effect factor

was set to 15. The imprinting factor is a measure of the magnitude of imprinting. For each gene

in a sample, a 95% confidence interval was determined around the Nt/Nd read ratio; the

imprinting factor is the low value of the high confidence interval divided by the high value of the

low confidence interval for the reciprocal cross (Gehring et al. 2011). The *cis*-effect factor is

calculated in a similar manner. In addition to these specifications, MEGs were required to have

a minimum of 70% maternal allele reads and PEGs were required to a maximum of 30%

maternal allele reads in both cross directions. In order for a gene to be considered as

consistently imprinted, it had to be called as imprinted in at least 75% (3 of 4) of pairwise

comparisons.

For correction of endosperm reads to account for potential maternal tissue

contamination, we mapped reads from endosperm samples (this study) and whole-seed

samples (Povilus and Friedman 2022) to the reorganized *N. thermarum* genome and proceeded

as described in (Tonosaki et al. 2024) (Supplementary Materials and Methods).

Differential gene expression analysis between endosperm of hybrid crosses and non-

hybrid endosperm was performed by mapping reads to the concatenated genomes as described

above, and then mapping uniquely mapping reads to their respective reorganized genome using

kallisto (v 0.46.1) (Bray et al. 2016). Differential gene expression analysis was performed using

DEseq2 (Love et al. 2014) using mostly default settings and filtering for loci with adjusted p

value less than or equal to 0.01 and mean TPM (of all samples) greater than or equal to 10.

GO enrichment analysis was performed using ShinyGO 0.77 (Ge et al. 2020), using

default settings. Putative Arabidopsis homologs of all transcripts in the test set were used, and

the set of putative Arabidopsis homologs of all transcripts expressed during seed development

(TPM > 1) (Povilus and Friedman 2022) was used as the background set.

In situ hybridizations and histology

In situ hybridizations were performed as previously described (Pignatta et al. 2018) (see

Supplementary Materials and Methods for probe information). Preparation of seed samples for

histological analysis was performed as previously described for seeds of *A. thaliana* (Pignatta et

al. 2018) and stained with toluidine blue, with adaptations of incubation times as necessary. All

samples were sectioned on a Leica RM 2065 rotary microtome at a thickness of 8 μm and

imaged using a Zeiss Axio Imager M2. Image tiling, color and brightness/contrast adjustments

and Smart Sharpen were applied to whole images, with particular attention to having even

contrast and white-balance across different images (Adobe Photoshop).

Acknowledgments

Funding was provided by the National Science Foundation grants MCB 2101337 to M. G., MCB

1453459 to M. G., IOS 1812116 to R.A.P, and a NSF Graduate Research Fellowship to C.A.M.

Sequencing was performed at the MIT BioMicro Center and the Whitehead Institute Genome

Technology Core.

References

Angeles-Núñez JG, Tiessen A. 2010. Arabidopsis sucrose synthase 2 and 3 modulate metabolic homeostasis and direct carbon towards starch synthesis in developing seeds. *Planta* 232(3): 701-718. 10.1007/s00425-010-1207-9.

Birchler JA. 1993. Dosage analysis of maize endosperm development. *Annu. Rev. Genet.* 27: 181-204. 10.1146/annurev.ge.27.120193.001145.

Borg M, Jacob Y, Susaki D, LeBlanc C, Buendía D, Axelsson E, Kawashima T, Voigt P, Boavida L, Becker J, et al. 2020. Targeted reprogramming of H3K27me3 resets epigenetic memory in plant paternal chromatin. *Nat. Cell Biol.* 22(6): 621-629. 10.1038/s41556-020-0515-y.

Borsch T, Löhne C, Mbaye M, Wiersema L. 2011. Towards a complete species tree of *Nymphaea*: shedding further light on subg. Brachyceras and its relationship to the Australian water-lilies. *Telopea* 13: 193-217. ISSN (Electronic): 2200-4025.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-sq quantification. *Nature Biotech.* 34: 525-547. 10.1038/nbt.3519.

Burkart-Waco D, Ngo K, Liberman M, Comai L. 2015. Perturbation of parentally biased gene expression during interspecific hybridization. *PLoS One* 10(2): e0117293. 10.1371/journal.pone.0117293.

Cabanettes F, Klopp C. 2018. D-GENIES: dot plot large genomes in an interactive, efficient and simple way. *PeerJ* 6 :e4958. 10.7717/peerj.4958.

Campbell MS, Holt C, Moore B, Yandell M. 2014. Genome Annotation and Curation Using MAKER and MAKER-P. *Curr. Protoc. Bioinformatics* 48: 4.11.1–4.11.39. 10.1002/0471250953.bi0411s48.

Coimbra S, Salema R. 1999. Ultrastructure of the developing and fertilized embryo sac of *Amaranthus hypochandriacus L*. *Ann. Bot.* 84: 781-789. 10.1006/anbo.1999.0978.

Florez-Rueda AM, Paris M, Schmidt A, Widmer A, Grossniklaus U, Städler T. 2016. Genomic
imprinting in the endosperm is systematically perturbed in abortive hybrid tomato seeds. *Mol.*
Biol. Evo. 33(11): 2935-2946. 10.1093/molbev/msw175.

Florez-Rueda AM, Sharmann M, de Souza LP, Fernie AR, Bachelier JB, Figueiredo DD. 2024.
Genomic imprinting in an early-diverging angiosperm reveals ancient mechanisms for seed
initiation in flowering plants. *bioRxiv*. 10.1101/2024.09.18.613646.

Friedman WE. 2006. Embryological evidence for developmental lability during early angiosperm
evolution. *Nature* 441: 337-340. 10.1038/nature04690.

Friedman WE. 2008. Hydatellaceae are water lilies with gymnospermous tendencies. *Nature*
453: 94-97. 10.1038/nature06733.

Ge SX, Jung D, Yao R. 2020. ShinyGo: a graphical gene-set enrichment tool for animals and
plants. *Bioinformatics* 36(8): 2628-2629. 10.1093/bioinformatics/btz931.

Gehring M, Huh JH, Hsieh T, Penterman J, Choi Y, Harada JJ, Goldberg RB, Fischer RL. 2006.
DEMETER RNA glycosylase established MEDEA polycomb gene self-imprinting by allele-
specific demethylation. *Cell* 124(3):495-506. 10.1016/j.cell.2005.12.034.

Gehring M, Bubb KL, Henikoff S. 2009. Extensive demethylation of repetitive elements during
seed development underlies gene imprinting. *Science* 324(5933): 1447-1451.
10.1126/science.1171609.

Gehring M, Missirian V, Henikoff S. 2011. Genomic analysis of parent-of-origin allelic
expression in Arabidopsis thaliana seeds. *PLoS ONE* 6(8): e23687l.
10.1371/journal.pone.0023687.

Gent JI, HIggins KM, Swentowsky KW, Fu F, Zeng Y, Kim DW, Dawe RK, Springer NM,
Anderson SN. 2022. The maize gene maternal derepression of r1 encodes a DNA glycosylase
that demethylates DNA and reduces siRNA expression in the endosperm. Plant Cell 34(10):
3685-3701. 10.1093/plcell/koac199.

Haig D, Westoby M. 1991. Genomic imprinting in endosperm: its effect on seed development in
crosses between species, and its implications for the evolution of apomixis. *Philos. Trans. R.*
Soc. B 333(1266): 1-13. 10.1098/rstb.1991.0057.

Hetzl J, Foerster AM, Raidl G, Scheid OM. 2007. CyMATE: a new tool for methylation analysis
of plant genomic DNA after bisulphite sequencing. *Plant J.* 51(3): 526-536. 10.1111/j.1365-
313X.2007.03152.x.

Hsieh T, Ibarra CA, Silva P, Zemach A, Eshed-Williams L, Fischer RL, Zilberman D. 2009.
Genome-wide demethylation of Arabidopsis endosperm. *Science* 324(5933): 1451-1454.
10.1126/science.1172417.

Inoue A, Jiang L, Falong L, Suzuki T, Zhang Y. 2017. Maternal H3K27me3 controls DNA
methylation-independent imprinting. *Nature* 547:419-424. 10.1038/nature23262.

Josefsson C, Dilkes B, Comai L. 2006. Parent-dependent loss of gene silencing during
interpsecies hybridization. *Curr. Biol.* 16(13): 1322-1328. 10.1016/j.cub.2006.05.045.

Kinoshita T., Miura A, Choi Y, Kinoshita Y, Cao X, Jacobsen SE, Fischer RL, Kakutani T. 2004. One-way control of FWA imprinting in Arabidopsis endosperm by DNA methylation. *Science* 303(5657): 521-523. 10.1126/science.1089835.

Klosinska M, Picard CL, Gehring M. 2016. Conserved imprinting associated with unique epigenetic signatures in the *Arabidopsis* genus. *Nat. Plants* 2: 16145. 10.1038/nplants.2016.145.

Koren S, Walenz BP, Berlin K, Miller JR, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res.* 27: 722-736. 10.1101/gr.215087.116.

Korf I. 2004. Gene finding in novel genomes. *BMC Bioinform.* 5: 59. 10.1186/1471-2105-5-59.

Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* 27(11): 1571-1572. 10.1093/bioinformatics/btr167.

Lersten NR. 2004. Endosperm. In: Lersten NR, editor. In Flowering Plant Embryology. Blackwell Publishing. p. 161–163.

Lewis A, Mitsuya K, Umlauf D, Smith P, Dean W, Walter J, Higgins M, Feil R, Reik W. 2004. Imprinting on distal chromosome 7 in the placenta involves repressive histone methylation independent of DNA methylation. *Nature Genet.* 36:1291-1295. 10.1038/ng1468.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34: 3094-3100. doi:10.1093/bioinformatics/bty191.

López-Fernández MP, Maldonado S. 2013. Programmed cell death during quinoa perisperm development. *J. Exp. Bot.* 64(11): 3313-3325. 10.1093/jxb/ert170.

Losada JM, Bachelier JB, Friedman WE. 2017. Prolonged embryogenesis in *Austrobaileya scandens* (Austrobaileyaceae): its ecological and evolutionary significance. *New Phyol.* 215(2): 851-864. 10.1111/nph.14621.

Love MI, Huber W, Anders S (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, **15**, 550. 10.1186/s13059-014-0550-8.

Moreno-Romero J, Del Toro-De León G, Yadav VK, Santos-González J, Köhler C. 2019. Epigenetic signatures associated with imprinted paternally expressed genes in the Arabidopsis endosperm. *Genom. Biol.* 20(1): 41.10.1186/s13059-019-1652-0.

Niederhuth CE, Bewick AJ, Ji L, Alabady MS, Kim KD, Li Q, Rohr NA, Rambani A, Burke JM, Udall JA, et al. 2016. Widespread natural variation of DNA methylation within angiosperms. *Genom. Biol.* 17: 194. 10.1186/s13059-016-1059-0.

Orban I, Bouharmont J. 1998. Megagametophyte development of *Nymphaea nouchali Burm. F.* (Nymphaeaceae). *Bot. J. Linn. Soc.* 126: 339-348. 10.1111/j.1095-8339.1998.tb01386.x.

Park K, Kim MY, Vickers M, Park J, Hyun Y, Okamoto T, Zilberman D, Fischer RL, Feng X, Choi Y, Scholten S. 2016. DNA demethylation is initiated in the central cells of Arabidopsis and rice. *Proc. Natl. Acad. Sci. U S A* 113(52): 15138-15143. 10.1073/pnas.1619047114.

Patten MM, Ross L, Curley JP, Queller DC, Bonduriansky R, Wolf JB. 2014. The evolution of
genomic imprinting: theories, predictions and empirical tests. *Heredity* 113: 119-128.
10.1038/hdy.2014.29.

Pei L, Zhang, L Li J, Shen C, Qiu P, Tu L, Zhang X, Wang M. 2019. Tracing the origin and
evolution history of methylation-related genes in plants. *BMC Plant Biol.* 19(1): 307.
10.1186/s12870-019-1923-7.

Pennington PD, Costa LM, Gutierrez-Marcos JF, Greenland AJ, Dickinson HG. 2008. When
Genomes collide; aberrant seed development following maize interploidy crosss. *Ann. Bot.* 101:
833-843. 10.1093/aob/mcn017.

Picard CL, Gehring M. 2020. Identification and comparison of imprinted genes across plant
species. In: Spillane C, McKeown P, editors. Plant Epigenetics and Epigenomics: Methods and
Protocols, Methods in Molecular Biology volume 2093. Springer Science+Business Media, LLC,
part of Springer Nature, p. 173-201. 10.1007/978-1-0716-0179-2_13.

Picard CL, Povilus RA, Williams BP, Gehring M. 2021. Transcriptional and imprinting complexity
in Arabidopsis seeds at single-nucleus resolution. *Nat. Plants* 7: 730-738. 10.1038/s41477-021-
00922-0.

Pignatta D, Erdmann RM, Scheer E, Picard CL, Bell GW, Gehring M. 2014. Natural epigenetic
polymorphisms lead to intraspecific variation in Arabidopsis gene imprinting. *Elife*, 3: e03198.
10.7554/eLife.03198.

Pignatta D, Gehring M. 2012. Imprinting meets genomics new insights and new challenges.
Curr. Opin. Plant Biol. 15(5): 530-535. 10.1016/j.pbi.2012.09.004.

Pignatta D, Novitzky K, Satyaki PRV, Gehring M. 2018. A variably imprinted epiallele impacts
seed development. *PLoS Genet.* 14(11): e1007469. 10.1371/journal.pgen.1007469.

Povilus RA, DaCosta JM, Grassa C, Satyaki PRV, Moeglein M, Jaenisch J, Xi Z, Mathews S,
Gehring M, Davis CC, Friedman WE. 2020. Water lily (*Nymphaea thermarum*) genome reveals
variable genomic signatures of ancient vascular cambium losses. *Proc. Natl. Acad. Sci. U S A*
117(15): 8649-8656. Doi.org/10.1073/pnas.1922873117.

Povilus RA, Diggle PK, Friedman WE. 2018. Evidence for parent-of-origin effects and
interparental conflict in seeds of an ancient flowering plant lineage. *Proc. Royal. Soc. B*
285(1872): 20172491. 10.1098/rspb.2017.2491.

Povilus RA, Friedman WE. 2022. Transcriptomes across fertilization and seed development in
the water lily *Nymphaea thermarum*: evidence for epigenetic patterning during reproduction.
Plant Reprod. 35(5): 161-178. 10.1007/s00497-022-00438-3.

Povilus RA, Gehring M. 2022. Maternal-filial transfer structures in endosperm: a nexus of
nutritional dynamics and seed development. *Curr. Opin. Plant Biol.* 65: 102121.
10.1016/j.pbi.2021.102121.

Povilus RA, Losada JM, Friedman WE. 2015. Floral biology and ovule and seed ontogeny of
Nymphaea thermarum, a water lily at the brink of extinction with potential as a model system for
basal angiosperms. *Ann. Bot.* 115: 211–226. 10.1093/aob/mcu235.

Raas MWD, Zijlmans DW, Vermeulen M, Marks H. 2022. There is another: H3K27-mediated
genomic imprnting. *Trends Genet.* 38(1):82-96. 10.1016/j.tig.2021.06.017.

Rodrigues JA, Ruan R, Nishimura T, Sharma MJ, Sharma R, Ronald PC, Fischer RL, Zilberman
D. 2013. Imprinted expression of genes and small RNA is associated with localized
hypomethylation of the maternal genome in rice endosperm. Proc. *Natl. Acad. Sci. U S A*
110(19): 7934-7939. 10.1073/pnas.1306164110.

Rudall PJ, Remizowa MV, Beer AS, Bradshaw E, Stevenson DW, Macfarlane TD, Tuckett RE,
Yadav SR, Sokoloff DD. 2008. Comparative ovule and megagametophyte development in
Hydatellaceae and water lilies reveal a mosaic of features among the earliest angiosperms.
Ann. Bot. 101(7): 941-956. 10.1093/aob/mcn032.

Santini L, Halbritter F, Titz-Teixeira F, Suzuki T, Asami M, Ma X, Ramesmayer J, Lackner A,
Warr N, Pauler F, Hippenmeyer S, Laue E, Karlik M, Bocj C, Beyer A, Perry ACF, Leeb M.
2021. Genomic imprinting in mouse blastocysts is predominantly associated with H3K27me3.
Nature Commun. 12:3804. 10.1038/s41467-021-23510-4.

Satyaki PRV, Gehring M. 2019. Paternally acting canonical RNA-directed DNA methylation
pathwa genes sensitize Arabidopsis endosperm to paternal genome dosage. *The Plant Cell*
31(7):1563-1578. 10.1105/tpc.19.00047.

Scott RJ, Spielman M, Bailey J, Dickinson HG, Parent-of-origin effects on seed development in
Arabidopsis thaliana. *Development* 125(17): 3329-3341. 10.1242/dev.125.17.3329.

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. 2006. AUGUSTUS: ab initio
prediction of alternative transcripts. *Nucleic Acids Res.* 34: W435–W439. 10.1093/nar/gkl200.

Tonosaki K, Ono A, Nagata H, Furuumi H, K Nonomura, Sato Y, Comai L, Hatakeyama K,
Kamakatsu T, Kinoshita T. 2024. Multi-layered epigenetic control of persistent and stage-
specific imprinted genes in rice endosperm. *Nat. Plants* 10: 1231–1245. 10.1038/s41477-024-
01754-4.

Waters AJ, Makarevitch I, Eichten SR, Swanson-Wagner RA, Yeh C, Xu W, Schnable PS,
Vaughn MW, Gehring M, Springer NM. 2011. Parent-of-origin effects on gene expression and
DNA methylation in the maize endosperm. *Plant Cell* 23(12): 4221-4233.
10.1105/tpc.111.092668.

Wick RR, Schultz MB, Zobel J, KE Holt. 2015. Bandage: interactive visualization of de novo
genome assemblies. *Bioinformatics* 31(20): 3350-3352. 10.1093/bioinformatics/btv383.

Williams JH, Friedman WE. 2002. Identification of diploid endosperm in an early angiosperm
lineage. *Nature* 415: 522-526. 10.1038/415522a.

Xin M, Yang R, Li G, Chen H, Laurie J, Ma C, Wang D, Yao Y, Larkins BA, Sun Q, et al. 2013.
Dynamic expression of imprinted genes associates with maternally controlled nutrient allocation
during maize endosperm development. *Plant Cell* 25(9): 3212-3227. 10.1105/tpc.113.115592.

Zhang M, Li N, He W, Zhang H, Yang W, Liu B. 2016. Genome-wide screen of genes imprinted
in sorghum endosperm, and the roles of allelic differential cytosine methylation. *Plant J.* 85(3):
424-436. doi: 10.1111/tpj.13116.

Zhang M, Shaojun X, Dong X, Zhao X, Zeng B, hen J, Li H, Yang W, Zhao H, Wang G, Chen Z,
SUn S. 2014. Genome-size high resolution parental-specific DNA and histone methylation maps
uncover patterns of imprinting regulation in maize. *Genome Res.* 24:167-176.
10.1101/gr.155879.113

Zhang Z, Yu S, Jing L, Zhu Y, Jiang S, Xia H, Zhou Y, Sun D, Liu M, Li C, et al. 2021.
Epigenetic modifications potentially controlling the allelic expression of imprinted genes in
sunflower endosperm. *BMC Plant Biol.* 21(1): 570. 10.1186/s12870-021-03344-4.

Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA genome
assembler. *Bioinformatics* 29(21): 2669-6977. 10.1093/bioinformatics/btt476.

733
734
735 **Figures and Supplementary Figures and Tables**

736
737

Figure 1

= maternal tissue (seed coat)

= maternal tissue (perisperm)

= endosperm

= embryo

= *Nymphaea thermarum (Nt)*

= *Nymphaea dimorpha (Nd)*

400 µm

Nt
self-fertilized

Nd
self-fertilized

Nt x *Nd*
cross

Nd x *Nt*
cross

740
741
Figure 1: Seed structure in water lilies and F1 hybrids. Young seeds (9-10 days after anthesis) of self-
fertilized *N. thermarum* (*Nt*) and *N. dimorpha* (*Nd*), as well as reciprocal crosses between the two
species. In all seeds, young embryos are surrounded by cellular, diploid endosperm, which in turn is
surrounded by a maternal nutrient storage tissue (perisperm).
746
747

Figure 2: Genetic imprinting in water lily endosperm. A) Principal component analysis of RNA-seq data
from young endosperm samples isolated from self-fertilized *N. thermaurm* (Nt) and *N. dimorpha* (Nd)
seeds, and reciprocal crosses of *N. thermaurm* and *N. dimorpha*. B) Plot of the ratio of maternal to
paternal allele reads in endosperm of one reciprocal cross comparison. Female in cross listed first. Genes
that pass the stringency cut-offs for being called as imprinted are highlighted in red (MEGs, maternally
expressed genes) or blue (PEGs, paternally expressed genes). Gray dots indicate genes that are not
called as imprinted. C) Upset plot showing the number and consistency of genes called as MEGs or PEGs
across comparisons of different replicates. MEGs and PEGs called in at least 75% of comparisons are
highlighted in blue (PEGs) or red (MEGs). D) Average expression and average log2 fold change of genes
expressed > 1 TPM, in the comparison of Nt self-fertilized and Nt x Nd hybrid endosperm. Genes are
color-coded according to whether they are significantly differentially expressed in no comparisons or
consistently (75% or more of comparisons), and/or were identified as a MEG in no samples, in any
samples, or consistently (75% or more of samples). Number of genes in each category is noted. Similar
graphs for individual comparison types are shown in Supplementary Figure 3. E) Average expression and
average percent maternal expression for all expressed genes (TPM > 1). Genes called as MEGs in at least
75% of replicate comparisons are shown in red, with putative *Arabidopsis thaliana* homology indicated
for some MEGs of interest. F) In situ hybridization of putative homolog of *GA2OX2*. Inset shows
magnification of endosperm of sample treated with antisense probe. Scalebars = 200 µm. G) In situ
hybridization of putative homolog of *SUS3*. Inset shows magnification of endosperm of sample treated
with antisense probe. Scalebars = 200 µm. H) GO-enrichment analysis (for biological process terms) of
genes consistently called as MEG in the endosperm, reported with FDR-adjusted p-value and fold-
enrichment.

773
774

Figure 3: DNA methylation in water lily endosperm. A) Average parental allele cytosine methylation

across genes or repeats in young endosperm of reciprocal crosses between *N. thermarum* and *N.*

dimorpha, aligned at either the 5' transcription start site or 3' transcription end site. Each track

represents the maternal or paternal alleles of an individual sampled cross (N=3 for *N. thermarum* x *N.*

dimorpha crosses, N = 2 for *N. dimorpha* x *N. thermarum* crosses). Tracks are grouped by species and

color-coded to indicate whether the track represents the maternally- or paternally-inherited alleles. B)

Average number of maternally- or paternally-hypermethylated DMRs that overlap a gene and/or a

repeat region, for each species. DMRs were called by comparing the genomes of each species when

they were maternally- or paternally-inherited , for hybrid endosperm and leaf tissue (only one cross

direction performed for leaf tissue). Red bars (above 0) indicate maternal hypermethylation, blue bars

(below 0) indicate paternal hypermethylation. Bars are further color-coded to indicate the proportion of

DMRs that overlap a gene, a gene and repeat, a repeat, or neither gene nor repeat. C) Number of genes

(+/- 2kb) that consistently overlap maternally- or paternally-hypermethylated DMRs (gene had to have

at least one DMR overlap of the indicated type in at least 75% of comparisons, as well as have an

overlapping DMR of the opposite type in no more than 25% of comparisons). D) Genome browser

snapshots of DNA methylation for homologs of *GA2OX2* and *TMO6*, showing examples of methylation

patterning (blue = CG, orange = CHG, and green = CHH) on maternal or paternal genomes, for each

species as the maternal and paternal genome, in endosperm and leaf tissue.

Figure 4: Two hypotheses regarding the evolution of genetic imprinting in endosperm, endosperm ploidy, and nutrient storage strategies in angiosperm seeds. Hypothesis 1 posits that MEGs and PEGs were originally present in endosperm of the last common ancestor of angiosperms, and that PEGs were lost in the water lily lineage in association with the origin of perisperm. Hypothesis 2 posits that MEGs evolved as a maternal response to the addition of a paternal genome complement in a nutrient-mediating tissue, while PEGs evolved as a response to the origin of triploid endosperm, which features the addition of a second maternal genome complement. Dashed lines indicate events hypothesized as a result of this study. Seed diagrams show diversity of mature seed structures, highlighting diversity in developmental origin of the primary site of nutrient storage (*Nymphaea* = maternal perisperm; *Zea* = endosperm; *Arabidopsis* = embryo).

(See Supplementary Information PDF) Supplementary Figure 1: Image of individuals of *N. thermarum*, *N. dimorpha*, and an F1 hybrid.
Top left = *N. thermarum*, top right = *N. dimorpha*, bottom = F1 hybrid (demonstrating viability of hybrid crosses). Identity of individuals was confirmed by genotyping (EM seq samples s104, 105, s106, see Supplementary Table 1).

(See Supplementary Information PDF) Supplementary Figure 2: Genome assemblies for *N. thermarum* and *N. dimorpha*. A) Bandage diagrams for the updated *N. thermarum* genome assembly and the novel *N. dimorpha* genome assembly. B) Alignment of the *N. thermarum* genome mapped against the *N. dimorpha* genome.

(See Supplementary Information PDF) Supplementary Figure 3: Differential gene expression analysis between hybrid and parental endosperm with MEGs highlighted. A) Comparison of fold change and averaged expression (TPM) for *N. dimorpha* self-fertilized endosperm vs. *N. thermarm* x *N. dimorpha* hybrid endosperm. B) Comparison of fold change and averaged expression (TPM) for *N. dimorpha* self-fertilized endosperm vs. *N. dimorpha* x *N. thermarum* hybrid endosperm. C) Comparison of fold change and averaged expression (TPM) for *N. thermarum* self-fertilized endosperm vs. *N. thermarm* x *N dimorpha* hybrid endosperm. D) Comparison of fold change and averaged expression (TPM) for *N. thermarum* self-fertilized endosperm vs. *N. dimorpha* x *N. thermarum* hybrid endosperm. E) Upset graph showing overlap of MEGs that are significantly DE for the the four different comparison types; the putative *Arabidopsis thaliana* homologs of the 5 consistently DE MEGs is listed.

(See Supplementary Information PDF) Supplementary Figure 4: In situ hybridization positive control for perisperm, and *SUS3* expression in seeds of different crosses and self-fertilizations. A) RNA *in situ* hybridization of putative homolog of a terpene synthase sub-family in *N. thermarum* seeds, showing that detection of gene expression in the perisperm is possible. Results of experiments performed with antisense and sense probes are shown. Scalebars = 200 μm. B) In situ hybridization of putative homolog of *SUS3*, performed in seeds of reciprocal crosses of *N. thermarum* and *N. dimorpha* and in seeds from *N. dimorpha* self-fertilizations. Black arrowheads indicate detection of signal, white arrowheads indicate absence of signal. Scalebars = 200 μm. Results of experiments performed with antisense and sense probes are shown.

(See Supplementary Information PDF) Supplementary Figure 5: DNA methylation, including leaf tissues from non-hybrid *N, thermarum* and *N. dimorpha* plants

A) Number of DMRs that are hyper- or hypomethylated in the *N. dimorpha* compared to *N. thermarum* genome, that overlap a gene and/or a repeat region, for each species, in leaf tissue. DMRs were called by comparing the genomes of each species within the same sample (hybrid tissue, one sample) or between samples of leaves of each parental species (one sample of each species). B) Genome browser snapshots of DNA methylation for homologs of *GA2OX2* and *TMO6*, showing examples of methylation patterning (blue = CG, red = CHG, and green = CHH) on the genome for each species , in endosperm and leaf tissue, including leaves of non-hybrid *N, thermarum* and *N. dimorpha* plants. Black notches indicate cytosines for which there was sufficient data to include, but were unmethylated.

Supplementary Table 1: Summaries for sequencing, mapping, and genome assemblies and annotations.

Supplementary Table 2: Results and summaries of imprinting calling tests, report of imprinted loci that overlap DMRs, and summary of imprinting status of *Nymphaea* homologs of conserved imprinted genes.

Supplementary Table 3: Results and summaries of differential gene expression analysis between intraspecies and interspecies crosses

Supplementary Table 4: Results and summaries of imprinting calling tests, when data is adjusted for potential maternal tissue contamination of endosperm samples (assuming 50% and 25% contamination).

Supplementary Table 5: Report of DMR overlap with genomic features, as well as report and summary of genes with DMR overlaps