1   **Title: Optimized CART Cell Therapy for Metastatic Aggressive Thyroid Cancer**

3   **Authors:**   Claudia Manriquez Roman[1,2,3†], Elizabeth L. Siegler[2,3†], Justyna J. Gleba[4†], Truc
4   N. Huynh[2,3], Grace E. DeFranco[2,5,6], Aylin Alasonyalilar Demirer[4], Matthew L.
5   Pawlush[4], Michael Redig[4], Skyeler M. Klinge[4], Long K. Mai[2,3], , James L. Miller[4],
6   Erin E. Miller[4], Brooke L. Kimball[2,3], Erin E. Tapper[2,3], R. Leo Sakemura[2,3], Carli M.
7   Stewart[2,6,7], Ismail Can[2,3], Olivia L. Sirpilla[2,3], Jennifer M. Feigin[2,6,8], Kun Yun[2,6,9],
8   Omar L. Gutierrez Ruiz[2,3], Hong Xia[2,3], Mehrdad Hefazi Torghabeh[2,3], Kendall J.
9   Schick[2], Ekene J. Ogbodo[2,3], Gloria Olivier[10], Yushi Qiu[4], Robert C. Smallridge[11],
10   Abba Zubair[12], Han W. Tun[13], John A. Copland[4,8,9*], and Saad S. Kenderian[1,2,3,5,6,9*]

Affiliations:
[1] Center for Regenerative Biotherapeutics, Mayo Clinic; Rochester, MN
[2] T Cell Engineering Laboratory, Mayo Clinic; Rochester, MN
[3] Division of Hematology, Mayo Clinic; Rochester, MN
[4] Cancer Biology Department, Mayo Clinic; Jacksonville, FL
[5] Department of Immunology, Mayo Clinic; Rochester, MN
[6] Mayo Clinic Graduate School of Biomedical Sciences; Rochester, MN
[7] Department of Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic;
Rochester, MN
[8] Department of Biochemistry and Molecular Biology, Mayo Clinic; Rochester, MN
[9] Department of Molecular Medicine, Mayo Clinic; Rochester, MN
[10] Department of Business Development, Mayo Clinic; Rochester, MN
[11] Division of Endocrinology, Internal Medicine Department, Mayo Clinic; Jacksonville, FL
[12] Department of Laboratory Medicine and Pathology, Mayo Clinic; Jacksonville FL
[13] Hematology/Oncology Division, Department of Medicine, Mayo Clinic; Jacksonville, FL
27

28   † These authors contributed equally to this work

29   * Corresponding authors: Saad S. Kenderian, Mayo Clinic Rochester, 200 First Street S.W.,
30   Rochester, MN Kenderian.saad@mayo.edu, *or* John A. Copland, Mayo Clinic Florida,
31   copland.john@mayo.edu

33   **Running Title: TSHR-CART cell therapy in thyroid cancers**
34

Acknowledgements: This study was partly funded through the following:
Mayo Clinic Center for Regenerative Biotherapeutics (JAC and SSK)
Mayo Clinic Comprehensive Cancer Center (SSK)
National Institutes of Health R37CA266344 and R01AI79974 (SSK)
Department of Defense grant CA201127 (SSK)
Generous support from Renee Cousins King, M.D.

Competing Interests: SSK is an inventor on patents in the field of CAR immunotherapy that are licensed to Novartis (through an agreement between Mayo Clinic, University of Pennsylvania, and Novartis). SSK and RLS are inventors on patents that are licensed to Humanigen/Taran (through Mayo Clinic). SSK is an inventor on patents that are licensed to MustangBio (through Mayo Clinic). ELS, EET, RLS, OLS, KJS, and SSK are inventors on patents that are licensed to Chymal (through Mayo Clinic). CMR, ELS, TNH, LKM, BLK, RLS, CMS, IC, OLS, JMF, KY, OGR, HX, MHT, JAC and SSK are inventors on patents licensed to Immix (through Mayo Clinic). SSK receives research funding from Kite, Gilead, Juno, BMS, Novartis, Humanigen/Taran, MorphoSys, Tolero, Sunesis/Viracta, LifEngine Animal Health Laboratories Inc, and Lentigen. SSK has participated in advisory meetings with Kite/Gilead, Calibr, Luminary Therapeutics, Humanigen, Juno/BMS, Capstan Bio, Carisma, and Novartis. SSK has served on the data safety and monitoring board with Humanigen. SSK has served as a consultant for Torque, Calibr, Novartis, Capstan Bio, and Humanigen

ABSTRACT

Most thyroid cancer deaths are attributed to a subset of poorly differentiated, metastatic tumors. To improve treatment options for aggressive thyroid cancers, we developed a novel thyroid-stimulating hormone receptor (TSHR)-targeted chimeric antigen receptor T (CART) cell therapy, which demonstrated antigen-specific activation and antitumor efficacy against TSHR$^+$ cell lines *in vitro* and *in vivo*. However, de-differentiated thyroid cancers downregulate TSHR. We therefore developed a potent treatment strategy by combining our novel TSHR-CART cells with mitogen-activated protein kinase (MAPK) inhibitors, which redifferentiate thyroid tumors and upregulate TSHR expression. In patient-derived anaplastic thyroid cancer xenografts, combination therapy of TSHR-CART cells and MAPK inhibitors led to increased TSHR expression on the tumor tissue and significantly enhanced antitumor efficacy and prolonged survival compared to TSHR-CART monotherapy. Based on our data, we are launching a phase I clinical trial for TSHR-CART cell therapy alone or in combination with MAPK inhibitors in patients with metastatic thyroid cancers.

STATEMENT OF SIGNIFICANCE

Poor target selection and antigen escape limit CART cell efficacy in solid tumors. We developed TSHR-CART cells to treat differentiated thyroid cancers but observed TSHR downregulation in dedifferentiated thyroid cancers. We found that MAPK inhibitors restored TSHR expression and sensitized these cancers to TSHR-CART cell therapy.

INTRODUCTION

Thyroid cancer is the most common endocrine cancer, and its incidence is rising in the US and globally.(1, 2) Thyroid cancer is projected to become the fourth leading type of cancer across the world, and the causes appear to be complex and multifactorial, including dietary changes and environmental exposures. Patients with thyroid cancer have widely different clinical outcomes depending on the pathological subtype and mutation profile. Metastatic differentiated thyroid cancer (DTC), poorly differentiated thyroid cancer (PDTC), and anaplastic thyroid cancer (ATC) account for a majority of thyroid cancer-related deaths, even though these subtypes only represent 2-3% of all thyroid cancer diagnoses.(1, 3, 4) Hurthle cell carcinoma (HCC, now termed oncocytic thyroid cancer(5)) is a subtype of DTC known to be aggressive, highly metastatic, and have lower rates of overall survival than follicular thyroid cancers. HCC is often resistant to radioiodine therapy and has no current FDA-approved treatment options.(6, 7) PDTC and ATC are aggressive and often refractory to radioiodine therapy and other standard therapies.(8, 9) SEER database analysis through 2015 reported a median overall survival of 4 months in patients with ATC, with disease-specific mortality of over 98%. A more recent analysis reported a median overall survival of 15.7 months in patients with ATC receiving newer targeted therapies.(10) Despite these advancements in treatment options, PDTC and ATC remain incredibly lethal.

Chimeric antigen receptor (CAR) T cell therapy has emerged as a potentially curative therapy in a subset of patients with hematological malignancies and has been FDA-approved for several B cell malignancies as well as for multiple myeloma.(11-15) With the advent of adoptively transferred engineered cellular therapies, there is a compelling rationale to apply CART cell therapy to treatment-resistant solid tumors. To date, the efficacy of CART cell therapy in solid tumors has been modest at best.(16-21) Due to a lack of tumor-specific targets for CART cell therapy in solid tumors, identifying an antigen that is both uniquely and universally expressed on cancer cells is critical for development of antigen-specific immunotherapy to avoid on-target off-tumor toxicities.(20, 22, 23)

Thyroid-stimulating hormone receptor (TSHR) is a surface glycoprotein receptor and a major

regulator of thyroid function and metabolism. The TSHR mediates the activating mechanism of

TSH to the thyroid gland, resulting in the growth and proliferation of thyrocytes as well as thyroid

hormone production, including thyroxine (T4) and triiodothyronine (T3).(*24*) TSHR is

predominantly expressed on the basolateral membrane of normal thyroid follicular cells.(*25*)

TSHR expression is largely limited to the thyroid gland, with low levels of expression in additional

tissues, including the thymus and testes (immune privileged) according to Human Protein Atlas

and NIH GenBank databases. TSHR is abundantly expressed on thyroid tumor cells.(*25, 26*)

Given its unique expression, TSHR is recognized as a compelling target for advanced thyroid

cancer diagnostics and therapy after thyroidectomy. Overall, there is limited risk of off-tumor

adverse effects of targeting TSHR. TSHR has been targeted using biologics including monoclonal

antibodies for the treatment of Graves' disease and thyroid cancer.(*27*) In this study, we

generated CART cells targeting TSHR (TSHR-CART) and showed potent, antigen-specific

antitumor efficacy against TSHR-expressing cell lines *in vitro* and *in vivo,* as well as in patient-

derived xenograft (PDX) mouse models of ATC.

Further, we identified mitogen-activated protein kinase (MAPK) inhibition as a strategy to

upregulate TSHR in dedifferentiated thyroid cancers and increase the therapeutic index of TSHR-

CART cell therapy. Downregulation of TSHR expression in dedifferentiated thyroid cancer is a

potential mechanism of resistance to TSHR-CART cell therapy. MAPK inhibition is an FDA-

approved treatment against thyroid cancer and is documented to upregulate TSHR expression in

patients with advanced metastatic thyroid cancer.(*28, 29*) Clinical trials for metastatic thyroid

cancers have demonstrated that MAPK inhibition induces re-expression of genes necessary for

iodine uptake and retention, including TSHR and sodium iodide symporter (NIS).(*30, 31*)

Additionally, many thyroid cancers have gene mutations involving the constitutive activation of

BRAF, which stimulates MAPK signaling.(*32-34*) Therefore, when BRAF activation is switched

off genetically, or its downstream signaling is inhibited with MEK inhibitors (MEKi) or BRAF

inhibitors (BRAFi), tumors re-express TSHR and NIS and reaccumulate iodine. This strategy has

been employed in the clinic to enhance the response to radioactive iodine therapy.(*29*)

We therefore sought to address antigen escape in ATC by testing the combination of MEK/BRAF

small molecule inhibitors with TSHR-CART cell therapy. We found that MEK/BRAF inhibition led

to restoration of TSHR expression on thyroid cancer cells and enhancement of CART cell

expansion and antitumor activity *in vivo*. This strategy is highly translatable to patients with

advanced thyroid cancers who have extremely limited treatment options to date.

RESULTS

TSHR as a target for CART cell therapy

To validate TSHR as a target candidate for CART cell therapy in thyroid cancer, we verified its

expression on thyroid tissue and lack of expression on other normal tissues by

immunohistochemistry (IHC) analysis of specimens from the Mayo Clinic Thyroid Cancer Biobank

and on publicly available databases.(*35*) IHC analysis showed that, while TSHR expression was

high in normal thyroid tissue as well as multiple thyroid cancer histotypes, ATC samples

demonstrated attenuated TSHR expression (**Fig. 1A, B**). We then assessed TSHR expression in

a healthy tissue microarray and confirmed that TSHR expression was high in the thyroid and low

or absent in other tissues including adult thymus and testis (**Fig. 1C, Supplementary Fig. S1**).

Having validated TSHR as a potential antigen to target with CART cell therapy, we designed and

generated TSHR-CART cells using a single chain variable fragment isolated from the TSHR

autoantibody clone K1-70 and cloned into a lentiviral CAR construct containing 4-1BB and CD3ζ

(**Supplementary Fig. S2A**). Healthy donor T cells transduced with CAR-encoded lentiviral

particles (**Supplementary Fig. S2B**) demonstrated high CAR expression as assessed by flow

cytometry (**Fig. 1D**). TSHR-CART cells displayed comparable CD4:CD8 ratio and phenotype

compared to control untransduced T cells (UTD) (**Supplementary Fig. S2C-D**).

TSHR-CART cells exhibit potent and specific antitumor activity *in vitro*

We next examined the antigen-specific activation of TSHR-CART cells *in vitro* and *in vivo*. TSHR-

CART cells or UTD were incubated with the follicular thyroid cancer cell line, FTC-133, which was

transduced to overexpress TSHR; TSHR was transduced in these cells as thyroid cancer cell

lines and patient samples downregulate TSHR expression after extended monolayer cell

culture.(*36-39*) While TSHR-CART cells were not activated against wildtype (TSHR-) FTC-133

cells, TSHR-CART cells demonstrated profound antigen-specific effector functions against

TSHR+ target cells *in vitro*, including cytotoxicity against target cells (**Fig. 1E**) as well as TSHR-

CART proliferation (**Fig. 1F-G, Supplementary Fig. S3A**), degranulation (**Fig. 1H**), and cytokine

production (**Fig. 1I-L**). Similar results were found for wildtype (TSHR-) and TSHR+ THJ-529, a

PDTC cell line (**Supplementary Fig. S3B-H**).

TSHR-CART cells show durable antitumor activity in xenograft HCC models

We then assessed TSHR-CART cell activity in a xenograft model of HCC. We first measured

TSHR expression via IHC on biopsies obtained from patients with normal thyroid tissue, primary

HCC with minimal or extensive invasion, locally recurrent HCC, and metastatic HCC. Strong

TSHR expression was maintained in recurrent and metastatic lesions as well as primary tumors

(**Fig. 2A-B**). Having verified that HCC retains TSHR expression in patient samples and thus

providing a strong rationale for the use of TSHR-CART cell monotherapy, we then assessed the

in vivo activity of TSHR-CART cells against the TSHR+ HCC cell line, XCT.UC1 as a proof of

concept. Non-obese diabetic severe combined immunodeficiency IL2rγ-/- (NSG) mice were

subcutaneously engrafted with 1 x 10^7 XTC.UC1, which retains high endogenous TSHR

expression *in vivo*, and treated with 1 x10^7 UTD or TSHR-CART cells when tumors reached

~100mm^3. Mice in remission were rechallenged with 1 x 10^7 XTC.UC1 cells subcutaneously in

the opposite flank (**Fig. 2C**). Serial caliper measurements revealed significant antitumor efficacy

of TSHR-CART cells compared to UTD control; by Day 50 post-engraftment, all UTD-treated mice

had reached humane endpoints, while all TSHR-CART-treated mice were in remission or had

small tumors (**Fig. 2D-E**). TSHR-CART-treated mice in remission were then rechallenged on Day

51 with XTC.UC1 in the opposite flank and were followed for tumor growth along with a new group

of untreated XTC.UC1-bearing mice as controls. Again, TSHR-CART cells displayed potent

antitumor activity; both original and rechallenged tumors were significantly suppressed compared

to untreated controls (**Fig. 2F**). Peripheral blood sampling at 21 days post-tumor rechallenge

revealed significantly higher numbers of human CD3$^+$ T cells in TSHR-CART-treated mice

compared to controls (**Fig. 2G**). These data demonstrated that TSHR-CART cells exerted durable

tumor control in HCC, an aggressive disease setting with no current standard of care.

TSHR-CART cells demonstrate strong antitumor effects in TSHR$^+$ PDX models of PDTC

Next, we evaluated the antitumor activity of TSHR-CART cells in a dose escalation model in NSG

mice. Here, we utilized aggressive, BRAF-mutant, patient-derived PDTC cells, THJ-529T, which

were transduced to overexpress TSHR. Mice were subcutaneously injected with 2.5 x 10^6

THJ-529T cells, and engraftment was confirmed by caliper measurement of tumor masses. When

tumors reached ~100mm^3, mice were randomized to treatment with 5 x 10^6 UTD, 5 x 10^6 TSHR-

CART cells, or 1 x 10^7 TSHR-CART cells (**Fig. 3A**). Treatment with TSHR-CART cells led to

significant and dose-dependent tumor control (**Fig. 3B-E, Supplemental Fig. S4A-C**), improved

survival outcomes (**Fig. 3F, Supplemental Fig. S4D**), increased tumor-infiltrated human T cells

(**Fig. 3G-H**), and increased human T cell proliferation in the peripheral blood (**Supplemental Fig.**

S4E). Having shown that the 1 x 10^7 TSHR-CART dose yielded superior antitumor activity

compared to the 5 x 10^6 TSHR-CART dose, we did not further pursue testing of the 5 x 10^6 TSHR-

CART dose.

TSHR is downregulated in ATC but is restored with MAPK inhibition

ATC cells have been associated with de-differentiation and downregulation of TSHR expression,

and we showed that ATC significantly downregulated TSHR expression compared to normal

thyroid tissues or more differentiated thyroid cancers (**Fig. 1A, B**). This observed downregulation

raises concerns for antigen escape and subsequent resistance to TSHR-CART cell therapy. We

first verified that the MEKi, trametinib, restored TSHR expression in an ATC PDX proof-of-concept

model. In this model, 5mm^3 Th-560 ATC PDX tissue was surgically implanted subcutaneously in

NSG mice. When tumors reached approximately 100mm^3, mice were randomized to daily

treatment with either vehicle control or MAPK inhibitors at the indicated doses until the

experimental endpoint (**Fig. 4A**). The administration of MEKi (trametinib) resulted in significant

dose-dependent slowing of tumor growth (**Fig. 4B**) as well as TSHR re-expression which inversely

correlated with pERK expression (**Fig. 4C**) after one week of daily trametinib treatment compared

to placebo treatment. Because many ATC tumors are BRAF-mutant, we performed a similar

experiment in which mice received either vehicle control or the dual MEKi/BRAFi R05126766 at

the indicated doses. Treatment with R05126766 resulted in a significant dose-dependent slowing

of tumor growth (**Fig. 4D**), and pERK expression inversely correlated with TSHR expression (**Fig.**

4E) after one week of daily R05126766 treatment compared to placebo treatment. Similarly,

treatment with combination of MEKi (trametinib) and BRAFi (dabrafenib) resulted in significantly

increased TSHR expression in ATC PDX models (**Fig. 4F-G**). While modest antitumor effects of

MEKi and/or BRAFi were observed, the main purpose of these experiments was to assess TSHR

upregulation upon MEKi and/or BRAFi treatment in PDX models of ATC. Collectively, these data

support the therapeutic strategy of MEKi and BRAFi treatment to re-express TSHR and thereby

reduce ATC resistance to TSHR-CART cell therapy.

MAPK inhibition upregulates TSHR and sensitizes TSHR-CART cell therapy to

dedifferentiated thyroid cancer *in vitro*

We then tested if pretreatment with MEKi or MEKi/BRAFi enhanced TSHR-CART cell activity

against dedifferentiated thyroid cancer cell lines *in vitro*. The PDTC cell line, THJ-529, was

cocultured with trametinib alone or trametinib plus dabrafenib (10nM or 1µM for trametinib or

dabrafenib). After four days, THJ-529 cells were washed and assessed for TSHR expression by

flow cytometry. THJ-529 cells demonstrated modest but consistently increased TSHR expression

after treatment with MEKi or MEKi/BRAFi compared to vehicle-treated cells (**Fig. 5A**). THJ-529

cells were then washed to remove residual MAPK inhibitor and plated with TSHR-CART cells for

five additional days, at which point TSHR-CART cell proliferation was assessed via flow

cytometry. TSHR-CART cells demonstrated significantly more proliferation in response to THJ-

529 cells which were pretreated with trametinib or trametinib plus dabrafenib compared to vehicle

control-treated THJ-529 cells (**Fig. 5B**). Treatment with higher or more prolonged MAPK inhibitor

doses to further increase TSHR expression resulted in tumor cell death to the point where post-

coculture analyses could not be conducted; however, the modest TSHR upregulation observed

with low doses of MAPK inhibitors resulted in significantly increased TSHR-CART response. We

then assessed the sensitivity of TSHR-CART cells to low expression levels of TSHR. We

generated a series of TSHR$^+$ FTC-133 cells with varied levels of TSHR expression. TSHR$^+$ FTC-

133 cells had TSHR levels ranging from ~40,000 to 300,000 TSHR per cell, compared to ~6,000

TSHR per cell for our ATC PDX model, THJ560, and ~1,000 TSHR per cell for wildtype FTC-133

(**Fig. 5C-D**). TSHR-CART cells demonstrated potent cytotoxicity against all TSHR$^+$ FTC-133 cell

lines, with only the wildtype FTC-133 cells showing resistance to TSHR-CART cell killing (**Fig.**

5E). These findings indicated that low to moderate TSHR upregulation via MAPK inhibition leads

to robust antitumor activity of TSHR-CART cells.

MAPK inhibition does not negatively impact TSHR-CART cell activity

Therefore, we aimed to evaluate MEK and BRAF inhibition as a strategy to upregulate TSHR

expression on thyroid cancer cells and increase the therapeutic index of TSHR-CART cell

therapy. Given that T cells are shown to utilize MAPK signaling, we first tested for potential

detrimental effects of MEKi or MEKi/BRAFi on CART cell activity. TSHR-CART proliferation (**Fig.**

6A), killing (**Fig. 6B**), degranulation (**Supplemental Fig. S5A**), and intracellular cytokine

production (**Supplemental Fig. S5B**) were not affected by the addition of trametinib alone or

trametinib plus dabrafenib compared to media control. We next assessed the effects of prolonged

MAPK inhibition on TSHR-CART function through an exhaustion assay developed with repeated

CART activation.(*40*) Here, TSHR-CART cells were repeatedly stimulated with irradiated TSHR$^+$

FTC-133 cells along with vehicle control, 10nM trametinib, or 10nM trametinib plus 10nM

dabrafenib. After one week of chronic stimulation, TSHR-CART cells were isolated and cocultured

with fresh TSHR$^+$ FTC-133 cells and vehicle control, 10nM trametinib, or 10nM trametinib plus

dabrafenib for proliferation and cytotoxicity assays (**Supplemental Fig. S5C**). Here as well, the

addition of MAPK inhibitors did not appear to negatively impact TSHR-CART cell proliferation

(**Fig. 6C, Supplemental Fig. S5D**) or killing (**Fig. 6D**).

Combination of TSHR-CART cells plus MAPK inhibitors yields superior TSHR-CART cell

activity

Having shown that the addition of MAPK inhibitors was not detrimental to CART function, we next

studied the kinetics of TSHR expression after MEKi or MEKi/BRAFi to determine the longevity of

TSHR re-expression after treatment with MAPK inhibitors. Mice were implanted with 5mm^3 Th-

560 ATC PDX tumors. When tumors reached ~100mm^3, mice were treated daily with either

vehicle control or with 1.5 mg/kg dual MEKi/BRAFi (R05126766) plus 1 mg/kg BRAFi

(dabrafenib). TSHR expression was assessed by IHC and was found to be significantly elevated

after 7 days of MAPK inhibitor treatment compared to vehicle control. However, TSHR expression

was subsequently lost within two days after stopping MAPK inhibitor treatment (**Supplemental**

Fig. S6A-B).

Given these results, we compared different dosing regimens of MEKi/BRAFi in combination with

TSHR-CART cell therapy in ATC PDX models to determine if sequential or concurrent MAPK

inhibitor administration yielded superior CART antitumor activity. Mice were subcutaneously

implanted with 5mm^3 Th-560 ATC PDX tumors. When tumors reached ~100mm^3, mice

were randomized to daily treatment with vehicle control, 0.25 mg/kg trametinib (concurrently

administered until experimental endpoints or stopped after 7 days), or 0.25 mg/kg trametinib plus

2 mg/kg dabrafenib (concurrently administered until experimental endpoints or stopped after 7

days). Mice treated with vehicle control were engrafted with tumor one week later than mice

treated with MAPK inhibitors to ensure equal tumor sizes across all groups when UTD or TSHR-

CART cells were administered. As a further dose escalation, mice were then given 2×10^7 UTD

or TSHR-CART cells via tail vein injection. Tumor volume, body weight, and body condition were

inspected to monitor disease progression and to assess for endpoint criteria (**Fig. 7A**). Mice

treated with concurrent MEKi or concurrent MEKi plus BRAFi in combination with TSHR-CART

cells (CART + MEKi and CART + MEKi/BRAFi, respectively) showed superior antitumor efficacy

(**Fig. 7B-C, Supplementary Fig. S7A-F, Supplementary Fig. S8A-F**). We then performed a

followup experiment using 1×10^7 UTD or TSHR-CART cells. Here, mice with concurrent MAPK

inhibitor administration plus TSHR-CART cells demonstrated durable antitumor control lasting

beyond two months (**Supplementary Fig. S8G**). Given these results, we used the 1×10^7 TSHR-

CART dose in subsequent experiments. These findings indicate that administering MAPK

inhibitors concurrently with TSHR-CART cells enhances the antitumor activity of TSHR-CART

cell therapy.

We next assessed how long MAPK inhibitors needed to be administered concurrently with TSHR-

CART cell therapy for enhanced antitumor efficacy. Here, mice treated with UTD or TSHR-CART

cells were given vehicle control, MEKi/BRAFi for two weeks after CART administration,

MEKi/BRAFi, for four weeks after CART administration, or continuous MEKi/BRAFi after CART

administration until experimental endpoints were reached (**Fig. 7D**). Mice receiving the combination

of TSHR-CART and MAPK inhibitors showed superior tumor control and survival, regardless of the

duration of concurrent MAPK inhibitors (**Fig. 7E-F, Supplementary Fig. S9A-F**). This indicates that

using MAPK inhibitors in combination with TSHR CART cells for 2 weeks is sufficient to induce

TSHR upregulation and potent TSHR activity. Peripheral blood analysis at day 14 post-CART

injection showed comparable human T cell expansion in mice treated with TSHR-CART cells plus

vehicle control compared to mice treated with TSHR-CART cells plus MAPK inhibitors (**Fig. 7G**).

DISCUSSION

Here, we describe the development and evaluation of a highly translational CART cell therapy for

the treatment of poorly differentiated, anaplastic, and metastatic thyroid cancers. TSHR is a

rational target for CART cell therapy, given that it is widely expressed in thyroid cancers and rarely

expressed outside the thyroid. We subsequently generated and tested TSHR-CART cells, which

demonstrated antigen-specific activity against TSHR-expressing target cells *in vitro* and *in vivo*.

HCC represents ~4% of thyroid cancers and is typically radioiodine-resistant for metastatic

disease, retains TSHR expression, and currently has no FDA-approved therapy.(*7*) Our data

demonstrate that TSHR-CART cell therapy resulted in durable remissions *in vivo* and may

represent a treatment option for this subset of patients.

De-differentiated and anaplastic thyroid cancers can exhibit significant downregulation of thyroid

developmental genes, including TSHR.(*41, 42*) This phenomenon poses a formidable hurdle to

effective TSHR-CART treatment. Increasing evidence indicates that TSHR downregulation and

de-differentiation is mediated by MAPK signaling.(*28, 29, 34, 43, 44*) The majority of thyroid

cancers have activating oncogenic gene mutations involving PI3K, RET, RAS, and BRAF.(*32*)

Constitutive activation of these proteins stimulates MAPK signaling, inhibiting the expression of

thyroid hormone biosynthesis genes.(*33, 34*) Therefore, when BRAF activation is switched off

genetically or its downstream signaling is inhibited with MEKi or BRAFi, tumors re-express TSHR

and reaccumulate iodine. Such a strategy has been employed in the clinic to enhance the

response to radioactive iodine therapy.(*30*) Recent findings from patients with thyroid cancer who

have been treated with MEKi indicate a dramatic upregulation of TSHR within both primary and

secondary thyroid cancer tumors.(*30, 45*) Additionally, MEKi, alone or in combination with

BRAFi, demonstrate significant antitumor activity in metastatic thyroid cancer and are FDA-

approved for ATC.(*46*) We therefore tested the combination of TSHR-CART cell therapy plus

MAPK inhibition to upregulate TSHR expression in the tumor and enhance CART efficacy,

demonstrating restoration of TSHR expression and improved CART antitumor effects in mice

receiving combination therapy compared to mice receiving CART monotherapy.

Major limitations to effective CART cell therapy in solid tumor settings include appropriate target

antigen selection, antigen-negative escape, and lack of T cell infiltration into the tumor site. The

treatment strategy we have outlined here has strong potential to overcome these limitations, as

TSHR is nearly uniquely and universally expressed in thyroid tumors, MAPK inhibition induces

TSHR re-expression, and TSHR-expressing advanced thyroid tumors are commonly infiltrated by

immune cells, including T cells.(*9*)

Regarding the potential for on-target off-tumor toxicity, consistent with publicly available datasets,

our tissue microarray demonstrates that TSHR expression is mainly limited to the thyroid gland.

Furthermore, our CAR is cross reactive with mouse TSHR and we did not observe any toxicities

in our mouse experiments. As patients with aggressive thyroid cancers will have undergone

previous thyroidectomy, there will be no healthy thyroid tissue remaining upon treatment with

TSHR-CART cells. There have also been reports of extra-thyroidal TSHR expression and

function. Aberrant TSHR expression has been documented in the orbital fibroblasts of patients

with Graves' disease in the context of thyroid associated ophthalmopathy, and TSHR-targeted

treatments have shown clinical responses.(*47-51*) However, thyroid cancers have not been

associated with thyroid associated ophthalmopathy. As such, it is not probable that orbital/ocular

toxicities will arise with TSHR CART cell therapy. We also have not oberved any orbital toxicity in

our mouse models. In addition, K1-70, the TSHR antibody from which we derived the scFv for our

TSHR-CAR, was found to be well-tolerated at all dose levels in a phase I trial for patients with

Graves' disease, (*52*) and in a patient with both follicular thyroid cancer and Graves' disease.(*53*)

These clinical studies, while still nascent, demonstrate a promising safety profile. Vigilant

monitoring for orbital toxicities will be incorporated into our phase I clinical trial.

CART cells targeted against ICAM-1(*54*) and GFRα4(*55*) have demonstrated efficacy in preclinical

thyroid cancer models. Recently, TSHR-CART cell therapy displayed potent activity against

human thyroid cancer models *in vivo*.(*38*) Furthermore, two patients with high metastatic burden

and iodine-131-resistant thyroid cancer demonstrated a complete response (CR) and a near CR

three months after TSHR-CART treatment. The first patient, who had papillary thyroid carcinoma,

sustained CR until one year post-TSHR-CART infusion, indicating potential for long-term

antitumor responses.(*56*) However, these studies focused on differentiated thyroid cancers with

high TSHR expression. One patient with iodine-refractory, metastatic PDTC was treated with

CART cells targeting both TSHR and CD19. This patient achieved stable disease at day 30 post-

CART infusion and partial response 90 days post-CART infusion. While IHC demonstrated strong

TSHR expression pre-CART infusion, it is unknown if TSHR was downregulated after the start of

TSHR-CART cell therapy, leading to underwhelming therapeutic response in this dedifferentiated

thyroid cancer setting.(*57*)

Several reports have shown conflicting results regarding the potential impacts of MAPK inhibitors

on T cell function. One study found that dabrafenib did not suppress human T cell function, but

trametinib led to an incomplete and transient inhibition of T cell proliferation and cytokine

production. The addition of dabrafenib to trametinib was found to offset the negative effects of

trametinib on T cell function *in vitro.* Further, trametinib did not appear to have detrimental effects

on T cells, but rather increased tumor infiltration of T cells *in vivo*.(*58*) Another study found that

BRAFi also enhanced tumor infiltration of adoptively transferred T cells in mice.(*59*) Other reports

showed that MEKi led to reprogramming of CD8 T cells to stem cell memory phenotypes and

thereby enhanced antitumor activity.(*60*) Another group reported that MEKi during chronic T cell

stimulation ameliorated terminal T cell exhaustion by slowing the transcription rates of effector-

and terminal- exhaustion-related genes, reducing nutrient uptake and mitochondrial ROS

accumulation, and ultimately enhancing the persistence of tumor-infiltrating lymphocytes *in*

vivo.(*61*) An early report of the impacts on MAPK inhibitors on CART cells showed that trametinib

plus dabrafenib across a range of concentrations inhibited GD2-targeted patient-derived CART

cell activation, cytokine production, and cytotoxicity; however, the authors suggested that strong

stimulation of the CAR with GD2^{high} tumor cells had the potential to overcome the detrimental

effects of MAPK inhibitors on CART functions.(*62*) Another GD2-CART study confirmed that

trametinib inhibited CART cytotoxic abilities and cytokine secretion *in vitro*, but *in vivo*, trametinib

in combination with GD2-CART resulted in enhanced tumor suppression compared to

monotherapy treatment groups.(*63*) Another group showed that, while the combination of

trametinib and dabrafenib did negatively impact CSPG4-targeted CART cell *in vitro* functions,

these effects were milder than those seen with other MAPK inhibitor combinations.(*64*) Recently,

it was reported that MEKi reduced CART cell activation, exhaustion, apoptosis, and terminal

differentiation across multiple different CAR constructs in both solid and liquid tumors through

downregulation of c-Fos and JunB.(*65*) While consensus has yet to be reached, more recent

studies demonstrate that potential negative impacts of MAPK inhibitors on T cell function are

nuanced and context-dependent, and MAPK inhibitors may actually enhance CART function in a

variety of settings.

We present TSHR-CART cell therapy in combination with MAPK inhibition as a sensitizing

modality, representing a major step forward in the development of a treatment strategy for PDTC

and ATC as well as DTCs which have become refractory to radioiodine therapy. Based on these

data, we are planning to launch a phase I clinical trial for TSHR-CART cell therapy as a

monotherapy in patients with TSHR^{high} metastatic thyroid cancers and in combination with MAPK

inhibitors in patients with TSHR^{low} metastatic thyroid cancers.

MATERIALS AND METHODS

Study design

This study aimed to assess the efficacy of TSHR-CART cell therapy as a monotherapy in TSHR^{high}

thyroid cancers and in combination with MAPK inhibitors in TSHR^{low} thyroid cancers. First, the

expression of TSHR was assessed and verified on patient clinical samples utilizing IHC and tissue

microarray tools. Second, we designed and engineered human CART cells targeting TSHR

(TSHR-CART), that were subjected to various *in vitro* and vivo *assays* to assess their effector

functions (cytotoxicity, proliferation, degranulation and cytokine production). Third, we tested the

addition of MAPK inhibitors *in vitro* to assess their upregulation of TSHR and sensitization prior

to TSHR-CART treatment. Lastly, we performed *in vivo* assays to assess CART cell antitumor

activity in combination with MAPK inhibitors in a concurrent or sequential fashion. For each

experiment, the number of independent experiments (*n*) is specified in the figure legends. For *in*

vivo assays, NGS mice were utilized, and they were randomly assigned to control and

experimental groups, and the study was not blinded. Data collection was stopped at the

predetermined times. Human endpoint criteria were determined when mice presented above 20%

lost weight and tumor exceeded 700 mm^3. No data was excluded from analysis. Sample sizes

are indicated in the figure legends.

Reagents

Detailed information on all reagents used in this article is listed in Supplementary Table S1.

Cell lines

FTC-133, XTC.UC1, and THJ-529T cell lines were graciously provided by Dr. Copland's lab in

Mayo Clinic, Jacksonville, Florida. The FTC-133 cell line is derived from a differentiated follicular

thyroid carcinoma with known mutations that include *FLCN, MSH6, NF1, PTEN, TERT*, and *TP53*

(https://www.cellosaurus.org/CVCL_1219). The THJ-529T cell line was derived from a PDTC

tumor containing *BRAF* and *TERT* mutations (https://www.cellosaurus.org/CVCL_9916).(*66*)

XTC.UC1 is the only characterized HCC cell line and the only known cell line to continue to

express endogenous TSHR *in vitro*.(*67*) All cell lines were DNA short tandem repeat (STR)

validated prior to use. For indicated experiments, THJ-529 and FTC-133 cell lines were

transduced with a firefly luciferase (Luc$^+$) vector containing a hygromycin resistance gene

(Promega, Madison, WI, USA) as well as a TSHR vector and a puromycin resistance gene

(Genecopoeia Inc., Rockville, MD, USA). These cells were then cultured in selection media

containing either hygromycin (1 μg/mL) or puromycin (1 μg/mL) to obtain a pure population as

previously described.(*68*) FTC-133 and XTC.UC1 cell lines were cultured in D10 medium made

with DMEM (Corning Inc., Corning, NY, USA), 10% FBS (Sigma, St. Louis, MO, USA), and 1%

PSG (Gibco, Gaithersburg, MD, USA). For specific *in vitro* assays, FTC-133 cell lines were

transduced with different MOIs of a TSHR vector containing puromycin resistance gene

(Genecopoeia Inc., Rockville, MD, USA), and TSHR absolute counts were determined by flow

cytometry using absolute counting beads (CountBright™, Invitrogen, Carlsbad, CA, USA).

Mycoplasma testing was performed and confirmed to be negative monthly.

CAR construct and CART cell production

The use of recombinant DNA in the laboratory was approved by the Mayo Clinic Institutional

Biosafety Committee (IBC), IBC number HIP00000252.20. Peripheral blood mononuclear cells

(PBMCs) were isolated from de-identified normal donor blood apheresis cones(*69*) obtained

under a Mayo Clinic IRB approved protocol, using SepMate tubes (STEMCELL Technologies,

Vancouver, BC, Canada) and Lymphoprep™ solution (STEMCELL Technologies, Vancouver,

BC, Canada). T cells were separated with negative selection magnetic beads using EasySep^{TM}

Human T Cell Isolation Kit (STEMCELL Technologies, Vancouver, BC, Canada). Second-

generation 4-1BB- costimulated TSHR CAR constructs were synthesized *de novo* (IDT, Coralville,

USA) and cloned into a third-generation lentivirus under the control of the EF-1α promotor. The

TSHR-targeted single chain variable fragment was derived from an autoantibody from a patient

with Graves' disease, clone K1-70.(*70*) TSHR-CART cells were then generated through the

lentiviral transduction of normal donor T cells as previously described(*71, 72*). Briefly, lentiviral

particles were generated through the transient transfection of plasmid into 293T virus-producing

cells (ATCC, Manassas, VA, USA) in the presence of Lipofectamine 3000 (Invitrogen, Carlsbad,

CA, USA), vesicular stomatitis virus G (VSV-G), and packaging plasmids pCMVR8.74 (Addgene,

Cambridge, MA, USA). The titers and multiplicity of infection (MOI) were analyzed and calculated

by flow cytometry as previously described.(*68*) T cells isolated from normal donors were stimulated

using anti-CD3/CD28 Dynabeads (Life Technologies, Oslo, Norway) at a 3:1 beads-to-cell ratio

and then transduced 24 hours after stimulation with lentivirus particles at a MOI of 3. T cells were

cultured in T cell medium containing X-VIVO™ 15 media (Lonza, Basel, Switzerland), 10% human

AB serum (Corning Inc., Corning, NY, USA), and 1% PSG (Gibco, Gaithersburg, MD, USA).

Magnetic bead removal and the evaluation of CAR expression on T cells by flow cytometry were

performed on day 6 by staining with a goat anti-human IgG H+L secondary antibody (Invitrogen,

Carlsbad, CA, USA). CART cells were harvested and cryopreserved on day 8 for future

experiments. Prior to cryopreservation, CART cells were stained and evaluated to assess T-cell

subset and phenotype expression patterns using the following antibodies: CD45 APC (BioLegend,

San Diego, CA, USA), CD3 BV605 (BioLegend), CD4 PE-Cy7 (BD Biosciences, Franklin Lakes, NJ,

USA), CD8 BV421 (BD Biosciences), CCR7 PE (BioLegend) and CD45RA APC-Cy7 (BioLegend).

CART cells were thawed and rested in T cell medium overnight at 37°C and 5% CO_2 prior to

experimental use.

Multi-parametric flow cytometry

Samples were prepared for flow cytometry as previously described.(*68, 71-74*) In all analyses,

the population of interest was gated based on forward vs side scatter characteristics, then by

singlet gating, and then by live cell gating using LIVE/DEAD™ Fixable Aqua Dead Cell Stain Kit

(Invitrogen, Carlsbad, CA, USA). Flow cytometry was performed on a three-laser CytoFLEX

(Beckman Coulter, Chaska, MN, USA). All analyses were performed using FlowJo v10.7.1

software (Ashland, OR, USA).

Immunohistochemistry (IHC)

481 IHC was performed on paraffin-embedded tissue. Thyroid tissue microarray (TMA) was made

482 from archival samples under Mayo Clinic IRB approval with sample size for each tumor at 1.5 mm.

483 Paraffin blocks were cut into 5 μm thick sections. Slides were deparaffinized, rehydrated with

484 decreasing concentrations of ethanol, and finally with water. Additionally, a normal human tissue

485 microarray was purchased (Bio SB, Santa Barbara, CA, USA), which included 23 different normal

human tissue types. Antigen retrieval was performed using Antigen Retrieval Solution (pH 6 or pH 9) (Dako, Glostrup, Denmark) for at least 15 minutes in an autoclave, and cooling for 30 minutes at room temperature. Slides were washed with 3% hydrogen peroxide (Fisher Scientific, Waltham, MA, USA) for 10 minutes to block the activity of endogenous peroxidases. Primary anti-human TSHR antibody was used for 60 minutes at room temperature at 1:500 (abcam, ab218108, Cambridge, UK). The infiltration of human T lymphocytes into tumors was assessed using anti-human CD3 at 1:500 (abcam, ab52959, Cambridge, UK). The primary antibody was detected using the Envision Labeled Kit (Dako, Glostrup, Denmark) according to the manufacturer's protocol. After washing, slides were counterstained with Mayer's hematoxylin (Sigma-Aldrich, St. Louis, MO, USA). Slides were dehydrated through ethanol and xylene and cover-slipped using a xylene-based mounting medium (Fisher Scientific). Samples were examined under bright-field illumination at 20x objective, and digital images were obtained using Aperio AT2 (Leica, Wetzlar, Germany). Results were processed using Aperio eSlide Manager and H-Score values were estimated using the Aperio ImageScope Software (both Aperio Technologies, Vista, CA, USA). The staining of the TMA punches was scored using an algorithm in the Imagescope Software (Aperio Technologies) created by a histologist based upon signal intensity (0, 1C, 2C, 3C). H score was then calculated based upon signal intensity and percentage: HZ(1C%!1)(2C%!2)(3C%!3). Cases were excluded from the study if a section could not be assigned a score due to insufficient quantity of tumor tissue present. Normal tissues were used as a positive and negative control.

For cell line preparation for IHC, cells were washed three times with PBS (Corning, NY, USA), scraped from the bottom of the plate using a cell lifter (Fisherbrand, Waltham, MA, USA), transferred to 50 mL polypropylene tubes and centrifuged at room temperature for 2 minutes at 500 x g. The supernatant was aspirated, and 10% neutral buffered formalin (Fisher Scientific) was added for 30 minutes at room temperature. HistoGel (Thermo Scientific) was heated in the microwave for 3 seconds at maximal power and converted to a liquid state. Cells were transferred

to the HistoScreen Tissue Cassettes (Thermo Scientific) and covered with liquid Histogel.

Samples were placed at room temperature and allowed to solidify. HistoGel blocks were

transferred into tissue embedding cassettes (Thomas Scientific, Swedesboro, NJ, USA),

dehydrated in increasing concentrations of ethanol, xylene, and paraffin embedding. Sections

were prepared as described in the immunohistochemistry section. Primary anti-human TSHR

antibody was used for 60 minutes at room temperature at 1:1500 (abcam, ab218108, Cambridge,

UK). Envision labeled polymer (Dako, Glostrup, Denmark) was used for 30 minutes as a

secondary antibody. Slides were stained with diaminobenzidine tetrahydrochloride (DAB)

chromogen (Dako) for 5 minutes at room temperature and counterstained with Mayer's

hematoxylin (Sigma-Aldrich, St. Louis, MO, USA). Control staining with hematoxylin and eosin

(Sigma-Aldrich) was also performed. Aperio AT2 scanner (Leica, Wetzlar, Germany) was used

for the digital image, at 20x objective. Images were analyzed using the Aperio ImageScope

software (Aperio Technologies).

Immunocytochemistry (ICC)

THJ-529 cells (2×10^4 cells/chamber) were plated in a 4-well chamber slide (Nunc, Thermo

Scientific, Waltham, MA, USA), incubated overnight. After 24 hours, media was aspirated, and

cells were washed three times with PBS (Corning, NY, USA). Then, 2% paraformaldehyde was

added for 20 minutes at room temperature. Again, cells were washed three times with PBS. Next,

ice-cold 100% methanol (Fisher Scientific) was added for 7 minutes at -20°C. Methanol was

aspirated, and samples were allowed to air dry for 10 minutes. Next, serum-free-blocking diluent

(Dako) was used for 30 minutes at room temperature. Primary anti-human TSHR antibody was

used for 60 minutes at room temperature at 1:1500 (abcam, ab218108, Cambridge, UK). Slides

were subsequently prepared as described in the immunohistochemistry section.

***In vitro* T cell function assays**

Cytotoxicity assays were performed as previously described.(*68, 71-74*) Wildtype (TSHR⁻) Luc⁺

FTC-133, TSHR$^+$ Luc$^+$ FTC-133, wildtype (TSHR$^-$) Luc$^+$ THJ-529, or TSHR$^+$ Luc$^+$ THJ-529 cells

were used as target cells. UTD or TSHR-CART cells were co-cultured with target cells at various

effector: target (E:T) ratios in T cell medium. Cytotoxicity was calculated by bioluminescence

imaging on the Promega GloMax Explorer (Promega Corporation, Fitchburg, WI, USA) as a

measure of residual live cells at indicated time points. Samples were treated with 1µL D-luciferin

(30µg/mL) per 100µL sample volume (Gold Biotechnology, St. Louis, MO, USA) prior to imaging.

Proliferation assays were performed utilizing two approaches as previously described.(*68, 71-74*)

Briefly, for the first approach, TSHR-CART cells were stained with CFSE (Invitrogen) cell dye on

day 0, then washed and counted prior to being plated at a 1:1 E:T ratio with target cell lines. Each

assay included T cells with media only as a negative control and T cells with PMA and ionomycin

as a positive control. On day 5, percent of CFSE CART cells was determined via flow cytometry

after staining with APC-H7 anti-human CD3 (BD Biosciences) and Zombie R718™ (BioLegend)

dye. Alternatively, the same assay was repeated but without the addition of CFSE and with

LIVE/DEAD™ Fixable Aqua as the viability dye.

For *in vitro* experiments testing the cytotoxic and proliferation effects of TSHR-CART cells

exposed to different levels of TSHR on target cells, TSHR-CART cells were co-cultured with FTC-

133 tumor cells expressing different levels of TSHR and cytotoxicity and proliferation was

measured as described above.

For cytokine assays, media was collected on day 3 of co-culture of TSHR-CART cells with target

cells, and supernatants were isolated and purified via centrifugation as previously described(*71*).

Supernatants were prepared following manufacturer's protocol for the MILLIPLEX® Human

Cytokine/Chemokine/Growth Factor Panel A 38 Plex Premixed Magnetic Bead Panel (Millipore

Sigma, Ontario, Canada). Cytokine data was collected using a Luminex LX200 (Millipore Sigma).

For degranulation assays (*68, 71*), UTD or TSHR-CART cells were incubated with various target

cells at an E:T ratio of 1:5, and antibodies against CD107a FITC (BD Biosciences), CD28 (BD

Biosciences), and CD49d (BD Biosciences), as well as monensin (BioLegend), were added at the

start of the incubation. After 4 hours, cells were harvested and stained with LIVE/DEAD™ Fixable

Aqua. Cells were then fixed and permeabilized (FIX & PERM Cell Fixation & Cell Permeabilization

Kit, Life Technologies, Oslo, Norway) and stained for CD3 APC (, eBioscience, San Diego, CA,

USA) and intracellular cytokines including IL-2 PE CF594 (BD Horizon), GM-SCF BV421 (BD

Horizon) and MIP-1β Pe-Cy7 (BD Pharmigen).

For *in vitro* assays performed to assess the impact of MAPK inhibitors on TSHR-CART cells,

TSHR-CART cells were cultured with target tumor cells or media alone, along with either vehicle

control (DMSO), trametinib (10nM) alone, or trametinib plus dabrafenib (10nM). Effector functions

of TSHR-CART cells (proliferation, cytotoxicity, degranulation and cytokine profile) were

assessed as described above. Alternatively, TSHR-CART cells were cocultured in an *in vitro*

model for CART cell exhaustion previously described by our laboratory.(*40*) Here, TSHR-CART

cells were cultured at a 1:1 ratio with irradiated target cells along with either vehicle control

(DMSO), trametinib (10nM) alone, or trametinib plus dabrafenib (10nM). Drug-containing media

and irradiated target cells were added to the TSHR-CART cell cocultures every other day for 7

days. Then, chronically stimulated TSHR-CART cells were isolated using CD4 (Miltenyi Biotec,

Auburn, CA, USA) and CD8 (Miltenyi Biotec) microbeads and were subsequently plated with

fresh, non-irradiated target cells to assess cytotoxic and proliferative capacities as described

above.

***In Vivo* Models**

Male and female 6-8-week-old NSG mice were purchased (Jackson Laboratories, Bar Harbor,

ME, USA) and maintained in an animal barrier space that is approved by the institutional Biosafety

Committee for BSL2+ level experiments at the Mayo Clinic animal facility (IBC

#HIP00000252.20). Mice were treated on an IACUC-approved protocol (A00001767).

XCT.UC1 model: The purpose of this model was to test initial TSHR-CART activity in a thyroid

cancer model which retains endogenous TSHR expression *in vivo*. XCT.UC1 is a Hurtle cell

carcinoma cell line which retains TSHR expression when passaged in mice but not *in vitro*. 1 x

10^7 XTC.UC1 cells were suspended in a 50% Matrigel basement medium and subcutaneously

injected into the rear flank of NSG mice. Serial caliper measurements were performed to confirm

tumor engraftment. Mice were then randomized based on tumor burden to receive 1 x 10^7 UTD

or TSHR-CART. Tumor growth was monitored by serial caliper measurements (width x height x

length x 0.523 = mm^3). Mice were euthanized once IACUC-approved endpoint criteria were met,

and remaining tumors were collected for IHC as indicated in each specific experiment.

TSHR$^+$ THJ-529 model: The purpose of this model was to assess a TSHR-CART dose

escalation. TSHR$^+$ THJ-529 is a poorly differentiated thyroid cancer cell line made to overexpress

TSHR. 2.5 x 10^6 TSHR$^+$ THJ-529 cells were suspended in a 50% Matrigel basement medium and

subcutaneously injected into the rear flank of NSG mice. Serial caliper measurements were

performed to confirm tumor engraftment. Mice were then randomized based on tumor burden to

receive 5 x 10^6 UTD, 5 x 10^6 TSHR-CART, or 1 x 10^7 TSHR-CART. Tumor growth was monitored

by serial caliper measurements (width x height x length x 0.523 = mm^3). Mice were euthanized

once IACUC-approved endpoint criteria were met, and remaining tumors were collected for IHC

as indicated in each specific experiment.

TH-560 model: The purpose of this model was to assess TSHR upregulation or TSHR-CART

activity after MAPKi administration. This model reflects the strategy of TSHR-CART and MAPK

inhibitor combination to be used in patients with ATC. TH-560 is an ATC PDX with downregulated

TSHR expression. All NSG mice were surgically engrafted with 5mm^3 ATC PDX TH-560 tumors

in the rear flank. In TSHR upregulation experiments, mice were treated with the indicated doses of

MEKi/BRAFi (trametinib, dabrafenib, or R05126766). Upon euthanasia, tumors were harvested for

IHC.

In the MAPK inhibitor and TSHR-CART combination experiments, upon tumor engraftment, mice were

treated with a sub-therapeutic dose of 1mg/kg trametinib and 1.5mg/kg dabrafenib as a sensitizing

regimen to induce TSHR expression on ATC tumor cells. On Day 7, additional mice were surgically

engrafted with 5mm^3 ATC PDX tumors in the same manner as the first arm but were treated with

placebo rather than MAPK inhibitors. In the first MAPK inhibitor and TSHR-CART combination

experiment, in which mice were treated sequentially or concurrently with MEKi/BRAFi and TSHR-

CART cells, all mice were randomized based on tumor burden when tumors reached ~100mm^3

and injected with 2 x 10^7 UTD or TSHR-CART; this higher TSHR-CART dose was a further dose

escalation experiment. In the second MAPK inhibitor and TSHR-CART combination experiment, in

which mice were treated concurrently with MEKi/BRAFi and TSHR-CART cells with various

MEKi/BRAFi treatment stopping points, all mice were randomized based on tumor burden when

tumors reached ~100mm^3 and injected with 1 x 10^7 UTD or TSHR-CART. This lower dose of

TSHR-CART was used since no further treatment benefit was observed with the higher 2 x 10^7

dose. For all mice in the MAPK inhibitor and TSHR-CART combination experiments, tumor growth

was monitored by serial caliper measurements (width x height x length x 0.523 = mm^3). Weekly

tail vein bleeding after injection of CART cells was performed to assess T cell expansion. Mouse

peripheral blood was lysed using BD FACS Lyse (BD Biosciences) and then analyzed with flow

cytometry as previously described.(*68*) Mice were euthanized at IACUC-approved endpoints.

Statistical analysis

GraphPad Prism (La Jolla, CA, USA) and Microsoft Excel (Microsoft, Redmond, WA, USA) were

used to analyze raw experimental data. Statistical tests are described in figure legends.

Author Contributions: CMR, ELS, and JJG wrote, reviewed, and edited the manuscript,

designed and performed experiments, and analyzed data. JJG, JLM, EEM, AAD, TNH, GED,

LKM, BLK, EET, RLS, ELS, MLP, AAD, CMS, IC, OLS, KY, JMF, OLGR, HX, MHT, KJS, and

EJO aided in data collection. GO aided in patent and intellectual property management. YQ, HWT,

RCS, and AZ aided in data collection and troubleshooting. JAC and SSK formulated the initial

concept, designed experiments, supervised the study, and wrote the manuscript. All authors

edited and approved the final version of the manuscript.

REFERENCES

1. C. M. Kitahara, J. A. Sosa, Understanding the ever-changing incidence of thyroid cancer. *Nat Rev Endocrinol* **16**, 617-618 (2020).

2. J. Kim, J. E. Gosnell, S. A. Roman, Geographic influences in the global rise of thyroid cancer. *Nat Rev Endocrinol* **16**, 17-29 (2020).

3. H. Lim, S. S. Devesa, J. A. Sosa, D. Check, C. M. Kitahara, Trends in Thyroid Cancer Incidence and Mortality in the United States, 1974-2013. *JAMA* **317**, 1338-1348 (2017).

4. R. C. Smallridge, J. A. Copland, Anaplastic thyroid carcinoma: pathogenesis and emerging therapies. *Clin Oncol (R Coll Radiol)* **22**, 486-497 (2010).

5. C. K. Jung, A. Bychkov, K. Kakudo, Update from the 2022 World Health Organization Classification of Thyroid Tumors: A Standardized Diagnostic Approach. *Endocrinol Metab (Seoul)* **37**, 703-718 (2022).

6. S. Ahmadi, M. Stang, X. S. Jiang, J. A. Sosa, Hurthle cell carcinoma: current perspectives. *Onco Targets Ther* **9**, 6873-6884 (2016).

7. D. G. McFadden, P. M. Sadow, Genetics, Diagnosis, and Management of Hurthle Cell Thyroid Neoplasms. *Front Endocrinol (Lausanne)* **12**, 696386 (2021).

8. D. Viola, L. Valerio, E. Molinaro, L. Agate, V. Bottici, A. Biagini, L. Lorusso, V. Cappagli, L. Pieruzzi, C. Giani, E. Sabini, P. Passannati, L. Puleo, A. Matrone, B. Pontillo-Contillo, V. Battaglia, S. Mazzeo, P. Vitti, R. Elisei, Treatment of advanced thyroid cancer with targeted therapies: ten years of experience. *Endocr Relat Cancer* **23**, R185-205 (2016).

9. J. D. French, Immunotherapy for advanced thyroid cancers - rationale, current advances and future strategies. *Nat Rev Endocrinol* **16**, 629-641 (2020).

10. A. Maniakas, R. Dadu, N. L. Busaidy, J. R. Wang, R. Ferrarotto, C. Lu, M. D. Williams, G. B. Gunn, M. C. Hofmann, G. Cote, J. Sperling, N. D. Gross, E. M. Sturgis, R. P. Goepfert, S. Y. Lai, M. E. Cabanillas, M. Zafereo, Evaluation of Overall Survival in Patients With Anaplastic Thyroid Carcinoma, 2000-2019. *JAMA Oncol* **6**, 1397-1404 (2020).

665   11.   C. H. June, M. Sadelain, Chimeric Antigen Receptor Therapy. **379**, 64-73 (2018).

666   12.   S. J. Schuster, M. R. Bishop, C. S. Tam, E. K. Waller, P. Borchmann, J. P. McGuirk, U.

667   Jäger, S. Jaglowski, C. Andreadis, J. R. Westin, I. Fleury, V. Bachanova, S. R. Foley, P.

668   J. Ho, S. Mielke, J. M. Magenau, H. Holte, S. Pantano, L. B. Pacaud, R. Awasthi, J. Chu,

669   Ö. Anak, G. Salles, R. T. Maziarz, Tisagenlecleucel in Adult Relapsed or Refractory

670   Diffuse Large B-Cell Lymphoma. **380**, 45-56 (2018).

671   13.   S. S. Neelapu, F. L. Locke, N. L. Bartlett, L. J. Lekakis, D. B. Miklos, C. A. Jacobson, I.

672   Braunschweig, O. O. Oluwole, T. Siddiqi, Y. Lin, J. M. Timmerman, P. J. Stiff, J. W.

673   Friedberg, I. W. Flinn, A. Goy, B. T. Hill, M. R. Smith, A. Deol, U. Farooq, P. McSweeney,

674   J. Munoz, I. Avivi, J. E. Castro, J. R. Westin, J. C. Chavez, A. Ghobadi, K. V. Komanduri,

675   R. Levy, E. D. Jacobsen, T. E. Witzig, P. Reagan, A. Bot, J. Rossi, L. Navale, Y. Jiang, J.

676   Aycock, M. Elias, D. Chang, J. Wiezorek, W. Y. Go, Axicabtagene Ciloleucel CAR T-Cell

677   Therapy in Refractory Large B-Cell Lymphoma. *N Engl J Med* **377**, 2531-2544 (2017).

678   14.   N. C. Munshi, L. D. Anderson, Jr., N. Shah, D. Madduri, J. Berdeja, S. Lonial, N. Raje, Y.

679   Lin, D. Siegel, A. Oriol, P. Moreau, I. Yakoub-Agha, M. Delforge, M. Cavo, H. Einsele, H.

680   Goldschmidt, K. Weisel, A. Rambaldi, D. Reece, F. Petrocca, M. Massaro, J. N. Connarn,

681   S. Kaiser, P. Patel, L. Huang, T. B. Campbell, K. Hege, J. San-Miguel, Idecabtagene

682   Vicleucel in Relapsed and Refractory Multiple Myeloma. *N Engl J Med* **384**, 705-716

683   (2021).

684   15.   S. L. Maude, T. W. Laetsch, J. Buechner, S. Rives, M. Boyer, H. Bittencourt, P. Bader, M.

685   R. Verneris, H. E. Stefanski, G. D. Myers, M. Qayed, B. De Moerloose, H. Hiramatsu, K.

686   Schlis, K. L. Davis, P. L. Martin, E. R. Nemecek, G. A. Yanik, C. Peters, A. Baruchel, N.

687   Boissel, F. Mechinaud, A. Balduzzi, J. Krueger, C. H. June, B. L. Levine, P. Wood, T.

688   Taran, M. Leung, K. T. Mueller, Y. Zhang, K. Sen, D. Lebwohl, M. A. Pulsipher, S. A.

689   Grupp, Tisagenlecleucel in Children and Young Adults with B-Cell Lymphoblastic

690   Leukemia. *N Engl J Med* **378**, 439-448 (2018).

16.   G. L. Beatty, E. K. Moon, Chimeric antigen receptor T cells are vulnerable to
immunosuppressive mechanisms present within the tumor microenvironment.
Oncoimmunology **3**, e970027 (2014).

17.   M. Castellarin, K. Watanabe, C. H. June, C. C. Kloss, A. D. Posey, Driving cars to the
clinic for solid tumors. *Gene Therapy* **25**, 165-175 (2018).

18.   S. Ma, X. Li, X. Wang, L. Cheng, Z. Li, C. Zhang, Z. Ye, Q. Qian, Current Progress in
CAR-T Cell Therapy for Solid Tumors. *Int J Biol Sci* **15**, 2548-2560 (2019).

19.   B. Moghimi, D. Barrett, CAR T Cells for Solid Tumors. *Current Stem Cell Reports* **3**, 269-
278 (2017).

20.   R. C. Sterner, R. M. Sterner, CAR-T cell therapy: current limitations and potential
strategies. **11**, 69 (2021).

21.   L. A. Kankeu Fonkoua, O. Sirpilla, R. Sakemura, E. L. Siegler, S. S. Kenderian, CAR T
cell therapy and the tumor microenvironment: Current challenges and opportunities. *Mol*
Ther Oncolytics **25**, 69-77 (2022).

22.   I. Can, M. J. Cox, E. L. Siegler, R. Sakemura, S. S. Kenderian, Challenges of chimeric
antigen receptor T-cell therapy in chronic lymphocytic leukemia: lessons learned. *Exp*
Hematol **108**, 1-7 (2022).

23.   R. Sakemura, M. Hefazi, E. L. Siegler, M. J. Cox, D. P. Larson, M. J. Hansen, C.
Manriquez Roman, K. J. Schick, I. Can, E. E. Tapper, P. Horvei, M. M. Adada, E. D.
Bezerra, L. A. Kankeu Fonkoua, M. W. Ruff, W. Nevala, D. K. Walters, S. A. Parikh, Y.
Lin, D. F. Jelinek, N. E. Kay, P. L. Bergsagel, S. S. Kenderian, Targeting Cancer-
Associated Fibroblasts in the Bone Marrow Prevents Resistance to CART-Cell Therapy in
Multiple Myeloma. *Blood*, (2022).

24.   M. Tuncel, Thyroid Stimulating Hormone Receptor. *Mol Imaging Radionucl Ther* **26**, 87-
91 (2017).

25.   C. W. Rowe, J. W. Paul, C. Gedye, J. M. Tolosa, C. Bendinelli, S. McGrath, R. Smith,

Targeting the TSH receptor in thyroid cancer. *Endocr Relat Cancer* **24**, R191-R202 (2017).

26. P. Carayon, C. Thomas-Morvan, E. Castanas, M. Tubiana, Human thyroid cancer: membrane thyrotropin binding and adenylate cyclase activity. *J Clin Endocrinol Metab* **51**, 915-920 (1980).

27. T. F. Davies, R. Latif, Targeting the thyroid-stimulating hormone receptor with small molecule ligands and antibodies. *Expert Opin Ther Targets* **19**, 835-847 (2015).

28. O. ElMokh, V. Taelman, P. Radojewski, M. A. Roelli, A. Stoss, R. A. Dumont, M. S. Dettmer, W. A. Phillips, M. A. Walter, R. P. Charles, MEK Inhibition Induces Therapeutic Iodine Uptake in a Murine Model of Anaplastic Thyroid Cancer. *J Nucl Med* **60**, 917-923 (2019).

29. D. Liu, S. Hu, P. Hou, D. Jiang, S. Condouris, M. Xing, Suppression of BRAF/MEK/MAP kinase pathway restores expression of iodide-metabolizing genes in thyroid cells expressing the V600E BRAF mutant. *Clin Cancer Res* **13**, 1341-1349 (2007).

30. A. L. Ho, R. K. Grewal, R. Leboeuf, E. J. Sherman, D. G. Pfister, D. Deandreis, K. S. Pentlow, P. B. Zanzonico, S. Haque, S. Gavane, R. A. Ghossein, J. C. Ricarte-Filho, J. M. Dominguez, R. Shen, R. M. Tuttle, S. M. Larson, J. A. Fagin, Selumetinib-enhanced radioiodine uptake in advanced thyroid cancer. *N Engl J Med* **368**, 623-632 (2013).

31. D. N. Hayes, A. S. Lucas, T. Tanvetyanon, M. K. Krzyzanowska, C. H. Chung, B. A. Murphy, J. Gilbert, R. Mehra, D. T. Moore, A. Sheikh, J. Hoskins, M. C. Hayward, N. Zhao, W. O'Connor, K. E. Weck, R. B. Cohen, E. E. Cohen, Phase II efficacy and pharmacogenomic study of Selumetinib (AZD6244; ARRY-142886) in iodine-131 refractory papillary thyroid carcinoma with or without follicular elements. *Clin Cancer Res* **18**, 2056-2065 (2012).

32. I. Landa, N. Pozdeyev, C. Korch, L. A. Marlow, R. C. Smallridge, J. A. Copland, Y. C. Henderson, S. Y. Lai, G. L. Clayman, N. Onoda, A. C. Tan, M. E. R. Garcia-Rendueles, J. A. Knauf, B. R. Haugen, J. A. Fagin, R. E. Schweppe, Comprehensive Genetic

Characterization of Human Thyroid Cancer Cell Lines: A Validated Panel for Preclinical Studies. *Clin Cancer Res* **25**, 3141-3151 (2019).

33. J. C. Ricarte-Filho, M. Ryder, D. A. Chitale, M. Rivera, A. Heguy, M. Ladanyi, M. Janakiraman, D. Solit, J. A. Knauf, R. M. Tuttle, R. A. Ghossein, J. A. Fagin, Mutational profile of advanced primary and metastatic radioactive iodine-refractory thyroid cancers reveals distinct pathogenetic roles for BRAF, PIK3CA, and AKT1. *Cancer Res* **69**, 4885-4893 (2009).

34. J. A. Knauf, J. A. Fagin, Role of MAPK pathway oncoproteins in thyroid cancer pathogenesis and as drug targets. *Curr Opin Cell Biol* **21**, 296-303 (2009).

35. M. Uhlen, L. Fagerberg, B. M. Hallstrom, C. Lindskog, P. Oksvold, A. Mardinoglu, A. Sivertsson, C. Kampf, E. Sjostedt, A. Asplund, I. Olsson, K. Edlund, E. Lundberg, S. Navani, C. A. Szigyarto, J. Odeberg, D. Djureinovic, J. O. Takanen, S. Hober, T. Alm, P. H. Edqvist, H. Berling, H. Tegel, J. Mulder, J. Rockberg, P. Nilsson, J. M. Schwenk, M. Hamsten, K. von Feilitzen, M. Forsberg, L. Persson, F. Johansson, M. Zwahlen, G. von Heijne, J. Nielsen, F. Ponten, Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).

36. W. C. van Staveren, D. W. Solís, L. Delys, L. Duprez, G. Andry, B. Franc, G. Thomas, F. Libert, J. E. Dumont, V. Detours, C. Maenhaut, Human thyroid tumor cell lines derived from different tumor types present a common dedifferentiated phenotype. *Cancer Res* **67**, 8113-8120 (2007).

37. T. Pilli, K. V. Prasad, S. Jayarama, F. Pacini, B. S. Prabhakar, Potential utility and limitations of thyroid cancer cell lines as models for studying thyroid cancer. *Thyroid* **19**, 1333-1342 (2009).

38. H. Li, X. Zhou, G. Wang, D. Hua, S. Li, T. Xu, M. Dong, X. Cui, X. Yang, Y. Wu, M. Cai, X. Liao, T. Zhang, Z. Yang, Y. Du, X. Li, CAR-T Cells Targeting TSHR Demonstrate Safety and Potent Preclinical Activity Against Differentiated Thyroid Cancer. *J Clin Endocrinol*

769   *Metab* **107**, 1110-1126 (2022).

770   39.   A. M. Meireles, A. Preto, A. S. Rocha, A. P. Rebocho, V. Máximo, I. Pereira-Castro, S.

771   Moreira, T. Feijão, T. Botelho, R. Marques, V. Trovisco, L. Cirnes, C. Alves, S. Velho, P.

772   Soares, M. Sobrinho-Simões, Molecular and genotypic characterization of human thyroid

773   follicular cell carcinoma-derived cell lines. *Thyroid* **17**, 707-715 (2007).

774   40.   C. M. Stewart, E. L. Siegler, R. L. Sakemura, M. J. Cox, T. Huynh, B. Kimball, L. Mai, I.

775   Can, C. Manriquez Roman, K. Yun, O. Sirpilla, J. H. Girsch, E. Ogbodo, W. Mohammed

776   Ismail, A. Gaspar-Maia, J. Budka, J. Kim, N. Scholler, M. Mattie, S. Filosto, S. S.

777   Kenderian, IL-4 drives exhaustion of CD8(+) CART cells. *Nat Commun* **15**, 7921 (2024).

778   41.   T. Liu, Q. Men, X. Su, W. Chen, L. Zou, Q. Li, M. Song, D. Ouyang, Y. Chen, Z. Li, X. Fu,

779   A. Yang, Downregulated expression of TSHR is associated with distant metastasis in

780   thyroid cancer. *Oncol Lett* **14**, 7506-7512 (2017).

781   42.   M. D'Agostino, M. Sponziello, C. Puppin, M. Celano, V. Maggisano, F. Baldan, M. Biffoni,

782   S. Bulotta, C. Durante, S. Filetti, G. Damante, D. Russo, Different expression of TSH

783   receptor and NIS genes in thyroid cancer: role of epigenetics. *J Mol Endocrinol* **52**, 121-

784   131 (2014).

785   43.   J. A. Fagin, S. A. Wells, Jr., Biologic and Clinical Perspectives on Thyroid Cancer. *N Engl*

786   *J Med* **375**, 2307 (2016).

787   44.   N. Jin, T. Jiang, D. M. Rosen, B. D. Nelkin, D. W. Ball, Dual inhibition of mitogen-activated

788   protein kinase kinase and mammalian target of rapamycin in differentiated and anaplastic

789   thyroid cancer. *J Clin Endocrinol Metab* **94**, 4107-4112 (2009).

790   45.   D. Chakravarty, E. Santos, M. Ryder, J. A. Knauf, X. H. Liao, B. L. West, G. Bollag, R.

791   Kolesnick, T. H. Thin, N. Rosen, P. Zanzonico, S. M. Larson, S. Refetoff, R. Ghossein, J.

792   A. Fagin, Small-molecule MAPK inhibitors restore radioiodine incorporation in mouse

793   thyroid cancers with conditional BRAF activation. *J Clin Invest* **121**, 4700-4711 (2011).

794   46.   V. Subbiah, R. J. Kreitman, Z. A. Wainberg, J. Y. Cho, J. H. M. Schellens, J. C. Soria, P.

Y. Wen, C. Zielinski, M. E. Cabanillas, G. Urbanowitz, B. Mookerjee, D. Wang, F. Rangwala, B. Keam, Dabrafenib and Trametinib Treatment in Patients With Locally Advanced or Metastatic BRAF V600-Mutant Anaplastic Thyroid Cancer. *J Clin Oncol* **36**, 7-13 (2018).

47. I. M. Wakelkamp, O. Bakker, L. Baldeschi, W. M. Wiersinga, M. F. Prummel, TSH-R expression and cytokine profile in orbital tissue of active vs. inactive Graves' ophthalmopathy patients. *Clin Endocrinol (Oxf)* **58**, 280-287 (2003).

48. R. S. Bahn, TSH receptor expression in orbital tissue and its role in the pathogenesis of Graves' ophthalmopathy. *J Endocrinol Invest* **27**, 216-220 (2004).

49. X. Cui, F. Wang, C. Liu, A review of TSHR- and IGF-1R-related pathogenesis and treatment of Graves' orbitopathy. *Front Immunol* **14**, 1062045 (2023).

50. H. Y. Choi, H. Jeon, J. W. Yang, J. H. Ahn, J. H. Jung, Thyroid-Stimulating Hormone Receptor Expression on Primary Cultured Human Extraocular Muscle Myoblasts. *Curr Eye Res* **43**, 1484-1488 (2018).

51. Y. Lu, S. J. Atkins, R. Fernando, A. Trierweiler, T. Mester, A. B. D. Grisolia, P. Mou, P. Novaes, T. J. Smith, CD34- Orbital Fibroblasts From Patients With Thyroid-Associated Ophthalmopathy Modulate TNF-α Expression in CD34+ Fibroblasts and Fibrocytes. *Invest Ophthalmol Vis Sci* **59**, 2615-2622 (2018).

52. J. Furmaniak, J. Sanders, P. Sanders, Y. Li, B. Rees Smith, TSH receptor specific monoclonal autoantibody K1-70TM targeting of the TSH receptor in subjects with Graves' disease and Graves' orbitopathy—Results from a phase I clinical trial. *Clinical Endocrinology* **96**, 878-887 (2022).

53. M. Ryder, M. Wentworth, A. Algeciras-Schimnich, J. C. Morris, J. Garrity, J. Sanders, S. Young, P. Sanders, J. Furmaniak, B. Rees Smith, Blocking the Thyrotropin Receptor with K1-70 in a Patient with Follicular Thyroid Cancer, Graves' Disease, and Graves' Ophthalmopathy. *Thyroid* **31**, 1597-1602 (2021).

54. I. M. Min, E. Shevlin, Y. Vedvyas, M. Zaman, B. Wyrwas, T. Scognamiglio, M. D. Moore,
W. Wang, S. Park, S. Park, S. Panjwani, K. D. Gray, A. B. Tassler, R. Zarnegar, T. J.
Fahey, 3rd, M. M. Jin, CAR T Therapy Targeting ICAM-1 Eliminates Advanced Human
Thyroid Tumors. *Clin Cancer Res* **23**, 7569-7583 (2017).

55. V. G. Bhoj, L. Li, K. Parvathaneni, Z. Zhang, S. Kacir, D. Arhontoulis, K. Zhou, B.
McGettigan-Croce, S. Nunez-Cruz, G. Gulendran, A. C. Boesteanu, L. Johnson, M. D.
Feldman, E. Radaelli, K. Mansfield, M. Nasrallah, R. S. Goydel, H. Peng, C. Rader, M. C.
Milone, D. L. Siegel, Adoptive T cell immunotherapy for medullary thyroid carcinoma
targeting GDNF family receptor alpha 4. *Mol Ther Oncolytics* **20**, 387-398 (2021).

56. L. Xiao, X. C. Liu, K. S. Zhou, Y. Liu, Y. Huang, C. F. Pu, Z. Y. Cao, R. H. Zhu, H. Y. Tang,
Z. P. Huang, H. Yang, X. Huang, Y. P. Song, R. B. Liu, Z. Wu, V. Lu, CoupledCAR
technology for treating thyroid cancer. *Journal of Clinical Oncology* **39**, (2021).

57. J. Ding, D. Li, X. Liu, H. Hei, B. Sun, D. Zhou, K. Zhou, Y. Song, Chimeric antigen receptor
T-cell therapy for relapsed and refractory thyroid cancer. *Exp Hematol Oncol* **11**, 59
(2022).

58. L. Liu, P. A. Mayes, S. Eastman, H. Shi, S. Yadavilli, T. Zhang, J. Yang, L. Seestaller-
Wehr, S. Y. Zhang, C. Hopson, L. Tsvetkov, J. Jing, S. Zhang, J. Smothers, A. Hoos, The
BRAF and MEK Inhibitors Dabrafenib and Trametinib: Effects on Immune Function and in
Combination with Immunomodulatory Antibodies Targeting PD-1, PD-L1, and CTLA-4.
Clin Cancer Res **21**, 1639-1651 (2015).

59. C. Liu, W. Peng, C. Xu, Y. Lou, M. Zhang, J. A. Wargo, J. Q. Chen, H. S. Li, S. S.
Watowich, Y. Yang, D. Tompers Frederick, Z. A. Cooper, R. M. Mbofung, M. Whittington,
K. T. Flaherty, S. E. Woodman, M. A. Davies, L. G. Radvanyi, W. W. Overwijk, G. Lizée,
P. Hwu, BRAF inhibition increases tumor infiltration by T cells and enhances the antitumor
activity of adoptive immunotherapy in mice. *Clin Cancer Res* **19**, 393-403 (2013).

60. V. Verma, N. Jafarzadeh, S. Boi, S. Kundu, Z. Jiang, Y. Fan, J. Lopez, R. Nandre, P. Zeng,

F. Alolaqi, S. Ahmad, P. Gaur, S. T. Barry, V. E. Valge-Archer, P. D. Smith, J. Banchereau,

M. Mkrtichyan, B. Youngblood, P. C. Rodriguez, S. Gupta, S. N. Khleif, MEK inhibition

reprograms CD8(+) T lymphocytes into memory stem cells with potent antitumor effects.

Nat Immunol **22**, 53-66 (2021).

61. T. Mitra, J. Rahman, M. Hwee, R. J. Jesus Faustino Ramos, H. Liu, T. Hartman, J. Cross,

M. de Jesus, M. Huse, V. Longo, P. Zanzonico, S. A. Vardhana, MEK-dependent

bioenergetic demand drives terminal CD8[+] T cell exhaustion. *bioRxiv*,

2025.2004.2030.651453 (2025).

62. T. Gargett, C. K. Fraser, G. Dotti, E. S. Yvon, M. P. Brown, BRAF and MEK inhibition

variably affect GD2-specific chimeric antigen receptor (CAR) T-cell function in vitro. *J*

Immunother **38**, 12-23 (2015).

63. A. Tomida, S. Yagyu, K. Nakamura, H. Kubo, K. Yamashima, Y. Nakazawa, H. Hosoi, T.

Iehara, Inhibition of MEK pathway enhances the antitumor efficacy of chimeric antigen

receptor T cells against neuroblastoma. *Cancer Sci* **112**, 4026-4036 (2021).

64. J. Dörrie, L. Babalija, S. Hoyer, K. F. Gerer, G. Schuler, L. Heinzerling, N. Schaft, BRAF

and MEK Inhibitors Influence the Function of Reprogrammed T Cells: Consequences for

Adoptive T-Cell Therapy. *Int J Mol Sci* **19**, (2018).

65. X. Wang, X. Tao, P. Chen, P. Jiang, W. Li, H. Chang, C. Wei, X. Lai, H. Zhang, Y. Pan, L.

Ding, Z. Liang, J. Cui, M. Shao, X. Teng, T. Gu, J. Wei, D. Kong, X. Si, Y. Han, H. Fu, Y.

Lin, J. Yu, X. Li, D. Wang, Y. Hu, P. Qian, H. Huang, MEK inhibition prevents CAR-T cell

exhaustion and differentiation via downregulation of c-Fos and JunB. *Signal Transduct*

Target Ther **9**, 293 (2024).

66. L. A. Marlow, S. D. Rohl, J. L. Miller, J. A. Knauf, J. A. Fagin, M. Ryder, D. Milosevic, B.

C. Netzel, S. K. Grebe, H. V. Reddi, R. C. Smallridge, J. A. Copland, Methodology, Criteria,

and Characterization of Patient-Matched Thyroid Cell Lines and Patient-Derived Tumor

Xenografts. *J Clin Endocrinol Metab* **103**, 3169-3182 (2018).

67.   A. Zielke, S. Tezelman, G. H. Jossart, M. Wong, A. E. Siperstein, Q. Y. Duh, O. H. Clark,

874   Establishment of a highly differentiated thyroid cancer cell line of Hurthle cell origin.

875   *Thyroid* **8**, 475-483 (1998).

876   68.   R. M. Sterner, R. Sakemura, M. J. Cox, N. Yang, R. H. Khadka, C. L. Forsman, M. J.

877   Hansen, F. Jin, K. Ayasoufi, M. Hefazi, K. J. Schick, D. K. Walters, O. Ahmed, D. Chappell,

878   T. Sahmoud, C. Durrant, W. K. Nevala, M. M. Patnaik, L. R. Pease, K. E. Hedin, N. E.

879   Kay, A. J. Johnson, S. S. Kenderian, GM-CSF inhibition reduces cytokine release

880   syndrome and neuroinflammation but enhances CAR-T cell function in xenografts. *Blood*

881   **133**, 697-709 (2019).

882   69.   A. B. Dietz, P. A. Bulur, R. L. Emery, J. L. Winters, D. E. Epps, A. C. Zubair, S. Vuk-

883   Pavlovic, A novel source of viable peripheral blood mononuclear cells from leukoreduction

884   system chambers. *Transfusion* **46**, 2083-2089 (2006).

885   70.   J. Sanders, R. N. Miguel, J. Furmaniak, B. R. Smith, TSH receptor monoclonal antibodies

886   with agonist, antagonist, and inverse agonist activities. *Methods Enzymol* **485**, 393-420

887   (2010).

888   71.   R. L. Sakemura, M. Hefazi, M. J. Cox, E. L. Siegler, S. Sinha, M. J. Hansen, C. M. Stewart,

889   J. M. Feigin, C. Manriquez Roman, K. J. Schick, I. Can, E. E. Tapper, P. Horvei, M. M.

890   Adada, E. D. Bezerra, L. A. Kankeu Fonkoua, M. W. Ruff, C. L. Forsman, W. K. Nevala,

891   J. C. Boysen, R. C. Tschumper, C. L. Grand, K. R. Kuchimanchi, L. Mouritsen, J. M.

892   Foulks, S. L. Warner, T. G. Call, S. A. Parikh, W. Ding, N. E. Kay, S. S. Kenderian, AXL

893   Inhibition Improves the Antitumor Activity of Chimeric Antigen Receptor T Cells. *Cancer*

894   *Immunol Res* **11**, 1222-1236 (2023).

895   72.   M. J. Cox, C. Manriquez Roman, E. E. Tapper, E. L. Siegler, D. Chappell, C. Durrant, O.

896   Ahmed, S. Sinha, R. Mwangi, N. S. Scott, M. Hefazi, K. J. Schick, P. Horvei, M. W. Ruff,

897   I. Can, M. Adada, E. Bezerra, L. A. Kankeu Fonkoua, S. A. Parikh, N. E. Kay, R.

898   Sakemura, S. S. Kenderian, GM-CSF disruption in CART cells modulates T cell activation

and enhances CART cell anti-tumor activity. *Leukemia*, (2022).

73.   M. J. Cox, F. Lucien, R. Sakemura, J. C. Boysen, Y. Kim, P. Horvei, C. Manriquez Roman,

M. J. Hansen, E. E. Tapper, E. L. Siegler, C. Forsman, S. B. Crotts, K. J. Schick, M. Hefazi,

M. W. Ruff, I. Can, M. Adada, E. Bezerra, L. A. Kankeu Fonkoua, W. K. Nevala, E. Braggio,

W. Ding, S. A. Parikh, N. E. Kay, S. S. Kenderian, Leukemic extracellular vesicles induce

chimeric antigen receptor T cell dysfunction in chronic lymphocytic leukemia. *Mol Ther* **29**,

1529-1540 (2021).

74.   R. Sakemura, A. Bansal, E. L. Siegler, M. Hefazi, N. Yang, R. H. Khadka, A. N. Newsom,

M. J. Hansen, M. J. Cox, C. Manriquez Roman, K. J. Schick, I. Can, E. E. Tapper, W. K.

Nevala, M. M. Adada, E. D. Bezerra, L. A. Kankeu Fonkoua, P. Horvei, M. W. Ruff, S. A.

Parikh, M. K. Pandey, T. R. DeGrado, L. Suksanpaisan, N. E. Kay, K. W. Peng, S. J.

Russell, S. S. Kenderian, Development of a Clinically Relevant Reporter for Chimeric

Antigen Receptor T-cell Expansion, Trafficking, and Toxicity. *Cancer Immunol Res* **9**,

1035-1046 (2021).

 FIGURES

Figure 1

Figure 1. TSHR is a rational target for CART cell therapy, and TSHR-CART cells demonstrate antigen-specific effector functions *in vitro*. A) Representative IHC staining shows TSHR expression across normal, benign, and malignant thyroid tissues. PTC: papillary

thyroid cancer; FTC: follicular thyroid cancer; HCC: Hurthle cell carcinoma; ATC: anaplastic

thyroid cancer. **B)** H-score quantitation of TSHR expression is shown. H-scores provided are

based on 0 – 3 IHC scoring and percentage of total areas each score (0x%0 + 1x%1 + 2x%2 +

3x%3 = H Score; 0 – 300 range with statistical analysis (mean and SEM; ****$p < 0.0001$, one-way

ANOVA; normal n = 11, benign n = 9, PTC n = 11, FTC n = 9, HCC n = 37, ATC n = 10). **C)** H-

score quantification of normal human tissue stained with TSHR (mean and SEM; **$p < 0.01$, ***p

< 0.001, one-way ANOVA; thyroid n = 7, all other tissues n = 2). **D)** Representative histogram

shows CAR expression on TSHR-CART cells as determined by flow cytometry. UTD T cells were

used as a negative control. **E)** Luciferase$^+$ TSHR$^+$ or TSHR$^-$ FTC-133 target cells were co-cultured

with either UTD or TSHR-CART cells at varying effector-to-target ratios, and cytotoxicity was

measured via bioluminescence after 48 hours (mean and SEM; ***$p < 0.001$, ****$p < 0.0001$, two-

way ANOVA; 3 biological replicates). **F-G)** CFSE-labeled UTD or TSHR-CART cells were co-

cocultured with media alone (negative control), PMA/ionomycin (positive control), or TSHR$^+$ or

TSHR$^-$ FTC-133 target cells. After five days, T cell proliferation was assessed by flow cytometry

using CFSE staining (F) or absolute counts of CD3$^+$ cells (G) (mean and SEM; ****$p < 0.0001$,

two-way ANOVA; 3 biological replicates). **H)** UTD or TSHR-CART cells were-cocultured with

TSHR$^+$ FTC-133 target cells at a 1:5 ratio for four hours, and degranulation was then assessed

by flow cytometry of CD3$^+$ CD107a$^+$ cells (mean and SEM; ****$p < 0.0001$, two-way ANOVA; 2

biological replicates). **I-K)** UTD or TSHR-CART cells were-cocultured with TSHR$^-$ or TSHR$^+$ FTC-

133 target cells at a 1:1 ratio for three days, and supernatants were analyzed for human cytokines

via a 38-multiplex assay (mean and SEM; **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, two-way

ANOVA; 2 biological replicates). **L)** Heat map of additional analytes from the 38-plex assay

(median; **$p < 0.01$, ****$p < 0.0001$, two-way ANOVA; 2 biological replicates).

Figure 2

Figure 2. TSHR-CART cells exhibit potent antitumor activity in vivo against Hurthle cell carcinoma. A) Representative IHC staining of TSHR expression across normal and HCC tissues. Hurthle cell carcinoma: HCC; primary tumor with minimal invasion: PMI; primary tumor with extensive invasion: PEI; recurrent tumor with extensive invasion: REI; metastatic tumors: M. **B)** TSHR expression was quantified using H-Score (based on 0 – 300 IHC scoring and percentage of total areas each score, 0 – 300 range with statistical analysis) (mean and SEM; **p < 0.01, ***p < 0.001, ****p <0.0001, one-way ANOVA. Normal n=11, HCC n=37, PMI n=26, PEI n=11, REI n=34, M n=16). **C)** Experimental schema of a HCC xenograft model. NSG mice were subcutaneously engrafted with TSHR$^+$ XTC.UC1 thyroid cancer cells in Matrigel. Three days later, after engraftment was confirmed, mice were randomized according to tumor volume to receive either UTD (10 X 10^6) or TSHR-CART cells (10 x 10^6) intravenously. Mice were monitored for well-being, tumor volume, and overall survival. At day 51, mice treated with TSHR-CART cells which achieved complete remission were rechallenged subcutaneously with XCT.UC1 cells in the opposite flank. Additionally, an untreated group was engrafted with XCT.UC1 cells and included as a negative control. Mice were monitored for well-being, tumor volume, T cell expansion, and overall survival. **D)** Tumor volume was assessed via serial caliper measurements (mean and SEM; ****p < 0.0001, two-way ANOVA; n=6 mice per group). **E)** Kaplan-Meier survival curve (***p < 0.001, Log-rank test; n=6 mice per group). **F)** Tumor volume as assessed by serial caliper measurements (mean and SEM; ****p < 0.0001, two-way ANOVA; n=6 mice per group). **G)** Peripheral blood was collected 21 days after tumor rechallenge and assessed by flow cytometry using absolute counts of hCD3$^+$ cells (mean and SEM; **p < 0.01, two-tailed unpaired t-test; n = 3 mice per group).

Figure 3

Figure 3. TSHR-CART cells exhibit dose-dependent antitumor activity in PDX models of TSHR⁺ PDTC. A) Experimental schema of a PDTC xenograft model. NSG mice were subcutaneously engrafted with TSHR$^+$ THJ-529 thyroid cancer cells in Matrigel. One week later, after engraftment was confirmed, mice were randomized according to tumor volume and received either UTD (5 x 10^6) or TSHR-CART cells (5 x 10^6 or 1 x 10^7) intravenously. Mice were monitored for well-being, tumor volume, and overall survival. **B)** Tumor volume was assessed via serial

caliper measurements (mean and SEM; **p < 0.01, ****p < 0.0001, two-way ANOVA; n=5-6 mice

per group). **C-E)** Individual tumor growth curves are shown for each treatment group, including

UTD (**C**), TSHR-CART^{low} (**D**), and TSHR-CART^{high} (**E**). N = 5-6 mice/group. **F)** Kaplan-Meier

survival curve (*p < 0.05, ***p < 0.001, Log-rank test; n=5-6 mice per group). **G)** Representative

IHC staining shows human CD3 expression in thyroid tumors excised from mice in each treatment

group. **H)** H-score quantitation of human CD3 expression is shown. H-scores provided are based

on 0 – 3 IHC scoring and percentage of total areas each score (0x%0 + 1x%1 + 2x%2 + 3x%3 =

H Score; 0 – 300 range with statistical analysis) (mean and SEM; **p < 0.01, one-way ANOVA; n

= 5-6 mice per group).

Figure 4

Figure 4. Treatment of ATC PDX tumors with MAPK inhibitors restores TSHR expression. A) Experimental schema of an ATC PDX model. NSG mice were subcutaneously engrafted with TH-560 ATC PDX tumors. When tumors reached approximately 100 mm^3, mice were orally treated with MAPK inhibitors daily and serially monitored for tumor volume. On day 8, mice were sacrificed, and tumors were harvested and assessed for TSHR expression by IHC. **B)** Tumor volumes were assessed via serial caliper measurements in mice treated with trametinib at the indicated doses (mean and SEM; **p < 0.01, two-way ANOVA; n=2 mice per group). **C)** Representative IHC staining (left) shows human TSHR and

pERK expression in thyroid tumors excised from mice in each treatment group. H-scores (right)

provided are based on 0 – 3 IHC scoring and percentage of total areas each score (0x%0 + 1x%1

+ 2x%2 + 3x%3 = H Score; 0 – 300 range with statistical analysis. Mean and SEM; n = 2 mice

per group). **D)** Tumor volumes were assessed via serial caliper measurements in mice treated

with RO5126766 at the indicated doses (mean and SEM; $*p < 0.05$, $**p < 0.01$, two-way ANOVA;

n=2 mice per group). **E)** Representative IHC staining (left) and H-score quantitation (right) show

human TSHR and pERK expression in thyroid tumors excised from mice in each treatment group.

H-scores provided are based on 0 – 3 IHC scoring and percentage of total areas each score

(0x%0 + 1x%1 + 2x%2 + 3x%3 = H Score; 0 – 300 range with statistical analysis. Mean and SEM;

n = 2 mice per group). **F)** Experimental schema of an ATC PDX model. NSG mice were

subcutaneously engrafted with Th-560 ATC PDX tumors. When tumors reached approximately

100 mm^3, mice were orally treated with MAPK inhibitors daily and serially monitored for tumor

volume. On Day 7, mice were sacrificed, and tumors were harvested and assessed for TSHR

expression by IHC. **G)** Representative IHC staining (top) and H-score quantitation (bottom) show

human TSHR expression in thyroid tumors excised from mice in each treatment group. H-scores

provided are based on 0 – 3 IHC scoring and percentage of total areas each score (0x%0 + 1x%1

+ 2x%2 + 3x%3 = H Score; 0 – 300 range with statistical analysis. Mean and SEM; $**p <0.01$,

unpaired two-tailed t-test, n = 3-4 mice per group).

Figure 5. TSHR-CART cells are sensitive to moderate upregulation of TSHR on target cells.

A) THJ-529 cells were cocultured with 10 nM or 1µM vehicle control, trametinib, or both trametinib and dabrafenib. After four days, THJ-529 cells were trypsinized and flowed for TSHR expression. Representative histogram shown. **B)** After four days of MAPK inhibitor co-culture, THJ-529 cells were trypsinized and plated with TSHR-CART cells at a 1:1 ratio and assessed for proliferation five days later as determined by flow cytometry of absolute number of CD3$^+$ cells (mean and SEM; *p < 0.05, **p < 0.01, two-way ANOVA; 1 biological replicate). **C)** Assessment of absolute TSHR expression levels in wildtype or lentivirally transduced TSHR$^+$ FTC-133 tumor cells and wildtype PDX tumors, as determined by flow cytometry using absolute counting beads (mean and SEM; ****p < 0.0001, one-way ANOVA; n= 3-9 technical replicates). **D)** Representative histogram shows different levels of TSHR expression on tumor cells as determined by flow cytometry. **E)** Percentage of killing measured via bioluminescent imaging after 48 hours of co-culture at different

1040   E:T ratios of TSHR-CART cells and luciferase⁺ FTC-133 tumor cells expressing various levels of

1041   TSHR (mean and SEM; ****$p < 0.0001$, two-way ANOVA; 2 biological replicates).

Figure 6

Figure 6. TSHR-
CART effector
functions remain
intact after the
addition of MAPK
inhibitors. A) TSHR-
CART cells were co-
cocultured with either
media alone
(negative control) or
TSHR⁺ FTC-133
target cells at a 1:1
ratio, along with
vehicle control, 10nM
trametinib, or 10nM

both trametinib and dabrafenib. After five days, T cell proliferation was assessed by flow cytometry

using absolute counts of CD3⁺ cells (mean and SEM; two-way ANOVA; 2 biological replicates).

B) TSHR-CART cells were-cocultured with luciferase⁺ TSHR⁺ FTC-133 target cells at various E:T

ratios, along with vehicle control, 10nM trametinib, or 10nM both trametinib and dabrafenib. After

48 hours, target cell killing was assessed by bioluminescence (mean and SEM; two-way ANOVA;

2 biological replicates). **C-D)** In chronic stimulation assays, TSHR-CART cells were co-cocultured

with either media alone (negative control) or irradiated TSHR⁺ FTC-133 target cells at a 1:1 ratio,

along with vehicle control, 10nM trametinib, or 10nM both trametinib and dabrafenib. Irradiated

target cells and MAPK inhibitor-containing media were replenished every other day for one week.

Then, TSHR-CART cells were isolated with CD4 and CD8 beads and plated in proliferation (**C**)

or cytotoxicity (**D**) assays as described above (mean and SEM; two-way ANOVA; 2 biological

replicates).

Figure 7

Figure 7. TSHR-CART cells show the most potent antitumor activity when combined with

MAPK inhibitor administration. A) Experimental schema of an ATC PDX model. NSG mice

were subcutaneously engrafted with Th-560 ATC PDX tumors. When tumors reached

approximately 100 mm^3, mice were orally treated with MAPK inhibitors (0.25 mg/kg trametinib

alone or with 2 mg/kg dabrafenib) daily for either seven days prior to TSHR-CART administration

or concurrently (until endpoints were reached) and serially monitored for tumor volume. Mice were

then randomized by tumor volume and treated with either UTD or TSHR-CART cells (20x10^6 i.v.).

Mice were monitored for well-being and tumor volume. **B)** Tumor volumes were assessed via

serial caliper measurements in mice treated with UTD or TSHR-CART with vehicle control, with

7 days of trametinib, or with concurrent trametinib at the indicated doses (mean and SEM; **p <

0.01, ****p < 0.0001, two-way ANOVA; n=7 mice per group). **C)** Tumor volumes were assessed

via serial caliper measurements in mice treated with UTD or TSHR-CART with vehicle control,

with 7 days of trametinib plus dabrafenib, or with concurrent trametinib plus dabrafenib at the

indicated doses (mean and SEM; *p < 0.05, **p < 0.01, ****p < 0.0001, two-way ANOVA; n=7

mice per group). **D)** Experimental schema of an ATC PDX model. NSG mice were subcutaneously

engrafted with Th-560 ATC PDX tumors. When tumors reached approximately 100 mm^3, mice

were orally treated with MAPK inhibitors (0.25 mg/kg trametinib plus 2 mg/kg dabrafenib) daily for

seven days monitored for tumor volume. After 7 days of MAPK inhibitor treatment, mice were

randomized by tumor volume to be treated with either UTD or TSHR-CART cells (10x10^6 i.v.).

After UTD or TSHR-CART injection, mice were separated to vehicle control or MAPK inhibitors

(for 2 weeks, for 4 weeks, or continuously until experimental endpoints were reached. **E)** Mice

treated with TSHR-CART cells in combination with MAPK inhibitors showed superior tumor

volume control regardless of the duration of MAPK inhibitor treatment. Tumor volumes were

assessed via caliper measurements in mice treated with UTD or TSHR-CART cells in combination

with vehicle control or MAPK inhibitors (mean and SEM; ns = not significant, *p < 0.05, ****p <

0.0001, two-way ANOVA; n=7 mice per group. **F)** Kaplan-Meier survival curve (**p < 0.01, ***p <

0.001, Log-rank test; n=7 mice per group). **G)** *In vivo* TSHR-CART cell expansion was measured

by absolute count of hCD45$^+$ hCD3$^+$ cells per µL of blood via flow cytometry analysis (mean and

SEM; ns = not significant, one-way ANOVA; n = 7-33 mice per group).

Figure 1

Figure 2

A.

B.

C.

Re-challenge model

D.

E.

F.

G.

Figure 3

A.
THJ529 thyroid cancer cells

Rear flank cancer engraftment in 50% Matrigel

NSG

Day 0

Tail vein injection of TSHR CART-cell treatment

Day 7

• Weight loss assessment
• Caliper measurements
• Monitoring for endpoint criteria

Day 3-40

Study conclusion & Data analysis

Day 40

• UTD- 5x10⁶ cells
• CART^{low}- 5X10⁶cells
• CART^{high} -10x10⁶cells

B.

UTD (5x10⁶)
TSHR-CART^{low} (5x10⁶)
TSHR-CART^{high} (1x10⁷)

C.

D.

E.

F.

UTD (5x10⁶)
TSHR-CART^{low} (5x10⁶)
TSHR-CART^{high} (1x10⁷)

G.

	Cross section	20X Mag.
UTD		
TSHR-CART^{low}		
TSHR-CART^{high}		

H.

UTD (5x10⁶)
TSHR-CART^{low} (5x10⁶)
TSHR-CART^{high} (1x10⁷)

Figure 4

A.

B.

Tumor Volume (mm³) vs Day Post-MAPKi Treatment

Legend: Control, 0.05 mg/kg, 0.1 mg/kg, 0.25 mg/kg, 0.5 mg/kg, 0.75 mg/kg

**

C.

TSHR | pERK

CONTROL
0.1 mg/kg
0.5 mg/kg

H-Score vs Dose (mg/kg)

TSHR, pERK

D.

Tumor Volume (mm³) vs Day Post-MAPKi Treatment

Legend: Control, 0.025 mg/kg, 0.05 mg/kg, 0.1 mg/kg, 0.25 mg/kg, 0.5 mg/kg

** *

E.

TSHR | pERK

CONTROL
0.1 mg/kg
0.5 mg/Kg

H-Score vs Dose (mg/kg)

TSHR, pERK

F.

Trametinib (MEKi)+ Dabrafenib (BRAFi)

G.

TSHR⁺ H-Score

**

Control, MEKi/BRAFi

TSHR

Vehicle | MEKi/BRAFi

200μm | 200μm

Figure 5

A. 10nM | 1uM
- Vehicle 1.92% | 1.93%
- MEKi 20.6% | 35.1%
- MEKi/BRAFi 21.4% | 35.4%

TSHR-APC

B. Absolute Number of CD3⁺ Cells
- Vehicle
- MEKi
- MEKi/BRAFi
- p=0.0669
- 10nM, 1μM

C. Absolute TSHR Expression/Cell — ****
- TSHR^{high}
- TSHR$^{high/medium}$
- TSHR$^{medium/low}$
- TSHR^{low}
- TH-560 (ATC PDX)
- TSHR^{negative}

D.
- 99.9% TSHR^{high}
- 99.9% TSHR$^{high/medium}$
- 99.7% TSHR$^{medium/low}$
- 93.0% TSHR^{low}
- 69.1% TH-560 (ATC PDX)
- 2.1% TSHR^{negative}

TSHR-APC

E. % Target Cell Killing vs E:T Ratio
- 2.5:1, 1.25:1, 0.625:1
- ****
- TSHR^{high}
- TSHR$^{high/medium}$
- TSHR$^{medium/low}$
- TSHR^{low}
- TSHR^{negative}

Figure 6

Figure 7

A.

B.

Tumor Volume (mm³) vs Day Post-CART Treatment

Legend:
- UTD + vehicle
- MEKi ⟶ UTD
- UTD + MEKi
- TSHR-CART + vehicle
- MEKi ⟶ TSHR-CART
- TSHR-CART + MEKi

C.

Tumor Volume (mm³) vs Day Post-CART Treatment

Legend:
- UTD + vehicle
- MEKi/BRAFi ⟶ UTD
- UTD + MEKi/BRAFi
- TSHR-CART + vehicle
- MEKi/BRAFi ⟶ TSHR-CART
- TSHR-CART + MEKi/BRAFi

D.

E.

Tumor Volume (mm³) vs Day Post-CART Treatment

2-week tram/dab stop
4-week tram/dab stop

Legend:
- UTD + vehicle
- UTD + MEKi/BRAFi
- TSHR-CART + vehicle
- TSHR-CART + MEKi/BRAFi (2 weeks)
- TSHR-CART + MEKi/BRAFi (4 weeks)
- TSHR-CART + MEKi/BRAFi (continous)

F.

Probability of Survival vs Day Post- CART treatment

p=0.0593

Legend:
- UTD + vehicle
- UTD + MEKi/BRAFi
- TSHR-CART + vehicle
- TSHR-CART + MEKi/BRAFi (2 weeks)
- TSHR-CART + MEKi/BRAFi (4 weeks)
- TSHR-CART + MEKi/BRAFi (continous)

G.

Absolute number of hCD3⁺ T cells/uL

ns

Legend:
- UTD + vehicle
- UTD + MEKi/BRAFi
- TSHR-CART + vehicle
- TSHR-CART + MEKi/BRAFi

Tissue core | Tissue core 20x | Tissue core | Tissue core 20x | Tissue core | Tissue core 20x | Tissue core | Tissue core 20x

CERVIX | BONE MARROW | BLADDER | ADRENAL
SALIVARY | FALLOPIAN TUBE | THYMUS | THYROID
SKIN | COLON | BREAST | SPLEEN
UTERUS | TONSIL | LIVER | KIDNEY
PANCREAS | PLACENTA | TESTIS | PROSTATE
LUNG | BRAIN | STOMACH | Tissue TMA

Sup. Fig. S2

A.

| VH region | VL region | CD8a Hinge | CD8 TM | 4-1BB | CD3-ζ |

B.

PBMC isolation

T cell isolation

T cell stimulation (CD3/CD28 beads)

Day 0

CART cell count

Feed CART cells

Days 3-5

Day 1

Lentiviral transduction with TSHR CAR

Beads removal

Flow cytometry to assess CAR %

Count and feed CART cells

Day 6

Freeze and cryopreserve CART cells

Day 8

C.

% of CD45⁺ CD3⁺

CD4
CD8

UTD   TSHR-CART

D.

% of CD45⁺ CD3⁺

T_cm
T_naive
T_emra
T_em

UTD   TSHR-CART

Sup. Fig. S3

A.
UTD   TSHR-CART

Day 0
Day 2
Day 5

CFSE FITC

B.
Absolute TSHR Expression/Cell

****

THJ-529^WT   THJ-529^TSHR

C.
% TSHR⁺ Live Cells

****

THJ-529^WT   THJ-529^TSHR

D.
% Target Cell Killing

E:T Ratio

2.5:1   1.25:1   0.625:1

E.
Absolute Number of CD3⁺ T cells

****

Media   THJ-529^WT   THJ-529^TSHR

● UTD
■ TSHR-CART

● UTD + THJ-529^WT
■ UTD + THJ-529^TSHR
◆ TSHR-CART + THJ-529^WT
▲ TSHR-CART + THJ-529^TSHR

****

F.
% CD107a⁺ of CD3⁺ T cells

****

● UTD
■ TSHR-CART

Media   THJ-529^WT   THJ-529^TSHR

G.
% IL-2⁺ of CD3⁺ T cells

****

● UTD
■ TSHR-CART

Media   THJ-529^WT   THJ-529^TSHR

H.
%TNFα⁺ of CD3⁺ T cells

****

● UTD
■ TSHR-CART

ns

Media   THJ-529^WT   THJ-529^TSHR

A.

%CD107a+ of CD3+ T cells

ns

- TSHR-CART + vehicle
- TSHR-CART + MEKi
- TSHR-CART + MEKi/BRAFi

Media FTC-133^TSHR

B.

% of CD3+ T cells

ns ns ns

IL-2 GM-CSF MIP1β

- TSHR-CART + vehicle
- TSHR-CART + MEKi
- TSHR-CART + MEKi/BRAFi

C.

irr. TSHR+ FTC-133 + TSHR-CART + DMSO control or 10 nM MEKi or MEKi/BRAFi → isolate TSHR-CART and replate in assays

0 2 4 6 7 Days

irr. TSHR+ FTC-133 + DMSO control or 10 nM MEKi or MEKi/BRAFi

D.

TSHR-CART + media + DMSO — 29.3%

TSHR-CART + media + MEKi — 37.7%

TSHR-CART + media + MEKi/BRAFi — 29.3%

TSHR-CART + FTC-133^TSHR + DMSO — 80.3%

TSHR-CART + FTC-133^TSHR + MEKi — 81.1%

TSHR-CART + FTC-133^TSHR + MEKi/BRAFi — 76.7%

SSC

CD3 APC-Cy7

Sup. Fig. S6

A.

B.

TSHR protein expression lost within 48 hours of treatment cessation

Prior to MEK/BRAF inhibitors	Day 0 post 7-day MEK/BRAF inhibitors
H-score = 10.5	H-score = 78.6
Day 2 post 7-day MEK/BRAF inhibitors	Day 4 post 7-day MEK/BRAF inhibitors
H-score = 14.6	H-score = 15.4

Sup. Fig. S7

Sup. Fig. S8

A.

B.

C.

D.

E.

F.

G.

- UTD + vehicle
- UTD + MEKi/BRAFi
- TSHR-CART + vehicle
- TSHR-CART + MEKi/BRAFi

Sup. Fig. S9