Characterising tandem repeat complexities across long-read sequencing platforms with TREAT and *otter*

Niccolo' Tesi,[1,2,3,*,#] Alex Salazar,[1,*,#] Yaran Zhang,[1] Sven van der Lee,[1,2]

Marc Hulsman,[1,2,3] Lydian Knoop,[1] Sanduni Wijesekera,[1] Jana Krizova,[1]

Anne-Fleur Schneider,[1] Maartje Pennings,[5] Kristel Sleegers,[4] Erik-Jan

Kamsteeg,[5] Marcel Reinders,[3] & Henne Holstege [1,2,3,#]

* Authors contributed equally

\# Corresponding authors

[1] Section Genomics of Neurodegenerative Diseases and Aging, Department of Clinical Genetics, Vrije

Universiteit Amsterdam, Amsterdam UMC, Amsterdam, The Netherlands

[2] Alzheimer Center Amsterdam, Department of Neurology, Amsterdam Neuroscience, Vrije Universiteit

Amsterdam, Amsterdam UMC, Amsterdam, The Netherlands

[3] Delft Bioinformatics Lab, Delft University of Technology, Delft, The Netherlands

[4] Complex Genetics of Alzheimer's Disease Group, Antwerp Center for Molecular Neurology, VIB,

Antwerp B-2610, Belgium

[5] Department of Genome Diagnostics, Radboud University Medical Center, Nijmegen, The Netherlands

Keywords

LRS Special Issue; Tandem repeats; Long-read sequencing; PacBio; Nanopore; Genotyping;

Cross-platform compatibility;

Abstract

Tandem repeats (TR) play important roles in genomic variation and disease risk in humans. Long-read sequencing allows for the accurate characterisation of TRs, however, the underlying bioinformatics perspectives remain challenging.

We present *otter* and TREAT: *otter* is a fast targeted local assembler, cross-compatible across different sequencing platforms. It is integrated in TREAT, an end-to-end workflow for TR characterisation, visualisation and analysis across multiple genomes.

In a comparison with existing tools based on long-read sequencing data from both Oxford Nanopore Technology (ONT, Simplex and Duplex) and PacBio (Sequel 2 and Revio), *otter* and TREAT achieved state-of-the-art genotyping and motif characterisation accuracy.

Applied to clinically relevant TRs, TREAT/*otter* significantly identified individuals with pathogenic TR expansions. When applied to a case-control setting, we significantly replicated previously reported associations of TRs with Alzheimer's Disease, including those near or within *APOC1* (p=2.63x10-9), *SPI1* (p=6.5x10-3) and *ABCA7* (p=0.04) genes.

We finally used TREAT/*otter* to systematically evaluate potential biases when genotyping TRs using diverse ONT and PacBio long-read sequencing datasets. We showed that, in rare cases (0.06%), long-read sequencing suffers from coverage drops in TRs, including the disease-associated TRs in *ABCA7* and *RFC1* genes. Such coverage drops can lead to TR mis-genotyping, hampering the accurate characterisation of TR alleles.

Taken together, our tools can accurately genotype TR across different sequencing technologies and with minimal requirements, allowing end-to-end analysis and comparisons of TR in human genomes, with broad applications in research and clinical fields.

1. Introduction

Roughly 30% of the human genome consists of tandem repeats (TR) characterised by one or more repeat motifs that are defined by their consecutive repetition.[1] This repetitive pattern often leads to DNA instability, facilitating not only expansions and contractions of the repeating motif sequence, but also allelic diversity within the sequence.[2,3] Several definitions of TRs have been introduced based on the motif length and size variability, including microsatellites, minisatellites, and macrosatellites. Microsatellites (or short tandem repeats, STR) are the most abundant TRs in the human genome, are characterised by a repetitive motif of less than 6 base pairs (bp), and tend to cluster in non-coding regions of the genome.[4] Minisatellites are characterised by a repetitive motif with a size ranging 7-100 bp, and they are highly enriched in the telomeric regions of the genome.[5] Macrosatellites are characterised by larger tandem repeat units (>100 bp), and are enriched in the telomeric and centromeric portions of the genome.[6]

TRs can disrupt gene-expression regulation and contribute to over 40 neurological disorders.[1,7,8] Pathogenic TR expansions, surpassing critical lengths, are linked to conditions like spinocerebellar ataxias, Huntington's disease, Fragile-X syndrome, Amyotrophic lateral sclerosis (ALS), and Myotonic Dystrophy.[7–9] For instance, Fragile-X syndrome results from a GGC repeat expansion in the *FMR1* gene, with affected individuals having up to 4,000 copies compared to less than 50 in healthy individuals.[10] Similarly, ALS is caused by an intronic hexa-nucleotide repeat expansion (GCCCCG) in the *C9orf72* gene, exceeding a critical length of more than 200 copies.[11] Beyond diseases-causing, TRs have been also identified as risk factor for complex human diseases: for example, the intronic TR in the *ABCA7* gene is associated with a 4.5-fold increased risk of Alzheimer's Disease (AD) when the TR exceeds 5720 base pairs.[12,13]

Traditionally, the evaluation of TR lengths and sequences has been challenging. Conventional

methods, such as repeat-primed polymerase chain reaction (RP-PCR) and Southern blot

assays, are time-consuming and limited in detecting TRs within PCR-based boundaries.

Short-read sequencing approaches offer an alternative, but their limited read lengths often fail

to span repetitive regions effectively. Despite heuristic methods and statistical modelling,[14–19]

accurately assessing clinically relevant TRs remains difficult. The advent of long-read

sequencing, particularly with PacBio's High Fidelity (HiFi) and Oxford Nanopore Technology's

(ONT) Duplex technology (10-20kb on average, >99% accuracy),[20,21] has significantly

improved TR evaluation by providing long and accurate sequencing fragments.

Characterising TRs with long-read sequencing technology currently has two main limitations.

First, there is the need to characterise TRs across different (long-read) sequencing

technologies and data-types.[22–24] This is critically important given the growing long-read

sequencing initiatives aiming to comprehensively assess TRs in large genomic datasets,[25]

spanning both population-wide and clinical contexts. For example, some existing tools are

constrained by predefined TR databases, hindering the identification of new TR features such

as novel motif sequences;[26] other tools are technology and data-type-dependent,[22] or do not

produce generalizable multi-sample outputs.[23,24]

Second, there is a lack of comprehensive studies that have investigated potential biases when

sequencing TRs. For example, DNA methylation has been previously shown to impact base-

calling accuracy in long-read sequencing data.[27–29] Similarly, the formation of secondary

structures due to TRs could impact enzyme efficiency (*e.g.* polymerase or nanopores),[30]

potentially reducing read-quality and sequencing throughput in current long-read sequencing

technologies. Furthermore, some technologies require the alignment of noisy reads to

generate high quality consensus sequences, which might be more difficult in case of repetitive

regions. These problems may impact genotyping accuracy and lead to incorrect assessments

of allele-sequences, including disease-associated TRs in patients.

Here, we present TREAT (Tandem REpeat Annotation Toolkit), a unified workflow for

characterising TRs across multiple genomes, cross-compatible with diverse long-read

technologies and data-types (*e.g.* read-alignments and *de novo* assemblies). TREAT employs

a novel generic targeted local assembler, *otter*, that can adapt to different sequencing

chemistries to accurately characterise TRs. We benchmarked TREAT and *otter* with currently

available tools for TR genotyping (PacBio's TRGT and LongTR)[22,31] in terms of genotyping

accuracy, motif identification, and running performances. We then showcase TREAT and *otter*

applicability in a population-, clinical-, and case-control setting. Finally, we performed a

systematic analysis of ~864K genome-wide TRs in CHM13 reference genome to evaluate

sporadic coverage drops that can affect TR genotyping accuracy. We did so using the well-

characterised HG002 genome based on long-read sequencing data from ONT (Duplex and

Simplex), HiFi and non-HiFi data from PacBio's Revio and Sequel 2 instruments.

2. Results

2.1 Cross-compatible workflow for characterising tandem repeats with *otter* and TREAT

We present *otter* and TREAT, two bioinformatic tools that enable tandem repeat (TR)

characterisation across different long-read sequencing technologies and data-types with

minimal input requirements (*Figure 1*). *Otter* is a stand-alone generic targeted local assembler

for long-read sequencing data, which automatically adapts to sequencing error-rates and

coverage levels per target region. TREAT integrates *otter* to enable end-to-end unified

workflow for *de novo* motif characterisation and downstream analysis, including TR

visualisation, outlier-based and case-control comparisons (see *Methods*). Both tools require

sequencing data aligned to a reference genome (.bam files), the reference genome used

(.fasta file), and the coordinates of the regions of interest (chromosome, start and end

positions encoded in a .bed file). TREAT/*otter* outputs a multi-sample gVCF (Genomic Variant

Call Format) file reporting genotyped alleles, their size and relative repeat content (motif and

number of copies), of each TR in each sample.

Otter is written in C++ and the source code is freely available at

https://github.com/holstegelab/otter.

TREAT is written in Python and R (for plots). The source code is freely available at

https://github.com/holstegelab/treat along with example datasets, documentation, a dedicated

Conda configuration file and a Docker image to ease the installation.

Figure 1: Schematic workflow of TREAT and otter. **A.** Shows TREAT workflow, highlighting the required

inputs, the main features, and the main outputs of the tool. The red box highlights the main genotyping

engine based on *otter.* **B.** Summarises the main algorithmic steps of *otter*, a novel targeted local

assembler for long-read sequencing data.

146 2.2 *Otter* and TREAT enable accurate characterisation of both

147 PacBio and ONT long-read data

We benchmarked TREAT and *otter* with TRGT and LongTR,[22,31] currently available tools to

characterise TRs in long-read sequencing data. We compared: (*i*) genotyping accuracy, *i.e.*

the accuracy of the predicted allele sequences for a TR, (*ii*) motif characterisation accuracy,

and (*iii*) computational resources. We varied different long-read sequencing technologies

(PacBio Sequel 2 and Revio, ONT Simplex and Duplex) as well as different coverages (5x,

10x, 15x, 20x, 25x, and 30x) of HG002.[32] We focussed on a set of 161,382 TRs from PacBio's

repeat catalogue (see *Methods*). Predicted TR alleles were compared to the *expected* alleles

based on the HG002 T2T assembly (see *Methods*).

In PacBio data, we found comparable genotyping accuracy between *otter* (TREAT genotyping

engine) and TRGT, for both Sequel 2 and Revio datasets, although *otter* generated more

accurate genotypes for larger TRs (*e.g.* >500bp), achieving average error-rates of 0.2-2.5%,

compared to 0.6-3.8% of TRGT. Both methods were more accurate when increasing the

coverage, although this was less pronounced for larger TRs (>500 bp). Notably, genotyping

accuracy for both *otter* and TRGT was higher for PacBio's Sequel 2 data in comparison with

Revio data (*Figure 2A* and *Supplementary Results*). In ONT data, *otter* was generally more

accurate than LongTR although differences for large TRs were less clear. For both tools, we

observed better accuracies for Duplex data in comparison to Simplex data (*Figure 2B* and

Supplementary Results). Altogether, our benchmark across all tools revealed that PacBio led

to more accurate genotypes for TRs <500 bp, with PacBio and ONT having similar

performances for TRs ranging 500-1000 bp, and ONT leading to more accurate genotypes for

TRs >=1000 bp (*see Figure 2A-B and Supplementary Results*).

The above observations remain when using different distance metrics and partitioned by

different TR-types. For example, we observed similar performances when using the raw edit

distance and correlation between observed and expected allele sizes (*Figure S1* and

Supplementary Results). Furthermore, we found that TRs characterised by dinucleotide

repeat motifs were on average less accurate than TRs with longer motifs (*Figure S2*). The

fraction of alleles perfectly genotyped (*i.e.* with an edit distance of 0), compared to expected

alleles, increased with higher coverage across all technologies and tools (*Figure S3*), with

Sequel 2 data having the largest fraction of alleles perfectly matched, and ONT Simplex having

the least. In PacBio Sequel 2 and Revio data, TRGT generated a slightly higher fraction of

perfectly matched alleles with respect to *otter* (max difference 2.8%). In ONT data, *otter*

outperformed LongTR in all settings.

Similarly, TREAT, which makes use of TR-genotypes from *otter*, achieved similar motif

characterisation accuracy relative to TRGT (*Figure 2C*). In the GRCh38 reference genome,

the motifs of the 161K TRs were mostly dinucleotide (49%), followed by tetranucleotide (22%)

and 16+ bp motifs (11%) (*Figure S4*). Because LongTR does not directly report the identified

TR motifs, we compared TR motifs between TREAT and TRGT. On average, TREAT identified

the same motifs as TRGT in 96% of cases (*Figure 2C*), and this did not change for different

technologies (Sequel 2 or Revio) or different coverages. We observed a higher concordance

in motif detection between tools for shorter motifs (*Figure 2C*). When looking at the motifs

identified by TREAT on the GRCh38 reference genome, these matched known TR annotations

in 91% of the cases.

Finally, we evaluated the computational performances of *otter* (stand-alone), TREAT

(integrated workflow with *otter*), TRGT and LongTR. When using four threads, TRGT and *otter*

had similar run-time performances, while both were slightly faster than the integrated workflow

from TREAT (*Figure 2D*). On the other hand, for the ONT data, *otter* and TREAT were faster

than LongTR. In terms of memory consumption, performances were comparable between

TREAT and LongTR, while *otter* and TRGT used significantly less memory (*Figure 2D*). When

evaluating the multithreading capabilities in TREAT, we saw that when increasing the number

of threads to 6, 8, 10 and 12, the running times decreased by 1.3-, 1.5-, 1.6- and 1.8-fold (on

average across the different technologies), compared to 4 CPU threads (*Figure S5*).

In addition to the high-quality HiFi data, PacBio can output non-HiFi data, *i.e.* reads that did

not pass PacBio's internal HiFi quality thresholds, and that constitute a significant fraction of

all sequenced data (45% in HG002). We explored whether integrating both HiFi and non-HiFi

data could improve *otter*'s capability to accurately characterise TR allele sequences. Because

Revio uses a subset of these non-HiFi reads (those with at least 90% read quality) to improve

throughput and accuracy via DeepConsensus,[33] we performed this analysis only for Sequel 2

data. We found that non-HiFi data improved accuracy across all TR-lengths. Specifically, when

integrating non-HiFi reads of at least 85-90% read quality, genotyping accuracy improved by

nearly two-fold (*Figure S6*).

*Figure 2: Benchmarking between TREAT/otter, TRGT, and LongTR. **A.** Genotyping accuracy of otter*

*and TRGT on PacBio Sequel 2 and Revio data, stratified by TR size and sequencing depth. **B.***

Genotyping accuracy of otter and LongTR on ONT Simplex and Duplex data, stratified by TR size and

*sequencing depth. **C.** Motif identification accuracy of TREAT and TRGT on PacBio Sequel 2 and Revio*

*data, showing the overlap of matching motifs, stratified by motif size and sequencing depth. **D.** Memory*

usage and running time of otter, TREAT, TRGT, and LongTR, across technologies and sequencing

coverages.

2.3 TREAT's unified workflow enables diverse characterisations of tandem repeats

We applied TREAT's unified workflow to characterise TRs in a population and clinical setting.

First, we genotyped the set of 161K TRs in 47 genomes from the Human Pangenome

Research Consortium (HPRC),[34] for which PacBio HiFi data was available. We then extracted

the top 20% most variable TRs (N=32,208, based on the coefficient of variation, see *Methods*),

and performed a principal component analysis (PCA, *Figure 3A*) on the joint allele sizes (*i.e.*

the sum of the maternal and paternal alleles). We found that PC1 explained 12% of the total

variance and genetically represented the African-American axis, while PC2 explained 3.5% of

variance and corresponded to the American-Asian axis. The explained variance was similar

to that of a PCA including 40/47 matching samples and 30,544 random common (minor allele

frequency >10%) Single Nucleotide Polymorphisms (SNPs) (PC1: 14%, PC2: 4%, *Figure S7*).

We then used TREAT's outlier analysis to detect and score extreme TR expansions or

contractions of 35 clinically relevant TRs (*Table S1*) in 47 genomes from the HPRC, as well

as two Dutch CANVAS patients and 10 parent-offspring duos (see *Methods*).[35,36] The two

CANVAS patients were previously characterised to harbour expansions in the intronic TR in

RFC1.[35] For all individuals, PacBio HiFi data were generated with Sequel 2 instrument. In total,

we identified 30 instances where the TR length in certain samples were significantly different

from the distribution of TR lengths across all 69 genomes. The most significant deviations

were observed for the two CANVAS patients in the TR intronic of *RFC1* gene ($p<2 \times 10^{-16}$ for

both patients, *Figure 3B-D*). The joint allele size for these samples was 78- and 89-fold higher

than the median TR size across all 69 genomes. Significant TR expansions were also found

in the TR in *ATXN8* gene (HG01123 sample, $p<2 \times 10^{-16}$, *Figure S8*), and in *DMD* gene

(HG02622 sample, $p=6.90 \times 10^{-3}$, *Figure S9*). Interestingly, in the TR intronic of *RFC1* gene,

we also observed a significant heterozygous expansion in one parent of the parent-offspring

duos ($p=1.7 \times 10^{-3}$ and $p=5.18 \times 10^{-11}$, respectively for the short and long alleles, *Figure 3B*).

Unexpectedly, the child reported a homozygous non-expanded genotype, suggesting a mis-

assembly or an allele dropout.

Finally, we applied TREAT to characterise unique TRs that are present in CHM13 reference

genome but absent in GRCh38 across the 47 HPRC genomes. We first curated a set of ~864K

genome-wide TRs in the CHM13 reference genome (see *Methods*). We evaluated genotyping

accuracy by applying TREAT/*otter* to CHM13-aligned long-read datasets of HG002 (PacBio's

Revio and Sequel 2 as well as ONT's Duplex and Simplex). We observed similar

performances as those observed when using ~161K TRs from GRCh38 (see *Figure S10* and

Supplementary Results). These results showcase *otter* and TREAT's ability to *de novo*

characterise TRs across different reference genomes, and without prior knowledge of TR motif

composition. Based on a CHM13-to-GRCH38 liftover procedure, we found 1017 unique TRs

present in CHM13 and absent in GRCh38, 37% of which overlapped coding sequences

(*Supplementary Methods* and *Table S2*). We used TREAT/*otter* to characterise these TRs

across the 47 HPRC genomes and found a mean TR size of 129 bp (median=45 bp), mainly

composed of trinucleotide motifs (42%), followed by homopolymers (26%), and 6+ nucleotide

motifs (22%, *Figure S11*).

*Figure 3: TREAT visualisation and analysis modules. **A.** The PCA of the ancestry-based analysis based*

*on the 20% most variable TRs across 47 HPRC genomes. **B.** The main TR in the RFC1 gene. Y-axis:*

individuals, X-axis: TR size (in bp). Blue dots: smaller allele, orange dots: larger allele, red dots;

homozygous genotypes. Dashed line: the allele in the reference genome GRCh38. The right side of the

plot reports, for each sample and each allele, the motif and relative number of copies. The TR length of

the two CANVAS patients were identified as significant outliers compared to the length-distribution of

*47 samples from the HPRC. **C.** The distribution of allele sizes for the TR in RFC1 gene. **D.** Motif*

representation in CANVAS patients, as produced with MotifScope.[37]

2.4 Tandem repeats may be sensitive to coverage dropouts in long-read sequencing

A closer investigation of PacBio long-read data revealed unexpected drops of coverage in clinical TRs, consequently leading to mis-genotyping of disease-associated TRs. One example is the CANVAS-associated intronic TR in *RFC1,* where the most common allele consists of an (AAAAG)11 motif, with a total size of ~55bp. In CANVAS patients, the TR can range from 2-10 Kbp in total length (*Figure 3B-D* and *Supplementary Results*). In one parent-child duo, we found that the parent harboured an expanded heterozygous version of the TR: a shorter allele with a total length of 244 bp with the (AAAAG)50 motif; and a longer allele with a total length of 2.49 Kbp, composed primarily of the (AAGGG)490 motifs (*Figure 4A*). Long-read sequencing of brain tissue from the same individual (PacBio Sequel 2) confirmed these results, although the longer allele was further expanded by 180 bp (36 additional motif-copies), suggesting a somatic expansion in the brain relative to blood (*Figure 4A*). However, long-read data from the child yielded a homozygous allele-sequence of 63 bp with the (AAAAG)12 motif (*Figure 4A*). This was unexpected as at least one of the two allele-sequences from the parent should be inherited in the child. A closer analysis of HiFi long-read-pileup overwhelmingly supported this genotype. However, we observed an abnormal coverage drop in both the parent and child for this TR, which was alleviated when including non-HiFi data (*Supplementary Results*). After merging HiFi and non-HiFi data of the child, TREAT/*otter* correctly assembled the expanded allele-sequence at 2.65 Kbp in size with (AAGGG)>374. Penta-repeat primed PCR (RP-PCR) confirmed that both parent and child harboured repeat expansions separately composed of the (AAAAG) and (AAGGG) motifs (*Figure S12*). Therefore, HiFi data alone failed to capture this expanded allele-sequence, which was recoverable when including the non-HiFi data.

We observed similar situations of abnormal coverage drops in PacBio data in a separate

intronic TR in *ABCA7*, previously associated with Alzheimer's disease (AD). We

experimentally validated the lengths of this TR using Southern Blotting in a subset of nine

centenarians for which long-read sequencing was performed (*Figure S13* and *Supplementary*

Methods). The local HiFi coverage for these individuals ranged 1-7x (*Figure 4B* and

Supplementary Results). The correlation between experimentally validated alleles and HiFi-

based alleles was 0.58 (Pearson correlation, *Figure 4B*). However, the inclusion of non-HiFi

data increased read-support by four-fold to an average coverage of 22x. As a result, the

correlation with experimentally validated allele sizes increased to 0.99 (*Figure 4B*). These

results highlight standing challenges of characterising TRs with long-read sequencing data,

and suggest systematic biases of long-read sequencing in certain genomic regions.

The above observations motivated us to systematically characterise genome-wide coverage

drops of TRs across long-read sequencing technologies. We did this by investigating coverage

drops in the curated set of ~864K genome-wide TRs in the CHM13 reference genome, using

both PacBio and ONT long-read datasets of HG002 at ~38x coverage (see *Supplemental*

Results and Methods). The average TR-length in this curated set was 93 bp, with motifs being

mostly 16+bp motifs (23%), followed by dinucleotide (18%), tetranucleotide (14%), and

homopolymers (13%, *Figure S4)*. For each TR, we defined the *coverage ratio* by dividing the

local TR coverage *vs.* global genome-wide coverage. We found the average *coverage ratio* to

be 1.01, 1.02, 0.99 and 1.03, respectively for Sequel 2, Revio, ONT Simplex and Duplex

technologies. This indicated generally no unexpected coverage-drops in TRs (*Figure S14A*).

However, 486 (0.06%) unique TRs had ratios below 0.25 (*i.e.* a four-fold lower coverage than

expected based on the global average coverage), of which 454 (93%) were present in the

HG002 T2T reference assembly (*Table S3*). The majority of the low-coverage TRs (294/454,

65%) overlapped gene annotations, potentially leading to mis-genotyping that may impact

biological interpretation. Furthermore, we observed that some of these TRs were within 5 Kbp

of each other, suggesting that coverage drops can extend across multi-Kbp regions. Overall,

we observe significantly more low-coverage TRs in PacBio datasets compared to ONT

(OR=9.4, p-value<2x10-16, Fisher's exact test), with N=437 TRs (89%) being specific to

PacBio datasets. Moreover, 22% of these TRs (N=98) had low coverage in both Sequel 2 and

Revio datasets, suggesting potential systematic challenges in both technologies (*Figure*

S14B-G). This included the intronic TR in *ABCA7*, previously associated with Alzheimer's

disease. Interestingly, the average number of non-HiFi reads in these TRs was 10, indicating

that although reads were generated for these TRs, most were flagged as low-quality during

HiFi data generation.

Within the ONT datasets, we observe significantly more low-coverage TRs in the Duplex

dataset relative to the Simplex dataset (OR=2.6, p-value=1.76x10-3, Fisher's exact test).

We characterised the sequences of all low-coverage TRs to investigate potential characteristic

features. When comparing the 454 low-coverage TRs with the remaining of ~864K genome-

wide TRs, we found that low-coverage TRs were longer (p-value = 8.68e-14; 493 bp longer

on average) and harboured higher GC-content (p-value = 2.28e-50; 17.4% higher on

average). A comparison of dinucleotide content revealed that AG, CC, CG, CT, and GG

dinucleotides were significantly enriched in the low-coverage TRs (*Figure S14H-I*). Moreover,

we found that G-quadruplex DNA secondary structures (G4s) were more likely to occur in low-

coverage TRs (p-value=2.48e-45; 3.76% higher, *Figure S14H* and *Supplementary Methods*).

Figure 4: Coverage drops in TR in RFC1 and ABCA7 genes, associated with CANVAS and Alzheimer's Disease. ***A.*** *Shows the genotyped TR alleles and relative motif characterisation in a parent-child duo using only HiFi data, and HiFi + non-HiFi data. Long-read data from the brain of the parent were also available. Adding non-HiFi data rescued the missing allele in the child.* ***B.*** *Shows the comparison between experimentally validated alleles in the TR intronic of ABCA7 gene, and genotyped alleles based on HiFi data alone, and HiFi + non-HiFi. Experimental validation of TR alleles was performed with Southern Blot assay, and was available for 9 individuals for which long-read data was also*

 available. When adding non-HiFi data, we could recover the expanded alleles in the child that were

missed by HiFi data alone.

2.5 Comparing tandem repeats across multiple genomes in a case-control setting

With the acquired knowledge about possible allele dropouts in TRs, we used TREAT/*otter* in a case-control setting to replicate the association of four TRs that were previously shown to associate with Alzheimer's Disease (AD) risk (*Table 1*, *Table S4*). We did so by using a set of 246 AD patients (mean age = 67.9±9.8, 70% females) and N=248 cognitively healthy centenarians (mean age = 101.2±2.5, 70% females) that were sequenced with PacBio Sequel 2 instrument (*Methods* and *Figure S15*).[36] Across all 494 genomes, we observed a median coverage (HiFi data) of 14, 15, 14, and 4, respectively for the TRs in *APOC1*, *SPI1*, *FERMT2*, and *ABCA7* (*Figure 5A*). The combined allele size (*i.e.* the sum of the maternal and paternal alleles) of the TR nearby *APOC1* (chr19:44921096-44921134) was significantly expanded in AD patients compared to cognitively healthy centenarians (beta=0.38, p=2.63x10-9, *Figure 5B* and *Table 1*). In contrast, the short allele of the TR within *SPI1* gene was significantly contracted in AD patients compared to cognitively healthy centenarians (beta=-0.03, p=6.5x10-3, *Figure 5B* and *Table 1*). The direction of effect of these TRs was in line with the original studies.[38,39] We could not replicate the association of the TR within *FERMT2* (beta=0.01, p=0.27, short allele) (*Figure 5B* and *Table 1*).

For the intronic TR in *ABCA7*, we found significant expansions in AD cases after integrating non-HiFi data (beta=8.63x10-5, p=0.04, joint allele size, *Figure 5C-D*). We note that 22 samples were omitted due to reduced coverage levels even after integrating HiFi and non-HiFi data. We then identified TR size boundaries in the centenarian controls corresponding to the 5th and 95th percentiles of the joint TR allele sizes (2.2 Kbp and 8.4 Kbp, respectively). The

number of centenarians with a TR size lower than the 5th percentile was three-fold higher than

that of AD cases (1-tailed Fisher's exact test p=0.023, OR=3.2, *Figure 5E*), and the number of

AD cases with a TR size larger than the 95th percentile was two-fold higher than that of

centenarians (1-tailed Fisher's exact test p=0.04, OR=2.0, *Figure 5E*). Given the difficulties in

correctly assessing the allele sequences of this TR, we cannot exclude that additional samples

suffer from allelic dropouts, especially for the larger expanded allele-sequences.

*Figure 5: Replication of the association with AD of TRs in APOC1, SPI1, FERMT2 and ABCA7. **A.** The*

*coverage distribution of the four TRs in AD patients and cognitively healthy centenarians. **B.** The TR*

size difference between AD patients and cognitively healthy centenarians in APOC1, SPI1 and

FERMT2. For the associations, we used logistic regression models using the TR size as predictor for

 *AD case-control status. **C.** HiFi and combined HiFi + non-HiFi coverage distribution of the TR intronic*

*of ABCA7 gene. **D.** Comparison of the joint allele size of ABCA7 TR between AD cases and cognitively*

healthy centenarians, respectively using HiFi data, non-HiFi data, and the merged dataset of HiFi and

*non-HiFi. **E.** Number of AD cases and cognitively healthy centenarians in the lower 5^{th} quantile and*

upper 95^{th} quantile. Quantiles were defined based on the distribution of the joint TR-allele size in the

centenarians. We tested for the differential enrichment of AD and centenarians in each quantile with

Fisher's exact tests.

Table 1: Replication of TR previously associated with Alzheimer's Disease (AD)

	TRs previously associated with AD			
Region	chr19:44921096-44921134	chr11:47775208-47775243	chr19:1049436-1050066	chr14:52832909-52832938
Gene	APOC1	SPI1	ABCA7	FERMT2
Best model	Joint alleles	Short alleles	Joint alleles	Short alleles
Beta (OR)	0.38 (1.46)	-0.03 (0.97)	8.63x10-5 (1.01)	0.01 (1.01)
P-value	2.6x10-9	6.5x10-3	0.041	0.27
Original study	38014121	37745545	29589097	37745545
Original OR	NA	-0.01 (0.99)	4.5	0.01 (1.01)
Original model	Longer allele	Joint alleles	Individuals with alleles >5720 bp	Joint alleles
Original method	Logistic regression	Mixed linear models	Fisher's exact	Mixed linear models
Original p-value	4.3x10-10	NA	0.008	NA
Original samples	1489 AD vs. 1492 controls	6328 AD vs. 6580 controls	275 AD vs. 177 controls	6328 AD vs. 6580 controls
Data type	Short read sequencing	Short read sequencing	Southern blot	Short read sequencing

Region: genomic coordinates of the TR with respect to GRCh38; Gene: the closest gene as reported in

the original publications; Best model: model that yielded the most significant association, in our

comparison: Short allele, Long allele or Joint alleles size; Beta (OR): effect size and relative Odds Ratio

with respect to AD: an increased TR size leads to increased AD risk for positive estimates; P-value: p-

value of association. We used logistic regression models using TR size (short allele, long allele and

combined allele size) as predictor for AD case-control status, using 246 AD patients (cases) and 248

cognitively healthy centenarians (controls); Original study: the Pubmed ID of the original study; Original

OR: the odds ratio as reported in the original study; Original model: model used for association in the

original study; Original method: method used for association in the original study; Original p-value: the

416   *p-value reported in the original study; Original samples: the number of AD cases and controls used in*

417   *the original study; Data type: the data on which the association were identified.*

3. Discussion

In this study, we provide novel contributions to better characterise tandem repeats (TRs) with

long-read sequencing data. First, we present our novel tools, *otter* and TREAT, that provide a

unified workflow to accurately characterise TRs using both Pacific Bioscience (PacBio) and

Oxford Nanopore Sequencing Technologies (ONT) datasets. This enabled us to characterise

genome-wide TRs in patients with neurodegenerative diseases and genomes from the Human

Pangenome Research Consortium (HPRC). Second, we show that in rare instances, long-

read sequencing technologies can suffer from abnormal coverage drops in TRs due to

potential systematic challenges, particularly in PacBio's HiFi technology. These coverage

drops can lead to TR mis-genotyping, as we observed in CANVAS and Alzheimer's disease

(AD)-associated TRs*.* Finally, we applied TREAT/*otter* to a case-control setting and replicated

TRs previously associated with AD across 494 long-read sequenced AD patients and

cognitively healthy centenarian genomes.

Our benchmark of *otter* and TREAT highlighted state-of-the-art performances of our tools in

terms of TR genotyping and motif identification accuracy. We showed that *otter,* TREAT, and

other existing tools provide generally accurate characterisations of TRs on both PacBio and

ONT datasets, and with improved accuracies at higher sequencing coverages. Across

technologies, our benchmark revealed that PacBio leads to generally more accurate

genotypes for relatively smaller TRs, with PacBio and ONT having similar performances for

TRs ranging 500-1000 bp, and ONT leading to more accurate genotypes for larger TRs. These

results remained when using other distance metrics as well as in a similar benchmark using

the CHM13 reference genome and a larger set of genome-wide TRs.

Our systematic analysis of coverage drops revealed that overall, coverage drops of TRs are

rare (0.6%), and do not impact the overall genotyping performances of TREAT/*otter* and other

tools. However, our analysis relied on HG002, a highly homozygous genome sequenced at

high coverage (38x). Hence, TR coverage drops may be more prevalent in other (low-

coverage) genomes that harbour expanded TR sequences, especially those with GC-rich

sequences. TRs with coverage drops were often large (>500 bp), high in GC-content, and with

higher densities of predicted G-quadruplex DNA secondary structures (G4s). G4s have been

previously reported to reduce polymerase efficiency.[40] As PacBio's HiFi technology relies on

multiple successful passes of a DNA polymerase in a circular DNA template,[20] we speculate

that the interference of G4s might reduce the number of passes in the circular template,

possibly leading to lower quality reads (non-HiFi reads). Altogether, incidents of TR coverage

drops were enriched in PacBio's Revio and Sequel 2 datasets, and to a lower extent in ONT's

Duplex and Simplex datasets, with ONT Simplex suffering the least. Although rare, we showed

and experimentally validated that coverage drops in TRs can occur at clinically relevant TRs,

requiring extra attention when characterising these TRs. To this end, we showed that local vs.

global coverage ratio is an effective way to identify such problematic regions, and that for

PacBio, these regions can be (in part) rescued by adding noisier non-HiFi data, as shown for

the TRs in *ABCA7* and *RFC1* genes.

TREAT and *otter* can be used to genotype and characterise potentially any type of repetitive

sequences. However, this remains challenging for very large TRs spanning several kilobases,

for example those in telomeric and centromeric regions of the genome. We also note that

regions where sequencing error-rates exceed inter-allele dissimilarities may still be difficult to

genotype. As the error rate in ONT Simplex data is relatively higher than PacBio and ONT

Duplex, this is likely driving the lower genotyping accuracy observed in ONT Simplex. These

limitations are not only specific to TREAT and *otter*, but extend to other existing tools. With

newer sequencing technologies bringing longer read lengths (*e.g.,* ONT ultra-long reads),

together with more complete reference genome assemblies, it might become possible to

genotype any satellite region (micro-, mini-, and macro-satellites) in the genome with TREAT

and *otter*.

We were able to replicate previously reported TRs associated with AD by comparing a cohort

of AD patients and cognitively healthy centenarians. We acknowledge that these TRs were

previously identified using different experimental methods (*e.g.* short-read sequencing,

southern blotting), and analyses strategies (logistic regressions, linear mixed models, fisher's

exact test).[12,38,39] While this heterogeneity hampers the direct comparison of the effect size

estimates, all associations we observed were in the same direction as the original studies. In

particular, the TR intronic of *ABCA7* was shown to carry an odds ratio for AD of 4.5 when one

allele was expanded >5.7 Kbp.[12] Similarly, we observed that individuals carrying larger allele-

sequences were significantly associated with AD. However, in our cohort, the effect was

mainly driven by cognitively healthy centenarians having a shorter joint-allele size (*i.e.* more

AD-protection), rather than AD cases having a more expanded TR-sizes. While we cannot

exclude that we have missed some expanded genotypes due to allele dropouts, the

centenarians that we included were previously shown to be enriched with the protective alleles

in the majority of Single Nucleotide Polymorphisms (SNPs) associated with AD.[41]

In summary, *otter* and TREAT are flexible and accurate bioinformatics tools compatible with

different sequencing platforms and requiring minimal input requirements, that enable end-to-

end analysis and comparisons of tandem repeats in human genomes with broad applications

in research and clinical fields.

4. Methods

4.1 TREAT

The main analysis is the *assembly* analysis, which uses *otter* for TR genotyping, and is

followed by TR content characterisation (identification of motif and number of copies) on the

individual TR alleles. In addition to the *assembly* analysis, TREAT implements a *reads* analysis. Here, TR genotyping is performed using an iterative clustering framework based on TR sizes (*Supplementary Methods*). This is followed by TR content characterisation, which is done on all individual reads (*Supplementary Methods* and *Supplementary Results*). This analysis may be preferred when information from all reads is needed, for example for performing a multiple sequence alignment, or when studying somatic instability.

In all cases, TR content characterisation is performed with *pytrf* (https://github.com/lmdu/pytrf). When multiple motif annotations for the same sequence are found by *pytrf*, a consensus representation of the repeat content is generated. Briefly, if the fraction of sequence annotated with a given motif is >95%, then the relative motif is regarded as the best motif describing the TR. In case two or more motifs are found, each describing a portion of the sequence, then the intersection is calculated by intersecting the motif-specific start and end positions. If the intersection is <90%, then the motifs and the relative number of copies are combined. For example, for sequence TGTGTGTGTGTGTGGAGAGAGAGAGAGA, *pytrf* identifies (*i*) 7 copies of TG (ranging positions 1-14, 50% of the sequence covered), and (*ii*) 7 copies of GA (ranging positions 15-28, 50% of the sequence covered). In this case, the combined sequence annotation will be TG+GA, repeated 7+7 times (see *Supplementary Methods*).

TREAT's analysis module consists of an outlier-detection framework, and a case-control analysis. The outlier-detection scores extreme variations in TR allele sizes across a set of samples. Outliers are detected using a normalised distance that quantifies how far each allele size is from the median allele size, scaled by the variability of the data (*Supplementary Methods*). A p-value for each individual is then calculated by comparing each data point's distance to a chi-squared distribution. The case-control analysis employs logistic regression models to compare allele sizes (short allele, long allele, and joint allele size) between cases and controls.

4.2 *Otter:* a stand-alone, fast, local assembler

Otter is a generic stand-alone method for generating fast local assemblies of a given region or genotyping whole-genome *de novo* assemblies. *Otter* in the main genotyping engine of TREAT *assembly* analysis. Briefly, given a region of interest, *otter* uses the htslib library to identify spanning reads (region of interest is fully contained in the reads) and non-spanning reads (only partially contained) in a given BAM file, and extracts the corresponding subsequence per read based on their alignment (*Figure 1B*).[42] When a reference genome is provided, it will perform local read-realignments on non-spanning reads if it detects a clipping-signal, which can indicate suboptimal mappings to due highly divergent sequences (*Figure 1B*). This is done by aligning (using WFA2-lib alignment library)[43] the flanking sequences of a region (100 bp by default, modifiable with '--flank-size' parameter) derived from the reference genome onto each read, which are then used to recalibrate the corresponding subsequence of the region of interest. Recalibrated non-spanning reads are reclassified as *spanning* if both flanking sequences are successfully aligned with a minimum length and sequence similarity (by default, 90% sequence similarity, modifiable with '--min-sim' parameter). In the context of TRs, this realignment procedure often correctly recalibrates the alignments of TRs with major length and/or motif-composition differences relative to a reference genome.

Otter identifies unique allele-sequences by clustering spanning-reads via pairwise-sequence alignment (*Figure 1B* and *Supplementary Methods*). To manage high somatic variation and/or sequencing errors, *otter* estimates local baseline error-rates per region using a gaussian-kernel density estimator. This produces a one-dimensional distribution of spanning pairwise-sequence distances. In single homozygous allele-sequences, the distribution is unimodal centred at 0. With multiple allele-sequences, the distribution is multimodal, where peaks represent sequence errors between reads from different allele-sequences. *Otter* identifies these peaks and performs hierarchical clustering, stopping when distances exceed the densest peak, partitioning reads into initial clusters. This procedure is followed by a curation

step to ensure sufficient read support, adapting to local coverage (*Figure 1B*). If no maximum

number of alleles (*α*) is enforced, *otter* outputs all clusters. Otherwise, clusters below the

coverage threshold are merged, and if clusters exceed *α*, hierarchical clustering continues

until *α* clusters remain. *Otter* then generates a final consensus sequence per cluster via

pseudo-partial order alignment procedure of spanning and non-spanning reads inspired from

Ye and Ma, 2016.[44]

555 4.3 Genomes included for testing

HPRC: Publicly available PacBio long-read HiFi data of 47 individuals from the Human

Pangenome Reference Consortium (HPRC) were downloaded (*Data Accession*).[34] For the

well characterised HG002 genome,[32] we also downloaded data generated with Oxford

Nanopore (ONT, Duplex and Simplex chemistries) and PacBio Revio technologies. Finally,

we generated long-read sequencing data for HG002 using the PacBio Sequel 2 instrument

across three SMRT cells, keeping both HiFi and non-HiFi data. ONT data was aligned to the

reference genomes (GRCh38 and CHM13) using minimap2 (2.21-r1071, specifying -x map-

ont).[45] PacBio data was aligned using pbmm2 (1.9.0, specifying –preset CCS and –preset

SUBREADS respectively for HiFi and non-HiFi data).[20]

100-plus Study cohort and Alzheimer Dementia Cohort: For the replication of TRs previously

associated with Alzheimer's Disease (AD), we used HiFi sequencing (Sequel 2) data from the

blood DNA of N=246 patients with AD from the Amsterdam Dementia Cohort (ADC),[36,46] and

N=248 cognitively healthy centenarians from the 100-plus Study cohort.[36,47] Ten cognitively

healthy centenarians were sequenced as a trio, including the blood-derived DNA from the

centenarian, the brain-derived DNA from the centenarian and blood-derived DNA from a child

of the centenarian. The combined set of a centenarian and child is referred to as parent-child

duo throughout the manuscript. Sequencing data pre-processing was conducted as previously

described (*Supplementary Methods*).[36]

CANVAS patients: We used the HiFi data (Sequel 2) of two patients diagnosed with CANVAS

(Cerebellar ataxia with neuropathy and vestibular areflexia syndrome), caused by a TR

expansion in *RFC1* gene.[35]

4.4 Evaluating *otter* and TREAT performances

Comparison with existing tools: We compared TREAT/*otter* to TRGT and LongTR.[22,31] For the

comparison, we used the HG002 genome and a set of 161,382 TRs from PacBio's repeat

catalogue (version 0.3.0, available at

https://github.com/PacificBiosciences/trgt/tree/main/repeats). We compared the tools'

genotyped alleles to the expected alleles from the T2T assembly of HG002. As metrics, we

used (*i*) normalised edit distance, (*ii*) raw edit distance, (*iii*) allele size correlation between the

observed and expected alleles, and (*iv*) fraction of perfectly genotyped alleles. In addition, we

evaluated motif identification accuracy, and computational resources.

TREAT/otter applications: We compared the performances of TREAT *assembly* and *reads*

analyses by correlating the estimated TR allele sizes with each other (*Supplementary*

Results). Then, we used TRs for a population stratification analysis: using the set of 161K

TRs, we selected the top 20% most variable TRs based on the coefficient of variation (ratio of

standard deviation to the mean TR joint allele size). Then we applied Principal Component

Analysis (PCA) based on the joint allele sizes. For 40/47 matching samples with Single

Nucleotide Polymorphisms (SNP) data from the 1000Genome project,[48] we also performed

PCA based on 30,544 randomly sampled common (minor allele frequency >10%) SNPs.

To evaluate clinical applicability, we applied the TREAT/*otter* outlier analysis module on the

combined dataset of 47 HPRC genomes plus the two CANVAS patients and the ten parent-

child duos. For this analysis, we focused on 35 clinically relevant TRs (*Table S1*), that were

previously associated with neurological diseases.[7,8,12] Finally, TREAT/*otter* case-control

analysis module was used to replicate the association of four TRs that were previously

associated with Alzheimer's Disease (AD).[12,38,39] The commands used for the outlier and case-

control analyses are available in *Supplementary Methods.*

606 4.5 Systematic analysis of allele dropouts in tandem repeats

Curated set of TRs in CHM13: We downloaded and curated repeat annotations for the CHM13

reference genome (version 2.0, https://github.com/marbl/CHM13, *Supplementary Methods*).

This curated dataset counted 864,424 TRs genome-wide. We extracted the corresponding

parental and maternal allele-sequences in HG002 for these TRs by aligning the HG002 T2T

assembly (version 0.7) to CHM13.[32]

TRs unique to CHM13: We first genotyped the 864K TRs using *otter* in HG002 from different

technologies (Sequel 2, Revio, Simplex and Duplex), and at different coverage levels (5x, 10x,

15x, 20x, 25x and 30x), and calculated the normalised edit distance between observed and

expected TR alleles (*Supplementary Results*). We then focussed on a set of TRs present in

CHM13 and absent in GRCh38, and used TREAT/*otter* to characterise the repeat content of

these TRs in 47 genomes from HPRC.

Evaluation of coverage drops in TR: Using HG002 data from Sequel 2, Revio, Simplex and

Duplex technologies (~30x coverage each), we calculated the ratio between local TR

coverage and average global coverage. TRs where this ratio was <0.25 were regarded as low-

coverage TRs. We then investigated sequence characteristics of low-coverage TR, including

average size, dinucleotide content, and propensity to form G-quadruplex DNA secondary

structures (G4s). For the latter, we used pqsfinder (v2.10.1) with 'min_score = 20' parameter.[49]

Data access

Human Pangenome Consortium data is publicly available and can be downloaded from

https://github.com/human-pangenomics/HPP_Year1_Data_Freeze_v1.0?tab=readme-ov-

file.

Long-read sequencing data generated with PacBio Sequel 2 for the 2 CANVAS patients as

well as 246 AD patients and 248 cognitively healthy centenarians is available upon submission

of a research proposal to the Alzheimer Genetics Hub (AGH, https://alzheimergenetics.org/).

Consent statement

636 The Medical Ethics Committee of the Amsterdam UMC and Radboud UMC approved all

637 studies. All participants and/or their legal representatives provided written informed consent

638 for participation in clinical and genetic studies.

Funding

Part of the work in this manuscript was carried out on the Cartesius supercomputer, which is

embedded in the Dutch national e-infrastructure with the support of SURF Cooperative.

Computing hours were granted to H. H. by the Dutch Research Council (100plus: project#

vuh15226, 15318, 17232, and 2020.030; Role of VNTRs in AD; project# 2022.31, Alzheimers

Genetics Hub project# 2022.38). N.T is appointed at ABOARD and H.H, M.R, S.L are

recipients of ABOARD, a public-private partnership receiving funding from ZonMW

(#73305095007) and Health~Holland, Topsector Life Sciences & Health (PPP-allowance;

#LSHM20106).[41,50] This work is supported by a VIDI grant from the Dutch Scientific Counsel

(#NWO 09150172010083) and a public-private partnership with TU Delft and PacBio,

receiving funding from ZonMW and Health~Holland, Topsector Life Sciences & Health (PPP-

allowance), and by Alzheimer Nederland WE.03-2018-07. S.L. is recipient of ZonMW funding

(#733050512). H.H. was supported by the Hans und Ilse Breuer Stiftung (2020), Dioraphte

16020404 (2014) and the HorstingStuit Foundation (2018). Acquisition of the PacBio Sequel

2 long read sequencing machine was supported by the ADORE Foundation (2022).

Conflicts of interest

657 All authors declare no conflict of interest.

Acknowledgements

660 We are grateful to all the reviewers involved in the peer-review process for their comments

661 which have largely improved our manuscript.

References

1. Hannan, A. J. Tandem repeats mediating genetic plasticity in health and disease. *Nat Rev*

Genet **19**, 286–298 (2018).

2. Lynch, M. *et al.* A genome-wide view of the spectrum of spontaneous mutations in yeast.

Proc Natl Acad Sci U S A **105**, 9272–9277 (2008).

3. Pearson, C. E., Edamura, K. N. & Cleary, J. D. Repeat instability: mechanisms of dynamic

mutations. *Nat Rev Genet* **6**, 729–742 (2005).

4. Subramanian, S., Mishra, R. K. & Singh, L. Genome-wide analysis of microsatellite

repeats in humans: their abundance and density in specific genomic regions. *Genome Biol*

4, R13 (2003).

5. Linthorst, J. *et al.* Extreme enrichment of VNTR-associated polymorphicity in human

subtelomeres: genes with most VNTRs are predominantly expressed in the brain. *Transl*

Psychiatry **10**, 369 (2020).

6. Dumbovic, G., Forcales, S.-V. & Perucho, M. Emerging roles of macrosatellite repeats in

genome organization and disease development. *Epigenetics* **12**, 515–526 (2017).

7. McMurray, C. T. Mechanisms of trinucleotide repeat instability during human

development. *Nat Rev Genet* **11**, 786–799 (2010).

8. Khristich, A. N. & Mirkin, S. M. On the wrong DNA track: Molecular mechanisms of repeat-

mediated genome instability. *J Biol Chem* **295**, 4134–4170 (2020).

9. Stevanovski, I. *et al.* Comprehensive genetic diagnosis of tandem repeat expansion

disorders with programmable targeted nanopore sequencing. *Sci Adv* **8**, eabm5386

(2022).

10. Yu, S. *et al.* Fragile X Genotype Characterized by an Unstable Region of DNA. *Science*

252, 1179–1181 (1991).

11. DeJesus-Hernandez, M. *et al.* Expanded GGGGCC hexanucleotide repeat in noncoding

region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* **72**, 245–256

(2011).

12. On Behalf of the BELNEU Consortium *et al.* An intronic VNTR affects splicing of ABCA7

and increases risk of Alzheimer's disease. *Acta Neuropathologica* **135**, 827–837 (2018).

13. De Roeck, A., Van Broeckhoven, C. & Sleegers, K. The role of ABCA7 in Alzheimer's

disease: evidence from genomics, transcriptomics and methylomics. *Acta Neuropathol*

138, 201–220 (2019).

14. Dolzhenko, E. *et al.* ExpansionHunter: a sequence-graph-based tool to analyze variation

in short tandem repeat regions. *Bioinformatics* **35**, 4754–4756 (2019).

15. Gymrek, M., Golan, D., Rosset, S. & Erlich, Y. lobSTR: A short tandem repeat profiler for

personal genomes. *Genome Res.* **22**, 1154–1162 (2012).

16. Kristmundsdóttir, S., Sigurpálsdóttir, B. D., Kehr, B. & Halldórsson, B. V. popSTR:

population-scale detection of STR variants. *Bioinformatics* **33**, 4041–4048 (2017).

17. Bakhtiari, M., Shleizer-Burko, S., Gymrek, M., Bansal, V. & Bafna, V. Targeted genotyping

of variable number tandem repeats with adVNTR. *Genome Res.* **28**, 1709–1719 (2018).

18. Gelfand, Y., Hernandez, Y., Loving, J. & Benson, G. VNTRseek-a computational tool to

detect tandem repeat variants in high-throughput sequencing data. *Nucleic Acids Res* **42**,

8884–8894 (2014).

19. Eslami Rasekh, M., Hernández, Y., Drinan, S. D., Fuxman Bass, J. I. & Benson, G.

Genome-wide characterization of human minisatellite VNTRs: population-specific alleles

and gene expression differences. *Nucleic Acids Research* **49**, 4308–4324 (2021).

20. Wenger, A. M. *et al.* Accurate circular consensus long-read sequencing improves variant

detection and assembly of a human genome. *Nat Biotechnol* **37**, 1155–1162 (2019).

21. Sereika, M. *et al.* Oxford Nanopore R10.4 long-read sequencing enables the generation

of near-finished bacterial genomes from pure cultures and metagenomes without short-

read or reference polishing. *Nat Methods* **19**, 823–826 (2022).

22. Dolzhenko, E. *et al.* Characterization and visualization of tandem repeats at genome scale.

Nat Biotechnol (2024) doi:10.1038/s41587-023-02057-3.

23. Chiu, R., Rajan-Babu, I.-S., Friedman, J. M. & Birol, I. Straglr: discovering and genotyping

tandem repeat expansions using whole genome long-read sequences. *Genome Biol* **22**,

224 (2021).

24. Masutani, B., Kawahara, R. & Morishita, S. Decomposing mosaic tandem repeats

accurately from long reads. *Bioinformatics* **39**, btad185 (2023).

25. Gustafson, J. A. *et al. Nanopore Sequencing of 1000 Genomes Project Samples to Build*

a   Comprehensive   Catalog   of   Human   Genetic   Variation.

http://medrxiv.org/lookup/doi/10.1101/2024.03.05.24303792   (2024)

doi:10.1101/2024.03.05.24303792.

26. Ren, J., Gu, B. & Chaisson, M. J. P. vamos: variable-number tandem repeats annotation

using efficient motif sets. *Genome Biol* **24**, 175 (2023).

27. Gouil, Q. & Keniry, A. Latest techniques to study DNA methylation. *Essays in Biochemistry*

63, 639–648 (2019).

28. Liu, Y. *et al.* DNA methylation-calling tools for Oxford Nanopore sequencing: a survey and

human epigenome-wide evaluation. *Genome Biol* **22**, 295 (2021).

29. Amarasinghe, S. L. *et al.* Opportunities and challenges in long-read sequencing data

analysis. *Genome Biol* **21**, 30 (2020).

30. Mirkin, S. M. Expandable DNA repeats and human disease. *Nature* **447**, 932–940 (2007).

31. Ziaei Jam, H. *et al.* LongTR: genome-wide profiling of genetic variation at tandem repeats

from long reads. *Genome Biol* **25**, 176 (2024).

32. Jarvis, E. D. *et al.* Semi-automated assembly of high-quality diploid human reference

genomes. *Nature* **611**, 519–531 (2022).

33. Baid, G. *et al.* DeepConsensus improves the accuracy of sequences with a gap-aware

sequence transformer. *Nat Biotechnol* (2022) doi:10.1038/s41587-022-01435-7.

34. Wang, T. *et al.* The Human Pangenome Project: a global resource to map genomic

diversity. *Nature* **604**, 437–446 (2022).

35. van de Pol, M. *et al.* Detection of the ACAGG Repeat Motif in RFC1 in Two Dutch Ataxia

Families. *Mov Disord* **38**, 1555–1556 (2023).

36. Salazar, A. *et al. An AluYb8 Retrotransposon Characterises a Risk Haplotype of*

TMEM106B Associated in Neurodegeneration.

http://medrxiv.org/lookup/doi/10.1101/2023.07.16.23292721   (2023)

doi:10.1101/2023.07.16.23292721.

37. Zhang, Y. *et al. MotifScope: A Multi-Sample Motif Discovery and Visualization Tool for*

Tandem Repeats. http://biorxiv.org/lookup/doi/10.1101/2024.03.06.583591 (2024)

doi:10.1101/2024.03.06.583591.

38. Guo, M. H., Lee, W.-P., Vardarajan, B., Schellenberg, G. D. & Phillips-Cremins, J.

Polygenic burden of short tandem repeat expansions promote risk for Alzheimer's

disease. *medRxiv* 2023.11.16.23298623 (2023) doi:10.1101/2023.11.16.23298623.

39. Wang, H. *et al.* Structural Variation Detection and Association Analysis of Whole-Genome-

Sequence Data from 16,905 Alzheimer's Diseases Sequencing Project Subjects. *medRxiv*

2023.09.13.23295505 (2023) doi:10.1101/2023.09.13.23295505.

40. Lago, S. *et al.* Promoter G-quadruplexes and transcription factors cooperate to shape the

cell type-specific transcriptome. *Nat Commun* **12**, 3885 (2021).

41. Tesi, N. *et al.* Cognitively healthy centenarians are genetically protected against

Alzheimer's disease. *Alzheimer's & Dementia* **20**, 3864–3875 (2024).

42. Bonfield, J. K. *et al.* HTSlib: C library for reading/writing high-throughput sequencing data.

Gigascience **10**, giab007 (2021).

43. Marco-Sola, S. *et al.* Optimal gap-affine alignment in O (s) space. *Bioinformatics* **39**,

btad074 (2023).

44. Ye, C. & Ma, Z. (Sam). Sparc: a sparsity-based consensus algorithm for long erroneous

sequencing reads. *PeerJ* **4**, e2016 (2016).

45. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

46. van der Flier, W. M. & Scheltens, P. Amsterdam Dementia Cohort: Performing Research

to Optimize Care. *Journal of Alzheimer's Disease* **62**, 1091–1111 (2018).

47. Holstege, H. *et al.* The 100-plus Study of cognitively healthy centenarians: rationale,

design and cohort description. *European Journal of Epidemiology* (2018)

doi:10.1007/s10654-018-0451-3.

48. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation.

Nature **526**, 68–74 (2015).

49. Hon, J., Martínek, T., Zendulka, J. & Lexa, M. pqsfinder: an exhaustive and imperfection-

tolerant search tool for potential quadruplex-forming sequences in R. *Bioinformatics* **33**,

3373–3379 (2017).

50. Dreves, M. A. E. *et al.* Rationale and design of the ABOARD project (A Personalized

Medicine Approach for Alzheimer's Disease). *Alzheimers Dement (N Y)* **9**, e12401 (2023).

781