TITLE:
Single-animal, single-tube RNA extraction for comparison of relative transcript levels via qRT-PCR in the tardigrade *Hypsibius exemplaris.*

AUTHORS AND AFFILIATIONS:
Molly J. Kirk[1]*, Chaoming Xu[1]* Jonathan Paules[1] , and Joel H. Rothman[1,2]
[1] Department of Molecular Cellular and Developmental Biology University of California, Santa Barbara, Santa Barbara, CA.
[2] Neuroscience Research Institute, University of California, Santa Barbara, Santa Barbara, CA.
mollyjanekirk@ucsb.edu
joelrothman@ucsb.edu
chaomingxu@ucsb.edu
jonathan_paules@ucsb.edu

KEYWORDS:
Single-tardigrade RNA extraction, Single-tube, qRT-PCR, Heat-shock

SUMMARY:
This work presents a rapid RNA extraction and transcript level comparison method for analyzing gene expression in the tardigrade *Hypsibius exemplaris.* Using physical lysis, this high-throughput method requires a single tardigrade as the starting material and results in robust production of cDNA for quantitative Reverse Transcription Polymerase Chain Reaction (qRT-PCR).

ABSTRACT:
The tardigrade *Hypsibius exemplaris* is an emerging model organism renowned for its ability to survive environmental extremes. To explore the molecular mechanisms and genetic basis of such extremotolerance, many studies rely on RNA-sequencing (RNA-seq), which can be performed on populations ranging from large cohorts to individual animals. Reverse Transcription Polymerase Chain Reaction (RT-PCR) and RNA interference (RNAi) are subsequently used to confirm RNA-seq findings and assess the genetic requirements for candidate genes, respectively. Such studies require an efficient, accurate, and affordable method for RNA extraction and measurement of relative transcript levels by quantitative RT-PCR (qRT-PCR). This work presents an efficient single-tardigrade, single-tube RNA extraction method (STST) that not only reliably isolates RNA from individual tardigrades but also reduces the required time and cost for each extraction. This RNA extraction method yields quantities of cDNA that can be used to amplify and detect multiple transcripts by quantitative PCR (qRT-PCR). The method is validated by analyzing dynamic changes in the expression of genes encoding two heat-shock-regulated proteins, Heat-Shock Protein 70 $\beta2$ (HSP70 $\beta2$) and Heat-Shock Protein 90α (HSP90α), making it possible to assess their relative expression levels in heat-exposed individuals using qRT-PCR. STST effectively complements existing bulk and single tardigrade RNA extraction methods, permitting rapid and affordable examination of individual tardigrade transcriptional levels by qRT-PCR.

INTRODUCTION:

Tardigrades are small multicellular animals renowned for their ability to survive
extreme conditions that are lethal to most other forms of life[1]. For example, these
animals can survive nearly 1000-times the dose of ionizing radiation that is lethal to
humans[2–10], nearly complete desiccation[11–15], freezing in the absence of added
cryoprotectants,[16–18] and, in their desiccated state, even the vacuum of space[19, 20].
Owing to their unique capacity for survival in extreme environments, these animals have
become foundational models for understanding extremotolerance in complex,
multicellular organisms[1, 21–23].
Stable genetic manipulation of these remarkable animals, including transgenesis
and germline gene modification, has remained elusive until recently[24, 25]. As such, most
experiments to reveal molecular mechanisms of extremotolerance are performed
through transcriptional profiling via RNA sequencing. Many valuable and informative
RNA sequencing data sets exist for tardigrades under various extreme conditions,
ranging from radiation[8, 9, 26–28], heat stress[29], freezing stress [12, 17, 29], and desiccation[27,
30–33]. Some of these studies have utilized bulk RNA extraction and purification methods
to illuminate our molecular understanding of extremotolerance. However, bulk extraction
of RNA transcripts from many animals prevents analysis of variation in gene expression
between individuals, thus missing the potential richness of more refined data sets.
Importantly, these studies often analyze heterogeneous populations of animals that
include both animals that survive environmental stressors and those that do not. As
such, these studies are confounded by averaging expression data from multiple and
potentially dramatically different response states. To address this issue, Arakawa et al.,
2016[34] developed an elegant low-input RNA-seq pipeline that applies an RNA extraction
kit followed by a linear PCR amplification step using single[34–36] or multiple[26–28, 30, 37, 38]
animals as input. These studies have been foundational to our understanding of
tardigrade extremotolerance[22]. Interestingly, this protocol has also been applied to qRT-
PCR using seven animals as starting material[39].
In most model organisms, having identified potential targets via RNA-seq, qRT-
PCR is then performed to confirm transcriptional changes identified by RNA-seq and
assess the expression time course of candidate genes in a high-resolution manner. To
test the function of identified genes, such studies are often followed by RNAi-mediated
knockdown of molecular targets[40, 41] and analysis of extremotolerant capacity[12, 42]. The
efficacy of each RNAi knockdown is typically confirmed by qRT-PCR by directly
monitoring the decrease in transcript abundance. However, RNAi is a labor-intensive
process in tardigrades as each dsRNA must be delivered via manual microinjection of
individuals[40, 41]. Owing to the low throughput nature of this strategy, a rapid, low-cost
RNA extraction method adapted for qRT-PCR from single animals would be highly
valuable for tardigrade research. Although previous methods have been developed to
extract RNA from single tardigrades, these protocols have not combined their extraction
with qRT-PCR, instead relying on optical density-based methods[12, 41, 42]. Motivated by
these challenges, we sought to develop a protocol that reliably yields RNA in quantity
and quality that can be used for qRT-PCR from single *H. exemplaris*.
Adapted from a single-animal RNA extraction protocol developed for
Caenorhabditis elegans[20], STST is optimized for *H. exemplaris*. The extraction method
consists of six rapid freeze-thaw steps, physically disrupting the cuticle, allowing RNA
extraction and subsequent cDNA synthesis. The STST method decreases extraction

time by more than 24-fold compared to bulk RNA extraction methods, as described by
Boothby, 2018[43], and by 30% compared to single tardigrade RNA extraction kits, as
described by Arakawa et al., 2016[44]. Further, the number of sample-experimenter
interactions is decreased from 5 to only 1 compared to RNA extraction kit preparations,
thus reducing the risk of contamination by exogenous ribonucleases. When querying for
highly expressed genes, the STST method produces sufficient cDNA for 25 quantitative
RT-PCR reactions per single tardigrade, requiring only 1 μL of the total 25 μL cDNA
volume per reaction. However, template concentrations need to be empirically
determined for lower abundance transcripts.
We evaluated the efficacy of the STST method for analyzing dynamic changes in
gene expression by investigating the differential expression of the genes encoding heat-
shock protein-90α (HSP90α) and heat-shock protein 70β2 (HSP70β2) in response to
short-term heat-shock at 35°C for 20 minutes. Both HSP70β2 and HSP90α in most
eukaryotic organisms are rapidly upregulated following short-term heat-shock exposure
(20 minutes)[45]. Analysis in *H. exemplaris* revealed that both the HSP70β2 and HSP90α-
encoding RNAs extracted from single heat-treated tardigrades showed statistically
significant increases in expression following short-term heat exposure. These findings
demonstrate that the STST protocol can be used to analyze dynamic changes in gene
expression in individual animals over time.
The STST extraction method should complement existing experimental methods
such as RNA-seq by facilitating rapid and inexpensive RNA extraction and subsequent
comparison of transcript levels by qRT-PCR. This method will also be valuable for
assessing the efficiency and penetrance of RNAi in manually injected individuals more
quantitatively than optical density alone. Finally, owing to their similar cuticular
structures and physical characteristics, it is likely that this method will also be effective
for analyzing gene expression in other tardigrade species[46].
119
120　**PROTOCOL:**
121　[Place **Figure 1** here]
122
123　For detailed tardigrade and algal culturing procedures, refer to McNuff et al. 2018[47–49].
124
125　**1. Sterilization of Spring Water**
126　　1.1. Pour 2 L of spring water from a 5-gallon water jug (see Materials Section for
127　　　　Specifics) into a 2 L autoclave-safe glass bottle.
128
129　　1.2. Place the cap on the autoclave-safe bottle and seal with a small amount of
130　　　　autoclave tape. Do not tighten the bottle; just place the cap on top.
131
132　　1.3. Autoclave the spring water for 50 minutes on a wet cycle with no drying step.
133
134　　1.4. Allow the water to come to room temperature and seal the cap firmly before
135　　　　storing it at room temperature.
136
2. Glass micropipette pulling (with a pipette puller)

2.1. Secure a glass micropipette (O.D. 1 mm, I.D. 0.58 mm, Length 10 cm) on a
micropipette puller. Avoid contact with the heating filament, as this will alter the
pipette shape and damage the filament. The pulling of the pipette will need to be
determined empirically for each filament and pipette puller. However, to serve as
a starting point for optimization, use 78°C and a single pull step of pull weight of
182.2 grams.
144
145　2.2. Allow the filament to heat and gravity to separate the glass micropipette into two
146　　　 glass micropipettes with sharp points **(Figure 1b)**.
147
148　2.3. Store these pulled glass micropipettes in a closed 100 mm petri dish with wax or
149　　　 clay to hold them in place and prevent the sharp tips from breaking.
150
151　3. **Glass micropipette pulling (without a pipette puller)**
152　3.1. Light a Bunsen burner or other controlled flame source on a low setting.
153
154　3.2. Take a glass micropipette with one end in each hand.
155
156　3.3. Hold the center of the glass micropipette over the flame until the glass begins to
157　　　 melt. Then, rapidly pull the two ends apart. This will create two very delicate sharp
158　　　 tips.
159
160　3.4. Lightly break the tip with a pair of sterile fine forceps.
161
162　3.5. Store these pulled glass micropipettes in a closed 100 mm petri dish with wax or
163　　　 clay to hold them in place and prevent the sharp tips from breaking.
164
165　**4. RNA extraction**
166　4.1. Obtain 0.5 L of liquid nitrogen in a cryo-safe container.
167
168　　　 *CAUTION: Liquid nitrogen is cryogenic and may cause burns if exposed to skin*
169　　　 *or eyes. When handling, use protective clothing, splash goggles, nitrile gloves,*
170　　　 *cryo-gloves, a lab coat, and closed-toed shoes. Ascertain that the container is*
171　　　 *liquid nitrogen safe before transporting the liquid. Using an ethanol-dry ice bath*
172　　　 *for this step may also be possible.*
173
174　4.2. Make cDNA synthesis master mix: a 10 µL solution containing 1 µL of random
175　　　 hexamer primer, 2 µL of DNase, 4 µL of 5x RT Buffer, 1 µL Enzyme Mix, 1 µL of
176　　　 H2O, and 1 µL of 10 mM dNTPs. Store this solution on ice.
177
178　4.3. Prepare Tardigrade Lysis Buffer (5 mM Tris (pH=8), 0.5% (v/v) Detergent 1, 0.5%
179　　　 (v/v) Detergent 2, 0.25mM EDTA in sterile nuclease-free Water). This solution can
180　　　 be stored on the bench top for 6 months. However, maintain sterility and avoid
181　　　 potential RNAse-contaminating sources.
182
183　4.4. Aliquot enough lysis buffer for extractions (2 µL/ tardigrade).

4.5. Add RNAse inhibitor to the Tardigrade Lysis Buffer solution to a final concentration of 4 units/µL (U/µL).

4.6. Vortex and spin down the solution at room temperature on a bench-top centrifuge at a speed of 2000 xg for 5 seconds before storing the solution on ice.

4.7. Remove as many tardigrades as needed for your experiment from a culture using a sterile filter-tipped P1000 pipette and place them in a sterile 35-mm Petri dish. Any number of tardigrades may be processed in this way. Usually, three tardigrades per condition are processed for extraction.

4.8. Wash the tardigrades three times, using 1 mL of autoclaved sterile spring water and a sterile filter-tipped P1000 pipette. Slowly pipetting them up and down helps to remove algal contaminants.

4.9. Using a dissecting microscope at 25x to 50x magnification, transfer a single tardigrade from this washed culture to a new sterile 35-mm petri dish using a sterile filter-tipped P10 pipette.

4.10. Use a sterile filter-tipped P200 pipette to wash the single tardigrade in 100 µL of sterile nuclease-free water. This wash step is used to further remove contaminants, including ribonucleases.

4.11. Transfer the washed tardigrade to the bottom of a clean, sterile PCR tube in 1-2 µl of sterile nuclease-free water using a sterile filter-tipped P10 pipette, carefully ensuring the tardigrade is not stuck to the side of the tip.

4.12. Visualize the tardigrade under a dissecting microscope at 25x magnification.

4.13. To facilitate water removal of water, break the tip of the pulled glass micropipette lightly outside of the tube. The bore should be big enough to pull up the water but not the tardigrade.

4.14. Using the capillary action of a pulled glass micropipette, remove water until the animal is surrounded by a small bubble of water approximately two tardigrade lengths in diameter.

4.15. Monitor the water removal process via the dissecting scope to ensure the water level is appropriate and the tardigrade remains hydrated. **Figure 1c** offers an example of how much water to remove.

*NOTE: This is a critical step. A small bubble of water will surround the tardigrade to prevent it from drying out, but as much excess water as possible should be removed to prevent dilution of the lysis buffer. For an example of the remaining water levels, please refer to **Figure 1c**.*

4.16. Immediately after removing the water, add 2 μL of Tardigrade Lysis Buffer to the bottom of the tube, briefly vortex, and centrifuge the tube at room temperature for 5 seconds at 2000 x g on a tabletop centrifuge.

4.17. Immediately place the samples containing the tardigrades into a PCR tube rack and ensure that they are held tight by the rack.

4.18. Grip the rack using a pair of long coarse forceps and gently dip the rack containing the samples into the liquid nitrogen until fully frozen. (**Figure 1d**).

4.19. Remove the rack from the liquid nitrogen and immediately place it on ice. Allow the sample to thaw (this should take 45 seconds to 1 minute total). Monitor the sample every 15 seconds by removing it from the ice and visibly inspecting it. Once the sample is visibly transparent, move on to the next step.

4.20. Repeat steps 4.18- 4.19 five more times. A total of six freeze-thaw cycles are required for maximal lysis and extraction. (**Figure 2a,b**)

4.21. Once the freeze-thaw is complete, place samples on ice and *immediately* progress to the next step. Samples should not be frozen at this point for storage as this will diminish available RNA for cDNA preparation.

5. cDNA Synthesis

5.1. Add 2 μL of cDNA synthesis master mix to the PCR tube containing tardigrade lysate. Briefly flick the tube and spin it down at room temperature at 2000 x g for 5 seconds with a tabletop centrifuge before replacing the samples on ice.

5.2. Place the samples in a thermocycler and incubate at 25°C for 10 minutes to anneal primers, at 55°C for 30 minutes to perform reverse transcription, and finally heat inactivate enzymes at 85°C for 5 minutes.

5.3. After the incubation, immediately place the tube on ice and dilute the sample to a total volume of 25 μL by adding 21 μL of sterile nuclease-free water. For low-copy number transcripts, this dilution step can be altered as determined empirically.

6. qPCR
6.1. The annealing temperature of the primer set should be determined using total RNA prepared from larger amounts of tardigrades, for example, the bulk extraction method presented in Boothby, 2018[43].

6.2. A PCR temperature gradient should be run to determine the optimal annealing temperature before running qRT-PCR (for all PCR settings used in this protocol, refer to **Tables 1 and 2.)**

276   6.3. Thaw one tube of indicator dye super mix on ice and isolate from light. Place a 96-
277   well qPCR plate on ice and place 5 µL super mix, 2µL of water, 1 µL of each
278   primer (10 µM), and 1µL of cDNA product in the number of desired wells.
279
280   6.4. Seal the PCR plate with plate seal and run the qRT-PCR using an annealing
281   temperature appropriate for the primer set. (For all qRT-PCR settings used in this
282   paper, refer to **Table 3.)**
283

284 **7. Quantification and Results Interpretation**
285   7.1. Results are compared quantitatively to one or more control housekeeping genes,
286   whose expression is expected to be constant over the imposed conditions. For
287   this study, the actin gene was used.
288

289   7.2. The C_t-values or cycle threshold for each well are obtained and compared to the
290   C_t values of the control housekeeping gene reactions. The fold change in gene
291   expression is calculated using the following equation:
292

$$\Delta C_t = C_t(gene\ of\ interest) - C_t(housekeeping\ gene)$$
$$\Delta\Delta C_t = \Delta C_t(Sample) - \Delta C_t(Control\ average)$$
$$fold\ gene\ expression = 2^{-(\Delta\Delta C_t)}$$

294   Fold gene expression is plotted for each transcript and tardigrade as a $2^{-(\Delta\Delta C_t)}$[50].
295

296   7.3. To obtain a rough estimate of the transcript number from the C_t-value, the
297   following equation was used:
298
299

$$N = 10x2^{(35-Ct)}$$

300
301   Where N is the number of transcripts, and 2 is the assumed PCR efficiency or
302   the fold increase in fluorescence per cycle of PCR[50].
303

RESULTS:
Development and optimization of single-tardigrade RNA extraction.
Adapting the protocol from Ly et al., 2015[45] for RNA extraction in tardigrades, the
STST system is optimized to maximize the quantity and quality of the preparation
(**Figure 1a**). RT-PCR was performed for actin transcripts, quantifying transcript yield by
amplifying a 527 bp region spanning exons 1 and 2. (Sequences for these primers can
be found in **Table 1)**. The optical density of the expected actin band was quantified as
fluorescence intensity with ImageJ/FIJI. All regions of interest quantified were of equal
areal size across each gel.
First, methods of mechanical lysis required for disrupting the cuticle of the
tardigrade were assessed. In contrast to a single-animal RNA extraction protocol
described for *C. elegans*[45] which reported that Proteinase K lysis and a heat-shock were
sufficient to lyse the animals, tardigrade extraction required a minimum of six freeze-
thaw cycles to achieve consistent and robust RNA extraction (**Figure 2a, Figure S1**).

These samples were vortexed at 2700 rotations per minute for 15 seconds between
each freeze-thaw cycle to promote cell lysis. To minimize time and potential
degradation, the freeze-thaw cycle number was kept at this minimal value throughout
the remainder of the preparations. At lower numbers of freeze-thaw cycles (1-5), higher
levels of genomic DNA (gDNA) products, as evidenced by the appearance of slower-
migrating bands than expected for the actin cDNA, were found. This suggests that the
freeze-thaw process prior to dsDNA removal by DNase is required for the complete
removal of gDNA. The lower number of freeze-thaw cycles may decrease the probability
that all nuclei are effectively permeabilized, thus preventing dsDNase from degrading all
gDNA during the 25°C incubation period. At higher temperatures, during the reverse
transcription and heat inactivation steps, the remaining nuclear DNA may be released
and serves as a template during subsequent PCR. (**Figure 2a, Figure S1**). Ultimately,
the consistency of extraction success and the yield, as measured by optical density,
increased with higher numbers of freeze-thaw cycles, with six cycles showing a three-
fold increase in cDNA yield compared to no freeze-thaw cycling (**Figure 2b**). [Place
Figure 2 Here]
Next, we tested whether the near-complete removal of excess water carried over
from the sample transfer was essential to achieve consistent lysis of single tardigrades.
The STST protocol uses a minimal volume of lysis buffer to maximize RNA
concentration. We were concerned that excess water might dilute the detergent and
EDTA in the lysis buffer and thus interfere with reliable lysis. **Figure 2c and Figure S2**
show the results of triplicate individual tardigrade extractions in the presence and
absence of residual water from the tardigrade transfer. These data indicate that
removing excess water is critical to success, as product was not observed in samples
containing excess water.
We sought to determine whether the order of lysis is important for achieving
robust extraction of RNA and found that performing the freeze-thaw step before the
Proteinase K and enzymatic lysis resulted in little or no detectable RNA by RT-PCR
(lanes 2-4 of **Figure 2d**). However, PCR products were readily obtained when chemical
lysis was performed before the freeze-thaw step (lanes 5-7 of **Figure 2d**). This suggests
either that Proteinase K digestion of the cuticle is required before mechanical lysis or
that prior freeze-thaw lysis extracts the RNA, leaving it exposed to endogenous RNAse
activity throughout the proteinase K treatment.
To assess whether proteinase K digestion is required for the isolation of RNA
from single tardigrades, RNA quality was assessed via high-sensitivity tape station.
Animals that experienced vortexing after proteinase K treatment showed high RNA
fractionation, while freeze-thaw cycling after proteinase K treatment resulted in
increased yet inconsistent RNA integrity scores (RIN) (**Figure 2e).** Intriguingly, RNA
extraction in the absence of the proteinase K and vortexing steps increased consistency
and quality significantly, suggesting that the proteinase K treatment is neither required
nor helpful for RNA extraction from *H. exemplaris* (**Figure 2e).** Eliminating this treatment
significantly reduced the time required for our protocol from 45 minutes with proteinase
K treatment to 7 minutes in our improved STST protocol. Further, increasing the RNAse
inhibitor concentration from 4U/μL to 8U/μl or 16 U/μL resulted in incomplete freezing of
the animals, yielding inconsistent results at 8U/μL and a complete lack of freezing and
subsequent lysis at 16 U/μL (**Figure 2f**). This effect is most likely attributable to the

RNAse storage solution containing 50% glycerol, which acts as a cryoprotectant. Also,
we found that thawing on ice resulted in moderately increased RIN scores (**Figure 2g).**
Finally, to assess the total yield of STST preparations, RNA quantity per µL was
measured via High Sensitivity RNA Qubit. This analysis revealed a yield of 7.11 +/- 1.43
ng/µL (**Figure 2h)**, suggesting that the STST preparation yields 14.24 +/- 2.88 ng per
single *Hypsibius exemplaris* animal (containing ~1400 cells) (**Figure 2i**). This result is
consistent with the yields observed with *C. elegans* adult (~3000 cells, including germ
cells), which yield ~ 35 ng per animal[45].
372
Comparison to existing tardigrade RNA extraction protocols
In developing the STST RNA extraction protocol, we aimed to limit the time
required to perform the technique, the number of sample-experimenter interactions, the
cost, and the number of animals required for robust RNA extraction. **Figures 3a and 3b**
depict a schematic contrasting a previously described single-tardigrade RNA extraction
protocol that has been used for RNA sequencing following linear PCR amplification
(**Figure 3a**) and the single-tube, single-tardigrade RNA extraction method (**Figure 3b**),
highlighting each method in terms of these four parameters. Previously, single-
tardigrade RNA extraction protocols required approximately ten minutes, three tubes,
and five experimenter interactions before cDNA synthesis; however, these methods
have not yet been assessed for qRT-PCR applications on single animals. Current
published RNA extraction protocols for qRT-PCR assessment of transcriptional changes
use a minimum of seven tardigrades[24]. The STST method described here decreases
the time to seven minutes with only one sample-experimenter interaction and a single
tube. Removing sample-experimenter interactions reduces the potential for RNAse
contamination, thus increasing the potential for high-quality extractions. [24]
To compare the efficacy of STST extractions with the RNA extraction kit, single-
tardigrade extractions were performed using both methods, and transcript numbers
were quantitatively assessed using qPCR. Amplifying a 150 bp region of the
housekeeping gene encoding actin, we used the C_t value, i.e., the cycle number
required for the fluorescence signal to be detected above the background, to indicate
cDNA quantity. The average C_t value of single tardigrades processed with the STST
protocol was 27.86 +/- 1.268 **(Figure 3c)**, while those extracted with the RNA extraction
kit were 37.67+/- 3.311 **(Figure 3c).** While both protocols yield a signal above
background fluorescence obtained from control qPCRs run without template cDNA, only
the STST protocol was statistically significant compared to background levels. This
finding indicates that although RNA was present in the RNA extraction kit samples, the
kit did not produce sufficient quantities of RNA to be consistently detected. Using these
C_t values, we were able to estimate the transcript number per µL of cDNA: STST
yielded ~1760+/-954.2 actin transcripts per µL, while the RNA Kit-based method yielded
~ 7.5+/- 12.29 actin transcripts per µL **(Figure 3d),** suggesting that as a single-
tardigrade extraction method, the STST system is > 200-fold more efficient. When
quantified, the STST protocol produced significantly more transcripts than the RNA
extraction kit, which was again statistically indistinguishable from background
measurements **(Figure 3d).** This suggests that the current RNA kit-based method for
single tardigrades does not sufficiently extract RNA or loses significant amounts of RNA
throughout the purification process, resulting in undetectable quantities of cDNA

downstream. To visualize this effect, qPCR samples were run on an analytical gel,
confirming minimal amplification of RNA kit-extracted products (**Figure 3e,** lanes 2-4)
and robust amplification in the STST-extracted samples (**Figure 3e,** lanes 5-8).
Finally, we compared the cost of RNA extraction and cDNA synthesis using the
RNA extraction kit with the STST method (**Table 4**). As the elution volume for the RNA
kit was three times that of the STST protocol, it requires three times the cDNA synthesis
material, accounting for the higher cost in cDNA synthesis for each whole tardigrade
extract before samples were diluted to equal volumes of 25μL. Further, as we had
previously determined (**Figure 3 c and d**) that a linear amplification step would be
required to yield quantities above background C_t values, we have included that step in
our cost estimates as directly described by Arakawa et al. 2016[44]; however, there may
be a more economical way to perform linear amplification of which we are unaware.
[Place Figure 3 here]
423
Application of the single-tube single-tardigrade extraction protocol for quantifying heat-
shock responses in H. exemplaris.
We next sought to explore whether this protocol can be applied to effectively
monitor transcriptional changes in *H. exemplaris*, turning to the canonical heat-shock
response pathway, which results in the upregulation of both HSP70β2 and HSP90α
(**Figure 4a**) following brief exposure to high temperatures[45]. Tardigrades in their active
state are highly sensitive to increases in temperature[29]; thus, we selected 35°C for 20
minutes as the heat-shock, as we found that this resulted in minimal lethality (**Figure
4b**). We observed that the response follows a similar pattern to that seen in other
organisms[45]. We first exposed tardigrades to either rearing temperature (23°C) or high
heat (35°C) for 20 minutes (**Figure 4c**). We then extracted RNA from individual animals,
and generated cDNA using the STST method by harvesting immediately or at 1 hour, 2
hours, 4 hours, and 6 hours after exposure to the heat-shock. We subsequently
quantified transcripts for HSP70β2 (**Figure 4d**) and HSP90α (**Figure 4e**) using qRT-
PCR. Significant upregulation of HSP70β2 (~11-fold) and HSP90α(~4-fold) was
observed after 1 hour, and expression slowly returned to baseline over the next several
hours. These results follow the expected patterns of heat-shock responses seen in *C.
elegans*[45].
Finally, to assess the quality of STST as an extraction method for qRT-PCR, we
compiled all actin C_t values across all the samples run for panels **4d** and **e.** The average
actin C_t values for an actin 150 bp amplicon and an actin 527 bp amplicon were not
statistically different (**Figure 4f**). Furthermore, the comparison of variance (standard
deviation) across C_t values derived from triplicates obtained from each individual extract
were consistently below the average quality cut-off of 0.5 cycles per individual extract[50]
(**Figure 4g).**
449
450   **FIGURE AND TABLE LEGENDS:**
451
Figure 1. Single-tube pipeline for RNA extraction from a single tardigrade. a)
Scheme showing the protocol for RNA extraction from a single tardigrade, including six
freeze-thaw cycles and subsequent cDNA synthesis. Samples may subsequently be
used for RT-PCR and qRT-PCR. **b)** Image of micropipette taper used for removal of

water. Scale, 2 mm. **c)** Bright field image of a tardigrade in a small volume of water
(dotted line). Removal of most water to the extent shown is required for successful
extraction and prevents dilution of lysis buffer. Scale, 50 μm. **d)** Image showing
immersion of samples in liquid nitrogen using long forceps to rapidly freeze-thaw the
samples safely. Some of the content was created in BioRender. Kirk, M. (2022)
BioRender.com/d93s511
462
Figure 2: Optimization of single-tardigrade RT-PCR using actin cDNA as a marker
for extraction quantity. a) Representative gel depicting results from single-tardigrade
RT-PCR extracted using 4 (lane 2-4), 5 (lane 5-7), and 6 (lane 8-10) freeze-thaw cycles
to enhance lysis after proteinase K treatment and heat-shock. Full gel-containing
samples from 1, 2, and 3 freeze-thaw cycles are shown in **Figure S1**. **b)** optical density
quantification of ethidium bromide staining of actin RT-PCR across various freeze-thaw
cycle numbers. Data represent optical density values of bands from three individual
extractions and PCR amplification of actin via RT-PCR per condition. One way ANOVA,
with Tukey's multiple comparisons post hoc 0 vs. 6, p= 0.020 and 3 vs. 6 p=0.022, error
bars represent S.D. (Standard deviation) **c)** Representative gel showing the effect
residual spring water removal from isolated tardigrades prior to the addition of lysis
buffer. Samples containing water (lanes 2-4) and samples where the water was
removed (lanes 5-*7*). **d)** Representative gel showing the effect of lysis order, with
freeze-thaw performed prior to chemical lysis (lanes 2-4) or chemical lysis with
proteinase K prior to freeze-thaw lysis (lanes 5-7). **e)** RNA integrity scores reported by
Agilent High Sensitivity Tape station of single tardigrade extracts animals in the
absence of freeze-thaw, in the absence of vortexing, and utilizing only freeze-thaws
without proteinase k digestion or vortexing. Each data point represents the RIN score
from one singular RNA extraction. One way ANOVA, with Tukey's multiple comparisons
post hoc, * p=0.036. **f)** RIN scores reported from single tardigrade extracts in the
presence of 4U/μL, 8U/μl and 16U/μL RNAse inhibitor. Brown-Forsythe ANOVA with
Dunnett's T3 multiple comparisons test, **** p=<0.0001 and *=0.0487. **h)** RIN scores
from single tardigrade extracts thawing on ice. **i)** RNA quantity in ng/μL as measured
using Qubit high sensitivity RNA kit. **j)** RNA quantity in ng per single tardigrade extract
using STST. 5 μL of PCR products were loaded per lane unless otherwise noted. All
error bars are reported S.D..
Some of the content was created in BioRender. Kirk, M. (2022) BioRender.com/d93s51
490
Figure 3. **Comparison of RNA extraction protocols.** Schematized time-course for **a)**
an existing phenol and guanidine isothiocyanate-based RNA extraction kit, requiring
~10 minutes for extraction (Scale bar is 30 minutes) and **b)** The single-tube, single-
tardigrade protocol that permits extraction in 7 minutes and requires one tube. The time
courses are drawn to scale. Scale bar is 30 minutes. **c)** Graph showing C_t values of
actin qRT-PCR reactions run in triplicate where each data point represents the average
Ct values of three technical replicates from one individual tardigrade extract. Single-
tardigrade-based extractions and the background control revealing background
fluorescence from samples run on qRT-PCR in the absence of template. One way
ANOVA, Tukey's multiple comparisons test ***, p=0.0003 and ****, p<0.0001, error bars
reflect S.D. **d)** Estimated transcripts per μL of sample derived from the C_t values

depicted in **(c)**.Kruskal-Wallis Test, Dunn's multiple comparisons test, *, p=0.0165 error bars reflect S.D. **e)** Representative gel showing actin RT-PCR from single-tardigrade cDNA samples extracted using RNA kit (lanes 2-4) and STST lanes (5-8). Gels were loaded with 6 μL of sample and 1 kb plus DNA Ladder

Figure 4. Expression of HSP-70 and HSP-90 transcripts. a) Representative gel showing amplification of HSP70 from genomic DNA (gDNA; lane 2) and cDNA (lane 3) and HSP90 from gDNA (lane 4) and cDNA (lane 5). Variance in gDNA and cDNA lengths reflects the primer design, which spans an intron. **b)** Survival of tardigrade post exposure to 23°C and 35°C for 20 minutes. Each data point represents percent survival from a cohort of six animals. T-test found no statistical difference. **c)** Conditions analyzed. **d)** Relative expression levels of HSP70β2 at the indicated time post-20-minute heat-shock at 35 °C. Kruskals-Wallis Test with Dunn's multiple comparisons analysis *, p=0.0339 (1 vs. 0)**, p=0.0043 (1 vs. 4), **, p=0.0097 (1 vs. 6). Each data point represents normalized expression levels from an individual tardigrade RNA extract averaged across three technical replicates. **e)** Relative expression levels of HSP90α at the indicated time post-20-minute 35°C heat-shock. Brown-Forsythe ANOVA with Dunnett's T3 multiple comparisons test*, p=0.0382 (1 vs. 0) *, p=0.0421 (1 vs. 4). Each data point represents normalized expression levels from an individual tardigrade RNA extract averaged across three technical replicates. **f)** Raw C_t values for actin qRT-PCR amplifying either a 150 bp or 527 bp amplicon from all samples presented in panels **d** and **e**. Each data point represents C_t from an individual tardigrade RNA extract averaged across three technical replicates; T-test found no statistical difference. **g)** Standard deviation in C_t values across technical replicates for each extract represented in panels **d** and **e**. Reactions amplified an actin amplicon of either 150 bp or 527 bp; T-test found no statistical difference. Some content was created in BioRender. Kirk, M. (2022) BioRender.com/d93s51

Table 1. Thermocycling conditions for the Actin and HSP90 PCR reaction

Table 1. PCR thermocycling protocol for non-quantitative PCR. The table describes the exact thermocycling procedure used for this study. Tm is denoted as the melting temperature for a reaction. The specific melting temperatures for each reaction can be found in **Table 2.**

Table 2. Melting temperature or Tm for each primer pair used in this study.

Table 2. Melting temperature and sequences for primers used in this study. The table shows the forward and reverse sequences of primers used for this study and their optimized melting temperature. It is important to note that HSP70 required a brief touch-down protocol consisting of 6 cycles before the amplification to ensure specificity.

Table 3. Thermocycling conditions for the qPCR

546 **Table 3. Thermocycling conditions for qPCR.** The table depicts the thermocycling
547 settings for qPCR reactions used in this study. The melting temperatures or Tm's were
548 identical to those shown in **Table 2.**
549
550 **Table 4: Cost breakdown for RNA extraction.**
551
552 **Table 4: Cost breakdown for RNA extraction.** The table describes the cost
553 breakdown for each RNA extraction method per single animal. Prices were obtained
554 from the vendor website on September 30th, 2024, and are subject to change.
555
DISCUSSION:
This study presents an efficient method for the extraction of RNA for single-
tardigrade qRT-PCR. Directly comparing the STST methodology to an existing single
tardigrade RNA extraction kit revealed that STST RNA extraction yields >200-fold
higher amounts of actin RNA transcripts, reduces the cost to less than one dollar per
sample, and reduces the time required for extraction by 30%. To apply STST to a
relevant biological question, we assessed the short-term heat-shock response
expression profile. We found that transcripts for both HSP70β2 and HSP90α, as
expected, were strongly upregulated 1 hour after heat exposure.
Although the STST protocol is a substantial improvement over previous methods,
several limitations present opportunities for improvement and further assessment. First,
we have not evaluated the ability of STST to detect transcripts expressed at very low
levels. Troubleshooting for appropriate template concentration may be required for
transcripts which are expressed at low levels. Secondly, while the STST method
showed an ~ 80% success rate, defined as the percent of extracts resulting in an actin
C_t values less than 31, there were occasions when the method failed to obtain usable
quantities of cDNA, perhaps because of RNAse activity or insufficient lysis. Optimizing
the inhibition of RNAse activity might help increase the success rate. Finally, RNA
quality could be improved in this method. As in this study, measures can be taken to
circumvent the effects of degradation and subsequent fragmentation of the RNA, as well
as to mitigate its impact on qRT-PCR results. Improvements in RNA quality would
further enhance the rapid assessment of RNA transcriptional changes, the analysis of
RNAi knockdown efficacy, and the ability to evaluate variance across populations of
many individual animals. It is important to note that all protocols for RNA extraction in
tardigrades require some form of mechanical lysis. Traditional phenol and guanidine
isothiocyanate protocols require three freeze-thaw cycles[43], while the RNA extraction kit
requires manual rupturing with a pipette[36, 44] Ultimately, different methods of mechanical
lysis may improve quality in the future.
We identified several critically important steps in the STST method. First,
removing most of the water from the tube prior to adding lysis buffer is critical: we leave
a bubble approximately two tardigrade lengths in diameter surrounding the animal.
Though we have not measured this minute volume, removing as much water as
possible will improve consistency and overall success (**Figure 1c**). Second, the
immediate transfer of the samples into a cDNA preparation reaction is important. Our
preparation does not include any RNA purification, so it is imperative that one does not
pause the procedure after extraction. It is possible that extracts could be successfully

stored at -80 °C; however, we have not assessed the effect of such storage on qRT-PCR quality. Third, adhering to the guidelines for performing qRT-PCR on mildly degraded samples is vital , i.e. analyzing short amplicons of < 200 bp using multiple housekeeping genes to normalize data whenever possible[51, 52].

Overall, our findings represent an important step in the growing repertoire of molecular tools available in tardigrade research. This method complements existing RNA extraction methods[43, 44], which have provided techniques for single tardigrade quantification of transcriptional changes via RNA-seq following a linear amplification step[43, 44]. STST allows for quantification of transcripts isolated from single tardigrades without the need for a scale-up step. As most tardigrade responses to extreme conditions, such as those triggered by desiccation, result in ~75-80% survival rates in *H. exemplaris*; bulk samples contain mixed populations of tardigrades, including both survivors and non-survivors. Such a mixed population has the potential to confound the analysis of transcriptional changes that occur during response and recovery from extreme stresses. In bypassing the need for bulk preparation, STST and the tardigrade RNA-seq method[36, 44] avoid the confounding problems inherent in a mixed population and allow for the full dynamic range of a response to be quantified. We envision that STST can be used not only to quantify the range of transcriptional responses and assess the efficacy of RNAi knockdowns, but also be used in many other applications in the future.

DATA AVAILABILITY:
All raw analytical gel data has been incorporated into the supplemental data of this manuscript. Transcript sequences were identified via Ensembl search at: https://metazoa.ensembl.org/Hypsibius_exemplaris_gca002082055v1/Info/Index?db=core HSP70β2 (BV898_04401), HSP90α (BV898_50798) and Actin(BV898_02877).

ACKNOWLEDGMENTS:
We would like to acknowledge the NIH Ruth Kirschstein Fellowship # 5F32AG081056-02 and the Errett Fisher Post-Doctoral Fellowship, which supported Dr. Molly J. Kirk, the Crowe Family Fellowship, which supported Chaoming Xu, and a University of California, Santa Barbara Academic Senate Grant, and NIH grants # R01GM143771 and #2R01HD081266, which supported these research efforts. The authors also acknowledge the use of the Biological Nanostructures Laboratory within the California NanoSystems Institute, supported by the University of California, Santa Barbara and the University of California, Office of the President.

DISCLOSURES : The authors declare no conflicts of interest to disclose.

REFERENCES:
1.   Møbjerg, N., Neves, R.C. New insights into survival strategies of tardigrades. *Comparative Biochemistry and Physiology -Part A : Molecular and Integrative Physiology*. 254 (2021).
2.   Jönsson, K.I., Harms-Ringdahl, M., Torudd, J. Radiation tolerance in the eutardigrade Richtersius coronifer. *Int. J. Radiat. Biol.* 81 (9), 649–656 (2005).

3. Horikawa, D.D. *et al.* Radiation tolerance in the tardigrade Milnesium tardigradum.
Int. J. Radiat. Biol. 82 (12), 843–848 (2006).
4. Bruckbauer, S.T., Cox, M.M. Experimental evolution of extremophile resistance to
ionizing radiation. *Trends in Genetics.* 37 (9), 830–845 (2021).
5. Jönsson, K.I., Hygum, T.L., Andersen, K.N., Clausen, L.K.B., Møbjerg, N.
Tolerance to Gamma Radiation in the Marine Heterotardigrade, Echiniscoides
sigismundi. *PLoS One.* 11 (12), e0168884 (2016).
6. Jönsson, K.I. Radiation tolerance in tardigrades: Current knowledge and potential
applications in medicine. *Cancers (Basel).* 11 (9), 1333 (2019).
7. Yoshida, Y. *et al.* RNA sequencing data for gamma radiation response in the
extremotolerant   tardigrade Ramazzottius varieornatus. *Data Brief.* 36
(107111), 107111 (2021).
8. Clark-Hachtel, C.M., Hibshman, J.D., De Buysscher, T., Stair, E.R., Hicks, L.M.,
Goldstein, B. The tardigrade Hypsibius exemplaris dramatically upregulates DNA
repair   pathway genes in response to ionizing radiation. *Curr. Biol.* 34 (9), 1819-
1830.e6 (2024).
9. Anoud, M. *et al.* Comparative transcriptomics reveal a novel tardigrade specific
DNA binding protein induced in response to ionizing radiation (2024).
10. Jönsson, K.I., Schill, R.O. Induction of Hsp70 by desiccation, ionising radiation
and heat-shock in   the eutardigrade Richtersius coronifer. *Comp. Biochem.*
Physiol. B Biochem. Mol. Biol. 146 (4), 456–460 (2007).
11. Boothby, T.C. Desiccation of hypsibius exemplaris. *Cold Spring Harbor Protocols.*
2018 (11), 871–873 (2018).
12. Boothby, T.C. *et al.* Tardigrades Use Intrinsically Disordered Proteins to Survive
Desiccation. *Molecular Cell.* 65 (6), 975-984.e5 (2017).
13. Horikawa, D.D., Higashi, S. Desiccation tolerance of the tardigrade Milnesium
tardigradum collected in   Sapporo, Japan, and Bogor, Indonesia. *Zoolog. Sci.*
21 (8), 813–816 (2004).
14. Halberg, K.A., Jørgensen, A., Møbjerg, N. Desiccation tolerance in the tardigrade
Richtersius coronifer relies on muscle mediated structural reorganization. *PLoS*
ONE. 8 (12) (2013).
15. Sørensen-Hygum, T.L., Stuart, R.M., Jørgensen, A., Møbjerg, N. Modelling
extreme desiccation tolerance in a marine tardigrade. *Sci. Rep.* 8 (1), 11495
(2018).
16. Lyons, A.M., Roberts, K.T., Williams, C.M. Survival of tardigrades (Hypsibius
exemplaris) to subzero temperatures depends on exposure intensity, duration,
and ice-nucleation-as shown by large-scale mortality dye-based assays.
17. Møbjerg, A. *et al.* Extreme freeze-tolerance in cryophilic tardigrades relies on
controlled ice formation but does not involve significant change in transcription.
Comparative Biochemistry and Physiology Part A: Molecular & Integrative
Physiology. 271, 111245 (2022).
18. Tsujimoto, M., Imura, S., Kanda, H. Recovery and reproduction of an Antarctic
tardigrade retrieved from a moss   sample frozen for over 30 years. *Cryobiology.*
72 (1), 78–81 (2016).
19. Jönsson, K.I. Tardigrades as a potential model organism in space research.
Astrobiology. 7 (5), 757–766 (2007).

20. Jönsson, K.I., Rabbow, E., Schill, R.O., Harms-Ringdahl, M., Rettberg, P.
Tardigrades survive exposure to space in low Earth orbit. *Current biology⬚: CB*.
18 (17), R729–R731 (2008).
21. Kasianchuk, N., Rzymski, P., Kaczmarek, Ł. The biomedical potential of
tardigrade proteins: A review. *Biomedicine and Pharmacotherapy*. 158 (2023).
22. Arakawa, K. Annual Review of Animal Biosciences Examples of Extreme Survival:
Tardigrade Genomics and Molecular Anhydrobiology (2022).
23. Hvidepil, L.K.B., Møbjerg, N. New insights into osmobiosis and chemobiosis in
tardigrades. *Front. Physiol.* 14, 1274522 (2023).
24. Tanaka, S., Aoki, K., Arakawa, K. In vivo expression vector derived from
anhydrobiotic tardigrade genome enables live imaging in Eutardigrada.
Proceedings of the National Academy of Sciences. 120 (5) (2023).
25. Kondo, K., Tanaka, A., Kunieda, T. Single-step generation of homozygous knock-
out/knock-in individuals in an extremotolerant parthenogenetic tardigrade using
DIPA-CRISPR.
26. Yoshida, Y., Hirayama, A., Arakawa, K. Transcriptome analysis of the tardigrade
Hypsibius exemplaris exposed to the DNA-damaging agent
bleomycin. *bioRxiv*. 2024.02.01.578372 (2024).
27. Yoshida, Y. *et al.* Time-series transcriptomic screening of factors contributing to
the cross-tolerance to UV radiation and anhydrobiosis in tardigrades. *BMC*
Genomics. 23 (1), 405 (2022).
28. Yoshida, Y. *et al.* RNA sequencing data for gamma radiation response in the
extremotolerant tardigrade Ramazzottius varieornatus. *Data in Brief*. 36, 107111
(2021).
29. Neves, R.C., Møbjerg, A., Kodama, M., Ramos-Madrigal, J., Gilbert, M.T.P.,
Møbjerg, N. Differential expression profiling of heat stressed tardigrades reveals
major shift in the transcriptome. *Comparative Biochemistry and Physiology -Part*
A⬚: Molecular and Integrative Physiology. 267 (2022).
30. Yoshida, Y. *et al.* Comparative genomics of the tardigrades Hypsibius dujardini
and Ramazzottius varieornatus. *PLOS Biology*. 15 (7), e2002266 (2017).
31. Wang, C., Grohme, M.A., Mali, B., Schill, R.O., Frohme, M. Towards decrypting
cryptobiosis--analyzing anhydrobiosis in the tardigrade   Milnesium tardigradum
using transcriptome sequencing. *PLoS One*. 9 (3), e92663 (2014).
32. Mali, B. *et al.* Transcriptome survey of the anhydrobiotic tardigrade Milnesium
tardigradum in comparison with Hypsibius dujardini and Richtersius coronifer.
BMC Genomics. 11 (1), 168 (2010).
33. Förster, F. *et al.* Transcriptome analysis in tardigrade species reveals specific
molecular   pathways for stress adaptations. *Bioinform. Biol. Insights*. 6, 69–96
(2012).
34. Arakawa, K., Yoshida, Y., Tomita, M. Genome sequencing of a single tardigrade
Hypsibius dujardini individual. *Scientific Data*. 3 (1), 160063 (2016).
35. Arakawa, K. Transcriptome assembly of Richtersius coronifer with annotated
BLAST result against Ramazzottius varieornatus. (2019).
36. Yoshida, Y., Konno, S., Nishino, R., Murai, Y., Tomita, M., Arakawa, K. Ultralow
input genome sequencing library preparation from a single   tardigrade
specimen. *J. Vis. Exp.* (137) (2018).

729 37. Murai, Y., Yagi-Utsumi, M., Fujiwara, M., Tomita, M., Kato, K., Arakawa, K.
730   Multiomics study of a heterotardigrade, Echinisicus testudo,
731   suggests convergent evolution of anhydrobiosis-related proteins in Tardigrada.
732   *bioRxiv*. 2020.10.27.358333 (2020).
733 38. Yoshida, Y., Sugiura, K., Tomita, M., Matsumoto, M., Arakawa, K. Comparison of
734   the transcriptomes of two tardigrades with different hatching coordination. *BMC*
735   *Developmental Biology*. 19 (1), 24 (2019).
736 39. Tanaka, S., Aoki, K., Arakawa, K. In vivo expression vector derived from
737   anhydrobiotic tardigrade genome enables live imaging in Eutardigrada.
738   *Proceedings of the National Academy of Sciences*. 120 (5), e2216739120 (2023).
739 40. Tenlen, J.R. Microinjection of dsRNA in tardigrades. *Cold Spring Harb. Protoc.*
740   2018 (11), db.prot102368 (2018).
741 41. Tenlen, J.R., McCaskill, S., Goldstein, B. RNA interference can be used to disrupt
742   gene function in tardigrades. *Development Genes and Evolution*. 223 (3), 171–
743   181 (2013).
744 42. Giovannini, I., Boothby, T.C., Cesari, M., Goldstein, B., Guidetti, R., Rebecchi, L.
745   Production of reactive oxygen species and involvement of bioprotectants during
746   anhydrobiosis in the tardigrade Paramacrobiotus spatialis. *Scientific Reports*. 12
747   (1) (2022).
748 43. Boothby, T.C. Total RNA extraction from tardigrades. *Cold Spring Harbor*
749   *Protocols*. 2018 (11), 905–907 (2018).
750 44. Arakawa, K., Yoshida, Y., Tomita, M. Genome sequencing of a single tardigrade
751   Hypsibius dujardini individual. *Sci. Data*. 3 (1), 160063 (2016).
752 45. Ly, K., Reid, S.J., Snell, R.G. Rapid RNA analysis of individual Caenorhabditis
753   elegans. *MethodsX*. 2, 59–63 (2015).
754 46. Czerneková, M., Vinopal, S. The tardigrade cuticle. *Limnological Review*. 21 (3),
755   127–146 (2021).
756 47. Goldstein, B. *Hypsibius dujardini collection notes and culture protocol from Bob*
757   *McNuff*. (2007).
758 48. McNuff, R. Laboratory culture of hypsibius exemplaris. *Cold Spring Harbor*
759   *Protocols*. 2018 (11), 867–870 (2018).
760 49. Gabriel, W.N. *et al.* The tardigrade Hypsibius dujardini, a new model for studying
761   the evolution of development. *Dev. Biol.* 312 (2), 545–559 (2007).
762 50. Ruiz-Villalba, A., Ruijter, J.M., van den Hoff, M.J.B. Use and misuse of cq in qpcr
763   data analysis and reporting. *Life*. 11 (6) (2021).
764 51. Antonov, J. *et al.* Reliable gene expression measurements from degraded RNA
765   by quantitative real-time PCR depend on short amplicons and a proper
766   normalization. *Laboratory Investigation*. 85 (8), 1040–1050 (2005).
767 52. Toussaint, J. *et al.* Improvement of the clinical applicability of the genomic grade
768   index through a qRT-PCR test performed on frozen and formalin-fixed paraffin-
769   embedded tissues. *BMC Genomics*. 10, 424 (2009).
770

a.

Collect Tardigrade → Remove Excess Water → Lysis Buffer → 6x Freeze Thaw Cycles → cDNA Synthesis

Quantitative RT-PCR

RFU

Cycles

b.

c.

d.

a. Previous Work: RNA Kit

RT PCR ~ 10 min / 1 Tardigrade/3 Tubes/5 Interactions

30 min

- Isolation of Tardigrades
- RNA extraction
- Washes

b. Current Work: STST

RT PCR ~ 7 min / 1 Tardigrade/1 Tube/ 1 Interaction

30 min

c.

****
***

Background RNA Kit STST

Ct value (Actin)

d.

*

Background RNA Kit STST 2.0

Copy Number/μL

e.

RNA Kit   STST

Name of Material/ Equipment	Company	Catalog Number	Comments/Description
Corning™ PYREX™ Reusable Media Storage Bottles	Thermo Fisher Scientific	06-414-1E	Refered to as 2 L Autoclave-safe Glass Bottle
Andwin Scientific 3M LEAD FREE AUTOCLAVE TAPE 1"	Thermo Fisher Scientific	NC0802040	Refered to as Autoclave Tape
Hypsibius exemplaris Z151 Strain	Carolina	133960	Refered to as Tardigrades or *H. exemplaris*
Chloroccoccum hypnosporum .	Carolina	152091	Refered to as Algae
Direct-zol Micro Prep	Zymo Research	R2060	Refered to as RNA extraction kit
RNasin® Ribonuclease Inhibitor	Promega	N2111	Refered to as RNAse inhibitor
Maxima H Minus First Strand cDNA Synthesis Kit	Thermo Fisher Scientific	K1651	Refered to as cDNA Synthesis Master Mix
SsoAdvanced™ Universal SYBR® Green Supermix	BIO RAD	1725271	Refered to as Indicator Dye Super mix
Triton™ X-100	Fluka	93443	Refered to as Detergent 1
Tris base	Fisher Scientific	T395-500	Refered to as Tris or Tris Base
TWEEN® 20	Sigma aldrich	P1379-500	Refered to as Detergent 2
EDTA	Fisher Scientific	S311-500	Refered to as EDTA
Water - PCR/RT-PCR certified, nuclease-free	Growcells	PCPW-0500	Refered to as Sterile Nuclease Free Water
Spring water	Nestle Pure Life	44221229	Refered to as Spring Water
RNAse-Free, 8 Strip 0.2mL PCR Tubes with caps	Invitrogen	AM12230	Refered to as Sterile PCR Tube
BioRadHard-Shell® 96-Well PCR Plate	BioRad	HSS9641	Refered to as PCR Plate
PETRI DISH, PS, 35/10 MM, WITH VENTS	Grenier	627102	Refered to as 35 mm Petri dish
Thermo Scientific™ Nunc™ Petri Dishes	Thermo Fisher Scientific	08-757-099	Refered to as 100 mm Petri dish
Supertek Scientific Tirrill Burners	Thermo Fisher Scientific	S09572B	Refered to as Bunsen Burner
Dumont 5 Biology Tweezers	Fine Science Tools	11254-20	Refered to as Fine Forceps
Fisherbrand™ Economy Impact Goggles	Fisher Scientific	19-181-501	Refered to as Splash Goggles
Nitrile Gloves	Fisher Scientific	17-000-314	Refered to as Nitrile Gloves
BULWARK FR Lab Coat:	Grainger	26CF64	Refered to as Lab Coat
Tempshield™ Cryo-Gloves™	Fisher Scientific	11-394-305	Refered to as Cryo Gloves
PIPETMAN P1000, 100-1000 µL, Metal Ejector	Gilson	F144059M	Refered to as P 1000 Pipette
PIPETMAN P200, 20-200 µL, Metal Ejector	Gilson	F144058M	Refered to as P 200 Pipette
PIPETMAN P10, 1-10 µL, Metal Ejector	Gilson	F144055M	Refered to as P 10 Pipette
Stereo-Microscope System w/optics and illumination	TriTech Research	SMT1	Refered to as Dissecting Microscope
Autoclave Tape	Thermo Fisher Scientific	AB1170	Refered to as PCR Plate Seals

Item	Manufacturer	Catalog #	Reference
Pound This 4-Color Modeling Clay	American Science Surplus	96517P001	Refered to as Clay
C1000 Touch^{TM} Bio-rad Thermocycler	BioRad	1851148	Refered to as Thermocycler
C1000 Touch^{TM} Bio-rad Thermocycler with CFX^{TM} Optics Module	BioRad	1845097	Refered to as qPCR thermocycler
Glass Micropipette O.D. 1mm ID 0.58, Length 10 cm	TriTech Research	GD-1	Reffered to as glass micropipette
Narishige Dual-Stage Glass Micropipette Puller	Tritech Research	PC-10	Refered to as micropipette puller
Filament for pippette Puller	Tritech Research	PC-10H	Refered to as Filament
Table Top Centrifuge	Qualitron	DW-41-115-NEW	Refered to as Table Top Centrifuge
Liquid Nitrogen Dewar 1L	Agar Scientific	AGB7475	Refered to as Cryo-safe container
Daigger & Company Vortex-Genie 2 Laboratory Mixer	Thermo Fisher Scientific	3030A	Refered to as Vortexer
4 Star Straight Strong Medium Point Tweezer	Excelta	00-SA-DC	Refered to as Long forceps
96-Well PCR Rack with Lid Assorted, 5 Racks/Unit	Genesee Scientific	27-202A	Refered to as PCR Rack
Prism v10.0	GraphPad	N/A	Refered to a Prism
Benchling v8	Benchling	N/A	Refered to as Benchling
FIJI v 2.14.0/1.54f	ImageJ,	N/A	Refered to as FIJI/ImageJ
1000µl Premium Pipet Tips, Low Binding, Racked, Sterile	Genesee Scientific	23-165RS	Refered to as Sterile Filter-Tipped P 1000 Pipette Tips
200µl Premium Barrier Tips Low Binding, Racked, Sterile	Genesee Scientific	23-412	Refered to as Sterile Filter-Tipped P 200 Pipette Tips
10µl Premium Barrier Tips Low Binding, Racked, Sterile	Genesee Scientific	23-401	Refered to as Sterile Filter-Tipped P 10 Pipette Tips

Table 1.

Step	Temp. (°C)	Duration	No. of cycles
Initial denaturation	95	60 s	1
Denaturation	95	15 s	
Annealing	Tm	30 s	35
Extension	72	1 min	
Final Extension	72	5　min	1
Hold	4	∞	-

Table 2.

Primer Set Name	Fwd Sequence	Rev Sequence	Tm
Actin (520bp)	CCTCAGAACAGTCGCAATGG	CCAGAGTCCAGCACGATAC	62
Actin (150 bp)	GTCATCAGGGTGTCATGGTC	CTTTTCCCTGTTGGCCTTGG	62
HSP90	CTTCACCATCCAGACGGACA	CCCAATCGTTAGTGAGGC	62
HSP70	ACGTGGTGAAGACTTGC	TGAAGCCAGCATTGAGA	65*
* 72°C (-2°C/cycle) for six cycles before a TM at 65 °C for non quantitative PCR			

Table 3.

Step	Temp. (°C)	Duration	No. of cycles
Initial denaturation	95	60 s	1
Denaturation	95	15 s	30
Annealing Extension	Tm	30 s	
Denaturation	95	10 s	1
Melt Curve Analysis	65 (+0.5°C)	10 s	60

Table 4.

RNA Kit	Cost/Prep	STST	Cost/Prep
Kit	$6.82	Detergent 1	3.6E-07
CDNA prep Cost	$2.07	Tris base	0.0002
Linear Amplificaiton PCR k	$99.32	Detergent 2	2.56E-06
		EDTA	5.5E-07
		RNAse inhibitor	0.23
		Glass Micropipette	0.07
		CDNA Prep Cost	0.57
Total	$108.21	Total	$0.87