**Title**

Endopiriform neurons projecting to ventral CA1 are a critical node for recognition memory

**Authors**

Naoki Yamawaki[1,2,3], Hande Login[1,2,3], Solbjørg Østergaard Feld-Jakobsen[1], Bernadett Mercedesz Molnar[1], Mads Zippor Kirkegaard[1], Maria Moltesen[1], Aleksandra Okrasa[1], Jelena Radulovic[1,2,3,4,5] and Asami Tanimura*[1,2,3]

**Affiliations**

1. Department of Biomedicine, Aarhus University, Aarhus C, Denmark
2. PROMEMO, The Center for Proteins in Memory, Aarhus University, Aarhus C, Denmark
3. DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus C, Denmark
4. Dominick P. Purpura Department of Neuroscience, Albert Einstein College of Medicine, New York, USA
5. Department of Psychiatry and Behavioral Sciences, Albert Einstein College of Medicine, New York, USA

*Correspondence: Asami Tanimura (asami.tanimura@biomed.au.dk)

**Abstract**

The claustrum complex is viewed as fundamental for higher order cognition; however, the circuit organization and function of its neuroanatomical subregions are not well understood. We demonstrated that some of the key roles of the claustrum complex can be attributed to the connectivity and function of a small group of neurons in its ventral subregion, the endopiriform (EN). We identified a subpopulation of EN neurons by their projection to the ventral CA1 (EN$^{\text{vCA1-proj.}}$ neurons), embedded in recurrent circuits with other EN neurons and the piriform cortex. Although the EN$^{\text{vCA1-proj.}}$ neuron activity was biased toward novelty across stimulus categories, their chemogenetic inhibition selectively disrupted the memory-guided but not innate responses of mice to novelty. Based on our functional connectivity analysis, we suggest that EN$^{\text{vCA1-proj.}}$ neurons serve as an essential node for recognition memory through recurrent circuits mediating sustained attention to novelty, and through feed forward inhibition of distal vCA1 neurons shifting memory-guided behavior from familiarity to novelty.

**Introduction**

The claustrum complex, an evolutionarily conserved brain region found across mammalian species, reptiles, and birds, is hypothesized to serve as a node for establishing higher order cognitive functions by coordinating neuronal activity on a global scale (*1-3*). This view is based on its extensive connections with many cortical and subcortical areas (*4, 5*). Some of these functions include sensory perception and attention, which may affect memory processing, including working memory, associative memory, and recognition memory (*6-8*). Consistent with this view, abnormalities of the claustrum complex is found in major cognitive disorders, including Alzheimer's disease, schizophrenia, and attention-deficit/hyperactivity disorders (*9-11*). However, microcircuitry and functional segregation of the individual constituents of the claustrum complex have remained unclear.

The cytoarchitecture and genetic expression broadly divide the claustrum complex into dorsal and ventral parts, with the dorsal part typically referred to as "claustrum" and ventral part having its own nomenclature: "endopiriform (EN)" in rodents (*12, 13*). Current evidence indicates that the claustrum forms reciprocal connections with dorsal cortices containing sensory, motor and

51  association areas generating diverse physiological effects depending on the cortical targets and
52  their activity patterns (*4, 5*). In contrast, EN primarily provides inputs to the piriform cortex and
53  limbic systems (*14, 15*), suggesting its distinct circuit organization and role in bridging olfactory
54  information processed by the piriform cortex and memory processed by the limbic systems (*16,*
55  *17*).
56
One of distinct targets of EN, the ventral CA1 (vCA1), is known to coordinate in a range of
behaviors related to exploration and recognition memory, and these functions are thought to be
regulated by its afferent inputs in a domain-specific manner (*18-24*). We, therefore, hypothesized
that EN is a key node for establishment of recognition memory. To address this, we set out to
characterize circuit and function of EN neurons defined by their projection to vCA1 (EN $^{\text{vCA1-proj.}}$
neurons) using genetic tools and mouse models of social and object recognition memory.
63
We found EN $^{\text{vCA1-proj.}}$ neurons innervated multiple components of the limbic system except
amygdala and prefrontal cortex and produced potent feedforward inhibitory control over vCA1
pyramidal neurons. During recognition memory test, activity of EN $^{\text{vCA1-proj.}}$ neurons were
condensed around conspecifics or objects where mice spent most time on. However, disruption of
EN $^{\text{vCA1-proj.}}$ activity only impaired memory-guided exploration of novel stimuli without affecting
innate exploration induced by novelty. These findings demonstrate that EN subserves some of the
key functions required for recognition memory governed by specific limbic system.
71
**Results**
**Endopiriform represents a major afferent of the ventral CA1**
To assess the significance of EN as a vCA1 afferents, we injected a retrograde tracer into the
vCA1 of mice and compared the number of the retrogradely labeled neurons in the EN and other
brain regions (**Fig. 1A, B**, **Methods**). We found labeled neurons in multiple areas including the
septum, entorhinal cortex, and basolateral amygdala (**Fig. 1C-F and Fig. S1A**) (*25, 26*). In
addition to these well-known vCA1 afferents, prominent labeling was observed in the area
defined as EN in Allen Brain Atlas (**Fig. 1G, Methods**). The number of presynaptic neurons in
EN was ~10 fold lower than entorhinal cortex (data not shown). However, the normalized count
of presynaptic neurons in EN was significantly greater than basolateral amygdala, lateral septum,
and medial septum (**Fig. 1H**).
The vCA1-projecting neurons in the EN were spread along antero-posterior axis with two peak
densities, one in anterior and another in posterior to the bregma (**Fig. 1I, J**). Number of labeled
neurons was greater in posterior EN (**Fig. 1J**). In contrast, the same analysis along dorso-ventral
axis showed a single dominant peak at the depth consistent with the location of EN (**Fig. S1B**).
To further confirm the location of these vCA1-projecting neurons, we injected a retrograde tracer
in one color into vCA1 and a retrograde tracer in another color into a cortical area known to
receive projections from EN or "claustrum" representing a dorsal part of claustrum complex (**Fig.**
**2A**). We targeted the prefrontal cortex to label EN, and the anterior cingulate cortex, the motor
cortex, and the dorsomedial entorhinal cortex to label claustrum (**Fig. 2C-F**) (*27-30*).
We found there is a major spatial overlap in labeling of vCA1-projecting neurons with prefrontal
cortex-projecting neurons, but not with other cortical projection neurons (**Fig. 2G-J**). Moreover,
around 30% of vCA1-projecting neurons were found to be double-labelled when a cortical
injection was made into the prefrontal cortex, suggesting some vCA1-projecting neurons send
axons to this cortical area (**Fig. 2K, L**).
To further determine the cortical and subcortical innervation pattern of vCA1-projecting EN
neurons, we used an intersectional approach to label their somata and axons with eGFP (**Fig.**
**2M**). Apart from vCA1, axons from vCA1-projecting neurons were found in various cortical
areas including prefrontal cortex, lateral entorhinal cortex, and piriform cortex (**Fig. 2N-P and**

**Fig. S2A-C**). Projection to prefrontal cortex was sparse relative to other areas, as expected based
on the retrograde labeling data (**Fig. 2K**). The connectivity of vCA1-projecting EN neurons to the
amygdala, which represents the major component of limbic systems, was also sparse relative to
vCA1 (**Fig. S2C**), indicating vCA1 is the main target of these EN neurons in this system.
Taken together, these findings indicate EN represents a major afferent of vCA1. Moreover, EN
neurons projecting their axons to vCA1 also send strong collaterals to lateral entorhinal cortex
and piriform cortex, but relatively weak to the prefrontal cortex or other limbic structures. Since
our subsequent study focused on EN neurons defined by their projection to vCA1, we refer to
them as $EN^{vCA1-proj.}$ neurons.
110
**Projection pattern and synaptic connectivity of EN axons in vCA1**
We next investigated projection pattern of EN axons in hippocampus by injecting an anterograde
tracer (AAV-GFP) into the EN (**Fig. 3A, B**). Analysis of different parts of hippocampal section
indicated EN axons were mostly confined in stratum lacunosum-moleculare (SLM) layer of distal
part of vCA1 (**Fig. 3C, D**). Projection to intermediate or dorsal CA1 was limited or undetectable
(**Fig. S3A-C**).
Since SLM mainly consists of GABAergic neurons (*31*), we hypothesized that EN axons form
synapses with this cell type. To test this, we first applied monosynaptic rabies tracing technique to
Vgat-Cre mice (**Fig. 4A, Methods**). The distribution of starter cells (identified by GFP and
mCherry co-labeling) was confirmed to be within the ventral-intermediate region of hippocampus
using AMaSiNe (**Fig. 4B, S4A**) (*32*). In these mice, the presynaptic neurons were consistently
observed in EN in addition to other expected areas (e.g., septum and entorhinal cortex) (**Fig. 4C**
and **d, S4B-F**) (*33, 34*). The density of the presynaptic neurons in EN was greater at the posterior
region to the bregma (**Fig. 4D**), consistent with our earlier observation (**Fig. 1J**).
To determine how EN inputs are integrated into the local circuit in vCA1, we systematically
assessed the connections between EN axons and GABAergic neurons in different layers of distal
vCA1. We expressed channelrhodopsin-2 (ChR2) into the EN axons by injecting AAV-ChR2-
mCherry into EN and performed whole-cell recording from GABAergic neurons in acute vCA1
slices in the presence of TTX (1 µM) and 4-AP (100 µM) to isolate monosynaptic connection
(*35*). To ensure the recording from GABAergic neurons, we used Vgat::Ai14 mice in which all
GABAergic neurons are labeled with tdTomato (**Fig. 4E-G**). We recorded from 3 to 4
GABAergic neurons located in different laminar position in distal vCA1 (Py-SR border, SR, SR-
SLM border, or SLM) in the same slice to construct a "laminar profile" of EN→vCA1 inputs
(**Fig. 4F and G**). Whole-field LED stimulation evoked excitatory postsynaptic current (EPSC) in
most GABAergic neurons recorded. However, contrary to the expectation, connection probability
and strength were lowest in neurons in SLM compared to neurons in other layers (**Fig. 4H-J**).
Taken together, these results indicate EN axons preferentially innervate the GABAergic neurons
spanning across the Py-SR border to SR-SLM border in distal vCA1.
139
**EN axons produce feedforward inhibition onto vCA1 pyramidal neurons**
SLM also contains the apical tuft dendrite of pyramidal neurons, which could be a target of EN
axons in addition to GABAergic neurons. To test this, we expressed ChR2 in EN axons but made
whole-cell recording from pyramidal neurons in distal vCA1 in the presence of TTX and 4-AP.
Since CA1 pyramidal neurons can be differentiated into superficial and deep layer neurons with
distinct anatomy and function (*36*), we recorded from neurons in each layer as well as nearby
GABAergic neurons at the Py-SR and SR-SLM borders for the comparison (**Fig. 5A**). We found
monosynaptic input to superficial and deep layer pyramidal neurons was similar, and
monosynaptic input to pyramidal neurons (superficial and deep pooled) and GABAergic neurons
was also comparable (**Fig. S5**).

Since EN axons innervated multiple GABAergic neurons across different layers, we hypothesized
they may exert stronger inhibition onto pyramidal neurons through feed-forward inhibition. To
test this, we recorded EPSC and inhibitory postsynaptic current (IPSC) from superficial and deep
pyramidal neurons using cesium-based internal solution in the brain slices expressing ChR2 in the
EN axons (**Methods**). At the command potential of -70 mV, photo-stimulation of EN axons
evoked EPSCs in majority of pyramidal neurons in both layers (7 out of 10 neurons in superficial
and 5 out of 10 neurons in deep) (**Fig. 5B**). At the command potential of +12 mV, the same
stimulation evoked outward postsynaptic current in the same neurons, including those without
EPSCs (10 out of 10 for superficial and deep) (**Fig. 5B**). This outward current was abolished by
application of TTX and 4-AP or gabazine (10 µM), indicating it is disynaptically driven and
mediated by inhibitory GABA$_A$ receptor (**Fig. 5B**). When relative strength of EPSCs and IPSCs
were compared for each neuron, the strength of IPSCs overwhelmed the EPSCs in all cases (**Fig.
5C**). This indicates EN axons disynaptically inhibit pyramidal neurons in vCA1 (**Fig. 5D**).

**EN$^{vCA1-proj.}$ neurons receive inputs from the piriform cortex**
We next examined afferents that may drive the EN→vCA1 circuit. To address this, retrograde
CAV-2-Cre was injected into vCA1 to express Cre into EN$^{vCA1-proj.}$ neurons, then applied
monosynaptic rabies tracing technique to specifically label its presynaptic neurons in wild type
mice (**Fig. 6A**). Starter cells and presynaptic neurons were spatially mapped and quantified using
AMaSiNe (**Fig. 6B and C, S6**). Since presynaptic neurons were sparse in contralateral
hemisphere, data from ipsilateral hemisphere was analyzed. Moreover, we applied minimum
labeling threshold to determine the brain region for further analysis (**Methods**). We found the
piriform cortex contained highest number of presynaptic neurons followed by EN (**Fig. 6D-F**).
Within piriform cortex, layer 2 neurons were the major source of afferent (**Fig. 6G**). These data
indicate EN$^{vCA1-proj.}$ neurons receive major input from piriform cortex and form a recurrent
connection with neurons in EN (**Fig. 6H**).

**The activity of EN$^{vCA1-proj.}$ neurons is correlated with the time mice spend in space**
The downstream target of EN, vCA1, is implicated in social, odor and object recognition memory
(*20-22*). To determine whether and how the activity of EN neurons is related to this function, we
expressed GCaMP8s in EN$^{vCA1-proj.}$ neurons and monitored their activity using fiber photometry
while video recording mice performing social recognition memory test consisting of pretest,
sociability test, and discrimination test (**Fig. 7A, B, Methods**). The nose position of the subject
mouse was tracked during tests using DeepLabCut (**Fig. 7C**) (*37, 38*), and cumulative time and
calcium event were mapped onto the arena space (**Fig 7C, S7**). We found these values were
correlated in space across all sessions (i.e., the more time the mice spent in a given space, the
more the EN$^{vCA1-proj.}$ activity), including in the open field with no object or in the test arena with
conspecifics or objects (**Fig. S7, S8A-B**).
Cumulative calcium events of EN$^{vCA1-proj.}$ neurons became high towards the space around objects
or conspecifics when they are present and was correlated with interaction times except for
sociability session (**Fig. S8C-D**). In the sociability session, the cumulative calcium events around
a novel conspecific and object were similar despite mice spending more time on conspecifics
(**Fig. 7E left, S8C**). In contrast, in the social discrimination test, both the cumulative time and
cumulative EN$^{vCA1-proj.}$ activity were higher for an unfamiliar than familiar conspecific (**Fig. 7D-
E, S8C**). As a result, EN$^{vCA1-proj.}$ activity and behavior had a greater degree of correlation in social
discrimination than sociability.
We also tested if a similar correlation occurs in a non-social context by recording EN$^{vCA1-proj.}$
activity and behavior in the object discrimination test (**Fig. 7F**). The cumulative time and calcium
events were significantly correlated across different sessions i.e., familiarization, and object

discrimination (**Fig. 7G, S8E**), largely consistent with the findings in the social discrimination
test.
These data indicate that the activity of $EN^{vCA1\text{-}proj.}$ neurons is generally correlated with the time
the mouse spends in a particular location at the basal level (i.e. open field) or task conditions.
However, the degree of correlation appears to vary depending on the session, such that it was
highest in the pretest/familiarization and social/object discrimination test, but lowest in the
sociability test.

**Inhibition of $EN^{vCA1\text{-}proj.}$ neurons impairs social/object recognition memory**

Although $EN^{vCA1\text{-}proj.}$ activity predicted the behavior in pretest/familiarization and discrimination
test better than sociability test, their causality is unclear. To address this, we inhibited the activity
of $EN^{vCA1\text{-}proj.}$ neurons using inhibitory DREADD (hM4Di) during social and object
discrimination tests (**Fig. 8A-C and h, S9A**) (*39*). To specifically express hM4Di in $EN^{vCA1\text{-}proj.}$
neurons, CAV-2-Cre and AAV-flox-hM4Di were bilaterally injected into vCA1 and EN,
respectively (**Fig. 8A-B**). hM4Di was replaced with tdTomato for control group. We made a
protocol that allows within-subject comparison, such that a mouse goes through a test with
chlozapine-N-oxide (CNO, 1 mg/ kg) treatment then the same test with saline treatment the next
day, or vice versa (**Fig. 8C and H**). In social discrimination test, CNO treatment did not affect the
pretest or sociability in hM4Di or control group, but specifically impaired social discrimination in
hM4Di group (**Fig. 8D-G, S9B-C**). A similar effect was also observed in novel object
discrimination (**Fig. 8I, S9D**).
Apart from recognition memory, vCA1 is implicated in anxiety and associative fear memory (*40,
41*), thus, we tested contribution of $EN^{vCA1\text{-}proj.}$ neurons to these functions. The anxiety level as
determined by locomotor activity and sociability was not affected by CNO treatment in hM4Di
group (**Fig. 8E, S9E**). Similarly, fear memory as assessed by the freezing response during training
(with trace fear conditioning) or recall to context or tone was not affected by CNO treatment in
hM4Di group (**Fig. S10**).
Taken together, these data indicate that $EN^{vCA1\text{-}proj.}$ activity is not causally related to innate
exploration behavior induced by novelty, or anxiety or fear memory. However, their activity is
necessary for mice to discriminate between familiar and unfamiliar conspecific or object,
suggesting their major role in general recognition memory.

**Discussion**

We characterized the circuit and function of EN neurons targeting specific subregions and layers
of vCA1. In contrast to other parts of claustrum complex, $EN^{vCA1\text{-}proj.}$ neurons were reciprocally
connected with the piriform cortex representing a major downstream target of the olfactory bulb
(*42*). This circuit motif may predict the predominant role of $EN^{vCA1\text{-}proj.}$ neurons in social
recognition memory, given the ability of social odor to engage multiple olfactory pathways
innervating the piriform cortex (*43*). However, we found that these neurons contributed to broader
domains of recognition memory. This is supported by the observations that the inhibition of
$EN^{vCA1\text{-}proj.}$ activity impaired both social and non-social discrimination.

In all phases of recognition memory test, $EN^{vCA1\text{-}proj.}$ activity was highly correlated with the time
mice spent in given locations and this correlation was biased towards novel stimuli when mice
were engaged in their exploration irrespective of the stimuli being social or non-social (**Fig. 9A**).
Taken together, these data suggest that the function of $EN^{vCA1\text{-}proj.}$ neurons was primarily related
to the detection of and attention to novel stimuli. The recurrent connections within EN, which is
another prominent feature of $EN^{vCA1\text{-}proj.}$ neurons' network, potentially support these processes to
help establish the recognition memory (*44*).
Attentional processes mediated by $EN^{vCA1\text{-}proj.}$ neurons can be considered of particular relevance
for memory-guided behavior, given that inhibition of $EN^{vCA1\text{-}proj.}$ activity selectively impaired

conspecific/object discrimination but not innate exploratory behavior provoked by novelty as seen in pretest, sociability, and familiarization (**Fig. 9A**). This functional specialization, likely associated with its unique circuit connectivity to the limbic system, could differentiate EN from claustrum.

Studies on EN are generally scarce, and the available evidence of EN projections to vCA1 suggests that these projections are sparse (*14*). Potential reasons include the difficulty in clearly delineating the EN from claustrum complex and non-uniform distribution of EN$^{vCA1-proj.}$ neurons along antero-posterior axis. By leveraging contemporary circuit analysis tools, we demonstrated that the EN→vCA1 circuit is non-trivial; EN$^{vCA1-proj.}$ neurons were more numerous than commonly studied vCA1 afferents, including those from the septum and amygdala. Moreover, these axons exhibited a potent disynaptic inhibition of distal vCA1 pyramidal neurons, indicating their major functional implications for vCA1 activity.

Our data indicated that although EN axons terminated at the SLM of distal vCA1, they exhibited stronger connections to GABAergic neurons spanning from the Py-SR border to the SR-SLM border than to SLM neurons. The detection of EN inputs from GABAergic neurons at the Py-SR border was surprising, considering the distance between soma position and the EN axons. However, there are several GABAergic cell types below SLM that extend their primary dendrites into SLM (*45, 46*), likely explaining the observed connectivity pattern. Among these are chandelier cells found at the Py-SR border, which preferentially inhibits the axon initial segments of pyramidal neurons, and are thus well suited to exert the powerful feedforward inhibition of vCA1 pyramidal neurons observed in our recordings (*47*).

There are several potential circuit effects by which EN inputs could affect vCA1 function. One is the induction of non-linear events in vCA1 pyramidal neurons by acting in concert with other afferents (e.g., lateral entorhinal cortex targeting SLM) (*34*) to generate a specific output pattern in pyramidal assemblies representing task-relevant stimuli. Alternatively, EN inputs may improve signal to noise ratio of the information conveyed through the lateral entorhinal cortex, CA2 (in the case of social context), and possibly other afferents (e.g., CA3) (*48*). Another scenario is tuning of vCA1 pyramidal neurons' response to familiar stimuli through dynamic combination of a weak monosynaptic excitatory input and strong disynaptic inhibitory input. Such mechanism suggests that at a milliseconds scale novelty (EN) and familiarity (vCA1) recognition might occur though alternating rather than simultaneous activity of brain microcircuits. Taken together, we propose a model for the role of EN$^{vCA1-proj.}$ neurons in recognition memory by balancing memory-guided attentional responses to familiarity and novelty through combination of feedforward inhibition of vCA1 (a node for recognition of familiarity) and the recurrent circuits that contribute to sustain attention to novelty (**Fig. 9B**).

The model of promoting novelty detection by suppressing familiarity responses is consistent with previous observations showing that vCA1 pyramidal neurons predominantly respond to familiar stimuli whereas novelty response can only occur through activation of vCA1 interneurons (*49, 50*). Nevertheless, alternating activation of EN and vCA1 could prove essential for the behavioral relevance of EN activity associated with recognition memory, without affecting innate exploratory behavior to novelty. Accordingly, the latter behavior is primarily associated with the prefrontal cortex and amygdala, areas that are only weakly targeted by EN$^{vCA1-proj.}$ neurons (**Fig. 2P, S2C**) (*51-53*).

In addition to advancing our understanding of the basic organization and function of brain circuits underlying higher cognitive processes, our findings suggest dysfunction of EN$^{vCA1-proj.}$ neurons

300 could be a key contributing factor to the deficits in claustrum complex function and recognition
301 memory found in neuropsychiatric disorders (*9, 10*). Discrete EN populations can thus emerge as
302 pathophysiological substrate, but also as important treatment targets of key symptoms of these
303 disorders.
304
**Materials and Methods**
**Mice**
All experiments were performed in accordance with standard ethical guidelines and were
approved by the Danish national animal experiment committee (License number: 2021-15-0201-
00801). Unless otherwise noted, all mice used were C57BL6/J mice or transgenic mice with
C57BL6/J backgrounds. They were 2-4 months old at the time of the experiment. Transgenic
mice used were B6.Cg-Gt (ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J mice (Ai14, JAX007914) and
B6J.129S6(FVB)-Slc32a1$^{tm2(cre)Lowl}$/MwarJ mice (Vgat-Cre, JAX028862). Similar number of
male and female mice were used for all experiments except for behavior tests. For behavior tests,
male mice were used.
**Viruses and retrograde tracers**
The information on viruses used in our experiments was the following. CAV-2-Cre (PVM);
AAV1-mDlx-HBB-chl-dlox-TVA_2A_oG(rev)-dlox (v271-1, VVF); AAV1-hSyn1-dlox-
TVA_2A_mCherry_2A_oG(rev)_dlox (v306-1, VVF); Rabies-GFP (NTNU viral core facility);
Rabies-mCherry (NTNU viral core facility); AAV5-CAG-GFP (37525-AAV5, Addgene); AAV5-
CAG.hChR2(H134R)-mCherry (10054-AAV5, Addgene); AAV8-hSyn-DIO-hM4Di (Gi) –
mCherry (44362-AAV8, Addgene); AAV5-hEF1a-dlox-EGFP(rev)-dlox (v217-5, VVF); and
retrograde AAVrg-CAG-GFP (37825-AAVrg, Addgene), AAV1-hSyn1-dlox-jGCaMP8s(rev)
(v627-1, VVF). We also used retrograde traces of red Retrobeads (Lumafluor) and Cholera toxin
subunit B conjugated with Alexa 647 (CTB647, Thermo Fisher).
**Stereotaxic injections**
Stereotaxic injections were performed using a stereotaxic frame (Model 940, Kopf). Mice were
anesthetized with isoflurane and were subcutaneously injected with buprenorphine (0.1 mg/kg)
and metacam (1.5 mg/Kg) for post-operative pain relief. After incising the scalp over the cranium,
a small hole was bored with a microdrill bit, and a beveled glass pipette (Wiretrol II, 5-000-2010,
Drummond Scientific Company) back-filled with mineral oil and front-filled with the material to
be injected was slowly inserted into a target coordinate. After injecting a small volume (50 - 100
nL) with a custom-made displacement injector (based on MO-10, Narishige), the pipette was left
in place for 3 to 5 mins before slow retraction. The incision was closed with a nylon suture. The
stereotaxic coordinates used were (relative to bregma, in mm): anteroposterior (AP) -3.0;
mediolateral (ML) 3.12; dorsoventral (DV) 3.7 for ventral CA1; AP +2.0; ML 0.2; DV 1.5 and
2.5 for prefrontal cortex; AP 0.0; ML 0.2; DV 1.0 and 1.5 for anterior cingulate cortex; AP 0.0;
ML 1.5; DV 0.3 and 0.7 for motor cortex; and AP -5.0; ML 3.0; DV 1.5 for dorsomedial
entorhinal cortex. For endopiriform injection, we injected two sites from the following
coordinates: AP -0.27, ML 3.2, DV 4.2; AP 0.0, ML 3.0, DV 4.2; or AP +1.0, ML 2.7, DV 4.0.
Mice were thermally supported with a feedback-controlled heating pad maintained at ~37 °C
(ThermoStar Homeothermic system, RWD). Mice were used for experiments 3-5 weeks post
injection.
**Implantation of the optic probe**
After stereotaxic injection of CAV-2-Cre into the vCA1 and AAV1-hSyn1-dlox-jGCaMP8s into
the EN (AP -0.27, ML 3.2, DV 4.2), a fiber optic cannula (400 μm core diameter, 4.5 mm length,
NA 0.39, R-FOC-BF-400C-39NA, RWD) was inserted towards EN. Once in the correct depth,
exposed brain surface around the implant was covered by Kwik-Cast (WPI) and was then fixed on
skull with dental adhesive resin cement (Super-Bond, SUN MEDICAL).
*Ex vivo* **electrophysiology**

Mice were deeply anesthetized with isoflurane and decapitated. Horizontal sections (300 μm)
containing ventral hippocampus were prepared by vibratome (VT1200S, Leica) in ice-cold
choline solutions containing (in mM): 25 $NaHCO_3$, 1.25 $NaH_2PO4-H_2O$, 2.5 KCl, 0.5 $CaCl_2$, 7
$MgCl_2$, 25 D-glucose, 110 Choline chloride, 11.6 Ascorbic acid, and 3.1 $C_3H_3NaO_3$. Slices were
subsequently incubated in artificial cerebrospinal fluid (ACSF) containing (in mM): 125 NaCl, 25
$NaHCO_3$, 1.25 $NaH_2PO4-H2O$, 2.5 KCl, 11 D-glucose, 2 $CaCl_2$, and 1 MgCl at 34 $^o$C for 30 mins
then at room temperature (~20 $^o$C) for at least 1 hour before the recording.
Whole-cell recording was performed with an upright microscope (BX51WI, Olympus) equipped
with a motor-controlled stage and focus (MP285A and MPC-200, Sutter instrument), differential
interference contrast, coolLED (model pE300), and a monochrome camera (Moment, Teledyne).
Neurons were visualized with a 60x lens (1.00 NA, LUMPlanFL N, Olympus) with the software
Micro-Manager-2.0 gamma (US National Institutes of Health). Pipettes (4-5 MΩ) were pulled
from thick-walled borosilicate capillary glass with a puller (model P-1000, Sutter instrument). For
a voltage-clamp recording, the internal solution contained (in mM): 135 $CsMeSO_3$, 10 HEPES, 4
Mg-ATP, 0.3 Na-GTP, 8 $Na_2$-Phosphocreatine, 3.3 QX-314 (pH was adjusted to 7.35 with
CsOH). In some experiments, a biocytin (4 mg/mL) and Alexa488/586 (50 μM) were further
added for morphological studies. Recordings and hardware control were performed with
Wavesurfer (Janelia Farm). Signals were amplified and Bessel filtered at 4 kHz with an amplifier
(MultiClamp 700B, Molecular Device), then sampled at 10 kHz with a data acquisition board
(USB-6343, National Instrument). Recording with access resistance change of > 20% from the
baseline (~30 MΩ or less) were discarded. Liquid junction potential was not corrected. All
recordings were performed at ~34 $^o$C maintained with the inline heating system (TC-324C,
Warner instrument).
To stimulated channelrhodopsin-2, a blue LED was delivered at a short pulse (5-msec) through a
4x objective lens (0.16 NA, UPlanSApo, Olympus). The light power (measured at the focal plane)
was 20 mW for the recordings in normal ACSF and 40 mW for the recordings in the presence of
tetrodotoxin and 4-aminopyridine. Photo-stimulation was repeated 3 to 6 times (at intervals of 10
or 20 sec) to obtain an average response trace. Photo-evoked responses less than 2 standard
deviations of the baseline were considered as a "zero" response. The mean amplitudes indicate
mean amplitudes between LED stimulus-onset to 50 ms of average responses.
**Quantification of presynaptic neurons**
Mice were deeply anaesthetized with a ketamine (120 mg/kg) and xylazine (24 mg/kg) mixture
and were transcardially perfused with 4% paraformaldehyde (PFA). The brains were extracted
and immersed in 4% PFA overnight followed by 20 and 30% sucrose solution for a
cryoprotection. The brains were subsequently embedded in O.C.T compound (Tissue TEK,
Sakura) in a mold and frozen on the dry ice. Coronal slices (100 μm) of entire brain were cut with
Cryostat (CM3050S, Leica) and were washed with phosphate-buffered saline (PBS) before DAPI
staining. All sections were mounted onto glass slides with a coverslip (thickness:0.13-0.17 mm,
Hounisen) and mounting media (DAKO, S3023, Agilent Technologies).
The injection sites and its specificity were verified by registering the images into Allen Brain atlas
using QuickNII and VisuAlign (*54*).
For quantification of presynaptic neurons and visualization of their distribution in 3D mouse
brain, epifluorescence images acquired with Slidescanner (4231 x 3462 pixels, 10x objective,
Olympus VS120) were analyzed with Matlab-based program AMaSiNe (*32*). For **Fig.1**, the
coronal section between AP + 3 to -4 mm were registered in AMaSiNe and analyzed.
For the monosynaptic rabies tracing study in **Fig. 6**, the brain areas containing over 1%
presynaptic cells of total presynaptic cells were proceeded further analysis. Fraction of
presynaptic cells in the different areas were calculated by dividing a number of presynaptic cells
in each brain areas by the average number of presynaptic neurons in all brain areas.

For the study in **Fig. 2**, the sections were immunostained with NeuN before imaging. Here,
sections were immersed with 5% normal goat serum and 0.2% Triton X for 30 mins at room
temperature before incubation with NeuN primary antibody (anti-rabbit, 1:1000) (ABN78,
Milipore) at 4 °C for overnight. The sections were then washed with PBS and were further
incubated with secondary antibody (goat anti-rabbit conjugated with Alexa 488 or 568, 1:500)
(A11008 or A11036, Thermo Fisher) for 1 hour at room temperature before mounting. A single
plane image of claustrum complex was acquired with confocal microscope (10x objective, 1024 x
1024 pixels, Zeiss). 2-4 images, range of AP: +1.4 to -0.2, were collected per brain. Fraction of
double-labelled neurons (with different retrograde tracers) were calculated for each brain by
dividing a total number of double-labeled neurons by the total number of vCA1-projecting
neurons (labeled with one color).
**Quantification of fluorescence signal from axons**
For **Fig. 3 and S3**, horizontal section (100 μm) images were taken with Sliderscanner (10x lens,
pixel: 2560 x 3072). All images from the same brain were taken with the same light intensity and
detection setting. After background subtraction, GFP signals in ventral, intermediate, and dorsal
CA1 column were normalized by the highest signal in ventral CA1 in the same brain.
For **Fig. 2**, coronal sections (100 μm) were immunostained with GFP antibody before imaging.
Here, sections were washed with PBS, the sections were incubated in blocking solution
containing 5% normal goat serum and 0.2% Triton X for 30 mins at room temperature before
incubation with GFP primary antibody (anti-rabbit, 1:1000) (NB600-308, MNovus Biologicals) in
blocking solution at 4 °C for overnight. The sections were then washed with PBS and were further
incubated with secondary antibody (goat anti-rabbit conjugated with Alexa 488, 1:300) (A11008,
Thermo Fisher) for 1 hour at room temperature before mounting. Images were taken with
Slidescanner (10x, pixel: 3072 x 3584). Signal per pixel in ROI was calculated for each projection
site after background subtraction.
**Social memory test with chemogenetic inhibition of EN$^{vCA1\text{-}proj.}$ neurons**
Social memory test is based on previous articles with slight modifications (*55*)$^,$ (*56*). All tested
mice were habituated to the handling for 3 days. Prior to the test day, we divided subject mice
into two groups. On the 1$^{st}$ test day (session 1), one group received chlozapine-N-oxide (CNO, 1
mg/kg, i.p.) and another group received a vehicle. After 30 mins, the subject mice were placed in
a three-chamber box for 5 mins (habituation), then the same box with two pencil chambers
containing the same object in opposite corners for 5 mins (pretest). After 5 mins break in their
home cage, the subject mice were again placed in the same box, but this time, an object in one
pencil chamber was replaced with different object and an object in another pencil chamber was
replaced with a conspecific (3-4 weeks old male, habituated to the handling and pencil chamber
for 3 days prior to the session 1). This "sociability test" lasted for 10 mins and was followed by
social discrimination test after 30 mins break in the home cage. The social discrimination test
lasted for 5 mins and consisted of the same setting as the sociability test except an object in a
pencil chamber, which was replaced with a new conspecific (i.e., a novel mice). In the following
day, CNO and saline treatments were swapped between the groups, and all mice went through the
same series of tests (session 2) for the within-subject comparison of the CNO effect. The position
of the pencil chamber including an object or a conspecific were counterbalanced. The tests were
performed under 6-10 lx room light and video was recorded at 25 frames/sec with GigE camera
(Basler ace acA1300-60gm, Basler) placed at the ceiling.
**Trace fear conditioning**
After one week of social memory test, subject mice were examined for their tone-associated fear
memory and extinction. One group received CNO while another group received vehicle 45 mins
before the test. In conditioning day, mice were placed in a box with a metal grid floor and ethanol
odor (context A) and after 3 mins were delivered with 20-sec pure tone (2 kHz, 20 dB) followed
by 18-sec time gap (trace) and 2-sec footshock (0.5 mA). The exposure to the same tone-trace-

shock sequence was repeated 2 more times with inter-trial interval of 2 mins. In the following
day, mice were re-exposed to the context A for 3 mins to assess their fear memory to the context.
From day 3, mice were placed in a new context, the context B (smooth floor, breach odor), and
went through the same protocol as in conditioning day but without shocks. The process was
repeated over 5 days to assess the extinction of the fear memory to the tone and trace. All test was
performed in dark (but infrared light was on), and behavior was video recorded at 25 frames/sec
with infrared GigE camera (Basler ace acA1300-60gm, Basler).

**Novel object recognition test**
The novel object recognition test is based on the previous article with slight modifications (*55*).
The subject mice were divided into two groups and CNO treatment was done as same as social
memory test. 40 mins after CNO/vehicle i.p. injection, the subject mice were placed in an open
field box (50 x 50 cm) for 5 mins (habituation). In following familiarization session, two same
shape objects are placed in the open field box, and then the subject mice explored for 10 mins.
After 1 hour break in their home cage, the subject mice were again placed in the same box, but
one object is replaced by a novel object. The subject mice are allowed to explore the box for 5
min in discrimination session.

**Social memory test and novel object recognition test with *in vivo* fiber photometry**
Optic probe implanted mice were habituated handling for 3 days prior the test. In test day, the
subject mouse attached with patch code were placed in an open field box (50 x 50 cm) for 5 min
(habituation). After habituation, the mouse was place in a home cage for resting and waiting
following session. For pretest, pencil chambers containing the same objects were placed at the
corners of the open field box. Pretest was tested for 5 min. For sociability test, a novel mouse (4
weeks old, male) or a novel object were placed in pencil chambers. Sociability test took 10 min.
After sociability test, the subject mouse return to a home cage and rest for 20 min. After 20 min
rest, the subject mouse was tested discrimination session for 5 min. In discrimination test, a novel
mouse (4 weeks old, male) or a familiar mouse that was used as the stimulus mouse in sociability
test were placed in pencil chambers. Both in sociability and discrimination test, positions of
pencil chambers were counterbalanced between subject mice. The tests were performed under 6-
10 lux room light and video was recorded at 60 frames/sec with GigE camera (Basler ace
acA1300-60gm, Basler) placed at the ceiling. Video acquisitions were controlled by Bonsai.
Simultaneous start of video recording and calcium signal recording was controlled by Raspberry
Pi.
After one week of social memory test, subject mice were examined for novel object recognition
test. The procedure, the arenas and objects were used the same as in the novel object recognition
test with chemogenetic inhibition of $EN^{vCA1\text{-}proj.}$ neurons.

**Behavior data acquisition and analysis**
Ethovision (version 16 or 17, Noldus) was used for the hardware control and data acquisition and
initial quantification for the behavior. Behavioral variables, including movement trajectory, times
spent in predefined areas, total distance traveled, travelling velocity, and freezing were extracted
and further analyzed/nested with custom functions in the Matlab (R2020b). For social memory
test, interaction zones were defined as an area between the edge of pencil chamber and 3 cm away
from the edge. When the nose of the subject mouse was in the zone, the time was counted as
interaction time. Sociability index was calculated by dividing total interaction time with mouse by
total interaction time with mouse and object. The discrimination index was calculated by dividing
total interaction time with a novel mouse by total interaction time with a familiar mouse and a
novel mouse. Data from mice were excluded if discrimination index was less than 0.5 during
saline treatment. For trace fear conditioning test, mice were considered "freezing" if pixel change
between frames was less than 0.1 % for more than 1 sec.
Deeplabcut (ver.2.3) was used for tracking of annotated body points of the subject mice
connected with patch code for calcium imaging of $EN^{vCA1\text{-}proj.}$ neurons during social memory test

and novel object recognition test. Nose points were used for further analysis using Matlab
(R2020b). To define interaction zone, reference points was set at the center of pencil chamber.
The areas that distance between nose point and reference point was less than 100 pixel for social
memory test and 50 pixel for novel object recognition test were dedicated to interaction zone.

**Fiber photometry and analysis**

Population activity of $EN^{vCA1\text{-}proj.}$ neurons were recorded using fiber photometry (Doric Lenses).
GCaMP8s was excited with 470 nm light at 20-30 µW measured at the tip (isosbestic point used
was 415 nm (15 µW)). Calcium-dependent and independent signals were collected using lock-in
mode. Signals were detected at 10x gain. Recorded signals were butterworth filtered (low-pass at
40Hz) and calcium signals (in z-score) were extracted using Doric neuroscience studio V6 (Doric
Lenses). Signals were further down sampled to 60 Hz using Matlab (R2020b) signal analyzer to
match the behavioral sampling data (60 fps). Z-score $> 2.56$ (alpha = 0.01) was determined to be
a calcium event. For heat map of cumulative time and calcium events, spatial binning was 50
pixels. To make correlation map (**Fig. S7 and S8A**), correlations between column of cumulative
time heat map and cumulative calcium event heat map were tested using Matlab corr function.

**Statistical analysis**

Data and statistical analysis were performed with MATLAB R2020b (Mathworks). Unless
otherwise noted, group data was presented as median, and first and third quartiles are shown for
dispersion. Statistics were performed using a non-parametric test (signed-rank test for paired data
and rank-sum test for unpaired data). One-way ANOVA (post hoc test: Tukey-Kramer test or
Kruskal-Wallis test) were applied for multi-group comparison. Two-way ANOVA were
performed for comparison of multi-factors. Pearson correlation coefficient was applied for
correlation analysis. Significance was defined as $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. Outlier was
detected by Grubbe test (GraphPad, significant level: 0.05). Sample number were indicated as
"n", animal number were indicated as "N".

**References**

1.    K. Narikiyo *et al.*, The claustrum coordinates cortical slow-wave activity. *Nat Neurosci* **23**, 741-753 (2020).
2.    H. Norimoto *et al.*, A claustrum in reptiles and its role in slow-wave sleep. *Nature* **578**, 413-418 (2020).
3.    F. C. Crick, C. Koch, What is the function of the claustrum? *Philos Trans R Soc Lond B Biol Sci* **360**, 1271-1279 (2005).
4.    J. B. Smith, A. K. Lee, J. Jackson, The claustrum. *Curr Biol* **30**, R1401-R1406 (2020).
5.    Y. Goll, G. Atlan, A. Citri, Attention: the claustrum. *Trends Neurosci* **38**, 486-495 (2015).
6.    J. M. Gardiner, A. J. Parkin, Attention and recollective experience in recognition memory. *Mem Cognit* **18**, 579-583 (1990).
7.    A. L. M. Siegel, A. D. Castel, The role of attention in remembering important item-location associations. *Mem Cognit* **46**, 1248-1262 (2018).
8.    K. Oberauer, Working Memory and Attention - A Conceptual Analysis and Review. *J Cogn* **2**, 36 (2019).
9.    C. Y. Chen *et al.*, The cognitive dysfunction of claustrum on Alzheimer's disease: A mini-review. *Front Aging Neurosci* **15**, 1109256 (2023).
10.   N. G. Cascella, G. J. Gerner, S. C. Fieldstone, A. Sawa, D. J. Schretlen, The insula-claustrum region and delusions in schizophrenia. *Schizophr Res* **133**, 77-81 (2011).
11.   X. Wang, Y. Jiao, T. Tang, H. Wang, Z. Lu, Altered regional homogeneity patterns in adults with attention-deficit hyperactivity disorder. *Eur J Radiol* **82**, 1552-1557 (2013).
12.   J. B. Smith *et al.*, The relationship between the claustrum and endopiriform nucleus: A perspective towards consensus on cross-species homology. *J Comp Neurol* **527**, 476-499 (2019).
13.   J. S. Grimstvedt *et al.*, A multifaceted architectural framework of the mouse claustrum complex. *J Comp Neurol* **531**, 1772-1795 (2023).

14. M. Behan, L. B. Haberly, Intrinsic and efferent connections of the endopiriform nucleus in rat. *J*
*Comp Neurol* **408**, 532-548 (1999).
15. K. Majak, J. Morys, Endopiriform nucleus connectivities: the implications for epileptogenesis and
epilepsy. *Folia Morphol (Warsz)* **66**, 267-271 (2007).
16. W. Fu, T. Sugai, H. Yoshimura, N. Onoda, Convergence of olfactory and gustatory connections
onto the endopiriform nucleus in the rat. *Neuroscience* **126**, 1033-1041 (2004).
17. G. D. R. Watson, J. B. Smith, K. D. Alloway, Interhemispheric connections between the
infralimbic and entorhinal cortices: The endopiriform nucleus has limbic connections that parallel
the sensory and motor connections of the claustrum. *J Comp Neurol* **525**, 1363-1380 (2017).
18. G. Pi *et al.*, Posterior basolateral amygdala to ventral hippocampal CA1 drives approach behaviour
to exert an anxiolytic effect. *Nat Commun* **11**, 183 (2020).
19. S. Ciocchi, J. Passecker, H. Malagon-Vina, N. Mikus, T. Klausberger, Brain computation.
Selective information routing by ventral hippocampal CA1 projection neurons. *Science* **348**, 560-
563 (2015).
20. M. R. Hunsaker, P. M. Fieldsted, J. S. Rosenberg, R. P. Kesner, Dissociating the roles of dorsal
and ventral CA1 for the temporal processing of spatial locations, visual objects, and odors. *Behav*
*Neurosci* **122**, 643-650 (2008).
21. T. Okuyama, T. Kitamura, D. S. Roy, S. Itohara, S. Tonegawa, Ventral CA1 neurons store social
memory. *Science* **353**, 1536-1541 (2016).
22. R. P. Kesner, M. R. Hunsaker, W. Ziegler, The role of the dorsal and ventral hippocampus in
olfactory working memory. *Neurobiol Learn Mem* **96**, 361-366 (2011).
23. T. Meira *et al.*, A hippocampal circuit linking dorsal CA2 to ventral CA1 critical for social
memory dynamics. *Nat Commun* **9**, 4163 (2018).
24. T. Tanimizu *et al.*, Functional Connectivity of Multiple Brain Regions Required for the
Consolidation of Social Recognition Memory. *J Neurosci* **37**, 4103-4116 (2017).
25. S. Tao *et al.*, Whole-Brain Mapping the Direct Inputs of Dorsal and Ventral CA1 Projection
Neurons. *Front Neural Circuits* **15**, 643230 (2021).
26. M. M. Gergues *et al.*, Circuit and molecular architecture of a ventral hippocampal network. *Nat*
*Neurosci* **23**, 1444-1452 (2020).
27. T. Kitanishi, N. Matsuo, Organization of the Claustrum-to-Entorhinal Cortical Connection in
Mice. *J Neurosci* **37**, 269-280 (2017).
28. H. Qadir *et al.*, Structural Connectivity of the Anterior Cingulate Cortex, Claustrum, and the
Anterior Insula of the Mouse. *Front Neuroanat* **12**, 100 (2018).
29. J. Jackson, M. M. Karnani, B. V. Zemelman, D. Burdakov, A. K. Lee, Inhibitory Control of
Prefrontal Cortex by the Claustrum. *Neuron* **99**, 1029-1039 e1024 (2018).
30. J. B. Smith, K. D. Alloway, Functional specificity of claustrum connections in the rat:
interhemispheric communication between specific parts of motor cortex. *J Neurosci* **30**, 16832-
16844 (2010).
31. K. A. Pelkey *et al.*, Hippocampal GABAergic Inhibitory Interneurons. *Physiol Rev* **97**, 1619-1747
(2017).
32. J. H. Song *et al.*, Precise Mapping of Single Neurons by Calibrated 3D Reconstruction of Brain
Slices Reveals Topographic Projection in Mouse Visual Cortex. *Cell Rep* **31**, 107682 (2020).
33. T. F. Freund, M. Antal, GABA-containing neurons in the septum control inhibitory interneurons in
the hippocampus. *Nature* **336**, 170-173 (1988).
34. O. M. Bilash, S. Chavlis, C. D. Johnson, P. Poirazi, J. Basu, Lateral entorhinal cortex inputs
modulate hippocampal dendritic excitability by recruiting a local disinhibitory microcircuit. *Cell*
*Rep* **42**, 111962 (2023).
35. L. Petreanu, T. Mao, S. M. Sternson, K. Svoboda, The subcellular organization of neocortical
excitatory connections. *Nature* **457**, 1142-1145 (2009).
36. N. B. Danielson *et al.*, Sublayer-Specific Coding Dynamics during Spatial Navigation and
Learning in Hippocampal Area CA1. *Neuron* **91**, 652-665 (2016).
37. A. Mathis *et al.*, DeepLabCut: markerless pose estimation of user-defined body parts with deep
learning. *Nat Neurosci* **21**, 1281-1289 (2018).

38. J. Lauer *et al.*, Multi-animal pose estimation, identification and tracking with DeepLabCut. *Nat*
*Methods* **19**, 496-504 (2022).
39. B. L. Roth, DREADDs for Neuroscientists. *Neuron* **89**, 683-694 (2016).
40. J. C. Jimenez *et al.*, Anxiety Cells in a Hippocampal-Hypothalamic Circuit. *Neuron* **97**, 670-683
e676 (2018).
41. J. C. Jimenez *et al.*, Contextual fear memory retrieval by correlated ensembles of ventral CA1
neurons. *Nat Commun* **11**, 3492 (2020).
42. J. M. Bekkers, N. Suzuki, Neurons and circuits for odor processing in the piriform cortex. *Trends*
*Neurosci* **36**, 429-438 (2013).
43. L. M. Feinberg, T. A. Allen, D. Ly, N. J. Fortin, Recognition memory for social and non-social
odors: differential effects of neurotoxic lesions to the hippocampus and perirhinal cortex.
*Neurobiol Learn Mem* **97**, 7-16 (2012).
44. Y. Shu, A. Hasenstaub, D. A. McCormick, Turning on and off recurrent balanced cortical activity.
*Nature* **423**, 288-293 (2003).
45. X. G. Li, P. Somogyi, J. M. Tepper, G. Buzsaki, Axonal and dendritic arborization of an
intracellularly labeled chandelier cell in the CA1 region of rat hippocampus. *Exp Brain Res* **90**,
519-525 (1992).
46. T. Klausberger, P. Somogyi, Neuronal diversity and temporal dynamics: the unity of hippocampal
circuit operations. *Science* **321**, 53-57 (2008).
47. B. Dudok *et al.*, Recruitment and inhibitory action of hippocampal axo-axonic cells during
behavior. *Neuron* **109**, 3838-3850 e3838 (2021).
48. M. C. Chiang, A. J. Y. Huang, M. E. Wintzer, T. Ohshima, T. J. McHugh, A role for CA3 in social
recognition memory. *Behav Brain Res* **354**, 22-30 (2018).
49. K. Tao *et al.*, Disrupted social memory ensembles in the ventral hippocampus underlie social
amnesia in autism-associated Shank3 mutant mice. *Mol Psychiatry* **27**, 2095-2105 (2022).
50. X. Deng, L. Gu, N. Sui, J. Guo, J. Liang, Parvalbumin interneuron in the ventral hippocampus
functions as a discriminator in social memory. *Proc Natl Acad Sci U S A* **116**, 16583-16592
(2019).
51. J. A. Wei *et al.*, Amygdala neural ensemble mediates mouse social investigation behaviors. *Natl*
*Sci Rev* **10**, nwac179 (2023).
52. T. B. Franklin *et al.*, Prefrontal cortical control of a brainstem social behavior circuit. *Nat Neurosci*
**20**, 260-270 (2017).
53. A. A. Shah, D. Treit, Excitotoxic lesions of the medial prefrontal cortex attenuate fear responses in
the elevated-plus maze, social interaction and shock probe burying tests. *Brain Res* **969**, 183-194
(2003).
54. M. A. Puchades, G. Csucs, D. Ledergerber, T. B. Leergaard, J. G. Bjaalie, Spatial registration of
serial microscopic brain images to three-dimensional reference atlases with the QuickNII tool.
*PLoS One* **14**, e0216796 (2019).
55. I. Mesic *et al.*, Double Dissociation of the Roles of Metabotropic Glutamate Receptor 5 and
Oxytocin Receptor in Discrete Social Behaviors. *Neuropsychopharmacology* **40**, 2337-2346
(2015).
56. B. Rein, K. Ma, Z. Yan, A standardized social preference protocol for measuring social deficits in
mouse models of autism. *Nat Protoc* **15**, 3464-3477 (2020).

**Acknowledgments**
We thank Wen-Hsien Hou, Chihiro Nakamoto, Ana Cicvaric, Hui Zhang, and Elizabeth Wood for
constructive discussion of this study. We thank Peter Bjerge, Bjark B. Brix, and Dennis Olesen
for technical help. We acknowledge the bioimaging core facility, Health, Aarhus University,
Denmark, for the use of equipment and support.
**Funding:** LF professorships, R310-2018-3611 (JR)
LF experiment, R436-2023-471 (NY)
LF experiment, R436-2023-855 (AT)
654

655      **Author contributions:**
656            Conceptualization: AT
657            Methodology: AT, NY, BMM
658            Investigation: AT, NY, HL, SJF, BMM, MZ, MM
659            Supervision: AT, NY, JR
660            Writing—original draft: AT, NY
661            Writing—review & editing: AT, NY, JR
662

663      **Competing interests:**
664      The authors declare no competing interests.
665

666      **Data and materials availability:**
667      All data needed to evaluate the conclusions in the paper are present in the paper and/or
668      Supplementary Materials.
669

670      **Figures and Tables**
671

672
673

**Figure 1. Endopiriform projects to the ventral CA1**
(**A**) Schematic diagram of a retrograde tracer injection performed. CAV-2-Cre or CTB647 was
used in combination with Ai14 or wild-type mice, respectively. (**B**) An example fluorescent
image of injection site. (**C**) Distribution of retrogradely labeled vCA1-projecting neurons in entire
brain as a result of injection in panel B. (**D-G**) Fluorescent images showing retrogradely labeled
vCA1-projecitng neurons in septum (SP), entorhinal cortex (ENT), basolateral amygdala (BLA),
and endopiriform (EN). Str: Striatum, CLA: Claustrum. (**H**) Box plots showing relative number
of vCA1-projecting neurons in different brain areas. Cell number in each brain area were
normalized by summed cell number of EN, BLA, SP (lateral), and SP (medial). Median value for
EN, BLA, SP (lateral), and SP (medial) is 0.45, 0.24, 0.08, and 0.21, respectively. Statistics: EN
vs BLA, SP (lateral), or SP (medial), p < 0.001 for all; BLA vs SP (lateral) or SP (medial): p =
0.002 and 0.694, respectively; SP (lateral) vs SP (medial) p = 0.023. One-way ANOVA, repeated
comparison with Tukey-Kramer test. N = 5. (**I**) An example fluorescent image of horizontal
section showing retrogradely labeled vCA1-projecting neurons in EN. A: anterior, P: posterior.

**(J)** Top: Example map showing distribution of vCA1-projecting neurons in entire EN in one mouse. Bottom left: A plot showing a quantified distribution at bin size of 1 mm. Each bin was normalized to total number of labeled neurons in EN. Data from each mouse is shown in grey. Median is indicated in black. Bottom right: Box plots comparing a number of vCA1-projecting neurons in anterior and posterior EN. Data were normalized to a total number of labeled neurons in EN. Median value for anterior vs posterior was 0.39 vs 0.54. Statistics: p = 0.008, Rank-sum.

**Figure 2. Axon branching of vCA1-projecting EN neurons**
**(A)** Schematic diagram of injection performed and two hypothesized outcomes. Yellow color indicates double-labeled neurons. PFC: Prefrontal cortex, ACC: Anterior cingulate cortex, MC: Motor cortex, ENT(dm): Dorsomedial entorhinal cortex. **(B-F)** Example fluorescent images of injection site for each region indicated. **(G-J)** Example fluorescent images of retrogradely labeled neurons in CLA complex. vCA1-projecting neurons were colored in red and cortex-projecting neurons were colored in green. **(K)** Box plots showing percentage of double-labeled vCA1-projecting neurons for each cortical injection. Medial value for PFC, ACC, MC, and ENT(dm) was 28.6, 4.6, 0, and 1.2, respectively. Statistics: PFC vs MC or ENT(dm), p = 0.015 and 0.047, respectively. One-way ANOVA with post-hoc Kruskal-Wallis test. Slice and animal number: 16 and 4 for PFC, 20 and 5 for ACC, 10 and 3 for MC, and 18 and 4 for ENT(dm). **(L)** Schematic diagram of projection pattern found in the experiment shown in A. **(M)** Schematic diagram of the intersectional approach performed to label vCA1-projecting EN neurons. **(N-O)** Example fluorescent images showing vCA1-projectig axons in different brain areas. All images were from the same mouse. **(P)** Magnified images showing vCA1-projecting axons in cortical layers. Magnified areas are indicated in **O** and fig. **S2A-C** as dashed squares. PIR: Piriform cortex

**Figure 3. EN projection pattern in vCA1**

(**A**) Schematic diagram of injection performed. (**B**) An example fluorescent image of horizontal brain section containing injection site in EN. (**C**) An example fluorescent image of ventral hippocampal slice with EN axons resulted from the injection in (**B**). Distal and proximal vCA1 where GFP signals were measured were indicated by dashed lines. (**D**) Normalized GFP signals across the layer. All signals were normalized by maximum GFP signal in the distal subregion. Median of area under the curve: distal; 11.12, proximal; 5.99. distal vs proximal, p = 0.016. Wilcoxon signed-rank test. n = 8 slices (N = 4 mice). SR: stratum radiatum, Py: pyramidal layer.

**Figure 4. EN$^{vCA1-proj.}$ neurons target vCA1 interneurons**

(**A**) Schematic diagram of injection performed for monosynaptic rabies tracing. (**B**) Location of starter cells in the hippocampus. Data from 5 brains were overlaid. (**C**) Example images showing presynaptic cells in EN (white arrow). (**D**) Top: Location of presynaptic EN neurons. Data from 5 brains were overlaid. Bottom: Plots indicate distribution of presynaptic cells in EN along AP axis. Cell counts in a given bin (1 mm) were normalized by total cell number in EN. Thin gray lines indicate individual data. A thick black line indicates median. Inset showing plot comparing cell number in anterior and posterior location as in Fig. 1J. Median: anterior vs posterior, 0.25 vs 0.72, p = 0.031; Wilcoxon signed-rank test. (**E**) Top: Schematic diagram of injection performed. Bottom: An example fluorescent image of injection site in EN. (**F**) Schematic diagram of recordings performed. Recordings were performed from identified GABAergic neurons sequentially in a random order. (**G**) An example confocal image of an acute vCA1 slice used for

recordings. Recorded GABAergic neurons were filled with biocytin and post-labeled with
Alexa488. White dashed lines indicate border of Py-SR and SR-SLM. (**H**) Median traces of
photo-evoked EPSCs recorded from neurons at different laminar position. (**I**) Connection
probability (CP) of EN axons and recorded neurons at different laminar positions. Numbers in
bars indicate number of response positive neurons /total recorded neurons. (**J**) Left: Peak
amplitudes of evoked EPSCs (normalized to total input from recorded slice). Right: the same as
left but for mean amplitude. Gray circles indicate each neuron. Black lines indicate median.
Median (peak amp): Py-SR vs SR vs SR-SLM vs SLM: 1.1 vs 1.0 vs 1.1 vs 0.0. [Py-SR,SR,
SR/SLM vs SLM], p = 0.076, 0.111, 0.04; one-way ANOVA, repeated comparison with Tukey-
Kramer test. Median (mean amp): Py-SR vs SR vs SR-SLM vs SLM: 1.0 vs 0.8 vs 1.4 vs 0.0.
[Py-Sr, SR, SR-SLM vs SLM], p = 0.14, 0.251, and 0.114; one-way ANOVA, repeated
comparison with Tukey-Kramer test. N = 5 mice.

**Figure 5. EN disynaptically inhibits vCA1 pyramidal neurons**
 (**A**) A confocal image of recorded pyramidal neurons in vCA1. Recorded pyramidal neurons
were filled with biocytin and post-labelled with Alexa488. Ori.: oriens, Super.: superficial layer
pyramidal neuron. Deep: deep layer pyramidal neuron. (**B**) Example traces of photo-evoked
EPSCs and IPSCs in superficial (black trace) and deep (gray trace) pyramidal neurons. Red trace:
after TTX and 4AP. Blue trace: after gabazine. (**C**) Scatter plots of peak amplitudes of EPSCs and
IPSCs for each neuron. Dashed line indicates a unitary line. Superficial neurons: empty Black
circles. Deep neurons: Gray circles. N = 4 mice. (**D**) A summary diagram of EN→vCA1 circuit.
In vCA1, EN axons innervate GABAergic neurons (IN), which in turn inhibits pyramidal neurons
(both superficial and deep).

766
767
**Figure 6. EN^vCA1-proj. neurons receive inputs from the piriform cortex and within EN**
(**A**) Schematic diagram of performed. (**B**) Location of starter cells in EN. Data from 6 brains were
overlaid. (**C**) An example map of presynaptic cells in one brain. Top: Horizontal view. Injections
were made in left hemisphere. Bottom: Front view. (**D**) Number of presynaptic cells in different
brain areas (normalized to total presynaptic cell number). Median for AIA, AON, PIR,
ENT(lateral), CLA, EN, Pallidum, and SI (in fraction): 0.7, 0.3, 3.8, 0.3, 0.5, 1.7, 0.3, and 0.2.
[AON, ENT(lateral), CLA, Pallidum, SI] vs PIR: p = 0.016, 0.008, 0.048, 0.006, and 0.003. EN
vs SI, p = 0.043. one-way ANOVA with post-hoc Kruskal-Wallis test. Full list of statistical tests
and p values are in Table1. (**E**) An example fluorescent image of coronal slice containing EN and
PIR and labeled presynaptic cells. (**F**) Example maps of presynaptic cells in PIR (top) and EN
(bottom). (**G**) Number of presynaptic cells in PIR in different layers (normalized to the total
number of labeled neurons in PIR). Median: L1 vs L2 vs L3, 0.11 vs 0.61 vs 0.27. L1 vs L2, p <
0.001; L1 vs L3, p = 0.005; L2 vs L3, p < 0.001; one-way ANOVA, repeated comparison with
Tukey-Kramer test. (**H**) Summary diagram showing input-output circuit of EN^vCA1-proj. neurons
(red circle).
783

784
785
**Figure 7. Activity of EN$^{vCA1\text{-}proj.}$ neurons predicts social/object discrimination performance**
(**A**) Right and Middle: Schematic diagrams of injection and optic cannula implantation
performed. Left: An example image showing jGCaMP8s expressing EN$^{vCA1\text{-}proj.}$ neurons and a
tract of optic cannula. (**B**) The diagram of social recognition memory test. A test consisted of
pretest (5 min), sociability test (10 min), and discrimination test (5 min). (**C**) Left: An example
image of a mouse with nose point marker tracked by DLC. Right: An example map of nose points
and calcium events in the arena during sociability test. (**D**) Nose distance vector (distance
between subject's nose point and a center point of novel/familiar mouse chambers) and
simultaneously acquired calcium signals during discrimination test. Gray dashed lines indicate the
border of interaction zones. Open red circles indicate the calcium event detected (threshold: Z >
2.58). A part of calcium signal trace was expanded at the bottom for clarity. (**E**) Left: Correlation
between calcium event ratios (calcium event number in the mouse interaction zone/ calcium event
number in the object interaction zone) and sociability index. p = 0.29, N = 8 mice. Right: same as
in left panel, but for calcium event ratios (calcium event number in the unfamiliar mouse
interaction zone/ calcium event number in the familiar mouse interaction zone) and discrimination
index. p = 0.003, N = 8. (**F**) The diagram of novel object recognition memory test. A test
consisted of habituation (5 min), familiarization (10 min), and discrimination test (5 min). (**G**)
Same as in (**E**), but for calcium event number ratios (calcium event number in the unfamiliar
object interaction zone/calcium event number in the familiar object interaction zone and
discrimination index. p = 0.193. N = 5 mice.
806

A

B

vHP-proj.
neurons

Str

0.5 mm

Str

vHP-proj.
neurons

EN   vCA1

flox-hM4Di
-mCherry    CAV2
-Cre

1 mm

C

CNO or
Saline (i.p.) → 30 m → Session 1 ⋯⋯ 1 day ⋯⋯ CNO or
Saline (i.p.) → 30 m → Session 2

Habituation → Pre test → 5 m → Sociabi. test → 30 m → Discrim. test

Object

Mouse
A    Novel
object

Mouse
A    Mouse
B

D

Sociabi.test: hM4Di mouse (#2131)

Saline              CNO

Mouse
A    Novel
object    Mouse
A    Novel
object

E

Sociability index

n.s.       n.s.

hM4Di
(N = 14)    tdTom.
(N = 8)

Saline  CNO    Saline  CNO

F

Discrim.test: hM4Di mouse (#2131)

Saline              CNO

Mouse
A    Mouse
B    Mouse
A    Mouse
B

G

Discrimination index

*       n.s.

hM4Di
(N = 14)    tdTom.
(N = 8)

Saline  CNO    Saline  CNO

H

CNO or
Saline (i.p.) → 40 m → Session 1 ⋯ 1 day ⋯ CNO or
Saline (i.p.) → 40 m → Session 2

Habituation   Familiarization   5 m   1 h   test

I

novel object recog. index

*

hM4Di
(N = 5)

Saline  CNO

**Figure 8. Chemogenetic inhibition of EN$^{vCA1-proj.}$ neurons specifically impairs recognition memory**

(**A**) Schematic diagram of injection performed. (**B**) Left: An example fluorescent image showing expression of hM4Di in EN$^{CA1-proj.}$ neurons. Right: Expansion of areas marked by white rectangle in image on the left. (**C**) The diagram of social recognition memory test. A session consisted of habituation (5 min), Pretest (5 min), Sociability test (10 min), and discrimination test (5 min). The group treated with a saline in session 1 was treated with CNO (1mg/ kg) in session 2 the following day. (**D**) Example cumulative times heat map in arena during sociability tests for the same mouse under two different treatments. (**E**) Plots comparing sociability index for hM4Di and tdTomato mice under two different treatments. Gray empty circles indicate each mouse. Gray thin lines show paired data from one mouse. Thick black bars indicate median. Median sociability index: hM4Di mice (Saline vs CNO), 0.75 vs 0.76, p = 0.391; signed-rank test; tdTomato mice (Saline vs CNO), 0.77 vs 0.75, p = 0.383; signed-rank test. (**F**) Example cumulative time heat maps in arena during discrimination tests for the same mouse under two different treatments. (**G**) Comparison of discrimination index for hM4Di and tdTomato mice under two different treatments. Gray empty circles indicate each mouse. Gray thin lines show paired data from one mouse. Thick black bars indicate median. Median discrimination index: hM4Di mice (Saline vs CNO), 0.66 vs 0.57, p = 0.042; signed-rank test; tdTomato mice (Saline vs CNO), 0.65 vs 0.63, p = 0.742; signed-rank test. (**H**) The diagram of novel object recognition memory test. A session consisted of habituation (5 min), familiarization (10 min), and discrimination test (5 min). CNO treatment was the same as in **C**. (**I**) Same as in **G**, but for novel object discrimination index. Median discrimination index: hM4Di mice (Saline vs CNO), 0.64 vs 0.5, p = 0.003; signed-rank test.

**Figure 9. The role of EN$^{vCA1-proj.}$ neurons in recognition memory. (A)** Summary of findings from *in vivo* experiments. (**B**) The model of EN$^{vCA1-proj.}$ circuit function in recognition memory. During memory-guided exploration, recurrent circuits in EN$^{vCA1-proj.}$ neurons maintain the attentional response to novel stimuli while regulating the degree of response to familiar stimuli in vCA1 via feedforward inhibition.

**Table 1. Related to Fig.6D**

| Compared brain areas | | p value |
|---|---|---|
| Agranular insular | AOB | 0.764 |
| Agranular insular | PIR | 0.617 |
| Agranular insular | ENT | 0.624 |
| Agranular insular | CLA | 0.921 |

| Agranular insular | EN | 0.971 |
|---|---|---|
| Agranular insular | Palidum | 0.588 |
| Agranular insular | Substantia innominata | 0.44 |
| AOB | PIR | 0.016 |
| AOB | ENT | 1 |
| AOB | CLA | 1 |
| AOB | EN | 0.156 |
| AOB | Palidum | 1 |
| AOB | Substantia innominata | 1 |
| PIR | ENT | 0.008 |
| PIR | CLA | 0.048 |
| PIR | EN | 0.994 |
| PIR | Palidum | 0.006 |
| PIR | Substantia innominata | 0.003 |
| ENT | CLA | 1 |
| ENT | EN | 0.091 |
| ENT | Palidum | 1 |
| ENT | Substantia innominata | 1 |
| CLA | EN | 0.317 |
| CLA | Palidum | 0.999 |
| CLA | Substantia innominata | 0.992 |
| EN | Palidum | 0.079 |
| EN | Substantia innominata | 0.043 |
| Palidum | Substantia innominata | 1 |

## Supplemental figures

**Fig. S1. Whole brain distribution of vCA1-projecing neurons**

(**A**) An example distribution map of vCA1-projecting neurons in SP (lateral and medial), BLA and ENT (lateral and medial) shown from front and top view. Neurons in different brain areas are color-coded. (**B**) The side view of the brain shown in Fig. 1J. Plots indicate distribution of vCA1-projecting neurons in EN along dorsal-ventral axis (DV). Data from different mice are shown in gray. Median is indicated in black.

**Fig. S2. Axon branching of vCA1-projecting EN neurons**

(**A-C**) Example fluorescent images of branching of vCA1-projecting neurons. All images were from the same mouse shown in Fig. 2N-O.

**Fig. S3. Projections of EN^vCA-proj. neurons to the hippocampus**

(**A**) Top: An example fluorescent images of intermediate hippocampal sections from the same brain as in Fig. 3B and C. Bottom: GFP signal profile at the distal and proximal region indicated in the top image. (**B**) The same as in (**A**) but for dorsal hippocampus. Median area under the curve (AUC): distal vs proximal (intermediate), 6.3 vs 3.5, p = 0.007; distal vs proximal (dorsal), 4.3 vs 3.0, p = 0.055; Wilcoxon signed-rank test. n = 8 slices (N = 4 mice). (**C**) Comparison of GFP signal (in AUC) between different hippocampal axis for distal or proximal vCA1. Distal; ventral vs intermediate, p = 0.086; ventral vs dorsal, p < 0.001; intermediate vs dorsal, p = 0.136. Proximal: ventral vs intermediate, p = 0.018; ventral vs dorsal, p = 0.001; intermediate vs dorsal, p = 0.695; Kruskal-Wallis test. n = 8 slices (N = 4 mice).

**Fig. S4. Presynaptic neurons of ventral hippocampal GABAergic neurons**

(**A**) Example fluorescent images of starter cells in vCA1. TVA_oG expressing neurons were labeled by GFP and rabies expressing neurons were labeled by mCherry. GFP and mCherry double expressing neurons were counted as starter cells. White allows in eight panel indicate starter cells. White dashed lines indicate the Py-SR and SR-SLM border. (**B**) An example map of presynaptic cells in a whole brain. Injection was made in the left hemisphere. (**C-F**) Fluorescent images of presynaptic cells in coronal sections from the same mouse brain. Presynaptic neurons were in EN (**C**), septum(SP) (**D**), nucleus reunion (RE) (**E**), contralateral vCA3 (vCA3) (**F**), ventral tegmental area (VTA) (**F**), and lateral ENT (**F**).

**Fig. S5. EN-mediated excitatory inputs to ventral hippocampal neurons**

(**A**) Plots comparing peak amplitude of photo-evoked EPSCs in superficial and depp pyramidal neurons. Thin gray lines indicate paired data. Median peak amplitude; superficial vs deep (in pA): 11.5 vs 5.9, p = 0.625; signed-rank test. (**B**) Plots comparing peak amplitude of photo-evoked EPSCs in pyramidal and GABAergic neurons. The data from superficial and deep pyramidal neurons were pooled. n = 10 for pyramidal neurons and 11 for GABAergic neurons. N = 4 mice.

896  Median EPSC peak amplitude: Pyramidal vs GABAergic (in pA): 7.4 vs 8.0, p = 0.86. Rank-sum
897  test.
898

899
900  **Fig. S6. Presynaptic neurons of EN$^{vCA1-proj.}$ neurons**
901  (**A**) An example fluorescent image of starter cells in EN. (**B-D**) Expanded images of the area
902  marked by white square in panel A. White allows indicate starter cells. (**E**) Example fluorescent
903  images of coronal sections in series. Labeled neurons (in green) indicate presynaptic cells of
904  EN$^{vCA1-proj.}$ neurons.

**Fig. S7. Nose point location and calcium signal data during social memory test**

Maps of nose points (filled gray circles) and calcium events (filled red circles) in arena. Heat
maps of the same data (but cumulative time and calcium event are separated) were also shown for
each mouse ( N = 8). Peason's correlation coefficients with $p < 0.05$ is indicated by asterisk. All p
values were presented in table S1.

**Fig. S8. Calcium signals during exploratory behavior in open field, pretest and familiarization test.**

(**A**) Example of maps of nose points and calcium events in open field arena. All p values in a correlation map were presented in table S2. (**B**) Correlation plots of event number and cumulative time in each pixels shown in A. p < 0.001, N = 7. (**C**) Box plots comparing a calcium event number when the mouse was in the interaction zones. In pretest, the object was aligned by longer interaction time (preferred object is plotted in right side). Median event muber: Pre (left vs right), 20 vs 63.5; Sociability (mouse vs object), 92.5 vs 77.5; Discrimination (familiar vs novel), 22.5 vs 63.5. p = 0.001, 0.385, and 0.006. Signed-rank test. N = 6. (**D** and **E**) Correlation plots of calcium event number ratios and preference index in pretest (**D**) and familiarization session (**E**). pretest, p < 0.001, N 0 8; familiarization, p = 0.014, N = 5.

**Fig. S9. The effect of chemoegentic inhibition of EN$^{vCA1-proj.}$ neurons on the pretest and locomotion**

(**A**) Frequency-current (FI) curve showing the effect of CNO on action potential number generated by current injection. Recording was made from hM4Di expressing EN$^{vCA1-proj.}$ neurons before and after bath application of CNO (100 nM). Black line indicates before and red line indicates after CNO application. Inset: Traces showing spike response to 100 pA (left) and 200 pA (right) current injection. Saline vs CNO; F (1, 4.77) = 0.38, p = 0.0329, two-way ANOVA. n = 4 neurons. N = 2 mice. (**B**) Same as in Fig. 8E, but for preference index in pretest. Median preference index: hM4Di mice (Saline vs CNO), 0.5 vs 0.5, p = 0.92; signed-rank test; tdTomato mice (Saline vs CNO), 0.48 vs 0.53, p = 0.227; signed-rank test. (**C**) Absolute change of preference index before and after CNO treatment in hM4Di and tdTomato mice. Median change of preference index: hM4Di mice 0.11 (N = 14); tdTomato 0.11 ( N = 8). hM4Di vs tdTomato, p = 0.517, ranksum test. (**D**) Same as in B, but for preference index in familiarization session of novel object recognition test. Median preference index: hM4Di mice (Saline vs CNO), 0.49 vs 0.5, p = 0.28; signed-rank test (**E**) Plots comparing locomotor activity (distance moved and velocity) for hM4Di and tdTomato mice under two different treatments. Median total distance: Pretest, hM4Di mice (Saline vs CNO), 17.17 vs 19.10, p = 0.426; tdTomato mice (Saline vs CNO), 20.60 vs 17.98, p = 0.25; Sociability test, hM4Di mice (Saline vs CNO), 34.72 vs 33.51, p = 0.855; tdTomato mice (Saline vs CNO), 38.59 vs 35.76, p = 0.547, Discrimination test, hM4Di mice (Saline vs CNO), 16.55 vs 18.58, p = 0.0676; tdTomato mice (Saline vs CNO), 18.73 vs 20.60, p = 0.945, all signed-rank test. Median velocity: Pretest, hM4Di mice (Saline vs CNO), 5.98 vs 6.55, p = 0.391; tdTomato mice (Saline vs CNO), 7.14 vs 6.21, p = 0.383; Sociability test, hM4Di mice (Saline vs CNO), 6.01 vs 5.76, p = 0.808; tdTomato mice (Saline vs CMO), 6.78 vs 6.13, p = 0.313; Discrimination test, hM4Di mice (Saline vs CNO), 5.88 vs 6.57, p = 0.104; tdTomato mice (Saline vs CNO), 6.79 vs 6.99, p = 0.945. All signed-rank test.

**Fig. S10. The effect of chemoegentic inhibition of EN$^{vCA1\text{-}proj.}$ neurons on trace fear conditioning**

(**A**) Schematic of trace fear conditioning protocol. (**B**) The effect of CNO on freezing behavior during conditioning. Black indicates saline treated group and red indicates CNO treated group. Data points of individual mice are indicated by empty circles. Thick horizontal bars indicate median values. Saline vs CNO, $F(1, 9) = 2.52$, $p = 0.115$; two-way ANOVA. (**C**) The effect of CNO on freezing behavior during recall to context "A". Median values of freezing %: Saline = 14.91, CNO = 30.13. Saline vs CNO, $p = 0.576$; one-way ANOVA. (**D**) The effect of CNO on freezing behavior during conditional stimulus recall in context "B". Saline vs CNO, $F(1, 9) = 0.38$, $p = 0.54$; two-way ANOVA