Functional associations of evolutionarily recent human genes exhibit sensitivity to the 3D genome landscape and disease

Katherine Fleck[1,2*], Victor Luria[3,4,5,6*], Nitanta Garag[1], Amir Karger[7], Trevor Hunter[1], Daniel Marten[4,5], William Phu[4,5], Kee-Myoung Nam[8], Nenad Sestan[3], Anne H. O'Donnell-Luria[4,5,9#], Jelena Erceg[1,2,10#]

[1]Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT 06269

[2]Institute for Systems Genomics, University of Connecticut, Storrs, CT 06269

[3]Department of Neuroscience, Yale School of Medicine, New Haven, CT 06510

[4]Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA 02115

[5]Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA 02142

[6]Department of Systems Biology, Harvard Medical School, Boston, MA 02115

[7]IT-Research Computing, Harvard Medical School, Boston, MA 02115

[8]Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06510

[9]Department of Pediatrics, Harvard Medical School, Boston, MA 02115

[10]Department of Genetics and Genome Sciences, University of Connecticut Health Center, Farmington, CT 06030

*K.F. and V.L. contributed equally to this work.

#To whom correspondence may be addressed. Email: odonnell@broadinstitute.org or jelena.erceg@uconn.edu.

Keywords: 3D genome organization, gene evolution, human accelerated regions (HARs), gene function/ontology, cell type-specific processes

Abstract

Genome organization is intricately tied to regulating genes and associated cell fate decisions. In this study, we examine the positioning and functional significance of human genes, grouped by their evolutionary age, within the 3D organization of the genome. We reveal that genes of different evolutionary origin have distinct positioning relationships with both domains and loop anchors, and remarkably consistent relationships with boundaries across cell types. While the functional associations of each group of genes are primarily cell type-specific, such associations of conserved genes maintain greater stability across 3D genomic features and disease than recently evolved genes. Furthermore, the expression of these genes across various tissues follows an evolutionary progression, such that RNA levels increase from young genes to ancient genes. Thus, the distinct relationships of gene evolutionary age, function, and positioning within 3D genomic features contribute to tissue-specific gene regulation in development and disease.

Main Text

Introduction

The genome is intricately folded and regulated in the nucleus, which may contribute to cell type diversity. Recent technological advances in mapping of chromosomal interactions and single-cell imaging have provided insights into the organization of chromatin at various levels (1-4). Domains (also termed topologically associating domains or TADs) with increased chromatin interactions may be insulated from neighboring regions by boundaries with decreased chromatin interactions (5-8). Formation of these domains is often mediated by extrusion of chromatin loops, which dynamically link distal anchor points (6, 9-12). However, the interplay between genome structure and function is still unclear (13-16). For instance, while some chromosomal disruptions and perturbations of regulators involved in genome folding may not lead to substantial effects on gene regulation, other structural disruptions at specific loci may affect contacts between regulatory regions, leading to cancer and congenital limb malformations (13, 14, 17-22). Moreover, communication between regulatory regions associated with gene activation is facilitated by direct chromatin looping or other hub-like structures (6, 13, 14, 23-34). Having proper regulatory interactions, including those between enhancers and promoters, is key for cell type-specific gene activity.

Understanding how the genome is organized and the extent to which its organization is conserved may illuminate the functional significance of genomic regions. For instance, perfectly conserved sequences between distantly related species that diverged 90 to 300 million years ago, named ultraconserved elements (UCEs) (35-38), have non-random relationships to 3D genome organization (39). Specifically, UCEs are enriched in domains and associated with renal and

neuronal processes when present in domains that are common to many cell types (39). Clusters of conserved noncoding elements (CNEs) around developmental genes, with 70-90% CNE conservation level, are also highly concordant with domains (40). As for human accelerated regions (HARs), sequences conserved in vertebrates but fast-evolving in the human lineage (41-49), they are associated with domains that have human-specific structural variants, suggesting potential rewiring of contacts between regulatory regions (50). Moreover, structural variants and mutations affecting conserved and fast-evolving regions are often related to diseases and neurodevelopmental disorders (37, 46, 51-55).

Protein-coding genes also experience varying levels of conservation, from extremely high sequence conservation across evolutionary taxa to *de novo* evolution in specific lineages. To understand how gene expression and function emerge and change with time, genes of different evolutionary ages can be compared once the minimal evolutionary age of each gene is evaluated (56-58). Novel protein-coding genes originating by non-copying mechanisms (*de novo* from intergenic regions; from long non-coding RNAs that become coding; or by overprinting, that is, translating an existing mRNA in a different frame) have been discovered since 2006 by studying naturally occurring gene variants, through mutagenesis screens and by proteogenomics in species ranging from plants to humans (56, 59-64). Such novel genes are expressed in different tissues (65, 66), have function (59, 67-71) and, in contrast to new genes arising from copying mechanisms (gene or genome duplication; retrotransposition; exon shuffling; or pseudogene resurrection), can have novel protein domains (70-72). The process of gene birth and death is continuous, and some genes are specific to particular lineages (73).

Here, we asked how the positionings of genes of different evolutionary ages in the 3D genome landscape relate to their expression and functional significance. Interestingly, while genes

of different evolutionary ages have distinct relationships with domains and loop anchors, their relationships with boundaries are consistent across cell types. Furthermore, evolutionarily older genes maintain stable functional associations regardless of genomic positioning and are expressed at higher levels than more recently evolved genes, which exhibit functional associations that are sensitive to the positioning of these genes within the 3D genome organization and disease. Together, the distinct 3D genomic positioning of genes as a function of their evolutionary age may be related to the regulation of tissue-specific programs in development and disease.

Results

Regions of varying sequence conservation have distinct relationships with 3D genomic features across cell types

To understand the relationship between gene age and features of 3D genome organization, we built a dataset with all human protein-coding genes and control intergenic sequences. First, we evaluated the evolutionary age of all human protein-coding genes (Fig. 1*A* and Fig. S1). Of these 21,436 single-isoform genes (only the oldest and longest protein isoform was kept), most (19,334) are annotated in Ensembl 89 (Fig. S1*A*). Additionally, 2,102 unannotated genes with translation evidence from proteogenomics studies (74-77) were also used (Fig. S1*B*). We estimated gene evolutionary age by phylostratigraphy. This method estimates the minimal evolutionary age of a gene by determining the most distant species in which a protein sequence sufficiently similar to the query sequence is detected, and inferring that the gene must be at least as ancient as the common ancestor of the query species and the most distant species (57, 78, 79). Thus, a human gene for which a sufficiently similar sequence exists in bacteria is Ancient (14,845 genes); a human gene also present in sponges but not in more distant species is specific to Metazoan (2253 genes); a human gene also present in lancelets is specific to Chordates (1584 genes); a human gene also present in platypus but not in more distant species is specific to Mammals (685 genes); a human gene also present in lemurs is specific to Primates (2069 genes) (Fig. 1*B*). We refer to these five categories (ancient, metazoan, chordate, mammal, primate) based on the evolutionary age of genes as *eras* and genes within them as *era genes*. As controls for human protein-coding genes of varying evolutionary age, we used intergenic open reading frames (Igen ORFs, 3924 sequences) that are potential protein-coding sequences with no known evidence of transcription or translation.

Additionally, we used human intergenic sequences that cannot encode proteins (Igen Non-ORFs, 3892 sequences) and are matched in length to Igen ORFs (Table S1). We find that there is an evolutionary progression from short young genes to long ancient genes (Fig. 1 *C* and *D*), as noted previously in multiple eukaryotic species (56, 57, 80). Interestingly, the youngest genes are very similar in size to control Igen ORFs (Fig. S1 *C* and *D*), suggesting young genes are picked at random from the myriad of ORFs present in the vast intergenic parts of the human genome.

Second, we included two datasets with different levels of sequence conservation. The first dataset is comprised of UCEs, genomic regions at least 200 base pairs (bp) in length that share 100% sequence identity between reference genomes of distantly related species. Comparisons between human, mouse, and rat; or human, dog, and mouse; or human and chicken genomes identified 896 UCEs in total (39, 53, 81). The second dataset is comprised of >3,100 HARs corresponding to sequences that are conserved in vertebrates but are rapidly evolving in the human genome (41-49).

We began by visualizing the placement of era genes, UCEs, and HARs in the 3D genome landscape as illustrated with an example region of the GM12878 genome (6, 82) (Fig. 1*E*). Then, to globally assess how genomic regions of varying sequence conservation relate to features of 3D genome organization such as domains, loop anchors, and boundaries, we compiled Hi-C datasets from various published studies encompassing a wide range of cell types (Table S1) (5, 6, 19, 83-93). Due to the variation in methodology between these studies (e.g. Hi-C protocol, sequencing depth, quantity of input material), we considered each dataset individually as well as grouped by 3D genomic feature to generate pooled datasets. Then, we performed correlation analyses as previously described (39, 53). Briefly, the genome was partitioned into equal bins of defined size, within which the proportion of sequence covered by each feature of interest was determined. The

global pairwise correlations between feature densities were performed to generate Spearman

correlation coefficients and p values. These analyses were carried out using a breadth of bin sizes

(20 kb, 50 kb, and 100 kb) to account for the differences among individual 3D genomic features

in terms of their number of regions, median region size, and proportion of genome covered (Table

S1).

We assembled 18 domain datasets from six studies (5, 6, 83, 85, 86, 93) covering 14 human

cell types, including human embryonic and induced pluripotent stem cells as well as fetal cells,

differentiated and cancer cell lines representing different germ layers (Table S1). Individually,

these domain datasets cover varying percentages of the genome, ranging from 40.07% (HMEC) to

87.72% (SK-N-SH). Pooling the datasets together and collapsing the overlapping intervals resulted

in a single dataset (hereafter named "pooled domains") spanning 98.06% of the genome (Table

S1). Pooled domains show predominantly significant, positive correlation with genes of most

evolutionary eras, UCEs, and HARs ($6.27 \times 10^{-35} \leq p \leq 0.022$), except mammal genes (p = 0.654)

(Fig. 1*F* and Table S2). When examined by individual cell types, ancient genes and UCEs continue

to display mainly significant, positive correlations with domains ($0 \leq p \leq 0.746$; Fig. S2 and Table

S2). This observation is aligned with previous findings that UCEs and clusters of highly conserved

noncoding elements, which may have regulatory function, coincide with domains (39, 40).

Notably, annotated Ensembl ENSP genes also closely match the trend in correlations between

ancient genes and domains ($0 \leq p \leq 0.078$; Fig. S2 and Table S2). Interestingly, less conserved era

genes (metazoan, chordate, mammal, and primate) and fast-evolving HARs have more variable

strengths of correlations with domains across cell types ($2.34 \times 10^{-92} \leq p \leq 0.988$; Fig. S2 and Table

S2). Taken together, these observations suggest that conserved elements have more consistent

relationships with domains, while less conserved elements show more variable relationships with domains across cell types.

We continued our analyses using 36 datasets of loop anchors from various studies (6, 19, 83, 84, 87-92). These datasets span 36 distinct cell types ranging from H1 human embryonic stem (ES) cells, ES-cell-derived lineages, fetal cells, to differentiated and cancer cell lines (Table S1). Loop anchors cover a smaller portion of the genome than domains; these individual datasets range from 1.97% (RPE1) to 9.83% (aorta). Again, we pooled the loop anchor datasets together and collapsed the overlapping intervals to generate a dataset named "pooled loop anchors," which covers 56.55% of the genome (Table S1). Similarly to pooled domains, pooled loop anchors display mostly significant, positive correlations with genes of most eras, UCEs, and HARs ($1.08 \times 10^{-172} \leq p \leq 0.003$), except for mammal genes, whose correlation is significantly negative (p $= 6.03 \times 10^{-5}$) (Fig. 1F and Table S2). Notably, HARs have previously shown enrichment in loops specific to human compared to rhesus macaque and mouse (94). While the observed significant, positive correlation in UCEs differs from a former study (39), which displayed a non-significant relationship between UCEs and loop anchors, this result can be explained by the increase in the number of datasets used in this analysis (36 datasets here compared to previously used eight datasets). Our observation that UCEs are positively correlated with pooled loop anchors is consistent with the regulatory activity of some UCEs (95-103) and with loops facilitating enhancer-promoter contacts (6). When separated by cell type, UCEs together with HARs and less conserved era genes (metazoan, chordate, mammal, primate) have variable relationships with loop anchors ($8.65 \times 10^{-118} \leq p \leq 0.965$; Fig. S3 and Table S2). On the other hand, ancient and ENSP genes show a robust, significant positive correlation with loop anchors across most cell types ($0 \leq p \leq 0.944$; Fig. S3 and Table S2). In addition, we noted that relationships of genes of different eras

and HARs with loop anchors remain persistent across some cell types, mostly corresponding to Hi-C datasets produced with the same methodology (Fig. S3 and Tables S1 and S2). Overall, ancient genes have the most consistent relationships, while younger genes, HARs, and UCEs have more variable relationships with loop anchors across cell types.

Finally, we turned to boundary datasets from multiple studies (5, 84, 88, 92). These boundary datasets cover 20 cell types and primary tissues (Table S1). Individual datasets cover portions of the genome more comparable to loop anchors than domains ranging from 2.61% (aorta) to 3.95% (hESC). A dataset of pooled boundaries covers 22.5% of the genome (Table S1). This pooled dataset is significantly positively correlated with genes of different eras ($2.62 \times 10^{-136} \le p \le 2.90 \times 10^{-9}$), consistent with transcription start sites being enriched at boundaries (5). Conversely, pooled boundaries are significantly negatively correlated with control Igen ORFs ($p = 9.06 \times 10^{-18}$) and Igen Non-ORFs ($p = 2.11 \times 10^{-14}$), UCEs ($p = 1.81 \times 10^{-6}$), and HARs ($p = 1.76 \times 10^{-25}$) (Fig. 1*F* and Table S2). These observations in UCEs and HARs are different from those found with pooled domains and loop anchors (Fig. 1*F*) and are consistent with the depletion of UCEs at boundaries (39). Furthermore, in distinct contrast to domains and loop anchors, genes of different eras, UCEs, and HARs all show remarkable consistency in their correlations with boundaries across cell types (Fig. S4 and Table S2). Strikingly, genes display predominantly significant, positive correlations with boundaries ($2.19 \times 10^{-110} \le p \le 0.306$), as opposed to HARs with entirely significant, negative correlations ($8.91 \times 10^{-12} \le p \le 0.009$) and UCEs with significant or non-significant negative correlations ($6.24 \times 10^{-4} \le p \le 0.136$) (Fig. S4 and Table S2).

Additional correlation analyses with different bin sizes (20 and 100 kb; Tables S3 and S4) provide similar results to 50-kb results (Table S2) for boundaries as well as domains and loop

anchors. Moreover, the control sets of Igen ORFs and Igen Non-ORFs, with no evidence of regulated transcription, support these findings with era genes by displaying similar correlations regardless of the inspected 3D genomic feature or cell type (Fig. 1 and Figs. S2–S4 and Table S2). Taken together, the relationships between genomic regions of varying sequence conservation and boundaries are consistent yet distinct, while the relationships with domains and loop anchors are variable across cell types. This variability, more prominent with evolutionarily younger genes and fast-evolving HARs, may suggest their potential role in tissue-specific programs.

Gene expression changes with evolutionary age

Due to the interplay between the 3D organization of the genome and gene expression, we measured the transcription of genes and intergenic control sequences using the Genotype to Tissue Expression (GTEx) database, which contains RNA sequencing data from 54 human adult tissues (Table S5) (104). We grouped the data into six categories: three germ layers (ectoderm, mesoderm and endoderm), germ line (ovary, testis), and the brain, since some evolutionarily novel genes function in neurons (59, 65, 70, 105). First, we examined the RNA expression of genes as a function of their evolutionary age (Fig. 2*A* Mean Counts, and Fig. S5*A* transcripts per million (TPM)). We found that gene expression changes with evolutionary era, such that, in all tissue categories, ancient genes are expressed at the highest levels while primate genes are expressed at significantly lower levels (Tables S5 and S6; for example, Ectoderm: Ancient 2709 ± 56, Primate 118 ± 117, average ± standard deviation, $p < 2.225 \times 10^{-308}$). Moreover, the expression of all genes is 100-1000 times higher than that of control Igen ORFs and Igen Non-ORFs, which reflect the

transcriptional background of the genome (Tables S5 and S6; for example, Ectoderm: Primate 118 ± 117, Igen Non-ORF 0.23 ± 0.04, average ± standard deviation, $p < 2.225 \times 10^{-308}$).

Second, we examined the RNA expression of genes within 6 tissue categories (Fig. 2*B* and Fig. S5 *B* and *C*) and found that within each category there is a similar evolutionary progression of gene expression, such that ancient genes are expressed at higher levels than evolutionarily novel genes. Interestingly, within the testis, the expression of genes is higher than in other tissues, as observed in other species such as fruit fly (66). We observed that, for most genes of all eras, the expression of genes in the brain is similar or lower than in other tissues (Fig. S5 *B* and *C*). Within tissue categories, gene expression can vary substantially between individual tissues (Fig. S6). Within the brain, cerebellar hemispheres express the highest RNA levels of ancient genes while the amygdala and *substantia nigra* express the lowest RNA levels of ancient genes (Fig. S6). Additionally, we observed that, while the expression of genes generally increases with gene length (short genes are expressed at lower levels than long genes), within each gene length decile the evolutionary progression from ancient genes expressed at high levels to young genes expressed at low levels is mostly maintained (Fig. S5*D*). Taken together, these data show that gene expression increases with evolutionary age and that all protein-coding genes are expressed at levels much higher than the transcriptional background of the genome.

Era genes are primarily associated with prominent cell type-specific processes

Given the distinct relationships between regions of varying sequence conservation and 3D genome structure across cell types, we next investigated the functional associations of those regions. Previous work by Domazet-Lošo and Tautz associated Ensembl annotated genes grouped

by their evolutionary age with specific gene ontology (GO) terms (106). Due to the complexity of our datasets, which comprise recently curated annotated genes and as yet unannotated genes, we used the Genomic Regions Enrichment of Annotations Tool (GREAT) (107) to assess GO term enrichments in genomic coordinates of interest. This approach allowed inspection of different regions including HARs, UCEs, and unannotated genes, particularly primate genes, without a need to rely on gene identifiers. We observed that metazoan genes are enriched for GO terms related to signaling pathways against all era genes as a background (Fig. S7*A*). Meanwhile, chordate genes are associated with immune response-related processes (Fig. S7*B*). As for mammal and primate genes, both are linked to keratinization and keratinocyte differentiation processes (Fig. S7 *C* and *D*). However, mammal genes are also associated with defense response to other organisms (Fig. S7*C*). Ancient genes do not show enrichment in GO terms, most likely since they form a majority (69.25%) of all era genes. Together, these observations are consistent with a previous report (106) and suggest that various categories of era genes are predominantly associated with distinct cell type-specific GO terms.

Evolutionarily older regions show stable functional associations within 3D genomic features

Having the corresponding functional associations of era genes, we examined whether these associations are related to positioning of era genes within 3D genomic features. Unsurprisingly, era genes in pooled domains are associated with the same GO terms (Fig. 3 *A-D*) as with the entire genome (Fig. S7). Pooling the domains across cell types may include regions that are domains in certain cell types but are boundaries and loop anchors in other cell types, and therefore incorporates more than 99% of the genes. In pooled loop anchors, metazoan and chordate genes still maintain

very similar enrichment of GO terms (Fig. S8 *A* and *B*) as noted with genes in pooled domains (Fig. 3 *A* and *B*). On the other hand, evolutionarily younger mammal and primate genes lose association with defense response to other organisms and peptide cross-linking, respectively (Fig. S8 *C* and *D*). As for pooled boundaries, we observed a notable shift in the gene eras that exhibit GO term enrichment. For instance, ancient genes in pooled boundaries appear mainly associated with protein localization to telomeres as well as various ncRNA processes (Fig. S9*A*), consistent with the enrichment of housekeeping genes at boundaries (5). Meanwhile, metazoan and chordate genes in pooled boundaries show stable association with signaling pathways and immune response-related processes, respectively (Fig. S9 *B* and *C*). Interestingly, evolutionarily recent mammal and primate genes in pooled boundaries do not display any GO term enrichments. Overall, the functional associations of older metazoan and chordate genes seem to be stabilized and not affected by their positioning within 3D genomic features. In contrast, functional associations of younger mammal and primate genes may change depending on 3D genomic feature these genes overlap, suggesting sensitivity to their positioning within genome architecture.

We next compared our functional associations of era genes in various 3D genomic features with those of UCEs and HARs. Previous studies suggest that specific subsets of UCEs are associated with RNA processing and developmental processes (35, 39, 108-111). In line with this, we observed that all UCEs in pooled domains, boundaries, or loop anchors maintain consistent stable enrichment for GO terms related to transcription and RNA metabolic processes (Figs. S8*E*, S9*D*, and S10*A*) (35, 110, 112). HARs in pooled domains, which comprise all but one HAR in our dataset, are associated with neuronal processes, cartilage formation, and thymus development (Fig. S10*B*), consistent with functional implications of HARs due to their increased divergence in humans (42-44, 46-48, 50, 52). As for HARs in pooled loop anchors, associations are minorly

altered to those with HARs in pooled domains (Fig. S8*F*). In pooled boundaries, HARs do not show enrichment for GO terms, which corresponds to the strong negative correlation between HARs and pooled boundaries (p = $1.76{\times}10^{-25}$; Fig. 1D). Together, these results point to sensitivity in positioning of HARs within 3D genome architecture similar to evolutionarily recent mammal and primate genes.

The functional associations of regions of varying sequence conservation are more sensitive to disease state in loop anchors than in domains

Due to potential changes in chromatin folding and gene regulation in disease states (17, 20, 113), we next asked whether healthy and diseased cancer states can impact the positioning and the functional associations of regions of varying sequence conservation within 3D genomic features. Correlations of pooled domains with metazoan, mammal, and primate genes as well as UCEs change from nonsignificant in one state to significant in the other ($2.29{\times}10^{-7} \leq p \leq 0.870$; Fig. 4*A* and Table S2). Despite this variation in 3D positional relationships, the functional associations of those regions seem mainly unaffected. Broadly, era genes, UCEs, and HARs in both pooled healthy domains (Fig. S11) and pooled cancer domains (Fig. S12) are enriched for very similar GO terms, as seen in all pooled domains (Fig. 3 and Fig. S10). For pooled loop anchors, we observed changes in the correlations for chordate and mammal genes along with HARs between healthy and cancer states (Fig. 4*A*). As for functional associations, although GO terms associated with era genes, UCEs, and HARs in pooled healthy loop anchors (Fig. 4*B* and Fig. S13) closely resemble those found with all pooled loop anchors (Fig. S8), GO term enrichments in pooled cancer loop anchors can show notable shifts depending on the class of region (Fig. S14). For

instance, older metazoan and chordate genes and UCEs in pooled loop anchors have largely similar

GO term enrichments in cancer (Fig. S14 *B*, *C*, and *D*) and healthy (Fig. S13 *A*, *B*, and *E*) states,

potentially suggesting the exploitation of signaling pathways by cancer cells (114). In contrast,

younger mammal and primate genes in pooled cancer loop anchors show no GO term enrichments.

As for HARs in pooled loop anchors, we noted changes in GO term enrichments between healthy

(Fig. 4*B*) and cancer states (Fig. 4*C*). While both states are still associated with neuronal-related

processes, the individual GO terms are often state-specific. These diverse associations of HARs

are in line with their implications in human neurodevelopment and disease (115, 116).   In

summary, the functional associations of evolutionarily recent mammal and primate genes in pooled

loop anchors seem to be lost in disease, suggesting that cancer cells may better exploit the stable

functional associations of older genes.

Discussion

Our study reveals that the relationships between genes of different evolutionary ages and

their positioning within domains and loop anchors are variable, while the correlations of those era

genes with boundaries display remarkable consistency across cell types. The transcription levels

of genes also increase with increasing evolutionary age in all tissue categories (Fig. 4*D*). As for

the functional associations of era genes, they are predominantly related to distinct cell type-specific

processes. The functional associations of more recently evolved genes as well as HARs are

sensitive to their 3D genomic positioning. This sensitivity is especially prominent in loop anchors

compared to domains in disease state (Fig. 4*E*). Loop formation has been implicated in

demarcating domains and facilitating contacts between enhancers and promoters (6, 9, 12).

Although domain and loop structures may be changing, at least half of them may be preserved across cell types (5, 6, 117). Ancient genes exhibit more consistent positive correlations with both domains and loop anchors between cell types, while less conserved, younger genes, with their marked increase in correlation variability, may be linked to cell type-specific chromatin organization. Furthermore, evolutionarily younger genes are associated with cell type-specific functions suggesting their potential role in tissue-specific programs. Interestingly, previous studies have shown that many loops involving enhancers and promoters may be tissue-specific during differentiation, and therefore related to cell type-specific gene regulation (118, 119). Hence, the relationships of these recent era genes with 3D genomic features may be implicated in evolution of tissue-specific genome activity. Moreover, modifications to the dynamic 3D regulatory landscape may even shape species-specific programs (120-125).

Notably, measuring transcriptional output with GTEx shows an evolutionary progression of gene expression: young genes are expressed at lower levels than ancient genes, yet at much higher levels that background genomic transcription. The output of gene regulation by way of using various gene regulatory elements, including 3D genome organization elements, is gene expression. Thus, the placement of 3D genome organization elements may be used to control gene expression at the RNA level, as transcriptional complexes must access DNA to produce RNA. Interestingly, for genes of all ages in the six tissue categories we considered, RNA levels are highest in testis, a tissue where DNA is loosely packed with protamines instead of histones (126), and are unexpectedly lowest in the brain, where gene expression is tightly regulated but there are many cell types. Indeed, some brain areas such as the cerebellum do have high RNA expression. The similar evolutionary progression of RNA levels in all tissues suggests RNA levels are regulated by the 3D genome organization elements that may be implicated in opening chromatin,

and thus in part independently of transcription factors – while a few transcription factors, like CTCF (22), change genome organization, most transcription factors regulate expression differently, by cooperatively aiding the assembly of the RNA polymerase II transcriptional complex. These two facets of gene regulation support transcription of a new gene, and are stabilized evolutionarily if the gene finds a function and perdures evolutionarily. Additionally, longer genes of all ages are expressed at higher levels than shorter genes of all ages, suggesting gene size, usually higher in neuronal genes (80, 127), is associated to gene expression level independently of gene age.

The functional associations found with genes of each era are intriguing in light of the biological processes that arose at different times in evolution (78, 106). In line with this notion, the SMAD-dependent and G-protein coupled receptor signaling pathways associated with metazoan genes are often involved in altering gene expression related to tissue differentiation, which is critical for development of multicellular animals (128). Chordate genes as well as mammal genes link to several immune processes which may mediate defense responses against pathogens (129, 130). Both mammal and primate genes are associated with keratin processes, possibly related to hair formation which may be integral to heat retention and homeothermy (131). Interestingly, these links to immune response and skin development are further supported by the functional associations of fast-evolving genes and regulatory landscape in mammals facilitating an organism's interactions with the environment (51, 132). Overall, the emergence of genes at different evolutionary times may be linked to novel biological processes gained at different levels of the phylogenetic hierarchy (78, 106).

Disruptions of the 3D genome architecture can play roles in various diseases, especially when interactions between regulatory regions are affected (13, 14, 17, 20, 119). In particular, the

identity and evolutionary age of the genes impacted by such alterations may be integral to understanding a disease. For instance, previous reports indicate a significant over-representation of disease-associated genes, including cancer-associated genes, that predate or originate with the emergence of Metazoa (106, 114). Interestingly, we observed little change in the functions associated with metazoan genes in loop anchors between healthy and cancer states, possibly related to the exploitation of signaling pathways by cancer cells in multicellular organisms (114). As for evolutionarily recent mammal and primate genes in loop anchors, both lose all functional GO term enrichments in cancer, consistent with the under-representation of such genes among cancer and other disease-associated genes (106, 114). Nevertheless, fast-evolving HARs in loop anchors maintain functional associations with neuronal processes in both healthy and cancer states, reminiscent of the other disease implications of HARs, including in neurological disease (115, 116).

In conclusion, we describe the functional significance of the 3D genomic positioning of genes of different evolutionary ages. The evolution of genes and their regulatory regions within the 3D genome may present different lenses through which the determinants of yet unresolved genetic diseases may be elucidated.

Materials and Methods

Human sequence datasets

As in other studies (80, 133), we retrieved 19,334 human annotated proteins from the Ensembl 89 database (www.ensembl.org), keeping the oldest and longest isoform per protein-coding gene. Additionally, we used 2,102 human unannotated protein sequences that are

proteogenomically confirmed (74-77). Furthermore, we used two kinds of control sequences. The first are Igen ORFs, localized in the non-genic non-repetitive part of the human genome, starting with a methionine (Met) codon and having at least 40 non-STOP codons, followed by a STOP codon. These sequences have no known evidence of transcription or translation but could in principle encode proteins, based on their DNA sequence. The second type of controls are intergenic DNA sequences that are random sequences from the non-repetitive part of the human genome that do not overlap with any selected ORFs or known protein-coding genes, do not start with Met, may have STOP codons and are not expected to encode open reading frames (Igen Non-ORFs). Igen Non-ORFs were chosen to have the same DNA sequence length distribution as Igen ORFs.

Gene age estimation

We estimated the evolutionary age of protein-coding genes by phylostratigraphy (78, 79), which classifies genes based on their taxonomic restriction (80). For genes encoding proteins of minimum 40 amino acids, the protein sequence was used to query the NCBI NR database (February 2017) and find the most taxonomically distant species in which a sufficiently similar protein sequence exists, such that the e-value of that most distant BLASTP hit is less that 10^{-3} (80). Sequence similarity was evaluated with BLASTP (134) and HMMER (135). Minimal gene age was assessed as the time when the most recent common ancestor of the query species and the most distant species lived. Within the human lineage, there are 31 taxonomical nodes, corresponding to 31 phylostrata (PS; NCBI Taxonomy, https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=9606), and their timing was estimated using TimeTree (136). The 31 phylostrata were grouped into 5 large-scale evolutionary

eras: Ancient (PS 1-3), Metazoan (PS 4-7), Chordate (PS 8-17), Mammal (PS 18-22), Primate (PS 23-31). Human genes also present in bacteria are in PS 1 and belong to the Ancient era, while human genes present only in humans are in PS 31 and thus belong to the Primate era. For each gene, only the oldest and longest protein isoform was kept.

UCE and HAR datasets

The UCE dataset includes 896 elements from comparisons between human-mouse-rat (HMR), human-dog-mouse (HDM), or human-chicken (HC) genomes as previously reported (39, 53, 81). The genomic coordinates for HARs were obtained from Girskis *et al*. (43) and converted to the hg19 genome assembly using the UCSC LiftOver tool (https://genome.ucsc.edu/cgi-bin/hgLiftOver). Any overlapping intervals were collapsed to avoid including the same region more than once.

Hi-C datasets

The coordinates for Hi-C annotated genomic features, including domains, boundaries, and loop anchors, were retrieved from published datasets as indicated in Table S1 (5, 6, 19, 83-93). The genomic coordinates were processed as previously described (39). Namely, when applicable, the Hi-C genomic annotations were lifted over to the hg19 genome assembly using LiftOver. To ensure that any given region was counted only once, overlapping intervals were collapsed to generate final pooled and individual datasets for domains, boundaries, and loop anchors that may differ from previously published original ones. Information about these final datasets, including number of regions, median region size (bp), coverage (bp), and proportion of genome covered (%), is provided in Table S1. Eres *et al*. (85) provided four iPSC datasets derived from four individuals,

while Dixon *et al*. (5) and Rao *et al*. (6) each provided one dataset of IMR90 using dilution and in situ Hi-C protocols, respectively. Rubin *et al*. (90) provided three Hi-C datasets corresponding to different stages of keratinocyte differentiation (days 0, 3, and 6). Dixon *et al*. 2012 (5) and Dixon *et al*. 2015 (84) each provided a Hi-C dataset for H1 human embryonic stem cells. Due to the methodological variations among these studies such as Hi-C protocol, quantity of input material, and sequencing depth, we examined each dataset individually, grouped by a Hi-C feature (e.g. Pooled Domains), and grouped additionally based on healthy or disease state (e.g. Pooled Cancer Domains).

Correlation analyses between 3D genomic features and regions of varying sequence conservation

Correlation analyses were performed using a previously established method (39, 53). Briefly, the genome was first split into bins of fixed size. In each bin, the proportion of sequence covered by a 3D genomic feature or regions of varying sequence conservation was calculated. Then, the global correlations among feature densities of interest were determined. The Spearman correlation coefficients and the corresponding p values were reported for a pairwise comparison between two features (e.g. between era genes, UCEs, or HARs and a 3D genomic feature) using three different bin sizes (20 kb, 50 kb, 100 kb; Tables S2, S3, and S4). The Spearman correlation coefficients were also indicated by a heatmap.

GTEx analysis

To determine the level of RNA expression of all human protein-coding genes encoding proteins of at least 40 amino acids and of control intergenic sequences, a RNA-Seq calling pipeline was used to calculate the number of reads per gene per sample broken down by tissue. GTF files

with all sequences (annotated and unannotated Genes, Igen controls) were run through the GTEx RNA-Seq calling pipeline of the Broad Institute (https://github.com/broadinstitute/gtex-pipeline). The control intergenic sequences inform on the background transcription of the human genome. Deseq2 normalization of means was used (https://github.com/broadinstitute/pyqtl) to account for sample-level sequencing depth between individuals and RNA composition. From this, we obtained a mean read count value for each sequence for each of the 54 tissues of GTEx. These tissues were aggregated into the following tissue categories: three germ layers (ectoderm, mesoderm, endoderm), two germline tissues (ovary, testis) and one tissue of special interest to the authors (brain). To avoid overweighting brain regions in ectoderm, only cortex and cerebellum were included (and these tissues were also included in brain, along with 11 other brain regions). Similar analyses were conducted with transcripts per million (TPMs) per gene, as available from the online resource of GTEx for annotated genes alone (v8, median TPM counts per gene). TPM data was downloaded directly from the GTEx v8 Browser's download page (https://storage.googleapis.com/adult-gtex/bulk-gex/v8/rna-seq/GTEx_Analysis_2017-06-05_v8_RNASeQCv1.1.9_gene_median_tpm.gct.gz), with median TPM counts per 54 tissues per Ensembl annotated gene. The purpose of the TPM analysis was checking if the changes with evolutionary age observed with Mean Counts remain the same with TPMs, which better accounts for transcript length but does not work as well for small genes.

Transcripts of human genes encoding proteins of at least 40 amino acids (19,334 annotated protein-coding genes, 2,102 unannotated proteogenomically confirmed genes) and control sequences were counted in 54 human adult tissues (GTEx database) using GRCh38 genomic coordinates for all sequences. Of 20,000 randomly picked Igen ORFs (ten sets of 2,000 each), for our analysis we used two sets of 2,000 each (4,000 sequences total), from which 3,916 Igen ORFs

passed GTEx filtering consisting of not duplicating or overlapping other Igen ORFs, and cleanly mapping to GRCh37. Of 20,000 randomly picked Igen Non-ORFs, for our analysis we used two sets of 2,000 each (4,000 sequences total), from which we used 3,889 Igen Non-ORFs that were filtered to not include duplicates, not overlap other control Igen ORFs, Igen Non-ORFs, or genes, and map cleanly to GRCh37. The same approach was applied to 20,000 GRCh37 Igen ORFs and 20,000 Igen Non-ORFs, resulting in 3,924 Igen ORFs and 3,892 Igen Non-ORFs, which cleanly mapped to GRCh38 and passed the filtering described previously. A small number of genes (less than 1.3% of all annotated genes) do not map fully between GRCh37 and GRCh38 (247 genes with the same age distribution as all genes; 179 ancient, 32 metazoan, 25 chordate, 8 mammal, 3 primate). The data were analyzed as a function of gene evolutionary age. Statistical comparisons between gene age categories (genes of different eras and the controls) and between tissue groups were done with the Mann-Whitney U test, with p values corrected for multiple comparisons with the Benjamini-Hochberg correction. P values of 0 indicate numerical values smaller than 2.225×10^{-308}. Data visualization was done using the Python library, Seaborn 0.12.2 (https://github.com/mwaskom/seaborn/tree/v0.12). Boxes are from the 25th to the 75th percentile; medians are indicated by horizontal line; whiskers indicate 1.5 times the interquartile range, with outliers not shown (but included in analysis). For visualization purposes, we applied a log10 data transformation, and handled 0.00 values by adding the smallest non-zero value to every log10 argument.

GREAT analysis

Genomic Regions Enrichment of Annotations Tool (GREAT, version 4.0.4) (107) analysis was performed to assess gene ontology (GO) terms associated with genes in each era, UCEs, or HARs. This approach of using genomic coordinates of interest allowed inspection of HARs, UCEs, and yet unannotated novel genes, especially in the category of primate genes, that may have otherwise gone amiss with more conventional approaches that rely on gene names or identifiers (107). Furthermore, subsets of regions of varying sequence conservation that overlap with 3D genomic feature of interest were determined using pybedtools wrapped bedtools intersect (137, 138). The argument u=True ensured that the same region was written only once if any overlaps were found with 3D genomic feature. Such regions that intersected a 3D genomic feature were assessed for their association with GO terms against the relevant background. For genes in each era, the total list of all era genes was used as a background. In the case of UCEs and HARs, the whole genome provided by GREAT was used as a background.

Data Availability

The publicly available datasets were obtained using accession numbers GSE71831 (83), GSE71072 (86), GSE63525 (6), GSE77565 (93), the 3D Genome Browser (3dgenome.org), dbGaP (GTEx, phs000424.v8.p2), and publications as provided in the Table S1.

Acknowledgments

We are grateful to C.-ting Wu and Marc Kirschner for discussions. We thank the Computational Biology Core within the Institute for Systems Genomics at the University of Connecticut for access to the high-performance computing cluster resources. We apologize to the authors whose work we

could not include due to space constraints. N.G. was supported by a Summer Undergraduate Research Fund (SURF) award. Work in J.E.'s laboratory was supported by the University of Connecticut and an award to J.E. from NIH/NIGMS (R35GM146922). V.L. and N.S. were supported by NIH grants NS095654, MH106934, MH116488, and MH110926 (N.S.), and the Simons Foundation (N.S.). D.M., W.P., and A.O.D.L. were supported by the Manton Center for Orphan Disease Research Scholar award (A.O.D.L.).

Author Contributions

K.F., V.L., A.O.D.L., and J.E. designed research. K.F., V.L., N.G., A.K., T.H., D.M., W.P., K.-M.N., N.S., A.O.D.L., and J.E. performed research and analyzed data. K.F., V.L., and J.E. wrote the paper. All authors approved the final version of this manuscript.

Competing Interest Statement

The authors declare no competing interests.

References

1. J. O. Davies, A. M. Oudelaar, D. R. Higgs, J. R. Hughes, How best to identify chromosomal interactions: a comparison of approaches. *Nat Methods* **14**, 125-134 (2017).
2. I. Jerkovic, G. Cavalli, Understanding 3D genome organization by multidisciplinary methods. *Nat Rev Mol Cell Biol* **22**, 511-528 (2021).
3. R. Kempfer, A. Pombo, Methods for mapping 3D chromosome architecture. *Nat Rev Genet* **21**, 207-226 (2020).

4.   A. Rao, D. Barkley, G. S. Franca, I. Yanai, Exploring tissue architecture using spatial transcriptomics. *Nature* **596**, 211-220 (2021).

5.   J. R. Dixon *et al*., Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376-380 (2012).

6.   S. S. Rao *et al*., A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* **159**, 1665-1680 (2014).

7.   E. P. Nora *et al*., Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* **485**, 381-385 (2012).

8.   T. Sexton *et al*., Three-dimensional folding and functional organization principles of the Drosophila genome. *Cell* **148**, 458-472 (2012).

9.   G. Fudenberg *et al*., Formation of Chromosomal Domains by Loop Extrusion. *Cell Rep* **15**, 2038-2049 (2016).

10.   M. Gabriele *et al*., Dynamics of CTCF- and cohesin-mediated chromatin looping revealed by live-cell imaging. *Science* **376**, 496-501 (2022).

11.   P. Mach *et al*., Cohesin and CTCF control the dynamics of chromosome folding. *Nat Genet* **54**, 1907-1918 (2022).

12.   A. L. Sanborn *et al*., Chromatin extrusion explains key features of loop and domain formation in wild-type and engineered genomes. *Proc Natl Acad Sci U S A* **112**, E6456-6465 (2015).

13.   Y. Ghavi-Helm, Functional Consequences of Chromosomal Rearrangements on Gene Expression: Not So Deleterious After All? *J Mol Biol* **432**, 665-675 (2020).

14.   A. M. Oudelaar, D. R. Higgs, The relationship between genome structure and function. *Nat Rev Genet* **22**, 154-168 (2021).

15.   B. van Steensel, E. E. M. Furlong, The role of transcription in shaping the spatial organization of the genome. *Nat Rev Mol Cell Biol* **20**, 327-337 (2019).

16.   R. P. McCord, N. Kaplan, L. Giorgetti, Chromosome Conformation Capture and Beyond: Toward an Integrative View of Chromosome Structure and Function. *Mol Cell* **77**, 688-708 (2020).

17.   K. C. Akdemir *et al*., Disruption of chromatin folding domains by somatic genomic rearrangements in human cancer. *Nat Genet* **52**, 294-305 (2020).

18.   T. S. Hsieh *et al*., Enhancer-promoter interactions and transcription are largely maintained upon acute loss of CTCF, cohesin, WAPL or YY1. *Nat Genet* **54**, 1919-1932 (2022).

19.   S. S. P. Rao *et al*., Cohesin Loss Eliminates All Loop Domains. *Cell* **171**, 305-320 e324 (2017).

20.   M. Spielmann, D. G. Lupianez, S. Mundlos, Structural variation in the 3D genome. *Nat Rev Genet* **19**, 453-467 (2018).

21.   W. Schwarzer *et al*., Two independent modes of chromatin organization revealed by cohesin removal. *Nature* **551**, 51-56 (2017).

22.   E. P. Nora *et al*., Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains from Genomic Compartmentalization. *Cell* **169**, 930-944 e922 (2017).

23.   J. M. Alexander *et al*., Live-cell imaging reveals enhancer-dependent Sox2 transcription in the absence of enhancer proximity. *Elife* **8** (2019).

24.   N. S. Benabdallah *et al*., Decreased Enhancer-Promoter Proximity Accompanying Enhancer Activation. *Mol Cell* **76**, 473-484 e477 (2019).

25. B. Bonev *et al.*, Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell* **171**, 557-572 e524 (2017).

26. P. Freire-Pritchett *et al.*, Global reorganisation of cis-regulatory units upon lineage commitment of human embryonic stem cells. *Elife* **6** (2017).

27. V. Y. Goel, M. K. Huseyin, A. S. Hansen, Region Capture Micro-C reveals coalescence of enhancers and promoters into nested microcompartments. *Nat Genet* **55**, 1048-1056 (2023).

28. H. L. Harris *et al.*, Chromatin alternates between A and B compartments at kilobase scale for subgenic organization. *Nat Commun* **14**, 3303 (2023).

29. T. Heist, T. Fukaya, M. Levine, Large distances separate coregulated genes in living Drosophila embryos. *Proc Natl Acad Sci U S A* **116**, 15062-15067 (2019).

30. A. M. Oudelaar *et al.*, Dynamics of the 4D genome during in vivo lineage specification and differentiation. *Nat Commun* **11**, 2722 (2020).

31. R. J. Palstra *et al.*, The beta-globin nuclear compartment in development and erythroid differentiation. *Nat Genet* **35**, 190-194 (2003).

32. K. S. M. Reed *et al.*, Temporal analysis suggests a reciprocal relationship between 3D chromatin structure and transcription. *Cell Rep* **41**, 111567 (2022).

33. D. Vernimmen, M. De Gobbi, J. A. Sloane-Stanley, W. G. Wood, D. R. Higgs, Long-range chromosomal interactions regulate the timing of the transition between poised and active gene expression. *EMBO J* **26**, 2041-2051 (2007).

34. S. Zhang, N. Ubelmesser, M. Barbieri, A. Papantonis, Enhancer-promoter contact formation requires RNAPII and antagonizes loop extrusion. *Nat Genet* **55**, 832-840 (2023).

35. G. Bejerano *et al.*, Ultraconserved elements in the human genome. *Science* **304**, 1321-1325 (2004).

36. N. Harmston, A. Baresic, B. Lenhard, The mystery of extreme non-coding conservation. *Philos Trans R Soc Lond B Biol Sci* **368**, 20130021 (2013).

37. D. Polychronopoulos, J. W. D. King, A. J. Nash, G. Tan, B. Lenhard, Conserved non-coding elements: developmental gene regulation meets genome organization. *Nucleic Acids Res* **45**, 12611-12624 (2017).

38. A. Sandelin *et al.*, Arrays of ultraconserved non-coding regions span the loci of key developmental genes in vertebrate genomes. *BMC Genomics* **5**, 99 (2004).

39. R. B. McCole, J. Erceg, W. Saylor, C. T. Wu, Ultraconserved Elements Occupy Specific Arenas of Three-Dimensional Mammalian Genome Organization. *Cell Rep* **24**, 479-488 (2018).

40. N. Harmston *et al.*, Topologically associating domains are ancient features that coincide with Metazoan clusters of extreme noncoding conservation. *Nat Commun* **8**, 441 (2017).

41. C. P. Bird *et al.*, Fast-evolving noncoding sequences in the human genome. *Genome Biol* **8**, R118 (2007).

42. E. C. Bush, B. T. Lahn, A genome-wide screen for noncoding elements important in primate evolution. *BMC Evol Biol* **8**, 17 (2008).

43. K. M. Girskis *et al.*, Rewiring of human neurodevelopmental gene regulatory programs by human accelerated regions. *Neuron* **109**, 3239-3251 e3237 (2021).

44. R. M. Gittelman *et al.*, Comprehensive identification and analysis of human accelerated regulatory DNA. *Genome Res* **25**, 1245-1255 (2015).

45. M. J. Hubisz, K. S. Pollard, Exploring the genesis and functions of Human Accelerated Regions sheds light on their role in human evolution. *Curr Opin Genet Dev* **29**, 15-21 (2014).

46. K. Lindblad-Toh *et al*., A high-resolution map of human evolutionary constraint using 29 mammals. *Nature* **478**, 476-482 (2011).

47. K. S. Pollard *et al*., Forces shaping the fastest evolving regions in the human genome. *PLoS Genet* **2**, e168 (2006).

48. S. Prabhakar, J. P. Noonan, S. Paabo, E. M. Rubin, Accelerated evolution of conserved noncoding sequences in humans. *Science* **314**, 786 (2006).

49. S. Prabhakar *et al*., Human-specific gain of function in a developmental enhancer. *Science* **321**, 1346-1350 (2008).

50. K. C. Keough *et al*., Three-dimensional genome rewiring in loci with human accelerated regions. *Science* **380**, eabm1696 (2023).

51. G. Andrews *et al*., Mammalian evolution of human cis-regulatory elements and transcription factor binding sites. *Science* **380**, eabn7930 (2023).

52. R. N. Doan *et al*., Mutations in Human Accelerated Regions Disrupt Cognition and Social Behavior. *Cell* **167**, 341-354 e312 (2016).

53. R. B. McCole, C. Y. Fonseka, A. Koren, C. T. Wu, Abnormal dosage of ultraconserved elements is highly disfavored in healthy cells but not cancer cells. *PLoS Genet* **10**, e1004646 (2014).

54. R. B. McCole *et al*., Structural disruption of genomic regions containing ultraconserved elements is associated with neurodevelopmental phenotypes. bioRxiv [Preprint] (2017). doi: 2010.1101/233197.

55. P. F. Sullivan *et al*., Leveraging base-pair mammalian constraint to understand genetic variation and human disease. *Science* **380**, eabn2937 (2023).

56. A. R. Carvunis *et al*., Proto-genes and de novo gene birth. *Nature* **487**, 370-374 (2012).

57. D. Tautz, T. Domazet-Loso, The evolutionary origin of orphan genes. *Nat Rev Genet* **12**, 692-702 (2011).

58. N. Vakirlis, Z. Vance, K. M. Duggan, A. McLysaght, De novo birth of functional microproteins in the human lineage. *Cell Rep* **41**, 111808 (2022).

59. Y. Shimomura *et al*., APCDD1 is a novel Wnt inhibitor mutated in hereditary hypotrichosis simplex. *Nature* **464**, 1043-1047 (2010).

60. M. T. Levine, C. D. Jones, A. D. Kern, H. A. Lindfors, D. J. Begun, Novel genes derived from noncoding DNA in Drosophila melanogaster are frequently X-linked and exhibit testis-biased expression. *Proc Natl Acad Sci U S A* **103**, 9935-9939 (2006).

61. T. J. Heinen, F. Staubach, D. Haming, D. Tautz, Emergence of a new gene from an intergenic region. *Current biology : CB* **19**, 1527-1531 (2009).

62. S. A. Slavoff *et al*., Peptidomic discovery of short open reading frame-encoded peptides in human cells. *Nature chemical biology* **9**, 59-64 (2013).

63. A. McLysaght, D. Guerzoni, New genes from non-coding sequence: the role of de novo protein-coding genes in eukaryotic evolutionary innovation. *Philos Trans R Soc Lond B Biol Sci* **370**, 20140332 (2015).

64. S. B. Van Oss, A. R. Carvunis, De novo gene birth. *PLoS Genet* **15**, e1008160 (2019).

65. J. Y. Chen *et al*., Emergence, Retention and Selection: A Trilogy of Origination for Functional De Novo Proteins from Ancestral LncRNAs in Primates. *PLoS Genet* **11**, e1005391 (2015).

66. L. Zhao, P. Saelao, C. D. Jones, D. J. Begun, Origin and spread of de novo genes in Drosophila melanogaster populations. *Science* **343**, 769-772 (2014).

67. D. M. Anderson *et al.*, A micropeptide encoded by a putative long noncoding RNA regulates muscle performance. *Cell* **160**, 595-606 (2015).

68. A. Kumar, J. W. Godwin, P. B. Gates, A. A. Garza-Garcia, J. P. Brockes, Molecular basis for the nerve dependence of limb regeneration in an adult vertebrate. *Science* **318**, 772-777 (2007).

69. D. Oz-Levi *et al.*, Noncoding deletions reveal a gene that is critical for intestinal function. *Nature* **571**, 107-111 (2019).

70. A. Vonica *et al.*, Apcdd1 is a dual BMP/Wnt inhibitor in the developing nervous system and skin. *Dev Biol* **464**, 71-87 (2020).

71. S. Yamamoto *et al.*, Structural characterization of human de novo protein NCYM and its complex with a newly identified DNA aptamer using atomic force microscopy and small-angle X-ray scattering. *Front Oncol* **13**, 1213678 (2023).

72. A. Garza-Garcia, R. Harris, D. Esposito, P. B. Gates, P. C. Driscoll, Solution structure and phylogenetics of Prod1, a member of the three-finger protein superfamily implicated in salamander limb regeneration. *PLoS One* **4**, e7123 (2009).

73. R. Neme, D. Tautz, Evolution: dynamics of de novo gene emergence. *Current biology : CB* **24**, R238-240 (2014).

74. A. A. Bazzini *et al.*, Identification of small ORFs in vertebrates using ribosome footprinting and evolutionary conservation. *EMBO J* **33**, 981-993 (2014).

75. J. R. Prensner *et al.*, Noncanonical open reading frames encode functional proteins essential for cancer cell survival. *Nat Biotechnol* **39**, 697-704 (2021).

76. A. Raj *et al.*, Thousands of novel translated open reading frames in humans inferred by ribosome footprint profiling. *Elife* **5** (2016).

77. B. Vanderperre *et al.*, Direct detection of alternative open reading frames translation products in human significantly expands the proteome. *PLoS One* **8**, e70698 (2013).

78. T. Domazet-Loso, J. Brajkovic, D. Tautz, A phylostratigraphy approach to uncover the genomic history of major adaptations in metazoan lineages. *Trends Genet* **23**, 533-539 (2007).

79. T. Domazet-Loso *et al.*, No Evidence for Phylostratigraphic Bias Impacting Inferences on Patterns of Gene Emergence and Evolution. *Mol Biol Evol* **34**, 843-856 (2017).

80. J. A. Weber *et al.*, The whale shark genome reveals how genomic and physiological properties scale with body size. *Proc Natl Acad Sci U S A* **117**, 20662-20671 (2020).

81. A. Derti, F. P. Roth, G. M. Church, C. T. Wu, Mammalian ultraconserved elements are strongly depleted among segmental duplications and copy number variants. *Nat Genet* **38**, 1216-1220 (2006).

82. N. C. Durand *et al.*, Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. *Cell Syst* **3**, 99-101 (2016).

83. E. M. Darrow *et al.*, Deletion of DXZ4 on the human inactive X chromosome alters higher-order genome architecture. *Proc Natl Acad Sci U S A* **113**, E4504-4512 (2016).

84. J. R. Dixon *et al.*, Chromatin architecture reorganization during stem cell differentiation. *Nature* **518**, 331-336 (2015).

85. I. E. Eres, K. Luo, C. J. Hsiao, L. E. Blake, Y. Gilad, Reorganization of 3D genome structure may contribute to gene regulatory evolution in primates. *PLoS Genet* **15**, e1008278 (2019).

86.   Y. Guo *et al*., CRISPR Inversion of CTCF Sites Alters Genome Topology and Enhancer/Promoter Function. *Cell* **162**, 900-910 (2015).

87.   J. H. I. Haarhuis *et al*., The Cohesin Release Factor WAPL Restricts Chromatin Loop Extension. *Cell* **169**, 693-707 e614 (2017).

88.   D. Leung *et al*., Integrative analysis of haplotype-resolved epigenomes across human tissues. *Nature* **518**, 350-354 (2015).

89.   D. H. Phanstiel *et al*., Static and Dynamic DNA Loops form AP-1-Bound Activation Hubs during Macrophage Development. *Mol Cell* **67**, 1037-1048 e1036 (2017).

90.   A. J. Rubin *et al*., Lineage-specific dynamic and pre-established enhancer-promoter contacts cooperate in terminal differentiation. *Nat Genet* **49**, 1522-1528 (2017).

91.   T. J. Salameh *et al*., A supervised learning framework for chromatin loop detection in genome-wide contact maps. *Nat Commun* **11**, 3428 (2020).

92.   A. D. Schmitt *et al*., A Compendium of Chromatin Contact Maps Reveals Spatially Active Regions in the Human Genome. *Cell Rep* **17**, 2042-2059 (2016).

93.   H. Won *et al*., Chromosome conformation elucidates regulatory relationships in developing human brain. *Nature* **538**, 523-527 (2016).

94.   X. Luo *et al*., 3D Genome of macaque fetal brain reveals evolutionary innovations during primate corticogenesis. *Cell* **184**, 723-740 e721 (2021).

95.   S. Bhatia *et al*., Disruption of autoregulatory feedback by a mutation in a remote, ultraconserved PAX6 enhancer causes aniridia. *Am J Hum Genet* **93**, 1126-1134 (2013).

96.   X. Lampe *et al*., An ultraconserved Hox-Pbx responsive element resides in the coding sequence of Hoxa2 and is active in rhombomere 4. *Nucleic Acids Res* **36**, 3214-3225 (2008).

97.   D. J. McBride, A. Buckle, V. van Heyningen, D. A. Kleinjan, DNaseI hypersensitivity and ultraconservation reveal novel, interdependent long-range enhancers at the complex Pax6 cis-regulatory region. *PLoS One* **6**, e28616 (2011).

98.   S. Pauls, S. F. Smith, G. Elgar, Lens development depends on a pair of highly conserved Sox21 regulatory elements. *Dev Biol* **365**, 310-318 (2012).

99.   L. A. Pennacchio *et al*., In vivo enhancer analysis of human conserved non-coding sequences. *Nature* **444**, 499-502 (2006).

100.   L. Poitras *et al*., An SNP in an ultraconserved regulatory element affects Dlx5/Dlx6 regulation in the forebrain. *Development* **137**, 3089-3097 (2010).

101.   F. Poulin *et al*., In vivo characterization of a vertebrate ultraconserved enhancer. *Genomics* **85**, 774-781 (2005).

102.   A. Visel *et al*., Ultraconservation identifies a small subset of extremely constrained developmental enhancers. *Nat Genet* **40**, 158-160 (2008).

103.   A. Woolfe *et al*., Highly conserved non-coding sequences are associated with vertebrate development. *PLoS Biol* **3**, e7 (2005).

104.   G. T. Consortium *et al*., Genetic effects on gene expression across human tissues. *Nature* **550**, 204-213 (2017).

105.   N. A. An *et al*., De novo genes with an lncRNA origin encode unique human brain developmental functionality. *Nat Ecol Evol* **7**, 264-278 (2023).

106.   T. Domazet-Loso, D. Tautz, An ancient evolutionary origin of genes associated with human genetic diseases. *Mol Biol Evol* **25**, 2699-2707 (2008).

107.   C. Y. McLean *et al*., GREAT improves functional interpretation of cis-regulatory regions. *Nat Biotechnol* **28**, 495-501 (2010).

108. L. F. Lareau, S. E. Brenner, Regulation of splicing factors by alternative splicing and NMD is conserved between kingdoms yet evolutionarily flexible. *Mol Biol Evol* **32**, 1072-1079 (2015).

109. L. F. Lareau, M. Inada, R. E. Green, J. C. Wengrod, S. E. Brenner, Unproductive splicing of SR genes associated with highly conserved and ultraconserved DNA elements. *Nature* **446**, 926-929 (2007).

110. J. Z. Ni *et al.*, Ultraconserved elements are associated with homeostatic control of splicing regulators by alternative splicing and nonsense-mediated decay. *Genes Dev* **21**, 708-718 (2007).

111. S. P. Pirnie, A. Osman, Y. Zhu, G. G. Carmichael, An Ultraconserved Element (UCE) controls homeostatic splicing of ARGLU1 mRNA. *Nucleic Acids Res* **45**, 3473-3486 (2017).

112. C. Rodelsperger *et al.*, Short ultraconserved promoter regions delineate a class of preferentially expressed alternatively spliced transcripts. *Genomics* **94**, 308-316 (2009).

113. K. C. Akdemir *et al.*, Somatic mutation distributions in cancer genomes vary with three-dimensional chromatin structure. *Nat Genet* **52**, 1178-1188 (2020).

114. T. Domazet-Loso, D. Tautz, Phylostratigraphic tracking of cancer genes suggests a link to the emergence of multicellularity in metazoa. *BMC Biol* **8**, 66 (2010).

115. S. Whalen, K. S. Pollard, Enhancer Function and Evolutionary Roles of Human Accelerated Regions. *Annu Rev Genet* **56**, 423-439 (2022).

116. S. Whalen *et al.*, Machine learning dissection of human accelerated regions in primate neurodevelopment. *Neuron* **111**, 857-873 e858 (2023).

117. J. Fraser *et al.*, Hierarchical folding and reorganization of chromosomes are linked to transcriptional changes in cellular differentiation. *Mol Syst Biol* **11**, 852 (2015).

118. T. Pollex *et al.*, Enhancer-promoter interactions become more instructive in the transition from cell-fate specification to tissue differentiation. *Nat Genet* 10.1038/s41588-024-01678-x (2024).

119. Z. Chen *et al.*, Widespread Increase in Enhancer—Promoter Interactions during Developmental Enhancer Activation in Mammals. bioRxiv [Preprint] (2022). doi: 2010.1101/2022.2011.2018.516017.

120. R. D. Acemel, D. G. Lupianez, Evolution of 3D chromatin organization at different scales. *Curr Opin Genet Dev* **78**, 102019 (2023).

121. L. Alvarez-Gonzalez *et al.*, Principles of 3D chromosome folding and evolutionary genome reshuffling in mammals. *Cell Rep* **41**, 111839 (2022).

122. F. Marletaz *et al.*, The little skate genome and the evolutionary emergence of wing-like fins. *Nature* 10.1038/s41586-023-05868-1 (2023).

123. E. McArthur *et al.*, Reconstructing the 3D genome organization of Neanderthals reveals that chromatin folding shaped phenotypic and sequence divergence. bioRxiv [Preprint] (2022). doi: 2010.1101/2022.2002.2007.479462.

124. F. M. Real *et al.*, The mole genome reveals regulatory rearrangements associated with adaptive intersexuality. *Science* **370**, 208-214 (2020).

125. M. Vietri Rudan *et al.*, Comparative Hi-C reveals that CTCF underlies evolution of chromosomal domain architecture. *Cell Rep* **10**, 1297-1309 (2015).

126. Y. Okada, Sperm chromatin structure: Insights from in vitro to in situ experiments. *Curr Opin Cell Biol* **75**, 102075 (2022).

127. M. J. McCoy, A. Z. Fire, Parallel gene size and isoform expansion of ancient neuronal genes. *Current biology : CB* 10.1016/j.cub.2024.02.021 (2024).
128. S. Jia, A. Meng, TGFbeta family signaling and development. *Development* **148** (2021).
129. T. Boehm, J. B. Swann, Origin and evolution of adaptive immunity. *Annu Rev Anim Biosci* **2**, 259-283 (2014).
130. K. Buchmann, Evolution of Innate Immunity: Clues from Invertebrates via Fish to Mammals. *Front Immunol* **5**, 459 (2014).
131. D. D. Wu, D. M. Irwin, Y. P. Zhang, Molecular evolution of the keratin associated protein gene family in mammals, role in the evolution of mammalian hair. *BMC Evol Biol* **8**, 241 (2008).
132. M. J. Christmas *et al.*, Evolutionary constraint and innovation across hundreds of placental mammals. *Science* **380**, eabn3943 (2023).
133. T. S. Klaric *et al.*, Human-specific features and developmental dynamics of the brain N-glycome. *Sci Adv* **9**, eadg2615 (2023).
134. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool. *J Mol Biol* **215**, 403-410 (1990).
135. S. El-Gebali *et al.*, The Pfam protein families database in 2019. *Nucleic Acids Res* **47**, D427-D432 (2019).
136. S. Kumar *et al.*, TimeTree 5: An Expanded Resource for Species Divergence Times. *Mol Biol Evol* **39** (2022).
137. R. K. Dale, B. S. Pedersen, A. R. Quinlan, Pybedtools: a flexible Python library for manipulating genomic datasets and annotations. *Bioinformatics* **27**, 3423-3424 (2011).
138. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010).

Figures

Fig. 1. Genomic regions of varying sequence conservation have distinct relationships with features of 3D genome organization.

(*A*) Human genes assigned an evolutionary age by phylogenomics and phylostratigraphy. Representative species used with TimeTree (136) to build a timetree: *Escherichia coli* (ancient), *Amphimedon queenslandica* (metazoan), *Branchiostoma floridae* (chordate), *Ornithorhynchus anatinus* (mammal), and *Homo sapiens* (primate). (*B*) Number of single-isoform human protein-coding genes in every evolutionary era (ancient, metazoan, chordate, mammal, primate). Most genes are ancient. (*C*) Protein length is indicated in number of amino acids (AA) and is largest for ancient genes. (*D*) The length of coding sequences (CDS) is indicated as the number of base pairs (bp) and is largest for ancient genes. (*E*) Visualization of example genomic regions of varying sequence conservation with the GM12878 Hi-C dataset (6) using the Juicebox tool (82). Domains (yellow squares) and peaks indicating loop presence (blue squares), as annotated in Juicebox. (*F*) Pooled domains, loop anchors, and boundaries are significantly positively correlated with era genes ($1.08 \times 10^{-172} \leq p \leq 0.022$), except for mammal genes in pooled domains and loop anchors ($6.03 \times 10^{-5} \leq p \leq 0.654$). UCEs and HARs are significantly positively correlated with pooled domains and loop anchors, but negatively correlated with pooled boundaries ($1.34 \times 10^{-29} \leq p \leq 0.017$). Spearman correlation analysis was performed by partitioning the genome into 50-kb bins. Spearman correlation coefficients are indicated by a heatmap. Non-significant p values are depicted by tildes. Control datasets include ENSP genes, Non-ENSP genes, Igen ORFs, and Igen Non-ORFs.

Fig. 2. Transcriptional levels vary with gene age and tissue of origin.

The expression of human protein-coding genes is measured as the log10 of the mean counts of RNA transcripts across 54 human tissues from the GTEx database (104). (*A*) RNA expression levels vary as a function of evolutionary category, increasing from youngest to oldest genes, and higher in all genes than in control non-genic sequences (Igen ORF, Igen Non-ORF). Testis expression is higher than expression in other tissues in control non-genic sequences. (*B*) RNA expression levels in 6 tissue categories: brain, 3 germ layers (ectoderm, mesoderm, endoderm),

and 2 germline tissues (ovary, testis). Within every tissue category, expression increases from youngest genes to oldest genes and is higher in genes than in control non-genic sequences.

Fig. 3. Genes of different evolutionary eras are associated with prominent cell type-specific processes in pooled domains.

GREAT analysis for different era genes that overlap pooled domains with GO terms associated with (*A*) signaling pathways for metazoan genes, (*B*) immune response-related processes for chordate genes, (*C*) skin development and defense response to other organisms for mammal genes, and (*D*) keratinocyte differentiation for primate genes. (*A-D*) Analysis performed against all era genes as a background.

A

B

GO Biological Process - HARs in Pooled Healthy Loop Anchors

C

GO Biological Process - HARs in Pooled Cancer Loop Anchors

D

Transcription Level

E

Fig. 4. The relationship between genomic regions of different sequence conservation and 3D genome organization in healthy versus disease states.

(*A*) Correlations of regions of varying sequence conservation with pooled domains and pooled loop anchors in healthy versus cancer states. Spearman correlation analysis was performed by partitioning the genome into 50-kb bins. Spearman correlation coefficients indicated by a heatmap. Non-significant p values are depicted by tildes. Control datasets include ENSP genes, Non-ENSP genes, Igen ORFs, and Igen Non-ORFs. (*B* and *C*) HARs that overlap pooled loop anchors in (*B*) healthy and (*C*) cancer states against whole genome background share associations with neuronal GO categories. However, such HARs in healthy state are enriched for GO terms associated with cartilage, pancreas, and gland development, while HARs in cancer are associated with pattern specification, eye, and kidney development. (*D*) Evolutionary progression of transcription levels: RNA expression increases with time from youngest genes (primate) to oldest genes (ancient) and is higher in genes than in control non-genic sequences (Igen ORF, Igen Non-ORF). (*E*) Schematic of the functional associations of genomic regions of varying sequence conservation and their changes with 3D genomic positioning and disease state. Change is indicated by a check mark and consistency is depicted by an X.

Supporting Information for

Functional associations of evolutionarily recent human genes exhibit

sensitivity to the 3D genome landscape and disease

Katherine Fleck[1,2*], Victor Luria[3,4,5,6*], Nitanta Garag[1], Amir Karger[7], Trevor

Hunter[1], Daniel Marten[4,5], William Phu[4,5], Kee-Myoung Nam[8], Nenad Sestan[3],

Anne H. O'Donnell-Luria[4,5,9#], Jelena Erceg[1,2,10#]

[1]Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT 06269

[2]Institute for Systems Genomics, University of Connecticut, Storrs, CT 06269

[3]Department of Neuroscience, Yale School of Medicine, New Haven, CT 06510

[4]Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA 02115

[5]Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge,
MA 02142

[6]Department of Systems Biology, Harvard Medical School, Boston, MA 02115

[7]IT-Research Computing, Harvard Medical School, Boston, MA 02115

[8]Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06510

[9]Department of Pediatrics, Harvard Medical School, Boston, MA 02115

[10]Department of Genetics and Genome Sciences, University of Connecticut Health Center, Farmington, CT 06030

[*]K.F. and V.L. contributed equally to this work.

[#]To whom correspondence may be addressed. Email: odonnell@broadinstitute.org or jelena.erceg@uconn.edu.

Fig. S1. Gene numbers and lengths for each evolutionary era.

(*A*) Number of single-isoform annotated human protein-coding genes in every evolutionary era (ancient, metazoan, chordate, mammal, primate). Most annotated genes are evolutionarily ancient.

(*B*) Number of single-isoform unannotated human protein-coding genes in every evolutionary era (ancient, metazoan, chordate, mammal, primate). Most unannotated genes are evolutionarily young. (*C*) Protein length is indicated in number of amino acids (AA). The evolutionarily youngest genes are much shorter than ancient genes and only slightly longer than control Igen ORF and Igen Non-ORF sequences. (*D*) The length of coding sequences (CDS) is indicated as the number of base pairs (bp). The evolutionarily youngest genes are shorter than ancient genes and similar to control Igen ORF and Igen Non-ORF sequences.

Fig. S2. The relationships between domains and various evolutionary genomic regions show stability or variation across cell types based on sequence conservation.

Ancient genes, annotated ENSP genes, and UCEs are primarily significantly positively correlated with domains ($0 \leq p \leq 0.746$). Conversely, less conserved genes and fast-evolving HARs show variable relationships with domains across cell types ($2.34 \times 10^{-92} \leq p \leq 0.988$). Spearman correlation analysis was performed by partitioning the genome into 50-kb bins. Spearman correlation coefficients are indicated by a heatmap. Non-significant p values are depicted by tildes. Control datasets include ENSP genes, Non-ENSP genes, Igen ORFs, and Igen Non-ORFs.

Fig. S3. The relationships between loop anchors and regions of various sequence conservation show different variation levels across cell types.

Ancient and annotated ENSP genes have more consistent relationships with loop anchors by cell type ($0 \leq p \leq 0.944$), whereas less conserved genes, HARs, and UCEs have more variable relationships ($8.65 \times 10^{-118} \leq p \leq 0.965$). Spearman correlation analysis was performed by partitioning the genome into 50-kb bins. Spearman correlation coefficients are indicated by a

heatmap. Non-significant p values are depicted by tildes. Control datasets include ENSP genes,

Non-ENSP genes, Igen ORFs, and Igen Non-ORFs.

Fig. S4. The relationship of boundaries with era genes, UCEs, and HARs is highly consistent across cell types.

Era genes are predominantly significantly positively correlated with boundaries ($2.19 \times 10^{-110} \leq p \leq 0.306$), while UCEs and HARs are non-significantly or significantly negatively correlated ($8.91 \times 10^{-12} \leq p \leq 0.136$). Spearman correlation analysis was performed by partitioning the genome into 50-kb bins. Spearman correlation coefficients are indicated by a heatmap. Non-significant p values are depicted by tildes. Control datasets include ENSP genes, Non-ENSP genes, Igen ORFs, and Igen Non-ORFs.

Fig. S5. Transcriptional level variation with gene age, tissue of origin and gene length.

As in Fig. 2, the expression of human protein-coding genes is measured for RNA transcripts across 54 human tissues from the GTEx database (1). (*A*) RNA expression levels, measured as log10 of TPMs, increase from youngest to oldest genes (Table S7). Within mammal and primate genes, expression in testis is higher than in other tissues. (*B*) RNA expression levels, measured as log10 of TPMs, in 6 tissue categories increase from youngest genes to oldest genes. (*C*) The raw mean counts highlight that expression is highest in testis and lowest in brain. (*D*) RNA expression levels, measured as log10 of mean counts, increase with the gene length, shown as CDS length. Within 7 CDS length deciles, ancient genes are expressed at highest levels while primate genes are expressed at lowest levels. Within 3 of the shorter CDS length deciles, the highest expression is that of metazoan genes or chordate genes. Only the first 4 deciles have control non-genic sequences, and their expression is 100-1000 times lower than that of genes.

Fig. S6. Gene expression in 54 human tissues and evolutionary eras.

Gene expression levels are measured as mean counts of RNA transcripts across 54 human tissues in 5 evolutionary eras, as in Fig. 2 and Fig. S5. Gene expression increases evolutionarily from primate to ancient genes. With each evolutionary era, testis expression is typically highest, followed by other tissues which differ between eras.

Fig. S7. Functional associations of era genes.

GREAT analysis for (*A*) metazoan genes, (*B*) chordate genes, (*C*) mammal genes, and (*D*) primate genes. Those GO terms are associated with very similar processes as for era genes in pooled domains (Fig. 3). (*A-D*) Analysis performed against all era genes as a background.

Fig. S8. Functional associations of era genes, UCEs, and HARs in pooled loop anchors.

Regions of varying sequence conservation overlapping pooled loop anchors are linked to GO terms associated with (*A*) signaling pathways for metazoan genes, (*B*) immune response-related processes for chordate genes, (*C*) skin development for mammal genes (note absence of defense response compared to Fig. 3*C*), (*D*) keratinocyte differentiation for primate genes (note absence of peptide cross-linking compared to Fig. 3*D*), (*E*) transcription and RNA metabolic processes for UCEs, and (*F*) neuronal processes, cartilage formation, pancreas development, and glandular epithelial cell differentiation for HARs. Backgrounds: all era genes for individual eras of genes; whole genome for UCEs and HARs.

Fig. S9. Functional associations of era genes and UCEs in pooled boundaries.

In pooled boundaries, enrichments for GO terms are related to (*A*) protein localization to telomeres

and ncRNA processes for ancient genes, (*B*) signaling pathways and regulation of reproductive

processes for metazoan genes, (*C*) immune response-related processes for chordate genes, and (*D*)

transcription and RNA metabolic processes for UCEs. Backgrounds: all era genes for individual eras of genes; whole genome for UCEs.

Fig. S10. Functional associations of UCEs and HARs in pooled domains.

GREAT analysis for UCEs and HARs that overlap pooled domains. (*A*) UCEs in pooled domains are associated with transcription and nucleic acid metabolism, and (*B*) HARs in pooled domains are associated with neuronal processes, cartilage formation, and thymus development. Background: whole genome.

Fig. S11. Era genes intersected with pooled domains in healthy state are associated with nearly identical processes as with all pooled domains.

GREAT analysis for regions of varying sequence conservation that overlap pooled healthy domains including (*A*) metazoan genes, (*B*) chordate genes, (*C*) mammal genes, (*D*) primate genes, (*E*) UCEs, and (*F*) HARs. Enrichments for GO terms are related to nearly identical processes as for era genes, UCEs, and HARs in all of the pooled domains (Fig. 3 and Fig. S10). Backgrounds: all era genes for individual eras of genes; whole genome for UCEs and HARs.

Fig. S12. Functional associations of era genes in pooled domains in cancer state are very similar to those in all of the pooled domains and pooled healthy domains.

When intersected with pooled cancer domains, different regions of varying sequence conservation including (*A*) metazoan genes, (*B*) chordate genes, (*C*) mammal genes, (*D*) primate genes, (*E*) UCEs, and (*F*) HARs are associated with very similar GO terms as those in all pooled domains (Fig. 3 and Fig. S10) and pooled healthy domains (Fig. S11), with the exception of gained association with the glycolytic process in primate genes (*D*). Backgrounds: all era genes for individual eras of genes; whole genome for UCEs and HARs.

Fig. S13. Functional associations of era genes and UCEs intersected with pooled loop anchors in healthy state closely resemble those in all pooled loop anchors.

In pooled loop anchors in healthy state, GO terms are associated with (*A*) signaling pathways for metazoan genes, (*B*) immune response-related processes for chordate genes, (*C*) skin development for mammal genes, (*D*) keratinocyte differentiation for primate genes, and (*E*) transcription and

nucleic acid metabolism for UCEs. These GO terms are largely similar to those seen with all pooled loop anchors (Fig. S8). Backgrounds: all era genes for individual eras of genes; whole genome for UCEs.

Fig. S14. Evolutionarily older genes and UCEs exhibit GO term enrichments when in pooled cancer loop anchors.

In pooled cancer loop anchors, GO terms are related to (*A*) a wide variety of processes for ancient genes, (*B*) signaling pathways for metazoan genes, (*C*) immune response-related processes for chordate genes, and (*D*) transcription and nucleic acid metabolism for UCEs. Note the absence of

GO terms for mammal and primate genes. Backgrounds: all era genes for individual eras of genes;

whole genome for UCEs.

Table S1. Information on datasets.

(*A*) Information on era genes. (*B*) Control dataset for Igen ORF. (*C*) Control dataset for Igen Non-ORF. (*D*) Hi-C datasets and related sources.

Table S2. Spearman correlation analysis of the relationship between regions of varying sequence conservation and 3D genomic features (domains, loop anchors, boundaries) by partitioning the genome into 50-kb bins.

(*A*) Spearman correlation coefficients using 50-kb bins. (*B*) p values using 50-kb bins. Non-significant p values (red).

Table S3. Spearman correlation analysis of the relationship between regions of varying sequence conservation and 3D genomic features (domains, loop anchors, boundaries) by partitioning the genome into 20-kb bins.

(*A*) Spearman correlation coefficients using 20-kb bins. (*B*) p values using 20-kb bins. Non-significant p values (red).

Table S4. Spearman correlation analysis of the relationship between regions of varying sequence conservation and 3D genomic features (domains, loop anchors, boundaries) by partitioning the genome into 100-kb bins.

(*A*) Spearman correlation coefficients using 100-kb bins. (*B*) p values using 100-kb bins. Non-significant p values (red).

Table S5. Mean counts of RNA transcripts in 6 adult human tissue categories in sequences from 7 evolutionary categories. There are 7 evolutionary categories of sequences: 5 evolutionary eras (Ancient, Metazoan, Chordate, Mammal, Primate), Igen ORFs, and Igen Non-ORFs. For every sequence, counts of RNA transcripts obtained using the GTEx database are shown, together with information about protein sequence, genomic localization, and gene evolutionary age.

Table S6. Significance of statistical comparisons of mean counts from different tissue categories and evolutionary categories. Statistical significance was calculated with the Mann-Whitney U test and corrected for multiple comparisons using the Benjamini-Hochberg correction.

Table S7. Counts of RNA TPM in 6 adult human tissue categories, in sequences from 7 evolutionary categories. As in Table S5, for every sequence, TPM counts from the GTEx database are shown, together with information about protein sequence, genomic localization, and gene evolutionary age.

SI References

1.   G. T. Consortium *et al.*, Genetic effects on gene expression across human tissues. *Nature* **550**, 204-213 (2017).