Large-scale composite hypothesis testing for omics analyses.

Annaïg De Walsche[1,2][✉], **Franck Gauthier**[2], **Alain Charcosset**[2], **and Tristan Mary-Huard**[1,2]

[1]MIA Paris-Saclay, INRAE, AgroParisTech, Université Paris-Saclay 91120 Palaiseau, France
[2]Génétique Quantitative et Evolution - Le Moulon, INRAE, CNRS, AgroParisTech, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

Abstract

Composite Hypothesis Testing (CHT) based on summary statistics has become a popular strategy to assess the effect of the same marker (or gene) jointly across multiple traits or at different omics levels. Although significant efforts have been made to develop efficient CHT procedures, most approaches face scalability constraints in terms of the number of traits/omics and markers to handle, or fail to account for potential correlations across traits efficiently. Methods relying on mixture models partially circumvent these limitations, but do not provide proper p-values, hampering the use of classical multiple testing procedures, graphical representations (e.g. Manhattan or QQ plots) and limiting their comparison with alternative approaches.

We introduce the qch_copula approach that combines the mixture model approach with a copula function to account for dependencies across traits/omics. The method comes with a p-value that is consistently defined for any composite hypothesis to be tested. By significantly reducing the memory size burden of the EM algorithm during inference, the method scales to the analysis of several (up to 20) traits and $10^5 - 10^6$ markers. We conducted a comprehensive benchmark study, comparing our approach with 6 state-of-the-art methods recently developed. The qch_copula procedure efficiently controls Type I error rate and yields substantial gain in detecting various patterns of joint associations. The interest of the method is further illustrated by the joint analysis of 14 association studies to detect pleiotropic genes involved in psychiatric disorders.

The proposed method is implemented in the R package qch, available on CRAN.

meta-analysis | composite hypothesis testing | mixture model | data integration | pleiotropy
Correspondence: *annig.de-walsche@inrae.fr*

Introduction

Consider a study where the goal is to assess the joint effect of a specific drug treatment in two different tissues. One aims at defining a test procedure that rejects hypothesis H_0 «the drug has no joint effect», when both hypotheses H_0^1 «the drug has no effect on tissue 1» and H_0^2 «the drug has no effect on tissue 2» are false. This corresponds to a particular case of composite hypothesis testing where the composite H_0 hypothesis to be tested is $H_0^1 \cup H_0^2$. A popular strategy to perform Composite Hypothesis Testing (CHT) is to combine the test statistics and/or p-values derived for each of the marginal hypotheses H_0^1 and H_0^2 into a single summary statistics. While the p-values related to H_0^1 and H_0^2 can be obtained using common statistical procedures, providing a suitable summary statistic

along with a valid rejection rule (that ensures the control of the false positive rate at the required nominal level) for the test of the composite H_0 hypothesis is not straightforward. The particular CHT problem of the form $H_0^1 \cup H_0^2$ was addressed as soon as the early 80s, with the seminal work of (1), and its generalization to the case of testing $H_0^1 \cup \ldots \cup H_0^Q$ with $Q \geq 2$ has been investigated by (2).

In the context of large omics studies CHT has rapidly become a standard procedure. In human genetics, it has been successfully used in the context of mediation analysis (3–6) or for the detection of genomic regions that are associated with two diseases, e.g. Prostate Cancer and Type 2 Diabete (7). In these omics applications CHT is typically performed at the gene or marker level, resulting in a large number of simultaneously tested composite hypotheses. A first consequence is the need for CHT methods to explicitly provide p-values in order to be combined with multiple testing correction procedures. A second consequence is the opportunity to exploit the large amount of available data to model and infer the relationship between the 2 series of p-values, and to account for this relationship in the CHT procedure to better control Type I error rate (7).

More recently, CHT has also been used in the context of integrative genomics to jointly analyse multiple omics traits, such as DNA methylation, copy number variation and gene expression (8–10). When applied to such studies, CHT typically requires the analysis of $Q > 2$ series of p-values (i.e. one serie per omics dataset) which led to the development of dedicated procedures. Most existing methods are based on a same model where the Q-uplet of p-value associated to each gene/marker is distributed as a multivariate mixture where each of the 2^Q components corresponds to a specific combination of H_0^q and H_1^q states (8–10). While versatile, the mixture approach suffers from several limitations. Although type I error rate can be evaluated through local FDR in a mixture model setting (11, 12), the approach does not yield p-values, limiting the range of multiple testing procedure to be applied, the use of sanity checks such as QQ-plots or histograms, and the comparison with alternative approaches. Moreover, the number of components of the mixture model restricts the application of method to $Q = 3$ (9) to ≈ 10 (8, 10). Based on the same mixture model approach, we propose a new method for CHT called **qch_copula** that explicitly accounts for the dependence structure across p-value series through a copula function (13). The new procedure comes with several important features. First we show how a consistent p-value

can be defined from the mixture model approach and show the connection between adaptive Benjamini-Hochberg FDR control (14) applied to these p-values and a local FDR control (11, 12) directly applied to the posteriors obtained from the mixture model. Second, we provide computational short-cuts that significantly alleviate the memory burden of EM inference procedure of the approach, extending its application to series of p-values as large as $n = 10^5$ genes/markers and $Q = 20$. Third, we show that the coupling of the non-parametric approach of (10) to estimate the marginal H_1^q distributions with the copula modelling for dependency yields excellent performance in terms of detection accuracy and Type I error rate control. Importantly, we provide the first benchmark study for the comparison of six recently developed CHT methods (5–10). It is shown that half of the methods do not correctly control for Type I error rate when the p-value series are correlated. Our approach is implemented in the R package **qch** available from the CRAN.

Materials and methods

In this section, we define the composite hypothesis (Section Composite hypothesis), then introduce our new method for CHT based on a mixture model approach (Section Model) and explain the inference (Section Inference) and the testing procedure (Section Testing composite hyprthesis). Section Computational shortcuts specifies the algorithmic optimisations that significantly alleviate the memory burden of inference procedure. We define our simulation framework for evaluate the performance of our method and existing methods (Section Simulation framework).

Composite hypothesis. Assume a collection of n items (e.g. genes or SNP) have been tested for their effects in Q conditions (e.g. traits, tissues or environments). We denote by H_0^q (resp. H_1^q) the null (resp. alternative) hypothesis corresponding to test q ($1 \leq q \leq Q$) and consider the set $\mathcal{C} := \{0,1\}^Q$ of all possible combinations of null and alternative hypotheses across the Q. For a given configuration $c := (c_1, ..., c_Q) \in \mathcal{C}$, the joint hypothesis $\mathcal{H}^c$ is defined as

$$\mathcal{H}^c := \left(\bigcap_{q:c_q=0} H_0^q \right) \cap \left(\bigcap_{q:c_q=1} H_1^q \right)$$

Considering two complementary subsets $\mathcal{C}_0$ and $\mathcal{C}_1$ of $\mathcal{C}$, we define the composite null and alternative hypotheses $\mathcal{H}_0$ and $\mathcal{H}_1$ as

$$\mathcal{H}_0 := \bigcup_{c \in \mathcal{C}_0} \mathcal{H}^c, \quad \mathcal{H}_1 := \bigcup_{c \in \mathcal{C}_1} \mathcal{H}^c$$

We aim at testing $\mathcal{H}_0$ versus $\mathcal{H}_1$ for each item i.

Model. Let P_i^q be the p-value obtained for test q on item i, and $Z_i^q = -\Phi^{-1}(P_i^q)$ its negative probit transform, where Φ stands for the standard Gaussian cdf. We note $Z_i := (Z_i^1, ..., Z_i^Q)$ the z-score profile of item i.

Each item i is associated with a vector $L_i := (L_i^1, ..., L_i^Q) \in \mathcal{C}$, where L_i^q is the binary variable being 0 if H_{0i}^q holds and 1

if H_{1i}^q holds. Assuming that the items are independent, each z-score profile arises from a mixture model with 2^Q components defined as follows:

$$Z_i \sim \sum_{c \in \mathcal{C}} w_c \psi^c. \tag{1}$$

with ψ^c the distribution of Z_i conditionally on $L_i = c$ and $w_c := \Pr\{L_i = c\}$.

The mixture model (1) involves 2^Q multivariate distributions ψ^c that need to be estimated, which may become quite cumbersome when Q becomes large. To reduce the computational burden, we assume that all cumulative distributions of ψ^c have the following form:

$$\Psi_\theta^c(Z_i) = C_\theta(F_{c_1}^1(Z_i^1), ..., F_{c_q}^q(Z_i^q), ..., F_{c_Q}^Q(Z_i^Q))$$

where F_0^q (resp. F_1^q) is the cumulative distribution of Z_i^q conditional on $L_i^q = 0$ (resp. $L_i^q = 1$) and C_θ is a copula of parameter θ accounting for the dependency structure between the Q z-scores. This yields:

$$\psi_\theta^c(Z_i) = c_\theta \left(F_{c_1}^1(Z_i^1), ..., F_{c_Q}^Q(Z_i^Q) \right) \prod_{q:c_q=0} f_0^q(Z_i^q) \prod_{q:c_q=1} f_1^q(Z_i^q) \tag{2}$$

where f_0^q (resp. f_1^q) is the distribution of Z_i^q conditional on $L_i^q = 0$ (resp. $L_i^q = 1$). Note that in this last expression one has $F_0^q = \Phi$ for all q, so that inferring the emission distributions of the mixture boils down to the estimation of Q univariate cumulative distributions F_1^q and parameter θ.

In practice one needs to choose the form of the copula distribution. Here we considered Gaussian copula as it allows the specification of different correlations for each pair of variables. The density function of the Gaussian copula is given by:

$$c_\theta(u) = |\theta|^{-\frac{1}{2}} \exp\left(-\frac{1}{2} \xi_u^T (\theta^{-1} - I) \xi_u \right)$$

$$\text{where } \xi_u = (\Phi^{-1}(u_1), ..., \Phi^{-1}(u_Q)) \quad \forall u \in [0,1]^Q$$

Here θ is the $(Q \times Q)$ correlation matrix associated to the Gaussian copula. Note that setting $\theta = I_Q$ corresponds to the qch method of (10).

Inference. The model parameters are the distributions $f_1^1, ..., f_1^Q$, the copula parameter θ and proportions w_c. Following (8) and (10), the inference procedure was split into two steps:

1. Get an estimate of each alternative distribution f_1^q,

2. Estimate proportions w_c and the copula parameter θ using an EM algorithm, after plug-in the estimates $\hat{f}_1^q$.

Marginal distributions. Combining Model (Eq. (1)) with the definition of the ψ_c Eq. (2), one has

$$Z_i^q \sim \pi_0^q f_0^q + (1 - \pi_0^q) f_1^q, \tag{3}$$

where f_0^q is the distribution of Z_i^q conditional on $L_i^q = 0$ and f_1^q its distribution conditional on $L_i^q = 1$. Since Z_i^q is defined

as the negative probit transform of a p-value, one has $f_0^q = \phi$ for all q's, with ϕ the standard Gaussian density function.

The null proportions can be directly derived by applying the following estimator:

$$\hat{\pi}_0^q = [n(1-\lambda)]^{-1}|\{i : P_i^q > \lambda\}|,$$

where $\lambda \in [0,1]$ is a tuning parameter that can be determined through bootstrap (15). Once the proportions of mixture Eq. (3) are known (or plugged-in after estimation), the alternative distribution f_1^q can be inferred through a non-parametric procedure using a kernel method (12). Both procedures (for the π_0^q and f_1^q estimations) are known to be consistent.

Configuration proportions and copula parameter. The kernel estimates $\hat{f}_1^q$ can be plugged directly into the mixture model, so that the only quantities to infer are the copula parameter θ and the weights w_c. Inference can be efficiently performed using a standard EM. Denoting $u_{ic} = (F_{c_1}(z_i^1),...,F_{c_Q}(z_i^Q))^T$ and $\xi_{ic} = (\Phi^{-1}(u_{ic}^1),...,\Phi^{-1}(u_{ic}^Q))^T$ one obtains the following explicit update equations (see Suppl. Mat. section 1 for details):

E step: $\quad \hat{\tau}_{ic} = \widehat{\Pr}\{L_i = c|Z_i; \hat{\theta}\} = \dfrac{\hat{w}_c \psi_{\hat{\theta}}^c(Z_i)}{\sum_{c' \in \mathcal{C}} \hat{w}_{c'} \psi_{\hat{\theta}}^{c'}(Z_i)}$

M step: $\quad \hat{w}_c = \dfrac{1}{n}\sum_i \hat{\tau}_{ic}$

$$\hat{\theta} = \frac{1}{n}\sum_{i=1}^{n}\sum_{c \in C} \hat{\tau}_{ic}(\xi_{ic}\xi_{ic}^T)$$

Testing composite hypothesis. The existing multivariate methods (8–10) are not strictly valid test procedures since they do not provide a p-value. We present here how consistent p-value can be defined from the mixture model approach.

Let c be the (unknown) configuration of the item under consideration, and $\mathcal{C}_0$ and $\mathcal{C}_1$ be any two complementary subsets of $\mathcal{C}$, and consider testing

$$\mathcal{H}_0 : \{c \in \mathcal{C}_0\} \quad \text{vs} \quad \mathcal{H}_1 : \{c \in \mathcal{C}_1\}$$

The mixture model Eq. (1) can be written in the following form:

$$Z \sim W_0 \psi^0 + W_1 \psi^1 := \psi$$

where $W_0 = \sum_{c \in \mathcal{C}_0} w_c$, $\psi^0 := W_0^{-1}\sum_{c \in \mathcal{C}_0} w_c\psi_c$ (resp. for W_1 and ψ^1). We consider the posterior

$$\tau = \Pr\{L \in \mathcal{C}_1|Z\} = \sum_{c \in \mathcal{C}_1}\Pr\{L = c|Z\} = \frac{W_1\psi^1(Z)}{W_0\psi^0(Z) + W_1\psi^1(Z)}$$

as a test statistic. The corresponding p-value is then:

$$\begin{aligned}\text{pval} &= \Pr_{\mathcal{H}_0}\{\tau(Z) > \tau\} \\ &= \int_{\mathbb{R}^Q} \mathbb{1}_{\{\tau(z)\geq\tau\}}\psi^0(Z)dZ \\ &= \frac{1}{W_0}\int_{\mathbb{R}^Q}\mathbb{1}_{\{\tau(z)\geq\tau\}}(1-\tau(z))\psi(Z)dZ\end{aligned}$$

where the last equality comes from the definition of τ. In practice, estimates of posteriors

$$\hat{\tau}_i = \frac{\widehat{W}_1\psi_{\hat{\theta}}^1(Z_i)}{\widehat{W}_0\psi_{\hat{\theta}}^0(Z_i) + \widehat{W}_1\psi_{\hat{\theta}}^1(Z_i)}$$

can be computed for each item i. Using these estimates and approximating the integral by its empirical counterpart, one gets:

$$\widehat{\text{pval}}_i = \frac{1}{n\widehat{W}_0}\sum_{j=1}^{n}\mathbb{1}_{\{\hat{\tau}_j > \hat{\tau}_i\}}(1-\hat{\tau}_j)$$

Having access to p-values provides several advantages over existing methods that rely on posteriors only, from combining the testing procedure with any correction method for multiple testing to checking the p-value distribution for quality control and providing graphical displays such as Vulcano or Manhattan plots.

The connection between correcting the p-values $\widehat{\text{pval}}_i$ for multiple testing using the adaptive Benjamini Hochberg procedure (14) and the estimation of lFDR from the posteriors that are provided in (11) and (12) is thoroughly established in Suppl. Mat. Section 2.

Computational shortcuts. When jointly analyzing a large number of conditions, the number of configurations $|\mathcal{C}| = 2^Q$ increases exponentially with the number of conditions Q and the memory burden of the procedure can be prohibitive. A classical EM implementation requires the computation and the storage of the matrix $T = (\tau_{ic})$ in the step (E), which becomes cumbersome whenever n and/or Q are large: assuming e.g. $n = 10^5$ and $Q = 15$, T requires 26 Go of storage. In the step (M), the update of the weights w_c at the iteration $(t+1)$ is given by $\hat{w}_c^{(t+1)} = \frac{1}{n}\sum_i \hat{\tau}_{ic} = \frac{1}{n}\sum_i \frac{\hat{w}_c^{(t)}\psi_{\hat{\theta}^{(t)}}^c(Z_i)}{S_i^{(t)}}$ with $S_i^{(t)} = \sum_{c \in \mathcal{C}} \hat{w}_c^{(t)}\psi_{\hat{\theta}^{(t)}}^c(Z_i)$. To reduce the memory burden, we proposed to compute and save only the quantities $S_i^{(t)}, 1 \leq i \leq n$ in the step (E). And in step (M), the $\hat{w}_c^{(t)}\psi_{\hat{\theta}^{(t)}}^c(Z_i)$ are calculated on the fly for updating the weights $\hat{w}_c^{(t+1)}$. The same procedure is used to update the copula parameter $\hat{\theta}^{(t+1)}$ (See Suppl. Mat. Section 3 for the details). This reduces the algorithm's memory footprint from $O(n \times 2^Q)$ for the matrix $T = (\tau_{ic})$ to $O(n + 2^Q)$ for S_i and

w_c. Storing only the quantities required allows to go from 26 Go to 1 Mo of storage for $n = 10^5$ and $Q = 15$, and allowing the application to a number of conditions as large as $Q = 20$ and $n = 10^5$. However, this shortcut involves the need to compute the $\psi_{\hat{\theta}(t)}^c(Z_i)$ for $i = 1, ..., n$ and $c \in \mathcal{C}$ three times, i.e. to compute of the $S_i^{(t)}$ in step (E) and to update the weights $\hat{w}_c^{(t+1)}$ and the copula parameter $\hat{\theta}^{(t+1)}$ in step (M), instead of one time at each iteration of the EM algorithm. This increases the algorithm's complexity resulting in a loss of efficiency in terms of computation time. In order to be efficient in terms of calculation time, the method is implemented in Rcpp and a parallelised version is available.

Simulation framework. We evaluated and compared the performance of our method with existing methods in various simulated scenarios. Hereafter, we will refer to items and conditions as "markers" and "traits", mimicking a multi-trait genome-wide association study. Each dataset consists of a $n \times Q$ matrix of z-scores generated as follows. Markers are clustered into $K + 1$ clusters. Each cluster is characterized by a configuration c, i.e. all markers of cluster k share the same configuration $c_k = (c_{k1}, ..., c_{kQ})$. One has

$$Z_i \sim \mathcal{N}(2c_k, \Sigma_\rho),$$

where Σ_ρ is a $Q \times Q$ correlation matrix where all covariances between traits are set to ρ. Note that $\mu_{iq} = 0$ if H_0^q holds for item i, and $\mu_{iq} = 2$ otherwise. Assuming that matrix Σ_ρ is common to all clusters, a given cluster k is characterized by its associated number of markers and associated configuration vector. We considered different values for Q (2,8,16), two scenarios: $sparse$ or $dense$, and two levels of correlation ($\rho = 0.3, 0.5$). In the $sparse$ scenario, a large proportion (i.e. ≥ 0.9) of the markers were associated with none of the traits, whereas in the dense case, half of the markers were associated with at least one trait. For each setting (i.e., a combination of a number of traits, a correlation level and a scenario), simulations were repeated 25 times. In total, $3 \times 2 \times 2 = 12$ settings were investigated (See Table 4 in Suppl. Mat. for details). All simulations were performed using $n = 10^5$.

We provide the details for generating one dataset in Setting 1: $Q = 2$, $\rho = 0.3$, $sparse$, and the other settings described in Suppl. Mat. Section 4. Twenty clusters of 50 markers associated with a single randomly chosen trait were simulated. Similarly, twenty clusters of 50 markers associated with the two traits were simulated. The process was repeated to generate ten clusters of 100 markers associated with one trait and ten clusters of 100 markers associated with two traits. A last cluster containing all remaining markers corresponding to the complete null configuration, i.e. $c = (0, 0)$, was generated. This last cluster included 96,000 markers, i.e., 96% of the total markers.

Note that this data generation corresponds to the model described by Equations 1 and 2 with the marginal alternative distributions f_1^q being the Gaussian distribution $\mathcal{N}(2, 1)$ and the copula parameter θ being Σ_ρ. As a result, our simulation framework may have an advantage in terms of the detection power of our method and IMIX (9) and Primo methods (8),

Method	Sparse	Dense
DACT_Efron	0 (0)	0 (0)
DACT_JC	0 (0)	**0.08 (0.017)**
DACT_NO	0.032 (0.03)	0.014 (0.015)
HDMT	**0.255 (0.018)**	**0.129 (0.008)**
PLACO	**0.148 (0.091)**	0.058 (0.027)
qch	**0.256 (0.018)**	**0.145 (0.008)**
IMIX	**0.595 (0.128)**	**0.222 (0.094)**
Primo	0.046 (0.021)	**0.079 (0.01)**
qch_copula	0.045 (0.021)	0.057 (0.012)

Table 1. False discovery rate of the procedures for the settings 1 and 2 ($Q = 2, \rho = 0.3$).
For each procedure the FDR (averaged over 25 runs) are displayed. Numbers in brackets correspond to standard errors. False discovery rate above 0.05 are in boldface.

which are also based on the same mixture model Equation 1 compared to other methods.

Results

Simulations. The methodology presented in Section Materials and methods and hereafter called qch_copula was compared to six different methods recently proposed: DACT (5), HDMT (6), PLACO (7), IMIX (9) Primo (8) and qch (10). The first three methods are based on the specific design of a test statistic that combines two p-values for each item and cannot handle a number of conditions Q larger than 2. IMIX and Primo handle the case $Q \geq 3$ and rely on distribution mixtures similar to the one of Equation 1. Except for DACT, HDMT and qch, all methods account for dependencies across p-values series. We first compared methods regarding Type I error (T1E) control. The methods that achieved control of T1E at the requested level were further compared in terms of detection power.

Type I error rate control. In settings 1-4, $n = 10^5$ markers were tested for $Q = 2$ traits. The goal was to identify markers associated with both traits. As Primo only provides an estimate of the FDR at a given threshold, we relied on the equivalence between lFDR estimation and p-values based on posteriors proven in Section Testing composite hypothesis to derive p-values for each marker for this method. FDR control at a nominal level of 5% was then applied through the adaptive BH procedure to obtain the final list of candidate markers per method. The results are presented in Table 1 for the case $\rho = 0.3$. Significant departures from the nominal T1E level were observed for most methods, DACT, Primo and qch_copula being the only ones to successfully control the FDR level, whatever the scenario. Note that the qch method, which assumes conditional independence, struggles to control the type I error rate effectively in the presence of correlation, illustrating the interest in taking dependence into account. Similar results were observed when the correlation between studies was high ($\rho = 0.5$, See Table 5 in Suppl. Mat.).

We further investigated TIE control in settings 5-8 and 9-12, where the number of traits was $Q = 8$ and $Q = 16$, re-

spectively, to identify markers associated with at least 2, 4 or 8 traits. Only Primo and qch_copula were considered, as the other methods can only handle cases where $Q = 2$. Although IMIX could theoretically handle cases with larger values of Q, in practice, the computational burden would be prohibitive, and the associated R package does not handle cases where $Q > 3$. The results are displayed in Table 2 for the case $\rho = 0.3$ (settings 5-6 and 9-10) and Table 6 in Suppl. Mat. for $\rho = 0.5$ (settings 7-8 and 11-12). In settings 5-8 ($Q = 8$), both Primo and qch_copula effectively controlled for T1E. Applied to settings 9-12 ($Q = 16$), Primo either ran out of memory or resulted in prohibitive computational cost (>24h). Our method efficiently controlled for T1E in the dense scenario. In the sparse scenario, one could observe slightly inflated T1E, the actual risk being lower than 8% in most cases.

Detection power. The results of the power detection analysis are detailed for settings 5-6 and 9-10, where the number of traits was $Q = 8$ and $Q = 16$, respectively, and $\rho = 0.3$ in Table 2. Overall, qch_copula and Primo yielded high power whatever the tested composite hypothesis. However, qch_copula exhibited a substantial power improvement - up to 45% - over Primo, regardless of the simulation setting. Similar results were observed when the correlation between traits was high ($\rho = 0.5$, settings 7-8 and 11-12, see Table 6 in Suppl. Mat.) and when $Q = 2$ (settings 1-4, not shown). The average computational time associated with qch_copula was 78 minutes for the model fitting step and $\approx$1mn for each tested composite hypothesis.

Application to the detection of pleiotropic genes. To illustrate our method, we performed a comprehensive analysis of 14 psychiatric disorders using data derived from genetic association studies conducted by the Psychiatric Genomics Consortium, see Section 7 in Suppl. Mat. for details. We used the pre-processed dataset of $n = 24,155$ genes described in (16). The analysis aimed at identifying pleiotropic genes simultaneously associated with multiple disorders.

In the initial analysis, the authors used the PLACO method to detect genes associated with two or more disorders, focusing on top genes detected in at least seven disorders. Since the number of traits allowed by PLACO is limited to 2, all pairwise analyses of 2 among the 14 traits were performed. For a given gene, the list of associated disorders corresponded to those found significant in at least one pairwise analysis. This approach suffers from several technical limitations. First, it may yield ambiguous or inconsistent results across pairwise analyses. As an example, gene TMX2 was detected in both the ADHD-AN and the AN-SCZ analyses but not in the ADHD-SCZ analysis. Similarly gene MIR2113, one of the two genes "that are identified in 10 psychiatric disorders", is declared significant in only 26 out of the 45 PLACO pairwise comparisons involving the 10 candidate disorders, with Anorexia Nevrosa (AN) being identified in only 1 out of its 13 associated pairwise comparisons. Second, aggregating the results of all pairwise analyses does not come with any statistical guarantee regarding false positive control for the final gene list.

The goal of detecting pleiotropic genes can be turned into a composite hypothesis testing procedure where the corresponding composite null hypothesis for gene i is defined as follows:

$$\mathcal{H}_{0i} : \{\text{Gene } i \text{ is associated with at most } q-1 \text{ disorders}\},$$

i.e. $\mathcal{H}_1$ genes are associated to at least q disorders. We performed CHT for $q \in 7, \ldots, 14$ with a nominal type-I error rate $\alpha = 0.05$ and compared the outcomes with those obtained using the PLACO pairwise analyses. The results are displayed in Table 3. Our method detected less candidate genes than PLACO. In particular, qch_copula did not identify candidate genes associated with more than 8 disorders. Focusing on the composite hypothesis test with $q = 7$, our method identified 60 genes (see Suppl. Mat. Table 8 for their description), of which 29 were in common with the PLACO approach. The lists of genes specifically detected by either PLACO or qch_copula were compared as follows. For a given list, the initial GWAS p-values associated to each gene were collected (i.e. 14 p-values per gene) and the 7 smallest ones were selected, resulting in e.g. 7×31 p-values corresponding to the genes specific to qch_copula. We then computed the proportion of these p-values that were higher than 0.05. As one aims at detecting genes associated with at least 7 disorders, this proportion is expected to be low. The PLACO proportion was 11.5% whereas the qch_copula was 3.2%. The higher number of candidates combined with a lower significance of the initial p-values for PLACO corroborate the findings of the simulation study regarding the difficulty for PLACO to control type I error rate.

Figure 1 (left) displays the Manhattan plot corresponding to the qch_copula method applied to the "at least 7 disorders" $\mathcal{H}_1$ hypothesis. The plot identifies 5 majors peaks with at least 4 detected genes on chromosomes 2, 11, 17, 19 and 20 respectively. The heatmap of the initial GWAS -log10(p-values) of the corresponding candidates genes (Fig. 1, right) reveals that most peaks are associated with the 3 alcohol related disorders (AUDIT-C, AUDIT-P, AUDIT-T), schizophrenia (SCZ) and bipolar disorder.

Conclusion

We have developed a novel approach called qch_copula for composite hypothesis testing based on a multivariate mixture model. Our method comes with a consistent p-value directly defined from the mixture model approach, which, to the best of our knowledge, is the first of its kind.

A key feature of qch_copula lies in its ability to model the emission distributions of the mixture components adaptatively. Previous published methods rely on fully parametric models, assuming both the f_0^q and f_1^q distributions of Equation 3 to be Gaussian density functions (9), or (a mixture of) χ^2 density functions, after data transformation (8). While such modellings allow one to account for correlations across traits explicitly, they significantly constrain the shape of f_1^q. In comparison, qch_copula performs a non-parametric esti-

Q	Scenario	Test	Primo		qch_copula	
			FDR	Power	FDR	Power
8	Dense	at least 2	0.008 (0.001)	0.143 (0.006)	0.051 (0.001)	0.609 (0.003)
		at least 4	0.005 (0.001)	0.126 (0.007)	0.044 (0.002)	0.534 (0.006)
		at least 8	0.048 (0.006)	0.125 (0.007)	0.05 (0.006)	0.186 (0.01)
	Sparse	at least 2	0.012 (0.004)	0.109 (0.008)	0.054 (0.004)	0.388 (0.007)
		at least 4	0.008 (0.004)	0.108 (0.009)	0.053 (0.004)	0.405 (0.014)
		at least 8	0.079 (0.019)	0.092 (0.009)	0.068 (0.023)	0.166 (0.019)
16	Dense	at least 2	-	-	0.052 (0.002)	0.646 (0.004)
		at least 4	-	-	0.057 (0.002)	0.678 (0.006)
		at least 8	-	-	0.057 (0.002)	0.659 (0.007)
	Sparse	at least 2	-	-	0.072 (0.004)	0.528 (0.008)
		at least 4	-	-	0.082 (0.004)	0.626 (0.01)
		at least 8	-	-	0.068 (0.006)	0.568 (0.018)

Table 2. Performance of the procedures Primo and qch_copula for the settings 5-6 ($Q = 8$) and 9-10 ($Q = 16$) with $\rho = 0.3$.
For each procedure the FDR and the detection power (averaged over 25 runs) are displayed. Numbers in brackets correspond to standard errors.

Fig. 1. Results of the composite hypothesis test identifying genes associated with at least 7 disorders. **Left:** -log10(p-values) of the composite hypothesis test along the chromosomes. Significant genes are in orange. **Right:** Initial GWAS -log10(p-values) of the 5 majors peaks with at least 4 candidate genes on chromosomes 2, 11, 17, 19 and 20 respectively. Stars correspond to p-values below 0.001.

At least	7	8	9	10	>10
PLACO	80	38	21	2	0
qch_copula	60	18	0	0	0

Table 3. Number of candidate genes identified by PLACO and qch_copula.

mation of the alternative marginal distribution while still accounting for dependencies through the use of the copula function. This adaptive estimation procedure allows for a better fit of the data, resulting in an efficient control of Type I errors and improved detection performance. In practice, the correlation matrix is inferred within the framework of the EM algorithm jointly with the prior proportions, enhancing our ability to capture the dependence more accurately compared to methods like Primo (8) and PLACO (7), where the correlation matrix is estimated upstream.

Our model assumes that the copula function is the same across all components, employing a single correlation matrix to model the dependence structure. While this may be seen as a limitation, it leads to practical benefits. Modelling the dependency componentwise, as in the IMIX procedure, can lead to a prohibitive computational burden and downgrade the results. This is because some components may be poorly represented in the dataset, compromising the inference of the associated correlation matrix. In contrast, qch_copula strikes a favourable balance between data fitting and computational efficiency in the inference procedure, offering a more practical and effective solution.

The procedure allows the testing of a broad range of composite hypotheses, as one can define any two complementary sets of configurations under $\mathcal{H}_0$ and $\mathcal{H}_1$. Note that a any series of composite hypotheses may be tested in parallel, since the testing step stands independently of the model inference step and does not require parameter re-estimation. The approach relies exclusively on series of p-values, making it amenable to data integration of studies with diverse outcomes, including continuous, binary, and time-to-event measurements. The exclusive reliance on p-values is also particularly advantageous when direct access to the raw data is constrained due to ethical considerations or data confidentiality. In conclusion, we strongly believe that the inherent flexibility of composite hypothesis testing has potential uses in various research settings, including multi-population or multi-environment association analyses in quantitative genetics and extended multi-omics data integration. The proposed methodolgy is implemented in the R package qch, available on CRAN.

ACKNOWLEDGEMENTS

The authors are grateful to the INRAE MIGALE bioinformatics facility (MIGALE, IN-RAE, 2020. Migale bioinformatics Facility, doi:10.15454/1.5572390655343293E12) for providing computing resources. The authors thank Psychiatric Genomics Consortium (PGC) for making the summary statistics of psychiatric disorders GWAS publicly available and are grateful of all the investigators and participants contributed to the studies.

FUNDING

This work was partially supported by INRAE's metaprogram DIGIT-BI0 and by KWS. The funders had no role in the conceptualization, design, data collection, analysis, publication decision, or manuscript preparation.

Bibliography

1. Michael E Sobel. Asymptotic confidence intervals for indirect effects in structural equation models. *Source: Sociological Methodology*, 13:290–312, 1982.

2. Roger L Berger. Multiparameter hypothesis testing and acceptance sampling. 24:295–300, 1982.

3. Yen-Tsung Huang. Joint significance tests for mediation effects of socioeconomic adversity on adiposity via epigenetics. 2018.

4. Yen-Tsung Huang. Genome-wide analyses of sparse mediation effects under composite null hypotheses. 2019.

5. Zhonghua Liu, Jincheng Shen, Richard Barfield, Joel Schwartz, Andrea A. Baccarelli, and Xihong Lin. Large-scale hypothesis testing for causal mediation effects with applications in genome-wide epigenetic studies. *Journal of the American Statistical Association*, 2021.

6. James Y. Dai, Janet L. Stanford, and Michael LeBlanc. A multiple-testing procedure for high-dimensional mediation hypotheses. *Journal of the American Statistical Association*, 117:198–213, 2022.

7. Debashree Ray and Nilanjan Chatterjee. A powerful method for pleiotropic analysis under composite null hypothesis identifies novel shared loci between type 2 diabetes and prostate cancer. *PLoS Genetics*, 16, 2020.

8. Kevin J. Gleason, Fan Yang, Brandon L. Pierce, Xin He, and Lin S. Chen. Primo: Integration of multiple gwas and omics qtl summary statistics for elucidation of molecular mechanisms of trait-associated snps and detection of pleiotropy in complex traits. *Genome Biology*, 21: 1–24, 2020.

9. Ziqiao Wang and Peng Wei. Imix: a multivariate mixture model approach to association analysis through multi-omics data integration. *Bioinformatics*, 36:5439–5447, 2021.

10. Tristan Mary-Huard, Sarmistha Das, Indranil Mukhopadhyay, and Stéphane Robin. Querying multiple sets of p-values through composed hypothesis testing. *Bioinformatics*, 38:141–148, 12 2021.

11. G. J. McLachlan, R. W. Bean, and L. Ben Tovim Jones. A simple implementation of a normal mixture approach to differential gene expression in multiclass microarrays. *Bioinformatics*, 22:1608–1615, 2006.

12. Stéphane Robin, Avner Bar-Hen, Jean Jacques Daudin, and Laurent Pierre. A semi-parametric approach for mixture models: Application to local false discovery rate estimation. *Computational Statistics & Data Analysis*, 51:5483–5493, 2007.

13. Abe Sklar. Random variables, joint distribution functions, and copulas. *Kybernetika*, 9(6): 449–460, 1973.

14. Yoav Benjamini and Yosef Hochberg. On the adaptive control of the false discovery rate in multiple testing with independent statistics. *Journal of Educational and Behavioral Statistics*, 25:60–83, 2000.

15. John D Storey, Jonathan E Taylor, and David Siegmund. Strong control, conservative point estimation and simultaneous conservative consistency of false discovery rates: a unified approach. *J. R. Statist. Soc. B*, 66:187–205, 2004.

16. Haojie Lu, Jiahao Qiao, Zhonghe Shao, Ting Wang, Shuiping Huang, and Ping Zeng. A comprehensive gene-centric pleiotropic association analysis for 14 psychiatric disorders with gwas summary statistics. *BMC Medicine*, 19, 2021.

17. Henrik Hult and Filip Lindskog. Multivariate extremes, aggregation and dependence in elliptical distributions. 34:587–608, 2002.

18. Laramie Duncan, Zeynep Yilmaz, Helena Gaspar, Raymond Walters, and et al. Significant locus and metabolic genetic correlations revealed in genome-wide association study of anorexia nervosa. *American Journal of Psychiatry*, 174:850–858, 2017.

19. T. Otowa, K. Hek, M. Lee, E. M. Byrne, and et al. Meta-analysis of genome-wide association studies of anxiety disorders. *Molecular psychiatry*, 21:1391, 10 2016.

20. Ditte Demontis, Raymond K. Walters, Joanna Martin, Manuel Mattheisen, and et al. Discovery of the first genome-wide significant risk loci for attention-deficit/hyperactivity disorder. *Nature genetics*, 51:63, 2019.

21. Jakob Grove, Stephan Ripke, Thomas D. Als, Manuel Mattheisen, and et al. Identification of common genetic risk variants for autism spectrum disorder. *Nature genetics*, 51:431, 2019.

22. Sandra Sanchez-Roige, Abraham A. Palmer, Pierre Fontanillas, Sarah L. Elson, and et al. Genome-wide association study meta-analysis of the alcohol use disorders identification test (audit) in two population-based cohorts. *American Journal of Psychiatry*, 176:107–118, 2019.

23. Eli A. Stahl, Gerome Breen, Andreas J. Forstner, Andrew McQuillin, and et al. Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nature genetics*, 51:793, 2019.

24. Joëlle A. Pasman, Karin J.H. Verweij, Zachary Gerring, Sven Stringer, and et al. Gwas of lifetime cannabis use reveals new risk loci, genetic overlap with psychiatric traits, and a causal influence of schizophrenia. *Nature neuroscience*, 21:1161, 2018.

25. Naomi R. Wray, Stephan Ripke, Manuel Mattheisen, Maclej Trzaskowski, and et al. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nature genetics*, 50:668, 2018.

26. Paul D. Arnold, Kathleen D. Askland, Cristina Barlassina, Laura Bellodi, and et al. Revealing the complex genetic architecture of obsessive-compulsive disorder using meta-analysis. *Molecular Psychiatry*, 23:1181–1181, 2018.

27. Caroline M. Nievergelt, Adam X. Maihofer, Torsten Klengel, Elizabeth G. Atkinson, and et al. International meta-analysis of ptsd genome-wide association studies identifies sex- and ancestry-specific genetic risk loci. *Nature Communications 2019 10:1*, 10:1–16, 2019.

28. Max Lam, Chia Yen Chen, Zhiqiang Li, Alicia R. Martin, and et al. Comparative genetic architectures of schizophrenia in east asian and european populations. *Nature genetics*, 51:1670–1678, 2019.

29. Dongmei Yu, Jae Hoon Sul, Fotis Tsetsos, Muhammad S. Nawaz, and et al. Interrogating the genetic determinants of tourette's syndrome and other tic disorders through genome-wide association studies. *American Journal of Psychiatry*, 176:217–227, 2019.

Supplementary Note 1: Estimation of the proportions w_c and the copula parameter θ using an EM algorithm.

Initialisation. The prior proportions are initialized using

$$\hat{w}_c = \prod_{q:c_q=0} \hat{\pi}_0^q \prod_{q:c_q=1} (1 - \hat{\pi}_0^q)$$

, where $\hat{\pi}_0^q$ corresponds to the marginal prior probability in model (3), estimated in the first step of the inference procedure. Parameter θ is initialized using the inversion of Kendall's tau estimator (17):

$$\hat{\theta}_{ij} = \sin \frac{\pi}{2} \tau_{ij}$$

where τ_{ij} is the Kendall's tau for the random vector $(Z_i, Z_j)^T$.

Maximum likelihood estimate for the Gaussian copula parameter. The complete likelihood of the observations $z_i = (z_i^1, ..., z_i^Q), 1 \leq i \leq n$ and the latent variables $l_{ic} = \mathbb{1}_{\{l_i=c\}}, 1 \leq i \leq n, c \in \mathcal{C}$ is given by:

$$L(w, \theta | z, l) = \prod_{i=1}^{n} \prod_{c \in \mathcal{C}} [w_c \psi_\theta^c(z_i)]^{l_{ic}}$$

$$= \prod_{i=1}^{n} \prod_{c \in \mathcal{C}} \left[w_c \left(\prod_{q:c_q=0} \phi(z_i^q) \prod_{q:c_q=1} f_1^q(z_i^q) \right) c_\theta(F_{c_1}(z_i 1^1), ..., F_{c_Q}(z_i^Q))) \right]^{l_{ic}}$$

Define $Q\left(w, \theta | w^{(t)}, \theta^{(t)}\right)$ as the expected value of the complete log-likelihood, with respect to the current conditional distribution of L given Z and the current estimates of the parameters $w^{(t)}$ and $\theta^{(t)}$:

$$Q(w, \theta | w^{(t)}, \theta^{(t)}) = \mathbb{E}_{L|Z, w^{(t)}, \theta^{(t)}} [\log L(w, \theta | z, l)]$$

We denote $u_{ic} = (F_{c_1}(z_i^1), ..., F_{c_Q}(z_i^Q))$ and $\xi_{ic} = (\Phi^{-1}(u_{ic}^1), ..., \Phi^{-1}(u_{ic}^Q)), 1 \leq i \leq n, c \in \mathcal{C}$. Then, we have:

 De Walsche *et al.* | Large-scale composite hypothesis testing.

$$Q(w,\theta|w^{(t)},\theta^{(t)}) = \sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}\left[\log(w_c) + \sum_{q:c_q=0}\log(\phi(z_i^q)) + \sum_{q:c_q=1}\log(f_1^q(z_i^q)) + \log(c_\theta(u_{ic}))\right]$$

$$= \sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}\log(c_\theta(u_{ic})) + \text{cste}$$

$$= \sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}\log\left[|\theta|^{-\frac{1}{2}}\exp\left(-\frac{1}{2}\xi_{ic}^T(\theta^{-1}-I)\xi_{ic}\right)\right] + \text{cste}$$

$$= \frac{n}{2}\log(|\theta^{-1}|) - \frac{1}{2}\sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}(\xi_{ic}^T(\theta^{-1}-I)\xi_{ic}) + \text{cste}$$

$$= \frac{n}{2}\log(|\theta^{-1}|) - \frac{1}{2}\sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}(\xi_{ic}^T\theta^{-1}\xi_{ic}) + \text{cste}$$

$$= \frac{n}{2}\log(|\theta^{-1}|) - \frac{1}{2}\sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}\text{tr}\left(\xi_{ic}^T\theta^{-1}\xi_{ic}\right) + \text{cste}$$

$$= \frac{n}{2}\log(|\theta^{-1}|) - \frac{1}{2}\sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}\text{tr}\left(\xi_{ic}\xi_{ic}^T\theta^{-1}\right) + \text{cste}$$

$$= \frac{n}{2}\log(|\theta^{-1}|) - \frac{1}{2}\text{tr}\left(\sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}(\xi_{ic}\xi_{ic}^T)\theta^{-1}\right) + \text{cste}$$

$$= \frac{n}{2}\log(|\Lambda|) - \frac{1}{2}\text{tr}(S\Lambda) + \text{cste}$$

where $\hat{\tau}_{ic} = \dfrac{w_c^{(t)}\psi_{\theta^{(t)}}^c(Z_i)}{\sum_{c\in\mathcal{C}}w_c^{(t)}\psi_{\theta^{(t)}}^c(Z_i)}$, $\Lambda = \theta^{-1}$ and $S = \sum_{i=1}^{n}\sum_{c\in\mathcal{C}}\hat{\tau}_{ic}(\xi_{ic}\xi_{ic}^T)$.

Differentiating the log-likelihood with respect to $\Lambda = \theta^{-1}$ leads to

$$\frac{\delta}{\delta\Lambda}\log L(\theta|x) = \frac{n}{2}\frac{\delta}{\delta\Lambda}\log(|\Lambda|) - \frac{1}{2}\frac{\delta}{\delta\Lambda}\text{tr}(S\Lambda)$$

$$= \frac{n}{2}(\Lambda^{-1})^T - \frac{1}{2}S^T$$

$$= \frac{n}{2}\theta^T - \frac{1}{2}\left(\sum_{i=1}^{n}\sum_{c\in C}\hat{\tau}_{ic}(\xi_{ic}\xi_{ic}^T)\right)^T$$

Setting the derivative at 0, one obtains

$$\frac{\delta}{\delta\Lambda}\log L(\theta|x) = 0 \Leftrightarrow \hat{\theta} = \frac{1}{n}\sum_{i=1}^{n}\sum_{c\in C}\hat{\tau}_{ic}(\xi_{ic}\xi_{ic}^T) \tag{4}$$

Note that the explicit expression of $\hat{\theta}$ obtained in Eq. (4) guarantees that the estimated matrix is symmetric, but not necessarily a positive-definite one with value 1 on the diagonal. In practice, one needs to check that $\hat{\theta}$ is properly a correlation matrix at each (M) step of the EM procedure, and whenever needed correct the initial estimate by i) setting all negative eigenvalues at 0 then ii) scaling the resulting covariance matrix into a correlation matrix.

Supplementary Note 2: Equivalence between Benjamini-Hochberg correction on p-values and the estimation of lFDR from the posteriors

In the classical setting of the test procedure, one can control the FDR at a nominal level of α by applying the Benjamini-Hochberg (BH) procedure. Assume n test were performed and let $p_{(1)} \leq ... \leq p_{(n)}$ be the set of ordered p-values. Define the value r such as

$$r = \arg\max_{k}\left\{k \ / \ \frac{n_0 p_{(k)}}{k} \leq \alpha\right\}.$$

The set of rejected hypotheses is then $S_{BH} = \{i_{(1)}, ..., i_{(r)}\}$, where $i_{(k)}$ is the index associated to p-value $p_{(k)}$. Here n_0 is the number of items for which $\mathcal{H}_0$ holds, which is usually unknown. In practice, n_0 can be upper bounded by n or adaptively obtained from the data, which will be designated hereafter by the adaptive BH procedure(14).

Alternatively, in the setting of a mixture model with two components 0 and 1 with emission distributions ψ_0, ψ_1 and priors W_0, W_1, one can define the posterior

$$\tau = \frac{W_1\psi_1(x)}{W_0\psi_0(x) + W_1\psi_1(x)},$$

The local FDR can then be approximately controlled at a nominal level of α by applying the following empirical procedure (11, 12). Assume n items were generated under the previous mixture model and let $\tau_{(1)} \leq ... \leq \tau_{(n)}$ be the set of ordered posteriors. Define the value m such as

$$m = \arg\max_k \left\{ k \; / \; \frac{1}{k} \sum_{1=1}^{k} (1 - \tau_{(i)}) \leq \alpha \right\}.$$

The set of items classified into class 1 is then $S_{lFDR} = \{i_{(1)}, ..., i_{(m)}\}$, where $i_{(k)}$ is the index associated to posterior $\tau_{(k)}$.

The connection between the two approaches in the context of CHT is established by the next proposition:

Proposition 1: Consider n observations drawn according to mixture model (1) of Section 2.2, and let $\hat{\tau}_i$ and $\widehat{pval}_i$ be the posterior and p-value associated to item i, respectively. Consider the adaptive BH procedure where n_0 is estimated by $n\hat{W}_0$, where $\hat{W}_0 = \sum_{c \in \mathcal{C}_0} \hat{w}_c$. Then $S_{BH} = S_{lFDR}$.

    Proof:

For a given item i, the p-value corresponding to the composite hypothesis test is constructed as follows:

$$\widehat{pval}_i = \frac{1}{nW_0} \sum_{j=1}^{n} \mathbb{1}_{\{\hat{\tau}_j > \hat{\tau}_i\}} (1 - \hat{\tau}_j) \tag{5}$$

Let $\widehat{pval}_{(1)} < ... < \widehat{pval}_{(n)}$ represent the ordered set of p-values, and $i_{(k)}$ is the index associated to p-value $\widehat{pval}_{(k)}$.

According to the adaptive BH procedure, the estimate of the False Discovery Rate when the set of rejected hypotheses is $S_{BH} = \{i_{(1)}, ..., i_{(k)}\}$ is given by:

$$\widehat{FDR}(k) = \frac{\hat{n}_0}{k} \widehat{pval}_{(k)} \tag{6}$$

with $\hat{n}_0$ an estimate of the number of true null hypotheses. Here, one can estimate the number of null hypotheses as $\hat{n}_0 = nW_0$. Hence, we have:

$$
\begin{aligned}
\widehat{FDR}(k) &= \frac{nW_0}{k} \widehat{pval}_{(k)} \\
&= \frac{nW_0}{k} \frac{1}{nW_0} \sum_{j=1}^{n} \mathbb{1}_{\{\hat{\tau}_j > \hat{\tau}_k\}} (1 - \hat{\tau}_j) \\
&= \frac{1}{k} \sum_{j=1}^{n} \mathbb{1}_{\{\hat{\tau}_j > \hat{\tau}_k\}} (1 - \hat{\tau}_j)
\end{aligned}
$$

By the definition of the p-value 5, $\widehat{pval}_i < \widehat{pval}_j \Leftrightarrow \hat{\tau}_j > \hat{\tau}_i$.

Consequently:

$$
\begin{aligned}
\widehat{FDR}(k) &= \frac{1}{k} \sum_{j=1}^{n} \mathbb{1}_{\{\widehat{pval}_j < \widehat{pval}_{(k)}\}} (1 - \hat{\tau}_j) \\
&= 1 - \frac{1}{k} \sum_{j=1}^{n} \mathbb{1}_{\{\hat{\tau}_j > \hat{\tau}_k\}} \hat{\tau}_j) \\
&= \widehat{lFDR}(\hat{\tau}_k)
\end{aligned}
$$

where $\widehat{lFDR}(\hat{\tau}_k)$ is the estimate of the false discovery rate when thresholding at the level $\hat{\tau}_k$ defined by (11) and (12).

Thus, applying the adaptive BH correction with $\hat{n}_0 = nW_0$ to our p-value 5 is equivalent to using the local FDR rejection rule defined by (11) and (12). ∎

Supplementary Note 3: Algorithm for the implicit EM

Here we present how the burden associated to the storage of the posterior matrix can be avoided in the EM algorithm. Quantities in red correspond to the ones that are actually stored in place of the posterior matrix for later use.

Algorithm 1 Proportions w_c and copula parameter θ inference using EM with memory handling

---

Inputs: f0Mat $\in \mathcal{M}_{n \times Q}$ containing the $f_0(Z_i^q)$.

         f1Mat $\in \mathcal{M}_{n \times Q}$ containing the $\hat{f}_1^q(Z_i^q)$.

         $\pi_0 \in \mathbb{R}^Q$ containing the proportions $\hat{\pi}_0^q$.

1:   $\hat{w}_c^{(t)} \leftarrow \prod_{q:c_q=0} \pi_0^q \prod_{q:c_q=1} (1 - \pi_0^q)$ ▷ initialisation

2:   $\hat{\theta}^{(t)} \leftarrow$ Kendall's tau estimate

3:

4:   **while** $||\hat{w}_c^{(t)} - \hat{w}_c^{(t+1)}|| > \epsilon$ or $||\hat{\theta}^{(t)} - \hat{\theta}^{(t+1)}|| > \epsilon$ **do**

5:      **Implicit step E:**

6:      **for** i in $1:n$ **do**

7:          $S_i^{(t)} \leftarrow 0$

8:          **for** $c \in \mathcal{C}$ **do**

9:              $S_i^{(t)} \leftarrow S_i^{(t)} + \hat{w}_c^{(t)} \psi_{\hat{\theta}^{(t)}}^c(Z_i)$ ▷ computation of $\psi_{\hat{\theta}^{(t)}}^c(Z_i)$ on the fly

10:          **end for**

11:      **end for**

12:

13:      **Step M:**

14:      **for** $c \in \mathcal{C}$ **do**

15:          $\hat{w}_c^{(t+1)} \leftarrow 0$

16:          **for** i in $1:n$ **do**

17:              $\hat{w}_c^{(t+1)} \leftarrow \hat{w}_c^{(t+1)} + \frac{\hat{w}_c^{(t)} \psi_{\hat{\theta}^{(t)}}^c(Z_i)}{S_i^{(t)}}$ ▷ computation of $\psi_{\hat{\theta}^{(t)}}^c(Z_i)$ on the fly

18:          **end for**

19:          $\hat{w}_c^{(t+1)} \leftarrow \frac{1}{n} \hat{w}_c^{(t+1)}$

20:      **end for**

21:

22:      $\hat{\theta}^{(t+1)} \leftarrow 0$

23:      **for** i in $1:n$ **do**

24:          **for** $c \in \mathcal{C}$ **do**

25:              $\hat{\theta}^{(t+1)} \leftarrow \hat{\theta}^{(t+1)} + \hat{\tau}_{ic}(\xi_{ic}\xi_{ic}^T)$ ▷ computation of $\psi_{\hat{\theta}^{(t)}}^c(Z_i)$ on the fly

26:          **end for**

27:      **end for**

28:      $\frac{1}{n}\hat{\theta}^{(t+1)} \leftarrow \hat{\theta}^{(t+1)}$

29:      $t \leftarrow t + 1$

30: **end while**

Outputs: $\hat{w}_c \in \mathbb{R}^{2^Q}$ containing the configuration proportions estimates.

         $\hat{\theta}$ the estimate of the copula parameter.

Supplementary Note 4: Simulation framework

Setting	Q	Correlation	Scenario
1		0.3	sparse
2			dense
3	2	0.5	sparse
4			dense
5		0.3	sparse
6			dense
7	8	0.5	sparse
8			dense
9		0.3	sparse
10			dense
11	16	0.5	sparse
12			dense

Table 4. Description of the 12 settings considered for the simulation study.

Details for generating each setting. Each dataset consists of a $n \times Q$ matrix of Z-scores generated as follows. Markers are clustered into $K+1$ clusters. Each cluster is characterized by a configuration c, i.e. all markers of cluster k share the same configuration $c_k = (c_{k1},...,c_{kQ})$. One has

$$Z_i \sim \mathcal{N}\left(2c_k, \Sigma_\rho\right),$$

where Σ_ρ is a $Q \times Q$ correlation matrix where all covariances between traits are set to ρ. Assuming that matrix Σ_ρ is common to all clusters, a given cluster k is characterized by its associated number of markers and associated configuration vector. We considered different values for Q (2,8,16), two scenarios: $sparse$ or $dense$, and two levels of correlation ($\rho = 0.3, 0.5$). Denoting $n_{50,\tilde{q}}$ and $n_{100,\tilde{q}}$ the number of clusters of 50 and 100 markers respectively, which are associated with exactly $\tilde{q}$ randomly chosen traits, the details for generating a dataset for each simulation setting are presented below.

Settings 1 to 4: $Q = 2$*.*

- Sparse scenario: $n_{50,1} = n_{50,2} = 20$ and $n_{100,1} = n_{100,2} = 10$, such that $w_{00} = 0.96$, $w_{01} + w_{10} = 0.02$ and $w_{11} = 0.02$.

- Dense scenario: $n_{50,1} = 400$, $n_{100,1} = 200$ and $n_{50,2} = 100$, $n_{100,2} = 50$, such that $w_{00} = 0.5$, $w_{01} + w_{10} = 0.4$ and $w_{11} = 0.1$.

Settings 5 to 8: $Q = 8$*.*

- Sparse scenario: $n_{50,j} = 20$ and $n_{100,j} = 10$ for $j \in \{1,2,4,6,8\}$, such that $w_0 = 0.90$ for the markers being associated with none of the traits and $w_j = 0.02$ for the markers being associated with exactly $j \in \{1,2,4,6,8\}$ traits.

- Dense scenario: $n_{50,j} = 100$ and $n_{100,j} = 50$ for $j \in \{1,2,4,6,8\}$, such that $w_0 = 0.50$ for the markers being associated with none of the traits and $w_j = 0.1$ for the markers being associated with exactly $j \in \{1,2,4,6,8\}$ traits.

Settings 9 to 12: $Q = 16$*.*

- Sparse scenario: $n_{50,q} = 20$ and $n_{100,q} = 10$ for $q \in \{1,2,4\}$, $n_{50,\tilde{q}} = 10$ and $n_{100,\tilde{q}} = 5$ for $\tilde{q} \in \{6,8,14,16\}$, such that $w_0 = 0.90$ for the markers being associated with none of the traits, $w_q = 0.02$ for the markers being associated with exactly $q \in \{1,2,4\}$ traits, and $w_{\tilde{q}} = 0.01$ for the markers being associated with exactly $\tilde{q} \in \{6,8,14,16\}$ traits.

- Dense scenario: $n_{50,q} = 100$ and $n_{100,q} = 50$ for $q \in \{1,2,4\}$, $n_{50,\tilde{q}} = 50$ and $n_{100,\tilde{q}} = 25$ for $\tilde{q} \in \{6,8,14,16\}$, such that $w_0 = 0.5$ for the markers being associated with none of the traits, $w_q = 0.1$ for the markers being associated with exactly $q \in \{1,2,4\}$ traits, and $w_{\tilde{q}} = 0.05$ for the markers being associated with exactly $\tilde{q} \in \{6,8,14,16\}$ traits.

Supplementary Note 5: Evaluation of the FDR control for composite hypothesis testing of two analysis

Method	Sparse	Dense
DACT_Efron	0 (0)	0 (0)
DACT_JC	0 (0)	0.026 (0.018)
DACT_NO	**0.104 (0.033)**	**0.055 (0.013)**
HDMT	**0.41 (0.017)**	**0.374 (0.015)**
PLACO	**0.3 (0.083)**	**0.052 (0.025)**
QCH	**0.437 (0.015)**	**0.239 (0.006)**
IMIX	**0.871 (0.021)**	**0.295 (0.079)**
Primo	**0.081 (0.022)**	**0.112 (0.009)**
QCH_copula	0.057 (0.025)	0.061 (0.008)

Table 5. False discovery rate of the procedures for settings 3 and 4 ($Q = 2$, $\rho = 0.5$).
For each procedure the FDR (averaged over 25 runs) are displayed. Numbers in brackets correspond to standard errors. False discovery rate above 0.05 are in boldface.

Supplementary Note 6: Detection power in large-scale composite hypothesis testing

Q	Scenario	Test	Primo		qch_copula	
			FDR	Power	FDR	Power
8	Dense	at least 2	0.005 (0.001)	0.144 (0.004)	0.058 (0.002)	0.617 (0.004)
		at least 4	0.006 (0.001)	0.147 (0.005)	0.064 (0.003)	0.549 (0.005)
		all	0.075 (0.006)	0.178 (0.006)	0.104 (0.007)	0.28 (0.006)
	Sparse	at least 2	0.008 (0.003)	0.115 (0.01)	0.063 (0.005)	0.422 (0.007)
		at least 4	0.01 (0.004)	0.12 (0.011)	0.067 (0.007)	0.415 (0.011)
		all	0.142 (0.031)	0.09 (0.01)	0.104 (0.023)	0.119 (0.012)
16	Dense	at least 2	-	-	0.057 (0.002)	0.722 (0.004)
		at least 4	-	-	0.067 (0.002)	0.756 (0.004)
		at least 8	-	-	0.083 (0.002)	0.681 (0.005)
		all	-	-	0.124 (0.011)	0.23 (0.014)
	Sparse	at least 2	-	-	0.08 (0.007)	0.647 (0.007)
		at least 4	-	-	0.094 (0.008)	0.724 (0.009)
		at least 8	-	-	0.1 (0.008)	0.582 (0.014)
		all	-	-	0.172 (0.062)	0.096 (0.023)

Table 6. Performance of the procedures Primo and qch_copula for the settings 7-8 ($Q = 8$) and 11-12 ($Q = 16$) with $\rho = 0.5$. For each procedure the FDR and the detection power (averaged over 25 runs) are displayed. Numbers in brackets correspond to standard errors.

Supplementary Note 7: Application to the detection of pleiotropic genes

Information of 14 psychiatric disorders analyzed in this study. We analyzed 14 psychiatric disorders obtained from the Psychiatric Genomics Consortium (PGC) [1]. The disorders studied are anorexia nervosa (AN), anxiety disorder (AD), attention-deficit/hyperactivity disorder (ADHD), autism spectrum disorder (ASD), alcohol use disorders (AUDIT-T alcohol use traits identification test based on total score, AUDIT-C alcohol use traits identification test based on consumption, AUDIT-P alcohol use traits identification test based on problematic consequences of drinking), bipolar disorder (BIP), cannabis use (CU[2]), major depression disorder (MDD), obsessive-compulsive disorder (OCD), post-traumatic stress disorder (PTSD), schizophrenia (SCZ) and Tourette's syndrome (TS). Table 7 details the information about each study.

Traits	N(case/control)	PGC study	Reference
AN	14,477 (3,495/10,982)	an2017	(18)
AD	17,526 (5,761/11,765)	anx2016	(19)
ADHD	53,293 (19,099/34,194)	adhd2019	(20)
ASD	46,350 (18,381/27,969)	asd2019	(21)
AUDIT-T	121,604	sud2019-alcuse	(22)
AUDIT-C	121,604	sud2019-alcuse	(22)
AUDIT-P	121,604	sud2019-alcuse	(22)
BIP	51,710 (20,352/31,358)	bip2019	(23)
CU	184,765 (53,180/131,585)		(24)
MDD	480,359 (135,458/344,901)	mdd2018	(25)
OCD	9,725 (2,688/7,037)	ocd2018	(26)
PTSD	174,659 (23,212/151,447)	ptsd2019	(27)
SCZ	77,096 (33,640/43,456)	scz2019asi	(28)
TS	14,307 (4,819/9,488)	ts2019	(29)

Table 7. Summary information of 14 psychiatric disorders analyzed in this study.

For all the disorders, we obtained their European-only summary statistics and performed the stringent quality controld described in (16): (i) excluded non-biallelic SNPs and those with strand-ambiguous alleles; (ii) excluded SNPs that had no rs label; (iii) removed duplicated SNPs or those not included in the 1000 GenomesProject or whose alleles did not match those there; (iv)

---

[1] as available from the PGC website: https://pgc.unc.edu/for-researchers/download-results/
[2] available from the following link: https://drive.google.com/drive/folders/1uFBeUwE-0UA0YmLTVQ9Ym_oU2HkHPdhk

excluded SNPs that were located within the region of major histocompatibility complex (chr6: 28.5–33.5 Mb) because of its complex LD structure [81]; (v) kept SNPs that had minor allele frequency (MAF)> 0.01.

We applied MAGMA [61] to aggregate a set of SNP-level associations into a single gene-level association signal relying on summary statistics. The file "G1000_EUR_gene_Annotated_VEGAS.genes.annot" defining the set of SNPs as those located within a given gene was provided by authors of (16). Then, the p-value of each gene was obtained by running this command for all the disorders:

```
./magma -bfile g1000_eur -gene-annot G1000_EUR_gene_Annotated_VEGAS.genes.annot
-pval [GWAS_result_file] N=[N] -genes-only -out [prefix]
```

Where g100_eur is the raw genotype data of the European panel of the 1,000 Genomes, which is available for download from the MAGMA website and serves as a reference for linkage disequilibrium.

Table 8. Candidate genes associated with at least 7 disorders identified by qch_copula.

Gene	CHR	Start	Stop	P-value
EIF2B4	2	27487218	27693324	0.0001603
SNX17	2	27493362	27700400	0.0001333
ZNF513	2	27500097	27703611	0.0001439
PPM1G	2	27504065	27732550	0.0001234
FTH1P3	2	27515489	27716443	0.0001494
NRBP1	2	27551472	27765124	0.0000756
KRTCAP3	2	27565232	27769348	0.0000836
IFT172	2	27567239	27812678	0.0000422
FNDC4	2	27614749	27818126	0.0000321
GCKR	2	27619705	27846550	0.0000298
C2orf16	2	27699388	27905589	0.0000535
ZNF512	2	27705835	27946082	0.0000602
CADM2	3	84908132	86223579	0.0000003
MIR5688	3	85334859	85534942	0.0000096
NFKB1	4	103322485	103638459	0.0000961
LINC00461	5	87736596	88080620	0.0000505
MIR9-2	5	87862670	88062757	0.0000107
TTC12	11	113085250	113337114	0.0001548
ANKK1	11	113158512	113371140	0.0000239
DRD2	11	113180316	113446001	0.0000222
MIR4301	11	113220744	113420810	0.0000175
ARHGAP27_1	17	43371267	43607663	0.0000795
ARHGAP27_2	17	43406717	43610282	0.0000716
PLEKHM1	17	43413265	43668146	0.0000368
MIR4315-1_1	17	43452728	43652801	0.0000449
MIR4315-2_1	17	43452728	43652801	0.0000449
LRRC37A4P	17	43483248	43697889	0.0000118
LOC644172_1	17	43577491	43779748	0.0000046
CRHR1	17	43597709	44013194	0.0000032
MGC57346	17	43597711	43815329	0.0000070
CRHR1-IT1	17	43616340	43823595	0.0000054
MAPT-AS1	17	43820721	44072879	0.0000025
SPPL2C	17	43822255	44024438	0.0000039
MAPT	17	43871747	44205699	0.0000019
MAPT-IT1	17	43873148	44076164	0.0000008
STH	17	43976615	44177060	0.0000014
KANSL1	17	44007281	44402740	0.0000062
KANSL1-AS1	17	44170938	44374089	0.0000087
LOC644172_2	17	44220929	44423154	0.0000146
LRRC37A	17	44272496	44515160	0.0000190
ARL17A_1	17	44276499	44539163	0.0000206
ARL17B	17	44276912	44539134	0.0000161
NSFP1_1	17	44350178	44600463	0.0000277
ARL17A_2	17	44486046	44757088	0.0001004
NSF	17	44568034	44934828	0.0000078
NSFP1_2	17	44568099	44818040	0.0000132
WNT3	17	44739871	44996126	0.0000258
WNT9B	17	44828967	45054437	0.0000344
FAM83E	19	49003856	49216694	0.0001283
RPL18	19	49018583	49222675	0.0000918
SPHK2	19	49022547	49233663	0.0000877
DBP	19	49033816	49240807	0.0001047
CA11	19	49041271	49249451	0.0001185
SEC1P	19	49041295	49285502	0.0001090
NTN5	19	49064663	49276264	0.0001385

TOP1	20	39557461	39853126	0.0001137
LOC101927117	20	39626633	39866640	0.0000639
PLCG1	20	39666160	39904357	0.0000394
MIR6871	20	39697662	39897718	0.0000565
ZHX3	20	39707088	40028739	0.0000677