Multi-way modelling of oral microbial dynamics and host-microbiome interactions during induced gingivitis

G.R. van der Ploeg[1], B.W. Brandt[2], B.J.F. Keijser[3], M.H. van der Veen[2,4], C.M.C. Volgenant[2,5], E. Zaura[2], A.K. Smilde[1], J.A. Westerhuis[1], A. Heintz-Buschart[1*]

[1] *Biosystems Data Analysis, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, The Netherlands*

[2] *Preventive Dentistry, Academic Centre for Dentistry, Vrije Universiteit Amsterdam and University of Amsterdam, Amsterdam, The Netherlands*

[3] *Microbiology and Systems Biology, TNO Healthy Living and Work, Leiden, The Netherlands*

[4] *Paediatric Dentistry, Academic Centre for Dentistry, Vrije Universiteit Amsterdam and University of Amsterdam, Amsterdam, The Netherlands*

[5] *Cariology, Academic Centre for Dentistry, Vrije Universiteit Amsterdam and University of Amsterdam, Amsterdam, The Netherlands*

*Author for correspondence: Anna Heintz-Buschart a.u.s.heintzbuschart@uva.nl

Geert Roelof van der Ploeg:	https://orcid.org/0009-0007-5204-3386
Monique van der Veen:	https://orcid.org/0000-0002-6844-4682
Bart Keijser:	https://orcid.org/0000-0003-1597-2948
Catherine Volgenant:	https://orcid.org/0000-0002-4049-2914
Bernd Brandt:	https://orcid.org/0000-0002-1155-1989
Egija Zaura:	https://orcid.org/0000-0003-1432-6194
Age Smilde:	https://orcid.org/0000-0002-3052-4644
Johan Westerhuis:	https://orcid.org/0000-0002-6747-9779
Anna Heintz-Buschart:	https://orcid.org/0000-0002-9780-1933

Abstract

Gingivitis - the inflammation of the gums - is a reversible stage of periodontal disease. It is caused by dental plaque formation due to poor oral hygiene. However, gingivitis susceptibility involves a complex set of interactions between the oral microbiome, oral metabolome and the host. In this

study, we investigated the dynamics of the oral microbiome and its interactions with the salivary

metabolome during experimental gingivitis in a cohort of 41 systemically healthy participants. We

use Parallel Factor Analysis (PARAFAC), which is a multi-way generalization of Principal

Component Analysis (PCA) that can model the variability in the response due to subjects, variables

and time. Using the modelled responses, we identified microbial subcommunities with similar

dynamics that connect to the magnitude of the gingivitis response. By performing high level

integration of the predicted metabolic functions of the microbiome and salivary metabolome, we

identified pathways of interest that describe the changing proportions of Gram-positive and Gram-

negative microbiota, variation in anaerobic bacteria, biofilm formation and virulence.

Key Words

Longitudinal data, Salivary metabolomics, Parallel Factor Analysis, Data integration, Pathway

enrichment analysis

Introduction

Microbial oral diseases, such as dental caries and gum inflammation, are a global public health

burden (Peres et al., 2019; Petersen et al., 2005; WHO, 2022). Together, they are the most prevalent

health condition world-wide, impacting quality of life and imposing a significant economic burden

with 5-10% of public health expenditure in most industrialized countries (WHO, 2022; Widström

et al., 2004). Prevention of oral diseases is therefore of critical importance.

Gingivitis is a reversible stage of periodontal disease marked by inflamed, bleeding gums (Chapple

et al., 2015; Curtis et al., 2020; Huang et al., 2014). It is caused by dental plaque accumulation due

to poor oral hygiene (Löe et al., 1965; Murakami et al., 2018) and is present in a substantial part of

the global population (Han et al., 2020; WHO, 2022). The pathogenesis of gingivitis involves a

complex set of interactions between the oral microbiome, the oral metabolome and the host

(Chapple et al., 2018; Huang et al., 2014). These are in turn dependent on many factors (Chapple et

al., 2018; Kilian et al., 2016), including host diet (Van Der Velden et al., 2011), life-style (Bergström

& Preber, 1986), and genetics (Nibali et al., 2017). Therefore, the oral microbiome alone does not

determine whether and to what extent gingivitis develops, as can be seen by the high degree of

individual variation in gingivitis susceptibility seen in experimental gingivitis models after

refraining from toothbrushing (Axelsson et al., 1991; Chapple et al., 2015, 2018; Guk et al., 2020;

Huang et al., 2014; van der Veen et al., 2016)

While there is no monocausal microbial agent in gingivitis, the gingival crevice microbiota

underneath the gums shifts to a dysbiotic state (Abusleme et al., 2021; Curtis et al., 2020; Diaz et al.,

2016). The depletion of Gram-positive bacteria, such as *Rothia dentocariosa,* and enrichment in

Gram-negative bacteria such as *Porphyromonas gingivalis*, *Prevotella* spp. and *Selenomonas* spp. is

frequently observed in case-control comparisons, despite this shift being less pronounced compared

to periodontitis progression (Curtis et al., 2020; Huang et al., 2014; Kistler et al., 2013; Schincaglia

et al., 2017). Additionally, increasing subgingival anaerobiosis during gingivitis progression

promotes the growth of pathobionts (Curtis et al., 2020; Diaz et al., 2002; Ter Steeg et al., 1988).

Being the medium through which the host and the microbiota interact, the salivary metabolome is

often studied as a non-invasive diagnostic of oral disease (Dawes & Wong, 2019; Proctor, 2016). A

well-known host-microbiome interaction is the relation between host sugar consumption and the

production of acids by the dental plaque microbiota. This lowers the local pH and shifts the

microbiota further towards dysbiosis (König & Navia, 1995; Marsh, 1991). However, many other

dynamic microbe-microbe and host-microbiome interactions that occur throughout gingivitis onset

and progression are not well understood.

Here, we investigate the spatially resolved oral bacterial microbiome and its interactions with the

salivary metabolome during experimental gingivitis to (1) identify microbial subcommunities with

common dynamics and (2) pinpoint potential impacts on biochemical pathways. Reversible

gingivitis was induced in a cohort of 41 systemically healthy participants by omission of oral hygiene

during a two-week gingivitis challenge (**Figure 1A**; (Prodan et al., 2016; van der Veen et al., 2016)).

Previously reported plaque levels, gingival bleeding upon probing, and red fluorescent plaque

quantification allowed grouping of the participants into identified low, mid and high responders

based on the last day of the intervention (**Figure 1B**; (van der Veen et al., 2016)). During a two-week

baseline period at two time points (day -14 and day 0), a challenge period at four time points (day 2

to day 14), and one week resolution period at one time point (day 21), the oral microbiome was

determined at six sites: tongue, saliva, supragingival plaque at the lower and upper lingual surfaces

and supragingival plaque at the lower and upper interproximal surfaces (**Figure 1C**). To identify the

individual variation in the time-resolved response of the oral microbiome to the challenge and

describe commonly responding (groups of) microbiota, we employ Parallel Factor Analysis

(PARAFAC; (Carroll & Chang, 1970; Harshman, 1970)), for modelling the subjects, time, and

microbial abundances at each site (**Figure 1D**). The salivary metabolome dynamics, assessed in

unstimulated saliva during the challenge, is likewise modelled and integrated with microbiome

models at the biochemical pathway level. These analyses should yield insights into microbial

processes that underlie individual gingivitis responses.

Figure 1: Overview of the study design and analysis. (a) Timeline of experimental gingivitis challenge with analyses; (b)
gingivitis responses: Three groups of responders were identified based on red fluorescent dental plaque (RF%) at day 14: low
(0-1.7%), mid (1.8-6.0%) and high (6.0%+); (c) oral sampling sites; (d) data analysis approach: Parallel factor analysis
(PARAFAC), a multi-way generalization of Principal Component analysis, decomposes a multi-way data cube into components
with three loading vectors each: one for the subjects, one for the variables (here: microbial relative abundances) and one for
time. This approach allows us to identify the individual variation in the time-resolved response to the challenge and to describe
commonly responding groups of microbiota.

Results

PARAFAC describes clinically relevant variation in oral microbiomes between subjects

To describe the microbial dynamics of each oral site for all subjects across time, we analysed 16S rRNA gene amplicon sequencing data for a total of 14,014 amplicon sequence variants (ASVs). After removal of sparse ASVs, we created separate PARAFAC models for each oral site, choosing the most appropriate number of PARAFAC components by inspecting the number of iterations needed to converge (Bro, 1997), the core consistency diagnostic (Bro & Kiers, 2003), the variation explained (Bro, 1997), and the tucker congruence coefficient per mode (Lorenzo-Seva & ten Berge, 2006; Tucker, 1951) (**Supplementary Data**). This yielded two-component models for the upper jaw lingual plaque (19.7% explained variation) and upper jaw interproximal plaque (16.5%), tongue (31.5%) and saliva (17.5%) microbiomes (**Figure 2**). Lower jaw lingual and interproximal plaque microbiomes were best represented by one-component models (explaining 11.4% and 7.2% of the variation, respectively). The model of the lower jaw interproximal plaque microbiome was not investigated further due to the limited amount of explained variation and due to the model not describing biologically meaningful information.

119
Figure 2: Overview of the PARAFAC models per sample type. In the rows, the PARAFAC models per sample type are described:
upper jaw lingual plaque, upper jaw interproximal plaque, lower jaw lingual plaque, tongue and saliva. Variance explained
per model are shown in the sample type labels. In the left column the loadings of the first and second component in the subject
mode are plotted against each other, with every subject having a unique character to identify them across the sample types.
In the middle column the loadings of the first and second component in the ASV mode are plotted against each other. We
identified microbial subcommunities (ASV clusters) with common responses per sampling site by selecting and then clustering
*well-modelled ASVs based on their fitted response (Methods, **Table 2**). ASVs are color-coded by cluster number (or shown in*
grey if removed prior to clustering). In the right column, the loadings per time point are shown per component. The model
corresponding to the lower jaw lingual plaque only has one component. The model of the lower jaw interproximal plaque
microbiome was not investigated further due to the limited amount of explained variation and due to the model not describing
biologically meaningful information.

To interpret the models, we tested the correlation between the time-resolved subject loadings of each component with the measured parameters of gingivitis. This approach revealed correlation of components with one or more clinical parameter for the three plaque microbiomes (**Table 1**). Overall, the PARAFAC models describe the variation that exists between subjects, rather than the variation within a subject (one-sided Wilcoxon rank-sum test: p=0.020, **Supplementary Figures A1 and A2**). Furthermore, subject age was found to be significantly described by the first component of the model corresponding to the tongue microbiome samples (p=0.035) and the second component of the model corresponding to the saliva microbiome samples (p=0.038) (**Supplementary Table B1**). Gender was not significantly described by any model (**Supplementary Table B1**). We also performed Pearson correlation tests of the time-resolved subject loadings per component with richness and evenness. All microbiome models were found to contain at least one component that significantly described richness or evenness (p≤0.05; **Table 1**). In conclusion, the PARAFAC models represented clinically and ecologically relevant parameters.

PARAFAC describes microbial subcommunities with common gingivitis responses

Building on the interpretability of the subject modes (**Table 1**), we identified microbial subcommunities with common responses per sampling site by selecting and then clustering well-modelled ASVs based on their fitted response (**Figure 2**, for cluster membership see **Table 2**). Per sampling site, at least one subcommunity containing mainly pathobionts or pro-inflammatory genera such as *Actinomyces*, *Campylobacter*, *Capnocytophaga*, *Fusobacterium*, *Leptotrichia* and *Porphyromonas* (Huang et al., 2014; Kirst et al., 2015; Kistler et al., 2013; Schincaglia et al., 2017; The Human Microbiome Project Consortium, 2012) and one subcommunity containing mainly commensal genera such as *Kingella oralis* (Abusleme et al., 2021), *Streptococcus* spp. (Caporaso et al., 2011; Zaura et al., 2009), and *Veillonella* spp. (Moore & Moore, 1994) have been identified.

The sum of the relative abundances of the ASVs in each cluster was determined to test the difference between the microbiomes of individuals in the low and high response groups at every time point per oral site (Benjamini-Hochberg corrected permutation test of mean difference: p≤0.05; **Supplementary Table B2)**. This approach revealed significant differences in the relative abundance of most plaque-associated ASV clusters in high responders compared to the low responders at baseline (day -14; **Figure 3**, **Supplementary Figure A3**). As such, the PARAFAC models of plaque microbiomes describe microbial subcommunities with common dynamics that connect to the magnitude of the gingivitis response.

We further investigated the robustness of the ecosystem during gingivitis onset and progression by

comparing the sum of the relative abundances per cluster at the start and end of the intervention

(**Supplementary Table B3**). We found significant differences between the relative abundances of

some ASV clusters at the start and end of the gingivitis challenge in lower jaw lingual (ASV cluster

1) and upper jaw interproximal (ASV cluster 4) plaque microbiomes in all response groups

(Benjamini-Hochberg corrected two-sided Wilcoxon rank sum test: $p \leq 7.2e\text{-}13$ and $p \leq 2.6e\text{-}6$,

respectively). For a further three ASV clusters (upper jaw lingual ASV cluster 1 and upper jaw

interproximal clusters 1 and 3), we found the sum of relative abundances to be significantly different

($p=0.002$, $p=0.0025$, $p=0.0039$, respectively) in high responders, but not in low responders. These

results suggest that ecosystem stability is different between high and low responders in several oral

niches.

174
Figure 3: Overview of the sums of relative abundances per ASV cluster and sample type, separating subjects by response group.
Error bars correspond to the standard error of the mean (SEM). The mean difference between the high and low response
groups per time point was tested using a Benjamini-Hochberg corrected permutation test of 999 iterations (*: p≤0.05; **:
p≤0.01; ***: p≤0.001). Mid responders and ASV clusters 3 and 4 of the upper jaw interproximal plaque samples are not shown
*for visual clarity (**Supplementary Figure A3** and **A4**, respectively). Please refer to **Table 2** for the list of identified species per*
ASV cluster.

Integration of the microbiome and metabolome PARAFAC models identifies host-microbiome

interactions at the pathway level

With the established connection between the clinical parameters of gingivitis and the variation in

the plaque microbiomes as modelled by PARAFAC, we investigated host-microbiome interactions

at the biochemical pathway level. We obtained functional predictions of the microbiome per oral

site using Tax4Fun2 (Aßhauer et al., 2015; Wemheuer et al., 2020). We then created separate

PARAFAC models of the salivary metabolomics data and of the functional prediction of the

microbiomes at each oral site. Despite correcting for correlation between most salivary metabolite

levels, likely due to variable water content of the saliva (see Methods), the subject loadings of both

components of the model corresponding to the salivary metabolome were found to correlate with

the correcting factor (p=2.0e-14 and p=2.0e-5, respectively). Regardless, we expected the salivary

metabolomics model to have sufficient freedom to also describe variation related to the gingivitis

response.

The well-modelled molecular functions (KEGG orthologous groups belonging to pathways that

produce or consume the salivary metabolites) and well-modelled salivary metabolites were

combined in a ranked list. KEGG-pathway enrichment was identified using SetRank, which is a

functional enrichment algorithm that corrects for multiple pathway membership (Simillion et al.,

2017). This revealed that the microbiome and metabolome responses to the gingivitis challenge

reflected differences in carbon and energy metabolism, as well as cell wall structure (**Figure 4,**

Supplementary Figure A5, Supplementary Table B4). For example, the tongue microbiome and

salivary metabolome were involved in a joint response in the biosynthesis of cofactors and carbon

metabolism (p=0.0045 and p=0.0052, respectively). The quorum sensing pathway was significantly

enriched in all plaque microbiomes (p≤0.05). The pathway enrichment results therefore reflect the

microbiome responses and highlight potential metabolic interactions with the human host.

Figure 4: Overview of the pathway enrichment results per combination of oral site microbiome and the salivary metabolome. Tax4Fun2 was used to create functional predictions from ASV data. Well-modelled metabolites and microbiome molecular functions were integrated by mapping them to KEGG pathway level. SetRank was then used to test for pathway enrichment and corrected for multiple-pathway membership. Pathways were filtered to be significant (p<=0.01) in at least one sample type and to contain at least two well-modelled KEGG molecular functions and two well-modelled metabolites. The red vertical line corresponds to p=0.05.

Discussion

We have shown that the unsupervised modelling approach of PARAFAC mainly described variation

between subjects, as expected from an algorithm describing the maximum amount of variation that

is present in the data. The modelled variation could be attributed to clinically relevant parameters.

In addition, we have observed that the model corresponding to the salivary metabolomics samples

partly describes the salivary dilution despite correcting for it using Probabilistic Quotient

Normalization (Dieterle et al., 2006). This might be an artifact due to an incorrect assumption that

correcting for water content of saliva can be done in the same way as for urine metabolomics

samples. Future work on this topic should focus on the appropriateness of using Probabilistic

Quotient Normalization prior to performing a decomposition analysis. Other multi-way approaches

exist to describe a specific type of variation (Bro, 1997; Carroll & Chang, 1970; Harshman, 1970).

For example, in N-way partial least squares (NPLS) the model is constrained to regress on an

outcome variable (Bro, 1996). Additionally, the PARAFAC models of the microbiome and

metabolome could be created in a linked fashion by keeping one of the modes equal to each other,

known as coupled matrix and tensor factorization (Acar et al., 2014, 2015; Singh & Gordon, 2008).

Each of these approaches might yield new insights as the models explicitly look at variation that is

common between the datasets. Further research is needed to compare such methods to PARAFAC

in the context of multi-omics data.

We observed that the microbiome PARAFAC models partly describe richness and evenness. This

result gives further evidence of a relationship between oral health and microbial diversity which

has been described before (Anderson et al., 2018; Schoilew et al., 2019; Thomas et al., 2014). While

the methodology used in this study targeted only bacteria, while other members of the oral

microbiome – such as viruses and fungi – might also be relevant in gingivitis onset and progression

(Baker et al., 2017; Nobbs & Jenkinson, 2015). We identified several bacterial subcommunities

within the oral sites that responded similarly to the intervention (**Table 2**). For example, a group of

ASVs was significantly more abundant in individuals in the high responder group compared to low

responders and represented species of known pathogenic or pro-inflammatory genera such as

Actinomyces, *Campylobacter*, *Capnocytophaga*, *Fusobacterium*, *Leptotrichia*, and *Porphyromonas*

(Huang et al., 2014; Kirst et al., 2015; Kistler et al., 2013; Schincaglia et al., 2017; The Human

Microbiome Project Consortium, 2012). Clusters of ASVs that were significantly more abundant in

low responders compared to high responders contained known commensal species related to oral

health, such as *Kingella oralis* (Abusleme et al., 2021), *Streptococcus* spp. (Caporaso et al., 2011;

Zaura et al., 2009), and *Veillonella* spp. (Moore & Moore, 1994). It has to be noted that sequencing

based only on the V4 region of the 16S rRNA gene may not provide sufficient resolution to correctly

assign pathogenicity or pro-inflammatory traits all ASVs. Similarly, some ASV have been assigned

to bacterial taxa that are not commonly observed in the oral cavity; this may indicate that these

bacteria were present in the sample due problematic hygiene in the host, or be due to classification

errors. Finally, we observed a difference in ecosystem stability between high and low responders in

some parts of the oral cavity. This suggests that in some subjects, the ecosystem can withstand the

ecological pressure of plaque accumulation and has some mechanisms that prevents healthy

microbiota from decreasing in abundance. Further research is needed to identify these mechanisms

and the microbiota involved.

By integrating the PARAFAC models of the salivary metabolomics data and of the predicted

metabolic functions from the microbiome data, we identified enriched pathways that reflected the

changing proportions of Gram-positive and -negative microbiota (cell membrane/wall component

pathways, such as lipopolysaccharide and peptidoglycan biosynthesis). Quorum sensing systems

were found to be significantly enriched in all plaque sites likely due to the presence of various

Streptococcus spp., that carry genes for oligopeptide binding protein quorum sensing system

(Fontaine et al., 2010; Gardan et al., 2009). These are known to regulate biofilm formation and

virulence (Hammer & Bassler, 2003; Kong et al., 2006; Preda & Săndulescu, 2019). The strong signal

in carbon metabolism and oxidative phosphorylation were likely reflective of the variation in

anaerobic bacteria, which accumulate during gingivitis (Chen et al., 2019; Diaz et al., 2002; Loesche,

1969). An interesting link between salivary metabolites and microbiome functional potential may

have been observed in the nucleoside/purine/pyrimidine biosynthesis pathways: while urea was one

of the best modelled salivary metabolites and is a potential biomarker for oral health (D'souza et al.,

2023; Gaál Kovalčíková et al., 2019; Nascimento et al., 2009; Osmani, 2023), a role of salivary

nucleosides in plaque formation or gingivitis is as yet unknown. While many of these functions can

be identified as housekeeping activities, we protected ourselves from wrongfully identifying such

generic activities as enriched by (1) using SetRank to correct for multiple pathway membership and

(2) creating a custom database of pathway elements and removing very large pathways that often

connect to such activities. Further research is needed to investigate the relevance of these enriched

pathways in the context of dental plaque formation and gingivitis.

In conclusion, we show that PARAFAC modelling of longitudinal oral microbiome and salivary

metabolomics data can be used to identify clinically relevant variation and similarly responding

microbial subcommunities related to gingivitis. By performing high level integration of predicted

microbiome functions and the salivary metabolome, we highlight biochemical pathways of oral

biofilm formation and maturation with a likely role in gingivitis.

Methods

Data description

The gingivitis challenge study was carried out at the Academic Centre for Dentistry Amsterdam

(ACTA) with 15 systemically healthy males and 26 systemically healthy females between the ages

of 18 and 55. Recruitment details and exclusion criteria were previously described (Prodan et al.,

2016). Reversible gingivitis was induced by omission of oral hygiene during the two-week gingivitis

challenge (day 0 to day 14), which was preceded by a two-week baseline period (day -14 to day 0)

and followed by a one-week resolution phase (day 14 to day 21; **Figure 1A**). Assessment of plaque

formation, bleeding and the collection of oral samples for microbiome were performed throughout

the baseline, challenge and resolution phases. Plaque and bleeding were assessed clinically in a half

mouth randomized contralateral model (Bentley & Disney, 1995). Oral samples for microbiome

analysis were taken at six sites in the mouth: supragingival plaque at the lower and upper jaw lingual

surfaces, supragingival plaque at the lower and upper jaw interproximal surfaces, tongue and saliva.

The salivary metabolome was only sampled during the challenge phase.

Red fluorescence imaging

Acquisition of red fluorescent plaque (RF) photographs has been described previously (Heinrich-

Weltzien et al., 2003; van der Veen et al., 2016; Volgenant et al., 2017). Briefly, fluorescence

photographs were taken of the vestibular aspect of the anterior teeth (cuspid to cuspid, upper and

lower jaw) in end-to-end position at every time point using a QLF-D camera (Inspektor Research

Systems BV, Amsterdam, the Netherlands). The photographs were assessed planimetrically for the

percentage red fluorescent protein (RFP) coverage using RFP analysis software (QA2 V1.25,

Inspektor Research Systems BV, Amsterdam, the Netherlands). Three response groups were

determined using the RF% values on day 14: low (0-1.7%), mid (1.8-6.0%) and high (6.0%+) (**Figure**

1B) (van der Veen et al., 2016).

DNA isolation and 16S rRNA gene amplicon sequencing

DNA was isolated from the samples using a previously described method (Zaura et al., 2017).

Samples were mixed with 300 µl lysis buffer (Agowa, Berlin, Germany), 500 µl phenol saturated

with tris-HCl (pH 8.0) and 500 µl zirconium beads (0.1 mm; BioSpec Products, Bartlesville, OK,

USA), and shaken in a bead beater for 3 min at 2800 oscillations/min. DNA released was purified

using magnetic beads (Agowa, Berlin, Germany) and used for amplicon sequencing.

The V4 hypervariable region of the 16S rRNA gene was targeted using primers F515 (5'- GTG CCA

GCM GCC GCG GTA A -3') and R806 (5'- GGA CTA CHV GGG TWT CTA AT -3'). The primers

included Illumina adapters and a unique 8-nucleotide sample index sequence key (Kozich et al.,

2013). PCR was performed using the Phusion Hot Start II High Fidelity PCR Master Mix (Thermo

Scientific, Waltham, MA, USA) with 100 pg template. The following amplification program was

used: initial denaturation at 98°C for 30 s; 30 cycles of 98°C for 10 s, 55°C for 30 s and 72°C for 30 s;

and final elongation at 72°C for 5 min. The amplicon libraries were pooled in equimolar amounts

and purified using the QIAquick Gel Extraction Kit (Qiagen, Valencia, CA, USA). Amplicon quality

and size were analysed on a Fragment Analyzer (Advanced Analytical Technologies Inc., Ankeny,

IA, USA). Amplicon sequencing was performed on the Illumina MiSeq platform (Illumina

Technologies, San Diego, CA, USA) using 2 x 200 cycle paired-end settings.

16S rRNA gene amplicon data pre-processing

The 16S rRNA gene sequencing data were pre-processed using DADA2 (version 1.14.0; (Callahan et

al., 2016)). All microbiome sampling locations except for saliva were pre-processed with default

DADA2 settings (truncLen = c(100, 100), maxN = 0, maxEE = c(2,2), truncQ = 2, rm.phix = TRUE,

compress = FALSE, matchIDs = TRUE, multithread = TRUE). The saliva samples were processed

with a shorter trimming setting (truncLen = c(200, 215)) due to lower quality reads. Taxonomic

classification of ASVs was performed using the SILVA NR99 (v138) and HOMD (v15.22) reference

databases.

Metabolomics sampling and profiling

Saliva sample collection and metabolite profiling were described previously (Prodan et al., 2016).

Unstimulated saliva was collected at the intervention time points. Participants were instructed to

allow saliva to accumulate on the floor of the mouth and to spit at 30 second intervals into a pre-

weighted 30 ml polypropylene tube. The collection period was 5 minutes. Non-targeted metabolite

profiling of the saliva samples for the gingivitis challenge time points was performed by Metabolon.

Samples were processed as described in Metabolon's standard method (Evans et al., 2009). Raw data

were extracted, peak-identified and quality controlled using Metabolon's hardware and software.

Compounds were identified by comparison with Metabolon's reference library (Lawton et al., 2008).

Metabolites were annotated with KEGG compound IDs (Kanehisa et al., 2021; Kanehisa & Goto,

2000).

Assessment of clinical parameters of gingivitis

Plaque and bleeding scores were clinically assessed by two experts in a half mouth randomized

contralateral model, as described previously (van der Veen et al., 2016). Plaque was quantified using

a modified Silness & Loë Plaque Index at six sites of the buccal and lingual aspects of all present

teeth (Silness & Löe, 1964). Gingival bleeding was quantified using the bleeding on marginal probing

index on six gingival areas on the buccal and lingual sides of all present teeth (Van der Weijden et

al., 1993).

Statistical Analysis - Microbiome data processing

The ASV data and taxonomic information were processed using MATLAB (version 2022a). The data

were separated by oral site. To limit the sparsity per dataset, ASVs were kept if the sparsity was

<50% in any response group (**Supplementary Figure A6**). All other ASVs were removed from the

data. By setting a response group-based sparsity-cutoff per ASV, we avoid removing biologically

relevant ASVs that would occur in only one or two response groups. ASVs were also removed if

they corresponded to chloroplast or mitochondrial sequences, as these were not relevant for the

study. An overview of the number of ASVs before and after filtering per sample type is reported in

Supplementary Table B5. Subsequently, a centered-log ratio transformation - using a pseudo-count

of 1 - was performed to correct for compositionality (Aitchison, 1982; Gloor et al., 2017). The

datasets were then converted to three-way arrays (**Figure 1D**). Missing samples were kept in the

data cube as a row of missing values (**Supplementary Table B6**). This is possible because PARAFAC

interpolates the missing data automatically in its alternating least-squares algorithm and maximises

the amount of information for the modelling procedure as the other samples of the subject do not

have to be removed entirely. Subsequently, centering across the subject mode and scaling within

the ASV mode - ignoring missing values - was performed to make the samples comparable at every

timepoint and the ASVs comparable for all time points (Bro & Smilde, 2003).

Statistical Analysis - Salivary metabolomics data processing

The salivary metabolomics data were processed using MATLAB (version 2022a). To limit the

sparsity per dataset, metabolites were excluded if they contained more than 25% values below the

detection limit (**Supplementary Figure A7**). Additionally, xenobiotic compounds were manually

assessed for occurrence across response groups and selected if they were prevalent in most subjects

(**Supplementary Data**). After feature selection, 400 of the 499 metabolites remained (**Supplementary**

Table B5). Values below the detection limit were imputed with a random value between 0 and the

detection limit per metabolite to preserve their distribution. To correct for the dilution caused by

the amount of water in the saliva samples, Probabilistic Quotient Normalization (PQN) was

performed using the median value of each metabolite as an artificial reference sample (Dieterle et

al., 2006). Next, the dataset was (natural) log transformed to stabilize the variance. The dataset was

then converted to a three-way array. Subsequently, centering across the subject mode and scaling

within the ASV mode - ignoring missing values - was performed to make the samples comparable

at every timepoint and the ASVs comparable for all time points (Bro & Smilde, 2003). All subjects

were fully sampled, except for subject 3CN8CB for whom no metabolome data was available

(**Supplementary Table B6**).

Functional profile prediction of microbiome data

A functional profile prediction based on the microbiome ASV count data was performed using the

Tax4Fun2 package in R (Tax4Fun2 version 1.1.5, R version 4.0.3) (Aßhauer et al., 2015; Wemheuer

et al., 2020). ASVs corresponding to Chloroplast or Mitochondria were removed. Samples were

rarefied to 10 000 reads to make them comparable (**Supplementary Figures A8** and **A9**). Samples with

fewer total reads were removed. This step removed 53 of 1 692 samples. ASVs without counts after

rarefying were removed. All steps together removed 6 903 out of 14 014 ASVs. Tax4Fun2 was run

with default settings against the included Ref99NR database (Wemheuer et al., 2020) and a custom

database based on the HOMD genomes (v9.14, accessed 2020-11-09) (Chen et al., 2010). Molecular

functions of the genomes were assigned using Tax4Fun2's DIAMOND wrapper (Buchfink et al.,

2015). HOMD-annotated 16S rRNA gene sequences were extracted from the genomes, cut to the

515F-806R fragment used in this study using cutadapt v1.18 (Martin, 2011). Exact duplication

within genomes were removed, and 16S rRNA gene sequence fragments were combined with the

functional assignments using Tax4Fun's generateUserDataByClustering function. The resulting

functional prediction profiles - which contain values between 0 and 1 for each function - were used

for subsequent analysis. The fraction of unused taxonomic units and the fraction of unused

sequences are reported per sample (**Supplementary Figures A10** and **A11**).

Statistical Analysis - Processing of functional profile predictions

The functionally predicted microbiome data were processed in MATLAB (version 2022a). The data

were separated by sample type. To make the data comparable to the salivary metabolomics data,

predicted KEGG orthologous groups (KOs) were removed if they did not belong to pathways that

the metabolites mapped to. This was done by mapping the KOs and metabolite compound IDs to

pathways using the KEGG API (Kawashima et al., 2003). To limit the sparsity per dataset, KOs were

removed if the number of zeroes for all response groups was >50% (**Supplementary Figure A12**).

Additionally, KOs were removed if the sum-of-squares was lower than 0.025% of the total sum-of-

squares of the dataset (**Supplementary Figure A13**). This latter filter was to ensure that the modelling

procedure focused on relevant variation in the data and made the total number of variables

comparable to the metabolomics data, while retaining most information. An overview of the

number of KOs remaining after feature selection is reported (**Supplementary Table B5**).

Subsequently a centered-log ratio transformation of each dataset was performed to correct for

compositionality (Aitchison, 1982; Gloor et al., 2017). The dataset was then converted to a three-

way array. Missing samples were kept in the data cube as a row of missing values (**Supplementary

Table B6**). Subsequently, centering across the subject mode and scaling within the ASV mode -

ignoring missing values - was performed to make the samples comparable at every timepoint and

the ASVs comparable for all time points (Bro & Smilde, 2003).

Statistical Analysis - Parallel Factor Analysis (PARAFAC)

Details on the creation of PARAFAC models for microbiome (Martino et al., 2021) and

metabolomics data (Bro, 1997) have been described elsewhere. The PARAFAC implementation

from the N-way toolbox (version 1.8.0.0) in MATLAB (version 2022a) was used to create PARAFAC

models for all datasets (Bro, accessed 26-10-2023). Similar to PCA, the correct number of

components of the PARAFAC model needed to be determined to create an optimal model (Bro &

Kiers, 2003; van der Ploeg et al., 2024). This was done by inspecting the number of iterations needed

to converge (Bro, 1997), the core consistency diagnostic (CORCONDIA) (Bro & Kiers, 2003), the

variation explained (Bro, 1997), and the Tucker congruence coefficient per mode (Lorenzo-Seva &

ten Berge, 2006; Tucker, 1951). Additionally, a jack-knife approach was used to determine the

stability of the PARAFAC modelling procedure. These metrics were inspected per component for every generated model (**Supplementary Data**). All generated models are reported in **Supplementary Figures A14-A26**. An overview of the model statistics also reported (**Supplementary Table B7**). Congruence loadings, which describe the relationships between the original variables of the dataset and the latent variables of the corresponding model, were calculated for every component in every sample type (Lorho et al., 2006). While the metrics above suggested a one-component PARAFAC model corresponding to the microbiome data at the lower jaw interproximal niche, the model described less than 10% of the variation in the data. Furthermore, manual inspection revealed that the model did not describe variation that we could explain biologically. Hence, we did not analyse this model further in subsequent steps.

A transformation of the subject loadings was required for comparison with the longitudinally measured clinical parameters of gingivitis. The ASV loading vectors were orthonormalized using the Gram-Schmidt orthonormalization procedure in the pracma R package (version 2.4.2) (Borchers & Borchers, 2022; Schmidt, 1989). The (same) transformation matrix is then applied to the Kronecker product of the subject and the time loadings to obtain interpretable loadings of every subject-time combination (Kiers, 2000). The correlation of the time-resolved subject loadings with the clinical parameters of gingivitis was then tested (**Table 1** and **Supplementary Table B1**).

Statistical Analysis - Microbiome ASV cluster analysis

Loading plots of the ASV mode and the subject mode for each of the microbiome PARAFAC models were created (**Figure 2**). ASVs were filtered out if they had a variation explained lower than the average of the model or a congruence loading lower than 0.4 (Lorho et al., 2006). ASVs were clustered based on their fitted abundances according to the PARAFAC model using the K-medoids algorithm from the cluster R package (version 2.1.4) with 50 random starts to be robust against outliers (Maechler, 2019). The number of clusters was determined using the within-cluster sum of squares, silhouette width and gap statistic metrics as reported by the factoextra R package (version 1.0.7, **Supplementary Figure A27**) (Kassambara & Mundt, 2017). An overview of the taxonomic information per cluster can be found in **Table 2**. Bacteria in **Table 2** were only reported if two criteria were met: (1) the taxonomic classification was resolved to species level by DADA2 using the SILVA (v138) and HOMD (v15.22) databases and (2) the annotation agreed with a separate best hit annotation run using the HOMD (v15.22) database. Cases where the classification was resolved to

species level by the DADA2 run but only to the genus level by the separate HOMD run are also

reported.

The response to the gingivitis intervention of the ASV clusters was shown using the relative

abundance sum per ASV cluster derived from the original count data (**Figure 3** and **Supplementary**

Figure A3). The mean of the summed relative abundances and standard error of the mean were

calculated per response group for a given sample type and time point. The mean difference between

the high and low responders was tested using a permutation analysis where the response group

membership of the subjects was permuted (**Supplementary Table B2**).

Statistical Analysis - Functional enrichment analysis

The PARAFAC models of the functionally profiled microbiome and salivary metabolomics data

were combined to perform a pathway enrichment test per sample type. This was done by calculating

the variance explained and congruence loadings per KO or metabolite in each model (Lorho et al.,

2006). If the PARAFAC model contained two components, the maximum of the two congruence

loadings per feature was used. The variance explained and congruences were normalized by the

maximum value per model prior to integration. The normalized variance explained and normalized

congruence per feature were averaged to obtain a ranking of pathway elements (KOs and

metabolites). This causes the best modelled KO and compound ID to be at the top of this list. The

ranked pathway element list was cut off at 75% of its length to ensure a good separation between

well-modelled and averagely-modelled pathway elements (**Supplementary Figure A28**).

The SetRank R package (version 1.1.0) was used to obtain pathways of interest (Simillion et al.,

2017). SetRank is a gene set enrichment algorithm that obtains a high sensitivity by correcting for

multiple pathway membership. A custom database of pathway elements was created by combining

the KO and compound ID to pathway mappings obtained through the KEGG API (**Supplementary**

Data). Pathways were removed from our database if they could not be performed by prokaryotes.

Generic, large pathways with more than 750 elements were removed to ensure that the enrichment

results were specific enough for interpretation (**Supplementary Figure A29**). Subsequently the

SetRank analysis was performed with standard settings per sample type. The multiple-pathway

corrected p-values are reported ($p \leq 0.05$, **Figure 4** and **Supplementary Table B4**)

Data Availability

The ASV abundance data is available at https://github.com/GRvanderPloeg/TIFN-multiway. Raw
sequencing data is available by request from the authors, in accordance with the informed consent
signed by the study participants.

Code Availability

The underlying code for this study is available on GitHub and can be accessed via
https://github.com/GRvanderPloeg/TIFN-multiway/releases/tag/v1.2.

Ethics statement

The study involving human participants was conducted in accordance with the ethical principles of the
64th WMA Declaration of Helsinki (October 2013, Brazil) and the Medical Research Involving Human
Subjects Act (WMO), approximating Good clinical Practice (CPMP/ICH/135/95) guidelines. The clinical
trial was approved by the Medical Ethical Committee of the VU Medical Center (2014.505) and
registered at the public trial register of the Central Committee on Research Involving Human Subjects
(CCMO) under number NL51111.029.14.

Author Contributions

G.R. van der Ploeg: formal analysis, methodology, writing - original draft; B.W. Brandt: investigation,
writing - review & editing; B.J.F. Keijser: investigation, project administration, writing - review &
editing; M.H. van der Veen: investigation, project administration, writing - review & editing; C.M.C.
Volgenant: investigation, writing - review & editing; E. Zaura: conceptualization, investigation, project
administration, funding acquisition, writing - review & editing; A.K. Smilde: conceptualization,
methodology, supervision, funding acquisition, writing - review & editing; J.A. Westerhuis:
conceptualization, methodology, supervision, formal analysis, writing - original draft; A. Heintz-
Buschart: conceptualization, formal analysis, supervision, project administration, writing - original
draft. All authors reviewed and approved the manuscript.

Competing interests

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

Acknowledgements

We want to thank Michelle van der Wurff (TNO) and Tim van den Broek (TNO) for their help in data
acquisition, processing and storing and Jesse Alderliesten (UU/UvA), Fred White (UvA), Cynthia

Albracht (UvA), Juan Pablo Bascur (CWTS) and Rianne Warmerdam (Aiden) for their useful discussions

and suggestions. The authors thank N.A.M. Rosema for coordinating the clinical study. For assistance

in capturing fluorescence photographs, the authors thank Y. Altindağ. For the clinical examinations the

authors thank J.M. Voll (clinical plaque assessment with Silness & Löe), and S. Bizzarro (assessment of

the bleeding on marginal probing).

This research is supported by the Dutch Technology Foundation STW (project number 10948) and the

Top Institute Food and Nutrition (TIFN), a public-private partnership on precompetitive research in

food and nutrition. Organizations supporting this project had no role in study design, data collection

and analysis, decision to publish, or preparation of the manuscript. QLF data was obtained with funding

from NWO (ZonMw-STW-NIG-program): "Project 10948: Seeing is believing? A novel tool for the

visualization of oral disease manifestations." GRvdP was funded by a grant from the University of

Amsterdam, Research Priority Area on Personal Microbiome Health.

527 References

Abusleme, L., Hoare, A., Hong, B., & Diaz, P. I. (2021). Microbial signatures of health, gingivitis, and

periodontitis. *Periodontology 2000*, *86*(1), 57–78. https://doi.org/10.1111/prd.12362

Acar, E., Bro, R., & Smilde, A. (2015). Data Fusion in Metabolomics Using Coupled Matrix and Tensor

Factorizations. *Proceedings of the IEEE*, *103*, 1602.

https://doi.org/10.1109/JPROC.2015.2438719

Acar, E., Papalexakis, E. E., Gürdeniz, G., Rasmussen, M. A., Lawaetz, A. J., Nilsson, M., & Bro, R. (2014).

Structure-revealing data fusion. *BMC Bioinformatics*, *15*(1), 239.

https://doi.org/10.1186/1471-2105-15-239

Aitchison, J. (1982). The Statistical Analysis of Compositional Data. *Journal of the Royal Statistical*

Society: Series B (Methodological), *44*(2), 139–160. https://doi.org/10.1111/j.2517-

6161.1982.tb01195.x

Anderson, A. C., Rothballer, M., Altenburger, M. J., Woelber, J. P., Karygianni, L., Lagkouvardos, I.,

Hellwig, E., & Al-Ahmad, A. (2018). In-vivo shift of the microbiota in oral biofilm in response to

frequent   sucrose   consumption.   *Scientific   Reports*,   *8*(1),   Article   1.

https://doi.org/10.1038/s41598-018-32544-6

Aßhauer, K. P., Wemheuer, B., Daniel, R., & Meinicke, P. (2015). Tax4Fun: Predicting functional profiles

from   metagenomic   16S   rRNA   data.   *Bioinformatics*,   *31*(17),   2882–2884.

https://doi.org/10.1093/bioinformatics/btv287

Axelsson, P., Lindhe, J., & Nyström, B. (1991). On the prevention of caries and periodontal disease.

Journal   of   Clinical   Periodontology,   *18*(3),   182–189.   https://doi.org/10.1111/j.1600-

051X.1991.tb01131.x

Baker, J. L., Bor, B., Agnello, M., Shi, W., & He, X. (2017). Ecology of the Oral Microbiome: Beyond

Bacteria. *Trends in Microbiology*, *25*(5), 362–374. https://doi.org/10.1016/j.tim.2016.12.012

Bentley, C. D., & Disney, J. A. (1995). A comparison of partial and full mouth scoring of plaque and

gingivitis   in   oral   hygiene   studies.   *Journal   of   Clinical   Periodontology*,   *22*(2),   131–135.

https://doi.org/10.1111/j.1600-051X.1995.tb00124.x

Bergström, J., & Preber, H. (1986). The influence of cigarette smoking on the development of

experimental   gingivitis.   *Journal   of   Periodontal   Research*,   *21*(6),   668–676.

https://doi.org/10.1111/j.1600-0765.1986.tb01504.x

Borchers,   H. W.,   &   Borchers,   M. H. W.   (2022).   Package   'pracma.'   *Accessed   On*,   *4*.

https://mirror.las.iastate.edu/CRAN/web/packages/pracma/pracma.pdf

Bro,   R.   (n.d.).   *The   N-way   Toolbox*.   Retrieved   October   26,   2023,   from

https://nl.mathworks.com/matlabcentral/fileexchange/1088-the-n-way-toolbox

Bro, R. (1996). Multiway calibration. Multilinear PLS. *Journal of Chemometrics*, *10*(1), 47–61.

https://doi.org/10.1002/(SICI)1099-128X(199601)10:1<47::AID-CEM400>3.0.CO;2-C

Bro, R. (1997). PARAFAC. Tutorial and applications. *Chemometrics and Intelligent Laboratory Systems*.

Bro, R., & Kiers, H. (2003). A new Efficient Method for Determining the Number of Components in

PARAFAC Models. *Journal of Chemometrics*, *17*, 274–286. https://doi.org/10.1002/cem.801

Bro, R., & Smilde, A. K. (2003). Centering and scaling in component analysis. *Journal of Chemometrics*,

17(1), 16–33. https://doi.org/10.1002/cem.773

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND.

Nature Methods, *12*(1), 59–60.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016).

DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*,

13(7), Article 7. https://doi.org/10.1038/nmeth.3869

Caporaso, J. G., Lauber, C. L., Costello, E. K., Berg-Lyons, D., Gonzalez, A., Stombaugh, J., Knights, D.,

Gajer, P., Ravel, J., Fierer, N., Gordon, J. I., & Knight, R. (2011). Moving pictures of the human

microbiome. *Genome Biology*, *12*(5), R50. https://doi.org/10.1186/gb-2011-12-5-r50

Carroll, J. D., & Chang, J.-J. (1970). Analysis of individual differences in multidimensional scaling via an

n-way generalization of "Eckart-Young" decomposition. *Psychometrika*, *35*(3), 283–319.

https://doi.org/10.1007/BF02310791

Chapple, I. L. C., Mealey, B. L., Van Dyke, T. E., Bartold, P. M., Dommisch, H., Eickholz, P., Geisinger, M.

L., Genco, R. J., Glogauer, M., Goldstein, M., Griffin, T. J., Holmstrup, P., Johnson, G. K., Kapila,

Y., Lang, N. P., Meyle, J., Murakami, S., Plemons, J., Romito, G. A., … Yoshie, H. (2018).

Periodontal health and gingival diseases and conditions on an intact and a reduced

periodontium: Consensus report of workgroup 1 of the 2017 World Workshop on the

Classification of Periodontal and Peri-Implant Diseases and Conditions. *Journal of*

Periodontology, *89*, S74–S84. https://doi.org/10.1002/JPER.17-0719

Chapple, I. L. C., Van der Weijden, F., Doerfer, C., Herrera, D., Shapira, L., Polak, D., Madianos, P.,

Louropoulou, A., Machtei, E., Donos, N., Greenwell, H., Van Winkelhoff, A. J., Eren Kuru, B.,

Arweiler, N., Teughels, W., Aimetti, M., Molina, A., Montero, E., & Graziani, F. (2015). Primary

prevention of periodontitis: Managing gingivitis. *Journal of Clinical Periodontology*, *42*(S16),

S71–S76. https://doi.org/10.1111/jcpe.12366

Chen, M., Cai, W., Zhao, S., Shi, L., Chen, Y., Li, X., Sun, X., Mao, Y., He, B., Hou, Y., Zhou, Y., Zhou, Q.,

Ma, J., & Huang, S. (2019). Oxidative stress-related biomarkers in saliva and gingival crevicular

fluid associated with chronic periodontitis: A systematic review and meta-analysis. *Journal of*

Clinical Periodontology, *46*(6), 608–622. https://doi.org/10.1111/jcpe.13112

Chen, T., Yu, W.-H., Izard, J., Baranova, O. V., Lakshmanan, A., & Dewhirst, F. E. (2010). The Human Oral

Microbiome Database: A web accessible resource for investigating oral microbe taxonomic and

genomic information. *Database*, *2010*.

https://academic.oup.com/database/article/doi/10.1093/database/baq013/405450?view=lo

ng

Curtis, M. A., Diaz, P. I., & Van Dyke, T. E. (2020). The role of the microbiota in periodontal disease.

Periodontology 2000, *83*(1), 14–25. https://doi.org/10.1111/prd.12296

Dawes, C., & Wong, D. T. W. (2019). Role of Saliva and Salivary Diagnostics in the Advancement of Oral

Health. *Journal of Dental Research*, *98*(2), 133–141.

https://doi.org/10.1177/0022034518816961

Diaz, P. I., Hoare, A., & Hong, B.-Y. (2016). Subgingival Microbiome Shifts and Community Dynamics in

Periodontal Diseases. *Journal of the California Dental Association*, *44*(7), 421–435.

https://doi.org/10.1080/19424396.2016.12221035

Diaz, P. I., Zilm, P. S., & Rogers, A. H. (2002). Fusobacterium nucleatum supports the growth of

Porphyromonas gingivalis in oxygenated and carbon-dioxide-depleted environments.

Microbiology, *148*(2), 467–472. https://doi.org/10.1099/00221287-148-2-467

Dieterle, F., Ross, A., Schlotterbeck, G., & Senn, H. (2006). Probabilistic Quotient Normalization as

Robust Method to Account for Dilution of Complex Biological Mixtures. Application in 1H NMR

Metabonomics. *Analytical Chemistry*, *78*(13), 4281–4290. https://doi.org/10.1021/ac051632c

D'souza, L. L., Lawande, S. A., Samuel, J., & Pinto, M. J. W. (2023). Effect of salivary urea, pH and

ureolytic microflora on dental calculus formation and its correlation with periodontal status.

Journal of Oral Biology and Craniofacial Research, *13*(1), 8–12.

Evans, A. M., DeHaven, C. D., Barrett, T., Mitchell, M., & Milgram, E. (2009). Integrated, Nontargeted

Ultrahigh Performance Liquid Chromatography/Electrospray Ionization Tandem Mass

Spectrometry Platform for the Identification and Relative Quantification of the Small-Molecule

Complement of Biological Systems. *Analytical Chemistry*, *81*(16), 6656–6667.

https://doi.org/10.1021/ac901536h

Fontaine, L., Boutry, C., De Frahan, M. H., Delplace, B., Fremaux, C., Horvath, P., Boyaval, P., & Hols, P.

(2010). A Novel Pheromone Quorum-Sensing System Controls the Development of Natural

Competence in *Streptococcus thermophilus* and *Streptococcus salivarius*. *Journal of*

Bacteriology, *192*(5), 1444–1454. https://doi.org/10.1128/JB.01251-09

Gaál Kovalčíková, A., Pančíková, A., Konečná, B., Klamárová, T., Novák, B., Kovaľová, E., Podracká, Ľ.,

Celec, P., & Tóthová, Ľ. (2019). Urea and creatinine levels in saliva of patients with and without

periodontitis. *European Journal of Oral Sciences*, *127*(5), 417–424.

https://doi.org/10.1111/eos.12642

Gardan, R., Besset, C., Guillot, A., Gitton, C., & Monnet, V. (2009). The Oligopeptide Transport System

Is Essential for the Development of Natural Competence in *Streptococcus thermophilus* Strain

LMD-9. *Journal of Bacteriology*, *191*(14), 4647–4655. https://doi.org/10.1128/JB.00257-09

Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., & Egozcue, J. J. (2017). Microbiome Datasets Are

Compositional: And This Is Not Optional. *Frontiers in Microbiology*, *8*, 2224.

https://doi.org/10.3389/fmicb.2017.02224

Guk, H.-J., Lee, E.-S., Jung, U.-W., & Kim, B.-I. (2020). Red fluorescence of Interdental plaque for

screening of gingival health. *Photodiagnosis and Photodynamic Therapy*, *29*, 101636.

https://doi.org/10.1016/j.pdpdt.2019.101636

Hammer, B. K., & Bassler, B. L. (2003). Quorum sensing controls biofilm formation in *Vibrio cholerae*.

Molecular Microbiology, *50*(1), 101–104. https://doi.org/10.1046/j.1365-2958.2003.03688.x

Han, L., Gao, X., Liao, M., Yu, X., Zhang, R., Liu, S., & Zeng, F. (2020). Hygiene practices among young

adolescents aged 12-15 years in low- and middle-income countries: A population-based study.

Journal of Global Health, *10*(2), 020436. https://doi.org/10.7189/jogh.10.020436

Harshman, R. A. (1970). *Foundations of the PARAFAC procedure: Models and conditions for an"*

explanatory"   multimodal   factor   analysis.

https://www.psychology.uwo.ca/faculty/harshman/wpppfac0.pdf

Heinrich-Weltzien, R., Kühnisch, J., Van Der Veen, M., De Josselin De Jong, E., & Stößer, L. (2003).

Quantitative light-induced fluorescence (QLF)—A potential method for the dental practitioner.

Quintessence International, *34*(3), 181–188.

Huang, S., Li, R., Zeng, X., He, T., Zhao, H., Chang, A., Bo, C., Chen, J., Yang, F., Knight, R., Liu, J., Davis,

C., & Xu, J. (2014). Predictive modeling of gingivitis severity and susceptibility via oral

microbiota. *The ISME Journal*, *8*(9), 1768–1780. https://doi.org/10.1038/ismej.2014.32

Kanehisa, M., Furumichi, M., Sato, Y., Ishiguro-Watanabe, M., & Tanabe, M. (2021). KEGG: Integrating

viruses   and   cellular   organisms.   *Nucleic   Acids   Research*,   *49*(D1),   D545–D551.

https://doi.org/10.1093/nar/gkaa970

Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids*

Research, *28*(1), 27–30. https://doi.org/10.1093/nar/28.1.27

Kassambara, A., & Mundt, F. (2017). Package 'factoextra.' *Extract and Visualize the Results of*

Multivariate   Data   Analyses,   *76*(2).   https://piyanit.nl/wp-

content/uploads/2020/10/factoextra.pdf

Kawashima, S., Katayama, T., Sato, Y., & Kanehisa, M. (2003). KEGG API: A Web Service Using

SOAP/WSDL   to   Access   the   KEGG   System.   *Genome   Informatics*,   *14*,   673–674.

https://doi.org/10.11234/gi1990.14.673

Kiers, H. A. L. (2000). Some procedures for displaying results from three-way methods. *Journal of*

Chemometrics,   *14*(3),   151–170.   https://doi.org/10.1002/1099-

128X(200005/06)14:3<151::AID-CEM585>3.0.CO;2-G

Kilian, M., Chapple, I. L. C., Hannig, M., Marsh, P. D., Meuric, V., Pedersen, A. M. L., Tonetti, M. S.,

Wade, W. G., & Zaura, E. (2016). The oral microbiome – an update for oral healthcare

professionals. *British Dental Journal*, *221*(10), Article 10.

https://doi.org/10.1038/sj.bdj.2016.865

Kirst, M. E., Li, E. C., Alfant, B., Chi, Y.-Y., Walker, C., Magnusson, I., & Wang, G. P. (2015). Dysbiosis and

Alterations in Predicted Functions of the Subgingival Microbiome in Chronic Periodontitis.

Applied and Environmental Microbiology, *81*(2), 783–793.

https://doi.org/10.1128/AEM.02712-14

Kistler, J. O., Booth, V., Bradshaw, D. J., & Wade, W. G. (2013). Bacterial Community Development in

Experimental Gingivitis. *PLoS ONE*, *8*(8), e71227.

https://doi.org/10.1371/journal.pone.0071227

Kong, K.-F., Vuong, C., & Otto, M. (2006). Staphylococcus quorum sensing in biofilm formation and

infection. *International Journal of Medical Microbiology*, *296*(2–3), 133–139.

König, K. G., & Navia, J. M. (1995). Nutritional role of sugars in oral health. *The American Journal of*

Clinical Nutrition, *62*(1), 275S-282S.

Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of a

Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data

on the MiSeq Illumina Sequencing Platform. *Applied and Environmental Microbiology*, *79*(17),

5112–5120. https://doi.org/10.1128/AEM.01043-13

Lawton, K. A., Berger, A., Mitchell, M., Milgram, K. E., Evans, A. M., Guo, L., Hanson, R. W., Kalhan, S.

C., Ryals, J. A., & Milburn, M. V. (2008). Analysis of the adult human plasma metabolome.

Pharmacogenomics, *9*(4), 383–397. https://doi.org/10.2217/14622416.9.4.383

Löe, H., Theilade, E., & Jensen, S. B. (1965). Experimental Gingivitis in Man. *Journal of Periodontology*,

36(3), 177–187. https://doi.org/10.1902/jop.1965.36.3.177

Loesche, W. J. (1969). Oxygen Sensitivity of Various Anaerobic Bacteria. *Applied Microbiology*, *18*(5),

723–727. https://doi.org/10.1128/am.18.5.723-727.1969

Lorenzo-Seva, U., & ten Berge, J. M. F. (2006). Tucker's congruence coefficient as a meaningful index

of factor similarity. *Methodology: European Journal of Research Methods for the Behavioral*

and Social Sciences, *2*(2), 57–64. https://doi.org/10.1027/1614-2241.2.2.57

Lorho, G., Westad, F., & Bro, R. (2006). Generalized correlation loadings: Extending correlation loadings

to congruence and to multi-way models. *Chemometrics and Intelligent Laboratory Systems*,

84(1), 119–125. https://doi.org/10.1016/j.chemolab.2006.04.023

Maechler, M. (2019). Finding groups in data: Cluster analysis extended Rousseeuw et al. *R Package*

Version, *2*(0), 242–248.

Marsh, P. D. (1991). Sugar, fluoride, pH and microbial homeostasis in dental plaque. *Proceedings of the*

Finnish Dental Society. Suomen Hammaslaakariseuran Toimituksia, *87*(4), 515–525.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet. Journal, *17*(1), 10–12.

Martino, C., Shenhav, L., Marotz, C. A., Armstrong, G., McDonald, D., Vázquez-Baeza, Y., Morton, J. T.,

Jiang, L., Dominguez-Bello, M. G., Swafford, A. D., Halperin, E., & Knight, R. (2021). Context-

aware dimensionality reduction deconvolutes gut microbial community dynamics. *Nature*

Biotechnology, *39*(2), 165–168. https://doi.org/10.1038/s41587-020-0660-7

Moore, W. E. C., & Moore, L. V. H. (1994). The bacteria of periodontal diseases. *Periodontology 2000*,

5(1), 66–77. https://doi.org/10.1111/j.1600-0757.1994.tb00019.x

Murakami, S., Mealey, B. L., Mariotti, A., & Chapple, I. L. C. (2018). Dental plaque–induced gingival

conditions. *Journal of Clinical Periodontology*, *45*(S20), S17–S27.

https://doi.org/10.1111/jcpe.12937

Nascimento, M. M., Gordan, V. V., Garvan, C. W., Browngardt, C. M., & Burne, R. A. (2009). Correlations

of oral bacterial arginine and urea catabolism with caries experience. *Oral Microbiology and*

Immunology, *24*(2), 89–95. https://doi.org/10.1111/j.1399-302X.2008.00477.x

Nibali, L., Di Iorio, A., Tu, Y., & Vieira, A. R. (2017). Host genetics role in the pathogenesis of periodontal

disease   and   caries.   *Journal   of   Clinical   Periodontology*,   *44*(S18).

https://doi.org/10.1111/jcpe.12639

Nobbs, A. H., & Jenkinson, H. F. (2015). Interkingdom networking within the oral microbiome. *Microbes*

and Infection, *17*(7), 484–492. https://doi.org/10.1016/j.micinf.2015.03.008

Odom, A. R. (2011). Five questions about non-mevalonate isoprenoid biosynthesis. *PLoS Pathogens*,

7(12), e1002323.

Osmani, F. (2023). Can the salivary urea and stimulated saliva concentration be a marker of periodontal

diseases   in   opioid   users?   A   case-control   study.   *Heliyon*,   *9*(7).

https://www.cell.com/heliyon/pdf/S2405-8440(23)04301-3.pdf

Peres, M. A., Macpherson, L. M. D., Weyant, R. J., Daly, B., Venturelli, R., Mathur, M. R., Listl, S., Celeste,

R. K., Guarnizo-Herreño, C. C., Kearns, C., Benzian, H., Allison, P., & Watt, R. G. (2019). Oral

diseases: A global public health challenge. *The Lancet*, *394*(10194), 249–260.

https://doi.org/10.1016/S0140-6736(19)31146-8

Petersen, P. E., Bourgeois, D., Ogawa, H., Estupinan-Day, S., & Ndiaye, C. (2005). The global burden of

oral diseases and risks to oral health. *Bulletin of the World Health Organization*.

Preda, V. G., & Săndulescu, O. (2019). Communication is the key: Biofilms, quorum sensing, formation

and prevention. *Discoveries*, *7*(3). https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7086079/

Proctor, G. B. (2016). The physiology of salivary secretion. *Periodontology 2000*, *70*(1), 11–25.

https://doi.org/10.1111/prd.12116

Prodan, A., Imangaliyev, S., Brand, H. S., Rosema, M. N. A., Levin, E., Crielaard, W., Keijser, B. J. F., &

Veerman, E. C. I. (2016). Effect of experimental gingivitis induction and erythritol on the

salivary metabolome and functional biochemistry of systemically healthy young adults.

Metabolomics, *12*(9), 147. https://doi.org/10.1007/s11306-016-1096-4

Schincaglia, G. P., Hong, B. Y., Rosania, A., Barasz, J., Thompson, A., Sobue, T., Panagakos, F., Burleson,

J. A., Dongari-Bagtzoglou, A., & Diaz, P. I. (2017). Clinical, Immune, and Microbiome Traits of

Gingivitis and Peri-implant Mucositis. *Journal of Dental Research*, *96*(1), 47–55.

https://doi.org/10.1177/0022034516668847

Schmidt, E. (1989). Über die Auflösung Linearer Gleichungen mit Unendlich Vielen Unbekannten. In A.

Pietsch (Ed.), *Integralgleichungen und Gleichungen mit unendlich vielen Unbekannten* (Vol. 11,

pp. 249–278). Springer Vienna. https://doi.org/10.1007/978-3-7091-9535-2_5

Schoilew, K., Ueffing, H., Dalpke, A., Wolff, B., Frese, C., Wolff, D., & Boutin, S. (2019). Bacterial biofilm

composition in healthy subjects with and without caries experience. *Journal of Oral*

Microbiology, *11*(1), 1633194. https://doi.org/10.1080/20002297.2019.1633194

Silness, J., & Löe, H. (1964). Periodontal Disease in Pregnancy II. Correlation Between Oral Hygiene and

Periodontal Condition. *Acta Odontologica Scandinavica*, *22*(1), 121–135.

https://doi.org/10.3109/00016356408993968

Simillion, C., Liechti, R., Lischer, H. E. L., Ioannidis, V., & Bruggmann, R. (2017). Avoiding the pitfalls of

gene set enrichment analysis with SetRank. *BMC Bioinformatics*, *18*(1), 151.

https://doi.org/10.1186/s12859-017-1571-6

Singh, A. P., & Gordon, G. J. (2008). Relational learning via collective matrix factorization. *Proceedings*

of the 14th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining,

650–658. https://doi.org/10.1145/1401890.1401969

Ter Steeg, P. F., Van Der Hoeven, J. S., De Jong, M. H., Van Munster, P. J. J., & Jansen, M. J. H. (1988).

Modelling the Gingival Pocket by Enrichment of Subgingival Microflora in Human Serum in

Chemostats. *Microbial Ecology in Health and Disease*, *1*(2), 73–84.

https://doi.org/10.3109/08910608809140185

The Human Microbiome Project Consortium. (2012). Structure, function and diversity of the healthy

human microbiome. *Nature*, *486*(7402), 207–214. https://doi.org/10.1038/nature11234

Thomas, A. M., Gleber-Netto, F. O., Fernandes, G. R., Amorim, M., Barbosa, L. F., Francisco, A. L. N.,

Guerra de Andrade, A., Setubal, J. C., Kowalski, L. P., Nunes, D. N., & Dias-Neto, E. (2014).

Alcohol and tobacco consumption affects bacterial richness in oral cavity mucosa biofilms. *BMC Microbiology*, *14*(1), 250. https://doi.org/10.1186/s12866-014-0250-2

Tucker, L. R. (1951). *A method for synthesis of factor analysis studies* (Vol. 984). Educational Testing Service Princeton, NJ. https://apps.dtic.mil/sti/pdfs/AD0047524.pdf

van der Ploeg, G. R., Westerhuis, J. A., Heintz-Buschart, A., & Smilde, A. K. (2024). *parafac4microbiome: Exploratory analysis of longitudinal microbiome data using Parallel Factor Analysis*. https://doi.org/10.1101/2024.05.02.592191

van der Veen, M. H., Volgenant, C. M. C., Keijser, B., ten Cate, J. (Bob) M., & Crielaard, W. (2016). Dynamics of red fluorescent dental plaque during experimental gingivitis—A cohort study. *Journal of Dentistry*, *48*, 71–76. https://doi.org/10.1016/j.jdent.2016.02.010

Van Der Velden, U., Kuzmanova, D., & Chapple, I. L. C. (2011). Micronutritional approaches to periodontal therapy. *Journal of Clinical Periodontology*, *38*(s11), 142–158. https://doi.org/10.1111/j.1600-051X.2010.01663.x

Van der Weijden, G. A., Timmerman, M. F., Nijboer, A., Lie, M. A., & Van der Velden, U. (1993). A comparative study of electric toothbrushes for the effectiveness of plaque removal in relation to toothbrushing duration: Timerstudy. *Journal of Clinical Periodontology*, *20*(7), 476–481. https://doi.org/10.1111/j.1600-051X.1993.tb00394.x

Volgenant, C. M. C., Zaura, E., Brandt, B. W., Buijs, M. J., Tellez, M., Malik, G., Ismail, A. I., ten Cate, J. M., & van der Veen, M. H. (2017). Red fluorescence of dental plaque in children—A cross-sectional study. *Journal of Dentistry*, *58*, 40–47. https://doi.org/10.1016/j.jdent.2017.01.007

Wemheuer, F., Taylor, J. A., Daniel, R., Johnston, E., Meinicke, P., Thomas, T., & Wemheuer, B. (2020). Tax4Fun2: Prediction of habitat-specific functional profiles and functional redundancy based on 16S rRNA gene sequences. *Environmental Microbiome*, *15*(1), 11. https://doi.org/10.1186/s40793-020-00358-7

WHO. (2022). *Global oral health status report: Towards universal health coverage for oral health by 2030*. WHO. https://www.who.int/publications/i/item/9789240061484

Widström, E., Eaton, K., & Vanobbergen, J. (2004). Oral healthcare systems in the Extended European

Union, partim:[Oral Health care system in] Belgium. *Oral Health & Preventive Dentistry*, *2*(3),

155–157.

Zaura, E., Brandt, B. W., Prodan, A., Teixeira de Mattos, M. J., Imangaliyev, S., Kool, J., Buijs, M. J.,

Jagers, F. L., Hennequin-Hoenderdos, N. L., Slot, D. E., Nicu, E. A., Lagerweij, M. D., Janus, M.

M., Fernandez-Gutierrez, M. M., Levin, E., Krom, B. P., Brand, H. S., Veerman, E. C.,

Kleerebezem, M., … Keijser, B. J. (2017). On the ecosystemic network of saliva in healthy young

adults. *The ISME Journal*, *11*(5), 1218–1231. https://doi.org/10.1038/ismej.2016.199

Zaura, E., Keijser, B. J., Huse, S. M., & Crielaard, W. (2009). Defining the healthy" core microbiome" of

oral microbial communities. *BMC Microbiology*, *9*(9).

804
805

Tables

Table 1: Overview of the correlation test results between the time-resolved subject loadings of the PARAFAC models and the clinical parameters of gingivitis and microbiome diversity metrics. Red fluorescence was assessed at every time point. Plaque and bleeding scores were only assessed at day -14, 0, 14 and 21 of the study. Plaque%: the percentage of sites in the mouth that were covered in plaque. Bleeding%: the percentage of sites in the mouth that bled upon probing. RF%: the percentage of sites that fluoresced red. Richness: the number of nonzero counts in a sample. Evenness: Shannon diversity divided by log(richness). Benjamini-Hochberg corrected p-values of Pearson correlation tests between the subject loadings of one component and the clinical parameters are reported (*: p≤0.05; **: p≤0.01; ***: p≤0.001).

Sample type	Component	Plaque%	Bleeding%	RF%	Richness	Evenness
Upper jaw, lingual	1	0.024*	0.49	8.5e-6***	0.015*	0.032*
	2	0.002**	0.26	4.2e-7***	0.78	1.1e-6***
Upper jaw, interproximal	1	0.0015**	0.032*	8.1e-4***	4.4e-9***	7.0e-9***
	2	0.53	0.15	0.95	0.61	0.94
Lower jaw, lingual	1	4.9e-4***	0.14	1.2e-8***	1.6e-17***	1.5e-12***
Tongue	1	0.78	0.48	0.30	0.024*	0.057
	2	0.85	0.057	0.15	5.3e-11***	0.028*
Saliva	1	0.66	0.20	0.11	0.61	0.78
	2	0.78	0.81	0.78	7.3e-14***	8.5e-6***

Table 2: Overview of all identified species per ASV cluster. ASVs within the same cluster show a similar response over time to the gingivitis intervention. Clustered ASVs are only reported here if two criteria were met: (1) the taxonomic classification was resolved to species level by DADA2 using the SILVA (v138) and HOMD (v15.22) databases and (2) the annotation agreed with a separate best hit annotation run using the HOMD (v15.22) database. Cases where the classification was resolved to species level by the DADA2 run but only to the genus level by the separate HOMD run are also reported.

Sample type	Cluster	Number of members	Identified species
Upper jaw, lingual	1	9	*Actinomyces naeslundii/oris/viscosus, Kingella oralis, Rothia dentocariosa, Streptococcus cristatis/sinensis, Veillonella dispar/parvula, Veillonella atypica/dispar, Veillonella parvula/rogosae/tobetsuensis*
	2	10	*Fusobacterium canifelinus/nucleatum, Fusobacterium massiliense, Granulicetella elegans, Prevotella nanceiensis, Veillonella massiliensis*
Upper jaw, interproximal	1	3	*Rothia dentocariosa, Veillonella atypica/dispar*
	2	12	*Abiotrophia defectiva, Aggregatibacter aphrophilus/kilianii, Capnocytophaga sputigena, Lachnoanaerobaculum cf., Lautropia mirabilis, Rothia aeria/dentocariosa*
	3	5	*Corynebacterium matruchotii, Leptotrichia hofstaii, Leptotrichia shahii/wadei*
	4	11	*Aggregatibacter segnis, Capnocytophaga granulosa, Capnocytophaga leadberri, Cardiobacterium hominis, Fusobacterium canifelinus/nucelatum, Fusobacterium nucleatum, Leptotrichia buccalis, Porphyromonas pasteri*
Lower jaw, lingual	1	21	*Actinomyces georgiae/pacaensis, Aggregatibacter segnis, Campylobacter concisus, Capnocytophaga granulosa, Capnocytophaga leadbetteri, Cardiobacterium hominis, Corynebacterium matruchotii, Fusobacterium canifelinus/nucleatum, Fusobacterium nucleatum, Fusobacterium hwasooki/nucleatum/periodonticum, Lachnoanaerobaculum cf., Leptotrichia hofstadii, Porphyromonas pasteri*
	2	1	*Streptococcus salivarius* (putative)
Tongue	1	21	*Actinomyces lingnae/marseillensis/pacaensis, Capnocytophaga leadbetteri, Fusobacterium periodonticum, Granulicatella adiacens/para-adiacens, Lachnoanaerobaculum cf., Oribacterium parvum, Oribacterium sinus, Peptostreptococcus stomatis, Porphyromonas pasteri, Prevotella nanceiensis, Veillonella parvula/rogosae/tobetsuensis*
	2	20	*Actinomyces graevenitzii, Alloprevotella rava, Atopobium parvulum, Lachnoanaerobaculum orale/saburreum, Prevotella histicola, Prevotella jejuni/melaninogenica, Prevotella salivae, Prevotella veroralis, Rothia mucilaginosa, Stomatobaculum longum*
Saliva	1	9	*Granulicetella adiacens/para-adiacens, Porphyromonas pasteri, Rothia aeria/dentocariosa*
	2	21	*Actinomyces graevenitzii, Atopobium parvulum, Campylobacter concisus, Lachnoanaerobaculum orale/saburreum, Prevotella histocola, Prevotella jejuni/melanogenica, Prevotella salivae, Rothia mucilaginosa, Stomatobaculum longum, Veillonella atypica/dispar*