A widespread family of ribosomal peptide metallophores involved in bacterial adaptation to copper stress

Laura Leprevost*[1], Sophie Jünger*[2], Guy Lippens[3], Céline Guillaume[2], Giuseppe Sicoli[4],

Lydie Oliveira[5], Alex Rivera-Millot[1&], Gabriel Billon[4], Céline Henry[5], Rudy Antoine[#1],

Svetlana Dubiley[#3], Séverine Zirah[#2], Yanyan Li[#2], Françoise Jacob-Dubuisson[#1]

[1] Univ. Lille, U1018-UMR9017-CIIL-Center for infection and Immunity of Lille, F-59000 Lille, FRANCE

[2] Unit Molecules of Communication and Adaptation of Microorganisms (MCAM), UMR 7245 CNRS, MNHN, Alliance Sorbonne University, 75005 Paris, FRANCE

[3] Toulouse Biotechnology Institute, CNRS/INRAE/INSA/UPS, 31077 Toulouse, FRANCE

[4] Univ. Lille, CNRS, UMR 8516 - LASIRE, Laboratoire de Spectroscopie pour les Interactions, la Réactivité et l'Environnement F-59000 Lille, FRANCE

[5] INRAE-AgroParisTech-Univ. Paris-Saclay, MICALIS, 78352 Jouy-en Josas, FRANCE

* contributed equally

for correspondence: rudy.antoine@pasteur-lille.fr; svetlana.dubiley@gmail.com; severine.zirah@mnhn.fr; yanyanli@mnhn.fr; francoise.jacob@ibl.cnrs.fr

& current address: INRS-Centre Armand-Frappier Santé Biotechnologie, Bacterial Symbionts Evolution, Laval, Quebec H7V 1B7, Canada

Abstract

Ribosomally synthesized, post-translationally modified peptides (RiPPs) are increasingly known to play prominent roles in bacterial adaptation. Herein, we identified and characterized a family of RiPP metallophores with rare 5-thiooxazole motifs, whose production is induced by transition metal stress in a number of pathogenic and environmental bacteria. Using *Caulobacter vibrioides* as a model, we demonstrated that the so-called bufferins confer protection to the bacteria from copper by complexing with the metal. Bufferin biosynthesis requires a multinuclear non-heme iron-dependent oxidase (MNIO, DUF692 family) and its partner protein to install thiooxazoles on Cys residues. By contrast, a hallmark feature of bufferin precursors, the N-terminal signal peptide sequence, is not essential for this transformation. Bufferin-like gene clusters are widespread in Eubacteria, including several pathogens, and the *C. vibrioides* epitomize the largest two families of RiPPs involving MNIO enzymes. Our study unveils a widespread but overlooked bacterial strategy mediated by a new type of RiPPs to cope with copper stress encountered in the environment and in the phagosomes of predatory protists or mammalian hosts.

Introduction

Microorganisms are endowed with an immense capacity of adaptation to their environments, both biotic and abiotic. One strategy is the production of specialised metabolites which play key roles in microbial adaptive processes, including defence against competing species, nutrient acquisition and resistance to stresses (1). Bacterial metallophores are prominent examples that can promote bacterial growth under different conditions (2-5). Several transition metals, including copper, are essential for living organisms (6, 7). However, the excess of copper is toxic to cells, because its high affinity for organic molecules results in the inactivation of essential pathways (7-9). Bacteria encounter copper in natural habitats and within phagocytic

cells, which use it as a microbicidal agent (10, 11). Therefore, they have developed an arsenal

of defence systems, including tight regulation of copper uptake, various export systems, and

sequestration/storage by copper-chelating molecules or proteins (12, 13). For pathogens, these

strategies are critical for host infection by enhancing their survival in phagocytes, and hence

they are linked to virulence (14).

Previous work on the whooping cough agent *Bordetella pertussis* has identified a four-gene

operon (BP2924-BP2927) controlled by BvgAS, the master two-component system that

regulates most virulence factors (15). This biosynthetic gene cluster (BGC) is similar to the *gig*

(gold-induced genes) operons of *Cupriavidus metallidurans* and *Legionella pneumophila* and

to a related operon of *Caulobacter vibrioides* (formerly *Caulobacter crescentus*) (16-18)

(Extended Fig. 1A). These operons except for that of *B. pertussis* are up-regulated upon

exposure to transition metals, notably copper (16-19). Although these data suggest a role for

gig-like genes in copper stress response, the function and the nature of derived biomolecules

have remained a mystery.

Ribosomally synthesized and post-translationally modified peptides (RiPPs) are natural

products with a tremendous variety of structures and biological activities (1, 20, 21). They are

synthesized from a gene-encoded precursor peptide, generally composed of an N-terminal

leader sequence and a core sequence where post-translational modifications (PTMs) occur. In

many cases, a RiPP recognition element (RRE) domain encoded in the same BGC interacts

with the leader sequence and serves as a docking site for the modification enzymes (22, 23).

RRE domains are thus a hallmark feature of RiPP biosynthesis. The vast structural diversity of

RiPPs is conferred by unlimited possibilities of chemical transformations. An emerging family

of PTM enzymes is the DUF692 family of multinuclear non-heme iron dependent oxidases

(MNIOs) that catalyse various reactions mostly but not exclusively on Cys residues, in the

biosynthesis of specific RiPPs (24-30). However, the full spectrum of reactions catalysed by

MNIO enzymes remains far from being fully appreciated.

Here we report the first functional and structural characterization of the molecules produced

by *gig*-like operons using *C. vibrioides* as a model. They represent a new family of RiPP

metallophores with 5-thiooxazole modifications installed by MNIOs and contribute to bacterial

adaptation to excess copper by complexing with this metal. We suggest to name these peptides

bufferins.

Results

Gig-like operons belong to putative RiPP BGCs. To gain insight into gene function, *in silico*

analyses were performed on the CCNA_3363-3366 operon of *C. vibrioides*, hereafter named

buf1. CCNA_3363 encodes a DUF2282-family protein (BufA1), whose members are 80-120

residues long and harbour four Cys residues and a predicted N-terminal, Sec-dependent signal

peptide (Fig. 1A; Extended Fig. 1A,B). The product of CCNA_3364 (BufB1) belongs to the

MNIO superfamily, whereas CCNA_3365 codes for a protein (BufC1) with an N-terminal

DUF2063 domain of unknown function. Importantly, the C-terminal domain of BufC1 is

predicted to be structurally related to RRE domains (Extended Fig. 1C). Finally, CCNA_3366

encodes a DoxX-type oxido-reductase (BufD1) (31, 32). The other *gig*-like operons share the

same gene organizations and predicted functions. In light of the recent discovery of MNIOs

involved in RiPP biosynthesis (24-30), we hypothesized that BufB1 and BufC1 would install

modifications on Cys residues of BufA1. Upstream of *bufA1*, CCNA_3362 and CCNA_3361

encoding the extracytoplasmic function (ECF)-family sigma factor SigF and the anti-sigma

protein NrsF were also found to be induced by copper stress (19) and are therefore part of the

buf1 BGC. Inspection of the *C. vibrioides* genome identified a second *buf2* BGC (CCNA_2999-

3001) which is also part of the SigF regulon (Fig. 1A). It encodes homologues of BufB1

(CCNA_3000) and BufC1 (CCNA_2999) but lacks a *doxX* gene. Its precursor peptide BufA2

(CCNA_3001) possesses a signal peptide, like BufA1, but its core sequence displays 6 Cys

residues and is without family signature (Extended Fig. 1C).

Bufferins are involved in adaptation to excess copper. In *C. vibrioides*, the *buf1* and *buf2*

BGCs belong to the SigF-controlled, core regulon of copper stress (17, 19, 33). The former was

also reported to respond to chromate and cadmium ions. These data suggest a link of the *buf*

operons with metal response. To test if they could be up-regulated by other transition metals,

we generated reporter fusions within *bufA1* and *bufA2* and confirmed that both are induced by

Cu^{2+} but not Fe^{2+}, Zn^{2+} or Mn^{2+} (Suppl. Fig. S1). As copper indirectly generates oxidative stress,

we also tested oxidants and identified HOCl as a weak inducer.

To determine if bufferins, the putative RiPP products of the *buf* BGCs, contribute to

adaptation to copper stress, we generated *C. vibrioides* deletion mutants that express one (Cv_{buf1}

and Cv_{buf2}) or neither BGC (Cv_{DKO}). Cv_{DKO} grew more slowly than the parental Cv_{WT} wild-

type strain when $CuSO_4$ was added to the growth medium, unlike Cv_{buf1} or Cv_{buf2} (Fig. 1B).

This indicates that bufferins enhance growth under copper stress and that they are functionally

redundant. To confirm that a single BCG is sufficient to protect bacteria from excess copper,

we performed a gain-of-function experiment in an *Escherichia coli* laboratory strain. For this,

we expressed the *buf1ABCD* operon under the control of a PT5-lac promoter upon IPTG

induction. Expression of *buf1ABCD* enhanced survival of *E. coli* exposed to a bactericidal

concentration of $CuSO_4$ (Fig. 1C).

We then asked whether the protection from copper would confer an advantage on *C.*

vibrioides in biotic interactions. As amoeba notably use copper against their bacterial preys, a

predation assay was performed (10). Attempts to measure survival of *C. vibrioides* after

ingestion by *Dictyostelium discoidum* were unsuccessful. Therefore, we tested if the presence

of the *buf* BGCs might slow down bacterial killing. We placed amoebal suspensions onto

bacterial lawns, and after incubation we determined bacterial densities in the contact zones by

densitometry scanning. Larger zones of complete bacterial lysis were observed for the Cv_{DKO}

than for the Cv_{WT} strain, consistent with a protective role of bufferins (Fig. 1D).

Features of buf BGCs required for function. To pinpoint key *buf* genes in the function of

bufferins, we constructed Cv_{buf1} derivatives that lack one or several genes ($Cv_{buf1(A)}$, $Cv_{buf1(ACD)}$,

$Cv_{buf1(ABD)}$, $Cv_{buf1(ABC)}$ and $Cv_{buf1(BCD)}$) and tested their growth phenotypes. Growth of all

deletion strains except for $Cv_{buf1(ABC)}$ was sluggish in the presence of $CuSO_4$, showing that

BufA1, BufB1 and BufC1 but not BufD1 are required for function in these growth conditions

(Fig. 2A). Similarly, the deletion of *bufB2* and *bufC2* in Cv_{buf2} impacted bacterial growth in the

presence of $CuSO_4$ (Fig. 2A).

As BufA1 harbours four conserved Cys residues (Cys^I to Cys^{IV}) that might be targets of

PTMs, we replaced them with Ser, separately or in combination and introduced the mutant

genes in the Cv_{buf1} chromosome by allelic exchange. All resulting strains $Cv_{buf1}^{CysISer+CysIVSer}$,

$Cv_{buf1}^{CysIISer+CysIIISer}$, $Cv_{buf1}^{CysIISer}$ and $Cv_{buf1}^{CysIIISer}$ grew more slowly in the presence of $CuSO_4$

than Cv_{buf1} (Fig. 2B), indicating the importance of the four Cys residues for function.

Characterization of PTMs on bufferins. To identify the PTMs on bufferins, we engineered an

IPTG-inducible version of SigF to control expression of their BGCs in *C. vibrioides*. The

metabolic profiles of IPTG-treated $Cv_{buf1}(pSigF)$ and $Cv_{buf2}(pSigF)$ were analysed by liquid

chromatography coupled to mass spectrometry (LC-MS) and compared to that of $Cv_{DKO}(pSigF)$

to reveal peptide species only present in the *buf*-expressing strains. In $Cv_{buf1}(pSigF)$, a major

species exhibited masses corresponding to BufA1 with the signal peptide cleaved off and a

mass shift of -10 Da (Fig. 3A; Suppl. Fig. S2; Suppl. Table S1). Interestingly, a second form

with a mass shift of -12 Da was also detected, but not systematically, and it disappeared upon

storage or freeze-thawing. The most stable product with a -10 Da mass shift is considered the

mature peptide product and named Bufferin 1 (Buf1). Similarly, Cv_{buf2}(pSigF) produced a

major peptide product with a mass shift of -18 Da, called bufferin 2 (Buf2), and another species

with a -20 Da mass shift (Extended Fig. 2).

Reduction and alkylation treatment yielded species carrying three or four alkylating groups

for Buf1 and four, five or six alkylating groups for Buf2 (Extended Fig. 3), suggesting that the

PTMs result in alkylation-sensitive groups. MS analyses of trypsin-digested, alkylated bufferins

revealed that in Buf1, Cys^{I} and Cys^{IV} had mass increments corresponding to alkylation,

indicative of unmodified residues (Extended Fig. 4). By contrast, in the central tryptic peptide

AGENDCKAGAGTSCAGTSK containing Cys^{II} and Cys^{III}, both Cys residues carried mass

increments corresponding to alkylation with additional -4.03 Da mass shifts (Fig. 3B), as

proposed by the untargeted PTM search solution SpecGlobX (34). Altogether, the data are

consistent with the presence of a disulfide (S-S) bond between Cys^{I} and Cys^{IV} (resulting in -2

Da) and the central Cys^{II} and Cys^{III} being targets of 4-electron oxidation reactions (-4 Da each)

that generate alkylation-sensitive groups. Consistently, substitution of Cys^{II} or Cys^{III} with Ser

in Buf1 yielded peptides with mass shifts of -6 Da (Extended Fig. 5). Similarly, in Buf2, the

central Cys residues carry -4.03 Da shifts, while Cys^{I} and Cys^{IV} would form a disulfide bond

(Suppl. Fig. S3). Based on these data, we interpret the -12 Da and -20 Da species of Buf1 and

Buf2 (which will be named $Buf1^{ox}$ and $Buf2^{ox}$), respectively, as forms harbouring labile S-S

bonds between two alkylation-sensitive groups, in equilibrium with the bufferins.

To facilitate bufferin purification and characterization we added C-terminal strep or twin-

strep tags to BufA1. The affinity-purified proteins showed -10 Da mass shifts relative to the

calculated masses (Suppl. Fig. S4). To probe if the modified Cys residues may be part of

heterocycles as with other MNIO enzyme-catalysed PTMs (24, 27), we recorded UV-visible

absorption spectra of purified Buf1 and the Buf1[CysIISer+CysIIISer] variant. The spectrum of the

former showed a broad absorbance band extending to 310 nm, unlike that of the latter (Fig. 3C).

Similarly, the spectrum of recombinant Buf2 showed an absorbance maximum at 305 nm.

(Suppl. Fig. S5). These results are consistent with the presence of aromatic heterocycles (35).

Identification of the PTMs of Bufferin 1 by NMR. To increase the yield of bufferin for

structural elucidation by NMR, we co-expressed *bufA1* carrying an N-terminal His$_6$-SUMO tag

with *bufB1* and *bufC1* in *E. coli* (Extended Fig. 6). Subsequent purification of modified BufA1

followed by trypsin digestion and preparative HPLC allowed to obtain the central 19-mer

peptide encompassing Cys[II] (Cys23) and Cys[III] (Cys31) at a milligram scale. Its ^{1}H spectrum

indicated two groups of anomalous resonances (Fig. 3D). By combining homonuclear ROESY

and TOCSY experiments (36) they were assigned to the NH protons of Lys24 and Ala32 that

follow the modified Cys23 and Cys31 residues, respectively, and to the Hα protons of Asp22

and Ser30 that precede these modified Cys residues (Extended Fig. 7A). The absence of both

Hα and NH protons for both Cys residues was verified on an ^{1}H, ^{15}N HSQC spectrum (Extended

Fig. 7B), indicating that their N and Cα atoms are part of heterocycles.

An ^{1}H, ^{13}C HMBC experiment showed correlations from the Asp22 and Ser30 Hα atoms

to carbon atoms at 161.1 ppm and 159.8 ppm, respectively (Extended Fig. 7C). Further

assignments were obtained with the uniformly ^{13}C,^{15}N-labeled 19-mer peptide and triple-

resonance NMR experiments (37, 38). In an HNCACB experiment we found no resonances

that could correspond to Lys24's upstream neighbour, confirming that the Cα of modified Cys[II]

does not resonate at a typical 50-ppm value (Extended Fig. 8A). With an HNCO experiment,

we found values around 184 ppm for the C=O carbon atoms of both Cys residues (Extended

Fig. 8B), far from 175-ppm and 200-ppm values expected for such atoms in random-coil or

thio-amide peptides, respectively (39, 40). By increasing the resolution and varying the offset

of the decoupling pulse during the C=O evolution period, we determined a 90-Hz coupling constant of these atoms with carbon atoms whose chemical shift is ~120 ppm, consistent with their absence in the HNCACB experiment (Extended Fig. 8C). Both parameters point to the C=O carbon atoms of the modified Cys connected to carbon atoms in aromatic rings (41).

The ^{13}C chemical shifts assigned on the modified peptide are in the same range as those for the ring carbon atoms of a 5-thiooxazole-containing natural product (35). Together with the anomalous chemical shift values for the Hα and NH protons of the flanking residues and taking into account the MS data, we conclude that the modification leads to cyclization of both Cys to form thiooxazole groups (Fig. 3E; Suppl. Text), consistent with their characteristic absorbance maximum.

Copper binding by Bufferin 1. Given that heterocycles are prone to copper binding (5), we interrogated if bufferins can complex with copper, which could be the basis for their protective effect. To test this, purified Buf1^{twstr} was incubated with CuCl$_2$ and analyzed by native MS. A 1:1 Buf1-Cu complex was detected (Suppl. Fig. S6), but the isotopic pattern suggested the occurrence of both Buf1 and Buf1^{ox}, and the complexed/apo forms ratio varied between experiments. We performed a spectroscopic binding assay with Buf1 to determine its affinity constant for copper (42). However, formation of the complex appeared to be very slow, suggesting an equilibrium between binding-competent and -incompetent bufferin forms (Suppl. Fig. S6).

Therefore, we investigated complex formation *in vivo* by treating cultures of *bufA1^{str}*-expressing *E. coli* with CuSO$_4$ for several hours prior to bufferin purification. Native MS analysis showed that Buf1^{str} thus produced was fully loaded with copper (Fig. 4A). This species was further investigated by electron paramagnetic resonance (EPR). Pulsed EPR spectra

confirmed that bufferin 1 binds paramagnetic Cu^{2+}, and spectral simulations indicated

coordinating N atoms (43) (Fig. 4B; Suppl. Tables S2 and S3; Suppl. Fig. S7).

To gain insight into the copper binding site, we produced ^{15}N- and $^{15}N,^{13}C$-labelled bufferin

samples and used 3-D spectra to assign the resonances. Simultaneously, we used the ^{1}H, ^{15}N

HSQC plane as a basis for HSQC-NOESY and TOCSY experiments to confirm the assignments

and gain some knowledge on structural aspects of bufferin. The sequential assignment was

interrupted by the modified Cys residues, as expected. Interestingly, amide resonances of the

tryptic peptide and of the same segment of the full-length bufferin do not correspond at all,

suggesting that the full-length sequence imposes structural constraints on the thiooxazole-

containing loop. Part of these likely come from the β sheet spanning two short strands in the

structural model, Thr9-Tyr13 and Trp44-Lys48 (Extended Fig. 9). The strong Hα(i-1) – HN(i)

NOE contacts between residues in these segments confirm their extended structure, and the

cross-strand NOE contacts between the amide protons of Lys45 and Cys12, and of Val47 and

Glu10, respectively, support their anti-parallel arrangement (Extended Fig. 9).

A comparison of the spectra of the bufferin produced with or without copper showed that

many peaks remained unchanged, others experienced a shift and some signal reduction, and yet

others disappeared altogether (Fig. 4C). The latter most probably concern residues close to the

paramagnetic Cu^{2+} ion. In particular, disappearance of the peaks corresponding to direct

neighbours of the thiooxazole heterocycles, Lys24, Ser30 and Ala32, strongly indicates that

these heterocycles are involved in Cu^{2+} coordination. Interestingly, signals of residues of the β-

sheet face oriented towards the thiooxazole-containing loop, Thr9, Trp44, Met46 and Lys48,

also disappeared in the spectrum of the complex. Whereas a full structural determination of

bufferin will require further experiments, our data support that the thiooxazole cycles and

residues of the β sheet are involved in copper binding.

In vitro reconstitution of MNIO enzymatic activity. The modification of Cys to 5-thiooxazole is a new type of reaction catalysed by MNIO enzymes. To investigate the function of BufB1, we purified recombinant SUMO-BufA1, BufB1 and BufC1 (Suppl. Fig. S8). Since MNIO enzymes require Fe ions for activity (26, 29), BufB1 was analysed by inductively coupled plasma - optical emission spectrometry, showing the presence of 2 Fe ions. Structural alignments of a BufB1 model with the crystal structure of MbnB of *Vibrio caribbeanicus*, an MNIO enzyme involved in methanobactin biosynthesis (26) confirmed that the Fe-coordinating residues are conserved in BufB1. Next, BufA1 with the SUMO-tag cleaved was incubated with BufB1 and/or BufC1 (Fig. 5A). LC-MS analyses showed that BufA1 was converted to a species with a −8 Da mass shift only when both BufB1 and BufC1 were present (Fig. 5B; Suppl. Fig. S9). Trypsin digestion of the *in vitro* modified BufA1 yielded the central 19-residue fragment harbouring the thiooxazole groups derived from Cys^{II} and Cys^{III} as confirmed by MS/MS analyses (Fig. 6B). Together with the gene inactivation data, these experiments established that BufB1 and BufC1 are required and sufficient to install the PTMs on BufA1.

Signal peptide sequence is not essential for heterocycle formation. The particularity of bufferin biosynthesis is that BufA precursors harbour N-terminal Sec-dependent signal peptide sequences. We investigated if this sequence is necessary for Cys transformation using the co-expression system of His-SUMO-*bufA1* with *BufB1* and *BufC1* in *E. coli* (Extended Fig. 6). The *bufA1* gene was mutated to generate variant precursors with the signal peptide truncated 8 residues from the N terminus (BufA1(-8aa)) or completely removed (BufA1(CP)). Subsequently, LCMS analysis of the purified BufA1 after SUMO cleavage indicated that both truncated precursors were modified with a 10 Da loss, consistent with the presence of two thiooxazoles and one S-S bond (Extended Fig. 10). Of note, BufA1(CP) was converted by BufB1/C1 in a less efficient manner, with the unmodified peptide also observed.

Large families of bufferin-like RiPPs in bacteria. To determine how widespread bufferins are among prokaryotes, we collected >14,000 MNIO enzyme-coding genes and found them to be widely distributed among Proteobacteria, Terrabacteria, Myxococcota, Fibrobacteres/Chlorobi/Bacterioidetes, Planctomycetes/Verrucomicrobia/Chlamydiota, and Acidobacteriota groups (detailed *in silico* analyses will be published separately). We analysed their genomic contexts and collected associated putative bufferin-like precursor genes to perform sequence similarity network (SSN) analyses of the corresponding proteins. The largest cluster corresponds to the DUF2282 family that includes *C. vibrioides* BufA1 (Fig. 6A). Most of the other proteins have no family signature, but they contain conserved Cys residues, consistent with their being potential RiPP precursors. BufA2 of *C. vibrioides* belongs to the second largest family of proteins characterized by 6 or 12 conserved Cys residues, which we named BUF_6/12Cys (Fig. 6A,B). Parallel SSN analyses of the MNIO enzymes superfamily (Fig. 6C) showed that those associated with DF2282 or BUF_6/12Cys bufferins belong to the largest sequence cluster, distinct from smaller clusters of enzymes involved in the biosynthesis of methanobactins, RiPPs from *Chryseobacterium*, TglA-type pearlins and amino-pyruvatides (24, 26-28, 30). Therefore, we have uncovered two large families of RiPPs whose biosynthesis involves MNIO enzymes, exemplified by the *C. vibrioides* bufferins, that are distinct from previously described MNIO enzyme-modified RiPPs.

Discussion

In this work, we unveiled the chemical nature and function of the products derived from heavy metal-induced *gig*-like operons found in various environmental and pathogenic bacteria (16-18). To the best of our knowledge, bufferins are the first genuine RiPP natural products harbouring 5-thiooxazole motifs. Using model bufferin systems in *C. vibrioides*, we

demonstrated that the 5-thiooxazoles are key for copper binding. Peptide natural products with

other heterocycles are known to bind transition metals and fulfil different functions (2-5). Most

prominently, methanobactins scavenge copper from the milieu to feed it to cuproenzymes, and

some of them also protect bacteria from toxic metals such as mercury (44). The defensive role

of bufferins is reminiscent of those of the non-ribosomal siderophore peptides anthrochelin and

yersiniabactin, which can passivate copper by chelation (2, 45). However, the fate of copper

ions bound by bufferins remains unknown. Periplasmic bufferins might provide specific

metallochaperones with copper for delivery to cuproproteins or alternatively facilitate its

export.

We thus discovered yet another bacterial strategy of adaptation to copper that makes use of

such a new type of RiPPs widespread in the phylogenetic tree. Bufferins are part of complex

and interwoven metal defence systems in bacteria, as suggested by the fact that they do not

provide a high level of protection by themselves, as shown in *C. metallidurans* (17, 46). They

are probably best seen as additional players that buttress active defence systems through copper

passivation under specific environmental conditions. Nonetheless, their presence in pathogenic

bacteria indicates that they are likely beneficial to the latter in host-pathogen interactions, likely

by helping bacteria deal with copper encountered in eukaryotic phagocytes. Whether all

putative bufferins found by genome mining function in metal adaptation similar to our model

bufferins remain to be addressed. If so, the nature of the metals, the stoichiometry and the

affinity of the bufferins will likely depend on their primary and tertiary structures.

The biosynthesis of bufferins involves several novel features. Transformation of Cys to 5-

thiooxazole is catalysed by the MNIO BufB and its partner protein BufC. A similar reaction

was previously reported to engage radical-SAM enzymes (35). Intense efforts have been made

recently to explore the chemical space of RiPPs involving MNIOs (24, 25, 28, 30, 47). To date,

members of this superfamily of enzymes are known to catalyse remarkable oxidation reactions,

including oxazolone coupled to thioamide formation, excision of cysteine β-methylene, Cys-involved macrocyclization, and C-C and C-N cleavages, as evidenced in the biosynthesis of methanobactins, TglA-type pearlins, RiPPs in *Chryseobacterium*, aminopyruvatides, and methanobactin-like peptides in *Vibrio*, respectively (24, 25, 27, 28, 30). This work thus further expands the reaction spectrum of MNIOs. In line with previous work on Cys-modifying MNIOs, BufB would similarly make use of the mixed-valent iron centres and proceed by an intermediate step of proton abstraction from the Cβ of cysteine (26, 27, 29, 30) (Fig. 7). Furthermore, bufferin precursor peptides commonly harbour signal peptides instead of leader peptides at the N-terminus. Herein we showed that the signal peptide is dispensable for heterocycle installation, which begs the question of the specific recognition between the precursor and the BufB/C complex but highlights the biotechnological potential of BufB/C.

Our bioinformatic analyses indicated that bufferins currently represent the largest two families of MNIO-associated RiPPs, widespread in different bacterial phyla. Putative bufferin systems feature original properties, including the N-terminal signal peptides of their precursors and the large sizes of their mature forms relative to most other RiPP precursor families. The use of signal-peptides, which determine Sec-dependent export and signal-peptidase-mediated processing of bufferins dispenses with the need for specific transporter and proteases genes in their BGCs, possibly explaining why bufferins had previously been overlooked by genome mining searches. The diversity in length and sequence of their cores provides almost infinite possibilities to build new peptide metallophores.

Methods

C. vibrioides constructs and growth conditions. *C. vibrioides* NA1000 was used. Allelic exchange was performed using pNPTS138 to construct deletion strains and to introduce modified versions of *bufA1* and *bufA2*. pSRK-Km was used for expression of SigF. Plasmids

were mobilized from *E. coli* S17 into *C. vibrioides* by conjugation. The presence of the expected deletions was verified by PCR. Chromosomal transcriptional fusions were generated using pFUS2, and β-gal activities were measured as in (48). The plasmids, recombinant strains, primers and synthetic genes used in this study are shown in Tables S4 to S6.

C. vibrioides was grown at 30°C in PYE medium with nalidixic acid (30 and 15 µg/ml in solid and liquid medium, respectively), and kanamycin (20 and 5 µg/ml) or gentamycin (5 and 0.5 µg/ml) when carrying specific plasmids. To record growth curves, cultures in the stationary phase were diluted to $OD_{600} = 0.05$ and inoculated in 96-well plates with continuous shaking at 30°C in a TECAN Spark plate reader, with OD_{600} measurements every 20 min for 28 h. $CuSO_4$ was added to 225 µM where indicated. The reporter strains were grown to early stationary phase in the presence of metal ions or ROS at the indicated concentrations. Xanthine oxidase and hypoxanthine were used to generate O_2^- and H_2O_2 (49). The production of ROS was quantified using bacterial cultures grown for 24 h starting from $OD = 0.1$ at 30°C under moderate shaking. The bacteria were then treated with 5 µM of CellROX Deep Red (Thermo fisher) for 30 min at 30°C, washed three times in 20 mM sodium phosphate (pH 7.5), and transferred into black-bottom 96-wells plates. The emitted fluorescence was measured at 690 nm after excitation at 640 nm with a TECAN Spark plate reader. Biological quadruplicates were obtained in all assays.

Genetic constructs in E. coli. pCA24 (50) was used for expression of the operon in *E. coli*. The entire *buf1* operon was PCR-amplified from genomic DNA and inserted after the T5-lac promoter of pCA24 using Gibson assembly (51). The sequence coding for BufA1 with a C-terminal Strep-tag II was PCR-amplified and used to replace the native *bufA1* gene, yielding pCA24-A^{str}BCD. For large-scale production of recombinant BufA1, the PCR-amplified gene was inserted into SspI-linearized pETHisSUMO by ligase-independent cloning, generating pETHisSUMO-A1. pACYC-Duet-based expression plasmids of N-terminally His$_6$-tagged

BufB1 and BufC1 were created using *in vivo* assembly (IVA) (52) yielding pACYC::HisC1 *and* pACYC::HisB1. For co-expression, *bufB1* was cloned into the MCS2 of pACYC::HisC1 by IVA cloning, yielding pACYC::HisC1B1Stag.

Gain-of function in E. coli. BL21(pCA24-*bufA1^{str}1BCD*) and BL21(pCA24-*psmCA*) (negative control) (53) were grown overnight at 37°C, diluted 100-fold in M9 medium supplemented with 1xBME vitamin solution (Sigma), 0.5% glycerol, 34 µg/mL chloramphenicol and 1 mM IPTG and grown at 37°C with vigorous shaking for 4 h. At $OD_{600} = 0.1$, one half of each culture was exposed to 0.3 mM $CuSO_4$. Incubation was pursued for 3 h before CFUs were counted.

Amoeba experiments. Exponentially growing *C. vibrioides* cultures were concentrated to $OD_{600} = 1$ in PYE medium and plated on PYE-agar with 1% glucose. 50-µL drops of *Dictyostelium discoidum* grown to 80% confluency in HL/5 medium at 20°C and diluted to 10^6/mL were placed on the plate centres, before incubation at 20°C for three days. ImageJ was used to measure the densities of bacteria in the zones of contact between bacteria and amoeba, divided by the area of these zones.

Native bufferin production and purification. Cultures of Cv$_{buf1}$^{TwStrep}(pSigF), Cvbuf1^{6His}(pSigF) and Cvbuf2^{6His}(pSigF) were started at $OD_{600} = 0.1$ in PYE supplemented with 100 µM IPTG. Bacteria were harvested after overnight growth, resuspended in 100 mM Tris-HCl (pH = 8), 150 mM NaCl and broken using a French press. Buf1^{TwStrep} was purified from the clarified lysate using a 1-mL StrepTrap XT column (Cytiva) following the manufacturer's instructions, concentrated using an Amicon ultrafiltration device and dialysed against 25 mM Hepes (pH 8) 150 mM NaCl. Buf1^{twstr} was truncated *in vivo* just after the first strep tag motif, as evidenced by native MS. The Buf1^{6His} and Buf2^{6His} proteins were purified on 1-mL Ni^{2+} columns using standard procedures.

For purification of isotopically labeled BufA1, *E. coli* BL21(pCA-A^{str}BCD) was grown in in M9 medium supplemented with 34 µg/mL chloramphenicol, 1xBME vitamin solution

(Sigma), 0.5% glycerol-$^{13}C_3$ and 0.1 g/L $^{15}NH_4Cl$ at 37 °C until OD600 ~ 0.6. Protein expression was induced with 1 mM IPTG, and the culture continued at 22°C overnight. $CuSO_4$ was added to 0.3 mM following overnight growth, and after 4 more hours bacteria were harvested by centrifugation. They were resuspended in 50 mM Tris-HCl buffer (pH 8.0), 300 mM NaCl and lysed by sonication. The clarified lysate was applied to a 1-mL Strep-Tactin® Superflow column (IBA Lifesciences, Germany), and after washing with 20 column volumes of the same buffer, BufA^{str} was eluted with 2.5 mM desthiobiotin, 50 mM Na-phosphate buffer (pH 7.0), 50 mM NaCl. The molar extinction coefficient used for Buf1^{str} was 38,158 $M^{-1}cm^{-1}$.

Protein production and purification for in vitro assays. N-His$_6$-BufB1 and N-His$_6$-BufC1 were each produced in *E. coli* BL21(DE3), in 1 L autoinduction ZYM-5052 medium (54) supplemented with 250 µM ferrous ammonium sulphate for BufB1. After 4 h of growth at 37°C, the temperature was lowered to 18°C and cultures were pursued for 16 h. N-His$_6$-tagged SUMO-BufA1 was produced in 1 L TB medium, with bacteria grown until OD_{600} = 0.6-0.7 before cooling down on ice. After adding IPTG to 0.1 mM, the cultures were pursued at 18°C for 16 h. Bacteria were resuspended in 50 mM Tris-HCl (pH 7.6), 300 mM NaCl, 10% glycerol, 10 mM imidazole, and lysed using French Press or by sonication in the case of BufA1. Recombinant proteins were purified by gravity using a 5-mL-column packed with Ni-NTA resin (Qiagen). After washes with 30 mM and 60 mM imidazole in 50 mM Tris-HCl (pH 7.6), 300 mM NaCl, the proteins were eluted with 50 mM Tris-HCl (pH 7.6), 300 mM NaCl, 250 mM imidazole, concentrated by ultrafiltration using Amicon centrifugal units, and buffer exchanged by PD10 columns (Cytiva) to 50 mM Tris-HCl (pH 7.6), 150 mM NaCl, 20% glycerol. Purified proteins were flash frozen and stored at -80°C. Purification followed the same procedure for all three proteins except that 10 mM of Tris(2-carboxyethyl)phosphine hydrochloride was added to all buffers for BufA1. The iron content of His$_6$-BufB1 was measured by using an inductively coupled plasma - optical emission spectrometer (ICP-OES

5110 VDV, Agilent Technologies) at $\lambda = 238.204$ nm calibrated using standard solutions. All

the solutions were acidified with HNO_3 (Fisher Scientific, 67–69%, Optima and trace metal

grade). Certified water Enviromat (SCP, N°140-025-038) was analysed for quality control

(certified value of 0.581 +/- 0.011 mg/L, experimental result of 0.59 +/- 0.02 mg/L). The limit

of detection was estimated to 0.005 mg/L and the average blank value was below this limit.

Large-scale production of SUMO-BufA1 for NMR studies. *E. coli* BL21(DE3,

pACYC::HisC1B1Stag, pETHisSUMO-A1) was grown in 2 L TB medium. For ^{13}C and ^{15}N-

labelling, protein expression was carried out in 1 L M9 medium supplemented with 1 g/L of

^{13}C-glucose and 5 g/L $^{15}NH_4Cl$. In both cases, cultures were grown until $OD_{600} = 0.6$-0.7 before

cooling down on ice. After adding ITPG to 0.1 mM, the cultures were pursued at 18°C for 16

h (TB medium) or 48 h (M9 medium), respectively. Purification of modified N-His$_6$-tagged

SUMO-BufA1 was performed as described above.

Peptide analyses by MS. Cell pellets and culture supernatants from 25-mL overnight *C.*

vibrioides cultures were resuspended in 1 mL 50 mM Tris HCl (pH 7.5) and submitted to fast-

prep lysis in Lysing Matrix B tubes (force 5, 30 sec, 2 times). The clarified lysates were

submitted to solid-phase extraction (SPE) on Sep-Pak tC2 Plus Light Cartridges (Waters)

conditioned with 5 mL methanol and 5 mL 0.01 % formic acid (FA). The cartridges were

washed with 5 mL 0.01 % FA and eluted with 2 mL methanol. The eluted fractions were split

in two, evaporated under vacuum and dissolved in 50 μL 0.01% FA/methanol 1:1 (v/v). One

was left intact (I), and the other was treated with 10 mM dithiothreitol (DTT) for 1 h at 50°C

and alkylated with 15 mM iodoacetamide for 1 h at 20°C in the dark. Half of that solution was

collected (R1) and the other half was treated with 0.01 μg/μL trypsin (Gold, Promega) overnight

at 37°C (R2). Alternatively, reduction-alkylation was performed using DTT together with 2-

bromoethylamine (55). R1 and R2 were desalted using Pierce C18 Spin columns. The purified

Buf1^{twstr} was analysed by LC-MS directly or after reduction-alkylation and digestion with

trypsin as above. I, R1 and R2 were analysed by ultra-high-performance liquid chromatography-MS on an Ultimate 3000-RSLC system (Thermo Scientific) connected to an electrospray ionization- quadrupole–time of flight instrument (Maxis II ETD, Bruker Daltonics). The separation was achieved using an Aeris widepore C4 column (3.6 μm, 200 Å, 2.1 × 100 mm, Phenomenex) for I and R1 samples and a RSLC Polar Advantage II Acclaim column (2.2 μm, 120 Å, 2.1 × 100 mm, Thermo Scientific) for R2, with a linear gradient (2-80 % in 15 min) of mobile phases A (FA 0.1%) and B (LC-MS grade acetonitrile (ACN) + FA 0.08%), at 300 μL/min. For I and R1, LC-MS detection was carried out in positive mode in the range *m/z* 250-2500. For R2 samples, data-dependent LCMS/MS acquisitions were conducted. They were also analyzed by nanoLC-MS on an UltiMate 3000 RSLCnano System coupled to an Orbitrap Fusion Lumos Tribrid (Thermo Fisher Scientific). The separation was achieved using a C18 Biosphere column (2 μm, 75 μm × 500 mm Neo Thermoscientific) with a linear gradient (2.5-45% B for 55 min, 45-98% for 10 min and 2.5% for 10 min) of phases A (FA 0.1%/ACN 98/2) and B (FA 0.1%/ACN 20/80) at 0.250 μL/min. The mass spectrometer was operated in positive mode in the range *m/z* 150-1200. Data-dependent LCMS/MS acquisition mode was used. Full MS scans were captured with a resolution of 120,000. Peptide identification was performed using X!TandemPipeline (56) or PEAKS Studio X Pro (Bioinfomatics Solutions Inc.), against the *Caulobacter* UniProtKB database (version 2023) with the following parameters set as variable modifications: one possible missed cleavage, Cys carboxyamidomethylation, and Met oxidation. Precursor mass and fragment mass tolerance were 10 ppm and 0.02 Da, respectively. Data filtering was achieved according to a peptide E value < 0.05. The false discovery rate at the peptide level was assessed from searches against a decoy database. MS2 spectra were analysed using BYONIC™ (Protein Metrics) together with i2MassChroQ 0.6.7 (http://pappso.inrae.fr/bioinfo/i2masschroq) and SpecGlobTool to highlight and localize unknown PTMs (34).

Native MS was carried out using flow injection with 50 μL/min 10 mM ammonium

acetate (pH 7). One μL of purified peptide at 20 μM in 10 mM ammonium acetate (pH 7) was

injected. For *in vitro* binding experiments, the peptide was previously incubated for 1 h with

CuCl$_2$ (1 or 2 equivalents) at room temperature.

NMR analyses. Purified His-SUMO-BufA1* was incubated with trypsin (1:100 w/w) at 37°C

overnight in 50 mM Tris-HCl (pH 7.6), 10 mM TCEP. Purification of the 19-residue central

trypsin fragment containing two modified Cys residues was conducted by HPLC (UltiMate^{TM}

3000 System, Thermo Scientific) on a preparative C18 column (Luna C18 (2), 5 μm, 100 Å,

250 x 10 mm, Phenomenex) at ambient temperature. Separation was performed using a linear

gradient (6 min at 5% B, 6-14 min from 5-30 % B, 14-18 min to 80% B, and 18-22 min at

80%B) of mobile phases A (0.1% FA) and B (ACN) at 3 mL/min, and monitored at 280 and

305 nm. Fractions containing the relevant peptide were pooled and dried by SpeedVac. A

sample of 3.4 mg of the tryptic peptide dissolved in 550 μL of 50 mM Na phosphate buffer (pH

6.5), 150 mM NaCl, with 25 μL D$_2$O and Trimethyl Silyl Propionate (TSP) as reference was

introduced in a standard 5-mm NMR tube. All spectra were recorded at 293 K on an 800-MHz

NEO Bruker spectrometer equipped with a QCP cryogenic probe head. TOCSY and ROESY

spectra were recorded for assignment of the peptide with 16 k x 512 points for a spectral width

of 12 ppm in both dimensions, with 16 and 64 scans per increment, respectively (total

measurement times of 4 h and 16 h, respectively). Scalar coupling transfer was accomplished

during a 60-ms DIPSI2 mixing time, whereas a 100-ms spin lock was used for the dipolar

magnetization transfer. The relaxation delay was set to 1 s for both spectra. The natural

abundance [1]H, [15]N HSQC spectrum was recorded with the Bruker hsqcf3gpph19 pulse

program, with 2 k x 128 points for a spectral width of 12 ppm x 36 ppm, centered on 4.7 ppm

and 119 ppm for the H and the N dimensions, respectively, and 256 scans per increment. Natural

abundance [1]H, [13]C HSQC and HMBC spectra were recorded with standard Bruker pulse

programs (hsqcetgpsisp.2 and hmbcedetgpl3nd), with 2 k x 256 points for a spectral window of 12 x 40 ppm and the ^{13}C carrier set at 80 ppm, and with 2 k x 256 points for a spectral window of 12 x 140 ppm and the ^{13}C window centered at 100 ppm, respectively, and 64 scans per increment for both. A HNCACB spectrum was recorded on the ^{15}N, ^{13}C labeled peptide as a cube of 12 x 30 x 70 ppm sampled with 2 k x 76 x 256 points centered at 4.7 x 119 x 70 ppm in the ^{1}H, ^{15}N and ^{13}C dimensions, respectively. For the HNCO spectrum, the ^{13}C carrier was shifted to 175 ppm, and the ^{13}CO window of 30 ppm was sampled with 64 points. In order to estimate the carbonyl frequencies of the modified Cys residues, we recorded the H(N)CO planes of the latter spectrum while introducing a selective pulse with varying offset during the ^{13}CO evolution period.

EPR experiments. Pulsed (EDFS) EPR experiments on the *in vivo*-formed Buf1^{twstr}-copper complex (200 μM) were carried out using an ELEXSYS E-580 spectrometer (Bruker) equipped with a Super-Q FTu bridge for X- (9 GHz) and Q- (35 GHz) band experiments. Samples were drawn into 3 × 4 mm quartz tubes, flash-frozen into liquid nitrogen and rapidly transferred into precooled dielectric resonators, ER 4118X-MD5W X-band and ER 5106QT-2w Q-band. Both X- and Q-band spectra were acquired at 20 K using a Bruker Cryogen-free (Cold-Edge) cooling system. Microwave pulse lengths $t_{\pi/2} = 10\text{-}12$ ns (Q-band) and $t_{\pi/2} = 16$ ns (X-band) were used with inter-pulse delays of $\tau = 172$ ns (Q-band) and $\tau = 132$ ns (X-band). A two-step phase-cycle was applied to remove all unwanted echoes on EDFS. Numerical simulation of the EPR spectra was conducted using the Matlab toolbox Easyspin 5.2.35.

Spectrophotometric copper binding assay. Copper binding to purified Buf1^{twstr} was assayed using competition experiments using a complex between 4-(2-pyridylazo)resorcinol (PAR) and Cu^{2+} ($\varepsilon_{505nm} = 41{,}500$ M^{-1} cm^{-1}) (42), with 4 μM Cu, 10 μM PAR and 5 to 40 μM Buf1^{twstr}. The visible spectra were recorded after increasing incubation periods at room temperature.

Enzymatic assays in vitro. Standard reactions were carried out in 100-µl solutions of 50 µM purified His-SUMO-BufA1 in 50 mM Tris-HCl (pH 7.6) and 10 mM TCEP. For removal of the affinity tag, this mixture was first incubated with in-house purified SUMO protease at a 1:20 ratio of protease/peptide overnight at 4°C. Subsequently, either or both purified N-His$_6$-BufB1 (25 µM) and N-His$_6$-BufC1 (25 µM) were added, and the mixtures were incubated at 30°C for 16 h. For control reactions BufB1 and BufC1 were inactivated by heating. Half of the reaction mixture was desalted using Zeba spin columns (Thermo Fisher Scientific) and directly analysed by LC-MSMS on a Aeris widepore C4 column. The other half was treated with trypsin (Promega) added to 1:100 (w/w) and incubated for 3 h at 37°C before LC-MSMS analysis on a RSLC Polar Advantage II Acclaim column. LC-MSMS measurements were performed on an Ultimate 3000-RSLC system (Thermo Scientific) connected to a high-resolution ESI-Q-TOF mass spectrometer (Maxis II ETD, Bruker Daltonics), as described above.

In silico analyses. The non-redundant NCBI database (release of Jan 2023) was searched for MNIO-coding genes. To identify new families of RiPPs, a script was designed to collect genes next to and in the same orientation as MNIO genes, coding for proteins less than 200 residues long. Predictions of signal peptides were performed using various softwares (Scotopus, Philius, Octopius, Scampi, Polyphobius, Topcons). Analyses by the Enzyme Function Initiative - Enzyme Similarity Tool (EFI-EST) were carried out with the predicted proteins (after removal of their signal peptides) to build sequence similarity networks (SSN). For the generation of a hmm profile for the BUF_6/12Cys family, the procedure described in (57) was followed. Briefly, a few dozen proteins were picked from the cluster of interest of the Representative Node Network, and their sequences were aligned with MAFFT (L-INS-i). After manual editing of the non-conserved mostly N- and C-terminal regions, a hmm profile was generated using this seed alignment.

References

1. Li Y, Rebuffat S. The manifold roles of microbial ribosomal peptide-based natural products in physiology and ecology. J Biol Chem. 2020;295(1):34-54.

2. Büttner H, Hörl J, Krabbe J, Hertweck C. Discovery and biosynthesis of anthrochelin, a growth-promoting metallophore of the human pathogen *Luteibacter anthropi*. ChemBioChem. 2023;24:e202300322.

3. Chaturvedi KS, Hung CS, Crowley JR, Stapleton AE, Henderson JP. The siderophore yersiniabactin binds copper to protect pathogens during infection. Nat Chem Biol. 2012;8(8):731-6.

4. Ghssein G, Brutesco C, Ouerdane L, Fojcik C, Izaute A, Wang S, et al. Biosynthesis of a broad-spectrum nicotianamine-like metallophore in Staphylococcus aureus. Science. 2016;352(6289):1105-9.

5. Kenney GE, Rosenzweig AC. Chalkophores. Annu Rev Biochem. 2018;87:645-76.

6. Andrei A, Ozturk Y, Khalfaoui-Hassani B, Rauch J, Marckmann D, Trasnea PI, et al. Cu Homeostasis in Bacteria: The Ins and Outs. Membranes (Basel). 2020;10(9).

7. Solioz M. Copper and Bacteria. Evolution, homeostasis and toxicity. Springer, editor. Cham, Switzerland: Springer Nature Switzerland AG; 2018.

8. Macomber L, Imlay JA. The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. Proc Natl Acad Sci U S A. 2009;106(20):8344-9.

9. Zuily L, Lahrach N, Fassler R, Genest O, Faller P, Seneque O, et al. Copper Induces Protein Aggregation, a Toxic Process Compensated by Molecular Chaperones. mBio. 2022:e0325121.

10. Hao X, Luthje F, Ronn R, German NA, Li X, Huang F, et al. A role for copper in protozoan grazing - two billion years selecting for bacterial copper resistance. Mol Microbiol. 2016;102(4):628-41.

11. Sheldon JR, Skaar EP. Metals as phagocyte antimicrobial effectors. Curr Opin Immunol. 2019;60:1-9.

12. Arguello JM, Raimunda D, Padilla-Benavides T. Mechanisms of copper homeostasis in bacteria. Front Cell Infect Microbiol. 2013;3:73.

13. Chandrangsu P, Rensing C, Helmann JD. Metal homeostasis and resistance in bacteria. Nat Rev Microbiol. 2017;15(6):338-50.

14. Hodgkinson V, Petris MJ. Copper homeostasis at the host-pathogen interface. J Biol Chem. 2012;287(17):13549-55.

15. Coutte L, Antoine R, Slupek S, Solans L, Derop J, Bonnefond A, et al. Combined RNAseq and ChIPseq Analyses of the BvgA Virulence Regulator of Bordetella pertussis. mSystems. 2020;5(3).

16. Jwanoswki K, Wells C, Bruce T, Rutt J, Banks T, McNealy TL. The Legionella pneumophila GIG operon responds to gold and copper in planktonic and biofilm cultures. PLoS One. 2017;12(5):e0174245.

17. Kohler C, Lourenco RF, Avelar GM, Gomes SL. Extracytoplasmic function (ECF) sigma factor sigmaF is involved in Caulobacter crescentus response to heavy metal stress. BMC Microbiol. 2012;12:210.

18. Wiesemann N, Mohr J, Grosse C, Herzberg M, Hause G, Reith F, et al. Influence of copper resistance determinants on gold transformation by Cupriavidus metallidurans strain CH34. J Bacteriol. 2013;195(10):2298-308.

19. Maertens L, Cherry P, Tilquin F, Van Houdt R, Matroule JY. Environmental Conditions Modulate the Transcriptomic Response of Both Caulobacter crescentus Morphotypes to Cu Stress. Microorganisms. 2021;9(6).

20. Arnison PG, Bibb MJ, Bierbaum G, Bowers AA, Bugni TS, Bulaj G, et al. Ribosomally synthesized
and post-translationally modified peptide natural products: overview and recommendations for a
universal nomenclature. Nat Prod Rep. 2013;30(1):108-60.

21. Montalban-Lopez M, Scott TA, Ramesh S, Rahman IR, van Heel AJ, Viel JH, et al. New developments
in RiPP discovery, enzymology and engineering. Nat Prod Rep. 2021;38(1):130-239.

22. Burkhart BJ, Hudson GA, Dunbar KL, Mitchell DA. A prevalent peptide-binding domain guides
ribosomal natural product biosynthesis. Nat Chem Biol. 2015;11(8):564-70.

23. Chekan JR, Ongpipattanakul C, Nair SK. Steric complementarity directs sequence promiscuous leader
binding in RiPP biosynthesis. Proc Natl Acad Sci U S A. 2019;116(48):24049-55.

24. Ayikpoe RS, Zhu L, Chen JY, Ting CP, van der Donk WA. Macrocyclization and Backbone
Rearrangement During RiPP Biosynthesis by a SAM-Dependent Domain-of-Unknown-Function 692.
ACS Cent Sci. 2023;9(5):1008-18.

25. Chioti VT, Clark KA, Ganley JG, Han EJ, Seyedsayamdost MR. N-Calpha Bond Cleavage Catalyzed
by a Multinuclear Iron Oxygenase from a Divergent Methanobactin-like RiPP Gene Cluster. J Am
Chem Soc. 2024.

26. Dou C, Long Z, Li S, Zhou D, Jin Y, Zhang L, et al. Crystal structure and catalytic mechanism of the
MbnBC holoenzyme required for methanobactin biosynthesis. Cell Res. 2022;32(3):302-14.

27. Kenney GE, Dassama LMK, Pandelia ME, Gizzi AS, Martinie RJ, Gao P, et al. The biosynthesis of
methanobactin. Science. 2018;359(6382):1411-6.

28. Nguyen DT, Zhu L, Gray DL, Woods TJ, Padhi C, Flatt KM, et al. Biosynthesis of macrocyclic
peptides with C-terminal beta-amino-alpha-keto acid groups by three different metalloenzymes.
BioRxiv. 2024.

29. Park YJ, Jodts RJ, Slater JW, Reyes RM, Winton VJ, Montaser RA, et al. A mixed-valent Fe(II)Fe(III)
species converts cysteine to an oxazolone/thioamide pair in methanobactin biosynthesis. Proc Natl
Acad Sci U S A. 2022;119(13):e2123566119.

30. Ting CP, Funk MA, Halaby SL, Zhang Z, Gonen T, van der Donk WA. Use of a scaffold peptide in
the biosynthesis of amino acid-derived natural products. Science. 2019;365(6450):280-4.

31. Müller FH, Bandeiras TM, Urich T, Teixeira M, Gomes CM, Kletzin A. Coupling of the pathway of
sulphur oxidation to dioxygen reduction: characterization of a novel membrane-bound
thiosulphate:quinone oxidoreductase. Mol Microbiol. 2004;53(4):1147-60.

32. Nambi S, Long JE, Mishra BB, Baker R, Murphy KC, Olive AJ, et al. The Oxidative Stress Network
of Mycobacterium tuberculosis Reveals Coordination between Radical Detoxification Systems. Cell
Host Microbe. 2015;17(6):829-37.

33. Alvarez-Martinez CE, Baldini RL, Gomes SL. A caulobacter crescentus extracytoplasmic function
sigma factor mediating the response to oxidative stress in stationary phase. J Bacteriol.
2006;188(5):1835-46.

34. Prunier G, Cherkaoui M, Lysiak A, Langella O, Blein-Nicolas M, Lollier V, et al. Fast alignment of
mass spectra in large proteomics datasets, capturing dissimilarities arising from multiple complex
modifications of peptides. BMC Bioinformatics. 2023;24(1):421.

35. Lewis JK, Jochimsen AS, Lefave SJ, Young AP, Kincannon WM, Roberts AG, et al. New Role for
Radical SAM Enzymes in the Biosynthesis of Thio(seleno)oxazole RiPP Natural Products.
Biochemistry. 2021;60(45):3347-61.

36. Wüthrich K. NMR of proteins and nucleic acids. New York: Wiley; 1986.

37. Sattler M. Heteronuclear multidimensional NMR experiments for the structure determination of
proteins in solution employing pulsed field gradients. Progr Nucl Magn Reson Spectr. 1999;34:93-158.

38. Yamazaki T, Lee W, Arrowsmith CH, Muhandiram DR, Kay LE. A Suite of Triple Resonance NMR
Experiments for the Backbone Assignment of 15N, 13C, 2H Labeled Proteins with High Sensitivity. J
Am Chem Soc 1994;116:11655-66.

39. Fiore KE, Patist MJ, Giannakoulias S, Huang CH, Verma H, Khatri B, et al. Structural impact of
thioamide incorporation into a beta-hairpin. RSC Chem Biol. 2022;3(5):582-91.

40. Wishart DS, Bigam CG, Holm A, Hodges RS, Sykes BD. 1H, 13C and 15N random coil NMR chemical
shifts of the common amino acids. I. Investigations of nearest-neighbor effects. J Biomol NMR.
1995;5(1):67-81.

41. NMR spectroscopy - Hans Reich NMR collection - content.

42. Novoa-Aponte L, Xu C, Soncini FC, Arguello JM. The Two-Component System CopRS Maintains
Subfemtomolar Levels of Free Copper in the Periplasm of Pseudomonas aeruginosa Using a
Phosphatase-Based Mechanism. mSphere. 2020;5(6).

43. Bennett B, Kowalski JM. EPR Methods for Biological Cu(II): L-Band CW and NARS. Methods
Enzymol. 2015;563:341-61.

44. Baral BS, Bandow NL, Vorobev A, Freemeier BC, Bergman BH, Herdendorf TJ, et al. Mercury binding
by methanobactin from Methylocystis strain SB2. J Inorg Biochem. 2014;141:161-9.

45. Koh EI, Robinson AE, Bandara N, Rogers BE, Henderson JP. Copper import in Escherichia coli by
the yersiniabactin metallophore system. Nat Chem Biol. 2017;13(9):1016-21.

46. Hirth N, Gerlach MS, Wiesemann N, Herzberg M, Grosse C, Nies DH. Full Copper Resistance in
Cupriavidus metallidurans Requires the Interplay of Many Resistance Systems. Appl Environ
Microbiol. 2023;89(6):e0056723.

47. Clark KA, Seyedsayamdost MR. Bioinformatic Atlas of Radical SAM Enzyme-Modified RiPP Natural
Products Reveals an Isoleucine-Tryptophan Crosslink. J Am Chem Soc. 2022;144(39):17876-88.

48. Antoine R, Alonso S, Raze D, Coutte L, Lesjean S, Willery E, et al. New virulence-activated and
virulence-repressed genes identified by systematic gene inactivation and generation of transcriptional
fusions in Bordetella pertussis. J Bacteriol. 2000;182(20):5902-5.

49. Rivera-Millot A, Slupek S, Chatagnon J, Roy G, Saliou JM, Billon G, et al. Streamlined copper defenses
make Bordetella pertussis reliant on custom-made operon. Commun Biol. 2021;4(1):46.

50. Kitagawa M, Ara T, Arifuzzaman M, Ioka-Nakamichi T, Inamoto E, Toyonaga H, et al. Complete set
of ORF clones of Escherichia coli ASKA library (a complete set of E. coli K-12 ORF archive): unique
resources for biological research. DNA Res. 2005;12(5):291-9.

51. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of Escherichia coli K-
12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol. 2006;2:2006 0008.

52. Garcia-Nafria J, Watson JF, Greger IH. IVA cloning: A single-tube universal cloning system exploiting
bacterial In Vivo Assembly. Sci Rep. 2016;6:27459.

53. Zyubko T, Serebryakova M, Andreeva J, Metelev M, Lippens G, Dubiley S, et al. Efficient in vivo
synthesis of lasso peptide pseudomycoidin proceeds in the absence of both the leader and the leader
peptidase. Chem Sci. 2019;10(42):9699-707.

54. Studier FW. Protein production by auto-induction in high density shaking cultures. Protein Expr Purif.
2005;41(1):207-34.

55. Serra A, Hemu X, Nguyen GK, Nguyen NT, Sze SK, Tam JP. A high-throughput peptidomic strategy
to decipher the molecular diversity of cyclic cysteine-rich peptides. Sci Rep. 2016;6:23005.

56. Langella O, Valot B, Balliau T, Blein-Nicolas M, Bonhomme L, Zivy M. X!TandemPipeline: A Tool
to Manage Sequence Redundancy for Protein Inference and Phosphosite Identification. J Proteome
Res. 2017;16(2):494-503.

57. Shelton KE, Mitchell DA. Bioinformatic prediction and experimental validation of RiPP recognition
elements. Methods Enzymol. 2023;679:191-233.

Acknowledgments. We thank J.-M. Saliou for preliminary MS analyses and J.-Y. Matroule

and P. Cherry for providing *C. vibrioides* strains, plasmids and protocols. This project was

funded by the ANR grant CuRiPP (ANR-22-CE44-0001-02) to FJD. L. Leprevost was

supported by a doctoral fellowship of Lille University. S. Jünger was supported by a PhD

fellowship of the French Ministry of Education (ED 227 MNHN-SU). For the EPR experiments

the financial support from the IR INFRANALYTICS FR2054 is acknowledged. ICP-OES

measurements were performed by V. Alaimo on the Chevreul Institute Platform (U-Lille /

CNRS). The Region Hauts de France and the French government are acknowledged for funding

this apparatus.

Fig. 1. Biosynthetic gene clusters and functions of bufferins. A. Gene composition of the *buf1* and *buf2* BGCs. The signal peptide-coding sequences and core sequences of the *bufA1* and *bufA2* genes are shown in pale and dark grey, respectively. In anti-sense of *buf1ABCD* are genes coding for the extracellular-type sigma factor SigF and the anti-sigma factor NrsF that regulate the expression of *buf1* and *buf2*. B. Effect of copper on bacterial growth. The Cv_{DKO}, $Cv_{buf1ABCD}$ (denoted *buf1*) and $Cv_{buf2ABC}$ (denoted *buf2*) strains were grown in liquid PYE medium without or with 225 μM copper. Their growth was measured in biological quadruplicates. C. Gain-of-function in *Escherichia coli*. Expression of *buf1* in *E. coli* BL21(pCA24-buf1A^{str}BCD) was induced by IPTG, and the bacteria were challenged with 0.3 mM CuSO4 for three hours before serial dilutions. BL21(pCA24-psmCA) expressing a non-relevant lasso peptide BGC was used as a negative control. D. Assay of bacterial lysis by *D. discoidum*. Bacterial densities of the lysis plaques were determined using ImageJ analyses. A non parametric Mann-Whitney test (two-sided) was used to analyze the differences between the Cv_{WT} strain (n=5) and the Cv_{DKO} strain (n=4). A P value of 0.05 was considered statistically significant (* indicates p = 0.0159). The medians are shown (WT = 419,226 and DKO = 344,516) with a confidence interval of 96.83%. The ranks calculated were WT = 35 and DKO = 10.

Fig. 2. Effect of Cu on the growth of *C. vibrioides*. A. Strains that express the indicated genes were grown in PYE medium or in PYE medium containing 225 uM copper. All deletions were performed in strains carrying a single *buf* BGC. B. Strains expressing the entire *buf1* BGC with the indicated substitutions in BufA1 were grown with or without copper as in A.

Fig. 3. Post-translational modifications on Bufferin 1. A. Sequence of Buf1 with the Cys residues labelled I to IV, and top-down LC-MS analyses of cell extracts of Cv_{buf1}(pSigF) compared with Cv_{DKO}(pSigF). Left panel: total ion chromatograms, right panels: deconvoluted mass spectra of the two compounds detected only for Cv_{buf1}(pSigF). Their molecular weights correspond to the core peptide (i.e. after removal of the signal peptide; calculated monoisotopic Mw=6978.35 Da), with mass shifts of -10 Da ($[M+H]^+$ at m/z 6969.32) and -12 Da ($[M+H]^+$ at m/z 6967.30). B. Bottom-up analysis of Buf1. MS/MS spectrum of the central tryptic peptide ($[M+2H]^{2+}$ at m/z 917.36). The two Cys carry carbamidomethyl groups resulting from alkylation with iodoacetamide (+ 57.02 Da), together with -4.03 Da mass shifts. C. UV-visible spectra of purified Buf1 and $Buf1^{CysIISer+CysIIISer}$. The broad absorbance band beyond 290 nm for the wt peptide suggests the presence of heterocycle(s) that are absent in the mutant peptide. D. ^{1}H NMR spectrum of the central tryptic peptide. Anomalous resonances are indicated by orange and green arrows. E. Tripeptide centered on Cys31 (= Cys^{III}) modified into a thio-oxazole. The values of the chemical displacements (in ppm) are given for various atoms. Salient NMR parameters leading to this assignment are the downfield shifted Hα proton of Ser30 (5.26 ppm, orange arrow in D), the ^{13}C assignments of 161.6 and 120 ppm for the ring carbons, the 90-Hz coupling of the latter carbon to its carbonyl resonating at 184 ppm, and the anomalous amide proton chemical shift of Ala32 (9.43 ppm) that follows modified Cys31. The latter signal (green arrow in D) clearly shows that only two Cys residues are modified.

Fig. 4. Analyses of the Bufferin 1-copper complex formed _in vivo_. A. Native mass spectrometry analysis of Buf1[str] purified from a copper-treated culture: enlargement of the isotopic patterns of the major charge state (6+). Upper two panels: purified Buf1[str] from a non-supplemented culture; lower two panels: purified Buf1[str] from _E. coli_ cultures supplemented with $CuSO_4$. The calculated isotopic patterns for $[M+6H]^{6+}$ and $[M+4H+CuII]^{6+}$ species are shown in green. B. Evidence for Cu^{2+} binding to Buf1[str] by EPR spectroscopy. Spectra of Echo-Detected Field-Swept (EDFS) of $CuSO_4$ (red line) and of the Bufferin 1-Cu^{2+} complex (black line) are shown. Differences with free $CuSO_4$ are generated by strongly coordinating ligands, i.e., in the first coordination sphere of Cu^{2+}. At the right, pseudo-modulation of the EDFS spectra. The bottom spectrum shows the four lines typical of free Cu^{2+} absent from the spectrum of the complex. C. Superposition of the ^1H, ^{15}N HSQC spectra of apo bufferin (black spots) and the _in-vivo_ formed copper complex (red spots).

Figure 5. *In vitro* reconstitution of enzymatic activity. A. Workflow of *in vitro* assay. The purified His$_6$-SUMO-BufA1 was first subjected to proteolysis by SUMO protease, generating the precursor BufA1 with an additional 11 aa (IEENLYFQSNA) at the N-terminus. This peptide was subsequently used as a substrate for BufB1 and BufC1, yielding modified BufA1* with two thiooxazoles. The orange box represents the 11-residue sequence. B. MS analysis of the reaction product. *Left panel*: Deconvoluted mass ion of BufA1* ([M+H]$^+$ at *m/z* 10535.1631) after incubation with BufB1 and BufC1 (blue) compared to that of the control reaction ([M+H]$^+$ at *m/z* 10543.1846) (grey)*,* showing a mass loss of 8 Da. The -10 Da and -12 Da forms are not seen here, notably because the reactions were performed in reducing conditions. *Right panel*: LCMS analysis of the 19-residue tryptic fragment from BufA1* generated in vitro. The extracted ion chromatogram (EIC) of [M+3H]$^{3+}$ (at *m/z* 573.8945) and the deconvoluted mass are shown.

Figure 6. *In silico* **analyses of the bufferin families.** A. SSN analyses of the putative signal peptide-containing bufferin-like precursors genetically associated with MNIO enzymes. In the representative node network DUF2282 proteins, including BufA1 of *C. vibrioides*, belong to the largest cluster (red dots). The second largest cluster (yellow dots) represents proteins of the BUF_6/12Cys family that includes BufA2 of *C. vibrioides*. The previously described MNIO enzyme-modified RiPPs of *Chryseobacterium* are found in two small clusters (in orange). B. Weblogos for the major two families of bufferins. C. SSN analyses of MNIO enzymes. In the representative node network (at 80% identity), MNIO enzymes genetically associated with DUF2282 domain-containing proteins, including BufB1 of *C. vibrioides,* and with BUF_6/12Cys proteins, including BufB2 of *C. vibrioides*, are in red and yellow, respectively. MNIO enzymes that modify other known RiPPs - methanobactins, *Chryseobacterium* RiPPs, TglA-type pearlins and aminopyruvatides - are found in small, distinct clusters numbered 1 to 4 and shown in purple, orange, green and dark blue dots, respectively.

Figure 7. Possible mechanism of BufB2 to convert Cys into a 5-thiooxazole motif involving a mixed valent di-iron center. This process starts with O_2 activation and abstraction of a β-H followed by cyclization and ring oxidation.

Extended Fig. 1. Main features of the *buf* BGCs. A. Gene organization of gold-induced genes (*gig*)-like operons found in the genomes of *C. vibrioides (C. crescentus)* (NC_011916.1), *B. pertussis* (NC_002929.2), *C. metallidurans* (NC_007974.2) and *L. pneumophilia* (NC_002942.5). Signal peptide-coding sequences are shown in pale grey in the DUF2282-coding genes. B. Sequence alignment of DUF2282-domain-containing proteins in *gig*-like operons. Predicted signal peptides are underlined and conserved cysteines are highlighted in orange. Consensus motifs are shown above the sequences. Sequence alignment was done by Clustal Omega. Signal peptides and cleavage sites were predicted using the SignalP 5.0 online tool. C. AlphaFold2 models for Buf1 and Buf2 (mature forms, after signal-peptide cleavage), BufB1, BufC1 and BufD1(DoxX) proteins of *C. vibrioides*. The BufC1 protein contains two domains, an all-α-helical DUF2063 domain (orange) and a RRE-like domain (green). In DoxX the 4-helix bundle is predicted to be inserted in the cytoplasmic membrane. N and C show the N and C termini of the proteins.

Extended Fig. 2. Top-down LC-MS analyses of bufferin 2. Top, sequence of Buf2 (core peptide) and bottom, LC-MS analyses of the cell extracts of Cv*buf2*(pSigF) compared with Cv$_{DKO}$(pSigF). Left panel: total ion chromatogram, right panel: deconvoluted mass spectra of the two main peaks detected only for Cv*buf2*. Two compounds were detected with masses corresponding to the expected core peptide (after removal of the signal peptide; calculated monoisotopic Mw 6294.96 Da), showing mass shifts of -18 Da ([M+H]⁺ at *m/z* 6277.79) and -20 Da ([M+H]⁺ at *m/z* 6275.78).

Extended Fig. 3. Reduction-alkylation of Buf1 and Buf2. Top-down LC-MS analyses of the cell extracts of Cv$_{buf1}$(pSigF) (A) or Cv$_{buf2}$(pSigF) (B) submitted to reduction and alkylation with 2-bromoethylamine. Total ion chromatograms compared to that of the untreated extracts (left panels), MS spectra of the main alkylated species detected (middle panels) and zooms on the major charge states (right panels). Minor species corresponding to additional unspecific alkylations (5 and 7 alkylations for Buf1 and Buf2, respectively) were also detected. These data indicate that the presumed PTMs of the Cys residues result in alkylation-sensitive groups.

Extended Fig. 4. Bottom-up analysis of Bufferin 1. MS/MS spectra of the tryptic peptides containing the Cys[I] and Cys[IV] residues, DEHKAGAQTEKCYGVSK ([M+3H]$^{3+}$ at *m/z* 636.63)(A) and GTCVTIKTPK ([M+2H]$^{2+}$ at *m/z* 552.81) (B). The two Cys residues carry each a carbamidomethyl group resulting from alkylation with iodoacetamide (+ 57.02 Da).

Extended Fig. 5 Analysis of the Cys^{II}Ser and Cys^{III}Ser variants of Buf1. A. Sequence of Buf1 and top-down LC-MS analyses of the cell extracts of Cv$_{buf1}$^{CysIISer}(pSigF) or Cv$_{buf1}$^{CysIIISer}(pSigF). Extracted ion chromatograms of the [M+7H]$^{7+}$ species of Buf1^{CysIISer} or Buf1^{CysIIISer} (*m/z* 995) (left), corresponding mass spectra (middle) and deconvoluted spectra (right). The measured *m/z* values (deconvoluted [M+H]$^{+}$ species at *m/z* 6957.36 and 6957.33 for Buf1^{CysIISer} and Buf1^{CysIIISer}, respectively) show -6 Da mass shifts from the masses expected for the proteins with no PTM (calculated monoisotopic Mw 6962.376951). This most likely corresponds to an S-S bond between Cys^{I} and Cys^{IV} and a - 4 Da mass shift for the remaining Cys residue (Cys^{II} or Cys^{III}). B. MS/MS spectra of the tryptic peptides AGEND**S**KAGAGTS**C**AGTSKVDYQGDAWK for Buf1^{CysIISer} ([M+4H]$^{4+}$ at *m/z* 707.56) and AGEND**C**KAGAGTS**S**AGTSKVDYQGDAWK for Buf1^{CysIIISer} ([M+3H]$^{3+}$ at *m/z* 943.08). For both variants, the unsubstituted Cys carry a carbamidomethyl group resulting from alkylation with iodoacetamide (+ 57.02 Da), together with a -4.03 Da mass shift.

Extended Fig. 6. *In vivo* production and purification of the central tryptic peptide. A. Modified BufA1* was produced through co-expression with BufB1 and BufC1 in *E. coli* and purified via Ni-affinity chromatography. The obtained peptide was digested with trypsin generating the 19-residue peptide 1 containing modifications on the two Cys residues (marked with *). B. Chromatogram of HPLC purification (UV = 280 nm). The peak at RT = 13.85 min was collected and corresponds to peptide 1. The inset shows the corresponding UV/vis scan exhibiting a new absorbance maximum at 305 nm. C. Mass spectrum of purified unlabelled (left panel) and ^{13}C- and ^{15}N-labelled peptide 1 (right panel). D. MS/MS spectrum of the central tryptic peptide ([M+H]$^{3+}$ at m/z 573.8945).

Extended Fig. 7 NMR experiments to assign the resonances of the 19-mer peptide (A18-K37). A. ROESY/Tocsy experiments. B. Natural abundance ^{1}H, ^{15}N HSQC spectrum. C and D. ^{1}H, ^{13}C HMBC experiments on the 19-mer peptide at natural isotopic abundance.

Extended Fig. 8. NMR experiments on the uniformly ^{13}C, ^{15}N labelled 19-mer peptide. A. HNCACB experiment. Sequential walk shown here for the sequence following Cys[II] (Lys24-Ala27). For Lys24 we did not find any resonances that could correspond to its upstream neighbour, whereas for downstream residues the amide resonance of a given residue Xi is coupled both to its own Cα/Cβ and the Cα/Cβ resonances of the preceding (Xi-1) residue. B. The HNOC plane shown at the ^{15}N frequency of the indicated residues. Whereas we find classical values around 175 ppm for all residues of the peptide, the residues downstream of a modified Cys resonate at 184 ppm (B). C. HNOC spectrum. Shown is the HNCO plane through the resonance of Ala32 (that immediately follows Cys[III]) while modifying the offset of the decoupling pulse during the C=O evolution period.

Extended Fig. 9. Structural characterization of bufferin 1. A. HSQC-NOESY and TOCSY experiments: sequential walks through residues of the first and second β strands, respectively. B. Strips from TOCSY-HSQC (blue) and NOESY-HSQC (red) spectra of bufferin 1. Characteristic Hα(i-1) – HN(i) NOE contacts are indicated by arrows. These data confirm the assignement and demonstrate the extended (β strand) nature of this segment. C. Strips through the TOCSY-HSQC (blue) and NOESY-HSQC (red) spectra of bufferin 1. These data show the inter-strand HN-HN NOE contacts, confirming the presence of an anti-parallel β sheet.

A

His-SUMO-MTRTATTAALAAAIALSAGVSIAAA-DEHKAGAQTEKCYGVSKAGENDCKAGAGTSCAGTSKVDYQGDAWKMVKKGTVCTIKTPKGMGSLTPKA

(-8 aa)   (CP)

B   **C**

Extended Fig. 10. The signal-peptide sequence is not essential for heterocycle formation. A. Schematic illustration of variant BufA1 with signal peptide truncations. The signal peptide is in orange, and the four Cys in the core peptide are highlighted in red. B. LCMS analysis of modified BufA1(-8aa). MS spectrum of the most abundant charge state [M+9H]$^{9+}$, corresponding to a deconvoluted monoisotopic mass [M+H]$^{+}$ of 8391.1281, is shown (blue) and compared to the theoretical spectrum of unmodified peptide (black). C. LCMS analysis of modified BufA1(CP). *Left panel*: The fully converted peptide was eluted at RT 9.6 min. The MS spectrum of the most abundant charge state [M+8H]$^{8+}$, corresponding to a deconvoluted monoisotopic mass [M+H]$^{+}$ of 8277.9398, is shown (blue) and compared to the theoretical spectrum of unmodified peptide (black). *Right panel*: unmodified peptide (green) was eluted with the modified one (blue) at RT 9.5 min. Its MS spectrum of the charge state [M+8H]$^{8+}$, corresponding to a deconvoluted monoisotopic mass [M+H]$^{+}$ of 8287.8888, is shown.