A widespread family of ribosomal peptide metallophores involved in bacterial adaptation to copper stress

Laura Leprevost[#1], Sophie Jünger[#2], Guy Lippens[3], Céline Guillaume[2], Giuseppe Sicoli[4],

Lydie Oliveira[5], Alex Rivera-Millot[1&], Gabriel Billon[4], Céline Henry[5], Rudy Antoine[*1],

Svetlana Dubiley[*3], Séverine Zirah[*2], Yanyan Li[*2], Françoise Jacob-Dubuisson[*1]

[1] Univ. Lille, U1018-UMR9017-CIIL-Center for infection and Immunity of Lille, F-59000 Lille, FRANCE

[2] Unit Molecules of Communication and Adaptation of Microorganisms (MCAM), UMR 7245 CNRS, MNHN, 75005 Paris, FRANCE

[3] Toulouse Biotechnology Institute, CNRS/INRAE/INSA/UPS, 31077 Toulouse, FRANCE

[4] Univ. Lille, CNRS, UMR 8516 - LASIRE, Laboratoire de Spectroscopie pour les Interactions, la Réactivité et l'Environnement F-59000 Lille, FRANCE

[5] INRAE-AgroParisTech-Univ. Paris-Saclay, MICALIS, 78352 Jouy-en Josas, FRANCE

[#] contributed equally

[*] for correspondence: rudy.antoine@pasteur-lille.fr; svetlana.dubiley@gmail.com; severine.zirah@mnhn.fr; yanyanli@mnhn.fr; francoise.jacob@ibl.cnrs.fr

[&] current address: INRS-Centre Armand-Frappier Santé Biotechnologie, Bacterial Symbionts Evolution, Laval, Quebec H7V 1B7, Canada

Abstract

Ribosomally synthesized and post-translationally modified peptides (RiPPs) are a structurally diverse group of natural products that bacteria employ in their survival strategies. Herein, we characterized the mode of action, structure, and biosynthetic pathway of bufferins, products of mysterious 'gold-induced genes' (*gig)*-like operons. We show that bufferins form a new family of RiPPs ubiquitous in Eubacteria. Using *Caulobacter vibrioides* as a model organism, we elucidate the important role of bufferins in protecting bacteria from copper-induced stress. The cleavable N-terminal signal peptide guides bufferins export to the periplasm, where they efficiently sequester the copper ions. The metal-binding propensity of bufferins relies on the post-translational transformation of cysteine residues in the precursor peptide into 5-thioxazoles. This rare modification is installed by a multinuclear non-heme iron-dependent oxidative enzyme (MNIO, DUF692 family) in conjunction with its partner protein belonging to the DUF2063 family. Our study broadens the landscape of known peptide modifications installed by MNIO enzymes and unveils a widespread but overlooked bacterial copper defense strategy.

Synopsis: New copper-chelating natural peptide products with 5-thiooxazole modification installed by mutinuclear non-heme iron dependent oxidative enzyme, involved in bacterial adaptation to excess copper

Introduction

Several transition metals including copper are essential for living organisms [1,2]. However, an excess of copper is toxic to the cell notably due to the mismetallation of non-cognate proteins that results in the inactivation of essential pathways [2-4]. Bacteria encounter copper in natural habitats and within eukaryotic cells, which use it as a bactericidal agent [5,6]. Therefore, they have developed an arsenal of defense systems, including tight regulation of copper uptake, various export systems, and sequestration/storage by copper-chelating molecules or proteins [7,8]. For pathogens, these strategies are critical for host infection by enhancing their survival in phagocytes, and hence they are linked to virulence [9].

Previous work on *Bordetella pertussis* has identified a four-gene operon controlled by the master regulator of virulence, BvgAS [10]. This biosynthetic gene cluster (BGC) is similar to the *gig* (gold-induced genes) operons of *Cupriavidus metallidurans* and *Legionella pneumophila* and to a related operon of *Caulobacter vibrioides* [11-13] (Fig. 1A; Suppl. Fig. S1). The latter operons are up-regulated upon exposure to transition metals, notably copper [11-14]. In particular, the *C. vibrioides* operon is part of the few genes induced by copper in both swarmer and stalked cells [14]. Although these data suggest a role for *gig*-like genes in copper stress response, the nature and the function of produced molecules has remained a mystery. The *gig*-like operons contain a gene encoding a DUF692 family protein, which was recently associated with the biosynthesis of several ribosomally synthetized post-translationally modified peptides (RiPPs) [15-21]. This suggests that the product of the *gig*-like operons could be also a post-translationally modified peptide.

RiPPs are natural products with a tremendous variety of structures and biological activities [22-24]. They are synthesized from a gene-encoded precursor peptide, generally composed of an N-terminal leader sequence and a core sequence where post-translational modifications (PTMs) occur. In many cases, a pathway-specific RiPP recognition element (RRE) domain interacts

with the leader sequence and mediates recruitment of the modification enzymes [25-27]. RRE

domains are thus a hallmark feature of RiPP biosynthesis. The vast structural diversity of RiPPs

is conferred by virtually unlimited possibilities of chemical transformations. An emerging

family of PTM enzymes is the DUF692 family of multinuclear non-heme iron dependent

oxidases (MNIOs) that catalyze various reactions in the biosynthesis of diverse RiPPs, mostly

but not exclusively on Cys residues [15-21]. However, the full spectrum of reactions catalyzed by

MNIOs remains far from being fully appreciated.

Here, using *C. vibrioides* as a model, we report the first functional and structural

characterization of the molecules produced by *gig*-like operons. Our study unveiled that they

represent a new family of RiPP metallophores, which contain a rare 5-thiooxazole modification

of cysteine residues installed by MNIO enzymes. Based on our finding that the products of *gig*-

like operons contribute to bacterial adaptation to toxic concentrations of copper by complexing

with the excess metal, we named these RiPPs bufferins.

Results

Gig-like operons belong to putative RiPP BGCs. To gain insight into the gene function of *gig*-

like operons, we performed an *in-silico* analysis on the CCNA_03363-03366 operon of *C.*

vibrioides, hereafter named *buf1*. CCNA_03363 encodes a DUF2282-family protein (BufA1),

whose members are 80-120 residues long and harbor four Cys residues (Cys^I to Cys^{IV}) and a

predicted Sec-dependent signal peptide (Fig. 1A). We hypothesized that BufA1 is an unusually

long precursor peptide whose translocation through the cytoplasmic membrane of the producing

cell would rely on the ubiquitous Sec export pathway rather than on a specialized export pump

encoded in its BGC as is often the case for other secreted RiPPs [28]. The AlphaFold2 model

predicts that the BufA1 core peptide is likely to contain a 2-stranded β sheet and some α-helical

structure (Suppl. Fig. S1). In the absence of PTMs, it would thus fold into a small protein, rather

Figure 1. *In silico* analyses of the bufferin family. A. Gene composition of the *buf1* and *buf2* BGCs. The sequences of BufA1 and BufA2 are shown with the Cys residues labeled (numbering according to the core peptides). The signal peptide and core peptide sequences are in pale and dark grey, respectively. In anti-sense of *buf1ABCD* are genes coding for an extracellular-type sigma factor, SigF, and an anti-sigma factor, NrsF, that regulate *buf1* and *buf2* expression. **B**. SSN analyses of the putative signal peptide-containing bufferin-like precursors genetically associated with MNIOs. In the representative node network DUF2282 proteins, including BufA1 of *C. vibrioides*, belong to the largest cluster (red dots). The second largest cluster (yellow dots) represents BUF_6/12Cys proteins, which include BufA2 of *C. vibrioides*. The previously described MNIO-modified RiPPs of *Chryseobacterium* have predicted signal peptides and are found in two small clusters (in orange). **C**. Weblogos for the DUF2282 and BUF_6/12Cys bufferins. **D**. SSN analyses of MNIOs. In the representative node network (at 80% identity), MNIOs genetically associated with DUF2282 domain-containing proteins, including BufB1 of *C. vibrioides,* and with BUF_6/12Cys proteins, including BufB2 of *C. vibrioides*, are in red and yellow, respectively. MNIOs that modify other known RiPPs - methanobactins, *Chryseobacterium* RiPPs, TglA-type pearlins and aminopyruvatides - are found in small, distinct clusters numbered 1 to 4 and shown in purple, orange, green and dark blue dots, respectively.

than being a fully unstructured peptide. The product of CCNA_03364 (BufB1) is predicted to belong to the MNIO family, whereas CCNA_03365 codes for a protein (BufC1) with an N-terminal DUF2063 domain of unknown function. The structural model of BufC1 revealed that its C-terminal domain is structurally related to RRE domains [25, 26, 27] (Suppl. Fig. S1). In light of the recent discovery of MNIO enzymes involved in RiPP biosynthesis [15-21], we hypothesized that BufB1 and BufC1 would install modifications on Cys residues of BufA1. Finally, CCNA_03366 encodes a putative DoxX-type oxido-reductase (BufD1) [29, 30]. The other known *gig*-like operons share the same gene organizations and predicted functions (Suppl. Fig. S1).

Inspection of the *C. vibrioides* genome identified a second operon, the *buf2* BGC (CCNA_02999-03001) (Fig. 1A) which encodes homologs of BufB1 (CCNA_03000) and BufC1 (CCNA_02999) but lacks a *doxX*-like gene. Its putative precursor peptide BufA2 (CCNA_03001) possesses a Sec-dependent signal peptide, like BufA1, but its core sequence contains six rather than four Cys residues (Cys^I to Cys^{VI}) and is without family signature (Fig. 1A). The predicted 3D structure of the BufA2 core peptide adopts a very similar fold to that of BufA1 (Suppl. Fig. S1).

To evaluate the prevalence of *gig*-like BGCs in prokaryotes, we collected >14,000 MNIO protein sequences from the NCBI non-redundant database. The retrieved sequences were widely distributed among Proteobacteria, Terrabacteria, Myxococcota, Fibrobacteres/Chlorobi/Bacterioidetes, Planctomycetes/Verrucomicrobia/Chlamydia, and Acidobacteriota groups (detailed *in silico* analyses will be published separately). We analyzed their genomic contexts and collected associated genes coding for putative precursor peptides, i.e., proteins smaller than 200 residues with Cys residues and a predicted N-terminal signal peptide. In a sequence similarity network (SSN) constructed with those proteins, the largest cluster corresponds to proteins with the DUF2282 signature, including *C. vibrioides* BufA1 (Fig. 1B). BufA2 of *C. vibrioides* belongs to the second largest cluster of proteins characterized

by six or twelve conserved Cys residues, which we named BUF_6/12Cys (Fig. 1B,C). Parallel

SSN analyses of the MNIO enzymes superfamily (Fig. 1D) showed that those associated with

DF2282 or BUF_6/12Cys bufferins belong to the most populated sequence cluster, distinct

from the smaller clusters of enzymes involved in the biosynthesis of methanobactins, RiPPs

from *Chryseobacterium*, TglA-type pearlins and amino-pyruvatides [15, 17-19, 21]. Together out

data indicate that bufferins currently form the largest group of RiPPs whose BGCs harbor

MNIO genes that are distinct from all previously described MNIO enzyme-modified RiPPs.

Bufferins are involved in adaptation to copper stress. In *C. vibrioides*, *buf1* and *buf2* BGCs

belong to the SigF-controlled, core regulon of copper stress [12, 14, 31]. To test if they could be up-

regulated by other biologically relevant transition metals, we generated reporter fusions within

bufA1 and *bufA2* and confirmed that both are induced by Cu^{2+} but not Fe^{2+}, Zn^{2+} or Mn^{2+} (Fig.

2A).

To determine if bufferins, the putative RiPP products of the *buf* BGCs, contribute to

adaptation to copper stress, we generated *C. vibrioides* deletion mutants that express a single

(Cv_{buf1} and Cv_{buf2}) or neither BGC (Cv_{DKO}). Cv_{DKO} grew more slowly than the parental Cv_{WT}

strain when $CuSO_4$ was added to the growth medium, unlike Cv_{buf1} or Cv_{buf2} (Fig. 2B). Thus,

bufferins enhance growth under copper stress and are functionally redundant. To confirm that

a single BGC is sufficient to protect bacteria from excess copper, we performed a gain-of-

function experiment in an *Escherichia coli* laboratory strain. For this, we expressed the

buf1ABCD operon under the control of a P_{T5-lac} promoter upon IPTG induction. Expression of

buf1ABCD enhanced survival of *E. coli* exposed to a bactericidal concentration of $CuSO_4$ (Fig.

2C).

We then asked whether the protection from copper would confer an advantage on *C.

vibrioides* in biotic interactions. As amoeba notably use copper against their bacterial preys [5],

Figure 2. Regulation and function of the bufferins. **A**. Reporter assays with *bufA1-lacZ* and *bufA2- lacZ* transcriptional fusions. Bacteria were grown for 16 h with the indicated concentrations of $CuSO_4$, $FeSO_4$, $ZnSO_4$ or $MnCl_2$. All experiments were performed in biological quadruplicates, with three technical replicates. Non-parametric, one-way analysis of variance Kruskal-Wallis (two-sided) tests followed by a Dunn's multiple-comparison tests were used to analyze the differences between treatment with various compounds and the control (n=4; * in the upper left graph indicates p=0.047). **B**. Effect of copper on bacterial growth. The Cv_{DKO}, $Cv_{buf1ABCD}$ (*buf1*) and $Cv_{buf2ABC}$ (*buf2*) strains were grown without or with 225 μM $CuSO_4$, in biological quadruplicates. **C**. Gain-of-function assay. Expression of *buf1* in *E. coli* BL21(pCA24-buf1A^{str}BCD) was induced by IPTG, and the bacteria were challenged with 0.3 mM $CuSO_4$ for three hours before serial dilutions. BL21(pCA24-psmCA) expressing a non-relevant RiPP BGC was used as a control. **D**. Assay of bacterial lysis by *D. discoidum*. Bacterial densities of the lysis plaques were determined using ImageJ analyses. A non-parametric Mann-Whitney test (two-sided) was used to analyze the differences between the Cv_{WT} strain (n=5) and the Cv_{DKO} strain (n=4; * indicates p = 0.0159). The medians are shown (WT = 419,226 and DKO = 344,516) with a confidence interval of 96.83%. The ranks calculated were WT = 35 and DKO = 10.

a predation assay was performed. However, our attempts to measure survival of *C. vibrioides*

after ingestion by *Dictyostelium discoidum* were unsuccessful. Therefore, we tested if the

presence of the *buf* BGCs might slow down bacterial killing. We placed amoebal suspensions

onto bacterial lawns, and after incubation we determined bacterial densities in the contact zones

by densitometry scanning. Larger zones of complete bacterial clearance were observed for the

Cc_{DKO} than for the Cc_{WT} strain, consistent with a protective role of bufferins (Fig. 2D).

Characterization of PTMs of bufferins. To identify the PTMs of bufferins, we engineered an

IPTG-inducible version of SigF to control expression of their BGCs in *C. vibrioides.* The

metabolic profiles of IPTG-treated Cv_{buf1}(pSigF) and Cv_{buf2}(pSigF) were analyzed by liquid

chromatography coupled to mass spectrometry (LC-MS) and compared to that of Cv_{DKO}(pSigF)

to reveal peptide species only present in the *buf*-expressing strains. In Cv_{buf1}(pSigF), a major

species exhibited a mass matching that of BufA1 with the Sec-dependent signal peptide cleaved

off and with a mass shift of -10 Da (Fig. 3A; Suppl. Fig. S2), suggesting that the core peptide

accumulates in the periplasm and contains post-translational modifications of one or more

residues. A second form with a mass shift of -12 Da was also detected, but not systematically,

and it disappeared upon freeze-thawing. The most stable product with a -10 Da mass shift is

considered the mature peptide product and named bufferin 1 (Buf1). Similarly, Cv_{buf2}(pSigF)

produced a major peptide product with a mass shift of -18 Da, called bufferin 2 (Buf2), and

another species with a -20 Da mass shift (Suppl. Figs. S2 and S3).

We speculated that mature bufferins contain modified Cys residues and disulfide (S-S)

bonds. To test this hypothesis, we performed reduction and alkylation treatment of the peptides.

This modification yielded species carrying three or four alkylated groups for Buf1 and four,

five or six alkylated groups for Buf2 (Suppl. Fig. S4), suggesting that the expected PTMs either

do not affect Cys residues or result in alkylation-sensitive groups. MS analyses of trypsin-

Figure 3. Post-translational modifications of bufferin 1. **A.** Top-down LC-MS analyses of cell extracts of Cv_{buf1}(pSigF) compared with Cv_{DKO}(pSigF). Left panel: total ion chromatograms, right panels: deconvoluted mass spectra of the two compounds detected only for Cv_{buf1}(pSigF). Their MW correspond to the core peptide (i.e. after removal of the signal peptide; calculated monoisotopic Mw=6978.35 Da), with mass shifts of -10 Da ([M+H]$^+$ at m/z 6969.32) and -12 Da ([M+H]$^+$ at m/z 6967.30). **B.** Bottom-up analysis of Buf1. MS/MS spectrum of the central tryptic peptide ([M+2H]$^{2+}$ at m/z 917.36). The two Cys residues carry carbamidomethyl groups resulting from alkylation with iodoacetamide (+ 57.02 Da), together with -4.03 Da mass shifts. **C.** UV/vis spectra of affinity-purified Buf1 and its Cys^{II}Ser+Cys^{III}Ser mutant version show an absorbance maximum at 305 nm for the first one only. **D.** NMR ROESY/TOCSY experiments to assign the resonances of the 19-mer peptide (Ala18-Lys36). **E.** HNCACB experiment on the uniformly ^{13}C, ^{15}N labeled 19-mer peptide: sequential walk for the sequence following Cys^{II} (Lys24-Ala27). For Lys24, we only identify its own Cα/Cβ carbons and no Cα/Cβ resonances of a preceding residue as is the case for the other residues. **F.** The HNCO planes at the ^{15}N frequency of the indicated residues show classical values around 175 ppm for all carbonyl resonances except for those residues downstream of a modified Cys, whose carbonyl carbons resonate at 184 ppm. **G.** HNCO planes through the resonance of Lys24 that immediately follows Cys23 (=Cys^{II}) while modifying the offset of the ^{13}C decoupling pulse during the C=O evolution period. Only when centered at 120 ppm, the 90-Hz carbon-carbon coupling is refocused. **H.** Measured NMR parameters and proposed structure for the bufferin 1 post-translational modifications (shown for Cys^{II}).

digested, alkylated bufferins revealed that in Buf1, Cys^I (Cys12) and Cys^{IV} (Cys52) (numbering

according to the core peptide, see Fig. 1) had mass increments corresponding to alkylation only,

indicative of unmodified residues (Suppl. Fig. S5). By contrast, in the central tryptic peptide

^{18}AGENDCKAGAGTSCAGTSK^{36} containing Cys^{II} (Cys23) and Cys^{III} (Cys31), both Cys

residues carried mass increments corresponding to alkylation with additional -4.03 Da mass

shifts (Fig. 3B), as proposed by the untargeted PTM search solution SpecGlobX [32].

To confirm that Cys residues are required for the bufferin function, we constructed strains

with the four conserved Cys residues of Buf1 replaced with Ser, separately or in combinations.

In line with the proposed modifications of Cys^{II} or Cys^{III}, the substitution of any of the two with

Ser in Buf1 yielded peptides with mass shifts of -6 Da relative to the calculated mass of the

mutated peptide (Suppl. Fig. S6). We tested the ability of the mutated BufA1 to protect *C.*

vibrioides from excess copper. As expected, in the presence of $CuSO_4$ mutant strains grew more

slowly than the parental Cv$_{buf1}$ (Suppl. Fig. S7), indicating the importance of both modified and

unmodified Cys residues for function. Altogether, the data are consistent with the presence of

an S-S bond between Cys^I and Cys^{IV} (resulting in -2 Da), and the central Cys^{II} and Cys^{III} being

targets of 4-electron oxidation reactions (-4 Da each) that generate alkylation-sensitive groups.

The minor Buf1 congener product with an overall mass shift of -12 Da would then correspond

to the formation of a labile S-S bond between the modified residues. Similarly, in Buf2, in

addition to the unmodified Cys^I (Cys24) and Cys^{VI} (Cys50) which might form an S-S bond

according to the AlphaFold2 model (Suppl. Fig. S1), the central Cys residues carry -4.03 Da

shifts (Suppl. Fig. S8).

To facilitate bufferin purification and further characterization of peptides, we added C-

terminal tags to BufA1. The affinity-purified tagged bufferin 1 was fully modified, displaying

-10 Da mass shifts relative to the calculated mass (Suppl. Fig. S9; Suppl. Table S1). To probe

if the modified Cys residues may be part of heterocycles as with other MNIO enzyme-catalyzed

PTMs [15, 18], we recorded UV-visible absorption spectra of purified Buf1 and the

Buf1[CysIISer+CysIIISer] variant. Indeed, consistent with the presence of aromatic heterocycles [33], the

spectrum of the former showed a broad absorbance peak extending to 310 nm, unlike that of

the latter (Fig. 3C). Similarly, the spectrum of recombinant Buf2 showed an absorbance

maximum at 305 nm (Suppl. Fig. S10).

Identification of the PTMs of bufferin 1 by NMR. To increase the yield of bufferin for

structural elucidation by NMR, we co-expressed *bufA1* carrying an N-terminal His$_6$-SUMO-tag

with *bufB1* and *bufC1* in *E. coli*. Subsequent purification of modified BufA1 followed by

trypsin digestion and preparative HPLC allowed to obtain the central 19-mer peptide

encompassing Cys[II] (Cys23) and Cys[III] (Cys31) at a milligram scale (Suppl. Fig. S11). Its [1]H

spectrum indicated two groups of anomalous resonances (Suppl. Fig. S12). By combining

homonuclear ROESY and TOCSY experiments [34], they were assigned to the amide protons of

Lys24 and Ala32 that follow the modified Cys[II] and Cys[III] residues, respectively, and to the Hα

protons of Asp22 and Ser30 that precede them (Fig. 3D). Further analysis of the same spectra

showed the absence of both Hα and NH protons for the modified Cys residues. The absence of

the latter was verified on a natural abundance [1]H, [15]N HSQC spectrum (Suppl. Fig. S12). These

data suggest that their N and Cα atoms are part of heterocycles.

A [1]H, [13]C HSQC experiment showed correlations from the Asp22 and Ser30 Hα atoms to

their own Cα carbons at 48.1 and 52.4 ppm, respectively, which is 6 ppm higher than their

random coil values [35]. The [1]H, [13]C HMBC spectrum connected the same Hα protons to carbon

atoms at 161.1 ppm and 159.8 ppm, respectively (Suppl. Fig. S12). We produced a uniformly

[13]C,[15]N-labeled modified BufA1 molecule in *E. coli* using the same expression system (Suppl.

Fig. S11), and we used its central tryptic peptide to obtain further carbon assignments on the

basis of triple-resonance NMR experiments [36, 37]. In the HNCACB experiment, we found no

resonances that could correspond to upstream neighbors for Lys24 or Ala32, confirming that

the Cα carbons of modified Cys[II] and Cys[III] do not resonate at the typical 50-ppm value

expected for Cys residues (Fig. 3E). In a HNCO experiment starting from the amide groups of

Lys24 or Ala32, we found values around 184 ppm for the C=O carbon atoms of both modified

Cys residues (Fig. 3F), far from 175-ppm and 200-ppm values expected for such atoms in

random-coil or thio-amide peptides, respectively [35, 38]. By increasing the resolution and varying

the offset of the decoupling pulse during the C=O evolution period, we determined a 90-Hz

coupling constant of both carbonyls with carbon atoms whose chemical shift is ~120 ppm,

consistent with their absence in the HNCACB experiment (Fig. 3G). Both parameters point to

the C=O carbon atoms of the modified Cys residues connected to carbon atoms in aromatic

rings [39].

When we compare the chemical shifts assigned on the modified BufA1 peptide with those

found for a 5-thiooxazole-containing natural product [33], we find the same 6-ppm upfield shift

of the Cα carbon of the preceding residue, the presence of similar carbon resonances at 120 and

160 ppm, and the 90-Hz coupling constant of the following carbonyl towards an aromatic ring

carbon. Together with the MS data and with the characteristic spectrophotometric absorbance

maximum at 305 nm [33], we conclude that the Buf1 modification implies cyclization of both its

central Cys residues to form thiooxazole groups (Fig. 3H).

Copper binding by bufferin 1. Given that heterocycles can be involved in copper binding [40],

we interrogated if bufferins can complex with copper, which could be the basis for their

protective effect. As our attempts to saturate bufferin 1 with copper *in vitro* proved to be

inefficient (Suppl. Fig. S13), we investigated complex formation *in vivo* by treating cultures of

bufA1-expressing *E. coli* with $CuSO_4$ for three hours prior to bufferin purification. Native MS

analysis showed that Buf1[str] thus produced was fully loaded with copper (Fig. 4A). Pulsed

Figure 4. Analyses of the bufferin 1-copper complex formed *in vivo*. A. Native mass spectrometry analysis of purified Buf1^{str} loaded with Cu *in vivo*: isotopic patterns of the major charge state (6+). The upper two and lower two panels show Buf1^{str} from a non-supplemented culture and from cultures supplemented with CuSO$_4$, respectively. The calculated isotopic patterns for [M+6H]$^{6+}$ and [M+4H+CuII]$^{6+}$ species are shown in green. **B.** Evidence for Cu^{2+} binding to Buf1^{str} by EPR spectroscopy. Spectra of Echo-Detected Field-Swept (EDFS) of CuSO$_4$ (red line) and of the Buf1-Cu^{2+} complex (black line) are shown. Pseudo-modulation of the EDFS spectra is shown underneath. The panels at the right show the Easyspin fits (in red) and the experimental spectra (in black) for the Buf1-Cu^{2+} complex (top) and CuSO$_4$ (bottom). **C.** Superposition of the ^{1}H, ^{15}N HSQC spectra of apo bufferin (black peaks) and the *in-vivo* formed Buf1-Cu^{2+} complex (red peaks). sc= side chain.

electron paramagnetic resonance (EPR) spectra confirmed that Buf1 binds paramagnetic Cu^{2+},

and spectral simulations indicated coordinating N atoms [41] (Fig. 4B; Suppl. Tables S2 and S3).

To gain further insight into the copper binding site, we produced a $^{15}N,^{13}C$-labeled Buf1

sample and recorded triple resonance spectra to assign its resonances. Simultaneously, we used

the ^{1}H, ^{15}N HSQC plane as a basis for HSQC-NOESY and TOCSY experiments to confirm the

assignments and gain some knowledge on structural aspects of the bufferin. The sequential

assignment was interrupted by the modified Cys residues, as expected. Interestingly, amide

resonances of the tryptic peptide and of the same segment in the full-length bufferin do not

correspond at all, suggesting that the full-length sequence imposes structural constraints on the

thiooxazole-containing loop. Part of these likely come from the β sheet spanning two short

strands in the structural model, Thr9-Tyr13 and Trp44-Lys48 (Suppl. Fig. S1), that we could

confirm experimentally through the strong Hα(i-1) – HN(i) NOE contacts between residues in

these segments and the cross-strand NOE contacts between the amide protons of Lys45 and

Cys12, and of Val47 and Glu10, respectively (Suppl. Fig. S14). PTMs of the Cys residues hence

do not interfere with the 3D structure predicted for the unmodified sequence.

A comparison of the spectra of the bufferin produced with or without copper showed that

whereas many peaks remained unchanged, some experienced a shift and some signal reduction,

and yet others disappeared altogether (Fig. 4C). The latter most probably concern residues close

to the paramagnetic Cu^{2+} ion. In particular, disappearance of the peaks corresponding to direct

neighbors of the thiooxazole heterocycles, such as Lys24, Ser30 and Ala32, confirms that the

heterocycles are involved in Cu^{2+} coordination. Interestingly, signals of Thr9, Trp44, Met46

and Lys48 predicted on the β-sheet face oriented towards the thiooxazole-containing loop also

disappeared in the spectrum of the Cu^{2+}-loaded Buf1. Whereas a full structural determination

of bufferin will require further experiments, our data support that the thiooxazole groups and

residues of the β sheet are involved in copper binding.

BufB1 and BufC1 are required for post-translational modification of bufferin. To dissect the role of individual *buf* genes in the function of bufferins, we constructed Cv$_{buf1}$ derivatives that lack one or several genes and tested their growth phenotypes in the presence of CuSO$_4$. All deletion strains except for that lacking *bufD1* displayed slow-growth phenotypes, showing that BufA1, BufB1 and BufC1, but not BufD1, are required for function in these conditions (Fig. 5A).

To reveal the function of BufB1 in the bufferin maturation, we reconstituted the modification reaction *in vitro* using the purified recombinant SUMO-BufA1, BufB1 and BufC1 proteins (Suppl. Fig. S15). Since MNIO enzymes require Fe^{II} and Fe^{III} ions for activity [17, 20], BufB1 was analyzed by inductively coupled plasma - optical emission spectrometry, showing the presence of two Fe ions. Structural alignments of a BufB1 model with the crystal structure of MbnB of *Vibrio caribbeanicus*, an MNIO enzyme involved in methanobactin biosynthesis [17] confirmed that the Fe-coordinating residues are conserved in BufB1. Next, BufA1 with the SUMO-tag cleaved was incubated with BufB1 and/or BufC1. LC-MS analyses showed that BufA1 was converted to a species with a -8 Da mass shift only when both BufB1 and BufC1 were present (Fig. 5B). Trypsin digestion of the *in vitro* modified BufA1 yielded the central 19-residue fragment harboring the thiooxazole groups derived from Cys^{II} and Cys^{III} as confirmed by MS/MS analyses (Fig. 5C). Together with the gene inactivation data, these experiments established that BufB1 and BufC1 are required and sufficient to install the PTMs on BufA1.

The signal-peptide sequence is dispensable for heterocycle formation. The particularity of bufferin biosynthesis is that BufA precursors harbor a cleavable Sec-dependent export signal peptide sequence at their N-terminus, which is an uncommon feature of bacterial RiPPs precursors. We investigated if this sequence also acts as a leader peptide, required for Cys

Figure 5. Biosynthesis of bufferin 1. A. Effect of Cu on the growth of *C. vibrioides* harboring various gene knock-outs in *buf1* BGC. All deletions were performed in Cv_{buf1}. Strains expressing the indicated genes were grown in PYE medium or in PYE medium containing 225 μM CuSO$_4$. **B.** *In vitro* reconstitution of the bufferin 1 biosynthesis. The deconvoluted mass spectra for the products of the *in vitro* reactions are shown, with the proteins added to the reactions indicated in the corresponding panels. The control reaction contained BufA1 and heat-denatured BufB1 and BufC1.
C. LC-MS analysis of the 19-residue tryptic fragment of the *in vitro* modified bufferin 1. The extracted ion chromatogram (EIC) of [M+3H]$^{3+}$ (at *m/z* 573.8945) and the deconvoluted mass are shown. **D**. Reconstitution of the system in heterologous host to address the role of the signal peptide for the PTMs. Schematic illustration of variant BufA1 with signal peptide truncations is shown on top. The signal peptide is in orange. *Left panel:* LC-MS analysis of modified BufA1(-8aa). MS spectrum of the most abundant charge state [M+9H]$^{9+}$, corresponding to a deconvoluted monoisotopic mass [M+H]$^+$ of 8391.1281 (blue) is compared to the theoretical spectrum of unmodified peptide (black). *Right panel*: LC-MS analysis of modified BufA1(CP). The MS spectrum of the most abundant charge state [M+8H]$^{8+}$, corresponding to a deconvoluted monoisotopic mass [M+H]$^+$ of 8277.9398 (blue) is compared to the theoretical spectrum of unmodified peptide (black).

transformation. Using the co-expression system of SUMO-*bufA1 with bufB1 and bufC1* in the *E. coli* cytoplasm, the *bufA1* gene was mutated to generate variant precursors with the signal peptide truncated eight residues from the N-terminus (BufA1(-8aa)) or completely removed (BufA1(CP)). Subsequent LC-MS analysis of the purified BufA1 after SUMO cleavage indicated that both truncated precursors were still modified with a 10 Da loss, consistent with the presence of two thiooxazoles and one S-S bond (Fig. 5D). Thus, this data shows that the signal peptide is dispensable for the installation of the PTMs.

Discussion

In this work, we solved a long-standing mystery by unveiling the chemical nature and function of the products derived from heavy metal-induced *gig*-like operons found in various environmental and pathogenic bacteria [11-13]. We showed that the products of *C. vibrioides gig*-like operons are post-translationally modified peptides, bufferins, which protect bacteria from copper stress by chelating excess metal. To the best of our knowledge, bufferins are the first RiPPs directly purified from the producing organism that harbor 5-thiooxazole modifications.

The biosynthesis of bufferins involves several novel features. Firstly, bufferin precursor peptides harbor Sec-dependent signal peptides instead of leader peptides at their N-terminus. Although peptides with bulky modifications or macrocycles would most likely be poor substrates for the Sec machinery due to the pore size of SecY [42], peptides with small modifications could be exported this way if fused to a Sec-dependent signal peptide [43]. To our knowledge, bufferins are the first RiPPs naturally exploiting Sec-dependent signals for efficient translocation to the periplasm. The BufA1 signal peptide is dispensable for the heterocycles installation in our heterologous system of expression, suggesting that determinants for the specific recognition of the precursor peptide by the BufB1C1 modification machinery are located within the core sequence. Secondly, transformation of Cys to 5-thiooxazole is catalyzed

by the MNIO enzyme BufB and its partner protein BufC, and not by a radical-SAM enzyme as was recently reported [33]. Intense efforts have been made recently to explore the chemical space of RiPPs involving MNIOs [15, 16, 19, 21, 44]. To date, members of this superfamily of enzymes are known to catalyze remarkable oxidation reactions, including oxazolone coupled to thioamide formation, excision of cysteine β-methylene, Cys-involved macrocyclization, and C-C and C-N cleavages, as evidenced in the biosynthesis of methanobactins, TglA-type pearlins, RiPPs in *Chryseobacterium*, aminopyruvatides, and methanobactin-like peptides in *Vibrio*, respectively [15, 16, 19-21]. Our present study even further expands the reaction spectra of MNIO enzymes. In line with previous work on Cys-modifying MNIO enzymes, BufB would similarly make use of the mixed-valent iron center and proceed by an intermediate step of proton abstraction from the Cβ of cysteine [17, 18, 20, 21] (Suppl. Fig. S16).

Using model bufferin systems, we demonstrated that the 5-thiooxazoles are key for the copper-protecting function of bufferins. Noteworthy, formation of the bufferin-Cu^{2+} complex was more efficient *in vivo* than *in vitro*. As expression of the bufferin BGC is triggered by the presence of copper in the milieu, the bufferin is likely exposed to copper as soon as it emerges in the periplasm. This can most likely not be faithfully reproduced *in vitro*.

Peptide natural products with other heterocycles are known to bind transition metals and fulfil different functions [45-48]. Most prominently, the RiPP methanobactins scavenge copper from the milieu to feed it to cuproenzymes, and some of them also protect bacteria from toxic metals such as mercury [49]. The defensive role of bufferins is reminiscent of those of the non-ribosomal peptides anthrochelin and yersiniabactin, which can passivate copper by chelation [50, 51]. The fate of copper ions bound by bufferins remains unknown. Periplasmic bufferins might provide specific metallochaperones with copper for delivery to cuproproteins or alternatively facilitate its export. Nevertheless, given the size and the diversity of the bufferin family, it is tempting to speculate that some representatives could bind other transition metals.

Bufferins are part of complex and interwoven metal defense systems in bacteria and are

probably best seen as additional players that buttress active defense systems through copper

passivation under specific environmental conditions. Our data suggest that they are significant

contributors to the copper defense strategy of *C. vibrioides*. The presence of bufferin BGCs in

pathogenic bacteria indicates that they are likely also beneficial to the latter in host-pathogen

interactions, where they might help bacteria deal with copper encountered in phagosomal

compartments [5, 6, 52].

Our bioinformatic analyses indicated that bufferin precursors currently represent the largest

family of MNIO substrates, widespread in different bacterial phyla. The presence of signal

peptides in the precursors and the large sizes of their mature forms relative to most other RiPP

families confer them original properties. The diversity in length and sequence of the bufferin

cores provides bacteria with almost infinite possibilities to build natural peptide products

involved in adaptation to metal or other stresses.

Supporting information

This contains the complete Materials and Methods; diagrams for selected BGCs; Alphafold2

models; LC-MS and MS/MS spectra of BufA1, BufA2 and mutants; growth curves; 1D, HSQC,

HMBC, TOCSY-HSQC, ROESY-HSQC NMR spectra; protein gels; a mechanistic model;

Tables of sequences; EPR parameters; plasmids; strains; oligonucleotides and synthetic genes;

and references for this section.

Acknowledgments. We thank J.-M. Saliou for preliminary MS analyses and J.-Y. Matroule

and P. Cherry for providing *C. vibrioides* strains, plasmids and protocols. This project was

funded by the ANR grant CuRiPP (ANR-22-CE44-0001-02) to FJD. L. Leprevost and S. Jünger

were supported by PhD fellowships of Lille University and the French Ministry of Education

(ED 227 MNHN-SU), respectively. S. Dubiley was supported as a visiting scientist by a

fellowship of the Collège de France. We thank the NMR facility of MetaToul

(www.metatoul.fr). Metatoul is part of the French National Infrastructure for Metabolomics and

Fluxomics MetaboHUB-AR-11-INBS-0010 (www.metabohub.fr), and is supported by the

Région Midi-Pyrénées, the ERDF, the SICOVAL and the French Minister of Education &

Research, who are all gratefully acknowledged. The LC-MS data were acquired at the MNHN

bioorganic mass spectrometry platform and the PAPPSO platform (http://pappso.inra.fr/en).

For the EPR experiments the financial support from the IR INFRANALYTICS FR2054 is

acknowledged. ICP-OES measurements were performed by V. Alaimo on the Chevreul

Institute Platform (U-Lille / CNRS). The Region Hauts de France and the French government

are acknowledged for funding this apparatus.

References

1.   Andrei, A.; Ozturk, Y.; Khalfaoui-Hassani, B.; Rauch, J.; Marckmann, D.; Trasnea, P. I.; Daldal,
F.; Koch, H. G., Cu Homeostasis in Bacteria: The Ins and Outs. *Membranes (Basel)* **2020,** *10* (9).

2.   Solioz, M., *Copper and Bacteria. Evolution, homeostasis and toxicity.* Springer Nature Switzerland AG:
Cham, Switzerland, 2018.

3.   Macomber, L.; Imlay, J. A., The iron-sulfur clusters of dehydratases are primary intracellular targets
of copper toxicity. *Proc Natl Acad Sci U S A* **2009,** *106* (20), 8344-9.

4.   Zuily, L.; Lahrach, N.; Fassler, R.; Genest, O.; Faller, P.; Seneque, O.; Denis, Y.; Castanie-
Cornet, M. P.; Genevaux, P.; Jakob, U.; Reichmann, D.; Giudici-Orticoni, M. T.; Ilbert, M., Copper
Induces Protein Aggregation, a Toxic Process Compensated by Molecular Chaperones. *mBio* **2022**,
e0325121.

5.   Hao, X.; Luthje, F.; Ronn, R.; German, N. A.; Li, X.; Huang, F.; Kisaka, J.; Huffman, D.;
Alwathnani, H. A.; Zhu, Y. G.; Rensing, C., A role for copper in protozoan grazing - two billion years
selecting for bacterial copper resistance. *Mol Microbiol* **2016,** *102* (4), 628-641.

6.   Sheldon, J. R.; Skaar, E. P., Metals as phagocyte antimicrobial effectors. *Curr Opin Immunol* **2019,** *60*,
1-9.

7.   Arguello, J. M.; Raimunda, D.; Padilla-Benavides, T., Mechanisms of copper homeostasis in
bacteria. *Front Cell Infect Microbiol* **2013,** *3*, 73.

8.   Chandrangsu, P.; Rensing, C.; Helmann, J. D., Metal homeostasis and resistance in bacteria. *Nat
Rev Microbiol* **2017,** *15* (6), 338-350.

9.   Hodgkinson, V.; Petris, M. J., Copper homeostasis at the host-pathogen interface. *J Biol Chem* **2012,**
287 (17), 13549-55.

10.   Coutte, L.; Antoine, R.; Slupek, S.; Solans, L.; Derop, J.; Bonnefond, A.; Hot, D.; Locht, C.,
Combined RNAseq and ChIPseq Analyses of the BvgA Virulence Regulator of Bordetella pertussis.
mSystems **2020,** *5* (3).

11.   Jwanoswki, K.; Wells, C.; Bruce, T.; Rutt, J.; Banks, T.; McNealy, T. L., The Legionella
pneumophila GIG operon responds to gold and copper in planktonic and biofilm cultures. *PLoS One* **2017,**
12 (5), e0174245.

12.   Kohler, C.; Lourenco, R. F.; Avelar, G. M.; Gomes, S. L., Extracytoplasmic function (ECF) sigma
factor sigmaF is involved in Caulobacter crescentus response to heavy metal stress. *BMC Microbiol* **2012,** *12,*
210.

13.   Wiesemann, N.; Mohr, J.; Grosse, C.; Herzberg, M.; Hause, G.; Reith, F.; Nies, D. H., Influence
of copper resistance determinants on gold transformation by Cupriavidus metallidurans strain CH34. *J*
Bacteriol **2013,** *195* (10), 2298-308.

14.   Maertens, L.; Cherry, P.; Tilquin, F.; Van Houdt, R.; Matroule, J. Y., Environmental Conditions
Modulate the Transcriptomic Response of Both Caulobacter crescentus Morphotypes to Cu Stress.
Microorganisms **2021,** *9* (6).

15.   Ayikpoe, R. S.; Zhu, L.; Chen, J. Y.; Ting, C. P.; van der Donk, W. A., Macrocyclization and
Backbone Rearrangement During RiPP Biosynthesis by a SAM-Dependent Domain-of-Unknown-Function
692. *ACS Cent Sci* **2023,** *9* (5), 1008-1018.

16.   Chioti, V. T.; Clark, K. A.; Ganley, J. G.; Han, E. J.; Seyedsayamdost, M. R., N-Calpha Bond
Cleavage Catalyzed by a Multinuclear Iron Oxygenase from a Divergent Methanobactin-like RiPP Gene
Cluster. *J Am Chem Soc* **2024.**

17.   Dou, C.; Long, Z.; Li, S.; Zhou, D.; Jin, Y.; Zhang, L.; Zhang, X.; Zheng, Y.; Li, L.; Zhu, X.;
Liu, Z.; He, S.; Yan, W.; Yang, L.; Xiong, J.; Fu, X.; Qi, S.; Ren, H.; Chen, S.; Dai, L.; Wang, B.;
Cheng, W., Crystal structure and catalytic mechanism of the MbnBC holoenzyme required for
methanobactin biosynthesis. *Cell Res* **2022,** *32* (3), 302-314.

18.   Kenney, G. E.; Dassama, L. M. K.; Pandelia, M. E.; Gizzi, A. S.; Martinie, R. J.; Gao, P.; DeHart,
C. J.; Schachner, L. F.; Skinner, O. S.; Ro, S. Y.; Zhu, X.; Sadek, M.; Thomas, P. M.; Almo, S. C.;
Bollinger, J. M., Jr.; Krebs, C.; Kelleher, N. L.; Rosenzweig, A. C., The biosynthesis of methanobactin.
Science **2018,** *359* (6382), 1411-1416.

19.   Nguyen, D. T.; Zhu, L.; Gray, D. L.; Woods, T. J.; Padhi, C.; Flatt, K. M.; MItchell, D. A.; van
der Donk, W. A., Biosynthesis of macrocyclic peptides with C-terminal beta-amino-alpha-keto acid groups
by three different metalloenzymes. *BioRxiv* **2024.**

20.   Park, Y. J.; Jodts, R. J.; Slater, J. W.; Reyes, R. M.; Winton, V. J.; Montaser, R. A.; Thomas, P.
M.; Dowdle, W. B.; Ruiz, A.; Kelleher, N. L.; Bollinger, J. M., Jr.; Krebs, C.; Hoffman, B. M.; Rosenzweig,
A. C., A mixed-valent Fe(II)Fe(III) species converts cysteine to an oxazolone/thioamide pair in
methanobactin biosynthesis. *Proc Natl Acad Sci U S A* **2022,** *119* (13), e2123566119.

21.   Ting, C. P.; Funk, M. A.; Halaby, S. L.; Zhang, Z.; Gonen, T.; van der Donk, W. A., Use of a
scaffold peptide in the biosynthesis of amino acid-derived natural products. *Science* **2019,** *365* (6450), 280-
284.

22.   Arnison, P. G.; Bibb, M. J.; Bierbaum, G.; Bowers, A. A.; Bugni, T. S.; Bulaj, G.; Camarero, J.
A.; Campopiano, D. J.; Challis, G. L.; Clardy, J.; Cotter, P. D.; Craik, D. J.; Dawson, M.; Dittmann, E.;
Donadio, S.; Dorrestein, P. C.; Entian, K. D.; Fischbach, M. A.; Garavelli, J. S.; Goransson, U.; Gruber,
C. W.; Haft, D. H.; Hemscheidt, T. K.; Hertweck, C.; Hill, C.; Horswill, A. R.; Jaspars, M.; Kelly, W. L.;
Klinman, J. P.; Kuipers, O. P.; Link, A. J.; Liu, W.; Marahiel, M. A.; Mitchell, D. A.; Moll, G. N.; Moore,
B. S.; Muller, R.; Nair, S. K.; Nes, I. F.; Norris, G. E.; Olivera, B. M.; Onaka, H.; Patchett, M. L.; Piel,
J.; Reaney, M. J.; Rebuffat, S.; Ross, R. P.; Sahl, H. G.; Schmidt, E. W.; Selsted, M. E.; Severinov, K.;
Shen, B.; Sivonen, K.; Smith, L.; Stein, T.; Sussmuth, R. D.; Tagg, J. R.; Tang, G. L.; Truman, A. W.;
Vederas, J. C.; Walsh, C. T.; Walton, J. D.; Wenzel, S. C.; Willey, J. M.; van der Donk, W. A., Ribosomally
synthesized and post-translationally modified peptide natural products: overview and recommendations for
a universal nomenclature. *Nat Prod Rep* **2013,** *30* (1), 108-60.

23.   Li, Y.; Rebuffat, S., The manifold roles of microbial ribosomal peptide-based natural products in
physiology and ecology. *J Biol Chem* **2020,** *295* (1), 34-54.

24.   Montalban-Lopez, M.; Scott, T. A.; Ramesh, S.; Rahman, I. R.; van Heel, A. J.; Viel, J. H.;
Bandarian, V.; Dittmann, E.; Genilloud, O.; Goto, Y.; Grande Burgos, M. J.; Hill, C.; Kim, S.; Koehnke,
J.; Latham, J. A.; Link, A. J.; Martinez, B.; Nair, S. K.; Nicolet, Y.; Rebuffat, S.; Sahl, H. G.; Sareen, D.;
Schmidt, E. W.; Schmitt, L.; Severinov, K.; Sussmuth, R. D.; Truman, A. W.; Wang, H.; Weng, J. K.;
van Wezel, G. P.; Zhang, Q.; Zhong, J.; Piel, J.; Mitchell, D. A.; Kuipers, O. P.; van der Donk, W. A.,
New developments in RiPP discovery, enzymology and engineering. *Nat Prod Rep* **2021,** *38* (1), 130-239.

25.   Burkhart, B. J.; Hudson, G. A.; Dunbar, K. L.; Mitchell, D. A., A prevalent peptide-binding domain
guides ribosomal natural product biosynthesis. *Nat Chem Biol* **2015,** *11* (8), 564-70.

26.   Chekan, J. R.; Ongpipattanakul, C.; Nair, S. K., Steric complementarity directs sequence
promiscuous leader binding in RiPP biosynthesis. *Proc Natl Acad Sci U S A* **2019,** *116* (48), 24049-24055.

27.   Kloosterman, A. M.; Shelton, K. E.; van Wezel, G. P.; Medema, M. H.; Mitchell, D. A., RRE-
Finder: a Genome-Mining Tool for Class-Independent RiPP Discovery. *mSystems* **2020,** *5* (5).

28.   Severi, E.; Thomas, G. H., Antibiotic export: transporters involved in the final step of natural
product production. *Microbiology (Reading)* **2019,** *165* (8), 805-818.

29.   Müller, F. H.; Bandeiras, T. M.; Urich, T.; Teixeira, M.; Gomes, C. M.; Kletzin, A., Coupling of
the pathway of sulphur oxidation to dioxygen reduction: characterization of a novel membrane-bound
thiosulphate:quinone oxidoreductase. *Mol Microbiol* **2004,** *53* (4), 1147-60.

30.   Nambi, S.; Long, J. E.; Mishra, B. B.; Baker, R.; Murphy, K. C.; Olive, A. J.; Nguyen, H. P.;
Shaffer, S. A.; Sassetti, C. M., The Oxidative Stress Network of Mycobacterium tuberculosis Reveals
Coordination between Radical Detoxification Systems. *Cell Host Microbe* **2015,** *17* (6), 829-37.

31.   Alvarez-Martinez, C. E.; Baldini, R. L.; Gomes, S. L., A caulobacter crescentus extracytoplasmic
function sigma factor mediating the response to oxidative stress in stationary phase. *J Bacteriol* **2006,** *188* (5),
1835-46.

32.   Prunier, G.; Cherkaoui, M.; Lysiak, A.; Langella, O.; Blein-Nicolas, M.; Lollier, V.; Benoist, E.;
Jean, G.; Fertin, G.; Rogniaux, H.; Tessier, D., Fast alignment of mass spectra in large proteomics datasets,
capturing dissimilarities arising from multiple complex modifications of peptides. *BMC Bioinformatics* **2023,**
24 (1), 421.

33.   Lewis, J. K.; Jochimsen, A. S.; Lefave, S. J.; Young, A. P.; Kincannon, W. M.; Roberts, A. G.;
Kieber-Emmons, M. T.; Bandarian, V., New Role for Radical SAM Enzymes in the Biosynthesis of
Thio(seleno)oxazole RiPP Natural Products. *Biochemistry* **2021,** *60* (45), 3347-3361.

34.   Wüthrich, K., *NMR of proteins and nucleic acids.* Wiley: New York, 1986.

35.   Wishart, D. S.; Bigam, C. G.; Holm, A.; Hodges, R. S.; Sykes, B. D., 1H, 13C and 15N random
coil NMR chemical shifts of the common amino acids. I. Investigations of nearest-neighbor effects. *J Biomol*
NMR **1995,** *5* (1), 67-81.

36.   Sattler, M., Heteronuclear multidimensional NMR experiments for the structure determination of
proteins in solution employing pulsed field gradients. *Progr. Nucl. Magn. Reson. Spectr.* **1999,** *34*, 93-158.

37.   Yamazaki, T.; Lee, W.; Arrowsmith, C. H.; Muhandiram, D. R.; Kay, L. E., A Suite of Triple
Resonance NMR Experiments for the Backbone Assignment of 15N, 13C, 2H Labeled Proteins with High
Sensitivity. *J. Am. Chem. Soc.* **1994,** *116*, 11655-11666.

38.   Fiore, K. E.; Patist, M. J.; Giannakoulias, S.; Huang, C. H.; Verma, H.; Khatri, B.; Cheng, R. P.;
Chatterjee, J.; Petersson, E. J., Structural impact of thioamide incorporation into a beta-hairpin. *RSC Chem*
Biol **2022,** *3* (5), 582-591.

39.   Wray, V., Progress in NMR Spectroscopy. 1979; Vol. 13, pp 177-256.

40.   Kenney, G. E.; Rosenzweig, A. C., Chalkophores. *Annu Rev Biochem* **2018,** *87*, 645-676.

41.   Bennett, B.; Kowalski, J. M., EPR Methods for Biological Cu(II): L-Band CW and NARS. *Methods*
Enzymol **2015,** *563,* 341-61.

42.   Bonardi, F.; Halza, E.; Walko, M.; Du Plessis, F.; Nouwen, N.; Feringa, B. L.; Driessen, A. J.,
Probing the SecYEG translocation pore size with preproteins conjugated with sizable rigid spherical
molecules. *Proc Natl Acad Sci U S A* **2011,** *108* (19), 7775-80.

43.   Kuipers, A.; Wierenga, J.; Rink, R.; Kluskens, L. D.; Driessen, A. J.; Kuipers, O. P.; Moll, G. N.,
Sec-mediated transport of posttranslationally dehydrated peptides in Lactococcus lactis. *Appl Environ*
Microbiol **2006,** *72* (12), 7626-33.

44.   Clark, K. A.; Seyedsayamdost, M. R., Bioinformatic Atlas of Radical SAM Enzyme-Modified RiPP
Natural Products Reveals an Isoleucine-Tryptophan Crosslink. *J Am Chem Soc* **2022,** *144* (39), 17876-17888.

45.   DiSpirito, A. A.; Semrau, J. D.; Murrell, J. C.; Gallagher, W. H.; Dennison, C.; Vuilleumier, S.,
Methanobactin and the Link between Copper and Bacterial Methane Oxidation. *Microbiol Mol Biol Rev* **2016,**
80 (2), 387-409.

46.   Kenney, G. E.; Rosenzweig, A. C., Methanobactins: Maintaining copper homeostasis in
methanotrophs and beyond. *J Biol Chem* **2018,** *293* (13), 4606-4615.

47.   Li, Y.; Ma, Y.; Xia, Y.; Zhang, T.; Sun, S.; Gao, J.; Yao, H.; Wang, H., Discovery and biosynthesis
of tricyclic copper-binding ribosomal peptides containing histidine-to-butyrine crosslinks. *Nat Commun*
2023, *14* (1), 2944.

48.   Semrau, J. D.; DiSpirito, A. A.; Obulisamy, P. K.; Kang-Yun, C. S., Methanobactin from
methanotrophs: genetics, structure, function and potential applications. *FEMS Microbiol Lett* **2020,** *367* (5),
fnaa045.

49.   Baral, B. S.; Bandow, N. L.; Vorobev, A.; Freemeier, B. C.; Bergman, B. H.; Herdendorf, T. J.;
Fuentes, N.; Ellias, L.; Turpin, E.; Semrau, J. D.; DiSpirito, A. A., Mercury binding by methanobactin
from Methylocystis strain SB2. *J Inorg Biochem* **2014,** *141,* 161-169.

50.   Büttner, H.; Hörl, J.; Krabbe, J.; Hertweck, C., Discovery and biosynthesis of anthrochelin, a
growth-promoting metallophore of the human pathogen *Luteibacter anthropi*. *ChemBioChem* **2023,** *24,*
e202300322.

51.   Koh, E. I.; Robinson, A. E.; Bandara, N.; Rogers, B. E.; Henderson, J. P., Copper import in
Escherichia coli by the yersiniabactin metallophore system. *Nat Chem Biol* **2017,** *13* (9), 1016-1021.

52.   Chaturvedi, K. S.; Henderson, J. P., Pathogenic adaptations to host-derived antibacterial copper.
Front Cell Infect Microbiol **2014,** *4,* 3.

537