Main manuscript for

A widespread family of ribosomal peptide metallophores involved in bacterial adaptation to metal stress

Laura Leprevost[#1], Sophie Jünger[#2], Guy Lippens[3], Céline Guillaume[2], Giuseppe Sicoli[4], Lydie Oliveira[5], Alex Rivera-Millot[1&], Gabriel Billon[4], Céline Henry[5], Rudy Antoine[1], Séverine Zirah[2], Svetlana Dubiley[*3], Yanyan Li[*2], Françoise Jacob-Dubuisson[*1]

[1] Univ. Lille, U1018-UMR9017-CIIL-Center for infection and Immunity of Lille, F-59000 Lille, FRANCE

[2] Unit Molecules of Communication and Adaptation of Microorganisms (MCAM), UMR 7245 CNRS, MNHN, 75005 Paris, FRANCE

[3] Toulouse Biotechnology Institute, CNRS/INRAE/INSA/UPS, 31077 Toulouse, FRANCE

[4] Univ. Lille, CNRS, UMR 8516 - LASIRE, Laboratoire de Spectroscopie pour les Interactions, la Réactivité et l'Environnement F-59000 Lille, FRANCE

[5] INRAE-AgroParisTech-Univ. Paris-Saclay, MICALIS, 78352 Jouy-en Josas, FRANCE

[#] contributed equally

[&] current address: INRS-Centre Armand-Frappier Santé Biotechnologie, Bacterial Symbionts Evolution, Laval, Quebec H7V 1B7, Canada

[*] correspondence: svetlana.dubiley@gmail.com; yanyanli@mnhn.fr; francoise.jacob@ibl.cnrs.fr

Author contributions: GL, RA, SD, SZ, YL, FJD designed research; LL, SJ, GL, CG, GS, LO, ARM, GB, CH, RA, SD performed research; LL, SJ, GL, CG, GS, LO, ARM, GB, CH, RA, SZ, SD, YL, FJD analyzed data; LL, SJ, GL, SD, SZ, YL, FJD wrote the paper.

Competing interest statement: The authors declare no competing interests.

Classification: Biological Sciences - Biochemistry - Microbiology

Keywords: RiPP, metallophore, MNIO enzyme

This PDF file includes: Main text and Figures 1 to 5

This article was deposited as a preprint in BioRxiv (BIORXIV/2024/585515)

Abstract

Ribosomally synthesized and post-translationally modified peptides (RiPPs) are a structurally diverse group of natural products that bacteria employ in their survival strategies. Herein, we characterized the structure, the biosynthetic pathway and the mode of action of a new RiPP family called bufferins. With thousands of homologous biosynthetic gene clusters throughout the eubacterial phylogenetic tree, bufferins form by far the largest family of RiPPs modified by multinuclear non-heme iron-dependent oxidases (MNIO, DUF692 family). Using *Caulobacter vibrioides* bufferins as a model, we showed that the conserved Cys residues of their precursors are transformed into 5-thiooxazoles, further expanding the reaction range of MNIO enzymes. This rare modification is installed in conjunction with a partner protein of the DUF2063 family. Bufferin precursors are the first examples of bacterial RiPPs found to feature an N-terminal Sec signal peptide and thus to be exported by the ubiquitous Sec pathway, a new paradigm in the RiPP field. Other original features of bufferins are their large size and protein-like fold, which blurs the line between modified peptides and proteins. We reveal that bufferins are involved in copper homeostasis, and their metal-binding propensity requires the thiooxazole heterocycles. Bufferins enhance bacterial growth under copper stress by sequestering excess metal ions in the periplasm. Our study thus describes a large family of RiPP metallophores and unveils a widespread but overlooked metal homeostasis mechanism in eubacteria likely to be relevant to One-Health strategies.

Significance statement

Copper is both essential and toxic in excess. Bacteria face copper in their environments, notably in phagocytes, hence they have developed several defense mechanisms. We discovered a widespread strategy of protection from copper, through the biosynthesis of natural products that we call bufferins. Bufferins are ribosomally synthesized post-translationally modified peptides

(RiPPs), natural products with key roles in bacterial physiology and ecology. Bufferins enhance

bacterial growth under copper stress by complexing with the metal using thiooxazole

heterocycles that result from enzymatic modification of cysteine residues. With thousands of

homologs throughout the eubacterial phylogenetic tree, bufferins represent a highly prevalent

strategy of adaptation to metal stress. They are larger in size than most RiPPs, expanding the

concept of RiPPs to modified proteins.

Main text

Introduction

Ribosomally synthesized and post-translationally modified peptides (RiPPs) are natural

products with a tremendous variety of structures and biological activities (1-3). They are

synthesized from a gene-encoded precursor peptide, generally composed of an N-terminal

leader sequence and a core sequence where post-translational modifications (PTMs) occur. In

many cases, a pathway-specific RiPP recognition element (RRE) domain interacts with the

leader sequence and mediates recruitment of the modification enzymes (4-6). RRE domains are

thus a hallmark of RiPP biosynthesis. The vast structural diversity of RiPPs is conferred by

virtually unlimited possibilities of chemical transformations. An emerging family of PTM

enzymes is the DUF692 family of multinuclear non-heme iron dependent oxidases (MNIOs)

that catalyze various reactions in the biosynthesis of diverse RiPPs, mostly but not exclusively

on Cys residues (7-13). These reactions include oxazolone coupled to thioamide formation,

excision of cysteine β-methylene, Cys-involved macrocyclization, and C-C and C-N cleavages

(7, 8, 11-13). However, the full spectrum of MNIO activities remains far from being fully

appreciated.

Four-gene *gig* operons (*gig* for *g*old-*i*nduced *g*enes) encoding DUF692-family proteins were

identified in the genomes of several microbial species (*SI* Fig. S1). In *Cupriavidus*

metallidurans, *Legionella pneumophila*, and *Caulobacter vibrioides*, the expression of these

operons is induced under various transition metal stresses that include gold, cadmium, copper,

etc., whereas in *Bordetella pertussis* it is controlled by the master regulator of virulence BvgAS

(14-18). Orthologous operons were notably found in methane-oxidizing bacteria (19).

However, despite genetic evidence for an involvement in the copper-resistant phenotype in *C.*

metallidurans (20), the function of these operons remains elusive. Based on their composition,

we speculated that they are biosynthetic gene clusters (BGCs) whose products could represent

a new RiPP family. Unlike all previously described RiPPs however, these putative precursors

have Sec signal peptides. They are also markedly larger than known MNIO-modified RiPP

precursors. Using genetic, structural, and bioinformatic analyses, we questioned the function,

the mode of action, and the prevalence of *gig*-like operons in bacteria.

We report the characterization of the products of the *C. vibrioide*s *gig*-like BGCs, the first

representative of bufferins, a new family of RiPPs. We revealed that bufferins are small protein

metallophores containing a rare 5-thiooxazole modification of cysteine residues. The joint

action of the DUF692 (MNIO) and DUF2063 proteins is required to install this post-

translational modification, which is new to the MNIO family of enzymes. We showed that the

C. vibrioides bufferins protect bacteria against toxic concentrations of copper by sequestering

the excess metal in the periplasm. Our bioinformatic study suggests that bufferins represent an

overlooked transition metal defense mechanism very common in Hydrobacteria and

Terrabacteria.

Results

Large new RiPP family involving MNIOs. We performed an *in-silico* analysis of the

CCNA_03363-03366 operon of *C. vibrioides*, hereafter named *buf1*, to gain insight into gene

functions. CCNA_03363 encodes a DUF2282-family protein (BufA1), whose members are 80-

120 residues long and harbor four Cys residues (Cys^I to Cys^{IV}) and a predicted Sec signal

peptide (Fig. 1A). We hypothesized that BufA1 is an unusually long precursor peptide whose

translocation across the cytoplasmic membrane would rely on the ubiquitous Sec export

pathway rather than on a specialized export system as is often the case for secreted RiPPs (21).

The AlphaFold2 model predicts that the 68-residue long BufA1 core peptide contains a 2-

stranded β sheet and some α-helical structure (*SI* Fig. S1). It thus folds into a small protein

rather than being a totally unstructured peptide.

The product of CCNA_03364 (BufB1) belongs to the MNIO family, whereas

CCNA_03365 codes for a protein (BufC1) with an uncharacterized N-terminal DUF2063

domain. The predicted structural model of BufC1 revealed that its C-terminal domain is related

to RRE domains (4-6) (*SI* Fig. S1). In light of recent discoveries of MNIOs involved in RiPP

biosynthesis (7-13), we hypothesized that BufB1 and BufC1 would install modifications on Cys

residues of BufA1. Finally, CCNA_03366 encodes a putative DoxX-type oxido-reductase,

BufD1 (22, 23).

Inspection of the *C. vibrioides* genome identified a second operon, the *buf2* BGC

(CCNA_02999-03001) (Fig. 1A) which encodes homologs of BufB1 (CCNA_03000) and

BufC1 (CCNA_02999) but lacks a *bufD1*-like gene. Like BufA1, its putative precursor peptide

BufA2 (CCNA_03001) possesses a Sec signal peptide. Its 61-residue core peptide contains six

Cys residues (Cys^I to Cys^{VI}) (Fig. 1A). Although its predicted 3D structure is very similar to

that of BufA1, this small protein does not have a corresponding profile in the databases of

protein families (*SI* Fig. S1).

To evaluate the abundance of homologous BGCs, we collected >14,000 MNIO protein

sequences from the NCBI non-redundant database. The retrieved sequences are widely

distributed　　　throughout　　　Pseudomonadota,　　　Terrabacteria,　　　Myxococcota,

Fibrobacteres/Chlorobi/Bacterioidetes,　Planctomycetes/Verrucomicrobia/Chlamydia,　and

Acidobacteriota. We analyzed their genomic contexts and collected associated genes coding for putative bufferin-like precursors, i.e., small proteins with Cys residues and a predicted Sec signal peptide. Thousands of these were collected, some of which are markedly larger than typical RiPP precursors and harbor numerous Cys residues (detailed *in silico* analyses will be published separately). In a sequence similarity network (SSN) constructed with these proteins, the largest cluster corresponds to the DUF2282 signature, including *C. vibrioides* BufA1 (Fig. 1B). BufA2 of *C. vibrioides* belongs to the second largest cluster of proteins characterized by six or twelve conserved Cys residues, which we named BUF_6/12Cys (Fig. 1B,C). Of note, BUF_6/12Cys proteins were previously identified in Antarctic soil metagenomes (24). Parallel SSN analyses of the MNIO superfamily (Fig. 1D) showed that those coded in the DUF2282 and BUF_6/12Cys BGCs belong to the most populated sequence cluster, distinct from small clusters of MNIOs involved in the biosynthesis of other RiPP types (7, 9-11, 13). Together our data indicate that bufferins currently form the largest group of MNIO-associated RiPPs.

Bufferins mediate adaptation to copper stress. In *C. vibrioides*, *buf1* and *buf2* BGCs belong to the SigF-controlled, core regulon of copper stress (15, 17, 25). To test if they could be up-regulated by other biologically relevant metals, we generated reporter fusions within *bufA1* and *bufA2* and confirmed that both are induced by high concentrations of Cu^{2+}, but not by Fe^{2+}, Zn^{2+} or Mn^{2+} (Fig. 2A; *SI* Fig. S2).

To determine if bufferins, the putative RiPP products of the *buf* BGCs, contribute to adaptation to copper stress, we prepared *C. vibrioides* deletion mutants that express a single (Cv_{buf1} and Cv_{buf2}) or neither BGC (Cv_{DKO}). Cv_{DKO} grew more slowly than the parental Cv_{WT} strain when $CuSO_4$ was added to the growth medium, unlike Cv_{buf1} or Cv_{buf2} (Fig. 2B). Thus, bufferins enhance growth under copper stress and are functionally redundant. To confirm that a single BGC is sufficient to protect bacteria from excess copper, we performed a gain-of-

function experiment. Expression of *buf1ABCD* in an *Escherichia coli* laboratory strain

enhanced survival of bacteria exposed to a bactericidal concentration of $CuSO_4$ (Fig. 2C).

We then asked whether the bufferins would confer an advantage on *C. vibrioides* in biotic

interactions. Since amoeba notably use copper against their bacterial preys (26), a predation

assay was performed. We attempted to measure survival of *C. vibrioides* after ingestion by

Dictyostelium discoidum, a well-established mode organism for the study of phagocytosis.

However, as these assays were unsuccessful, we tested if the presence of the *buf* BGCs might

slow down bacterial killing. We placed amoebal suspensions onto bacterial lawns, and after

incubation we determined bacterial densities in the contact zones by densitometry scanning.

Larger zones of bacterial clearance were observed for the Cc_{DKO} than for the Cc_{WT} strain,

consistent with a protective role of bufferins in this context (Fig. 2D).

Characterization of bufferin PTMs. To identify the PTMs of bufferins, we engineered an

IPTG-inducible version of SigF to control expression of their BGCs in *C. vibrioides*. The

metabolic profiles of IPTG-treated Cv_{buf1}(pSigF) and Cv_{buf2}(pSigF) were analyzed by liquid

chromatography coupled to mass spectrometry (LC-MS) and compared to that of Cv_{DKO}(pSigF)

to reveal peptide species only present in the *buf*-expressing strains. In Cv_{buf1}(pSigF), a major

species exhibited a mass matching that of BufA1 with the Sec signal peptide cleaved off and a

-10 Da mass shift (Fig. 3A; *SI* Fig. S3 and Table S1), suggesting that the core peptide

accumulates in the periplasm and contains post-translational modifications of one or more

residues. A second form with a -12 Da mass shift was also detected, but not systematically, and

it disappeared upon freeze-thawing of cell extracts. The most stable product with a -10 Da mass

shift is considered the mature peptide product and was named bufferin 1 (Buf1). Similarly,

Cv_{buf2}(pSigF) produced a major peptide product with a -18 Da mass shift, called bufferin 2

(Buf2), and a minor species with a -20 Da mass shift (*SI* Fig. S3 and Table S1).

To test the hypothesis that mature bufferins contain modified Cys residues and disulfide

(S-S) bonds, we performed reduction and alkylation treatment of the peptides. This yielded

species carrying three or four alkylated groups for Buf1 and four, five or six alkylated groups

for Buf2 (*SI* Fig. S4), suggesting that the expected PTMs do not affect Cys residues or result in

alkylation-sensitive groups. MS analyses of trypsin-digested, alkylated bufferins revealed that

Cys^I (Cys12) and Cys^{IV} (Cys52) in Buf1 (numbering according to the core peptide, see Fig. 1)

had mass increments corresponding to alkylation only, indicative of unmodified residues (*SI*

Fig. S5). By contrast, in the central tryptic peptide containing Cys^{II} (Cys23) and Cys^{III} (Cys31),

both Cys residues carry mass increments corresponding to alkylation with additional -4.03 Da

mass shifts (Fig. 3B), as proposed by the untargeted PTM search solution SpecGlobX (27).

To confirm that Cys residues are required for Buf1 function, we constructed strains with

the conserved Cys residues replaced with Ser, separately or in combinations. Consistent with

the proposed modifications of Cys^{II} or Cys^{III}, the substitution of either of them yielded peptides

with mass shifts of -6 Da relative to the calculated mass of the mutated peptide (*SI* Fig. S6). We

tested the ability of the mutated BufA1 to protect *C. vibrioides* from excess copper. All mutant

strains grew more slowly than the parental Cv$_{buf1}$ in the presence of CuSO$_4$ (*SI* Fig. S7),

indicating the importance of modified and unmodified Cys residues for function. Altogether,

the data are consistent with the presence of an S-S bond between Cys^I and Cys^{IV} and Cys^{II} and

Cys^{III} being targets of 4-electron oxidation reactions that generate alkylation-sensitive groups.

The Buf1 congener product with a -12 Da mass shift would correspond to a form with a labile

S-S bond between the modified residues. Similarly, in Buf2, in addition to the unmodified Cys^I

(Cys24) and Cys^{VI} (Cys50) likely forming an S-S bond (*SI* Fig. S1), the central Cys residues

carry -4.03 Da shifts (*SI* Fig. S8).

To facilitate bufferin purification and characterization, we added C-terminal tags to BufA1

(*SI* Table S1). The affinity-purified tagged Buf1 was fully modified (*SI* Fig. S9). To probe if

the modified Cys residues may be part of heterocycles as in other MNIO-modified RiPPs (7,

10), we recorded UV-visible absorption spectra of Buf1 and the Buf1[CysIISer+CysIIISer] variant.

Consistent with the presence of aromatic heterocycles (28), the spectrum of the former showed

a broad absorbance peak extending to 310 nm, unlike that of the latter (Fig. 3C). The spectrum

of recombinant Buf2 showed a similar absorbance maximum around 305 nm (*SI* Fig. S10).

PTM identification by NMR. To increase the yield of bufferin for structural elucidation by

NMR, we co-expressed *bufA1* encoding an N-terminal His$_6$-SUMO-tag with *bufB1* and *bufC1*

in *E. coli*. Purification of modified BufA1 followed by trypsin digestion and preparative HPLC

allowed the preparation of the Cys[II]- and Cys[III]-containing 19-mer peptide at a milligram scale

(*SI* Fig. S11). Its [1]H spectrum indicated two groups of anomalous resonances (*SI* Fig. S12). By

combining homonuclear ROESY and TOCSY experiments (29), they were assigned to the

amide protons of Lys24 and Ala32 that follow the modified Cys[II] and Cys[III] residues,

respectively, and to the Hα protons of Asp22 and Ser30 that precede them (Fig. 3D). Further

analysis of these spectra showed the absence of both Hα and NH protons for the modified Cys

residues. The absence of the latter was verified on a natural abundance [1]H, [15]N HSQC spectrum

(*SI* Fig. S12). These data suggest that their N and Cα atoms are part of heterocycles.

A [1]H, [13]C HSQC experiment showed correlations from the Asp22 and Ser30 Hα atoms to

their own Cα carbons at 48.1 and 52.4 ppm, respectively, 6 ppm higher than their random coil

values (30). The [1]H, [13]C HMBC spectrum connected the same Hα protons to carbon atoms at

161.1 and 159.8 ppm, respectively (*SI* Fig. S12). We produced a uniformly [13]C,[15]N-labeled

modified BufA1 molecule in *E. coli* and used its central tryptic peptide to obtain further carbon

assignments on the basis of triple-resonance NMR experiments (31, 32). In the HNCACB

experiment, we found no resonances that could correspond to upstream neighbors for Lys24 or

Ala32, confirming that the Cα carbons of modified Cys[II] and Cys[III] do not resonate at the typical

50-ppm value expected for Cys residues (Fig. 3E). In a HNCO experiment starting from the

amide groups of Lys24 or Ala32, we found values around 184 ppm for the C=O carbon atoms

of both modified Cys residues (Fig. 3F), far from 175-ppm and 200-ppm values expected for

such atoms in random-coil and thio-amide peptides, respectively (30, 33). By increasing the

resolution and varying the offset of the decoupling pulse during the C=O evolution period, we

determined a 90-Hz coupling constant of both carbonyls with carbon atoms whose chemical

shift is ~120 ppm, consistent with their absence in the HNCACB experiment (Fig. 3G). Both

parameters point to the C=O carbon atoms of the modified Cys residues connected to carbon

atoms in aromatic rings (34).

Comparisons of the chemical shifts assigned on the modified BufA1 peptide with those

found for a 5-thiooxazole-containing peptide (28) show the same 6-ppm upfield shift of the Cα

carbon of the preceding residue, the presence of similar carbon resonances at 120 and 160 ppm,

and the 90-Hz coupling constant of the following carbonyl towards an aromatic ring carbon.

Together with the MS data and the characteristic spectrophotometric spectrum (28), we

conclude that the Buf1 modification implies cyclization of its central Cys residues to form

thiooxazole groups (Fig. 3H).

Copper binding by bufferin 1. Since heterocycles can be involved in copper binding (35), we

interrogated if bufferins can complex with copper, which could be the basis for their protective

effect. As attempts to saturate Buf1 with copper *in vitro* were inefficient (*SI* Fig. S13), we

investigated complex formation *in vivo* by treating cultures of *bufABCD1*-expressing *E. coli*

with $CuSO_4$ for three hours prior to bufferin purification. Native MS analysis showed that

purified Buf1 thus produced was fully loaded with copper (Fig. 4A). Pulsed electron

paramagnetic resonance (EPR) spectra confirmed that Buf1 binds paramagnetic Cu^{2+}, and

spectral simulations indicated coordinating N atom(s) (36) (Fig. 4B; *SI* Tables S2 and S3).

To gain further insight into the binding site, we produced a $^{15}N,^{13}C$-labeled Buf1 sample

and recorded triple resonance spectra to assign its resonances. Simultaneously, we used the ^1H,

^{15}N HSQC plane as a basis for HSQC-NOESY and TOCSY experiments to confirm the

assignments and gain some knowledge on structural aspects of Buf1. The sequential assignment

was interrupted by the modified Cys residues, as expected. Interestingly, amide resonances of

the tryptic peptide and of the same segment in the full-length bufferin do not correspond at all,

suggesting that the full-length sequence imposes structural constraints on the thiooxazole-

containing loop. Part of these likely come from the β sheet spanning two short strands in the

structural model, Thr9-Tyr13 and Trp44-Lys48 (*SI* Fig. S1), that we confirmed experimentally

through the strong Hα(i-1) – HN(i) NOE contacts between residues in these segments and the

cross-strand NOE contacts between the amide protons of Lys45 and Cys12, and of Val47 and

Glu10, respectively (*SI* Fig. S14). PTMs of the Cys residues hence do not interfere with the 3D

structure predicted for the unmodified sequence.

A comparison of the spectra of the bufferin produced with or without copper showed many

peaks unchanged, some experiencing a shift and a signal reduction, and yet others disappearing

altogether (Fig. 4C). The latter most probably concern residues close to the paramagnetic Cu^{2+}

ion. In particular, disappearance of the peaks corresponding to direct neighbors of the

heterocycles, such as Lys24, Ser30 and Ala32, confirms that the thiooxazoles are involved in

Cu^{2+} coordination. Interestingly, signals of Thr9, Trp44, Met46 and Lys48 predicted on the β-

sheet face oriented towards the thiooxazole-containing loop also disappeared. Our data thus

support that the thiooxazole cycles and residues of the β sheet are involved in copper binding.

BufB1 and BufC1 are required for post-translational modification of bufferin. To dissect the

role of individual *buf* genes, we constructed Cv_{buf1} derivatives that lack one or several genes

and tested their growth phenotypes in the presence of $CuSO_4$. All deletion strains except for

that lacking *bufD1* displayed slow-growth phenotypes, showing that BufA1, BufB1 and BufC1,

but not BufD1, are required for the function (Fig. 5A).

Since MNIO enzymes require Fe^{II} and Fe^{III} ions for activity (9, 12), purified BufB1 was

analyzed by inductively coupled plasma - optical emission spectrometry, showing the presence

of two Fe ions. To reveal the role of BufB1 and BufC1 in the bufferin maturation, we

reconstituted the modification reaction *in vitro* using the purified proteins (*SI* Fig. S15). LC-

MS analyses showed that BufA1 was converted to a species with a -8 Da mass shift only when

BufB1 and BufC1 were present (Fig. 5B). Trypsin digestion of the *in vitro* modified BufA1

yielded the 19-residue peptide harboring the thiooxazole groups as confirmed by MS/MS

analyses (Fig. 5C). Together with the gene inactivation data, these experiments established that

BufB1 and BufC1 are required and sufficient to install the PTMs on BufA1.

The signal-peptide sequence is dispensable for heterocycle formation. A notable feature of

bufferin biosynthesis is that the precursor peptide harbors a Sec signal peptide, which is

unprecedented in bacterial RiPPs precursors. We investigated if this sequence also acts as a

leader peptide required for Cys transformation. Using the SUMO-*bufA1/bufB1/bufC1 E. coli*

co-expression system, the *bufA1* gene was mutated to generate variant precursors with the signal

peptide truncated eight residues from the N-terminus (BufA1(-8 aa)) or completely removed

(BufA1(CP)). Subsequent LC-MS analysis of purified BufA1 after SUMO-tag cleavage

indicated that both truncated precursors were still modified with a -10 Da shift, consistent with

the presence of two thiooxazoles and one S-S bond (Fig. 5D). Thus, the signal peptide is

dispensable for the installation of the PTMs, suggesting that main recognition determinants of

the precursor peptide by the BufB1C1 machinery are located within the core sequence.

Discussion

Here we report a widespread bacterial strategy for metal homeostasis based on the synthesis of periplasmic RiPP metallophores. Incidentally, we also solved the conundrum of the gold-induced genes found in environmental and pathogenic bacteria (14-16). Bufferins form a new RiPP family with original features and a new function. We showed that they harbor a rare modification, 5-thiooxazole, installed by an MNIO enzyme. With thousands of homologs, bufferins are highly prevalent in eubacteria and represent the largest family of MNIO-modified RiPPs.

The biosynthesis of bufferins involves unique features. To our knowledge they are the first RiPPs naturally exploiting Sec signals for efficient translocation to the periplasm. Peptides with bulky modifications or macrocycles would be poor substrates of the Sec machinery due to the pore size of SecY (37). Although bufferins harbor an S-S bond formed macrocycle, the latter is most likely installed in the oxidizing periplasmic environment after export. The use of 'generalist' systems, Sec and signal-peptidase, for export and proteolytic maturation is a new paradigm in the RiPP field.

The precursors and mature forms of bufferins are large compared to those of other MNIO-modified RiPPs, and our *in-silico* analyses identified yet larger bufferin-like precursors of more than 200 residues with numerous Cys residues. Thus, the bufferin family appears to harbor a continuum from modified peptides to proteins, which expands the concept of *bona fide* RiPPs to what could be 'RiPProteins'.

The conversion of Cys to 5-thiooxazole represents a new reaction catalyzed by MNIOs, further highlighting their chemical versatility. Intense efforts have been made recently to explore the chemical space of RiPPs undergoing MNIO-catalyzed transformations (7, 8, 11, 13, 38). In line with previous work on Cys-modifying MNIO enzymes, BufB would similarly make

use of the mixed-valent iron center and proceed by an intermediate step of proton abstraction

from the Cβ of cysteine (9, 10, 12, 13) (SI Fig. S16).

We demonstrated that model bufferins are involved in copper homeostasis by complexing

with the metal in the periplasm. This novel function for RiPPs contrasts with that of other

MNIO-modified metallophores, methanobactins, a phylogenetically restricted family of RiPPs

which essentially scavenge copper ions in the extracellular milieu for its provision to

cuproprotein clients (39-42). The mode of binding of bufferins also differs from that of

methanobactins. Notably, *in vivo* formation of the bufferin-Cu^{2+} complex most likely occurs

while the unfolded bufferin emerges from the Sec machinery in the periplasm and folds around

copper, which explains that our *in vitro* loading attempts were not successful. The defensive

role of bufferins is reminiscent of those of non-ribosomal peptides, anthrochelin and

yersiniabactin (43, 44), although the putative multiple metal binding sites of large bufferins

suggest that some of them might serve as intracellular metal reservoirs. Future studies on this

large, diverse family will likely also unveil roles for bufferins in the homeostasis of other

transition metals.

Copper is a major metal pollutant from anthropic sources (45, 46), and bufferins represent

a previously unknown but highly prevalent mechanism of bacterial adaptation likely relevant

in the environment, agriculture and healthcare. In *C. vibrioides*, their expression is induced at

high copper concentrations, suggesting that they come into play after other defense systems are

overwhelmed. Genetic determinants that provide environmental species with protection against

transition metals can contribute to the emergence of opportunistic pathogens (47), as

exemplified by *Legionella pneumophila* which harbors up to six bufferin BGCs. Bufferins are

most likely beneficial to bacteria in host-pathogen interactions notably in phagosomal

compartments, which might account for their presence in *Bordetella pertussis, Neisseria*

gonorrheae, Haemophilus influenzae and *Pseudomonas aeruginosa* (26, 48, 49). From a more

applied perspective, it should be possible in the long term to engineer bufferin variants for

bioremediation purposes (3).

Materials and methods

Genetic constructs and growth conditions. Construction details, plasmids, recombinant strains,

primers and synthetic genes are described in *SI* Text and Tables S4 to S6, respectively.

Functional experiments. *C. vibrioides* growth curves were recorded using a TECAN Spark

plate reader, with OD_{600} measurements every 20 min. The gain-of-function experiment was

performed in *E. coli* expressing the *buf*1 operon under the control of an IPTG-inducible

promoter. The bacteria were challenged with 0.3 mM $CuSO_4$ for 4 hours. For the predation

experiments, *Dictyostelium discoidum* was placed in the center of bacterial lawns before

incubation at 20˚C for three days, and the densities of bacteria in the contact zones were

determined with ImageJ. Experimental details are in *SI* Text.

Bufferin production and purification. For Buf1 or Buf2 production in *C. vibrioides*, strains

harboring the pSigF plasmid were grown overnight in PYE medium containing 100 μM IPTG,

and the bufferins were purified on Streptactin or Ni-NTA columns using standard procedures.

Production of isotopically labeled Buf1 was performed in *E. coli* BL21. For the *in vitro* assays,

N-His_6-SUMO-BufA1, N-His_6-BufB1 and N-His_6-BufC1 were produced in BL21(DE3),

purified on Ni-NTA columns, with 10 mM Tris(2-carboxyethyl)phosphine hydrochloride added

to all buffers for BufA1. Production of the trypsic peptide for NMR studies was performed in

BL21(DE3) co-expressing Sumo-BufA1, BufB1 and BufC1. The iron content of His_6-BufB1

was measured using an inductively coupled plasma - optical emission spectrometer (ICP-OES

5110 VDV, Agilent Technologies). Experimental details are provided in *SI* Text.

Peptide analyses by MS. All experimental details are given as *SI* text.

NMR analyses. The central tryptic peptide was purified by preparative HPLC. All NMR spectra were recorded at 293 K on an 800-MHz NEO Bruker spectrometer equipped with a QCP cryogenic probe head. Experimental details are in *SI* Text.

EPR experiments. Pulsed EPR experiments were performed on the Buf1^{twstr}-copper complex using an ELEXSYS E-580 spectrometer (Bruker). Experimental details are in *SI* Text. Numerical simulation of the spectra was conducted using the Matlab toolbox Easyspin 5.2.35.

Spectrophotometric copper binding assay. This was performed as described (50).

Enzymatic assays in vitro. Purified His-SUMO-BufA1 was digested with SUMO protease before adding either or both N-His$_6$-BufB1 and N-His$_6$-BufC1. After incubation the mixtures were analyzed by LC-MSMS. Details are presented as *SI* Text.

In silico analyses. The search for putative bufferin precursors and the SSN analyses are detailed in *SI* Text.

Acknowledgments. We thank E. Lesne and G. Roy for initiating this work long ago, J.-M. Saliou for preliminary MS analyses and J.-Y. Matroule and P. Cherry for providing *C. vibrioides* strains, plasmids and protocols. This project was funded by the ANR grant CuRiPP (ANR-22-CE44-0001-02) to FJD. L. Leprevost and S. Jünger were supported by PhD fellowships of Lille University and the French Ministry of Education (ED 227 MNHN-SU), respectively. S. Dubiley was supported as a Visiting Scientist by a fellowship of the Collège de France. We thank the NMR facility of MetaToul (www.metatoul.fr). Metatoul is part of the French National Infrastructure for Metabolomics and Fluxomics MetaboHUB-AR-11-INBS-0010 (www.metabohub.fr), and is supported by the Région Midi-Pyrénées, the ERDF, the SICOVAL and the French Minister of Education & Research, who are all gratefully acknowledged. The LC-MS data were acquired at the MNHN bioorganic mass spectrometry platform and the PAPPSO platform (http://pappso.inra.fr/en). For the EPR experiments the

financial support from the IR INFRANALYTICS FR2054 is acknowledged. ICP-OES

measurements were performed by V. Alaimo on the Chevreul Institute Platform (U-Lille /

CNRS). The Region Hauts de France and the French government are acknowledged for funding

this apparatus.

References

1.   P. G. Arnison *et al.*, Ribosomally synthesized and post-translationally modified peptide natural
products: overview and recommendations for a universal nomenclature. *Nat Prod Rep* **30**, 108-160
(2013).

2.   Y. Li, S. Rebuffat, The manifold roles of microbial ribosomal peptide-based natural products in
physiology and ecology. *J Biol Chem* **295**, 34-54 (2020).

3.   M. Montalban-Lopez *et al.*, New developments in RiPP discovery, enzymology and engineering.
Nat Prod Rep **38**, 130-239 (2021).

4.   B. J. Burkhart, G. A. Hudson, K. L. Dunbar, D. A. Mitchell, A prevalent peptide-binding domain
guides ribosomal natural product biosynthesis. *Nat Chem Biol* **11**, 564-570 (2015).

5.   J. R. Chekan, C. Ongpipattanakul, S. K. Nair, Steric complementarity directs sequence promiscuous
leader binding in RiPP biosynthesis. *Proc Natl Acad Sci U S A* **116**, 24049-24055 (2019).

6.   A. M. Kloosterman, K. E. Shelton, G. P. van Wezel, M. H. Medema, D. A. Mitchell, RRE-Finder:
a Genome-Mining Tool for Class-Independent RiPP Discovery. *mSystems* **5** (2020).

7.   R. S. Ayikpoe, L. Zhu, J. Y. Chen, C. P. Ting, W. A. van der Donk, Macrocyclization and Backbone
Rearrangement During RiPP Biosynthesis by a SAM-Dependent Domain-of-Unknown-Function
692. *ACS Cent Sci* **9**, 1008-1018 (2023).

8.   V. T. Chioti, K. A. Clark, J. G. Ganley, E. J. Han, M. R. Seyedsayamdost, N-Calpha Bond Cleavage
Catalyzed by a Multinuclear Iron Oxygenase from a Divergent Methanobactin-like RiPP Gene
Cluster. *J Am Chem Soc* 10.1021/jacs.3c11740 (2024).

9.   C. Dou *et al.*, Crystal structure and catalytic mechanism of the MbnBC holoenzyme required for
methanobactin biosynthesis. *Cell Res* **32**, 302-314 (2022).

10. G. E. Kenney *et al.*, The biosynthesis of methanobactin. *Science* **359**, 1411-1416 (2018).

11. D. T. Nguyen *et al.*, Biosynthesis of macrocyclic peptides with C-terminal beta-amino-alpha-keto
acid groups by three different metalloenzymes. *Acs Central Science* 10.1021/acscentralsci.4c00088
(2024).

12. Y. J. Park *et al.*, A mixed-valent Fe(II)Fe(III) species converts cysteine to an oxazolone/thioamide
pair in methanobactin biosynthesis. *Proc Natl Acad Sci U S A* **119**, e2123566119 (2022).

13. C. P. Ting *et al.*, Use of a scaffold peptide in the biosynthesis of amino acid-derived natural products.
Science **365**, 280-284 (2019).

14. K. Jwanoswki *et al.*, The Legionella pneumophila GIG operon responds to gold and copper in
planktonic and biofilm cultures. *PLoS One* **12**, e0174245 (2017).

15. C. Kohler, R. F. Lourenco, G. M. Avelar, S. L. Gomes, Extracytoplasmic function (ECF) sigma
factor sigmaF is involved in Caulobacter crescentus response to heavy metal stress. *BMC Microbiol*
12, 210 (2012).

16.   N. Wiesemann *et al.*, Influence of copper resistance determinants on gold transformation by
449   Cupriavidus metallidurans strain CH34. *J Bacteriol* **195**, 2298-2308 (2013).

450   17.   L. Maertens, P. Cherry, F. Tilquin, R. Van Houdt, J. Y. Matroule, Environmental Conditions
451   Modulate the Transcriptomic Response of Both Caulobacter crescentus Morphotypes to Cu Stress.
452   *Microorganisms* **9** (2021).

453   18.   L. Coutte *et al.*, Combined RNAseq and ChIPseq Analyses of the BvgA Virulence Regulator of
454   Bordetella pertussis. *mSystems* **5** (2020).

455   19.   A. T. Tveit *et al.*, Widespread soil bacterium that oxidizes atmospheric methane. *Proc Natl Acad Sci*
456   *U S A* **116**, 8515-8524 (2019).

457   20.   N. Hirth *et al.*, Full Copper Resistance in Cupriavidus metallidurans Requires the Interplay of Many
458   Resistance Systems. *Appl Environ Microbiol* **89**, e0056723 (2023).

459   21.   E. Severi, G. H. Thomas, Antibiotic export: transporters involved in the final step of natural
460   product production. *Microbiology (Reading)* **165**, 805-818 (2019).

461   22.   F. H. Müller *et al.*, Coupling of the pathway of sulphur oxidation to dioxygen reduction:
462   characterization of a novel membrane-bound thiosulphate:quinone oxidoreductase. *Mol Microbiol*
463   **53**, 1147-1160 (2004).

464   23.   S. Nambi *et al.*, The Oxidative Stress Network of Mycobacterium tuberculosis Reveals Coordination
465   between Radical Detoxification Systems. *Cell Host Microbe* **17**, 829-837 (2015).

466   24.   V. Waschulin *et al.*, Biosynthetic potential of uncultured Antarctic soil bacteria revealed through
467   long-read metagenomic sequencing. *ISME J* **16**, 101-111 (2022).

468   25.   C. E. Alvarez-Martinez, R. L. Baldini, S. L. Gomes, A Caulobacter crescentus extracytoplasmic
469   function sigma factor mediating the response to oxidative stress in stationary phase. *J Bacteriol* **188**,
470   1835-1846 (2006).

471   26.   X. Hao *et al.*, A role for copper in protozoan grazing - two billion years selecting for bacterial copper
472   resistance. *Mol Microbiol* **102**, 628-641 (2016).

473   27.   G. Prunier *et al.*, Fast alignment of mass spectra in large proteomics datasets, capturing
474   dissimilarities arising from multiple complex modifications of peptides. *BMC Bioinformatics* **24**, 421
475   (2023).

476   28.   J. K. Lewis *et al.*, New Role for Radical SAM Enzymes in the Biosynthesis of Thio(seleno)oxazole
477   RiPP Natural Products. *Biochemistry* **60**, 3347-3361 (2021).

478   29.   K. Wüthrich, *NMR of proteins and nucleic acids* (Wiley, New York, 1986).

479   30.   D. S. Wishart, C. G. Bigam, A. Holm, R. S. Hodges, B. D. Sykes, 1H, 13C and 15N random coil
480   NMR chemical shifts of the common amino acids. I. Investigations of nearest-neighbor effects. *J*
481   *Biomol NMR* **5**, 67-81 (1995).

482   31.   M. Sattler, Heteronuclear multidimensional NMR experiments for the structure determination of
483   proteins in solution employing pulsed field gradients. *Progr. Nucl. Magn. Reson. Spectr.* **34**, 93-158
484   (1999).

485   32.   T. Yamazaki, W. Lee, C. H. Arrowsmith, D. R. Muhandiram, L. E. Kay, A Suite of Triple Resonance
486   NMR Experiments for the Backbone Assignment of 15N, 13C, 2H Labeled Proteins with High
487   Sensitivity. *J. Am. Chem. Soc.* **116**, 11655-11666 (1994).

488   33.   K. E. Fiore *et al.*, Structural impact of thioamide incorporation into a beta-hairpin. *RSC Chem Biol*
489   **3**, 582-591 (2022).

34.   V. Wray, "Progress in NMR Spectroscopy". (1979), vol. 13, pp. 177-256.

35.   G. E. Kenney, A. C. Rosenzweig, Chalkophores. *Annu Rev Biochem* **87**, 645-676 (2018).

36. B. Bennett, J. M. Kowalski, EPR Methods for Biological Cu(II): L-Band CW and NARS. *Methods*
Enzymol **563**, 341-361 (2015).

37. F. Bonardi *et al.*, Probing the SecYEG translocation pore size with preproteins conjugated with
sizable rigid spherical molecules. *Proc Natl Acad Sci U S A* **108**, 7775-7780 (2011).

38. K. A. Clark, M. R. Seyedsayamdost, Bioinformatic Atlas of Radical SAM Enzyme-Modified RiPP
Natural Products Reveals an Isoleucine-Tryptophan Crosslink. *J Am Chem Soc* **144**, 17876-17888
(2022).

39. A. A. DiSpirito *et al.*, Methanobactin and the Link between Copper and Bacterial Methane
Oxidation. *Microbiol Mol Biol Rev* **80**, 387-409 (2016).

40. G. E. Kenney, A. C. Rosenzweig, Methanobactins: Maintaining copper homeostasis in
methanotrophs and beyond. *J Biol Chem* **293**, 4606-4615 (2018).

41. Y. Li *et al.*, Discovery and biosynthesis of tricyclic copper-binding ribosomal peptides containing
histidine-to-butyrine crosslinks. *Nat Commun* **14**, 2944 (2023).

42. J. D. Semrau, A. A. DiSpirito, P. K. Obulisamy, C. S. Kang-Yun, Methanobactin from
methanotrophs: genetics, structure, function and potential applications. *FEMS Microbiol Lett* **367**,
fnaa045 (2020).

43. H. Büttner, J. Hörl, J. Krabbe, C. Hertweck, Discovery and biosynthesis of anthrochelin, a growth-
promoting metallophore of the human pathogen *Luteibacter anthropi*. *ChemBioChem* **24**, e202300322
(2023).

44. E. I. Koh, A. E. Robinson, N. Bandara, B. E. Rogers, J. P. Henderson, Copper import in
Escherichia coli by the yersiniabactin metallophore system. *Nat Chem Biol* **13**, 1016-1021 (2017).

45. B. M. Staehlin, J. G. Gibbons, A. Rokas, T. V. O'Halloran, J. C. Slot, Evolution of a Heavy Metal
Homeostasis/Resistance Island Reflects Increasing Copper Stress in Enterobacteria. *Genome Biol*
Evol **8**, 811-826 (2016).

46. Z. Yu, L. Gunn, P. Wall, S. Fanning, Antimicrobial resistance and its association with tolerance to
heavy metals in agriculture production. *Food Microbiol* **64**, 23-32 (2017).

47. C. Baker-Austin, M. S. Wright, R. Stepanauskas, J. V. McArthur, Co-selection of antibiotic and
metal resistance. *Trends Microbiol* **14**, 176-182 (2006).

48. K. S. Chaturvedi, J. P. Henderson, Pathogenic adaptations to host-derived antibacterial copper.
Front Cell Infect Microbiol **4**, 3 (2014).

49. J. R. Sheldon, E. P. Skaar, Metals as phagocyte antimicrobial effectors. *Curr Opin Immunol* **60**, 1-9
(2019).

50. L. Novoa-Aponte, C. Xu, F. C. Soncini, J. M. Arguello, The Two-Component System CopRS
Maintains Subfemtomolar Levels of Free Copper in the Periplasm of Pseudomonas aeruginosa
Using a Phosphatase-Based Mechanism. *mSphere* **5** (2020).

Figure 1. *In silico* analyses of the bufferin family. A. Gene composition of the *buf1* and *buf2* BGCs and sequences of BufA1 and BufA2 with the Cys residues labeled (numbering according to the core peptides). The signal peptide and core peptide sequences are in pale and dark grey, respectively. In the vicinity of *buf1ABCD* are genes coding for SigF and the anti-sigma factor NrsF that regulate *buf1* and *buf2* expression. **B**. SSN analysis of bufferin-like precursors genetically associated with MNIOs. DUF2282 and BUF_6/12Cys proteins, including BufA1 and BufA2 of *C. vibrioides*, belong to the largest two clusters (red and yellow dots), respectively. Small clusters in orange contain the RiPPs of *Chryseobacterium* (7) that are predicted to harbor signal peptides. **C**. Weblogos for the DUF2282 and BUF_6/12Cys bufferins. **D**. SSN analysis of MNIOs (at 80% identity). MNIOs genetically associated with DUF2282 and BUF_6/12Cys proteins, including BufB1 and BufB2 of *C. vibrioides*, are in red and yellow, respectively. MNIOs associated with PTMS on methanobactins, *Chryseobacterium* RiPPs, TglA-type pearlins and aminopyruvatides are found in small clusters numbered 1 to 4.

Figure 2. Regulation and function of the bufferins. **A**. Reporter assays with *bufA1-lacZ* and *bufA2-lacZ* transcriptional fusions. Bacteria were grown for 16 h with the indicated concentrations of $CuSO_4$, $FeSO_4$, $ZnSO_4$ or $MnCl_2$. Non-parametric, one-way analysis of variance Kruskal-Wallis (two-sided) tests followed by a Dunn's multiple-comparison tests were used to analyze the differences between treated cultures and the control (n=4; * in the upper left graph indicates p=0.047). **B**. Effect of copper on bacterial growth. The Cv_{DKO}, $Cv_{buf1ABCD}$ (*buf1)* and $Cv_{buf2ABC}$ (*buf2*) strains were grown without or with 225 μM $CuSO_4$. **C**. Gain-of-function assay. Expression of *buf1* in *E. coli* BL21(pCA24-buf1A^{str}BCD) was induced by IPTG, and the bacteria were challenged with 0.3 mM $CuSO_4$ for three hours before serial dilutions. BL21(pCA24-psmCA) expressing a non-relevant RiPP BGC was used as a control. **D**. Assay of bacterial lysis by *D. discoidum*. Bacterial densities of the lysis plaques were determined using ImageJ analyses. A non-parametric Mann-Whitney test (two-sided) was used to analyze the differences between the Cv_{WT} strain (n=5) and the Cv_{DKO} strain (n=4; * indicates p = 0.0159). The medians are shown.

Figure 3. Post-translational modifications of bufferin 1. **A.** Top-down LC-MS analyses of cell extracts of $Cv_{buf1}(pSigF)$ compared with $Cv_{DKO}(pSigF)$. Left panel: total ion chromatograms, right panels: deconvoluted mass spectra of the compounds detected for $Cv_{buf1}(pSigF)$. Their Mw correspond to the core peptide (*i.e.*, without signal peptide; calculated monoisotopic Mw=6978.35 Da), with mass shifts of -10 Da ($[M+H]^+$ at *m/z* 6969.32) and -12 Da ($[M+H]^+$ at *m/z* 6967.30). **B.** Bottom-up analysis of Buf1. MS/MS spectrum of the central tryptic peptide ($[M+2H]^{2+}$ at *m/z* 917.36). The two Cys residues carry carbamidomethyl groups resulting from alkylation with iodoacetamide (+ 57.02 Da), together with -4.03 Da mass shifts. **C.** UV/vis spectra of affinity-purified Buf1 and the $Cys^{II}Ser+Cys^{III}Ser$ mutant. **D.** NMR ROESY/TOCSY experiments to assign the resonances of the 19-mer peptide. **E.** HNCACB experiment on the uniformly ^{13}C, ^{15}N-labeled peptide: sequential walk for the sequence following Cys^{II}. **F**. The HNCO planes at the ^{15}N frequency of the indicated residues show classical ~175-ppm values for all carbonyl resonances, except for that downstream of a modified Cys, whose carbonyl carbon resonates at 184 ppm. **G**. HNCO planes through the resonance of Lys24 that follows Cys^{II} while modifying the offset of the ^{13}C decoupling pulse during the C=O evolution period. Only when centered at 120 ppm is the 90-Hz carbon-carbon coupling refocused. **H.** Measured NMR parameters and proposed structure for the Buf1 post-translational modifications (shown for Cys^{II}).

Figure 4. Analyses of the bufferin 1-copper complex formed *in vivo*. A. Native MS analysis of the purified complex: isotopic patterns of the major charge state (6+) species. The spectra in black and red show Buf1 from a non-supplemented culture and from cultures supplemented with $CuSO_4$, respectively. The calculated isotopic patterns for $[M+6H]^{6+}$ and $[M+4H+CuII]^{6+}$ species are shown in green. **B.** Evidence for Cu^{2+} binding to Buf1 by EPR spectroscopy. Spectra of Echo-Detected Field-Swept of $CuSO_4$ (red) and of the Buf1-Cu^{2+} complex (black) are shown with the pseudo-modulation of the spectra underneath. The right panels show the Easyspin fits (red) and the experimental spectra (black) for the Buf1-Cu^{2+} complex (top) and for $CuSO_4$ (bottom). **C.** Superposition of the ^1H, ^{15}N HSQC spectra of apo Buf1 (black) and the Buf1-Cu^{2+} complex (red). sc= side chain.

Figure 5. Biosynthesis of bufferin 1. A. Effect of Cu on the growth of *C. vibrioides* strains expressing the indicated genes. **B.** *In vitro* reconstitution of Buf1 biosynthesis. The deconvoluted mass spectra for the products of the *in vitro* reactions are shown, with the added proteins indicated in the corresponding panels. Heat-denatured BufB1 and BufC1 were used for the control reaction. **C.** LC-MS analysis of the 19-residue peptide of *in vitro* modified SUMO-Buf1. The extracted ion chromatogram (EIC) of $[M+3H]^{3+}$ (at *m/z* 573.8945) and the deconvoluted mass are shown. **D**. Role of the signal peptide for the PTMs. A schematic representation of BufA1 variants with signal peptide truncations is shown on top. LC-MS analyses of modified BufA1(-8 aa) (left) and modified BufA1(CP) (right): the MS spectra of the most abundant charge states $[M+9H]^{9+}$ and $[M+8H]^{8+}$, corresponding to deconvoluted monoisotopic masses $[M+H]^+$ of 8391.1281 and 8277.9398 (blue), respectively, are compared to the theoretical spectra of the unmodified peptides (black).