2-oxoglutarate triggers assembly of active dodecameric *Methanosarcina mazei*
glutamine synthetase

Eva Herdering[1], Tristan Reif-Trauttmansdorff[2], Anuj Kumar[2], Tim Habenicht[1],
Georg Hochberg[2, 3, 4], Stefan Bohn[5], Jan Schuller[2], and Ruth A. Schmitz[1]

[1] Institute for General Microbiology, Christian Albrechts University, Am Botanischen Garten 1-9, D-
24118 Kiel, Germany;
[2] Center for Synthetic Microbiology (SYNMIKRO) Research Center and Department of Chemistry,
Philipps-Universität Marburg, Karl-von-Frisch Straße 14, 35043 Marburg
[3] Evolutionary Biochemistry Group, Max Planck Institute for Terrestrial Microbiology, Marburg, Karl-
von-Frisch Straße 10, 35043 Marburg
[4] Department of Chemistry, Philipps-Universität Marburg, Hans-Meerwein-Straße 4, 35043 Marburg
[5] Institute of Structural Biology, Helmholtz Zentrum Munich, Ingolstädter Landstrasse 1, 85764
Neuherberg, Germany
17
18
19 ***** Correspondence: **Ruth Anne Schmitz**
20 rschmitz@ifam.uni-kiel.de, phone: +49-(0)431-880-4334

Abstract

Glutamine synthetases (GS) are central enzymes essential for the nitrogen metabolism across all domains of life. Consequently, they have been extensively studied for more than half a century. Based on the ATP dependent ammonium assimilation generating glutamine, GS expression and activity are strictly regulated in all organisms. In the methanogenic archaeon *Methanosarcina mazei*, it has been shown that the metabolite 2-oxoglutarate (2-OG) directly induces the GS activity. Besides, modulation of the activity by interaction with small proteins (GlnK$_1$ and sP26) has been reported. Here, we show that the strong activation of *M. mazei* GS (GlnA$_1$) by 2-OG is based on the 2-OG dependent dodecamer assembly of GlnA$_1$ by using mass photometry (MP) and single particle cryo-electron microscopy (cryo-EM) analysis of purified strep-tagged GlnA$_1$. The dodecamer assembly from monomers/dimers occurred without any detectable intermediate oligomeric state and was not affected in the presence of GlnK$_1$. The 2.39 Å cryo-EM structure of the dodecameric complex in the presence of 12.5 mM 2-OG demonstrated that 2-OG is binding between two monomers. Thereby, 2-OG appears to induce the dodecameric assembly in a cooperative way. Furthermore, the active site is primed by an allosteric interaction cascade caused by 2-OG-binding towards an adaption of the transition state catalytic conformation. In the presence of additional glutamine, strong feedback inhibition of GS activity was observed. Since glutamine dependent disassembly of the dodecamer was excluded by MP, feedback inhibition most likely relies on an allosteric binding of glutamine to the catalytic site. Based on our findings, we propose that under nitrogen limitation the induction of *M. mazei* GS into a catalytically active dodecamer is not affected by GlnK$_1$ and crucially depends on the presence of 2-OG.

Introduction

Nitrogen is one of the key elements in life and it is essentially required in form of ammonium for biomolecules such as proteins or nucleic acids. Two major pathways of ammonium assimilation in bacteria and archaea are known. Under nitrogen (N) sufficiency glutamate dehydrogenase (GDH) is active and generates glutamate from oxoglutarate and ammonium (reviewed in van Heeswijk et al., 2013). However, under N limitation low ammonium concentrations lead to inactive GDH as a result of its low ammonium affinity, whereas the expression of glutamine synthetase (GS) is strongly induced in response to N limitation (Bolay et al., 2018; Gunka and Commichau, 2012; Stadtman, 2001). Consequently, under low ammonium conditions GS together with glutamate synthase (GOGAT) are responsible for ammonium assimilation via the GS/GOGAT pathway, one of the major intersections in central carbon and N metabolism. Accordingly, GS present across all domains of life plays a central role in cellular N assimilation under low N availability. The enzyme, its structure and regulation has been investigated in detail in different organisms for more than half a century (e.g. Dos Santos Moreira et al., 2019; Stadtman, 2001; Woolfolk and Stadtman, 1967).

Most of the GS are grouped into three major classes based on their monomeric size and oligomerization properties (overview in Dos Santos Moreira et al., 2019). GSI and GSIII, both found in bacteria and archaea mostly form dodecamers, whereas GSII found in Eukaryotes form decamers of smaller subunits (Dos Santos Moreira et al., 2019; He et al., 2009; Valentine et al., 1968; van Rooyen et al., 2011). The GSI class can be further grouped into Iα-type GS and Iβ-type GS based on their amino acid sequence and respective molecular mechanisms of activity regulation. Iß-type GS contain a conserved adenylylation site (Tyr397 residue near the active site) that allows for covalent modification of Iβ-type GS and leads to inactivation of the enzyme (Brown et al., 1994; Magasanik, 1993; Shapiro and Stadtman, 1970), whereas Iα-type GS are not covalently modified and mainly show feedback inhibition by end products of the glutamine metabolism including glutamine (Fisher, 1999; Gunka and Commichau, 2012).

GS regulation on transcriptional level

Since in contrast to GDH, GS catalyzed generation of glutamine requires ATP, most organisms strictly regulate the expression of GS in response to the nitrogen availability on the transcriptional level. In gram negative bacteria mainly transcriptional activation of the coding gene (*glnA*) under low nitrogen availability occurs via a transcriptional activator (e.g. NtrC in *Escherichia coli* (Jiang et al., 1998)). For several gram positive bacteria however, the mechanism of regulation is a de-repression of *glnA* transcription under N limitation, which has also been shown for methanoarchaea (Cohen-Kupiec et al., 1999; Fedorova et al., 2013; Fisher, 1999; Fisher and Wray, 2008; Hauf et al., 2016; Weidenbach et al., 2010, 2008). Whereas in gram positives the signal perception is complex and often also

involves protein interactions of GS with transcriptional regulators (reviewed in Gunka and
Commichau, 2012), signal perception and transduction in methanoarchaea occurs directly via the
small effector molecule 2-oxoglutarate (2-OG), which increases under N limitation. It has been shown
that binding of 2-OG to the global N repressor protein NrpR significantly changes the repressor
conformation resulting in dissociation from its respective operator (Lie et al., 2007; Weidenbach et
al., 2010; Wisedchaisri et al., 2010). In addition to expression regulation the activity of GS is also
strictly regulated in all organisms in response to changing N availabilities, however the underlying
molecular mechanism(s) of inhibition significantly differ for the various GS classes and in various
organisms (Reitzer, 2003)

Regulation of GS activity: highly diverse and often complex in various organisms

An extensive repertoire of cellular control mechanisms regulating GS activity in response to N
availability has been observed in different organisms. Inhibitory mechanisms in response to an N
upshift range from feedback inhibition by e.g. glutamine or other end products of the glutamine
metabolism (e.g. *E. coli* (Stadtman, 2004), *Bacillus subtilis* (Deuel et al., 1970), yeast (Legrain et al.,
1982)), proteolytic degradation (yeast, (Legrain et al., 1982), covalent modification by adenylylation
of the 1ß-type GS subunits (e.g. enterobacteriaceae), thiol-based GS regulation (e.g. in soybean
nodules (Masalkar and Roberts, 2015)), inhibition by regulatory proteins (e.g. in gram positive
bacteria (Travis et al., 2022a)), inhibition by interactions with small proteins (e.g. inhibitory factors in
Cyanobacteria (García-Domínguez et al., 1999; Klähn et al., 2018, 2015)), to directly effecting the
activity through the presence or absence of the small metabolite 2-OG, which has been shown for
the first time for *Methanosarcina mazei* (Ehlers et al., 2005). Moreover, often complex regulations
for GS activity including several of the different regulatory mechanisms are reported for one
organism. For example, yeast GS (ScGS) is regulated via feedback inhibition by glutamine and is
susceptible to proteolytic degradation under N starvation. It was also found to assemble into
nanotubes (He et al., 2009) and under advanced cellular starvation into inactive filaments (Petrovska
et al., 2014). In *E. coli*, the activity of the Iß-type GS (EcGS) is controlled by cumulative feedback
inhibition and covalent modification (reviewed in Reitzer, 2003). It has been shown that each of the
12 subunits can be modified by adenylylation (Tyr397) resulting in an inactivation of the respective
subunit (Stadtman, 1990). Moreover, the adenlylylation of single subunits in the complex in addition
makes the other subunits more susceptible to cumulative feedback inhibition by various substances
(Stadtman, 1990). These substances either bind the glutamine-binding pocket or have an allosteric
binding site (Liaw et al., 1993; Woolfolk and Stadtman, 1967). The dodecameric structure of EcGS has
been known for a long time (Almassy et al., 1986; Yamashita et al., 1989). However, when artificially
exposed to divalent cations (Mn^{2+}, Co^{2+}) it randomly aggregates and produces long hexagonal tubes
(paracrystalline aggregates) (Valentine et al., 1968). The detailed structural information on the

mechanisms of this GS-filament formation to an inactive form of EcGS, often associated with stress

responses, as well as the reversion into individual active dodecamers has only recently been

described by cryo-electron microscopy (cryo-EM) analysis (Huang et al., 2022). The *B. subtilis* GS has

been shown to be feedback regulated. In addition, binding of the transcriptional repressor GlnR to

the feedback inhibited complex not only activates the transcription repression function of GlnR

(Fisher and Wray, 2008) but also stabilizes the inactive GS conformation potentially changing from a

dodecamer into a tetradecameric structure (Travis et al., 2022a).

In *M. mazei,* a mesophilic methanoarchaeon, which is able to fix N_2, regulation of the central N

metabolisms has been studied extensively on the transcriptional and post-transcriptional level (Jäger

et al., 2009; Prasse and Schmitz, 2018; Veit et al., 2005). A central role of 2-OG for the perception of

changes in N availabilities has been proposed as has been demonstrated for cyanobacteria

(Forchhammer, 1999; Herrero et al., 2001). The activity of *M. mazei* GS encoded by *glnA₁*, is

regulated by several different mechanisms. GS /GlnA$_1$) is not covalently modified in response to N

availability and thus represents a Iß-type-GS (Ehlers et al., 2005). It has been demonstrated to get

directly activated under N starvation due to the high intracellular concentrations of the metabolite 2-

OG which directly induces GlnA$_1$ activity (Ehlers et al., 2005). 2-OG represents the internal signal for

N limitation since under N starvation the internal 2-OG level significantly increases due to missing

consumption by GDH (*M. mazei* contains the oxidative TCA part, anabolic). The increased cellular 2-

OG concentration has been shown to be directly perceived by the GlnA$_1$ most likely by direct binding

resulting in strong activation (Ehlers et al., 2005). Besides, we showed first evidence that in addition

two small proteins interact with *M. mazei* GlnA$_1$, the PII-like protein GlnK$_1$ and small protein sP26

comprising 23 amino acids (Ehlers et al., 2005; Gutt et al., 2021). The presence and potential

interaction of both small proteins show small effects on the GlnA$_1$ activity, however compared to the

strong 2-OG stimulation only to a very low extend, which might be neglectable and due to the

indirect GS activity assay. Moreover, based on initial complex formation analysis using size exclusion

chromatography (SEC) and pull-down approaches first indications were obtained that in the absence

of 2-OG the GlnA$_1$/GlnK$_1$ complexes are more stable than in the presence of high 2-OG. This led to

the conclusion that due to the shift to N sufficiency after a period of N limitation, GlnA$_1$ activity is

reduced due to the lower 2-OG concentration but also due to the inhibitory protein interaction with

GlnK$_1$ (Ehlers et al., 2005). Very recently the first structural analysis of *M. mazei* GlnA$_1$ was reported,

showing first GS complexes with GlnK$_1$ (Schumacher et al. 23). Based on their findings Schumacher et

al. propose a regulation of GlnA$_1$ activity by oligomeric modulation, with GlnK$_1$ stabilizing the

dodecameric structure and the formation of GlnA$_1$ active sites. Since this proposed model is entirely

missing the effects of 2-OG on GlnA$_1$ activity, we here aimed to study the obtained effects regulating

M. mazei GlnA$_1$ activity in more detail by evaluating oligomerization and complex formation between

GlnA$_1$, GlnK$_1$ and sP26 in dependence of 2-OG employing mass photometry (MP) allowing molecular

weight distribution of single complexes in solution and by high resolution cryo-EM.

RESULTS

2-OG is responsible for GlnA$_1$-dodecamer formation in M. mazei

The strep-tagged purified GlnA$_1$ was analyzed by SEC in the presence of 12.5 mM 2-OG

demonstrating that GS is exclusively present in a dodecameric structure, no other oligomers were

detectable (suppl. Fig. S1). To investigate the effects of 2-OG on *M. mazei* GlnA$_1$ in more detail, we

employed MP, a method that allows to measure the molecular weight distribution in solution. Strep-

tagged purified GlnA$_1$ (after SEC) was dialyzed into a 2-OG free HEPES buffer (see Materials and

Methods) and subsequently analyzed by MP, demonstrating that in the presence of low 2-OG

concentrations (0.1 mM) all of the *M. mazei* GlnA$_1$ was exclusively present as monomers/dimers with

no higher molecular weight complexes present. After addition of 12.5 mM 2-OG, the size distribution

shifted towards a higher molecular weight complex of 630-700 kDa (calculated based on the

measured dimer-size in each measurement; expected molecular weight of dodecamer: 634 kDa) (Fig.

1A, B). This molecular weight corresponds to a fully assembled dodecamer species, the same

oligomeric structure that is adapted in GS from other prokaryotes. Using 2-OG concentrations varying

between 0.1 and 12.5 mM, complex analysis showed that up to 62 % of all particles were assembled

in a dodecamer. This further allowed to determine the binding affinity of GlnA$_1$ to 2-OG to be K$_D$ =

0.75 ± 0.01 mM 2-OG (based on two biological replicates, calculated with the percentage of

dodecamer) as described in Materials and Methods, and verified that no other intermediate

oligomeric complexes were detectable during dodecameric assembly (Fig. 1A, C, suppl. Fig. S2A, B).

Activity measurements of Strep-GlnA$_1$ in the presence of increasing 2-OG concentrations showed a

strong increase of the activity from 0.0 U/mg in the absence of 2-OG up to 7.8 ± 1.7 U/mg in the

presence of 12.5 mM 2-OG (six independent protein purifications, Fig. 1D). Thus, we conclude that 2-

OG acts as a trigger for dodecameric assembly of *M. mazei* GlnA$_1$, setting it apart from other bacterial

and eukaryotic enzyme variants. Moreover, most likely in addition to the dodecameric assembly, 2-

OG is additionally required for a further 2-OG induced conformational switch of the active site, since

saturated GlnA$_1$ activities are not reached in the presence of 5 mM 2-OG, when most of the GlnA$_1$ is

in a dodecameric structure. For full activity, the presence of 12.5 mM 2-OG is required.

Figure 1

GlnK$_1$ has no detectable effects on GS dodecamer assembly or activity under the tested conditions

Previous studies have shown protein interactions between *M. mazei* $GlnA_1$ and $GlnK_1$ as well as $GlnK_1$

induced effects on $GlnA_1$ activity. Consequently, we next tested the effects of $GlnK_1$ presence on the

$GlnA_1$ oligomerization in the presence of 2-OG. Performing the MP analysis under the tested

conditions as before but in the presence of purified $GlnK_1$ demonstrated that (i) in the absence of 2-

OG varying ratios between $GlnA_1$ and $GlnK_1$ (20:1, 2:1, 2:10 based on monomers) did not result in any

dodecamer assembly of $GlnA_1$ (Fig. 2A, B), (ii) no difference in the $GlnA_1$ dodecameric assembly in the

presence of 2-OG was obtained in the presence of purified $GlnK_1$ (2:1), (iii) nor was binding of $GlnK_1$

to $GlnA_1$ detected by a respective increase in the mass of the higher oligomeric complex (Fig. 2B, C).

Moreover, the presence of $GlnK_1$ (2:1) neither had an influence on the 2-OG affinity (K_D (- $GlnK_1$) =

1.06 mM 2-OG; K_D (+ $GlnK_1$) = 1.02 mM 2-OG, K_D calculated based on the dodecamer/dimer ratio),

nor in any ratio on the specific activity of $GlnA_1$ (Fig. 2 D, C: exemplarily showing 2:1; suppl. Fig. S2C,

D). Consequently, we conclude that under the conditions tested using purified proteins, $GlnA_1$

dodecamer assembly occurs independently of $GlnK_1$ and no binding of $GlnK_1$ to the dodecameric

$GlnA_1$ occurs. However, we cannot exclude that cellular components/metabolites not present in

these experiments are crucial for a $GlnA_1$-$GlnK_1$ interaction.

Figure 2

Structural basis of oligomer formation by 2-OG

To now unravel the structural mechanism underlying *M. mazei* $GlnA_1$ activation by 2-OG, we

employed cryo-EM and single-particle analysis. Treating freshly purified Strep-$GlnA_1$ with 12.5 mM 2-

OG, effectively shifted the equilibrium towards fully assembled homo-oligomers as depicted in the

MP experiments. In the micrographs, fully assembled ring-shaped particles are visible. However,

initial attempts to obtain a 3D reconstruction were hindered by the pronounced preferred

orientation of particles within the ice, a challenge which has been overcome by introducing low

concentrations of CHAPSO (0.7 mM). In our final dataset, all particles exhibited well-distributed

oligomers in diverse orientations. Leveraging this dataset, we aligned the particles to a 2.39 Å

resolution structure, revealing well resolved side chains that facilitated seamless model building (Fig.

3, suppl. Fig. S3, suppl. Tab. S2). Consequently, we achieved a structure demonstrating excellent

geometry and density fitting.

The detailed structural analysis uncovered that $GlnA_1$ assembles into a dodecamer characterized by

stacked hexamer rings. A single $GlnA_1$ protomer is composed of 15 β-strands and 15 α-helices and is

split in into a larger C-domain and an N-domain by helix α3. The dodecameric arrangement is

achieved through two distinct interfaces, the hexamer interfaces and inter-hexamer interfaces.

Hexamer interfaces are situated between subunits within each ring, while inter-hexamer interfaces

occur between subunits derived from adjacent rings (Fig. 4A, B, C). The structures are highly similar

to Gram-positive bacterial GS structures (PDB: 4lnn, Murray et al., 2013),with root mean squared

deviations (rmsds) of 0.5–1.0 Å.

A closer inspection of the density reveals the density for the bound 2-OG at an allosteric site localized

at the interface between two $GlnA_1$ protomers in vicinity of the $GlnA_1$ catalytic site (Fig. 4B, D).

Several residues are contributing to its binding. R172' and S189' coordinate the γ-Carboxy-group.

Additionally, two tightly bound water molecules are detectable in the binding site. One is interacting

with the γ-Carboxy group while being stabilized by another water that is coordinated by S38 and R26.

Latter arginine is coordinating the α-Keto-group and, together with R87 and R173', the α-Carboxy

group of 2-OG (Fig. 4B). Notably, F24 stabilizes the 2-OG via stacking with its phenyl ring. This binding

contribution from two $GlnA_1$ protomers at the intersubunit junction enhances activation by boosting

readiness and the rate of full complex assembly. It operates akin to molecular glue that facilitate the

observed cooperative assembly.

A comparison with the substrate-bound $GlnA_1$ structure (PDB: 8tfk, Schumacher et al. 2023) revealed

that the catalytically important residues in *M. mazei* are the aspartic acid (D57) that abstracts the

proton from ammonium and the catalytic glutamic acid, Glu307. The active site of *M. mazei* $GlnA_1$ is

formed at the interface between two subunits in the hexamer and formed by five key catalytic

elements surrounding the active site: the E flap (residues 303–310), the Y loop (residues 369–377),

the N loop (residues 235–247), the Y^* loop (residues 152–161) and the D50' loop (residues 56-71).

The latter one is the only one that originates from adjacent neighboring protomer (Fig 5C, E).

Superposition of our structure with the apo- *M. mazei* X-ray structure (PDB: 8tfb, Schumacher et al.,

2023) reveals that 2-OG binding also triggers further movements that lead to structural changes in

the substrate binding pocket (Fig. 5A, B, D). R87' and its loop undergo a dramatic flip to coordinate 2-

OG and D170 of helix α3 (residues 167-181) (Fig. 5A). This, combined with the action of other

coordinating residues, initiates a motion that is propagated through the entire protein. Notably, helix

α3 shifts forward, causing F184 to flip over and facilitate a T-shaped aromatic interaction with F202.

The resulting pull on F202 causes F204 to flip, allowing π-stacking with the purine moiety of ATP (Fig.

5B). This series of structural changes primes the active site for ATP binding by already adopting the

side chain conformations that are observed in analogue (Met-Sox-P-ADP)-bound structure (transition

state) (PDB: 8tfk, Schumacher et al., 2023), thus facilitating nucleotide binding (Fig. 5C, E).

Additionally, the D50' loop adopts a position similar to the transition state in a catalytic competent

conformation. This involved a remodeling of the loop, leading to the positioning of key catalytic

residues in a catalytic competent configuration. Compared to the apo structure (Schumacher et al.,

2023), R66 flips out of the catalytic pocket, now accommodating R319 which participates in

phosphoryl transfer catalysis (Fig. 5D). In addition, Asp 57' moves deeper into the binding site,

facilitating the proton abstraction of NH_4^+ and preparing for its attack on the phosphorylated

glutamate. Similar to the ATP/ADP binding site, these catalytic elements are primed to ideally

stabilize the tetrahedral transition state. This is illustrated by the superposition of the inhibitor-

bound, transition-state locked structure (Schumacher et al., 2023) (Fig. 5C, E).

Figure 5

Feedback inhibition by glutamine does not affect the dodecameric M. mazei *$GlnA_1$ structure*

For bacteria it is known, that GS can be feedback inhibited. Very recently, the first feedback inhibition

of an archaeal GS by glutamine has been reported for *Methermicoccus shengliensis* GS (Müller et al.,

2023). The specific arginine residue identified to be relevant for the feedback inhibition is R66.

Consequently, we generated the respective *M. mazei* $GlnA_1$ mutant protein changing the conserved

arginine to alanine (R66A) (see also Fig. 5D, E) and compared the purified strep-tagged mutant

protein with the wildtype (wt) protein. In the presence of 12.5 mM 2-OG, the mutant protein showed

the same specific activity as obtained for the wt. However, when supplementing 5 mM glutamine,

exclusively the wt was strongly feedback inhibited, whereas the R66A mutant protein was not

significantly affected (Fig. 6A). In *B. subtilis*, R62 is responsible for feedback inhibition. The

superposition of the apo-BsGS structure (PDB: 4lnn, Murray et al., 2013) with our 2-OG-bound $GlnA_1$

reveals a similar positioning of the respective *M. mazei* R66 (Fig. 6B) indicating a similar mechanism.

Moreover, we can rule out an effect on the oligomeric structure of $GlnA_1$ by MP analysis, clearly

showing that glutamine does not induce disassembly of the dodecameric wt $GlnA_1$ (Fig. 6C). Instead,

this effect can be explained with the role of R66 being an important residue to bind to glutamine in

the product state of the enzyme.

Figure 6

DISCUSSION

2-OG is crucially required for M. mazei GS assembly to an active dodecamer and induces the
conformational state towards an active open state

In *M. mazei* increased 2-OG concentrations act as central N starvation signal (Ehlers et al., 2005).

Here we demonstrated the importance of 2-OG as the major regulator of *M. mazei* $GlnA_1$ activity by

using independent methods, MP and cryo-EM, to detect and structurally characterize single
complexes with high resolution and quantify the different oligomeric complexes. We have found
mono- and dimeric $GlnA_1$ (apo $GlnA_1$) to be inactive and crucially require 2-OG to form an active
dodecameric complex. Moreover, this dodecameric conformation is the only active state of $GlnA_1$. In
the first step, 2-OG assembles the dodecamer by binding at the interface of two subunits (Fig. 4B)
and functions as a molecular glue between neighbouring subunits. The assembly upon 2-OG addition
observed using MP appears to be cooperative, fast and without any detectable intermediate states
(Fig. 1B, C). Only immediately after thawing a frozen purified $GlnA_1$ preparation and in case that no
additional SEC was performed prior to MP analysis, samples showed additional octameric complexes
in MP with low abundance (suppl. Fig. S4). However, octameric complexes were never observed in
cryo-EM or detected by SEC analysis of frozen purified $GlnA_1$ samples. Consequently, octamers are
most likely broken or disassembled $GlnA_1$-dodecamers or dead-ends in assembly with no
physiological function, rather than an incomplete dodecamer during assembly. Thus, our findings are
contrary to the assembly model proposed by Schuhmacher et al. (Schumacher et al., 2023).

As a second step of activation, the allosteric binding of 2-OG causes a series of conformational
changes in $GlnA_1$ protomers, which prime the active site for the transition state and hence catalysis
of the enzyme. This conformational change of the ATP-binding pocket of the dodecameric $GlnA_1$
upon 2-OG binding goes hand in hand with the observed increased activity at higher 2-OG
concentrations (Fig. 1). Comparing our 2-OG-bound $GlnA_1$ dodecameric structure and the
dodecameric *M. mazei* $GlnA_1$ transition state (PDB: 8tfk) and apo structures (PDB: 8ftb) reported by
Schumacher et al. (Schumacher et al., 2023), clearly demonstrates that 2-OG transfers $GlnA_1$ into its
open transition state conformation (Fig. 5). The conformation of our 2-OG-bound dodecamer
resembled the transition state conformation (ADP-Met-Sox-bound complex) reported by Schumacher
et al., even though in our case no ATP was added (Fig. 5E). A reconfiguration of the active site upon
2-OG-binding has also been reported for GS in *Methanothermococcus thermolithotrophicus* (Müller
et al., 2024). In this report, which does not delineate dodecamer assembly at all, it was demonstrated
that binding of 2-OG in one protomer-protomer interface of a dodecameric GS causes a cooperative
domino effect in the hexameric ring of *M. thermolithotrophicus* GS (Müller et al., 2024). A 2-OG
bound protomer undergoes a conformational change and thereby induces the same shift in its
neighbouring protomer (Müller et al., 2024). This is comparable to our observed cooperativity of *M.*
mazei dodecamer assembly at low 2-OG concentrations (K_D = 0.75 mM, percentage of dodecamer).
On the other hand, *M. mazei* $GlnA_1$ reaches maximal activity only at much higher 2-OG
concentrations and likely requires a fully 2-OG-occupied dodecamer for maximal activity. The here
obtained high activities by 2-OG saturation (up to 9 U/mg) in comparison with previously described
M. mazei $GlnA_1$ activities in the absence of 2-OG in the significantly lower range (mU/mg) (Gutt et

al., 2021; Schumacher et al., 2023) support our conclusion that 2-OG is substantial for the GlnA$_1$

active state.

GlnA$_1$ activity is further regulated by feedback inhibition, small proteins and possibly filament

formation

M. mazei GlnA$_1$ belongs to the group of Iα-type GS, which are known to be feedback inhibited. We

confirmed a strong feedback inhibition by a genetic approach and found R66 to be the key residue

for this inhibition (Fig. 6) as suggested in Müller et al. 2024. The mechanism of feedback inhibition

has been described in detail for *B. subtilis* GS (Murray et al., 2013). There, R62 plays the central role

by binding glutamine and inducing a well ordered inactive structure at the substrate-binding pocket

upon glutamine-binding (Murray et al., 2013). The homologous *M. mazei* R66 likely conveys a similar

way of inhibition to *B. subtilis* GS (Fig. 6B, alignment in suppl. Fig. S5).

Further regulations by the two small proteins sP26 and the PII-like protein GlnK$_1$ have previously

been reported for *M. mazei* (Ehlers et al., 2005; Gutt et al., 2021; Schumacher et al., 2023). However,

in the present study neither an interaction with GlnK$_1$, nor GlnK$_1$ effects on GlnA$_1$ complex formation

analysed by MP, nor an effect of GlnK$_1$ on GlnA$_1$ activity was detectable under the conditions used at

varying 2-OG concentrations (0.1 to 12.5 mM) and ratios of GlnK$_1$ to GlnA$_1$ (20:1, 2:1, 2:10) (Fig. 2).

Moreover, the addition of GlnK$_1$ did not result in a change of the K$_D$ for 2-OG for the dodecamer

GlnA$_1$ assembly (Fig. 2D). In previous reports, GlnK$_1$ was shown to interact with GlnA$_1$ in vivo after a

nitrogen upshift by pull-down approaches (Ehlers et al., 2005), pointing towards an inhibitory

function of GlnK$_1$ under shifting conditions from N limitation to N sufficiency. Similarly, we could not

determine a cryo-EM structure including sP26 despite adding large excess of the small protein either

obtained by co-expression or by addition of a synthetic peptide. Because these attempts were

unsuccessful, we speculate that yet unknown cellular factor(s) might be required for an interaction of

GlnA$_1$ with both small proteins, GlnK$_1$ and sP26, which however is difficult to simulate under in vitro

conditions with purified proteins. Taken this into account, we speculate about a potential function of

the two small proteins beyond GlnA$_1$ inactivation or activation. Since the GlnA$_1$ reaction is coupled to

the GOGAT reaction (GS/GOGAT system) and the products of the two reactions replenish the

substrates for one other, it is tempting to speculate that GlnA$_1$ and GOGAT experience metabolic

coupling by sP26 and/or GlnK$_1$ e.g. by being involved in recruiting or separating GOGAT from GlnA$_1$.

Finally, higher oligomeric states of GS enzymes have been known for a long time for organisms like

yeast and *E. coli* (He et al., 2009; Huang et al., 2022; Petrovska et al., 2014; Valentine et al., 1968).

Interestingly, dependent on the ice thickness and on higher concentrated areas of the grids, we could

also observe filament-like structures of *M. mazei* GlnA$_1$ in cryo-EM and resolved their structure at a

resolution of 6.9 Å (suppl. Fig. S6). Such GlnA$_1$ filaments are also detectable in the cryo-EM images of

Schumacher et al. 2023 but were not reported. Their interface is much alike the previously reported

E.Coli GS filament structures (Huang et al., 2022). The physiological relevance of filamentation in *M.*

mazei however remains unresolved and raises the question, whether an additional rapid modulation

of $GlnA_1$ activity through higher oligomeric states exists, as described e.g. for yeast GS most

depending on stress conditions (Petrovska et al., 2014).

M. mazei *$GlnA_1$ shows unique properties*

Overall, we have confirmed 2-OG to be the central activator of $GlnA_1$ in *M. mazei,* which assembles

the active dodecamer and induces a conformational switch towards an active open state. Though 2-

OG has previously been reported as an on-switch for (methano)archaeal GS activity (Ehlers et al.,

2005; Müller et al., 2024; Pedro-Roig et al., 2013), the 2-OG-triggered assembly is novel and

described exclusively for *M. mazei* $GlnA_1$. In this respect, the way of GS regulation in *M. mazei* is

unique across all prokaryotic GS studied so far. Neither in cyanobacteria, enterobacteria or *Bacillus* is

2-OG a direct activator, nor is complex (dis-)assembly a mode of regulating GS activity in any other of

these model organisms. This is further supported by the absence of up to three of those four

arginines - coordinating 2-OG in *M. mazei* $GlnA_1$ - in these organisms (suppl. Fig. S5). The

cyanobacterial, enterobacterial and gram positive GS are present in the cell as active dodecamers

(Almassy et al., 1986; Bolay et al., 2018; Deuel et al., 1970). However, these dodecamers are

inactivated upon sudden N sufficiency through very different mechanisms: *Synechocystis* GS is

blocked by small proteins (inhibitory factors), the enterobacterial GS experiences gradual

adenylylation of subunits which abolishes the enzyme activity and *B. subtilis* GS is feedback inhibited

by glutamine and further inhibited by binding of the transcription factor GlnR (Almassy et al., 1986;

Bolay et al., 2018; Klähn et al., 2018, 2015; Stadtman, 2001; Travis et al., 2022b) (see Fig. 7).

The direct 2-OG activation and glutamine feedback inhibition of *M. mazei* GS are two fast, reversible

and very direct ways of reacting towards the changing N status of the cell. We propose that the

direct activation through 2-OG without any required additional protein as it is the case for all other

regulations, is a more simple and direct regulation of GS. Due to the evolutionary placement of

methanoarchaea and haloarchaea, where a direct 2-OG regulation has been found exclusively, this

may represent an ancient regulation.

MATERIALS AND METHODS

Strains and plasmids

For heterologous expression and purification of Strep-tagged $GlnA_1$ (MM_0964), the plasmid

pRS1841 was constructed. The $glnA_1$-sequence along with the sP26-sequence (including start-codon:

ATG) were codon-optimized for *Escherichia coli* expression and commercially synthesized by Eurofins

Genomics on the same plasmid (pRS1728) (Ebersberg, Germany). Polymerase chain reaction (PCR)

was performed using pRS1728 as template and the primers (Eurofins Genomics, Ebersberg,

Germany) GlnAopt_NdeI_for (5'TTTCATATGGTTCAGATGAAAAAATG3') and GlnA1opt_BamHI_rev

(5'TTTGGATCCTTACAGCATGCTCAGATAACGG3'). The resulting $GlnA_1$_opt PCR-product and vector

pRS375 were restricted with NdeI and BamHI (NEB, Schwalbach, Germany); the resulting pRS375

vector fragment and the $GlnA_1$ fragment were ligated resulting in pRS1841. For heterologous

expression of Strep-$GlnA_1$, pRS1841 was transformed in *E. coli* BL21 (DE3) cells (Thermo Fisher

Scientific, Waltham, Massachusetts) following the method of Inoue (Inoue et al., 1990). For

generating the Arg66Ala-mutant, a site-directed mutagenesis was performed. pRS1841 was PCR-

amplified using primers sdm_GlnA_R66A_for (5'ATTGAAGAAAGCGATATGAAACTGGCGC3') and

sdm_GlnA_R66A_rev (5'**CGC**GGTAAAGCCCTGAATGCTGCTACC3') by Phusion High-Fidelity polymerase

(Thermo Fisher Scientific, Waltham, Massachusetts) followed by religation resulting in plasmid

pRS1951. For heterologous expression, pRS1951 was transformed into *E. coli* BL21 (DE3).

In order to co-express sP26 along with Strep-$GlnA_1$, the construct pRS1863 was generated. pRS1728

with the codon-optimized sP26-sequence and pET21a (Novagen, Darmstadt, Germany) were

restricted with NdeI and NotI and the resulting untagged sP26_opt was ligated into the pET21a

backbone yielding pRS1863. pRS1863 was co-transformed with pRS1841 into *E. coli* BL21 (DE3) cells

(Thermo Fisher Scientific, Waltham, Massachusetts) selecting for both Kanamycin (pRS1841 derived)

and Ampicillin (pRS1863 derived) resistance.

The plasmid pRS1672 was constructed for producing untagged $GlnK_1$. The $GlnK_1$ gene was PCR-

amplified using primers GlnK1_MM0732.for (5'ATGGTTGGCTATGAAATACGTAATTG3') and

GlnK1_MM0732.rev (5'TCAAATTGCCTCAGGTCCG3') and cloned into pETSUMO by using the

Champion™ pET SUMO Expression System (Thermo Fisher Scientific, Waltham, Massachusetts)

according to the manufacturer's protocol. pRS1672 was then transformed into *E. coli* DH5α and BL21

(DE3) pRIL (suppl. Tab. S1).

Heterologous expression and protein purification: Strep-$GlnA_1$ and $GlnK_1$

Heterologous expression of Strep-$GlnA_1$-variants (pRS1841 and pRS1951) and Strep-$GlnA_1$-sP26-

coexpression (pRS1841 + pRS1863) were performed in 1 l Luria Bertani medium (LB,

Carl Roth GmbH + Co. KG, Karlsruhe, Germany). *E. coli* BL21 (DE3) containing pRS1841, pRS1841 and

pRS1863 or pRS1951 was grown to an optical turbidity at 600 nm (T_{600}) of 0.6 - 0.8, induced with 25

µM isopropylβ-d-1-thiogalactopyranoside (IPTG, Carl Roth GmbH + Co. KG, Karlsruhe, Germany) and

further incubated over night at 18 °C and 120 rpm. The cells were harvested (6,371 x g, 20 min, 4 °C)

and resuspended in 6 ml W-buffer (100 mM TRIS/HCl, 150 mM NaCl, 2.5 mM EDTA, (chemicals from

Carl Roth GmbH + Co. KG, Karlsruhe, Germany), 12.5 mM 2-oxoglutarate (2-OG, Sigma-Aldrich, St.

Louis, Missouri), pH 8.0). After the addition of DNase I (Sigma-Aldrich, St. Louis, Missouri), cell

disruption was performed twice using a French Pressure Cell at 4.135 x 10^6 N/m^2 (*Sim-Aminco*

Spectronic Instruments, Dallas, Texas) followed by centrifugation of the cell lysate for (30 min (13,804

x g, 4 °C). The supernatant was incubated with 1 ml equilibrated (W-buffer) Strep-Tactin sepharose

matrix (IBA, Gottingen, Germany) at 4°C for 1 h at 20 rpm. Strep-tagged GlnA$_1$ was eluted from the

gravity flow column by adding E-buffer (W-buffer + 2.5 mM desthiobiotine (IBA, Gottingen,

Germany)). Strep-GlnA$_1$ was always purified and stored in the presence of 12.5 mM 2-OG, either in E-

buffer or 50 mM HEPES, pH 7.0 at 4 °C for a few days or with 5 % glycerol at -80 °C (chemicals from

Carl Roth GmbH + Co. KG, Karlsruhe, Germany).

His$_6$-SUMO-GlnK$_1$ was expressed similarly using *E. coli* BL21 (DE3) pRIL + pRS1672. Expression was

induced with 100 µM IPTG, incubated at 37 °C, 180 rpm for 2 h and harvested. The pellet was

resuspended in phosphate buffer (50 mM phosphate, 300 mM NaCl, pH 8 (chemicals from

Carl Roth GmbH + Co. KG, Karlsruhe, Germany)) and the cell extract was prepared as described

above. His-tag-affinity chromatography-purifcation was performed with a Ni-NTA agarose (Qiagen,

Hilden, Germany) gravity flow column, the protein was purified by stepwise-elution with 100 and 250

mM imidazole (SERVA, Heidelberg, Deutschland) in phosphate buffer. SUMO-protease (Thermo

Fisher Scientific, Waltham, Massachusetts) was used according to the manufacturer's protocol to

cleave the His$_6$-SUMO-GlnK$_1$ and obtain untagged GlnK$_1$ by passing through the Ni-NTA-column after

the cleavage. Elution fractions of protein purifications were analyzed on 12 % SDS-PAGE gels and the

protein concentrations were determined by Bradford (Bio-Rad Laboratories, Hercules, California) or

Qubit protein assay (Thermo Fisher Sceintific, Waltham, Massachusetts).

Determination of glutamine synthetase activity

The glutamine synthetase activity was determined by performing a coupled optical assay (Shapiro

and Stadtman, 1970). The assay was performed as described in Gutt et al. 2021 with modifications.

Modifcations included the use of 50 mM HEPES, the adjustment of ATP-pH to 7.0 and the use of 5

mM glutamine (Sigma-Aldrich, St. Louis, Missouri) in some assays. The assays were performed with

four technical replicates per condition including two concentrations of GnA$_1$ (2 x 10 µg and 2 x 20 µg

of Strep-GlnA$_1$). Strep-GlnA$_1$ was stored in E-buffer or 50 mM HEPES containing 12.5 mM 2-OG which

was dialysed against 50 mM HEPES pH 7 using Amicon® Ultra catridges with 30 kDa filters

(MilliporeSigma, Burlington, Massachusetts) for the enzyme assays in the absence of 2-OG.

Mass photometry

The molecular weight of protein complexes was analysed by mass photometry (MP) using a Refeyn

twoMP mass photometer with the AcquireMP software (Refeyn Ltd., Oxford, UK). All measurements

were performed in 50 mM HEPES, 150 mM NaCl pH 7.0 (MP-buffer, chemicals from

Carl Roth GmbH + Co. KG, Karlsruhe, Germany) on 1.5 H, 24 x 60 mm microscope coverslips with

Culture Well Reusable Gaskets (GRACE BIO-LABS, Bend, Oregon). Strep-GlnA$_1$ and untagged GlnK$_1$

were prepared as described above. Prior to MP experiments, a size exclusion chromatography (SEC)

was performed with GlnA$_1$ in the presence of 12.5 mM 2-OG on a Superose™ 6 Increase 10/300 GL

column (Cytiva, Marlborough, Massachusetts) with a flow rate of 0.5 ml/min. Only the dodecameric

fraction was used for MP experiments and dialysed against MP buffer using Amicon® Ultra catridges

with 30 kDa filters (MilliporeSigma, Burlington, Massachusetts) beforehand. The Gel Filtration HMW

Calibration Kit (Cytiva, Marlborough, Massachusetts) was used as a standard in SEC. 75 – 200 nM

monomeric Strep-GlnA$_1$ were used in the MP measurements, GlnK$_1$ was added accordingly in the

desired ratio calculated based on monomers. The analysis of the acquired data was performed with

the DiscoverMP software by applying a pre-measured standard (Refeyn Ltd., Oxford, UK). Counts

were visualized in mass histograms as relative counts, which were calculated for the Gaussian fits of

the measured peaks. For the determination of K_D-values and creating sigmoidal fitted curves, RStudio

(RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL) was

used.

Cryo-electron sample preparation and Data collection:

Purified GS at a concentration of 1.5 mg/mL was rapidly applied to glow-discharged Quantifoil grids,

blotted with force 4 for 3.5 s, and vitrified by directly plunging in liquid ethane (cooled by liquid

nitrogen) using Vitrobot Mark IV (Thermo Fisher Scientific, Waltham, Massachusetts) at 100%

humidity and 4 °C. To overcome prefererred orientation bias, 0.7 mM CHAPSO was added to prevent

water-air interface interactions, consequently the concentration of the protein was increased to

6mg/ml. We added purified commercially synthesized sP26 (Davids Biotechnologie, Regensburg,

Germany) to all samples, but the peptide did not stably bind under the observed conditions. Data

was acquired with EPU in EER-format on an FEI Titan Krios G4 (Cryo-EM Platform, Helmholtz Munich)

equipped with a Falcon IVi detector (Thermo Fisher Scientific, Waltham, Massachusetts) with a total

electron dose of ~55 electrons per Å^2 and a pixel size of 0.76 Å. Micrographs were recorded in a

defocus range of -0.25 to -2.0 µm. For details see suppl. Table S2.

Cryo EM - Image processing, classification and refinement

All data was processed using Cryosparc (Punjani et al., 2017). Micrographs were processed on the fly
(motion correction, CTF estimation). Using blob picker, 878,308 particles were picked, 2D-classified
and used for ab initio reconstruction. Iterative rounds of ab initio and heterogenous refinement were
used to clean the particle stacks. The final refinements yielded models with an estimated resolution
of 2.39 Å sets at the 0.143 cutoff (suppl. Fig. S3).

An initial model was generated from the protein sequences using alphaFold (Jumper et al., 2021),
and thereupon fitted as rigid bodies into the density using UCSF Chimera (Pettersen et al., 2021). The
model was manually rebuilt using *Coot* (Emsley et al., 2010). The final model was subjected to real-
space refinements in PHENIX (Liebschner et al., 2019). Illustrations of the models were prepared
using UCSF ChimeraX (Pettersen et al., 2021). The structure is accessible under PDB: 8s59. For details
see suppl. Table S2.

Acknowledgements.

We thank the members of our laboratories for useful discussions on the experiments, as well as
Claudia Kiessling for technical assistance. This work was supported by the German Research Council
(DFG) priority program (SPP) 2002 'Small proteins in Prokaryotes, an unexplored world'
[Schm1052/20-2]. We acknowledge the contribution of the CryoEM Facility of the Philipps University
of Marburg. J.M.S. acknowledges the DFG for an Emmy Noether grant (SCHU 3364/1-1) cofunded by
the European Union (ERC, TwoCO2One, 101075992). Views and opinions expressed are, however,
those of the author(s) only and do not necessarily reflect those of the European Union or the
European Research Council. Neither the European Union nor the granting authority can be held
responsible for them. We thank Sandra Schuller for useful discussions and help in preparing
manuscript figures. GKAH was supported by the Max Planck Society.

REFERENCES

Almassy, R.J., Janson, C.A., Hamlin, R., Xuong, N.-H., Eisenberg, D., 1986. Novel subunit—subunit interactions in the structure of glutamine synthetase. Nature 323, 304–309. https://doi.org/10.1038/323304a0

Bolay, P., Muro-Pastor, M.I., Florencio, F.J., Klähn, S., 2018. The Distinctive Regulation of Cyanobacterial Glutamine Synthetase. Life 8, 52. https://doi.org/10.3390/life8040052

Brown, J.R., Masuchi, Y., Robb, F.T., Doolittlel, W.F., 1994. Evolutionary relationships of bacterial and archaeal glutamine synthetase genes. J. Mol. Evol. 38, 566–576. https://doi.org/10.1007/BF00175876

Cohen-Kupiec, R., Marx, C.J., Leigh, J.A., 1999. Function and regulation of glnA in the methanogenic archaeon Methanococcus maripaludis. J. Bacteriol. 181, 256–261. https://doi.org/10.1128/JB.181.1.256-261.1999

Deuel, T.F., Ginsburg, A., Yeh, J., Shelton, E., Stadtman, E.R., 1970. Bacillus subtilis glutamine synthetase. Purification and physical characterization. J. Biol. Chem. 245, 5195–5205.

Dos Santos Moreira, C.D., Ramos, M.J.R.N., Fernandes, P.M.A.A., 2019. Glutamine synthetase structure-catalysis relationship—Recent advances and applications. WIREs Comput. Mol. Sci. 9, e1399. https://doi.org/10.1002/wcms.1399

Ehlers, C., Weidenbach, K., Veit, K., Forchhammer, K., Schmitz, R.A., 2005. Unique mechanistic features of post-translational regulation of glutamine synthetase activity in *Methanosarcina mazei* strain Gö1 in response to nitrogen availability: Regulation of glutamine synthetase activity in *M. mazei*. Mol. Microbiol. 55, 1841–1854. https://doi.org/10.1111/j.1365-2958.2005.04511.x

Emsley, P., Lohkamp, B., Scott, W.G., Cowtan, K., 2010. Features and development of *Coot*. Acta Crystallogr. D Biol. Crystallogr. 66, 486–501. https://doi.org/10.1107/S0907444910007493

Fedorova, K., Kayumov, A., Woyda, K., Ilinskaja, O., Forchhammer, K., 2013. Transcription factor TnrA inhibits the biosynthetic activity of glutamine synthetase in Bacillus subtilis. FEBS Lett. 587, 1293–1298. https://doi.org/10.1016/j.febslet.2013.03.015

Fisher, S.H., 1999. Regulation of nitrogen metabolism in Bacillus subtilis: vive la différence! Mol. Microbiol. 32, 223–232. https://doi.org/10.1046/j.1365-2958.1999.01333.x

Fisher, S.H., Wray, L.V., 2008. Bacillus subtilis glutamine synthetase regulates its own synthesis by acting as a chaperone to stabilize GlnR-DNA complexes. Proc. Natl. Acad. Sci. U. S. A. 105, 1014–1019. https://doi.org/10.1073/pnas.0709949105

Forchhammer, K., 1999. The PII Protein in Synechococcus PCC 7942 Senses and Signals 2-Oxoglutarate Under ATP-Replete Conditions, in: Peschek, G.A., Löffelhardt, W., Schmetterer, G. (Eds.), The Phototrophic Prokaryotes. Springer US, Boston, MA, pp. 549–553. https://doi.org/10.1007/978-1-4615-4827-0_63

García-Domínguez, M., Reyes, J.C., Florencio, F.J., 1999. Glutamine synthetase inactivation by
protein–protein interaction. Proc. Natl. Acad. Sci. 96, 7161–7166.
https://doi.org/10.1073/pnas.96.13.7161

Gunka, K., Commichau, F.M., 2012. Control of glutamate homeostasis in Bacillus subtilis: a complex
interplay between ammonium assimilation, glutamate biosynthesis and degradation. Mol. Microbiol.
85, 213–224. https://doi.org/10.1111/j.1365-2958.2012.08105.x

Gutt, M., Jordan, B., Weidenbach, K., Gudzuhn, M., Kiessling, C., Cassidy, L., Helbig, A., Tholey, A.,
Pyper, D.J., Kubatova, N., Schwalbe, H., Schmitz, R.A., 2021. High complexity of Glutamine synthetase
regulation in Methanosarcina mazei: Small protein 26 interacts and enhances glutamine synthetase
activity. FEBS J. n/a. https://doi.org/10.1111/febs.15799

Hanahan, D., 1983. Studies on transformation of Escherichia coli with plasmids. J. Mol. Biol. 166,
557–580. https://doi.org/10.1016/s0022-2836(83)80284-8

Hauf, K., Kayumov, A., Gloge, F., Forchhammer, K., 2016. The Molecular Basis of TnrA Control by
Glutamine Synthetase in Bacillus subtilis*. J. Biol. Chem. 291, 3483–3495.
https://doi.org/10.1074/jbc.M115.680991

He, Y.-X., Gui, L., Liu, Y.-Z., Du, Y., Zhou, Y., Li, P., Zhou, C.-Z., 2009. Crystal structure of
Saccharomyces cerevisiae glutamine synthetase Gln1 suggests a nanotube-like supramolecular
assembly. Proteins Struct. Funct. Bioinforma. 76, 249–254. https://doi.org/10.1002/prot.22403

Herrero, A., Muro-Pastor, A.M., Flores, E., 2001. Nitrogen Control in Cyanobacteria. J. Bacteriol. 183,
411–425. https://doi.org/10.1128/jb.183.2.411-425.2001

Ho, C.-M., Li, X., Lai, M., Terwilliger, T.C., Beck, J.R., Wohlschlegel, J., Goldberg, D.E., Fitzpatrick,
A.W.P., Zhou, Z.H., 2020. Bottom-up structural proteomics: cryoEM of protein complexes enriched
from the cellular milieu. Nat. Methods 17, 79–85. https://doi.org/10.1038/s41592-019-0637-y

Huang, P.-C., Chen, S.-K., Chiang, W.-H., Ho, M.-R., Wu, K.-P., 2022. Structural basis for the helical
filament formation of Escherichia coli glutamine synthetase. Protein Sci. 31, e4304.
https://doi.org/10.1002/pro.4304

Inoue, H., Nojima, H., Okayama, H., 1990. High efficiency transformation of Escherichia coli with
plasmids. Gene 96, 23–28. https://doi.org/10.1016/0378-1119(90)90336-p

Jäger, D., Sharma, C.M., Thomsen, J., Ehlers, C., Vogel, J., Schmitz, R.A., 2009. Deep sequencing
analysis of the Methanosarcina mazei Gö1 transcriptome in response to nitrogen availability. Proc.
Natl. Acad. Sci. 106, 21878–21882. https://doi.org/10.1073/pnas.0909051106

Jiang, P., Peliska, J.A., Ninfa, A.J., 1998. Reconstitution of the signal-transduction bicyclic cascade
responsible for the regulation of Ntr gene transcription in Escherichia coli. Biochemistry 37, 12795–
12801. https://doi.org/10.1021/bi9802420

Joo, H.K., Park, Y.W., Jang, Y.Y., Lee, J.Y., 2018. Structural Analysis of Glutamine Synthetase from
Helicobacter pylori. Sci. Rep. 8, 11657. https://doi.org/10.1038/s41598-018-30191-5

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A.,
Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E.,
Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O.,
Senior, A.W., Kavukcuoglu, K., Kohli, P., Hassabis, D., 2021. Highly accurate protein structure
prediction with AlphaFold. Nature 596, 583–589. https://doi.org/10.1038/s41586-021-03819-2

Klähn, S., Bolay, P., Wright, P.R., Atilho, R.M., Brewer, K.I., Hagemann, M., Breaker, R.R., Hess, W.R.,
2018. A glutamine riboswitch is a key element for the regulation of glutamine synthetase in
cyanobacteria. Nucleic Acids Res. https://doi.org/10.1093/nar/gky709

Klähn, S., Schaal, C., Georg, J., Baumgartner, D., Knippen, G., Hagemann, M., Muro-Pastor, A.M.,
Hess, W.R., 2015. The sRNA NsiR4 is involved in nitrogen assimilation control in cyanobacteria by
targeting glutamine synthetase inactivating factor IF7. Proc. Natl. Acad. Sci. 112.
https://doi.org/10.1073/pnas.1508412112

Legrain, C., Vissers, S., Dubois, E., Legrain, M., Wiame, J.-M., 1982. Regulation of Glutamine
Synthetase from Saccharomyces cerevisiae by Repression, Inactivation and Proteolysis. Eur. J.
Biochem. 123, 611–616. https://doi.org/10.1111/j.1432-1033.1982.tb06576.x

Liaw, S.H., Pan, C., Eisenberg, D., 1993. Feedback inhibition of fully unadenylylated glutamine
synthetase from Salmonella typhimurium by glycine, alanine, and serine. Proc. Natl. Acad. Sci. U. S. A.
90, 4996–5000. https://doi.org/10.1073/pnas.90.11.4996

Lie, T.J., Dodsworth, J.A., Nickle, D.C., Leigh, J.A., 2007. Diverse homologues of the archaeal repressor
NrpR function similarly in nitrogen regulation. FEMS Microbiol. Lett. 271, 281–288.
https://doi.org/10.1111/j.1574-6968.2007.00726.x

Liebschner, D., Afonine, P.V., Baker, M.L., Bunkóczi, G., Chen, V.B., Croll, T.I., Hintze, B., Hung, L.-W.,
Jain, S., McCoy, A.J., Moriarty, N.W., Oeffner, R.D., Poon, B.K., Prisant, M.G., Read, R.J., Richardson,
J.S., Richardson, D.C., Sammito, M.D., Sobolev, O.V., Stockwell, D.H., Terwilliger, T.C., Urzhumtsev,
A.G., Videau, L.L., Williams, C.J., Adams, P.D., 2019. Macromolecular structure determination using X-
rays, neutrons and electrons: recent developments in Phenix. Acta Crystallogr. Sect. Struct. Biol. 75,
861–877. https://doi.org/10.1107/S2059798319011471

Magasanik, B., 1993. The regulation of nitrogen utilization in enteric bacteria. J. Cell. Biochem. 51,
34–40. https://doi.org/10.1002/jcb.240510108

Masalkar, P.D., Roberts, D.M., 2015. Glutamine synthetase isoforms in nitrogen-fixing soybean
nodules: distinct oligomeric structures and thiol-based regulation. FEBS Lett. 589, 215–221.
https://doi.org/10.1016/j.febslet.2014.11.048

McMullan, G., Naydenova, K., Mihaylov, D., Yamashita, K., Peet, M.J., Wilson, H., Dickerson, J.L.,
Chen, S., Cannone, G., Lee, Y., Hutchings, K.A., Gittins, O., Sobhy, M.A., Wells, T., El-Gomati, M.M.,
Dalby, J., Meffert, M., Schulze-Briese, C., Henderson, R., Russo, C.J., 2023. Structure determination by
cryoEM at 100 keV. Proc. Natl. Acad. Sci. 120, e2312905120.
https://doi.org/10.1073/pnas.2312905120

Müller, M.-C., Lemaire, O.N., Kurth, J.M., Welte, C.U., Wagner, T., 2024. Differences in regulation
mechanisms of glutamine synthetases from methanogenic archaea unveiled by structural
investigations. Commun. Biol. 7, 1–14. https://doi.org/10.1038/s42003-023-05726-w

Müller, M.-C., Lemaire, O.N., Kurth, J.M., Welte, C.U., Wagner, T., 2023. Differences in the regulation
mechanisms of the glutamine synthetase from methanogenic archaea unveiled by structural
investigations. https://doi.org/10.1101/2023.04.07.535877

Murray, D.S., Chinnam, N., Tonthat, N.K., Whitfill, T., Wray, L.V., Fisher, S.H., Schumacher, M.A.,
2013. Structures of the Bacillus subtilis glutamine synthetase dodecamer reveal large intersubunit
catalytic conformational changes linked to a unique feedback inhibition mechanism. J. Biol. Chem.
288, 35801–35811. https://doi.org/10.1074/jbc.M113.519496

Nilsson, M.T., Krajewski, W.W., Yellagunda, S., Prabhumurthy, S., Chamarahally, G.N., Siddamadappa,
C., Srinivasa, B.R., Yahiaoui, S., Larhed, M., Karlén, A., Jones, T.A., Mowbray, S.L., 2009. Structural
basis for the inhibition of Mycobacterium tuberculosis glutamine synthetase by novel ATP-
competitive inhibitors. J. Mol. Biol. 393, 504–513. https://doi.org/10.1016/j.jmb.2009.08.028

Pedro-Roig, L., Camacho, M., Bonete, M.-J., 2013. Regulation of ammonium assimilation in Haloferax
mediterranei: interaction between glutamine synthetase and two GlnK proteins. Biochim. Biophys.
Acta 1834, 16–23. https://doi.org/10.1016/j.bbapap.2012.10.006

Petrovska, I., Nüske, E., Munder, M.C., Kulasegaran, G., Malinovska, L., Kroschwald, S., Richter, D.,
Fahmy, K., Gibson, K., Verbavatz, J.-M., Alberti, S., 2014. Filament formation by metabolic enzymes is
a specific adaptation to an advanced state of cellular starvation. eLife 3, e02409.
https://doi.org/10.7554/eLife.02409

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H., Ferrin,
T.E., 2021. UCSF ChimeraX: Structure visualization for researchers, educators, and developers.
Protein Sci. Publ. Protein Soc. 30, 70–82. https://doi.org/10.1002/pro.3943

Prasse, D., Schmitz, R.A., 2018. Small RNAs Involved in Regulation of Nitrogen Metabolism. Microbiol.
Spectr. 6. https://doi.org/10.1128/microbiolspec.RWR-0018-2018

Punjani, A., Rubinstein, J.L., Fleet, D.J., Brubaker, M.A., 2017. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. Nat. Methods 14, 290–296.
https://doi.org/10.1038/nmeth.4169

Reitzer, L., 2003. Nitrogen Assimilation and Global Regulation in Escherichia coli. Annu. Rev.
Microbiol. 57, 155–176. https://doi.org/10.1146/annurev.micro.57.030502.090820

Schumacher, M.A., Salinas, R., Travis, B.A., Singh, R.R., Lent, N., 2023. M. mazei glutamine synthetase
and glutamine synthetase-GlnK1 structures reveal enzyme regulation by oligomer modulation. Nat.
Commun. 14, 7375. https://doi.org/10.1038/s41467-023-43243-w

Shapiro, B.M., Stadtman, E.R., 1970. [130] Glutamine synthetase (Escherichia coli), in: Methods in
Enzymology. Elsevier, pp. 910–922. https://doi.org/10.1016/0076-6879(71)17305-3

Stadtman, E.R., 2004. Regulation of Glutamine Synthetase Activity. EcoSal Plus 1,
10.1128/ecosalplus.3.6.1.6. https://doi.org/10.1128/ecosalplus.3.6.1.6

Stadtman, E.R., 2001. The story of glutamine synthetase regulation. J. Biol. Chem. 276, 44357–44364.
https://doi.org/10.1074/jbc.R100055200

Stadtman, E.R., 1990. [60] Discovery of glutamine synthetase cascade, in: Deutscher, M.P. (Ed.),
Methods in Enzymology, Guide to Protein Purification. Academic Press, pp. 793–809.
https://doi.org/10.1016/0076-6879(90)82062-7

Travis, B.A., Peck, J.V., Salinas, R., Dopkins, B., Lent, N., Nguyen, V.D., Borgnia, M.J., Brennan, R.G.,
Schumacher, M.A., 2022a. Molecular dissection of the glutamine synthetase-GlnR nitrogen
regulatory circuitry in Gram-positive bacteria. Nat. Commun. 13, 3793.
https://doi.org/10.1038/s41467-022-31573-0

Travis, B.A., Peck, J.V., Salinas, R., Dopkins, B., Lent, N., Nguyen, V.D., Borgnia, M.J., Brennan, R.G.,
Schumacher, M.A., 2022b. Molecular dissection of the glutamine synthetase-GlnR nitrogen
regulatory circuitry in Gram-positive bacteria. Nat. Commun. 13, 3793.
https://doi.org/10.1038/s41467-022-31573-0

Valentine, R.C., Shapiro, B.M., Stadtman, E.R., 1968. Regulation of glutamine synthetase. XII.
Electron microscopy of the enzyme from Escherichia coli. Biochemistry 7, 2143–2152.
https://doi.org/10.1021/bi00846a017

van Heeswijk, W.C., Westerhoff, H.V., Boogerd, F.C., 2013. Nitrogen Assimilation in Escherichia coli:
Putting Molecular Data into a Systems Perspective. Microbiol. Mol. Biol. Rev. 77, 628–695.
https://doi.org/10.1128/mmbr.00025-13

van Rooyen, J.M., Abratt, V.R., Belrhali, H., Sewell, T., 2011. Crystal structure of Type III glutamine
synthetase: surprising reversal of the inter-ring interface. Struct. Lond. Engl. 1993 19, 471–483.
https://doi.org/10.1016/j.str.2011.02.001

Veit, K., Ehlers, C., Schmitz, R.A., 2005. Effects of nitrogen and carbon sources on transcription of
soluble methyltransferases in Methanosarcina mazei strain Go1. J. Bacteriol. 187, 6147–6154.
https://doi.org/10.1128/JB.187.17.6147-6154.2005

Weidenbach, K., Ehlers, C., Kock, J., Ehrenreich, A., Schmitz, R.A., 2008. Insights into the NrpR regulon
in Methanosarcina mazei Gö1. Arch. Microbiol. 190, 319–332. https://doi.org/10.1007/s00203-008-
0369-3

Weidenbach, K., Ehlers, C., Kock, J., Schmitz, R.A., 2010. NrpRII mediates contacts between NrpRI and
general transcription factors in the archaeon Methanosarcina mazei Gö1. FEBS J. 277, 4398–4411.
https://doi.org/10.1111/j.1742-4658.2010.07821.x

Wisedchaisri, G., Dranow, D.M., Lie, T.J., Bonanno, J.B., Patskovsky, Y., Ozyurt, S.A., Sauder, J.M.,
Almo, S.C., Wasserman, S.R., Burley, S.K., Leigh, J.A., Gonen, T., 2010. Structural Underpinnings of
Nitrogen Regulation by the Prototypical Nitrogen-Responsive Transcriptional Factor NrpR. Structure
18, 1512–1521. https://doi.org/10.1016/j.str.2010.08.014

Woolfolk, C.A., Stadtman, E.R., 1967. Regulation of glutamine synthetase: III. Cumulative feedback
inhibition of glutamine synthetase from Escherichia coli. Arch. Biochem. Biophys. 118, 736–755.
https://doi.org/10.1016/0003-9861(67)90412-2

Yamashita, M., Almassy, R., Janson, C., Cascio, D., Eisenberg, D., 1989. Refined atomic model of
glutamine synthetase at 3.5 Å resolution. J. Biol. Chem. 264, 17681–90.
https://doi.org/10.2210/pdb2gls/pdb

 FIGURE LEGENDS

Figure 1: **GlnA$_1$-dodecamer-assembly is induced by 2-OG without detectable oligomeric**
intermediates. Oligomerisation states of purified strep-tagged GlnA$_1$ were assessed in dependence of
2-OG by mass photometry as described in MM using a Refeyn twoMP mass photometer (Refeyn Ltd.,
Oxford, UK). Mass spectra are shown with relative counts (number of counts in relation to the total
counts) plotted against the molecular weight. **A**: 75 nM GlnA$_1$ were preincubated in the presence of
varying 2-OG concentrations (0 to 25 mM) for ten min at room temperature and kept on ice until
measurement. The percentage of dodecamer considering the total number of counts was plotted
against the 2-OG concentration. One out of two independent biological replicates with each three
technical replicates is shown exemplarily. The molecular masses shown above the peaks correspond
to a Gaussian fit of the respective peak (Gaussian fit not shown) and the K$_D$ is indicated in green. **B**:
Exemplary mass spectra of GlnA$_1$ oligomers in the presence of 0.1 and 12.5 mM 2-OG. **C**: Mass
spectra of the three technical replicates (different green colors) of GlnA$_1$-oligomers at 0.39, 0.78 and
1.56 mM 2-OG, excluding the presence of intermediates. **D**: The specific activity of purified strep-
tagged GlnA$_1$ was determined as described in MM in the presence of varying 2-OG concentrations (0,
1.25, 5 and 12.5 mM). The standard deviation of four technical replicates is indicated for one out of
two biological replicates.

Figure 2: GlnA$_1$-dodecamer-assembly and activity are not influenced by GlnK$_1$ under the conditions
tested. Purified strep-tagged GlnA$_1$ and tag-less GlnK$_1$ were incubated in the absence or presence of
2-OG in varying concentrations for ten min at RT. Oligomerisation states were assessed by mass
photometry. Mass spectra are shown with relative counts (see Fig. 1). **A**: The obtained ratio of GlnA$_1$
dodecamer/dimer of three technical replicates are shown for varying ratios between GlnA$_1$ and GlnK$_1$
(20:1, 2:1, 2:10, ratios relating to monomers) in the absence of 2-OG. **B, C**: Exemplary mass spectra of
GlnA$_1$ incubated in the absence and presence of GlnK$_1$ (2:1) at 2-OG concentrations of 0 mM (B) and
12.5 mM (C). The molecular masses shown above the peaks correspond to a Gaussian fit of the
respective peak (Gaussian fit not shown). **D**: 200 nM monomeric GlnA$_1$ were preincubated with GlnK$_1$
(in a 2:1 ratio) in the presence of varying 2-OG concentrations (0.19 to 12.5 mM) for ten min at RT.
The percentage of GlnA$_1$ dodecamer considering the total number of counts was plotted against the
2-OG concentration. One biological replicate with three technical replicates was performed. The ratio
of GlnA$_1$ dodecamer/dimer was plotted against the 2-OG concentration and the KD is indicated in
green (●, - GlnK$_1$) and yellow (●, + GlnK$_1$). **E**: The specific activity of purified strep-tagged GlnA$_1$ in the
absence and presence of GlnK$_1$ (ratio 2:1) was determined as described in MM in the presence of

varying 2-OG concentrations (0, 0.78, 6.25 and 12.5 mM). The standard deviations of four technical

replicates of one biological replicate are indicated.

Figure 3: Structure of *M. Mazei* GlnA$_1$ with 2-OG. A: Three-dimensional segmented cryo-EM density

of the dodecameric complex colored by subunits. **B**: Corresponding views of the GlnA$_1$ atomic model

in cartoon representation.

Figure 4: Dimeric Interface and 2-OG binding site of dodecameric GlnA$_1$. A: Surface representation

of the *M. mazei* GlnA$_1$ 2-OG dodecamer with three GlnA$_1$ protomers fitted in cartoon representation

into the dodecamer as vertical (blue and ochre) and horizontal (blue and green) dimers. **B**: Horizontal

dimers and close-up of 2-OG binding site. Important residues are shown as atomic stick

representation, primed labels indicate neighboring protomer. 2-OG and water molecules important

for ligand binding fitted into density are shown in grey. Dotted lines represent polar interactions

between 2-OG, waters and residues. **C** Vertical dimers and close-up of dimerization site. C-terminal

helices H14/15 and H14'/ H15' of two neighboring protomers lead to tight interaction, mediated by

hydrophobic and polar interactions. **D**: Top-view of GlnA$_1$ hexamer, 2-OG and substrate binding sites

are depicted for one horizontal dimer.

Figure 5: Comparison of 2-OG and substrate binding site of 2-OG bound, apo and TS structures

(Schumacher et al., 2023)**.** Atomic models in cartoon, important residues shown in stick

representation. Colors: blue/green, purple/ochre and red/yellow represent *M. mazei* GlnA$_1$ 2-OG, *M.*

mazei GlnA$_1$ apo (PDB: 8tfb, Schumacher et al., 2023) and *M. mazei* GlnA$_1$ Met-Sox-P·ADP (PDB: 8tfk,

Schumacher et al., 2023) transition state (GlnA$_1$ TS), respectively. **A** *left:* GlnA$_1$ 2-OG dimer in

superposition with GlnA$_1$ apo showing large scale movements upon 2-OG binding. **A** *right:* Close-up

of 2-OG binding site of GlnA$_1$ 2-OG in superposition with GlnA$_1$ apo. Dramatic movement of Helix α3

(residue 167-181) and R87 loop show effect of 2-OG binding. **B**: Close-up of substrate binding site of

GlnA$_1$ 2-OG in superposition with GlnA$_1$ apo and ADP ligand from GlnA$_1$ TS. Helix α3 movement upon

2-OG binding leads to a cascade of conformational changes of the phenylalanines F184, F202 and

F204 that lead to a priming of the active site for ATP binding. **C**: Close-up of substrate binding site of

GlnA$_1$ 2-OG in superposition with GlnA$_1$ TS shows high similarity between 2-OG bound and transition

state structure. **D**: Close-up of substrate binding site of GlnA$_1$ 2-OG in superposition with GlnA$_1$ apo

and Met-Sox-P ligand from GlnA$_1$ TS. Large structural changes of the D50-loop with ejection of the

R66 key-residue shown. Flipping of the loop allows R319 and D57 to move in further and catalyze

phosphoryl-transfer and attack of NH_4^+, respectively. **E**: Close-up of the substrate binding site of

GlnA$_1$ 2-OG in in superposition with GlnA$_1$ TS reveals strong similarity between 2-OG bound and

transition state structure in the active site.

Figure 6: Feedback inhibition of GlnA$_1$ by glutamine. A: Specific activity of purified strep-tagged

GlnA$_1$ (wt) and the respective R66A-mutant protein was determined as described in Materials and

Methods in the presence of 12.5 mM 2-OG and after additional supplementation of 5 mM glutamine.

For wt and the R66A-mutant one out of two biological independent replicates are exemplarily

shown, the deviation indicates the average of four technical replicates. **B:** Superposition GS

structures without glutamine of *M. mazei* (blue, green) and *B. subtilis* (orange, pink; PDB: 4lnn,

Murray et al., 2013): substrate binding-site including R'66 (R'62, respectively), which are responsible

for feedback inhibition. **C**: Exemplary mass spectra of Strep-GlnA$_1$ with 12.5 mM 2-OG in presence

and absence of 5 mM glutamine. The molecular masses shown above the peaks correspond to a

Gaussian fit of the respective peak (Gaussian fit not shown).

Figure 7: Model of the various molecular mechanisms of glutamine synthetase activity regulation.

Comparison of the regulation of glutamine synthetase activity in *E. coli /Salmonella typhimurium*, and

B. subtilis, Synechocystis and *M. mazei*. GS are in general active in a dodecameric, unmodified

complex under nitrogen limitation. Upon an ammonium upshift, GS are inactivated by feedback

inhibition (BcGS, *E. coli),* covalent modification (adenylylation, EcGS) or binding of (small) inactivating

proteins (*Synechocystis*, BsGS). *M. mazei* GS on the contrary is regulated via the assembly of the

active dodecamer upon 2-OG-binding and furthermore is strongly feedback inhibited by glutamine.

(Bolay et al., 2018; Klähn et al., 2018, 2015; Stadtman, 2001; Travis et al., 2022b)**.** Created with

BioRender.com

FIGURE LEGENDS SUPPLEMENT

Figure S1: Affinity-purified Strep-GlnA$_1$ and size-exclusion-chromatography (SEC) of Strep-GlnA$_1$

after purification. A: 1.5 µg (lane 1) and 3 µg (lane 2) Strep-GlnA$_1$ on a coomassie-stained 12 % SDS-

Gel. **B**: Elution profile of Strep-GlnA$_1$ (black) and size standard (dashed line, molecular weights in

italics). Size exclusion chromatography was performed on a Superose™ 6 Increase 10/300 GL column

(Cytiva, Marlborough, USA) with a flow rate of 0.5 ml/min.

Figure S2: Sigmoidal fitted curves for mass photometry measurements of Strep-GlnA$_1$ with varying

concentrations of 2-OG. The curves were fitted and K$_D$-values calculated using RStudio (RStudio

Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL). **A, B**: Two

replicates for 2-OG titration, formation of dodecamer is shown in percent. **C, D**: 2-OG titration in the

absence (C) and presence of GlnK$_1$ (D), formation of dodecamer is shown as a ratio of

dodecamer/dimer.

Figure S3: Cryo-EM Data processing workflow A: Representative motion-corrected micrograph

showing different orientations of the GlnA$_1$ particles **B**: Cryo-EM processing tree used for obtaining

the high-resolution structure of GlnA$_1$. The map obtained is coloured by resolution, where the global

resolution was estimated using GSFSC **C**: Different regions of GlnA$_1$ encased around the cryo-EM

density.

Figure S4: Mass photometry of purified and thawed Strep-GlnA$_1$ before and after SEC. Mass spectra

of Strep-GlnA$_1$ samples with 0 and 12.5 mM after affinity-purification (blue ●, 0 mM and green ●,

12.5 mM 2-OG) and after SEC (0 mM 2-OG, grey ●).

Figure S5: Amino-acid sequence alignment of different model organism glutamine synthetases.

(Alignment tool: COBALT, visualization in SnapGene) Conserved amino-acids are highlighted in green.

The relevant residues in *M. mazei* for 2-OG- and substrate-binding, as well as the arginine

responsible for the feedback inhibition by glutamine are highlighted by coloured boxes (blue ●,

orange ● and purple ●, respectively).

Figure S6: *M. mazei* **GlnA$_1$ filaments. A**: Representative motion-corrected micrograph showing GlnA$_1$

filaments **B**: Reference-free 2D classes showcasing filament orientations **C, D**: 3D reconstructed map

of GlnA$_1$ filament and its model.

 SUPPLEMENTARY TABLES

 Table S1: Strains and plasmids.

	Properties	Reference
Strains		
E. coli DH5α	General cloning strain	(Hanahan, 1983)
E. coli BL21 (DE3)	Strain for protein expression	Thermo Fisher Scientific, Waltham, USA
E. coli BL21 (DE3) + pRIL	Strain for protein expression of genes with unusual codons/ Cm^R	Stratagene, La Jolla, USA
Plasmids		
pET21a	General cloning vector providing a C-terminal His_6-tag	Novagen/Merck, Darmstadt, Germany
pETSUMO	Expression vector providing an N-terminal His_6-SUMO-tag	Thermo Fisher Scientific, Waltham, USA
pRS375	pET28a/Strep + $GlnA_1$	(Gutt et al., 2021)
pRS1672	pETSUMO + $GlnK_1$	This work
pRS1728	pEX-A258 + codon-optimized $GlnA_1$ and sP26	Eurofins Scientific, Ebersberg, Germany
pRS1841	pRS375 + codon-optimized $GlnA_1$	This work
pRS1863	pET21a + codon-optimized sP26	This work
pRS1951	pRS1840mut: $R66AGlnA_1$	This work

840 **Table S2:** Cryo-EM data collection, refinement and validation statistics

	Dodecamer Gln complex
	(EMDB- 19730) (PDB 8S59)
Data collection and processing	
Magnification	165,000 x
Voltage (kV)	300
Electron exposure (e–/Å^2)	55.0
Defocus range (μm)	0.25-2.0
Pixel size (Å)	0.72
Symmetry imposed	D6
Initial particle images (no.)	1,243,001
Final particle images (no.)	878,308
Map resolution (Å)	2.39
FSC threshold	0.143
Map resolution range (Å)	2.3-3.0
Refinement	
Initial model used (PDB code)	*de novo,* AlphaFold
Model resolution (Å)	2.5
FSC threshold	0.5
Model resolution range (Å)	2.2-2.5
Map sharpening B factor (Å^2)	-80.1
Model composition	
Non-hydrogen atoms	46968
Protein residues	5352
Ligands	AKG: 12
B factors (Å^2)	
Protein	16.58
Ligand	15.64
R.m.s. deviations	
Bond lengths (Å)	0.003
Bond angles (°)	0.602
Validation	
MolProbity score	1.29
Clashscore	5.32
Poor rotamers (%)	0.46
Ramachandran plot	
Favored (%)	98.22
Allowed (%)	1.56
Disallowed (%)	0.23

Figure 1

Figure 2:

Figure 3

Figure 4

Figure 5

Figure 6:

Figure 7: