Title: Deciphering a beetle clock: individual and sex-dependent variation in daily

activity patterns

Running title: Locomotor activity rhythms in _T. castaneum_

Authors: Reshma R[1], Tobias Prüser[1], Nora K. E. Schulz[1], Paula M. F. Mayer[1], Maite

Ogueta[2], Ralf Stanewsky[2,3#], Joachim Kurtz[1,3*#]

[1]Institute for Evolution and Biodiversity, University of Münster, Hüfferstr. 1, 48149

Münster, Germany

[2]Institute of Neuro- and Behavioural Biology, University of Münster, 48149 Münster,

Germany

[3] Joint Institute for Individualisation in a Changing Environment, University of Münster

and Bielefeld University

[#]Shared senior author

*Corresponding author: Prof. Dr. Joachim Kurtz, Hüfferstr. 1, 48149 Münster,

joachim.kurtz@uni-muenster.de, T: +49 251 83-24661, F: +49 251 83-24668

Keywords

Circadian rhythm, diurnal, light entrainment, temperature entrainment, locomotor

activity patterns, individual variation, *Clock*, RNAi, cryptozoic

Abstract

Circadian clocks are inherent to most organisms, including cryptozoic animals

that seldom encounter direct light, and regulate their daily activity cycles. A conserved

suite of clock genes underpins these rhythms. In this study, we explore the circadian

behaviors of the red flour beetle *Tribolium castaneum*, a significant pest impacting

stored grain globally. We report on how daily light and temperature cues synchronize

distinct activity patterns in these beetles, characterized by reduced morning activity and

increased evening activity, anticipating the respective environmental transitions.

Although less robust, rhythmicity in locomotor activity is maintained in constant dark

and constant light conditions. Notably, we observed more robust rhythmic behaviors in

males than females with individual variations exceeding those previously reported for

other insect species. RNA interference targeting the *Clock* gene disrupted locomotor

activity rhythms. Our findings demonstrate the existence of a circadian clock and of

clock-controlled behaviors in *T. castaneum*. Furthermore, they highlight substantial

individual differences in circadian activity, laying the groundwork for future research on

the relevance of individual variation in circadian rhythms in an ecological and

evolutionary context.

Introduction

Virtually all lifeforms from bacteria to humans express daily patterns in their

physiological and behavioral processes, known as circadian rhythms, which enable them

to accommodate the recurring changes in their environment (Saunders, 2002). These

rhythms are not only reactions to the environment but are endogenously controlled by

molecular pacemaker systems called circadian clocks, which can be entrained to the

external periodic cues such as light and temperature, which are collectively known as

zeitgebers (Bell-Pedersen et al., 2005). Coordinating activities in anticipation to the

periodic environmental changes has obvious adaptive functions such as escaping

adverse environmental conditions, ensuring proper availability of food and mates,

synchronizing behaviors with conspecifics and reducing competition or predation risks

(Favreau et al., 2009; Vaze and Sharma, 2013). Apart from these extrinsic advantages, a

circadian clock confers an intrinsic advantage of maintaining a stable and optimal

temporal compartmentalization of biological processes regardless of any irregular

fluctuations in the environment. Even animals living in aperiodic conditions are known

to have a functioning molecular clock (Trajano and Menna-Barreto, 1996) which points

towards its intrinsic adaptive values.

Circadian clocks operate similarly across different taxa. Their cellular mechanism

is comprised of transcription-translation feedback loops of clock genes and their protein

products, capable of producing self-sustained circadian oscillations (Doležel, 2023;

Helfrich-Förster, 2004). Detailed knowledge about the molecular mechanisms of animal

clocks is available mainly from the studies in mouse and fruit fly as representatives of

mammals and insects, respectively. In the main circadian feedback loop of fruit flies, the

clock proteins PERIOD (PER) and TIMELESS (TIM) represent the transcriptional

repressors, whereas in mammals CRYPTOCHOROMES (CRY1 and CRY2) dimerize with the

PERIOD proteins (mPER1, mPER2 and mPER3) to perform the equivalent role. This

transcriptional inhibition occurs by blocking of their transcriptional activators CLOCK and

CYCLE (BMAL1 is the mouse homolog of CYCLE). A striking difference between the

mammalian and *Drosophila* clock lies in the mechanism by which light resets and

synchronizes the clock with the environment. Light induces TIM degradation in flies

whereas it leads to the transcriptional activation of the *Period* genes in mice (Helfrich-

Förster, 2004).

Insect clocks are best documented in *Drosophila*. However, given the enormous

diversity of insect species and their habitats, it is plausible that their molecular clock

mechanisms have diversified in response to their specific environmental conditions

(Kotwica-Rolinska et al., 2022; Tomioka and Matsumoto, 2015). Diversities in some clock

components are already known, for instance, certain insects like beetles and honeybees

possess the mammalian light-insensitive cryptochrome, CRY2, unlike *Drosophila*, which

possesses only the photosensitive CRY1. On the other hand, mosquitoes and monarch

butterflies are known to possess both CRY1 and CRY2 (Yuan et al., 2007). This

underscores the significance of investigating a broad range of insect models for gaining

a comprehensive understanding of various aspects of the circadian clock.

Tribolium castaneum, the red flour beetle, is a widely found pest of stored grains

and an important model organism of increasing relevance in many fields such as classical

genetics, population genetics, insect development and epigenetics (Campbell et al.,

2022). These flour beetles are cryptozoic, meaning that they are animals living mostly in

darkness (Brown et al., 2009) They have very little exposure to light in their habitat of

stored grains or behind tree barks, their supposed original environment (Good, 1936;

Sokoloff, 1974)**.** Their cryptozoic adaptations include loss of photoreceptors sensitive to

blue light (Jackowska et al., 2007) and an increase in odorant and gustatory receptors

(Richards et al., 2008). Even though the adults are normally known for being negatively

phototactic (Park, 1934), they crawl to flour surfaces from time to time where they can

get direct light exposure as observed in laboratory populations (Misra and Englert,

1985). Copulation usually occurs on the flour surface as well (Park, 1934). They are also

likely to experience temperature variations in their natural habitats. Despite spending

most of their life in the absence of light, we know that they still possess the genes of a

functional molecular clock (Li et al., 2018). However, whether *T. castaneum* indeed

displays rhythmic activity profiles, driven by a functional circadian clock, has not been

convincingly demonstrated.

T. castaneum has homologs for all clock genes, known in *D. melanogaster*

(Richards et al., 2008). First studies on the molecular clock in *T. castaneum* revealed that

Clock (*Clk*), *cycle* (*cyc*), *period* (*per*) and *timeless* (*tim*) are all rhythmically expressed (Li

et al., 2018). Instead of the *Drosophila*-type cryptochrome, CRY1, these beetles possess

the light-insensitive CRY2, the function of which remains unclear. It has been suggested

that CRY2 plays a role in inhibiting CLK and CYC, thus acting as a negative element of the

feedback loop as known in other insects such as Hymenoptera and Heteroptera

(Kotwica-Rolinska et al., 2022; Yuan et al., 2007). However, *T. castaneum* also possesses

the *Drosophila*-type TIM (Kotwica-Rolinska et al., 2022) along with the mammalian CRY2

making it a unique model to study insect clocks, also possibly in an evolutionary context

of transition from the *Drosophila*-type to mammalian-type clocks.

In insects, functional circadian clocks, and the rhythms they produce are

traditionally evaluated by measuring an individual's variation in locomotor activity

during the course of the day (Chiu et al., 2010). It has been shown that *T.castaneum*

exhibits rhythmic locomotor activity under light dark cycles (LD; Rath et al., 2021) as well

as in the absence of external cues in constant darkness (DD; Abe et al., 2021). However,

both studies report only average data of individually kept beetles, without determining

the overall rhythmicity (i.e., how many beetles behaved rhythmically in a certain

condition), nor showing individual actograms, which would allow to estimate the

individual variation within the population. It is important to investigate these behavioral

variations as an organism reacts to and modifies its environment through its behavior

(Duckworth, 2009) and therefore having sufficient behavioral variation can readily help

populations to adapt faster to changing environments (Bolnick et al., 2003).

Furthermore, to our knowledge, there are also no studies exploring whether the

circadian clock in these beetles can be synchronized by temperature fluctuations.

In the present study, we monitored the locomotor activity patterns of *T.*

castaneum to characterize in detail the circadian profiles in their movements. We

studied light as well as temperature entrainment and the effect of constant conditions

on an individual level to characterise the behavioral variations existing within the

population. We also performed RNAi to study the changes in activity patterns upon

knockdown of the clock gene encoding the key circadian transcription factor CLK. As *T.*

castaneum is an important model organism, knowledge about their biological clock and

how it influences their behavior and physiology can add to the knowledge in their proper

laboratory rearing conditions and proper timing of experiments. It can serve as an

exceptional model system for investigating the molecular clock mechanisms of

Coleoptera, where our current understanding is comparatively limited and thereby also

advancing our overall comprehension of insect clocks.

Materials and Methods

Beetle cultures

For this study, the Croatia 1 (Cro1) strain, collected in Croatia in 2010 and a

commonly used lab-adapted strain San Bernardino (SB), from San Bernadino, California,

USA, established by Sokoloff in 1960's were used (Milutinović et al., 2013). The

populations were kept at standard laboratory conditions of 30 °C, 70% humidity and a

12 h:12 h light: dark cycle (lights on at 08:00 and lights off at 20:00; light from

flourescent bulbs, 4000 K, cool white, 3350 lumen) in plastic boxes or glass jars with

foam stoppers to allow air circulation. The standard beetle diet consisted of heat-

sterilized (75 °C for at least 24 h) organic wheat flour (Bio Weizenmehl Type 550, dm -

Drogerie Market) mixed with 5% brewer's yeast powder (referred to as "flour" from now

on).

Locomotor activity

To monitor locomotor activity patterns, we used Drosophila Activity Monitoring

System (DAM5H-4, Trikinetics Inc., Waltham, MA, USA; referred to as "DAM5 system" in

the following). Individual beetle's movements interrupt one of four infrared light beams

intersecting the glass tubes housing single beetles, at equal distances from each other

(Rosato and Kyriacou, 2006, Fig S1) which enables the recording of an individual's

movements. Sexually matured virgin beetles of known sex were placed into an individual

glass tube (diameter: 5mm, length:65mm) with a flour-disc (40μl of flour/water solution

(0.15g/ml) dried overnight) as food inside on one end and sealed with cotton plugs on

both ends. The glass tubes were assembled into the DAM5 systems which further were

placed into incubators for the recording of activity. The standard rearing conditions were

maintained in the incubators throughout the period of recording unless stated

otherwise. Before the first day of recording, beetles acclimatized for at least six hours in

the activity monitors.

Circadian activity patterns under light entrainment followed by constant darkness (LD

DD)

For the general characterization of the circadian activity patterns under 12:12

Light/Dark (LD) and subsequent constant darkness (DD), we used the Cro1 population

(the beetles used belong to the handling control of a knockdown experiment). After sex

discrimination we individualized the pupae in wells of a 96-well flat-bottom micro

reaction plate (Type F, Sarstedt) containing 0.08 g of flour. 32 adults per sex, which were

at least two-week-old post eclosion were used for monitoring locomotor activity for an

initial seven days during LD at 30°C (normal rearing temperature) at a light intensity of

20 lux followed by seven days of DD.

Circadian activity patterns in constant light after light entrainment (LD LL)

To study the effect of constant light on the circadian activity patterns, we

monitored beetles of the Cro1 population in constant light (LL) after entrainment to LD,

at two different light intensities: 20 lux and 100 lux. Around two weeks post eclosion,

we used 32 adults per sex, which were individualized since sex discrimination at pupal

stage, for each light intensity. We recorded their activity for an initial seven days in the

LD followed by six days in LL.

Comparison of activity profiles between two laboratory populations

For comparing the circadian activity patterns of the two laboratory populations

Cro1 and SB, we separated the pupae according to sex and kept them in groups of 24

individuals in petri dishes (Ø10 cm, Waldeck) with *ad libitum* flour. For each population,

24 adults per sex which were at least 11 days post eclosion (SB population develops

slower than Cro1) were used. The activity recording was carried out in LD (20 lux) for 7

days followed by another 7 days of DD. During LD, a technical issue caused the

temperature and humidity to drop from 30°C to 25°C and from 70% to 50%, respectively

for 8 hours (from midnight until lights on at 8am).

Circadian activity patterns under temperature entrainment

To test if daily temperature cycles can entrain the locomotor activity patterns in

the absence of light, we recorded the beetles' behavior under oscillating temperatures

as the zeitgeber. Beetles from the Cro1 population were reared on flour-discs prepared

in 96-well plates from day 15 post oviposition onwards. During this time, they were not

exposed to light until 21 days post oviposition (as they also served as a control group in

a separate experiment, which is not described here). Around two weeks post eclosion,

beetles were sexed and 24 adults per sex were used for the activity recording for an

initial seven days of temperature cycling (12h 25°C/12h 30°C), followed by another

seven days of constant temperature, 30°C in DD. The activity recording was carried out

in complete darkness in both phases to make sure that the temperature change is the

only zeitgeber. The experiment was repeated a second time with the same conditions

(n=48 per sex during temperature entrainment and n=32 per sex during constant

temperature, sample size reduced due to recording error) except that the temperature

change was set to ramp for two hours to create a more natural condition of temperature

variations.

Activity profiles after knockdown of *Clk*

To test, if rhythmic behavior depends on the canonical molecular clock

mechanism, we performed RNAi-knockdown (Tomoyasu et al., 2008) of the key

circadian clock gene *Clk* at the pupal stage using the Cro1 population.

After sex discrimination, pupae were glued with their last abdominal segment

onto a glass slide for the dsRNA injection. The pupae were injected between the second

and third lowest segment of their abdomen with dsRNA of either *Clk* or jellyfish eGFP ,

which served as the control for the dsRNA injection treatment (Schulz et al., 2022). The

dsRNA construct of eGFP was produced in our lab (via cloning followed by PCR and *in*

vitro transcription using the T7 MEGAscript Kit (Ambion by Life TechnologiesTM GmbH,

Darmstadt, Gemany)). The dsRNA construct for *Clk* was obtained from EupheriaBiotech

(Dresden, Germany) and has previously been used in the ibeetle RNAi screen (Schmitt-

Engel et al., 2015, Table S1). A third treatment group served as the pure handling control

where the pupae glued on to the glass slides did not receive any injection. dsRNA was

injected at a concentration of 1000 ng/μL dissolved in Phosphate-buffered saline (PBS)

using a microliter injector (FemtoJet, Eppendorf AG, Hamburg, Germany) and

borosilicate glass capillaries (100 mm length, 1.0 mm outside diameter, 0.021 mm wall

thickness; Hilgenberg GmbH, Malsfeld, Deutschland). Upon eclosion, the beetles were

kept individually in 96-well plates with 0.08 g flour until the start of the activity

measurements in the DAM5 system.

The activity measurements started two weeks after eclosion. The activity of

beetles from all three groups were monitored in LD (light intensity of 100 lux) for seven

days followed by seven days in DD.

To confirm the successful downregulation following the RNAi treatment, we

measured *Clk* expression by qPCR. Fourteen days after eclosion, we shock froze the

individuals in liquid nitrogen. Furthermore, we usually pooled 3-4 whole bodies per

replicate. All samples were stored at -80°C until the RNA extraction.

We extracted total RNA by combining a TRIZOL-chloroform approach with a spin

column kit (SV Total RNA Isolation System, Promega (Peuß et al., 2016)). In short, we

ground the TRIZOL-submerged samples with a sterile pestle until no larger tissue parts

were visible. Samples then went into an ultrasound bath to further ensure disintegration

of tissue. After ten minutes of incubation, samples were spun down and the supernatant

was mixed with chloroform. Another centrifugation step ensured formation of phase

separation. We then transferred the upper aqueous phase mixed with lysis buffer onto

the spin column and proceeded according to manufacturer's protocol. Finally, RNA was

eluted with 80μl nuclease free water and stored at -80°C until further use.

For the qPCR, we first reverse transcribed RNA to cDNA using the RevertAid First

Strand cDNA Synthesis Kit (Thermo Scientific) and the provided oligo(dT)$_{18}$ primer,

following the manufacturer's instructions. Before the cDNA was used in the qPCR it was

diluted 1:5 with nuclease free water.

We performed qPCRs with technical duplicates and two housekeeping genes

rp49 and RpL13 a to normalize target gene expression. Amplification efficiency of

primers was previously established (Table S2) as well as the uniform expression of the

housekeeping genes across sexes and treatments. The qPCR ran on a LightCycler 480

System (Roche) with KAPA SYBR Fast mastermix (Sigma-Aldrich).

For each RNAi treatment and control, we measured gene expression separately

for males and females in four biological replicates. We included samples in the

downstream analysis if the standard deviation between the technical replicates

remained below 0.5 and the melting curves showed only one clear peak.

Behavior analyses

After two weeks of recording, we extracted the raw data of locomotor activity in

activity bins of 30 minutes using DAMFileScan software (TriKinetics). We looked for

activity patterns from the average histograms during different entrainment and

constant conditions, and differences in average activity levels, and proportion of

rhythmic beetles using data from individual beetles.

For the average comparison of diel patterns in movement, average histograms

of mean activity in 30-minute intervals for all individuals belonging to the same

treatment and sex group across the days of the same environmental condition, were

generated using the Fly-toolbox (Levine et al., 2002b).

Individual beetles were scored to be rhythmic or arrhythmic based on a

combination of qualitative and quantitative measures using double plotted actograms,

autocorrelation (Levine et al., 2002b) and χ^2-periodograms (Sokolove and Bushell, 1978)

generated using Fly-toolbox in MATLAB (MATLAB and Statistics Toolbox Release 2021b).

Double plotted actograms were used for the visual analysis to identify rhythmic

patterns. Each row in a double plotted actogram displays the activity in two consecutive

days for which the activity of the second day is replotted in the left part of the next row

to inspect for the activity patterns horizontally and vertically. A beetle is categorized as

rhythmic if activity patterns are repeated for at least half of the days of activity

recording. To reduce the subjectivity from the visual detection of rhythmicity,

quantitative thresholds from autocorrelation and periodograms were also used in

combination with the visual inspection of actograms (Levine et al., 2002b).

Autocorrelation graphs identify patterns in the data by examining the correlation

between the movement signal and a time-shifted version of itself. This analysis helps

detect both the strength of the rhythm (RS-value) and the duration of the identified

period (p). The χ2-periodogram assesses rhythmicity by testing various potential periods

within the data and evaluating their fit, represented by the Qp value. The dominant

signal period is determined based on the highest Qp value, while a 95% confidence

interval is used to evaluate the significance of the chosen period. Additionally, the h-

value indicates to which extend the Qp value of the selected period exceeds this

confidence interval. A beetle was categorized as rhythmic only if it met all three criteria:

having a clear rhythmic pattern in the actogram, RS>1 in the autocorrelation and h > 10

in the periodogram.

The circular phase plots showing the phase of individual beetles (Levine et al.,

2002b) were also plotted using MATLAB. For the analysis of activity levels, activity in 1-

minute intervals were averaged over all monitoring days of a specific condition for

individual beetles. The bar graphs, plotting proportion of rhythmic beetles and boxplots

for the average activity were generated using ggplot2 (Wickham, 2016) in RStudio

version 2021.09.1 (RStudio Team, 2021) under R version 4.1.2 (R Core Team, 2021). We

did some final editing of supplementary figure panels using Inkscape (Developers,

Inkscape Website, 2020)

Statistics

All the statistical analysis for the rhythmicity and average activity levels were

performed using R Studio employing additional package glmmTMB (Brooks et al., 2017).

The analysis for rhythmicity was done on binary data indicating either presence

or absence of consistent rhythmic patterns for individual beetles. The analysis for the

activity levels was done on the activity in 1-min bin averaged over all the days of a

condition for individual beetles. The differences in the proportion of rhythmic beetles

and activity levels were analyzed in generalized linear mixed effect models (GLMMs)

(Table S3) with the according error distributions. The final models were derived after p-

value based stepwise model simplification.

For the statistical analysis of gene expression after RNAi treatment, we used the

Relative Expression Software Tool (REST2009, Pfaffl et al., 2002). The tool is designed to

compare the gene expression between two groups – in our case, expression in the

knockdown treatment was compared to the expression in the RNAi control group. REST

employs a mathematical model, which takes the PCR efficiencies of the primers into

account when calculating expression ratios from the mean crossing point deviation. It

then performs a randomization test to determine statistically significant differences

between the treatment groups.

Results

To foster our understanding of the workings of the molecular clock in flour beetles and to further characterize the circadian patterns in their behavior, we first monitored and analysed the locomotor activity rhythms of an outbred population (Cro1) of *T. castaneum.* We tested the beetles' behavior under different light and temperature cycles for both sexes. Additionally, we compared the outbred strain (Cro1) to an inbred strain (SB), which has adapted to laboratory conditions for decades, to infer strain specific behavioral differences. Finally, we downregulated *Clk* gene expression via RNAi to investigate the role of this key circadian clock gene in generating the behavioral patterns.

Locomotor activity patterns under light entrainment

We monitored the locomotor activity patterns of beetles from the Cro1 population under LD (20 lux) conditions followed by DD. The average activity profiles of beetles showed a clear and repeatable diurnal rhythm (Fig 1 a-d). The daily average activity patterns were characterized by a consistent reduction in activity levels just before the onset of morning light in all the experiments, which we refer to as 'morning dip' (Figure 1 a). They reacted with a startle response of increased activity observed in response to the onset of the morning light stimulus (Figure 1a). This response fades

quickly within 30 minutes and is not observed under DD conditions. Upon more natural

like conditions with a linear ramping, extending the lights-on transition to 2 hours, the

startle response either disappeared or appeared at the end of the ramping period, but

as a lower peak compared to the non-ramping conditions (Fig S2). Further, beetles

gradually increased their activity levels throughout the late afternoon, a behavior which

we accordingly (and in analogy to a similar pattern observed in *D. melanogaster*

(Hamblen-Coyle et al., 1992)) call evening anticipation (Figure 1a). With the start of the

dark phase in LD, the beetles slightly reduced their activity levels from their highest peak

during the evening anticipation but remained highly active throughout the night until

they reduced their activity considerably during the morning dip (Figure 1 a and c).

Figure 1: Locomotor activity patterns of *T. castaneum*. a-d: Average histograms

showing mean activity of Cro1 beetles over the 7 days of recording in 30-minute bins

under LD (light entrainment of 20 lux) and DD; e: Boxplots showing activity in 1-minute

bin for individual beetles averaged over 7 days in LD and DD (males: n = 31, females:

n=32).

Locomotor activity patterns under constant darkness

In DD, the beetles' average activity profiles mirrored those observed during the LD except for the obvious absence of the startle response. Before the start of the subjective morning, there was a clear reduction of activity similar to the morning dip and a gradual increase of activity throughout the subjective late afternoon that resembled the evening anticipation (Fig 1 b and d). In addition to the generally higher levels of activity observed in females compared to males, the average histograms also illustrate the irregular and noisy patterns exhibited by the females (Fig 1 a-d). The analysis of individual average activity in 1-minute bins also reveals a significantly lower average activity for males (GLMM, estimate = -0.588, $p < 0.001$, Fig 1e). In terms of individual rhythmicity, 52% of males and 22% of females showed rhythmic patterns in LD whereas the proportion of rhythmic beetles was similar (32% for males and 28% for females) in DD conditions (Table 1). However, upon statistical analysis of individual rhythmicity, there was no significant effect for the sex (GLMM, estimate = 0.338 $p = 0.665$) or the light condition (GLMM, estimate = -0.491 $p=0.489$) whereas there was a weak trend for the interaction between the effects of sex and light condition (GLMM, estimate = 1.727, $p = 0.089$)

Variations in locomotor activity patterns

A substantial amount of individual variation can be seen in the rhythmic patterns

in LD (Figure 2 a-f). Among the rhythmic beetles, not all show identical patterns. Activity

peaks of many beetles start just before the lights turn off, and among these, some have

their peak of activity for a short duration before the lights turn off (Fig 2 c) or extending

further throughout the dark phase of the LD (Fig 2 a). The individual variations in phase

of their activity can also be seen from the circular phase plot (Fig 2 g). The females have

more coherent phase compared to the males. There was variation in terms of whether

the beetles could consistently maintain the rhythmicity throughout both light conditions

as well (Fig 3 a-d). Not all beetles that were rhythmic in LD were able to maintain their

rhythmicity during DD. 9 out of 31 males and 2 out of 32 females were rhythmic in LD

but showed arrhythmic behavior in DD. There was a small proportion of beetles (3 out

of 31 for males and 4 out of 32 for females) that showed rhythmic behavior only in DD

(Fig 3c, e).

Figure 2: Variation among individuals. a-f Variation in behavioral patterns among

individuals during LD; g: circular phase plot showing phases of beetles in LD based on

data from the first 3 days. The lengths of the line from the centre represents the

coherence of phases within a group.

Figure 3: Maintenance of rhythmicity under LD and DD. a- d: Individual actograms

showing different categories of beetles according to their status of being rhythmic (R)

or arrhythmic (Ar) under the two conditions of LD (first 7 days) and DD (last 7 days) (R/R

- rhythmic in LD and DD, R/Ar – rhythmic in LD but not in DD, Ar/R – Rhythmic in DD but

not in LD, Ar/Ar – Arrhythmic in both LD and DD), e: stacked bar plot showing the

proportion of these categories separately for males and females.

Locomotor activity patterns under constant light conditions

When LD was followed by constant light conditions (LL), the average activity

profiles of beetles resembled those observed in DD, however, overall rhythmicity

compared to DD was reduced under both light intensities tested (20 lux and 100 lux; Fig

4 and Fig S3, respectively, Table 1). When examining individual rhythmicity, males

showed a significantly higher proportion of rhythmic individuals compared to females

(GLMM, estimate = 1.711, p <0.001, Fig 4e). Furthermore, the proportion of rhythmic

beetles experienced a significant decrease under constant light conditions compared to

LD (GLMM, estimate = -3.193 p < 0.001, Fig 4e). There was an overall significant effect

of light intensity on the rhythmicity (GLMM, estimate = -1.002, p = 0.03, Fig 4e). There

was a significant interaction between light intensity and the light condition, with a

sharper drop in rhythmicity from LD to LL for the higher light intensity (GLMM, estimate

= 1.772, p= 0.026, Fig 4e).

Figure 4: Rhythmicity under LL conditions. a-d Average histograms showing mean

activity for all the beetles over the 7 days of recording under LD (light entrainment of 20

lux) and LL (20 lux) (Average histograms for 100 lux given in Fig S3), e: Proportion of

rhythmic beetles in LD and LL for two light intensities;20 lux and 100 lux (males: n = 32

per light intensity, females: n=32 per light intensity)

Population differences in locomotor activity and behavioral patterns

Both Cro1 and SB populations had similar behavioral patterns (Fig 5 a-h), but the

average activity was significantly lower for the SB population during LD (GLMM, estimate

= -0.667, p < 0.001, Fig 5 j) and DD (GLMM, estimate = -0.368, p = 0.04, Fig 5k). The

average activity of females was significantly higher than that of males in LD (GLMM,

estimate = -0.624, p < 0.001, Fig 5 j) but not in DD (GLMM, estimate = -0.118, p = 0.512,

Fig 5k). While examining individual rhythmicity, the SB population showed a significantly

higher proportion of rhythmic beetles in LD (GLMM, estimate = 1.116, p =0.006, Fig 5i).

The effect of light condition was also significant with a higher proportion of beetles

observed in LD (GLMM, estimate = 1.761, p <0.001, Fig 5 i). However, there was no

significant effect of sex on rhythmicity (GLMM, estimate = 0.413, p = 0.279, Fig 5 i).

Figure5: Population differences in rhythmicity. a-h: Average histograms showing mean

activity for all the beetles over the 7 days of recording under LD (light entrainment of 20

lux) and DD for Cro1 and SB populations, i: Proportion of rhythmic beetles in LD (Cro1

males: n = 23 , Cro1 females: n=24, SB males: n=24, SB females: n=23) and DD (Cro1

males: n = 20 , Cro1 females: n=19, SB males: n=20, SB females: n=22) j and k: Boxplots

showing activity in 1-minute bin for individual beetles averaged over 7 days in LD (j,

Cro1males: n = 23, Cro1 females: n=24, SB males: n = 24, SB females: n = 23) and DD (k,

Cro1 males: n = 22, Cro1 females: n=20, SB males: n = 21, SB females: n = 22)

Locomotor activity patterns during and after temperature entrainment

Similar to the behavior in LD, the beetles showed activity rhythms in a 12 hr: 12

hr temperature cycle of 25^0C: 30^0C in constant darkness (DD) (Fig 6 a-d, Fig S4). However

rhythmic behavioral patterns, apart from the prominent peaks of activity at the start of

temperature change in the subjective morning and evening, were less prominent

compared to LD. The peaks of activity at the temperature transitions resembled the

startle response observed after lights-on in LD and were not retained in constant

temperature conditions (Fig 6 a-d).

After the temperature cycles, only males showed a clear rhythmic activity

pattern under constant temperature conditions (Fig 6 c). The proportion of rhythmic

beetles significantly differed between sexes (GLMM, estimate = 1.418, p = 0.018, Fig 6

e).

Upon the more natural-like temperature cycling in constant darkness with a

ramping of temperature change within 2 hours, the activity patterns in the entrainment

phase appeared stronger compared to the entrainment with a square-wave of

temperature cycling (Fig S5). Both the startling peaks disappeared and there was a clear

dip in the morning activity as well as a stronger evening anticipation in the entrainment

phase resembling the pattern in LD (Fig S5). In the subsequent constant temperature

conditions, both males and females retained these activity patterns. There was no

difference in the proportion of rhythmic beetles between the sexes (GLMM, estimate =

0.185, p = 0.635, Fig S4 E) as well as between temperature cycles and constant

temperature conditions (GLMM, estimate = -0.427, p = 0.283, Fig S4 e).

Figure 6: Temperature entrainment. a-d Average histograms showing mean activity for

all the beetles over the 7 days of recording under temperature cycling and subsequent

constant temperature, both under constant darkness, e: Proportion of rhythmic beetles

in cycling temperature and constant temperature (males: n = 24 per temperature

condition, females: n=24 per temperature condition)

Locomotor activity patterns upon knockdown of *Clk*

We downregulated the key circadian clock gene *Clk* using RNAi in the pupal stage

to study the subsequent changes in activity profiles.

Clk gene expression was significantly different between the sexes (Table 2) in the

RNAi control. There was a significant reduction in *Clk* gene expression compared to the

RNAi control, indicating that dsRNA-mediated knockdown was efficient (Table 3).

Following the *Clk* knockdown, there was also a significant reduction in rhythmicity

(GLMM, estimate = -1.369, p = 0.005) in comparison to the RNAi control beetles,

showing that *Clk* function is important for behavioral rhythmicity (Fig 7). There was a

significant interaction effect of sex and light condition as well on the rhythmicity (GLMM,

estimate = 1.391, p = 0.04) (Fig 7).

Figure 7: Rhythmicity upon knockdown of *Clk*. Proportion of rhythmic individuals in

adults following the knockdown treatment at pupal stage for *Clk* in males (a) and females

(b) (LD: n =24 per sex and treatment groups, DD: n=24 per sex and treatment groups

except for RNAi control males (n = 23))

Table 1: Proportion of rhythmic beetles in different experiments

Experiment		Percentage of rhythmic beetles			
		Entrainment		Constant condition	
		Males	Females	Males	Females
LD (20 lux)/DD		51.6%	21.9%	32.3%	28.1%
LD LL	20 lux	50.0%	18.8%	18.8%	9.4%
	100 lux	75.0%	31.3%	12.5%	3.1%
Population difference in LD DD	Cro 1	52.2%	29.2%	5.0%	5.3%
	SB	58.3%	56.5%	30.0%	27.3%
Temperature entrainment		45.8%	25.0%	45.8%	16.7%

Temperature entrainment with ramping		35.4%	33.3%	28.1%	21.9%
Knockdown of *Clk*	Naïve	45.8%	20.8%	16.7%	12.5%
	RNAi control	58.3%	33.3%	17.4%	29.2%
	Clk knockdown	33.3%	16.7%	8.3%	4.2%

470 **Table 2: Sex difference in gene expression of *Clk***

Relative gene expression of males compared to females	95% C.I.	n	p value
0.312	0.158 - 0.795 ctrl vs ctrl (male vs male): 0.381 - 2.642	4	**0.02**

Table 3: Knockdown of *Clk*

Sex	Relative expression compared to RNAi control	95% C.I.	n	p value
Male	0.463	0.227 - 1.273 ctrl vs ctrl: 0.381 - 2.642	4	**0.031**
Female	0.531	0.279 - 0.988 ctrl vs ctrl: 0.539 - 1.858	4	**0.028**

Discussion

We used the Drosophila Activity Monitoring system (DAM) to perform a detailed characterization of the average as well as individual locomotor activity profiles of adult *T. castaneum* extending its application to study behavioral patterns in another important model organism. We recorded the activity of lab-adapted populations with high temporal resolution and described the activity patterns under light entrainment, constant dark conditions and for the first time to our knowledge, under thermal oscillations and constant light conditions. The beetles anticipated environmental transitions and even though less robust, maintained their rhythmicity in DD and LL. We showed that a substantial amount of individual variation exists in the diel patterns of

this cryptozoic species. The knockdown of *Clk* significantly reduced rhythmicity, strongly

supporting the existence of circadian clock in *T. castaneum* and its control of behaviors.

The finding that temperature cycling acts as a zeitgeber for the beetle clock, in

addition to light, is consistent with our expectations, considering that beetles are

exposed to cycling temperatures in their natural habitats of grain silos (Ziegler et al,

2021) as well as in their supposed original habitats under tree bark (Good, 1936;

Sokoloff, 1974). Studies in many other insects including fruit flies (Glaser and Stanewsky,

2005) and honeybees (Giannoni-Guzmán et al., 2021) have documented the role of light-

dark cycles as well as temperature oscillations as zeitgebers. The notable patterns of the

average activity profiles under both zeitgebers included a consistent drop of activity just

before the change of conditions in the morning which we refer to as 'morning dip' and

the gradual increase of activity throughout the late afternoon known as evening

anticipation. These patterns emerged even before the zeitgeber conditions change as

well as appeared consistently in constant conditions, indicating an anticipatory response

to the predicted changes resulting from their entrained molecular clock. Other insects

such as *D. melanogaster* (Dubowy and Sehgal, 2017) and *Plutella xylostella* (D. Wang et

al., 2021) also show such gradual increase in activity anticipating the transitions of

environmental conditions. However, a gradual decrease of activity (morning dip)

towards the time of dark-light transition is rather atypical.

The beetles reacted with a startle response of an exceptionally heightened

activity to the light turning on in the morning as well as to the change of temperature in

the subjective morning and evening. The absence of these startle responses in constant

conditions as well as their disappearance or strong reduction under more natural-like

conditions of ramping temperature or light conditions confirms that they are not

mediated by the circadian clock but are instead reactions to sudden changes in

environmental conditions. Similar startle responses immediately following the dark-light

transition were observed in *Ae. aegypti*, and, consistent with our study, these responses

were not present in constant darkness (Gentile et al., 2009) .

Average histograms consistently exhibited evening anticipation in constant

conditions as well as following the entrainment phase for both light and temperature as

zeitgebers. A steep evening anticipation was characteristic of the average activity

profiles under light entrainment, especially for males. The evening anticipation aligns

well with the time of day when mating rates of the *T.castaneum* reach their maximum

(Z. Wang et al., 2023). In red flour beetles, males initiate the copulatory behavior and

must mate multiple times to ensure fertilization success, especially because of the last

male sperm precedence (Fedina and Lewis, 2008). An increased activity of males in

search of mates could be a potential reason why males possess a strong and clear

evening anticipation peak compared to the females. Similarly, in *P. xylostella*, males

were shown to have better rhythmic features than females under constant darkness in

low temperature conditions (D. Wang et al., 2021). Such sex-specific differences in

activity profiles are commonly observed in many insect species and are often associated

with their mating behavior. For example, in *Anopheles gambiae*, males have an earlier

flight activity peak compared to females, consistent with the fact that males aggregate

to form swarms to which females subsequently enter (Rund et al., 2012). In addition to

the difference in evening anticipation, the overall activity levels of females were

significantly higher than those of males, as was also reported recently (Benita et al.,

2024).   we also observe a large inter-individual variation in phase for the males

compared to females. Additionally, males generally exhibit a higher proportion of

rhythmicity, which may be attributed again to the necessity for males to initiate

copulation behavior, thereby requiring precise timing of such behaviors.

In addition to the individual variation in phase, there was substantial variation in

the individual activity patterns including differences in the duration of peak activity,

levels of activity at different times, and the ability to maintain rhythmic behavior under

constant conditions following entrainment. A study in the cave beetle species,

Laemostenus latialis also showed similar variability in individual behavioral patterns

(Pasquali and Sbordoni, 2014). Another study observed significant diversities in

circadian behaviors in social insect species compared to the solitary species like fruit flies

(Giannoni-Guzmán et al., 2014). Since *T.castaneum* is a non-social yet group living insect,

one could speculate that the dual role of the pheromone, 4,8-dimethyldecanal (DMD)as

an aggregation as well as a sex attractant for females (Levinson and Levinson, 1995)

enables individuals to rely on their conspecifics for timing their own mating rather than

a strict reliance on their circadian clock. Another possibility is that variability in their

peak activity timings could help in avoiding any possible male-male competition.

The proportion of rhythmic beetles considerably decreased in DD conditions

especially for males **.** The presence of all clock gene homologs known in *Drosophila*

(Richards et al., 2008) indicates that *T. castaneum* possesses a functional circadian clock.

Therefore, the decrease in rhythmicity in constant darkness compared to that under

light entrainment could possibly indicate a weak link or uncoupling between the

endogenous clock and the locomotor behavior under constant darkness (Abhilash et al.,

2017) as seen in the case of blind mole rats (Beale et al., 2016) and in *Drosophila* species

inhabiting higher latitudes with weakly rhythmic environments (Bertolini et al., 2019).

The inter-individual variability seen in locomotor activity profiles could also possibly be

the result of a weak behavioral regulation by the endogenous clock (Beale et al., 2016).

One could speculate that such a weak coupling or even uncoupling might be due to the

absence or reduced selective pressure for circadian functions in their possible ancestral

natural dark habitats such as rotting logs and tree barks as well as in their recent, also

largely dark habitats of grain silos (Good 1936; Sokoloff 1974). The possible chance of

relying on olfactory cues from conspecifics for mating initiation might have also resulted

in a weak selection pressure for the evolution of a strong endogenous clock or a strong

coupling of the clock with behaviors.

There was a significant decrease in the proportion of rhythmic beetles in

constant light conditions, more pronounced under higher light intensity, even if weak

rhythmic patterns were observable in the average histograms. In wild-type D.

melanogaster, strong constant light leads to arrhythmic behavior (Konopka et al., 1989,

Matsumoto et al., 1994) because of the light-dependent TIM degradation by CRY1

which conveys photic signals to the circadian clock (Ceriani et al., 1999). The beetles are

not expected to have such an arrhythmic behavior as they possess only the light-

insensitive cryptochrome (CRY 2) (Yuan et al., 2007). However, most likely retinal and/or

potentially also unknown extraretinal photoreceptors play a role (Helfrich-Förster,

2020) in photic entrainment, which could then possibly affect the rhythmicity under

such constant light conditions. A similar reduction of rhythmicity under constant light

conditions was reported for *Nasonia vitripennis*, which also possesses only the light-

insensitive CRY 2 (Bertossa et al., 2014), whereas some other closely related *Nasonia*

species retained their rhythmicity (Bertossa et al., 2013). A study in linden bug, which

also possesses only CRY 2, revealed a deteriorating effect of constant light on

rhythmicity even though around 40% of individuals showed strong rhythmicity under

such constant light conditions (Kaniewska et al.,, 2020)

To test if the observed behavioral rhythms depend on a molecular circadian

clock, we attempted the knockdown of the key circadian clock gene *Clk* in the pupal

stage and subsequently monitored locomotor activity profiles at the adult stage. Indeed,

rhythmicity in *Clk* dsRNAi animals was significantly reduced compared to the control *GFP*

dsRNAi beetles, strongly supporting a role for a canonical CLK-dependent

transcriptional/translational circadian feedback loop in *T. castaneum*. In the future,

double knockdowns of different clock genes could be performed to induce more severe

clock disruption and phenotypes.

Since *T.castaneum* is a non-social but gregarious species, further experiments to

study their circadian rhythms in a group setting would be the next step. The social

influence on circadian rhythms has not only been demonstrated in social insects, but

also for a solitary species like *Drosophila* for which this is most likely mediated by

olfactory cues (Levine et al., 2002a). Studies of circadian rhythms of *T.castaneum* in a

social setting are especially important, because it is already known that an exposure to

the male aggregation pheromone increases the expression of the *timeless* clock gene in

these beetles (Rath et al., 2021).

As *T. castaneum* is a ubiquitous pest, knowledge about its diel patterns in

behavior under different external environmental cues can be useful in pest management

strategies, for instance in light traps. Recently, circadian clock research is gaining more

importance in immune system homeostasis (Haspel et al., 2020). Improved knowledge

about the circadian clock and the rhythms it controls in *T. castaneum,* which is an

excellent model to study insect immunity (Altincicek et al., 2008; Ferro et al., 2019), can

be beneficial for the emerging research in the intersection of circadian function and

immune regulation which has potential clinical relevance. It is remarkable that despite

the absence of strong light cues in their original habitats, the molecular clock of flour

beetles still shows a strong entrainment to light as well as that many beetles can retain

this rhythmicity under constant conditions. The considerable variation seen in different

aspects of the diel patterns along with its status as a group living yet non-social insect,

renders this organism a unique model for investigating circadian clocks and the different

behavioral and physiological parameters they control. This complements studies in

other insect groups, contributing to a comprehensive understanding of circadian

rhythms across different insect species.

Acknowledgments

We thank Hannah Goldgräbe for helping with the experiments.

Conflict of interest statement

The authors have no potential conflicts of interest.

Note

Supplementary material for this article including the final data files is available online.

Author contributions

JK and RS conceived the ideas. JK, RS, MO, RR, NS and TP planned the

experimental procedure. RR, TP, NS and PM carried out the experiments and RR carried

out the statistical analyses together with NS, MO, TP and PM. RR wrote the first draft

which was revised by NS and TP. RR and TP made the figure panels. All authors

contributed to writing of the article in further revisions and approved the submitted

version.

Funding

This research was funded by the German Research Foundation (DFG) as part of

the SFB TRR 212 (NC3, Project number 316099922) – Project number 396780003 (to

Joachim Kurtz)

References

Abe, M. S., Matsumura, K., Yoshii, T., and Miyatake, T. (2021). Amplitude of circadian rhythms becomes weaken in the north, but there is no cline in the period of rhythm in a beetle. *PLOS ONE*, 16: e0245115.

Abhilash, L., Shindey, R., and Sharma, V. K. (2017). To be or not to be rhythmic? A review of studies on organisms inhabiting constant environments. *Biological Rhythm Research*, 48: 677–691.

Altincicek, B., Knorr, E., and Vilcinskas, A. (2008). Beetle immunity: Identification of immune-inducible genes from the model insect Tribolium castaneum. *Developmental & Comparative Immunology*, 32: 585–595.

Beale, A. D., Whitmore, D., and Moran, D. (2016). Life in a dark biosphere: A review of circadian physiology in "arrhythmic" environments. *Journal of Comparative Physiology. B, Biochemical, Systemic, and Environmental Physiology*, 186: 947–968.

Bell-Pedersen, D., Cassone, V. M., Earnest, D. J., Golden, S. S., Hardin, P. E., Thomas, T. L., and Zoran, M. J. (2005). Circadian rhythms from multiple oscillators: Lessons from diverse organisms. *Nature Reviews Genetics*, 6: 544–556.

Benita, M., Menahem, A., Scharf, I., and Gottlieb, D. (2024). Inter- and intraspecific

female behavioral plasticity drive temporal niche segregation in two Tribolium

species. *Behavioral Ecology*, 35: arad116.

Bertolini, E., Schubert, F. K., Zanini, D., Sehadová, H., Helfrich-Förster, C., and Menegazzi,

P. (2019). Life at High Latitudes Does Not Require Circadian Behavioral

Rhythmicity under Constant Darkness. *Current Biology*, 29: 3928-3936.e3.

Bertossa, R. C., Dijk, J. van, Diao, W., Saunders, D., Beukeboom, L. W., and Beersma, D.

G. M. (2013). Circadian Rhythms Differ between Sexes and Closely Related

Species of Nasonia Wasps. *PLOS ONE*, 8: e60167.

Bertossa, R. C., van de Zande, L., Beukeboom, L. W., and Beersma, D. G. M. (2014).

Phylogeny and oscillating expression of period and cryptochrome in short and

long photoperiods suggest a conserved function in Nasonia vitripennis.

Chronobiology International, 31: 749–760.

Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., and

Forister, M. L. (2003). The Ecology of Individuals: Incidence and Implications of

Individual Specialization. *The American Naturalist*, 161: 1–28.

Brooks, M. E., Kristensen, K., Benthem, K. J. van, Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J. Mächler, M., and Bolker, B. M. (2017). glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling.

The R Journal, 9: 378–400.

Brown, S. J., Shippy, T. D., Miller, S., Bolognesi, R., Beeman, R. W., Lorenzen, M. D.,

Bucher, G., Wimmer, E. A., and Klingler, M. (2009). The red flour beetle,

Tribolium castaneum (Coleoptera): A model for studies of development and pest

biology. *Cold Spring Harbor Protocols*, 2009: pdb. emo126.

Campbell, J. F., Athanassiou, C. G., Hagstrum, D. W., and Zhu, K. Y. (2022). Tribolium

castaneum: A Model Insect for Fundamental and Applied Research. *Annual*

Review of Entomology, 67: 347–365.

Ceriani, M. F., Darlington, T. K., Staknis, D., Más, P., Petti, A. A., Weitz, C. J., and Kay, S.

A. (1999). Light-dependent sequestration of TIMELESS by CRYPTOCHROME.

Science (New York, N.Y.), 285: 553–556.

Chiu, J. C., Low, K. H., Pike, D. H., Yildirim, E., and Edery, I. (2010). Assaying Locomotor

Activity to Study Circadian Rhythms and Sleep Parameters in Drosophila. *JoVE*

(Journal of Visualized Experiments), e2157.

Developers,Inkscape Website. (2020). Draw Freely | Inkscape. Retrieved from

https://inkscape.org./.

Doležel, D. (2023). Molecular Mechanism of the Circadian Clock. In H. Numata and K.

Tomioka (Eds.), *Insect Chronobiology*, Singapore: Springer Nature (pp. 49–84).

Dubowy, C., and Sehgal, A. (2017). Circadian Rhythms and Sleep in Drosophila

melanogaster. *Genetics*, 205: 1373–1397.

Duckworth, R. A. (2009). The role of behavior in evolution: A search for mechanism.

Evolutionary Ecology, 23: 513–531.

Favreau, A., Richard-Yris, M.-A., Bertin, A., Houdelier, C., and Lumineau, S. (2009). Social

influences on circadian behavioural rhythms in vertebrates. *Animal Behaviour*,

77: 983–989.

Fedina, T. Y., and Lewis, S. M. (2008). An integrative view of sexual selection in Tribolium

flour beetles. *Biological Reviews*, 83: 151–171.

Ferro, K., Peuß, R., Yang, W., Rosenstiel, P., Schulenburg, H., and Kurtz, J. (2019).

Experimental evolution of immunological specificity. *Proceedings of the National*

Academy of Sciences, 116: 20598–20604.

Gentile, C., Rivas, G. B. S., Meireles-Filho, A. C. A., Lima, J. B. P., and Peixoto, A. A. (2009).

Circadian expression of clock genes in two mosquito disease vectors: Cry2 is

different. *Journal of Biological Rhythms*, 24: 444–451

Giannoni-Guzmán, M. A., Avalos, A., Perez, J. M., Loperena, E. J. O., Kayım, M., Medina,

J. A., Massey, S. E., Kence, M., Kence, A., Giray, T., et al (2014). Measuring

individual locomotor rhythms in honey bees, paper wasps and other similar-sized

insects. *Journal of Experimental Biology*, 217: 1307–1315.

Giannoni-Guzmán, M. A., Rivera-Rodriguez, E. J., Aleman-Rios, J., Melendez Moreno, A.

M., Pérez Ramos, M., Pérez-Claudio, E., Loubriel, D., Moore, D., Giray, T., and

Agosto-Rivera, J. L. (2021). The Role of Colony Temperature in the Entrainment

of Circadian Rhythms of Honey Bee Foragers. *Annals of the Entomological Society*

of America, 114: 596–605.

Glaser, F. T., and Stanewsky, R. (2005). Temperature Synchronization of the Drosophila

Circadian Clock. *Current Biology*, 15: 1352–1363.

Good, N. E. (1936). The flour beetles of the genus Tribolium. *USDA Technical Bulletin*,

27–28.

Hamblen-Coyle, M. J., Wheeler, D. A., Rutila, J. E., Rosbash, M., and Hall, J. C. (1992).

Behavior of period-altered circadian rhythm mutants ofDrosophila in light: Dark

cycles (Diptera: Drosophilidae). *Journal of Insect Behavior*, 5: 417–446.

Haspel, J. A., Anafi, R., Brown, M. K., Cermakian, N., Depner, C., Desplats, P., Gelman, A.

E., Haack, M., Jelic, S., Kim, B. S., et al. (2020). Perfect timing: Circadian rhythms,

sleep, and immunity — an NIH workshop summary. *JCI Insight*, 5: e131487

Helfrich-Förster, C. (2004). The circadian clock in the brain: A structural and functional

comparison between mammals and insects. *Journal of Comparative Physiology*

A, 190: 601–613.

Helfrich-Förster, C. (2020). Light input pathways to the circadian clock of insects with an

emphasis on the fruit fly Drosophila melanogaster. *Journal of Comparative*

Physiology A, 206: 259–272.

Jackowska, M., Bao, R., Liu, Z., McDonald, E. C., Cook, T. A., and Friedrich, M. (2007).

Genomic and gene regulatory signatures of cryptozoic adaptation: Loss of blue

sensitive photoreceptors through expansion of long wavelength-opsin

expression in the red flour beetle Tribolium castaneum. *Frontiers in Zoology*, 4:

24.

Kaniewska, M. M., Vaněčková, H., Doležel, D., and Kotwica-Rolinska, J. (2020). Light and

Temperature Synchronizes Locomotor Activity in the Linden Bug, Pyrrhocoris

apterus. *Frontiers in Physiology*, *11*. Retrieved from

https://www.frontiersin.org/articles/10.3389/fphys.2020.00242

Konopka, R. J., Pittendrigh, C., and Orr, D. (1989). Reciprocal Behaviour Associated with

Altered Homeostasis and Photosensitivity of Drosophila Clock Mutants. *Journal*

of Neurogenetics, 6: 1–10.

Kotwica-Rolinska, J., Chodáková, L., Smýkal, V., Damulewicz, M., Provazník, J., Wu, B. C.-

H., Hejníková, M., Chvalová, D., and Doležel, D. (2022). Loss of Timeless Underlies

an Evolutionary Transition within the Circadian Clock. *Molecular Biology and*

Evolution, 39: msab346.

Levine, J. D., Funes, P., Dowse, H. B., and Hall, J. C. (2002a). Resetting the Circadian Clock

by Social Experience in Drosophila melanogaster. *Science*, 298: 2010–2012.

Levine, J. D., Funes, P., Dowse, H. B., and Hall, J. C. (2002b). Signal analysis of behavioral

and molecular cycles. *BMC Neuroscience*, 3:1.

Levinson, A., and Levinson, H. (1995). Reflections on structure and function of

pheromone glands in storage insect species. *Anzeiger Für Schädlingskunde,*

Pflanzenschutz, Umweltschutz, 68: 99–118.

Li, C.-J., Yun, X.-P., Yu, X.-J., and Li, B. (2018). Functional analysis of the circadian clock

gene timeless in Tribolium castaneum. *Insect Science*, 25: 418–428.

Matsumoto, A., Motoshige, T., Murata, T., Tomioka, K., Tanimura, T., and Chiba, Y.

(1994). Chronobiological analysis of a new clock mutant, Toki, in Drosophila

melanogaster. *Journal of Neurogenetics*, 9: 141–155.

Milutinović, B., Stolpe, C., Peuß, R., Armitage, S. A. O., and Kurtz, J. (2013). The Red Flour

Beetle as a Model for Bacterial Oral Infections. *PLOS ONE*, 8: e64638.

Misra, R. G., and Englert, D. C. (1985). Differential effects of light upon fecundity upon

tribolium castaneum. *Transactions of the Illinois State Academy of Science*, 105–

109.

Park, T. (1934). Observations on the general biology of the flour beetle, Tribolium

confusum. *Quarterly Review of Biology*, 9: 36–54.

Pasquali, V., and Sbordoni, V. (2014). High variability in the expression of circadian

rhythms in a cave beetle population. *Biological Rhythm Research*, 45: 925–939.

Peuß, R., Wensing, K. U., Woestmann, L., Eggert, H., Milutinović, B., Sroka, M. G. U.,

Scharsack, J. P., Kurtz, J., and Armitage, S. A. O. (2016). Down syndrome cell

adhesion molecule 1: Testing for a role in insect immunity, behaviour and

reproduction. *Royal Society Open Science*, 3: 160138.

Pfaffl, M. W., Horgan, G. W., and Dempfle, L. (2002). Relative expression software tool

(REST©) for group-wise comparison and statistical analysis of relative

expression results in real-time PCR. *Nucleic Acids Research*, 30: e36.

R Core Team. (2021). *R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.* Retrieved from

https://www.R-project.org/

Rath, A., Benita, M., Doron, J., Scharf, I., and Gottlieb, D. (2021). Social communication

activates the circadian gene Tctimeless in Tribolium castaneum. *Scientific

Reports*, 11: 16152.

Richards, S., Gibbs, R. A., Weinstock, G. M., Brown, S. J., Denell, R., Beeman, R. W., Gibbs,

R., Bucher, G., Friedrich, M., Grimmelikhuijzen, C., et al. (2008). The genome of

the model beetle and pest Tribolium castaneum. *Nature*, 452: 949–955.

Rosato, E., and Kyriacou, C. P. (2006). Analysis of locomotor activity rhythms in

Drosophila. *Nature Protocols*, 1: 559–568.

RStudio Team. (2021). *RStudio: Integrated Development Environment for R. RStudio,*

PBC,Boston, MA. Retrieved from http://www.rstudio.com/

Rund, S. S. C., Lee, S. J., Bush, B. R., and Duffield, G. E. (2012). Strain- and sex-specific

differences in daily flight activity and the circadian clock of *Anopheles gambiae*

mosquitoes. *Journal of Insect Physiology*, 58: 1609–1619.

Saunders, D. S. (2002). *Insect Clocks*. Elsevier.

Schmitt-Engel, C., Schultheis, D., Schwirz, J., Ströhlein, N., Troelenberg, N., Majumdar,

U., Dao, V. A., Grossman, D., Richter, T., Tech, M., et al. (2015). The iBeetle large-

scale RNAi screen reveals gene functions for insect development and physiology.

Nature Communications, 6:7822.

Schulz, N. K. E., Mohamed, F. F., Lo, L. K., Peuß, R., de Buhr, M. F., and Kurtz, J. (2022).

Paternal knockdown of tRNA(cytosine-5-)-methyltransferase (Dnmt2) increases

offspring susceptibility to infection in red flour beetles. *Insect Molecular Biology*,

31: 711–721.

Sokoloff, A. (1974). *The Biology of Tribolium: With Special Emphasis on Genetic Aspects.*

(Vol. 2). Clarendon Press, Oxford.

Sokolove, P. G., and Bushell, W. N. (1978). The chi square periodogram: Its utility for

analysis of circadian rhythms. *Journal of Theoretical Biology*, 72: 131–160.

Tomioka, K., and Matsumoto, A. (2015). Circadian molecular clockworks in non-model

insects. *Current Opinion in Insect Science*, 7: 58–64.

Tomoyasu, Y., Miller, S. C., Tomita, S., Schoppmeier, M., Grossmann, D., and Bucher, G.

(2008). Exploring systemic RNA interference in insects: A genome-wide survey

for RNAi genes in Tribolium. *Genome Biology*, 9: R10.

Trajano, E., and Menna-Barreto, L. (1996). Free-Running Locomotor Activity Rhythms in

Cave-Dwelling   Catfishes,   Trichomycterus   sp.,   from   Brazil   (Teleostei,

Siluriformes). *Biological Rhythm Research*, 27: 329–335.

Vaze, K. M., and Sharma, V. K. (2013). On the Adaptive Significance of Circadian Clocks

for Their Owners. *Chronobiology International*, 30: 413–433.

Wang, D., Yang, G., and Chen, W. (2021). Diel and Circadian Patterns of Locomotor

Activity in the Adults of Diamondback Moth (Plutella xylostella). *Insects*, 12: 727.

Wang, Z., Zhao, Y., Wang, Y., and Lu, Y. (2023). Population divergence in pheromone

communication for mate choice of Tribolium castaneum and its association with

gut   bacteria.   *Journal   of   Pest   Science*.   Retreived   from

https://doi.org/10.1007/s10340-023-01632-6

Wickham, H. (2016). *Ggplot2*. Cham: Springer International Publishing.

Yuan, Q., Metterville, D., Briscoe, A. D., and Reppert, S. M. (2007). Insect Cryptochromes:

Gene Duplication and Loss Define Diverse Ways to Construct Insect Circadian

Clocks. *Molecular Biology and Evolution*, 24: 948–955.

Ziegler, V., Paraginski, R. T., and Ferreira, C. D. (2021). Grain storage systems and effects

of moisture, temperature and time on grain quality—A review. *Journal of Stored*

Products Research, 91: 101770.

822
823