Developmental Regulation of Alternative Polyadenylation in an Adult Stem Cell Lineage

Lorenzo Gallicchio[1], Neuza R. Matias[1], Fabian Morales-Polanco[2,3], Iliana Nava[1], Sarah Stern[1], Yi Zeng[3], Margaret T. Fuller[1,3,C]

[1]Department of Developmental Biology, Stanford University School of Medicine, Stanford USA.

[2]Department of Biology, Stanford University School of Humanities and Sciences, Stanford USA.

[3]Department of Genetics, Stanford University School of Medicine, USA.

[C]Correspondence to mtfuller@stanford.edu

Running Title: Developmental regulation of alternative polyadenylation

Keywords: Alternative Polyadenylation, Development, Cellular Differentiation, Spermatogenesis, mRNA processing, Cleavage factor Complex II.

Abstract

Co-transcriptional alternate processing of nascent mRNA molecules can make major

contributions to cell type specific gene expression programs as proliferating precursor cells

initiate terminal differentiation. Alternative Cleavage and Polyadenylation (APA) can result in the

production of mRNA isoforms from the same gene locus with either longer or shorter 3'UTRs. In

Drosophila spermatogenesis, approximately 500 genes undergo APA as proliferating

spermatogonia differentiate into spermatocytes, producing transcript isoforms with shortened

3'UTRs, and resulting in profound stage specific changes in the proteins expressed. The molecular

mechanisms that specify usage of upstream polyadenylation sites in spermatocytes are thus key

to understanding the changes in cell state. Here, we show that PCF11 and Cbc, the two

components of Cleavage factor II (CFII), orchestrate APA switching during *Drosophila*

spermatogenesis. Knockdown of *PCF11* or *cbc* in spermatocytes caused dysregulation of APA, with

many transcripts normally cleaved at a proximal site in spermatocytes now cleaved at their distal

site, as in spermatogonia. Although *PCF11* is widely expressed, *cbc* is strongly upregulated in

spermatocytes. Our findings reveal a developmental mechanism where changes in activity of

specific cleavage factors can direct cell type specific APA at selected genes, presenting CFII as a

key developmental regulator of APA during spermatogenesis.

Introduction

Alternative cleavage and polyadenylation (APA) of mRNAs results from changes in the site at which the 3' end cut that terminates nascent transcripts is made (Tian and Manley 2016; Gallicchio *et al.* 2023). If alternative polyadenylation sites are located in the 3'UTR of a genetic locus, APA results in production of alternate mRNA isoforms that differ in 3'UTR length and content, with potentially profound effects on transcript fate, including translation, stability, and localization (An *et al.* 2008; Mayr and Bartel 2009; Mansfield and Keene 2012; Berry *et al.* 2022). Since transcripts from more than 70% of human genes are alternatively polyadenylated under one condition or another, APA is widespread (Gruber and Zavolan 2019). The most intriguing cases are those where the choice of 3' end cut site is developmentally regulated, leading, for example, to production of transcripts with longer 3'UTRs from specific genes in neurons or shorter 3'UTRs in differentiating male germ cells compared to precursor cells (Hilgers *et al.* 2012; Miura *et al.* 2013; Shan *et al.* 2017; Vallejos Baier *et al.* 2017; Grassi *et al.* 2019; Berry *et al.* 2022). Changes in 3'UTR length due to APA have also been observed in many pathological states, such as the 3'UTR shortening of many transcripts observed in some cancers (Mayr and Bartel 2009; Morris *et al.* 2012; Venkat *et al.* 2020). Because these changes in 3'UTR content can have dramatic effects on protein expression and so cell state, cell type specific regulation of APA is emerging as an important developmental mechanism to promote robust cell fate decisions.

During cleavage and polyadenylation, the last step of mRNA synthesis, the nascent mRNA molecule is cleaved and a string of adenosines is added by nuclear PolyA polymerase to protect the newly made transcript from degradation and facilitate its export from the nucleus (Jing *et al.* 1999). Most mRNAs undergo cleavage and polyadenylation, with a notable exception being histone transcripts, which are not polyadenylated (Marzluff *et al.* 2008). Cleavage and polyadenylation is mediated by the cleavage machinery, which is composed of 4 well conserved multi-protein-complexes: the core complex CPSF (Cleavage and Polyadenylation Factor), which contains the endonuclease (CPSF73) that makes the actual cut that forms the transcript 3' end, CstF (Cleavage and Stimulation Factor), CFI (Cleavage Factor I) and CFII (Cleavage Factor II). The cleavage machinery is also associated with several non-complex proteins: Symplekin (a scaffolding protein), PABP2 and the nuclear PolyA polymerase (Figure 1 A) (Gallicchio *et al.* 2023).

The site at which 3' end cleavage will take place is guided by recognition by CPSF through the

components CPSF30 and WDR33 of the polyadenylation signal (PAS), an hexamer (most

commonly AAUAAA or a close variant) in the nascent mRNA usually ~15-30 nucleotides upstream

of the cleavage site (Clerici *et al.* 2018; Gallicchio *et al.* 2023). While the central role of the core

complex CPSF in recognizing the PAS and cleaving the nascent mRNA molecule is relatively clear,

the specific contributions of the other complexes and associated factors in specifying whether,

where, on what specific gene products, and in which cell types a cut is made is under investigation

(Vallejos Baier *et al.* 2017; Zhu *et al.* 2018; Turner *et al.* 2020).

Developmentally regulated changes in the PAS site utilized can have profound effects on

the suite of proteins expressed as cells progress from one state to the next in a developmental

sequence. For example, during spermatogenesis in flies and mice, many genes express mRNA

isoforms with long 3'UTRs in proliferating spermatogonia, but isoforms with shorter 3'UTRs at

later stages of germ cell development (Li *et al.,* 2016; Lee *et al.,* 2022; Berry *et al.,* 2022). In

Drosophila melanogaster testes (Figure 1 B), over 500 genes express transcripts cleaved at a distal

PAS in proliferating spermatogonia but processed at a more proximal PAS in differentiating

spermatocytes, resulting in production of mRNA isoforms with much shorter 3'UTRs (Figure 1 C).

Spermatogenesis in *Drosophila* starts with an asymmetric cell division that causes one stem cell

daughter to detach from hub cells, stop self-renewing and start mitotic proliferation, founding a

line of interconnected spermatogonia enclosed in two cysts cells. After 4 rounds of mitosis the

spermatogonia in the cyst start differentiation, beginning with pre meiotic S phase, and becoming

spermatocytes. After roughly 3.5 days of meiotic prophase, during which the germ cells grow 25

times in volume, the spermatocytes undergo meiosis I and II, producing a cyst of 64 haploid

spermatids. These then undertake the dramatic morphological changes that lead to production

of mature and functional sperm (Figure 1 B). The stage specific 3'UTR shortening due to APA

observed in spermatocytes compared to spermatogonia (Figure 1 C) resulted in a dramatic shift

in the proteins expressed, with transcripts from at least 200 genes switching from co-migrating

with polysomes in spermatogonia to not associated with ribosomes in young spermatocytes. In

addition, transcripts from another 50 genes were not translated in spermatogonia but associated

with polysomes in spermatocytes (Berry *et al.,* 2022). Thus, the developmental program(s) that

specify whether nascent transcripts are cleaved at proximal 3'UTR sites in spermatocytes rather

than the distal sites utilized in spermatogonia can switch expression of many proteins from on in

spermatogonia to off in spermatocytes or vice versa, with potential impacts on cell state.

Here we show that cell type specific production of transcripts with short 3'UTRs from

specific genes due to APA in spermatocytes is regulated by one of the 4 complexes involved in 3'

end cleavage and polyadenylation. We found that the two components of CFII, *PCF11* and

crowded by cid (cbc), are up regulated in spermatocytes compared to spermatogonia. When

levels of either *PCF11* or *cbc* were knocked down (KD) in spermatocytes by cell type specific RNAi,

over 230 genes (41%) for which nascent transcripts would normally be cleaved at a proximal PAS

in spermatocytes resumed producing transcripts cleaved at the distal PAS, utilized in

spermatogonia. Lowering protein levels of components of the cleavage machinery associated

complexes CFI or CstF had much less effect on APA at these loci, suggesting that CFII plays a special

role in regulating APA during this differentiation sequence. The PCF11 and Cbc proteins interact

structurally: immunoprecipitation of Cbc from testis extracts brought with it a >250KD form of

PCF11. Reduced expression of *PCF11* protein in spermatocytes due to RNAi resulted in

concomitant loss of Cbc, suggesting that Cbc protein was not stable without its binding partner

in CFII. However, the reciprocal was not true. Plentiful PCF11 protein persisted in the nuclei of

spermatocytes in which expression of *cbc* had been knocked down by RNAi. Strikingly, *PCF11*

mRNA is broadly expressed in many cell types in adult flies, while *cbc* expression is low in most

adult tissues but up regulated in spermatocytes. These observations suggest a model where a

developmentally regulated increase in expression of *cbc* in spermatocytes leads to greater activity

of cleavage factor CFII. This in turn facilitates recognition of an upstream PAS, and consequent

3'UTR shortening by APA of nascent transcripts from specific genes, indicating CFII as a key player

in developmental regulation of alternative cleavage and polyadenylation.

Results

Specific cleavage factor proteins are upregulated in spermatocytes.

Immunostaining of whole mount *Drosophila* testes revealed that some but not all cleavage factor proteins are up-regulated in spermatocytes (Figure 1 D-J). Relative expression levels of six protein components of various sub-complexes of the cleavage factor machinery were measured by immunostaining using available antibodies against endogenous proteins (for CstF50, CstF64, PABP2 and PCF11), against HA (for a CRISPR *Drosophila* line tagged with HA at the endogenous *cbc* locus by CRISPR, *HA:cbc*), or against GFP (for a *Drosophila* line carrying a Fosmid based transgenic copy of CPSF6 tagged with sfGFP, (Sarov *et al.* 2016)). In comparing nuclear signal in spermatogonia and spermatocytes, each measurement was divided by the area of the corresponding nucleus to account for the great difference in nuclear size between the two cell types. Immunofluorescence signal from antibodies recognizing two members of the CstF complex, CstF50 (Figure 1 E) and CstF64 (Figure 1 F), decreased slightly or remained almost the same in nuclei in spermatocytes compared to spermatogonia (quantified in Figure 1 D). In contrast, immunofluorescence signal from antibodies recognizing PABP2 (Figure 1 G), the CFI component CPSF6 (Figure 1 H) and the two members of CFII, Cbc (Figure 1 I) and PCF11 (Figure 1 J) increased in nuclei from spermatocytes compared to spermatogonia (quantified in Figure 1 D).

Down-regulation of CFII components in spermatocytes shifted APA to more distal sites.

To knock down (KD) expression of cleavage factor components in spermatocytes, females carrying a *bamGal4* expression driver coupled with *UAS>dicer2* were crossed with males carrying RNAi constructs under control of a *UAS* array with an *hsp70* promoter. *bamGal4* flies express Gal4 under control of the *bag-of-marble (bam)* promoter, which fires in mid to late spermatogonial and early spermatocyte stages. The crosses were incubated at 25°C for three days, then the progeny shifted to 29°C to facilitate effective transcriptional activation by Gal4.

Immunofluorescence staining revealed that *bamGal4* driven KD using the *VDRC103710* RNAi line lowered levels of *PCF11* protein detected in spermatocyte nuclei to a level similar to that detected in spermatogonia (Figure 2 B). Testes from *UAS>PCF11-RNAi-VDRC-103710;bamGal4* males viewed in live squash preparations by phase contrast microscopy had

plentiful spermatocytes but few elongating spermatid bundles (Figure 2 E). An RNAi line directed
against a different region of the PCF11 transcript, *UAS*>PCF11-RNAi-*NIG-10228R-4*, also lowered
the level of anti-PCF11 immunofluorescence signal in spermatocytes when driven using *bamGal4*
(Figure 2 C). Consistent with the less complete KD, testes from *UAS*>PCF11-RNAi-*NIG-10228R-*
4;bamGal4 males viewed by phase contrast microscopy had some elongating spermatid bundles
(Figure 2 F). In both cases, PCF11 protein was still detected in early spermatogonia and in nuclei
of the somatic cyst cells that accompany spermatocyte cysts, as expected since the *bamGal4*
driver is not expressed in these cells (Figure 2 B, C, arrows).

KD of *cbc*, the partner of PCF11 in CFII, also caused defects in spermatogenesis.
Immunofluorescence staining of testes from males heterozygous for a *cbc* allele tagged at the
endogenous locus with the HA epitope (*HA:cbc*) showed strong anti-HA signal in spermatocyte
nuclei, compared to little or no signal detected in spermatogonia. However, for two RNAi lines
directed against different regions of *cbc*, KD under control of *bamGal4* reduced the level of anti-
HA signal in spermatocytes to below the level of detection, as in spermatogonia (Figure 2 G-H-I).
In both cases, analysis of unfixed squash preparations by phase contrast microscopy showed that
the KD testes had abundant spermatocytes but lacked elongating spermatid bundles (Figure 2 K-
L).

Hybridization chain reaction fluorescence *in situ* hybridization (HCR-FISH) revealed that
KD of *PCF11* or *cbc* in spermatocytes by RNAi affected APA at both the *Kpc2* (also named
isopeptidase-T-3) and *Red* genes (Figure 3). *Kpc2* and *Red* had both been found to express
transcripts with a long 3'UTR in testis filled with spermatogonia but a much shorter 3'UTR in
testes filled with mid-stage differentiating spermatocytes (Figure 3 A, E, J). Comparing signal in
the same sample from two different sets HCR-FISH probes (SOM table 1), one recognizing the
common coding sequence and one recognizing the 3'UTR extension (Figure 3 A), revealed the
proportion of the total transcripts from a given gene (recognized by the common probe set) that
contained the 3' UTR extension (recognized by the 3'UTR extension probe). For both *Kpc2* and
Red, KD of either *PCF11* or *cbc* in spermatocytes under control of *bamGal4* dramatically increased
the level of signal from the 3'UTR extension detected in spermatocytes (Figure 3 C, D, H, I)
compared to testes from control flies raised under the same temperature regimen (Figure 3 B, G).

Quantification of the images confirmed that the ratio of fluorescence signal from probes
recognizing the long 3'UTR versus from probes against the coding sequence (CDS) for each gene
was significantly higher in the *PCF11* or *cbc* KD spermatocytes than in spermatocytes from control
testes (Figure 3 F, K).

Analysis of cleavage site usage by 3'seq from testes in which either *PCF11* or *cbc* had been
knocked down in spermatocytes by RNAi under control of *bamGal4* showed that the increase in
signal from the 3'UTR extension observed in KD spermatocytes was due to a shift from cleavage
at the proximal site normally utilized in spermatocytes to the more distal cut site normally utilized
in spermatogonia for both *Kpc2* and *Red* (Figure 4 A, B). The effect of KD of *cbc* was strongest,
while the effect of knock down of *PCF11* with the *UAS>PCF11-RNAi-NIG-10228R-4* construct was
weakest (Figure 3 A, B), consistent with the less efficient reduction in PCF11 protein levels
observed with that particular line (Figure 2 C).

3'seq data revealed that loss of function of *cbc* or *PCF11* affected APA at many loci. For
roughly half of the >500 genes previously identified by Berry *et al.* (2022) as undergoing 3'UTR
shortening by APA in spermatocytes compared to spermatogonia, KD of either *PCF11* or *cbc* in
spermatocytes resulted in increased usage of the distal cleavage site normally used in
spermatogonia (Figure 4 C). 3'seq data was analyzed via PolyA miner (Yalamanchili *et al.* 2020)
providing as input the list of pairs of proximal PAS (pPAS) and distal PAS (dPAS) from the >500
genes most changed in spermatogonia versus spermatocytes called by Berry *et al.* (2022). Orange
indicates that the frequency of the gene being cleaved at the distal PAS in a KD sample was higher
than in control testes, while blue indicates that the frequency of the gene being cleaved at the
proximal PAS in a KD sample was higher than in control testes. Noticeably, a negative PolyA index
does not indicate a shortening event but indicates a gene for which the proximal PAS was
overrepresented in a KD compared to a control sample. This could be caused either by different
expression of the gene compared to the control or by the different cell composition, as generally
KDs do not have as many elongating spermatids as a control testes. Again, KD of *cbc* in
spermatocytes had the strongest effect on the APA switch, affecting the highest number of genes.
Importantly, >94% of the genes where APA was affected (PolyA index > 0.5) by KD of *PCF11* also
showed a switch in APA with KD of *cbc,* consistent with these two CFII components acting

together in the choice of where to make the 3' end cut for transcripts from specific genes. Analysis
of RNAseq libraries from testes samples in which *cbc* or *PCF11* where knocked down under
bamGal4 revealed that in general the genes that had a PolyA index either greater than 0.5 or
smaller than negative 0.5 were not expressed at dramatically different levels in KD versus control
testes (SOM Figure 3).

3'seq also revealed that lowering the levels of the nuclear protein PABP2 in
spermatocytes also frequently affected APA (SOM Figure 4). Immunofluorescence staining
showed effective KD of PABP2 protein to below detection in spermatocytes from flies carrying
UAS>PABP2-RNAi-VDRC-106466;bamGal4, although PABP2 protein was still abundant in nuclei of
somatic cyst cells, as expected (SOM Figure 4 B). For two different RNAi lines, testes from *UAS-*
PABP2-RNAi;bamGal4 flies had plentiful spermatocytes but lacked elongating spermatid bundles
(SOM Figure 4 D-E). 3'seq revealed that in testes from *UAS>PABP2-RNAi-VDRC-106466;bamGal4*,
a substantial percentage of transcripts from the *Kpc2* and *Red* loci switched 3' end cleavage from
the proximal site normally utilized in spermatocytes to the more distal site normally utilized in
spermatogonia (Figure 4 A, B). Global analysis of the 3'seq data showed that for the >500 genes
identified previously as undergoing APA in spermatocytes compared to spermatogonia, for >86%
of the genes that switched toward the more distal PAS (PolyA index > 0.5) after KD of *PABP2*, APA
was similarly affected (PolyA index > 0.5) by KD of either *PCF11* or *cbc* (Figure 4 C, F).

In contrast, KD of any of the three components of the CstF complex (*CstF64*, *CstF50*, and
su(f)) or a component of CFI (*CPSF6*) in spermatocytes had much less effect on the observed APA
profile of mRNAs expressed in testes assessed by 3'seq (Figure 4 C). Immunofluorescence staining
with antibodies against CstF64 or CstF50 showed that levels of the proteins were substantially
reduced in spermatocytes in testes in which expression of the respective genes had been KD by
RNAi under control of *bamGal4* (SOM Figure 5). In both cases, signal remained in the nuclei of
early spermatogonia and cyst cells, as expected from the cell type specificity of the *bamGal4*
expression driver. No antibody reagents against *su(f)* were available to assess effectiveness of the
RNAi construct directed against *su(f)*. However, the effects of knocking down *su(f)* on protein
levels of CstF64 and CstF50 in spermatocytes nuclei (SOM Figure 6), and the defects in
spermatogenesis in *su(f)-RNAi;bamGal4* testes observed by phase contrast microscopy (SOM

Figure 5), suggested that the KD of *su(f)* was effective. Levels of *CstF64* protein were substantially

reduced and levels of *CstF50* protein were below the level of detection by immunofluorescence

staining in spermatocytes from males carrying an RNAi construct directed against *su(f)* expressed

under control of *bamGal4* (SOM Figure 6 C, D). KD of *CstF64* under control of *bamGal4* also

lowered the levels of CstF50 in spermatocyte nuclei to below detection (SOM Figure 6 F). Notably,

when either *su(f)* or *CstF64* were KD in spermatocytes, nuclear CstF50 staining disappeared,

although some signal remained in the cytoplasm. Together the protein expression data suggest

that components of the CstF complex may depend on each other for protein stability and/or

nuclear localization, as previously reported (Grozdanov *et al.* 2018). Consistent with the KDs

affecting function, squash preparations of testes from *UAS>su(f)-RNAI-BL-65693;bamGal4*

showed abundant spermatocytes but no elongating spermatid stages (SOM Figure 5 G). Testes

from *UAS>CstF64-RNAi-BL65987; bamGal4* males also showed defective spermatogenesis, with

plentiful spermatocytes but extremely aberrant elongation stages (SOM Figure 5 F). Surprisingly,

even though CstF50 protein was clearly strongly reduced in spermatocytes from *UAS>CstF50-*

RNAi-BL77377;bamGal4   males, spermatogenesis appeared largely normal by phase contrast

microscopy (SOM Figure 5 C).

KD of either *CstF64* or *su(f)* in spermatocytes shifted the site of 3' end formation toward

the more distal site utilized in spermatogonia at a very small percentage of the >500 genes

previously identified as subject to 3'UTR shortening in spermatocytes due to developmentally

regulated APA (Figure 4 C). These tended to be among the same set of genes where APA was

affected by KD of *cbc* or *Pcf11* (components of CFII): >88% of the genes affected by KD of *CstF64*

were also affected by KD of either *PCF11, cbc,* or both (Figure 4 G). However, the effects of loss of

function of CstF components on APA in spermatocytes was much less than the effect of loss of

CFII components: only 14 of the 235 genes where APA was strongly affected by KD of either *cbc*

or *PCF11* in spermatocytes also showed a switch toward the more distal cut site in testes where

function of *CstF64* had been knocked down in spermatocytes (Figure 4 G). All the genes affected

(PolyA index > 0.5) by *CstF64* KD were also affected either by *CstF50* or *su(f)* KDs. Interestingly, of

the few genes affected by *CstF50* KD, for almost all the choice of 3' end cut site was not affected

by *CstF64* or *su(f)* KDs (Figure 4 H). KD under control of *bamGal4* of *CPSF6*, a component of the

CFI complex, which binds nascent transcripts upstream the PAS, caused extensive defects in spermatid differentiation for the two RNAi lines tested, suggesting loss of function (SOM Figure 7). Analysis of cut sites utilized by 3' end seq indicated that for the >500 genes previously identified as undergoing APA in spermatocytes, the effect of *CPSF6* KD on APA was mild compared to KDs of *cbc* or *PCF11,* although *CPSF6* KD affected APA at many of the same genes affected by loss of function of CstF components (Figure 4 C).

De novo genome wide calling of PAS sites located on annotated 3'UTRs revealed that in addition to many of the >500 genes from Berry *et al.* (2022), KD of components of the cleavage machinery affected APA at additional loci (SOM Figure 8-9). For this analysis we only considered genes in which the most up regulated PAS and the most down regulated PAS in a KD compared to a WT sample were located in the annotated gene 3'UTR. Again, the strongest effects were detected in *cbc* KD testes, followed by *PCF11* KD and *PABP2* KD testes (SOM Figure 8), while *CstF64*, *CstF50*, *su(f)* and *CPSF6* KDs had a much milder effect on the global APA profile (SOM Figure 9). When *cbc* or *PCF11* or *PABP2* were knocked down in spermatocytes many more genes showed lengthening of 3'UTR, suggesting that expression levels of components of CFII and/or PABP2 in spermatocytes helped functional cleavage at the proximal rather than the distal site.

Of the components of the cleavage machinery tested, the 3'seq data indicated that the 3'UTR shortening of transcripts from selected genes due to APA in spermatocytes was most strongly dependent on levels of CFII (composed of *cbc* and *PCF11*), or *PABP2.* For almost half of the >500 genes previously identified as undergoing the developmentally regulated APA, lowering function of one or the other of these proteins in spermatocytes resulted in a shift toward usage of the more distal 3' end cut site typical of spermatogonia. A number of genes seemed to be more affected by KD of *cbc* than by KD of *PCF11*, for example *oaf* (Figure 4 D). However, over half of the genes previously identified as subject to 3'UTR shortening in spermatocytes compared to spermatogonia due to APA were not substantially affected by KD of any of the cleavage factor machinery we tested, *nudE* for example (Figure 4 E and SOM Figure 2).

Transcription of the CFII component *cbc* is strongly upregulated in spermatocytes.

According to Fly Base, the *PCF11* locus can give rise to 7 different RNA isoforms (Figure 5 A), grouped in 3 classes for simplicity: a short isoform (*PCF11-RC*) of 819 nucleotides predicted to encode a small protein (~10 KDa), a medium isoform (*PCF11-RB*) of 2647 nucleotides encoding a ~63 KDa protein, and a group (*PCF11-RD-RE-RF-RG-RH*) of ~6000 nucleotide transcripts predicted to encode a family of >200 KDa proteins that have subtle differences in a central region. All the transcript isoforms share the same 5'UTR but differ in their 3'UTR (as well as in exon combinations). PCF11 has a CID domain (POLII C-terminal binding domain) that is present in both the PCF11-long and the 63 KDa PCF11-RB isoforms, but only partially present in the short protein encoded by *PCF11-RC*. In addition, PCF11 have been shown to bind Cbc in mammalian and yeast cells, through a region close to the C-terminus of PCF11 (Sadowski *et al.* 2003; Schäfer *et al.* 2018). Notably, while the *PCF11-RNAi-NIG10228-R4* line targets a region common to all *PCF11* mRNA isoforms, *PCF11-RNAi-VDRC107310* targets a region present only in the *PCF11-long* isoforms. KD with *PCF11-RNAi-VDRC107310;bamGal4* in spermatocytes had a mildly stronger effect on the APA profile detected by 3'seq than the KD with *PCF11-RNAi-NIG10228R-4;bamGal4*. The *cbc* locus is simpler, with only one predicted transcript 1654 nucleotides long, encoding a 46.8 KDa predicted protein (Figure 5 A).

The two components of CFII show substantially different transcription patterns in adult flies. In the Fly Cell Atlas single nuclear RNA-seq analysis of tissues from adult flies (Li *et al.,* 2022), transcription of *PCF11* was detected at generally high levels in many cell types (Figure 5 B). In contrast, *cbc* expression was detected at substantial levels in only a small subset of nuclei from adult tissues, with the clusters showing high *cbc* transcription belonging to the testes sample (Figure 5 C, D). Plotting expression on the UMAP of snRNA-Seq data from testis plus seminal vesicle revealed that *PCF11* is transcribed in many somatic cell types in the testis, including terminal epithelia, seminal vesicle, nuclei from testis muscle cells and prominently in the somatic cyst cells that enclose germ line cysts. *PCF11* transcripts were also expressed in germ cells, with the level increasing in spermatocytes compared to spermatogonia (Figure 5 E). In contrast, although transcription of *cbc* increased substantially in spermatocytes compared to spermatogonia, little signal from *cbc* was detected in somatic cells in the testis.

HCR-FISH on testes revealed that long form transcript(s) from *PCF11*, were up-regulated

in spermatocytes compared to spermatogonia (Fig. 5 H-H'-H''). In contrast, the intermediate sized

transcript isoform *PCF11-RB*, detected with a set of probes recognizing the 3'UTR of *PCF11-RB*,

which derives from an intronic region of the *PCF11-long* isoforms so should not recognize fully

processed *PCF11-long* isoforms in the cytoplasm, was extremely low in both cell types (Fig. 5 I-I'-

I''). Single nuclear RNAseq from the Fly cell Atlas confirmed that while the collective *PCF11-long*

isoforms are highly expressed in the germline and in other cell lineages in the testis cluster

(including somatic cells and muscle cells, Figure 5 K) the *PCF11-RB* transcript is very lowly

expressed in the germline, and modestly expressed in the somatic and muscle lineages (Figure 5

L). Similar HCR-FISH using a set of probes against the *cbc* protein coding region confirmed that

expression of *cbc* transcripts was strongly up regulated in spermatocytes compared to

spermatogonia (Fig. 5 J-J'-J'').

PCF11 and Cbc co-immunoprecipitate and knock down of *PCF11* decreases Cbc protein in

spermatocytes nuclei.

Analysis of proteins expressed in wild type testes by polyacrylamide gel electrophoresis

(PAGE) followed by Western blot probed with antibodies against PCF11 showed a predominant

high molecular weight (>250 KDa) band (Figure 6 A), consistent with the expression of *PCF11-long*

mRNA isoforms detected by HCR-FISH and snRNA-seq (Figure 5 H, K). Anti-PCF11 also detected a

~70 KDa protein isoform in testis extracts, as predicted from the *PCF11-RB* mRNA isoform. The

antibody used was raised against amino acids 1-283 of *Drosophila* PCF11 (Zhang and Gilmour

2006), however no protein product corresponding to PCF11-RC was observed, despite a

seemingly high level of the *PCF11-RC* transcript isoform detected by snRNAseq of *Drosophila*

testis tissue (SOM Figure 10). Western Blots of testes from homozygous flies carrying the *HA:cbc*

allele probed with anti-HA showed expression of the expected ~50 KDa isoform (Figure 6 B).

Immunoprecipitation with anti-HA antibodies from extracts of testes from flies

homozygous for *HA:cbc* co-immunoprecipitated the >250 KDa band from the PCF11 isoforms,

consistent with the Cbc and PCF11 proteins interacting in a complex (Figure 6 C). The 70 KDa

isoform of PCF11 was not detected in the immunoprecipitation, consistent with the prediction

that the long but not the medium or short PCF11 isoforms contain the Cbc binding site. Consistent

with interaction between Cbc and PCF11, accumulation of Cbc protein in spermatocytes required

PCF11. Analysis by RNA-seq showed that when PCF11 expression was knocked down in late

spermatogonia and spermatocytes by RNAi under control of *bamGal4*, *cbc* transcript was still

expressed in *PCF11* KD testes at a similar level compared to control testes (Figure 6 D). Likewise,

in testes undergoing KD of *cbc* by RNAi under control of *bamGal4*, transcripts from the *PCF11*

locus were still expressed (Figure 6 E). However, the situation was different at the level of protein

accumulation in spermatocytes. Simultaneous immunostaining of testis from *HA:cbc/+;bamGal4*

males with anti-PCF11 and anti-HA antibodies showed both proteins present in spermatocyte

nuclei from control males (Figure 6 F). However, for two different RNAi lines directed against

PCF11, KD of *PCF11* under control of *bamGal4* resulted in loss of the signal from HA:Cbc (Figure

6 G'-H', quantified in K). In contrast, KD of *HA:*cbc by RNAi under control of *bamGal4* did not cause

loss of signal from anti-PCF11 (Figure 6 I''-J'', quantified in L).

Discussion

Our results indicate that cell type specific differential expression of components of the

Cleavage Factor complex II (CFII) is a key developmental mechanism regulating alternative

polyadenylation as cells switch from proliferation to onset of differentiation in an adult stem cell

lineage. Cell type specific knock down (KD) experiments suggest that the developmentally

specified increase in expression of PCF11 and Cbc, the two components of CFII, is important for

the shift to 3' end formation at a more proximal cleavage site in spermatocytes compared to

spermatogonia at over 200 loci during *Drosophila* spermatogenesis. Knock down of several other

cleavage factors had much milder effects on shifting 3' end cleavage from the proximal site

normally utilized in spermatocytes to the more distal site normally utilized in spermatogonia,

singling out CFII for a key role in the developmentally regulated 3'UTR shortening due to APA in

normal *Drosophila* spermatocytes.

Of the two components of CFII, reducing levels of Cbc expressed in spermatocytes had the

most dramatic effect on PAS choice, effecting more than 40% of the >500 genes previously

identified as subject to APA leading to 3' end cleavage at a more proximal PAS in spermatocytes.

Expression of both *PCF11* and *cbc* is developmentally regulated, with mRNA and protein levels

considerably higher in spermatocytes compared to spermatogonia. Analysis of the single nuclear

RNAseq data from the Fly Cell Atlas project (Li *et al.,* 2022) revealed that expression of *cbc* is much

more cell type specific, being upregulated strongly in spermatocytes compared to other cell types

and tissues. In contrast, *PCF11* was expressed much more broadly, with transcription detected in

every fly tissue analyzed. This, plus our finding that knock down of *PCF11* in spermatocytes

resulted in failure of Cbc protein to accumulate, while PCF11 protein was still present in

spermatocytes where *cbc* was knocked down, suggest a model where PCF11 carries out functions

independent of Cbc, but when Cbc protein is upregulated in spermatocytes it binds some of the

PCF11 and recruits it into CFII, where it can bind and activate the core 3' end cleavage machinery

to cut at a more proximal PAS site in specific transcripts that are normally skipped in

spermatogonia. Although previous studies involving knock down of CLP1 (the mammalian

homolog of Cbc) in mammalian cells did not show remarkable effect on APA (Li *et al.* 2015;

Ogorodnikov *et al.* 2018), our findings raise the possibility that activity of Cbc in the choice of 3'

end cut site may be more important in some cell types than in others, and on certain specific

transcripts rather than in general.

Consistent with the model that PCF11 may carry out functions independent of CFII, PCF11

has been found to bind the CTD of RNA POLII *in vitro* and in yeast (Sadowski *et al.* 2003; Zhang

and Gilmour 2006), where it acts as a termination factor (Guéguéniat *et al.* 2017; Schäfer *et al.*

2018; Kamieniarz-Gdula *et al.* 2019). CHIP-seq of POLII in control versus *PCF11* knock down

human cells showed that POLII fails to terminate efficiently when PCF11 is reduced (Kamieniarz-

Gdula *et al.* 2019). Accordingly, downregulating *PCF11* expression levels has been shown to

induce 3'UTR lengthening in cancer cells and zebrafish (Li *et al.* 2015; Ogorodnikov *et al.* 2018;

Kamieniarz-Gdula *et al.* 2019). However, if/how skipping cleavage sites might be linked with POLII

termination, and what other functions PCF11 might carry out in association with RNA POLII still

need to be elucidated.

Cbc is a much smaller protein than PCF11, does not have a recognized RNA binding

domain, and immunoprecipitation of cbc *in vitro* failed to detect bound RNA (Schäfer *et al.* 2018).

On the other hand, PCF11 has been shown to bind RNA *in vitro*, with a preference for G-rich

regions. Our findings raise the possibility that Cbc might act as a developmentally regulated factor

that, when upregulated in certain cell types might bind to PCF11 to form CFII, utilize the ability of

PCF11 to recognize RNA, and modulate PCF11 activity to enhance 3' end cleavage at specific sites.

Developmentally regulated switches in the site at which nascent mRNAs are terminated,

such as those we have observed dependent on components of CFII, may serve as a mechanism

to quicky change the menu or levels of proteins expressed as cells progress through sequential

steps in differentiation. By effecting the length and content of 3'UTRs, APA can change whether

mRNAs are translated, blocked from translation, differentially localized, stabilized, or degraded.

Since APA is a co-transcriptional process that does not require opening or closing of chromatin or

re-initiation of transcription, it may provide a more rapid way to turn expression of some proteins

on and other proteins off, so changing cell state and canalizing cell fate decisions. As such,

understanding how the alternative choices in 3' end cut site are regulated is key for understanding

developmental regulation of cell states.

Materials and Methods

Fly Strains and husbandry.

Flies were grown on standard molasses media and kept at 25 °C, 29 °C or room temperature as indicated. For RNAi induced knockdowns (KD), fly crosses (females *Bamgal4* driver X males RNAi line) were kept at 25 °C for three days. Adults were then shifted to a new media bottle and kept at 25 °C while eggs and early larvae were shifted to 29 °C to increase KD efficiency. To further increase KD penetrance, we used a Gal4 driver that also expressed *UAS > dicer2*. Fly lines used in this study: *W1118* (BL #3605), *Bam Gal4* (Chen and McKearin 2003), *CPSF6:GFP* (VDRC #318105), *HA:Cbc* (N-terminal CRISPR tag, kindly gifted from the Wang lab, (Wu *et al.* 2021)), *PCF11*-RNAi_1 (VDRC #103710) *PCF11*_RNAi_2 (NIG #10228R4), *cbc*-RNAi_1 (BL #77426), *cbc*-RNAi_2 (VDRC #100686), *CPSF6*-RNAi_1 (BL #34804), *CPSF6*-RNAi_2 (VDRC #107147), *CstF64*-RNAi_1 (BL #65987), *CstF50*-RNAi_1 (BL #77377), *PABP2*-RNAi_1 (VDRC #106466), *PABP2*-RNAi_2 (VDRC #33499), *su(f)*-RNAi_1 (BL 55693).

Immunofluorescence (IF) staining and signal quantification.

Testes from 0-3 days old were removed by dissecting in 1X PBS on a cyclops dissecting dish at room temperature over 30 minutes or less, then testes were immediately fixed in an Eppendorf with 5% formaldehyde for 30 minutes, with rocking at room temperature. If needed, fixed samples were stored in 1X PBS at 4 °C for up to one week. Testes were then permeabilized in 0.3% Triton-X-100 and 0.6% sodium deoxycholate for 30 minutes at room temperature and rinsed twice in 1X PBS + 0.05% Tween-20 (PBT), blocked for 30 minutes in 1X Western Blocking Reagent (WBR, Roche #) in PBT at room temperature, then incubated with primary antibody mixed in 1X WBR+PBT overnight at 4 °C. After primary antibody incubation, testes were rinsed twice in PBT, blocked again for 30 minutes in 1X WBR+PBT at room temperature, then incubated with secondary antibody mixed in 1X WBR+PBT for 2 hours at room temperature in a dark container. Testes were then rinsed twice in PBT and mounted on a glass slide in 10 uL of ProLong diamond anti-fade with DAPI (ThermoFisher, CAT #P36962). Primary antibodies used: guinea pig anti-CstF50 (1:300, kindly gifted from John T. Lis (Ni *et al.* 2008)), rabbit anti-PABP2 (1:500, kindly gifted

from Martine Simonelig (Benoit *et al.* 1999)), goat anti-Vasa (1:200, Santa Cruz Biotechnology sc-

26877, discontinued), rat anti-VASA (1:10 for supernatant or 1:100 for the concentrated version,

both from DSHB, antibody ID 760351) rabbit anti-CstF64 (1:400, kindly gifted from Zbigniew

Dominski (Skrajna *et al.* 2018)), rabbit anti-PCF11 (1:200, kindly gifted from David Scott Gilmour

(Zhang and Gilmour 2006)), rabbit anti-HA (1:500, Cell Signalling Technologies, #3724), mouse

anti-HA (1:500, CAT???), chicken anti-GFP (1:100, cat???). Secondary antibodies used: donkey or

goat anti-rabbit, anti-mouse, anti-guineapig, anti-chicken, anti-rat and anti-goat, all used at 1:500

and conjugated with either Alexa 488, Alexa 568, Alexa 594 or Alexa 647 depending on the

combination of antibodies used in a given experiment (all from the ThermoFisher catalog).

Imaging of IF experiments was performed on a Leica SP8 Confocal microscope. Protein levels were

quantified in Fiji by manually delineating the cell nucleui and applying the formula:

Integrated_Density – (Area X Background_Mean).

HCR^{TM}-RNA-FISH.

Testes were dissected from 0-3 days old male flies in 1X PBS on a cyclops dissecting dish at room

temperature over a maximum of 30 minutes, then fixed and permeabilized as done for

immunofluorescence. The protocol for "generic sample in solution" from the Molecular

Instruments website was followed, hybridization buffer, wash buffer, amplification buffer and

hairpins are all from Molecular Instruments (HCR™ RNA-FISH Bundle). After permeabilization,

samples were incubated with 200 uL of probe hybridization buffer for 30 minutes at 37 °C. Probe

mix were prepared mixing 3.2 uL of 1 uM probes (one per gene or transcript) in 200 uL of probe

hybridization buffer. Hybridization buffer was removed from samples and probe mix added.

Samples were incubated overnight in probe mix at 37 °C. The following day samples were washed

4 times for 15 minutes each in 200 uL of probe wash buffer at 37 °C and then washed 3 times 5

minutes each in 5X SSCT (5X sodium chloride sodium citrate (SSC) and 0.1% Tween-20 in H$_2$O) at

room temperature. Samples were incubated in 500 uL of amplification buffer for 30 minutes at

room temperature. In the meantime hairpin mix was prepared: 30 pmol per sample (5 uL) of

harpin h1 and hairpin h2 (each fluorophore has its own harpin combination, in our case since we

always imaged 2 transcripts at the same time, we used 2 fluorophores and therefore 4 hairpins)

were heated at 95 °C for 90 seconds in a thermocycler in the dark and cool down at room

temperature in a drawer for 30 minutes (each hairpin in heat up and cooled in a separate tube).

Hairpin mix was prepared mixing 5 uL of each harpin to 250 uL of amplification buffer, and

samples were incubated in harpin mix at room temperature in the dark overnight. The following

day samples were washed with 500 uL 5X SSCT 2 times for 5 minutes each, then 2 times 30

minutes each and finally 1 time 5 minutes, always in the dark and at room temperature. Samples

were then mounted on a glass slide in 10 uL of ProLong diamond anti-fade with DAPI

(ThermoFisher, CAT #P36962). Probes were designed as in Bedbrook *et al.* 2023 and ordered from

Integrated DNA technologies as Oligo Pools (oPools™ 50 pmol/oligo). Harpins H1 and H2 were

ordered conjugated with either 488 or 546 fluorophores from Molecular Instruments. Probes

sequences are listed in supplementary material. Imaging was performed on a Leica SP8 confocal

microscope. For each testis, 5 areas from spermatocytes cytoplasm were quantified on Fiji from

both probe channels, subtracted by the background and then the ratio was calculated. The

following formula was used: *(Area_Mean_p2 – Back_Mean_p2) / (Area_Mean_p1 –*

Back_Mean_p1) where p2 is the probes set recognizing 3'UTR extension while p1 is the probe set

recognizing the gene CDS. Area's size varied between pictures but was constant between the 5

measurements from the same image.

3' end sequencing and analysis.

Testes were dissected from 0-3 days old male flies in S2 media in batches of 50-80 flies on a

cyclops dissecting dish at room temperature over a maximum of 1 hour per batch. Testes were

then immediately transfer in an Eppendorf containing 1X PBS, 1X PBS was removed and the testes

were snap-frozen in liquid nitrogen and stored at -80 °C. RNA was extracted from ~150 testes pairs

using the RNAeasy Mini kit (Qiagen, 74104) following the manufacturer directions with minor

adjustments. Frozen tissue was dissociated using a 1mL syringe with a 27G needle in RNAeasy

Mini kit lysis buffer supplied with 1:100 β-mercaptoethanol as suggested by the kit manual.

Quality of extracted RNA was assessed via Bioanalyzer, and the concentration measured by

Nanodrop and Qubit RNA HS Assay Kit (Life Technologies, Q32852). 3' end RNAseq libraries were

prepared using the Quant Seq (FWD) kit from Lexogen using 500 ng of total extracted RNA per

replicate. DNA libraries quality was checked via bioanalyzer and concentration measured with

Qubit dsDNA HS Assay Kit (Life Technologies, Q32854). Sequencing was performed by the

Stanford Genomics facility on an Illumina NextSeq 500 platform as in Berry *et al.* 2022. Each

biological sample was performed in duplicates. Raw reads were trimmed to remove adaptor

sequences and aligned to the *Drosophila* dm6 genome using bowtie 0.12.8 with a value of e=5000

to allow for the expected mismatched in the PolyA tail sequence, as in Berry *et al.* 2022.

Quantification of PAS usage of the ~500 genes identified in Berry *et al.* 2022 was performed using

PolyA miner (Yalamanchili *et al.* 2020), using as input a bed file containing the proximal and distal

PAS of the ~500 that are cleaved at a proximal PAS in spermatocytes compared to spermatogonia.

We generated this list by merging the proximal and distal cleavage sites from 3 different lists: bam^{-}

$^{/-}$ vs. 32H (very early spermatocytes), bam$^{-/-}$ vs. 48H (polar spermatocytes) and bam$^{-/-}$ vs 72H (mid-

stage maturing spermatocytes) to consider various stages of spermatocytes (32H, 48H and 72H

indicate hours after heat shock, which causes a pulse of bam expression in bam$^{-/-}$ flies, inducing

differentiation, as explained in Kim *et al.* (2017). As in this lists the coordinates of the cleavage

sites did not match perfectly, some genes had more than 2 cleavage sites considered. However,

PolyA miner was built to consider genes with multiple polyA sites.

***De novo* PAS calling and APA analysis**.

For analysis of genome wide PAS usage changes between WT and KD samples, we adapted the

analysis pipeline from Zeng *et al.* 2024. The sequenced 3' end-seq libraries were adapter-

trimmed, and quality filtered using bbduk. The filtered reads were mapped to dm6 *Drosophila*

genome using STAR and then extracted for unique-mapping reads using Samtools. If a read was

mapped to a region that is immediately upstream of six consecutive As or of a 10-bp region with

at least 60% of As, it was considered as a mis-primed read and removed. The resulting filtered

reads were used for *de novo* PAS calling using LAPA[61] that used the default setting, except

requiring a replication rate cutoff at 0.95. For each gene, lengthening or shortening was quantified

by considering the ratio between the 2 highest changing PAS: the PAS that is more up-regulated

in the KD and the PAS that is more down-regulated in the KD. For this analysis we considered only

genes for which both PASs located in the 3'UTR.

RNAseq and analysis.

The same total RNA extracted for 3'seq was also used for preparation of short read RNAseq library.

1 ug of total extracted RNA was used to prepare libraries following the protocol from the NEBNext

Ultra II Direction RNA Library Prep Kit for Illumina. Each biological sample was performed in

duplicate. Quality and concentration of libraries were measured via Bioanalyzer and Qubit, as for

3'seq libraries. Sequencing was performed by Novagene on a NOVAseq, PE 150 Illumina Platform.

Raw reads were trimmed to remove the Illumina adapters using trim_galore, then aligned to the

Drosophila dm6 genome using STAR. Differential expression analysis was performed using the

package DEseq2.

Fly Cell Atlas single nuclear RNAseq analysis

Using the Fly Cell Atlas "Testis" polydT-primed 10x single-nuclear RNA sequencing dataset (Li *et*

al. 2023), we aggregated all reads from nuclei in each of the 111 Leiden 6.0 clusters (as in Berry

et al. 2022). For each cluster, reads that uniquely mapped to one of the isoform specific *PCF11*

3'UTRs were counted and normalized by the length of that 3'UTR as well as the total number of

reads derived from that cluster. This produced an isoform-specific rpkm value for each Leiden 6.0

cluster that was then log-normalized and plotted using Seurat. gene expression and cell type

annotation plots across all the adult *Drosophila* tissues analyzed in the Fly Cell Atlas project were

generated in Seurat using published code modified to visualize relevant genes and cell identities

of interest (Raz *et al.* 2023).

Western Blot (WB).

Testes from 20-50 0-3 days old male flies were quickly dissected into 1X PBS on a cyclops

dissecting dish at room temperature, transferred to an Eppendorf and snap froze in liquid

nitrogen and stored at -80 °C until needed. Frozen tissue was lysed in 40 uL of cold Lysis Buffer

(either RIPA buffer [50 mM TRIS pH 7.5, 250 mM NaCl, 0.1% SDS, 0.5% Sodium deoxycholate to

which was added one tablet per 50 uL of protease inhibitor cocktail (cOmplete ULTRA Tablets,

Mini, EDTA-free, EASY pack, Roche, REF 05 892 791 001)] or JUME Buffer [10 mM MOPS, 10 mM

EDTA, 8M UREA, 1% SDS]). Tissue was dissociated using a 1 mL syringe with a 27G needle, then

incubated rocking at 4 °C for 20 minutes. Following incubation, lysate was centrifuged at max

speed for 5 minutes at 4 degrees. Supernatant was transfered to a new tube, mixed with 4X Laemli

Buffer with β-mercaptoethanol, boiled for 7 minutes at 85 °C, then run on a Mini-PROTEAN TGX

Precast Protein Gel (4–15% Mini-PROTEAN® TGX™ Precast Protein Gels, 10-well, 50 µl #4561084).

The separated proteins were transferred from the gel to a PVDF membrane (BIO-RAD, Cat.#

1620177) overnight at 4 °C. The following day, the membrane was blocked overnight at 4 °C in 5%

powder nonfat milk dissolved in PBST (1X PBS with 0.1% Tween-20). After blocking, the

membrane was incubated with primary antibody mix in 2% milk for 1 hour at room temperature,

washed 3 times 20 minutes each with PBST at room temperature, incubated with secondary

antibody mix in 2% powder nonfat milk for one hour at room temperature and washed again 3

times 20 minutes each at room temperature. Stainings were then revealed using Revvity Health

Sciences Inc Western Lightning Plus-ECL, Enhanced Chemiluminescence Substrate (PerkinElmer).

Primary antibodies used for Western blots: Rabbit anti-HA (1:5000 or 1:1000, Cell Signalling

Technologies, #3724), Rabbit anti-PCF11 (1:10.000, kindly gifted from David Scott Gilmour (Zhang

and Gilmour 2006)). Secondary antibodies used for Western blots: Peroxidase AffiniPure™ Goat

Anti-Rabbit IgG (H+L) (1;10.000, Jackson ImmunoResearch AB_2313567).

Immunoprecipitation (IP).

For Immunoprecipitation (IP) and co-IP experiments, testes were dissected into 1X PBS on a

cyclops dissecting dish at room temperature from batches of 50 flies for a maximum of 30 minutes

per batch, snap-frozen in liquid nitrogen and stored at -80 °C. For one sample ~100 pairs of testes

were used. The day before the actual IP, 2 uL per sample of rabbit anti-HA antibodies (Cell

Signalling Technologies, #3724) were crosslinked to magnetic beads as described in Baker *et al.*

2023. The day of the experiment, beads crosslinked with antibody were rinsed 1 time in 200 mM

glycine pH 2.5 and washed once for 5 minutes in 200 mM glycine pH 2.5 to remove non-

crosslinked antibodies, then blocked for 1 hour in 10% BSA in 50 mM Tris pH 8, and re-suspended

in 20 uL of RIPA buffer and stored at 4 °C while cell lysate was prepared. Frozen tissue was

dissociated in RIPA buffer using a 1mL syringe and a 27G needle as in WB experiments. The lysate

was rocked at 4 °C for 20 minutes, then centrifuged 5 minutes at max speed in an Eppendorf tube

at 4 °C to separate lysate from tissue debris. Supernatant was transferred to a new Eppendorf tube and was incubated 45 minutes at 4 °C with beads with no antibody. Tubes containing lysate and beads were placed on a magnetic rack, 40 uL of lysate were saved as input sample and the remaining volume (160 uL) was pipetted into the tubes containing beads crosslinked to antibody and left incubating for 4 hours at 4 °C rocking, volume was brought back to 220 uL by adding 40 uL of cold RIPA buffer. Since we noticed that HA:Cbc in cell lysate tends to form aggregates (see SOM Figure 11), we precipitated the input sample and resuspended in JUME Buffer. 4X Laemli Sample Buffer (BIO-RAD #1610747) mixed with β-mercaptoethanol was added to the input sample, incubated 7 minutes at 85 °C and frozen. After 4 hours rocking at 4 °C, IP samples were placed on a magnet to separate beads from lysate, 40 uL of the supernatant were saved as supernatant sample, mixed with 4X Laemli Sample Buffer with β-mercaptoethanol incubated 7 minutes at 85 °C and frozen; the rest of supernatant was tossed. The beads with attached protein were immediately resuspended in RIPA buffer and washed with RIPA buffer 2 times for 5 minutes rocking at room temperature. Tubes were placed on a magnet to separate beads from RIPA buffer, then RIPA buffer was removed and beads were resuspended in 40 uL Elution Buffer (1X cOmplete ULTRA Tablets, 1% SDS, 10 mM EDTA, 50 mM Tris pH 8), vortexed, and incubated for 30 minutes at 70 °C with frequent (every 5 minutes) vortexing or shaking. Tubes were then centrifuged at max speed for 10 seconds at room temperature, placed on the magnet to separate beads from the eluate. 40 uL of eluate were moved to another tube (IP-sample) to which was added 4X Laemli Sample Buffer with β-mercaptoethanol, incubated 7 minutes at 85 °C and frozen. The beads were then mixed with 40 uL of elution buffer, mixed with 4X Laemli Sample Buffer with β-mercaptoethanol, incubated 7 minutes at 85 °C and frozen. Samples were then run on a Mini-PROTEAN TGX Precast Protein Gel and the protocol for WB above followed.

If the same membrane needed to be re-stained with another antibody, the membrane was stripped by incubation at 55 °C for 30 minutes in 62.5 mM Tris pH 6.8, 2% SDS and 100 mM β-mercaptoethanol, washed 3 times for 10 minutes each at room temperature in PBST and either stored in 1X PBS or blocked again before being re-stained as above.

Protein precipitation.

4 volumes of ice-cold methanol are added to 1 volume of cell lysate, vortexed, and 1 volume of

chloroform added. The sample was vortexed again, mixed with 3 volumes of H_2O, vortexed and

centrifuged in 1.5 mL Eppendorf tubes at 4 ^oC max speed for 5 minutes. After centrifugation, 2

liquid layers formed, with proteins at the interface. Upper liquid layer was removed, and 3

volumes of ice-cold methanol added to the sample. After a quick vortex, the sample was

centrifuged at 4 ^oC max speed for 10 minutes to pellet the proteins. After centrifugation all liquid

was removed, and the white protein pellet was air-dried and resuspended in 40 uL of JUME buffer.

    Figure Legends

Figure 1. Expression of Cleavage Factors during *Drosophila* spermatogenesis. (A) Schematic

representation of the cleavage machinery, adapted from Gallicchio *et al.* (2023). Grey: CPSF =

Cleavage and Polyadenylation Factor complex. Orange: CstF = Cleavage and Stimulation Factor

complex. Blue: CFI = Cleavage Factor I. Green: CFII = Cleavage Factor II. UE: Upstream element.

DE: Downstream element. PAS = Polyadenylation signal (B) Schematic representation of a WT

Drosophila testes showing key stages of germ cell differentiation. (C) Diagram of the

developmentally regulated APA in the *Drosophila* testis. (D) Quantification of relative

immunofluorescence signal per nuclear area from (E-J). Each dot corresponds to a single nucleus.

For each testis 20 spermatogonial nuclei and 20 spermatocyte nuclei were quantified, N=5 testes

were quantified per staining, for a total of 100 spermatogonia and 100 spermatocytes. Statistical

significance was measured using the Kolmogorov-Smirnov t-test: (****) p-value < 0.0001; (*) p =

0.0366. (E-J) Immunofluorescence images of testis apical regions. (Magenta) anti-Vasa marks

germ cells. (Cyan) cleavage factor proteins: (E-G, J) Wild type (*w1118*) testes stained with anti-

(E) CstF50, (F) CstF64, (G) PABP2 and (J) PCF11. (H) Testes from *CPSF6:GFP* transgenic flies stained

with anti-GFP, (I) Testes from *HA:cbc/HA:cbc* flies stained with anti-HA. Dotted yellow line: border

between proliferating spermatogonia (SG) and differentiating spermatocytes (SC). Scale bars = 50

uM.

Figure 2. Knock down of *PCF11* and *cbc* in spermatocytes. (A-B-C) Apical testes tip stained with

anti-PCF11. (A) Control: *bamGal4*, (B) *PCF11-RNAi-VDRC-103710;bamGal4* and (C) *PCF11-RNAi*

NIG-10228R-4;bamGal4. (D-E-F) Phase contrast microscopy images of whole mount live testes

from *Drosophila* lines in (A-B-C) respectively. (G-H-I) Apical testes tip stained with anti-HA. (G)

Control *HA:cbc/Cyo;bamGal4*, (H) *HA:cbc/+,cbc-RNAi-BL-77426;bamGal4* and (I) *HA:cbc/+,cbc-*

RNAi-VDRC-100686;bamGal4. (J-K-I) Phase contrast microscopy images of whole mount live

testes. (J) Control: *bamGal4*, (K) *cbc-RNAi-BL-77426;bamGal4* and (L) *cbc-RNAi-VDRC-*

100686;bamGal4. Arrows indicate somatic cysts cells. Scale bars are 50 uM (A-B-C-G-H-I) and 200

uM (D-E-F-J-K-L). Note that in all the immunofluorescence experiments on *HA:cbc/+* flies, the

level of immunofluorescence signal with anti-HA in spermatogonia was under the detection limit

of the antibody, as the flies used in this experiment were heterozygous for the *HA:cbc/+*, while

the testes quantified in Fig.1-I were homozygous *HA:cbc/HA:cbc* (for comparison between

homozygous and heterozygous testes see SOM Figure 1).

Figure 3. Knockdown of either PCF11 or cbc reduces cleavage of the proximal PAS in

spermatocytes. (A) Schematic representation of a gene body (either *Kpc2* or *Red*) to illustrate

design of the two probe sets. (B-C-D, B''-C''-D'') Confocal microscopy images of (B, B'') *bamGal4*

(control), (C, C'') *PCF11-RNAi-VDRC-103710;bamGal4* and (D, D'') *cbc-RNAi-BL-77426;bamGal4*

testes stained against probes recognizing the CDS of the gene *Kpc2* (in cyan) and DAPI (B-C-D),

and against probes recognizing the 3'UTR extension of the gene *Kpc2* (in orange) (B''-C''-D''). (B'-

C'-D', B'''-C'''-D''') zooms in spermatocytes of corresponding regions from (B-C-D, B''-C''-D'')

respectively. (E) 3'seq pile reads for the gene *Kpc2* from a sample enriched for SG = spermatogonia

(16H PHS = post heat shock, Kim *et al.* 2017; Berry *et al.* 2022) and a sample enriched for

spermatocytes SC = spermatocytes (72H PHS) to show the APA switch from the distal PAS (dPAS)

used in spermatogonia and the proximal PAS (pPAS) used in spermatocytes. (F) Quantification of

fluorescence ratio between fluorescence coming from the probes recognizing the *Kpc2* 3'UTR

extension (orange) and fluorescence recognizing the *Kpc2* CDS (cyan) in spermatocytes. For each

genotype N=3 testes were quantified. (G-K) Analogous to (A-F) for the gene *Red*. Scale bars are

150 uM (B-B''-C-C''-D-D''-G-G''-H-H''-I-I'') and 50 uM (B'-B'''-C'-C'''-D'-D'''-G'-G'''-H'-H'''-I'-I'''). p

values are (****) = p<0.0001 and (***) = p = 0.0002. For statistical analysis we used the Welch's

t-test.

Figure 4. 3' end RNAseq libraries comparing effects on APA of knockdowns in spermatocytes of

different components of the cleavage machinery. (A-B) Integrative Genomics Viewer (IGV) plots

of 3' end RNA sequencing libraries for the genes (A) *Kpc2* and (B) *Red*. Regions of peaks that

change in color show mismatches between the PolyA tail and the genome to highlight the 3' end

of the mRNA. (C) Heatmap of PolyA index quantified with PolyA Miner across 3'seq libraries. Each

row represents one of the >500 genes that switch to a proximal PAS in WT spermatocytes

compared to spermatogonia from Berry *et al.* (2022). Each column represents a component of

the cleavage machinery that was KD in spermatocytes. The interval chosen to plot the PolyA index

(-1.5,1.5) does not represent the maximum and minimum PolyA index observed. In this heatmap

genes that showed a statistically non-significant PolyA index in all the samples or a negative PolyA

index in all the samples were removed, and in all other cases PolyA indexes that were non-

significant were put equal to zero (another heatmap with these genes showed is reported in SOM

Figure 2) (D-E) Integrative Genomics Viewer (IGV) plots of 3' end RNA sequencing libraries for the

genes (D) *oaf* and (E) *nudE*. (F-G-H) Venn diagrams representing number of genes whose polyA

index is > 0.5 in *cbc* KD libraries (dark green), *PCF11* VDRC KD libraries (light green), *PABP2* KD

libraries (red) and components of CstF complex (orange).

Figure 5. *PCF11* and *cbc* transcripts are both up-regulated in spermatocytes with *cbc* particularly

highly expressed in testis compared to other tissues. (A) Diagram adapted from FlyBase of *PCF11*

and *cbc* loci. Light blue boxes represent 5'UTRs, grey boxes represent exons and pink boxes

represent 3'UTRs, while introns are represented with a straight black line. In yellow are

represented the transcripts regions against which we designed HCR-FISH probes for RNA *in situ*.

Green and dark green boxes on *PCF11* locus represent the CID domain and a predicted Cbc

binding domain respectively. (B-G) Scope diagrams of Fly Cell Atlas single nuclear RNAseq (Li *et*

al. 2023) data showing: (B) *PCF11* and (C) *cbc* expression across all the analyzed tissues, (D) Umap

highlighting the testis cluster across all analyzed tissues (in purple), (E) *PCF11* and (F) *cbc*

expression in the testis sample, (G) Umap highlight germ-line cell clusters in the testis sample (in

orange). (H-H'-H'') Confocal microscopy of WT (*w1118*) testes stained with probes against *PCF11*

long isoforms (white) and DAPI (blue). (I-I'-I'') Confocal microscopy of WT (*w1118*) testes stained

with probes against *PCF11*-RB isoform (white) and DAPI (blue). (J-J'-J'') Confocal microscopy of

WT (*w1118*) testes stained with probes against *cbc* (white) and DAPI (blue). In (H-I-J) the boxes

highlight a region of spermatogonia zoomed in (H'-I'-J') and a region of spermatocytes zoomed

in (H''-I''-J''). (K-L) Umaps from Fly Cell Atlas single nuclear data reporting (K) *PCF11-long* and (L)

PCF11-RB expression across the different clusters of the testis plus seminal vesicle sample.

Scalebars are 50 uM in (H-I-J) and 20 uM in (H'-H''-I'-I''-J'-J'').

Figure 6. *PCF11* KD causes loss of Cbc. (A) Western Blots of WT (*w1118*) testis lysate stained against endogenous PCF11. (B) Western Blots of *HA:cbc* homozygous testes lysate stained against HA. (C) Immunoprecipitation (IP) of *HA:cbc* in WT (*w1118*) and *HA:cbc* homozygous testes followed by western blots stained against endogenous PCF11 and against HA. Input = 20% of lysate before IP. (D-E) IGV plots of (D) *cbc* and (E) *PCF11* loci of RNAseq data from *bamGal4* (control) samples (in grey), *PCF11-RNAi-VDRC-103710;bamGal4* samples (magenta) and *cbc-RNAi-BL-77426;bamGal4* samples (teal). (F-J) Immunofluorescence images showing apical tips of testes stained against Vasa to mark germ cells, HA and PCF11. (F-F'-F'') *HA:cbc/cyo;bamGal4* (control), (G-G'-G'') *HA:cbc/+,PCF11-RNAi-VDRC-103710;bamGal4*, (H-H'-H'') *HA:cbc/+,PCF11-RNAi-NIG-10228R-4;bamGal4*, (I-I'-I'') *HA:cbc/+,cbc-RNAi-BL-77426;bamGal4* and (J-J'-J'') *HA:cbc/+,cbc-RNAi-VDRC-100686;bamGal4.* (K-L) Quantification of Fluorescence/Area coming from HA antibodies (teal) and from PCF11 antibodies (magenta). For each testis 10 spermatogonia and 10 spermatocytes nuclei were quantified simultaneously for PCF11 and HA fluorescence and then averaged. For each genotype N=3 testes were quantified. Statistical analysis was performed using the Welch's t-test. p-values are: *bamGal4_PCF11* = 0.0011 (**), b*amGal4_cbc* = 0.0005 (***), *PCF11-RNAi-NIG10228R-4_PCF11* = 0.0341 (*), *cbc-RNAi-BL77427_PCF11* < 0.0001 (****), *cbc-RNAi-VDRC100686_PCF11* = 0.0303 (*). Scalebars are 50 uM (F-J''). Entire gels for western blots and co-IP are reported in SOM Figure 11.

Acknowledgements

We thank all the members of the Fuller lab for suggestions and comments on this manuscript

and Dr. Nicole Martinez and Dr. Lauren Goins for their valuable feedback on the manuscript and

suggested experiments. We thank Dr. Catherine Baker and Benjamin Bolival for their help in

maintaining fly crosses. We thank Dr. Jingxun Chen from the Brunet lab at Stanford for her help

in designing HCR-FISH probes and for sharing some of the reagents, the Wang lab for the *HA:cbc*

CRISPR tag flies, the Lis lab for anti-CstF50, the Simonelig lab for anti-PABP2, the Dominski lab for

anti-CstF64 and the Gilmour lab for anti-PCF11 antibodies. We thank Daniel Mokhtari for his help

parsing the Fly Cell Atlas dataset. We thank the Cell Science Imaging facility at Stanford,

supported, in part, by Award Number 1S10OD010580-01A1 from the National Center for

Research Resources (NCRR) (The contents of this manuscript are solely the responsibility of the

authors and do not necessarily represent the official views of the NCRR or the National Institutes

of Health), particularly the CSIF facility manager Kitty Lee. We thank the Stanford Fly Media

Center as well and all the staff working in the Developmental Biology department. L.G. was

supported by an AICF (American Italian Cancer Foundation) postdoctoral fellowship (2021-2023).

This research was supported by National Institutes of Health grant R35 GM136433) and funds

from the Katherine D. McCormick and Stanley McCormick Memorial Professorship and the Reed-

Hodgson Professorship in Human Biology to M.T.F. This manuscript is dedicated to Cuchuflí.

Author Contributions

LG and MTF conceived the project and wrote the manuscript. LG designed the

experimental work and performed or supervised all experimental and computational procedures.

NRM contributed to experimental design and bioinformatic analysis. FMP contributed to

experimental design and performed molecular cloning. IN performed immunofluorescence

experiments. SS performed bioinformatic analysis of the Fly Cell Atlas data. YZ performed

bioinformatic analysis to perform *de novo* mapping of PASs. Funding was acquired by LG and MTF.

References

An, J. J., K. Gharami, G. Y. Liao, N. H. Woo, A. G. Lau *et al.*, 2008 Distinct Role of Long 3' UTR BDNF mRNA in Spine Morphology and Synaptic Plasticity in Hippocampal Neurons. Cell.

Baker, C. C., L. Gallicchio, N. R. Matias, D. F. Porter, L. Parsanian *et al.*, 2023 Cell-type-specific interacting proteins collaborate to regulate the timing of Cyclin B protein expression in male meiotic prophase. Development dev.201709.

Bedbrook, C. N., R. D. Nath, R. Nagvekar, K. Deisseroth, and A. Brunet, 2023 Rapid and precise genome engineering in a naturally short-lived vertebrate (L. Moons, D. Y. R. Stainier, L. Moons, & P. Abitua, Eds.). Elife 12: e80639.

Benoit, B., M. Simonelig, A. Nemeth, N. Aulner, U. Kühn *et al.*, 1999 The Drosophila poly(A)-binding protein II is ubiquitous throughout Drosophila development and has the same function in mRNA polyadenylation as its bovine homolog in vitro. Nucleic Acids Res. 27: 3771–3778.

Berry, C. W., G. H. Olivares, L. Gallicchio, G. Ramaswami, A. Glavic *et al.*, 2022 Developmentally regulated alternate 3' end cleavage of nascent transcripts controls dynamic changes in protein expression in an adult stem cell lineage. Genes Dev. 36: 916–935.

Chen, D., and D. M. McKearin, 2003 A discrete transcriptional silencer in the bam gene determines asymmetric division of the Drosophila germline stem cell. Development 130: 1159–1170.

Clerici, M., M. Faini, L. M. Muckenfuss, R. Aebersold, and M. Jinek, 2018 Structural basis of AAUAAA polyadenylation signal recognition by the human CPSF complex. Nat. Struct. Mol. Biol. 25: 135–138.

Gallicchio, L., G. H. Olivares, C. W. Berry, and M. T. Fuller, 2023 Regulation and function of alternative polyadenylation in development and differentiation. RNA Biol. 20: 908–925.

Grassi, E., R. Santoro, A. Umbach, A. Grosso, S. Oliviero *et al.*, 2019 Choice of alternative polyadenylation sites, mediated by the rna-binding protein Elavl3, plays a role in differentiation of inhibitory neuronal progenitors. Front. Cell. Neurosci.

Grozdanov, P. N., E. Masoumzadeh, M. P. Latham, and C. C. MacDonald, 2018 The structural basis of CstF-77 modulation of cleavage and polyadenylation through stimulation of CstF-

64 activity. Nucleic Acids Res. 46: 12022–12039.

Gruber, A. J., and M. Zavolan, 2019 Alternative cleavage and polyadenylation in health and

disease. Nat. Rev. Genet.

Guéguéniat, J., A. F. Dupin, J. Stojko, L. Beaurepaire, S. Cianférani *et al.*, 2017 Distinct roles of

PCF11 zinc-binding domains in pre-mRNA 3-end processing. Nucleic Acids Res.

Hilgers, V., S. B. Lemke, and M. Levine, 2012 ELAV mediates 3' UTR extension in the Drosophila

nervous system. Genes Dev. 26: 2259–2264.

Jing, Z., H. Linda, and M. Claire, 1999 Formation of mRNA 3' Ends in Eukaryotes: Mechanism,

Regulation, and Interrelationships with Other Steps in mRNA Synthesis. Microbiol. Mol.

Biol. Rev. 63: 405–445.

Jongmin Kim, Chenggang Lu, Shrividhya Srinivasan, Stephan Awe, Alexander Brehm *et al.*, 2017

Blocking promiscuous activation at cryptic promoters directs cell type – specific gene

expression. Science (80-.).

Kamieniarz-Gdula, K., M. R. Gdula, K. Panser, T. Nojima, J. Monks *et al.*, 2019 Selective Roles of

Vertebrate PCF11 in Premature and Full-Length Transcript Termination. Mol. Cell 74: 158-

172.e9.

Lee, S., Y.-C. Chen, A. E. Gillen, J. M. Taliaferro, B. Deplancke *et al.*, 2022 Diverse cell-specific

patterns of alternative polyadenylation in Drosophila. Nat. Commun. 13: 5372.

Li, H., J. Janssens, M. De Waegeneer, S. S. Kolluru, K. Davie *et al.*, 2023 Fly Cell Atlas: A single-

nucleus transcriptomic atlas of the adult fruit fly. Science (80-.). 375: eabk2432.

Li, W., J. Y. Park, D. Zheng, M. Hoque, G. Yehia *et al.*, 2016 Alternative cleavage and

polyadenylation in spermatogenesis connects chromatin regulation with post-

transcriptional control. BMC Biol.

Li, W., B. You, M. Hoque, D. Zheng, W. Luo *et al.*, 2015 Systematic Profiling of Poly(A)+

Transcripts Modulated by Core 3' End Processing and Splicing Factors Reveals Regulatory

Rules of Alternative Cleavage and Polyadenylation. PLOS Genet. 11: e1005166.

Mansfield, K. D., and J. D. Keene, 2012 Neuron-specific ELAV/Hu proteins suppress HuR mRNA

during neuronal differentiation by alternative polyadenylation. Nucleic Acids Res. 40:

2734–2746.

Marzluff, W. F., E. J. Wagner, and R. J. Duronio, 2008 Metabolism and regulation of canonical

histone mRNAs: Life without a poly(A) tail. Nat. Rev. Genet.

Mayr, C., and D. P. Bartel, 2009 Widespread Shortening of 3'UTRs by Alternative Cleavage and

Polyadenylation Activates Oncogenes in Cancer Cells. Cell 138: 673–684.

Miura, P., S. Shenker, C. Andreu-Agullo, J. O. Westholm, and E. C. Lai, 2013 Widespread and

extensive lengthening of 39 UTRs in the mammalian brain. Genome Res.

Morris, A. R., A. Bos, B. Diosdado, K. Rooijers, R. Elkon *et al.*, 2012 Alternative cleavage and

polyadenylation during colorectal cancer development. Clin. Cancer Res. 18: 5256–5266.

Ni, Z., A. Saunders, N. J. Fuda, J. Yao, J.-R. Suarez *et al.*, 2008 P-TEFb Is Critical for the

Maturation of RNA Polymerase II into Productive Elongation In Vivo. Mol. Cell. Biol. 28:

1161–1170.

Ogorodnikov, A., M. Levin, S. Tattikota, S. Tokalov, M. Hoque *et al.*, 2018 Transcriptome 3'end

organization by PCF11 links alternative polyadenylation to formation and neuronal

differentiation of neuroblastoma. Nat. Commun.

Raz, A. A., G. S. Vida, S. R. Stern, S. Mahadevaraju, J. M. Fingerhut *et al.*, 2023 Emergent

dynamics of adult stem cell lineages from single nucleus and single cell RNA-Seq of

Drosophila testes (M. Buszczak, U. Banerjee, M. Buszczak, & T. G. Nystul, Eds.). Elife 12:

e82201.

Sadowski, M., B. Dichtl, W. Hübner, and W. Keller, 2003 Independent functions of yeast Pcf11p

in pre-mRNA 3' end processing and in transcription termination. EMBO J. 22: 2167–2177.

Sarov, M., C. Barz, H. Jambor, M. Y. Hein, C. Schmied *et al.*, 2016 A genome-wide resource for

the analysis of protein localisation in Drosophila. Elife 5:.

Schäfer, P., C. Tüting, L. Schönemann, U. Kühn, T. Treiber *et al.*, 2018 Reconstitution of

mammalian cleavage factor II involved in 3' processing of mRNA precursors. RNA.

Shan, L., C. Wu, D. Chen, L. Hou, X. Li *et al.*, 2017 Regulators of alternative polyadenylation

operate at the transition from mitosis to meiosis. J. Genet. Genomics 44: 95–106.

Skrajna, A., X. Yang, M. Dadlez, W. F. Marzluff, and Z. Dominski, 2018 Protein composition of

catalytically active U7-dependent processing complexes assembled on histone pre-mRNA

containing biotin and a photo-cleavable linker. Nucleic Acids Res. 46: 4752–4770.

Tian, B., and J. L. Manley, 2016 Alternative polyadenylation of mRNA precursors. Nat. Rev. Mol.
Cell Biol.

Turner, R. E., L. M. Henneken, M. Liem-Weits, P. F. Harrison, A. Swaminathan *et al.*, 2020
Requirement for cleavage factor IIm in the control of alternative polyadenylation in breast
cancer cells. RNA 26: 969–981.

Vallejos Baier, R., J. Picao-Osorio, and C. R. Alonso, 2017 Molecular Regulation of Alternative
Polyadenylation (APA) within the Drosophila Nervous System. J. Mol. Biol.

Venkat, S., A. A. Tisdale, J. R. Schwarz, A. A. Alahmari, H. Carlo Maurer *et al.*, 2020 Alternative
polyadenylation drives oncogenic gene expression in pancreatic ductal adenocarcinoma.
Genome Res. 30: 347–360.

Wu, J., X. Li, Z. Gao, L. Pang, X. Liu *et al.*, 2021 RNA kinase CLP1/Cbc regulates meiosis initiation
in spermatogenesis. Hum. Mol. Genet. 30: 1569–1578.

Yalamanchili, H. K., C. E. Alcott, P. Ji, E. J. Wagner, H. Y. Zoghbi *et al.*, 2020 PolyA-miner:
Accurate assessment of differential alternative poly-adenylation from 3'Seq data using
vector projections and non-negative matrix factorization. Nucleic Acids Res.

Zeng, Y., A. Lovchykova, T. Akiyama, C. Liu, C. Guo *et al.*, 2024 TDP-43 nuclear loss in FTD/ALS
causes widespread alternative polyadenylation changes. bioRxiv 2024.01.22.575730.

Zhang, Z., and D. S. Gilmour, 2006 Pcf11 Is a Termination Factor in Drosophila that Dismantles
the Elongation Complex by Bridging the CTD of RNA Polymerase II to the Nascent
Transcript. Mol. Cell 21: 65–74.

Zhu, Y., X. Wang, E. Forouzmand, J. Jeong, F. Qiao *et al.*, 2018 Molecular Mechanisms for CFIm-
Mediated Regulation of mRNA Alternative Polyadenylation. Mol. Cell 69: 62-74.e4.

Fig.1

Fig.2

Control: b*amGal4*	PCF11 RNAi VDRC 103710 ; b*amGal4*	*PCF11* RNAi NIG 10228R-4 ; b*amGal4*

PCF11 — A, B, C

Phase — D, E, F

Control: *HA:cbc/CyO ; bamGal4*	*HA:cbc/+,cbc RNAi ; bamGal4*

HA:cbc — G, H (BL 77426), I (VDRC 100686)

Control: *bamGal4*	*cbc RNAi BL 77426 ; bamGal4*	*cbc RNAi VDRC 100686 ; bamGal4*

Phase — J, K, L

Fig.3

Fig.4

Fig.5

Fig.6