A single theory for the evolution of sex chromosomes and the two rules of speciation

Authors: Thomas Lenormand[1]*, Denis Roze[2,3]

Affiliations:

[1] CEFE, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France.

[2] CNRS, UMR 7144, Roscoff, France.

[3] Sorbonne Université, Station Biologique de Roscoff, France.

*Correspondence to: thomas.lenormand@cefe.cnrs.fr

Short title: Regulatory evolution can explain both sex chromosome evolution and the two rules of speciation

Abstract

Three major empirical patterns involving sex chromosomes have been observed in higher eukaryotes: Y (or W) chromosomes are often non-recombining and degenerate; when two species hybridize, but one sex is sterile or inviable among hybrid offspring, it is most often the heterogametic sex (XY or ZW)—the so-called Haldane's rule; and the X (or Z) plays a disproportionately large effect on reproductive isolation compared to autosomes—the so-called large X effect. Each observation has generally received its own tailored explanation involving multiple genetic and evolutionary causes[1–3]. Here, we show that these empirical patterns all emerge from a single theory for sex chromosome evolution incorporating the co-evolution of cis and trans-acting regulators of gene expression, and leading to systematic and rapid misexpression of dosage-compensated genes in the heterogametic F1 hybrids, for young or old sex-chromosomes. This theory links the evolution of sex chromosomes to the patterns of post-zygotic isolation observed in hybrids.

Main text

Sex chromosomes play a prominent role in the process of speciation. This is captured in the "two rules of speciation"[4]. The first, 'Haldane's rule' (HR), is named after the British scientist who observed that it was more frequently the heterogametic sex that suffered more in hybrids[5]. Over a century of work has confirmed the generality of this observation in nature[6–10]. Additionally, there is often an asymmetry of the effect observed between reciprocal crosses[11], termed 'Darwin's corollary' to HR. The second, the 'Large X effect' (LX), refers to the observation that X chromosomes (and Z chromosomes in ZW species) disproportionately affect hybrid incompatibilities more than autosomes of equivalent size[4,12–15]. These rules, among the few law-like generalizations in biology, have been extensively tested and studied and the subject of intense theoretical investigation to understand their origin[6,8–10,16,17].

Presently, the consensus is that HR and LX are composite phenomena with multiple genetic and
evolutionary causes [1–3,6,8–10,16–19]. While some theories have received more support than others, none
offers a general solution—each failing to account for some observations. For example, a prominent
explanation for the two rules is the Dominance theory, first suggested by Muller [20] and later formalized
[21,18,22], based on the idea that genetic incompatibilities involving at least one gene located on the X
chromosome (or Z, in species with female heterogamety) may more strongly affect the fitness of the
heterogametic sex if these incompatibilities are on average partially recessive. However, the theory
does not explain why incompatibilities should be, on average, recessive [21,23], although fitness landscape
models propose possible solutions [24–26] (sup. mat 3.1). Furthermore, it does not explain well why HR
often involves sterility rather than viability (as many more genes impact viability than fertility) [2,27], why
it is observed in groups lacking a hemizygous X [3], or in groups where XX females only express one X, as
in marsupials [28] or placentals (although in the latter case, both Xs may be expressed at the level of
tissues [18]). Other theories better accounting for the importance of hybrid sterility in HR have their own
major limitations. The "faster male theory" explains well why male sterility often occurs in hybrid
crosses in some groups [2,18,27] and why it may occur in species lacking hemizygosity [3], but critically fails
to account for HR in species where females are the heterogametic sex [16–18] (in these cases, a "faster
female" type of explanation would be required). The "meiotic drive theory" explains well why sex
chromosomes could play a major role in the sterility of heterogametic hybrids [29–31]. It has received
some empirical support [31–33] but does not offer a convincing explanation for HR for viability, since the
driving mechanism will only manifest itself when F1 hybrids undergo meiosis and attempt to produce
gametes, but not before[12,16]. Lastly, while sex chromosome degeneration is another globally observed
and intensely studied phenomenon, none of these theories consider the processes leading to
degenerate sex chromosomes and their subsequent evolution to be related to the emergence of either
HR or LX (Table 1).

We recently proposed a new theory for the evolution of non-recombining, degenerated, and dosage-
compensated sex chromosomes based on XY regulatory divergence and the early emergence of dosage
compensation (DC) [34–36]. We show here that the same theory also predicts Haldane's rule and the large
X effect. In the present study, we followed the simulated independent evolution of 20 species using
this previously described model [34] (Methods, Sup. Mat 1, Fig S1a). It consists of individual-based
stochastic simulations of a population of diploid individuals, with XY males and XX females (all the
arguments below also apply to ZZ / ZW systems and large Z effect, but for simplicity we only discuss
the XY case), incorporating deleterious mutations occurring at many genes, the evolution of
recombination and the evolution of cis and trans regulation of gene expression [34]. In this model, "lucky"
Y inversions fix because they capture a portion of the Y with a lower-than-average load of deleterious
mutation in the population (while also including the sex-determining locus), leading to recombination
arrest between the X and Y chromosomes. Then cis-regulatory divergence occurs together with an
accumulation of deleterious mutations, and emerging DC. This emerging DC creates sex-antagonistic
effects that prevent the reestablishment of recombination, leading to a new degenerate, silenced and
dosage compensated Y stratum. Once such a new stratum has evolved, cis- and trans-regulators
continue to coevolve to maintain optimal expression in both males and females. We estimated the
rate of occurrence and the pattern of hybrid incompatibilities by measuring the fitness of F1 hybrids
among the species at different time steps, under scenarios of sex chromosomes at different stages of
their evolution (see Methods).

Figure 1 shows the decrease in fitness in male and female F1 hybrids relative to the fitness of male and
female offspring from within species crosses. HR is rapidly observed in all cases, with strong
asymmetries between the fitness of male and female F1 hybrids. LX also occurs in all cases compared
to autosomes of equivalent size (compare Fig 1A to Fig 1B-E). Genes with an effect limited to the

heterogametic sex (i.e., involved in fertility rather than viability) also contribute to HR and LX, and in
many cases, very strongly (Fig 1E, Sup mat 2.4). Interestingly, the case of sex-limited genes can also
predict the occasional occurrence of Y-autosomes incompatibilities, as observed in some cases (Sup
mat 2.5). In Sup. Mat 1.5, we show that with larger population size and stronger intensity of stabilizing
selection on dosage, the fitness asymmetry between male and female F1 offspring tends to be larger,
but the overall decline in hybrid fitness is slower. In Sup. Mat. 1.6, we show that asymmetries between
reciprocal crosses often occur, as observed in "Darwin's corollary" to HR [11] (Fig S6). In this model,
autosomes also contribute to hybrid breakdown, but less strongly and without generating an
asymmetry between the homo and heterogametic sex (Fig 1A, sup. mat 2.1). This outcome results from
the coevolution of cis- and trans-regulators of gene expression, and their divergence among species
(Fig S2), which has repeatedly been emphasized as a mechanism generating hybrid incompatibilities
[37–49].

In this model, regulators on sex chromosomes evolve rapidly due to recombination arrest,
chromosomal degeneration and the emergence of dosage compensation [34]. As a consequence,
differences in regulatory traits can rapidly evolve between species and cause dysfunctional regulation
of dosage compensated genes in hybrids (Fig S2 shows the regulatory divergence observed in the
different scenarios). But why would dysfunctional regulation disproportionately impact the
heterogametic sex (HR) and cause a large X effect? For the latter, the answer is straightforward: if
dysfunctional regulation of dosage compensated genes inordinately reduces hybrid fitness in the
heterogametic sex, then the X chromosome, which is the only chromosome evolving DC, will
necessarily have a disproportionate effect on hybrid fitness compared to autosomes (both in terms of
the number and impact of genes involved in hybrid incompatibilities). Regarding HR, the disruption of
the regulation of dosage compensated genes can be caused by the portion of the Y that is degenerate
and compensated in only one of the two hybridizing species (Fig 1B, 1C) or by the portion of the Y that
is degenerate and compensated in both species (Fig 1C, 1D), provided they exhibit some regulatory
divergence. The effect per gene is on average larger in the first case (sup. mat. 1.4), but the total effect
depends on also on the number of genes in these different portions of the Y, so that both effects can
contribute significantly to HR. We detail each case in turn.

First, consider the case where species a evolves a new non-recombining stratum on the Y not present
in species b. If a gene g is degenerate and compensated in species a but not in species b, then hybrid
males will suffer from under-expression if they are $g_a^Y g_b^X$, and overexpression if they are $g_a^X g_b^Y$,
assuming codominant effects of the trans-acting factors. In the first case, these hybrids miss half the
trans-acting factor required to fully achieve DC with a degenerated g_a^Y copy. In the second case, the
hybrids inherit trans-regulators from species a increasing expression of gene g, despite having two
functional copies of the g gene. In all cases, however, expression is closer to the optimum in females,
regardless of the direction of the cross. This applies to the case where DC evolves by doubling
expression of the gene g in males through a male-specific trans-acting factor (as with the drosophila
DC mechanism, Fig S7a) or where X chromosomes have double expression, one being randomly
silenced in females (as in the mammalian DC mechanism, Fig S7b). This argument applies more
generally when evaluated quantitatively in a model encompassing these cases and their intermediates
(see Sup. Mat. 2). The male/female fitness ratio is decreased proportionally to the number of genes
on the stratum, the intensity of stabilizing selection on gene expression, and a constant that depends
on the DC mechanism. The effect described here is similar to the verbal argument of Filatov [50], although
clarifying the role of regulators.

Consider now the case of two species that inherited from their common ancestor a portion of the Y
that is non-recombining, degenerate, and dosage-compensated. In this case, misexpression of dosage

compensated genes continues to cause a major fitness decrease in the heterogametic F1. In this
scenario, the fitness decrease occurs when the regulatory traits, for a focal dosage compensated gene,
have diverged between the two species. In F1 hybrids, these compensations will average out in females
(who receive half the autosomal trans-acting factors and half the X cis-acting factors from each species)
but not in males, who will have cis-acting factors on the X from a single species, but averaged trans-
acting factors on autosomes (Fig S8). Males will consistently strongly under or overexpress compared
to females. A quantitative model shows that the reduction in male/female F1 fitness is proportional to
the number of genes on the stratum, the intensity of stabilizing selection on expression levels, and the
between-species variance in the trait governing DC (see Sup. Mat. 2). This difference in averaging
between sexes is similar to the mechanism generating HR in models of quantitative traits coded by
multiple loci with additive effects [25,26]. However, while these previous models may be seen as a
particular case of the dominance theory (as they assumed a mutation on the X has the same effect
when hemizygous in males and when homozygous in females), this is not the case in our model, since
a mutation on a cis-regulator on the X does not interact with the same trans-regulators in males and
females (see Sup. Mat. 3.1 and Fig S12).

Sex-limited fertility genes can also contribute to the decrease in hybrid fitness of the heterogametic
sex, with some interesting specificities. Female-limited genes on the X retain a standard diploid
expression. The coevolution of their cis and trans regulators can slowly diverge between species, but
this effect is relatively weak and comparable to what is observed on autosomes (see Sup. Mat. 2). The
effect is much larger for male-limited genes, and even larger than for genes expressed in both sexes
(Fig 1E). We also need to distinguish between cases where male-limited genes are present on a recently
derived non-recombining portion of the Y in one species or are ancestral to both. In the first case, gene
expression divergence between X and Y-linked copies of genes present in the new stratum will
generate a fitness cost for male hybrids in a similar way as for the non-sex-limited case (Fig S7a).
However, a difference with this non-sex-limited case is that either the X or the Y-linked copy may be
silenced following recombination arrest (while extinction of the X copy could not occur for genes
expressed and required in both sexes). In the second case, male-limited genes may already have been
silenced (either on the X or Y chromosome) after recombination arrest and before the split between
the two species, in which case the divergence of their regulatory traits between species will generate
a fitness cost for F1 males (Fig S9a). However, genes retaining diploid expression at the time of the
split may strongly reduce the fitness of F1 males. Indeed, if the X-linked copy is subsequently silenced
in one species while the Y-linked copy is subsequently silenced in the other, F1 males will either show
a complete lack of expression or major overexpression of that gene (Fig S9b). A single essential gene
in this situation could therefore cause male sterility in the heterogametic F1.

Hence, HR, its "Darwin's corollary" for reciprocal crosses (Sup. Mat. 2.6), and LX could be caused by
misregulation of sex-linked gene expression in hybrids. Although the idea of DC disruption has been
repeatedly discussed [4,6,10,19,50–54], and is close to the initial suggestion of Muller [55], it has never been
quantitatively formalized, and usually considers the possibility that the DC mechanism is deficient in
hybrids, not, more precisely, that the expression of dosage-compensated gene is deficient in hybrids.
While most authors have been cautious about completely rejecting the DC disruption hypothesis, it
has been usually dismissed as an important explanation [8,9,16] based on various arguments that we list
and critically evaluate in Sup. Mat 3.2. Most of the 13 criticisms we list seem either questionable or
outdated (for instance, the idea that ZZ/ZW systems do not have DC while exhibiting HR). In most cases
they focus only on the direct disruption of the DC pathway, and not on the disruption of expression of
dosage-compensated genes. Two arguments in this list remain relatively strong and require scrutiny.

The first states that in species with an ancestral 'global' (i.e., chromosome-wide) DC mechanism, there
may be little scope for divergence in DC, for any focal gene, in recently diverged species. This would
make a theory based on diverging DC less effective, especially in pairs of species where the same
portion of the Y is non-recombining and where the cis-regulators involved in global DC do not seem to
have diverged. However, DC disruption for a focal gene may still occur if target cis-regulatory
sequences do not change but move location on the X (in systems like *Drosophila* with high-affinity sites
'HAS' targeted by the MSL complex [56]) or if some genes escape the global DC mechanism. Current
evidence indicates that chromosome-wide DC is the exception, not the norm, and even in these cases,
many genes may escape global DC [57–60]. As our theory shows, these genes should be scrutinized for
their role in decreased hybrid fitness. Furthermore, global somatic DC is often absent from the
germline, where some DC nevertheless occurs for some genes [61–65]. This is observed independently of
the mechanism of meiotic sex chromosome inactivation (MSCI), which might have specifically evolved
to control sex-chromosome meiotic drive during early meiosis [66,67], and disruption of which has also
been suggested to contribute to HR [13,68–70]. This lack of a global DC mechanism for germline-limited
genes in the heterogametic sex could explain why male fertility genes are major contributors to HR
(interestingly male X-linked genes in *Drosophila melanogaster* are compensated independently of the
Male-Specific Lethal complex [71]). Indeed, without being tied to a global chromosomal level DC system,
their local regulation could much more easily diverge between recently diverged species. The fact that
male-limited genes can degenerate on the X and remain on the Y could also be a potent factor
preventing somatic DC from being "used" in the germline: upregulating the X is certainly not fitting
compensation for Y-linked genes expressed in the male germline. This could account for the
observation that different DC systems evolve in somatic vs. germinal tissues[65,72]. Interestingly, this
would lead to the prediction that HR should most often be based on hybrid sterility in groups with
global DC, while more often based on viability in groups lacking it — species evolving a global somatic
DC mechanism will maintain local DC only for sex-limited fertility genes, while species without global
somatic DC will have many viability genes with local DC. Indeed, birds and butterflies lacking global
somatic DC [73] show more cases of HR through inviability compared to mammals and Diptera [74], an
observation, that has otherwise never received any explanation. Hence, the present theory could
account for HR in species with global somatic DC, and also explain the contrasted prevalence of HR on
sterility vs inviability in different groups, based on whether they evolved a global or local DC
mechanism.

The second argument concerns species with recombining sex chromosomes. In species without
hemizygosity, it has been shown that HR is weaker but present compared to species with a more
degenerated Y. This is the case, for example, in *Aedes* compared to *Anopheles* mosquitoes [3]. This
example has been used to support the "faster male" theory and to rule out the dominance theory
alone could explain HR [3]. The same argument would also argue against the theory described here.
However, as recent evidence points out, in *Ae. aegyptii* the sex "locus" is a 1.5 Mb region with 30 genes
in a 100 Mb non-recombining region encompassing the centromere of chromosome 1 and showing
some divergence between males and females [75–77]. Hence, these species may not entirely lack
hemizygosity. Like above, genes in this region should be scrutinized for their role in decreased hybrid
fitness. The other argument favoring the "faster male" theory is the overrepresentation of male
sterility for HR, which is not a pattern directly following the dominance theory [2,27]. However, our results
also show that fertility genes can play a disproportionate role on HR, without invoking a "faster male"
effect.

Beyond the faster male theory, the most strongly supported explanation for HR is the dominance
theory. Support for the dominance theory's predictions comes particularly from experiments in
Drosophila involving unbalanced females with attached X chromosomes. However, our theory makes

similar predictions regarding these crosses (see Sup. Mat. 2.3), meaning they cannot discriminate
between theories. How do the dominance and our theories compare beyond this? In marsupials and
placentals, dosage compensation works by inactivating one X (the paternal or a random one,
respectively). This is a difficulty for the dominance theory for viability, as females are somatically
hemizygous (like males) in these cases. This is less of a concern for sterility, if the responsible genes
are expressed in the germline where the inactive X is reactivated in mammals [78]. This is not a major
concern in our theory, where, as explained above, HR may result from the divergence in the global DC
mechanism or from the few genes escaping global DC, whose regulation can more easily diverge
between species (including genes only expressed in the male germline), but where the majority of
genes subject to the same global DC will not cause any fitness reduction in hybrids. In the dominance
theory, recessive incompatibilities occurring on genes escaping global DC may also contribute to HR.
However, if such genes are rare, this contribution would be negligible: the vast majority of
incompatibilities will concern genes subject to global somatic DC, so the fitness reduction in males and
females will tend to be similar. A second point relates to the underlying mechanism. Contrary to the
dominance theory, which simply poses that a fraction of genetic incompatibilities are only expressed
when one of the underlying loci is homo- or hemizygous (without providing a biological mechanism
that would generate this type of interaction), our theory is based on a biological model of cis and trans
regulator evolution, with underlying additive traits. Furthermore, direct empirical evidence is
accumulating showing early regulatory evolution on sex chromosomes[79–82] and a role of misregulation
of dosage compensated genes in hybrid dysfunction. For instance, in *Drosophila,* the key elements for
DC—the MSL complex and the MSL-binding sites on the X—are fast evolving [52,53,83] as are *cis* and *trans*
regulators of X expression [84]. Indeed, Y-degeneration and DC appear sufficiently rapid and species-
specific in related species with young gene-rich sex chromosomes for the DC theory to work [50].
Evidence is also accumulating linking the misregulation of sex chromosomes to HR in various hybrids
[13,70,85–93]. Misregulation of gene expression in hybrids has already been extensively studied[37–49,86,87,90,93],
and improving the precision of techniques to measure gene expression will be crucial to critically
evaluate our theory, and determine whether F1 hybrid fitness is indeed reduced by the misregulation
of dosage compensated genes.

Overall, and perhaps more importantly, HR has been viewed as a composite phenomenon, produced
by different causes [1,2,2–4,6,8–10,16–19], possibly distinct from the large X effect [13,18]. As Coyne once wrote
about HR and LX, "biology is not physics" [17], emphasizing that given the complexity of biological
systems, unifying theories are unlikely. This conclusion may be true, but we suggest that it should
perhaps be re-evaluated regarding HR and LX given the explanatory power of the theory presented
here (Table 1). We show that a single model not only explains (1) recombination arrest on sex
chromosomes, (2) degeneration of the Y, (3) the evolution of dosage compensation, as we previously
showed [34,35], but also why, upon hybridization, (4) the heterogametic sex suffers more than the
homogametic one, and (5) the X plays a disproportionate role in speciation. It may also explain why (6)
HR is often asymmetrical between reciprocal crosses, (7) HR often involves fertility genes, (8) somatic
and germline DC often differ, and why, (9) misregulation of gene expression on autosomes should
follow HR in time and eventually lead to complete reproductive isolation at a later stage of speciation.
This theory therefore deserves to be further investigated empirically.

Methods
264

Our baseline model of sex chromosome evolution is the same as in ref. 34. In this study, we consider
a population of 10000 individuals (unless specified otherwise). Each individual has a pair of sex
chromosomes with a sex-determining locus and 500 genes that are subject to partially recessive
deleterious mutations. These genes are regularly positioned along a chromosome (Fig S1a). The

recombination rate between consecutive genes (R_g) is set to 0.0005. Recombination events are
simulated by drawing the number of crossing over at meiosis from a Poisson distribution (whose mean
is equal to the total map length) and drawing their position uniformly along the chromosome. Gene
expression is regulated by cis-regulatory sequences (affecting expression only on the same
chromosome as themselves, and whose strength is measured by a quantitative trait noted *c*)
interacting with autosomal trans regulators (affecting expression of gene copies on both homologs and
whose strength is measured by a quantitative trait noted *t*) (Fig S1b). The recombination rate between
each *cis*-regulator and its gene (R_c) is 0.00005. All these elements mutate: mutation add a unbiased
Gaussian deviate $N(0, \sigma = 0.2)$ to regulatory traits *c* and *t* (at rates $U_t = U_c/2 = 10^{-4}$, negative trait
values are counted as zero). Mutation on coding sequences (occurring at rate $U_g = 2.10^{-4}$) are
deleterious with an effect drawn in an exponential distribution with mean $\bar{s}$ = 0.05. These deleterious
mutations can accumulate additively on each gene, up to a maximal value s_{max} = 0.3 representing the
fitness effect of loosing that gene altogether. In order to permit dosage compensation on a gene-by-
gene basis while maintaining symmetry in the model for males and females, it is assumed that each
gene is controlled by one male- and one female-expressed trans regulators (the model does not include
the question of evolving a global DC mechanism). We assume that the overall expression level of each
gene *Q* is subject to stabilizing selection around an optimal level Q_{opt} = 2 with intensity *I* (see Eq. 1 in
sup. mat 2 and Fig S1d, we use *I* = 0.1 unless specified otherwise). For proper scaling, we ensure that,
irrespectively of *I*, evolving zero expression has the same fitness effect as loosing the gene altogether
(i.e. at *Q* = 0, the fitness reduction is s_{max}), which results in an asymmetry in the function describing this
stabilizing selection (see Eq 1 and ref. 35). Furthermore, we assume that the dominance level of a
deleterious mutation occurring in a coding gene is modified by the relative expression of the two copies
of that gene [34,35]. Naturally, a deleterious mutation occurring in a less expressed gene copy is assumed
to be less harmful than one in a more highly expressed copy. Specifically, and arbitrarily denoting with
a 1 or 2 subscript two alleles of a given gene, we assume that the fraction of the protein expressed
from alleles "1" at a given locus is $\phi = c_1/(c_1 + c_2)$. This ratio measures the degree of allele-specific
expression. With $\phi = 1/2$ (i.e. with equally strong cis-regulators on both homologs), alleles are co-
expressed, while a departure from ½ indicates that one allele is relatively more expressed than the
other. Supposing that allele "1" is deleterious, then its dominance is given by $h_1 = \phi^{-ln(h)/ln(2)}$ (see
Fig S1c), where *h* is the parameter measuring the baseline dominance of the fitness effect of
deleterious mutations in a heterozygote when both alleles are equally expressed (with $\phi = 1/2$, we
have $h_1 = h$). We use *h* = 0.25, corresponding to the average value observed across species [94]. Fitness
effects combine multiplicatively across loci (no epistasis).

We also assume that mutations occur which suppress recombination on a segment of the Y
chromosome. For simplicity, these mutations are referred to as inversions, although they could
correspond to other mechanisms causing recombination arrest. Inversions of any size are possible
(their size is drawn in a uniform distribution within the non-recombining part of the Y), but we follow
only those on the Y that include the sex-determining locus, which will necessarily be confined to males
and cause recombination arrest. We assume that these inversions can accumulate, so that new
inversions can occur on chromosomes carrying a previous inversion and thus extend the non-
recombining region of the Y. Finally, we assume that reversions restoring recombination can occur,
and for simplicity, that such reversions cancel only the most recent inversion [34]. We consider that
inversions occur at a rate U_{inv} = 10^{-5}, and that these reversions occur 10 times less frequently. Table
S1b recapitulates all parameter used and their values.

We consider three main scenarios. In the first ("neo-Y"), the sex chromosome pair has just acquired
the sex-determination locus. We follow the independent evolution of this recent Y in different replicate
species (we call them species for simplicity, but these are, at least initially, only independently evolving

populations). After a given number of generations, F1 hybrids are generated between these independently evolved species, in the two directions of the cross (M x F and F x M), and their fitness is compared to that of male and female offspring produced within species. The second scenario ("half-degenerate case") corresponds to the case of a partially degenerated Y chromosome: it is equivalent to the first scenario, except that 50% of the Y is already non-recombining, fully degenerated, and fully dosage compensated in the common ancestor of the diverging species (with Drosophila like DC, i.e. with the trans-acting factors expressed in males t_m = 2). Finally, the third scenario ("XO") assumes that 100% of the Y is already non-recombining, fully degenerated, and fully dosage compensated in the ancestor (this is equivalent to considering diverging species with a XX / XO sex determination system, we also use t_m = 2 as the initial condition). These scenarios allow us to compare hybrid fitness between species at different stages of sex chromosome evolution.

We also investigate the case of genes with male- or female-limited effects to represent genes involved in fertility rather than viability. In this case, we suppose that the sex chromosomes are not recombining. We performed simulations with male and female-limited genes independently. When a simulation runs with only male-limited genes, the fitness of females stays at 1 throughout, and reciprocally for female-limited genes. For the comparison, we therefore randomly paired male-limited and female-limited simulations. For female-limited genes, all genes are initially non-degenerate and with fair diploid expression. Degeneration does not occur on the X. Whether degeneration occurs on the Y is irrelevant since the genes are all female-limited and therefore not expressed in males. For male-limited genes (assuming a XY system, the reverse situation with female-limited case would occur for a ZW system), we considered two initial conditions: (1) all genes are initially non-degenerate and with fair diploid expression. During the course of these simulations, each male-limited gene will degenerate and become silenced on either the X or the Y, independently in different populations. We term this initial condition the "unsorted" case since it is initially not decided which gene will be lost on the Y or on the X. Results show that c.a. 90% of genes end up being degenerate on the Y (and 10% on the X, Fig S11). In the second initial condition we suppose that 50% of the genes are degenerate on the Y (and dosage compensated on the X with t_m = 2) while 50% are degenerate on the X (and dosage compensated on the Y with t_m = 2). We term this initial condition the "sorted case", meaning that it is already decided which genes are degenerate on the X or Y (and it is the same for all replicated species).

In addition to these scenarios, we performed control simulations. We consider simulations with a fully recombining autosome instead of sex chromosome pair (without including new inversions on this autosome but including the possibility of regulatory evolution, with one male and one female trans-acting factor per gene). Note that in these simulations, the sex of each newly formed individual is drawn randomly (with an expected 1:1 sex ratio). Another control involves simulations with sex-chromosomes, but without considering new inversions (in the XO scenario, this control is irrelevant since the whole Y is already non-recombining). Finally, we also investigated the effect of varying population size and the intensity of stabilizing selection on dosage on the outcome (see Sup. mat 1.5).

References

1. Orr, H. A. Haldane's rule has multiple genetic causes. *Nature* **361**, 532–533 (1993).
2. Wu, C. I. & Davis, A. W. Evolution of postmating reproductive isolation: the composite nature of Haldane's rule and its genetic bases. *Am Nat* **142**, 187–212 (1993).
3. Presgraves, D. C. & Orr, H. A. Haldane's rule in taxa lacking a hemizygous X. *Science* **282**, 952–954 (1998).
4. Coyne, J. A. & Orr, H. A. Two rules of speciation. in *Speciation and its Consequences* 180–207 (Sinauer, Sunderland, MA, 1989).
5. Haldane, J. B. S. Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics* **12**, 101–109 (1922).

6.   Laurie, C. C. The weaker sex is heterogametic: 75 years of Haldane's rule. *Genetics* **147**, 937–
951 (1997).
7.   Coyne, J. A. & Orr, A. H. The evolutionary genetics of speciation. *Phil. Trans. R. Soc. Lond. B* **353**,
287–305 (1998).
8.   Schilthuizen, M., Giesbers, M. C. W. G. & Beukeboom, L. W. Haldane's rule in the 21st century.
Heredity **107**, 95–102 (2011).
9.   Delph, L. F. & Demuth, J. P. Haldane's Rule: Genetic Bases and Their Empirical Support. *JHERED*
107, 383–391 (2016).
10. Cowell, F. 100 years of Haldane's rule. *J of Evolutionary Biology* **36**, 337–346 (2023).
11. Turelli, M. & Moyle, L. C. Asymmetric postmating isolation: Darwin's corollary to Haldane's rule.
Genetics **176**, 1059–1088 (2007).
12. Coyne, J. A. Genetics and speciation. *Nature* **355**, 511–515 (1992).
13. Masly, J. P. & Presgraves, D. C. High-resolution genome-wide dissection of the two rules of
speciation in *Drosophila*. *PLOS Biology* **5**, e243 (2007).
14. Presgraves, D. C. Sex chromosomes and speciation in *Drosophila*. *Trends in Genetics* **24**, 336–
343 (2008).
15. Irwin, D. E. Sex chromosomes and speciation in birds and other ZW systems. *Molecular Ecology*
27, 3831–3851 (2018).
16. Coyne, J. A. & Orr, H. A. *Speciation*. (Sinauer Associates, Sunderland (Massachusetts), 2004).
17. Coyne, J. A. "Two rules of speciation" revisited. *Molecular Ecology* **27**, 3749–3752 (2018).
18. Turelli, M. & Orr, H. A. Dominance, epistasis and the genetics of postzygotic isolation. *Genetics*
154, 1663–1679 (2000).
19. Johnson, N. A. & Lachance, J. The genetics of sex chromosomes: evolution and implications for
hybrid incompatibility. *Annals of the New York Academy of Sciences* **1256**, E1–E22 (2012).
20. Muller, H. J. Isolating mechanisms, evolution and temperature. *Biol Symp* **6**, 71–125 (1942).
21. Orr, H. A. & Turelli, M. Dominance and Haldane's rule. *Genetics* **143**, 613–616 (1996).
22. Orr, H. A. A mathematical model of Haldane's rule. *Evolution* **47**, 1606–1611 (1993).
23. Sawamura, K. Maternal effect as a cause of exceptions for Haldane's Rule. *Genetics* **143**, 609–
611 (1996).
24. Barton, N. H. The role of hybridization in evolution. *Molecular ecology* **10**, 551–568 (2001).
25. Fraïsse, C., Gunnarsson, P. A., Roze, D., Bierne, N. & Welch, J. J. The genetics of speciation:
insights from Fisher's geometric model. *Evolution* **70**, 1450–1464 (2016).
26. Schneemann, H., Munzur, A. D., Thompson, K. A. & Welch, J. J. The diverse effects of
phenotypic dominance on hybrid fitness. *Evolution* **76**, 2846–2863 (2022).
27. Wu, C.-I., Johnson, N. A. & Palopoli, M. F. Haldane's rule and its legacy: Why are there so many
sterile males? *Trends in Ecology & Evolution* **11**, 281–284 (1996).
28. Watson, E. T. & Demuth, J. P. Haldane's Rule in marsupials: What happens when both sexes are
functionally hemizygous? *Journal of Heredity* **103**, 453–458 (2012).
29. Hurst, L. D. & Pomiankowski, A. Causes of sex ratio bias may account for unisexual sterility in
hybrids: A new explanation of Haldane's rule and related phenomena. *Genetics* **128**, 841–858
(1991).
30. Frank, S. A. Haldane's rule: A defense of the meiotic drive theory. *Evolution* **45**, 1714–1717
(1991).
31. Patten, M. M. Selfish X chromosomes and speciation. *Mol Ecol* **27**, 3772–3782 (2018).
32. Meiklejohn, C. D. & Tao, Y. Genetic conflict and sex chromosome evolution. *Trends in Ecology*
and Evolution **25**, 215–223 (2010).
33. Presgraves, D. C. & Meiklejohn, C. D. Hybrid sterility, genetic conflict and complex speciation:
lessons from the *Drosophila simulans* clade species. *Frontiers in Genetics* **12**, (2021).
34. Lenormand, T. & Roze, D. Y recombination arrest and degeneration in the absence of sexual
dimorphism. *Science* **375**, 663–666 (2022).
35. Lenormand, T., Fyon, F., Sun, E. & Roze, D. Sex chromosome degeneration by regulatory
evolution. *Current Biology* **30**, 3001-3006.e5 (2020).

36.   Lenormand, T. & Roze, D. Can mechanistic constraints on recombination reestablishment
explain the long-term maintenance of degenerate sex chromosomes? *Peer Community Journal*
4, e17 (2024).
37.   Johnson, N. A. & Porter, A. H. Rapid speciation via parallel, directional selection on regulatory
genetic pathways. *Journal of Theoretical Biology* **205**, 527–542 (2000).
38.   Ranz, J. M., Namgyal, K., Gibson, G. & Hartl, D. L. Anomalies in the expression profile of
interspecific hybrids of *Drosophila melanogaster* and *Drosophila simulans*. *Genome Res.* **14**,
373–379 (2004).
39.   Wittkopp, P. J., Haerum, B. K. & Clark, A. G. Evolutionary changes in cis and trans gene
regulation. *Nature* **430**, 85–8 (2004).
40.   Landry, C. R. *et al.* Compensatory cis-trans evolution and the dysregulation of gene expression
in interspecific hybrids of *Drosophila*. *Genetics* **171**, 1813–1822 (2005).
41.   Haerty, W. & Singh, R. S. Gene regulation divergence is a major contributor to the evolution of
Dobzhansky–Muller incompatibilities between species of *Drosophila*. *Molecular Biology and*
Evolution **23**, 1707–1714 (2006).
42.   Ortíz-Barrientos, D., Counterman, B. A. & Noor, M. A. F. Gene expression divergence and the
origin of hybrid dysfunctions. *Genetica* **129**, 71–81 (2006).
43.   Tulchinsky, A. Y., Johnson, N. A. & Porter, A. H. Hybrid incompatibility despite pleiotropic
constraint in a sequence-based bioenergetic model of transcription factor binding. *Genetics*
198, 1645–1654 (2014).
44.   Mack, K. L., Campbell, P. & Nachman, M. W. Gene regulation and speciation in house mice.
Genome Res **26**, 451–461 (2016).
45.   Mack, K. L. & Nachman, M. W. Gene regulation and speciation. *Trends in Genetics* **33**, 68–80
(2017).
46.   Sánchez-Ramírez, S., Weiss, J. G., Thomas, C. G. & Cutter, A. D. Widespread misregulation of
inter-species hybrid transcriptomes due to sex-specific and sex-chromosome regulatory
evolution. *PLOS Genetics* **17**, e1009409 (2021).
47.   Scarpino, S. V. *et al.* Evolution of a genetic incompatibility in the genus *Xiphophorus*. *Molecular*
Biology and Evolution **30**, 2302–2310 (2013).
48.   Meiklejohn, C. D., Coolon, J. D., Hartl, D. L. & Wittkopp, P. J. The roles of cis- and trans-
regulation in the evolution of regulatory incompatibilities and sexually dimorphic gene
expression. *Genome Research* **24**, 84–95 (2014).
49.   Wittkopp, P. J., Haerum, B. K. & Clark, A. G. Regulatory changes underlying expression
differences within and between Drosophila species. *Nature genetics* **40**, 346–50 (2008).
50.   Filatov, D. A. The two "rules of speciation" in species with young sex chromosomes. *Molecular*
Ecology **27**, 3799–3810 (2018).
51.   Orr, H. A. Does postzygotic isolation result from improper dosage compensation? *Genetics* **122**,
891–894 (1989).
52.   Rodriguez, M. A., Vermaak, D., Bayes, J. J. & Malik, H. S. Species-specific positive selection of
the male-specific lethal complex that participates in dosage compensation in *Drosophila*.
Proceedings of the National Academy of Sciences **104**, 15412–15417 (2007).
53.   Bachtrog, D. Positive selection at the binding sites of the male-specific lethal complex involved
in dosage compensation in *Drosophila*. *Genetics* **180**, 1123–1129 (2008).
54.   Vicoso, B. & Bachtrog, D. Progress and prospects toward our understanding of the evolution of
dosage compensation. *Chromosome Research* **17**, 585–602 (2009).
55.   Muller, H. J. Bearings of the 'Drosophila' work on systematics. in *The new systematics* 185–268
(Clarendon, Oxford, 1940).
56.   Quinn, J. J. *et al.* Rapid evolutionary turnover underlies conserved lncRNA–genome
interactions. *Genes Dev* **30**, 191–207 (2016).
57.   Carrel, L. & Willard, H. F. X-inactivation profile reveals extensive variability in X-linked gene
expression in females. *Nature* **434**, 400–404 (2005).

58. Disteche, C. M. Dosage compensation of the sex chromosomes and autosomes. *Seminars in Cell*
& Developmental Biology **56**, 9–18 (2016).
59. Chandler, C. H. When and why does sex chromosome dosage compensation evolve? *Annals of*
the New York Academy of Sciences **1389**, 37–51 (2017).
60. Gu, L. & Walters, J. R. Evolution of sex chromosome dosage compensation in animals: A
beautiful theory, undermined by facts and bedeviled by details. *Genome Biology and Evolution*
9, 2461–2476 (2017).
61. Tabuchi, T. M. *et al.* Chromosome-biased binding and gene regulation by the *Caenorhabditis*
elegans DRM complex. *PLOS Genetics* **7**, e1002074 (2011).
62. Meiklejohn, C. D., Landeen, E. L., Cook, J. M., Kingan, S. B. & Presgraves, D. C. Sex chromosome-
specific regulation in the *Drosophila* male germline but little evidence for chromosomal dosage
compensation or meiotic inactivation. *PLOS Biology* **9**, e1001126 (2011).
63. Strome, S., Kelly, W. G., Ercan, S. & Lieb, J. D. Regulation of the X chromosomes in
Caenorhabditis elegans. *Cold Spring Harb Perspect Biol* **6**, a018366 (2014).
64. Taxiarchi, C. *et al.* High-resolution transcriptional profiling of Anopheles gambiae
spermatogenesis reveals mechanisms of sex chromosome regulation. *Sci Rep* **9**, 14841 (2019).
65. Mahadevaraju, S. *et al.* Dynamic sex chromosome expression in *Drosophila* male germ cells. *Nat*
Commun **12**, 892 (2021).
66. Tao, Y. *et al.* A sex-ratio meiotic drive system in *Drosophila simulans*. II: An X-linked distorter.
PLoS Biology **5**, 2576–2588 (2007).
67. Lenormand, T., Engelstädter, J., Johnston, S. E. S. E., Wijnker, E. & Haag, C. R. Evolutionary
mysteries in meiosis. *Philosophical Transactions of the Royal Society B-Biological Sciences* **371**,
20160001 (2016).
68. Lifschytz, E. & Lindsley, D. L. The role of X-chromosome inactivation during spermatogenesis.
Proceedings of the National Academy of Sciences **69**, 182–186 (1972).
69. Turner, J. M. A. Meiotic sex chromosome inactivation. *Development* **134**, 1823–1831 (2007).
70. Larson, E. L., Keeble, S., Vanderpool, D., Dean, M. D. & Good, J. M. The composite regulatory
basis of the large X-effect in mouse speciation. *Molecular Biology and Evolution* **34**, 282–295
(2017).
71. Philip, P. & Stenberg, P. Male X-linked genes in *Drosophila melanogaster* are compensated
independently of the Male-Specific Lethal complex. *Epigenetics & Chromatin* **6**, 35 (2013).
72. Landeen, E. L., Muirhead, C. A., Wright, L., Meiklejohn, C. D. & Presgraves, D. C. Sex
chromosome-wide transcriptional suppression and compensatory cis-regulatory evolution
mediate gene expression in the *Drosophila* male germline. *PLOS Biology* **14**, e1002499 (2016).
73. Mank, J. E. The W, X, Y and Z of sex-chromosome dosage compensation. *Trends in Genetics* **25**,
226–233 (2009).
74. Schilthuizen, M., Giesbers, M. C. W. G. & Beukeboom, L. W. Haldane's rule in the 21st century.
Heredity **107**, 95–102 (2011).
75. Juneja, P. *et al.* Assembly of the genome of the disease vector *Aedes aegypti* onto a genetic
linkage map allows mapping of genes affecting disease transmission. *PLOS Neglected Tropical*
Diseases **8**, e2652 (2014).
76. Matthews, B. J. *et al.* Improved reference genome of *Aedes aegypti* informs arbovirus vector
control. *Nature* **563**, 501–507 (2018).
77. Aryan, A. *et al.* Nix alone is sufficient to convert female *Aedes aegypti* into fertile males and
myo-sex is needed for male flight. *Proceedings of the National Academy of Sciences* **117**,
17702–17709 (2020).
78. Sangrithi, M. N. & Turner, J. M. A. Mammalian X Chromosome Dosage Compensation:
Perspectives From the Germ Line. *BioEssays* **40**, 1800024 (2018).
79. Veltsos, P. *et al.* Early sex-Chromosome evolution in the diploid dioecious plant *Mercurialis*
annua. *Genetics* **212**, 815–835 (2019).

80. Wei, K. H. C. & Bachtrog, D. Ancestral male recombination in *Drosophila albomicans* produced geographically restricted neo-Y chromosome haplotypes varying in age and onset of decay. *PLoS Genetics* e1008502 (2019) doi:10.1371/journal.pgen.1008502.

81. Hough, J., Hollister, J. D., Wang, W., Barrett, S. C. H. & Wright, S. I. Genetic degeneration of old and young Y chromosomes in the flowering plant *Rumex hastatulus*. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 7713–8 (2014).

82. Martin, H. *et al.* Evolution of young sex chromosomes in two dioecious sister plant species with distinct sex determination systems. *Genome Biology and Evolution* **11**, 350–361 (2019).

83. Levine, M. T., Holloway, A. K., Arshad, U. & Begun, D. J. Pervasive and largely lineage-specific adaptive protein evolution in the dosage compensation complex of *Drosophila melanogaster*. *Genetics* **177**, 1959–1962 (2007).

84. Coolon, J. D. *et al.* Molecular mechanisms and evolutionary processes contributing to accelerated divergence of gene expression on the *Drosophila* X chromosome. *Mol Biol Evol* **32**, 2605–2615 (2015).

85. Lu, X. *et al.* Genome-wide misexpression of X-linked versus autosomal genes associated with hybrid male sterility. *Genome Res* **20**, 1097–1102 (2010).

86. Good, J. M., Giger, T., Dean, M. D. & Nachman, M. W. Widespread over-expression of the X chromosome in sterile F1 hybrid mice. *PLoS Genet* **6**, e1001148 (2010).

87. Oka, A. & Shiroishi, T. Regulatory divergence of X-linked genes and hybrid male sterility in mice. *Genes Genet Syst* **89**, 99–108 (2014).

88. Turner, L. M., White, M. A., Tautz, D. & Payseur, B. A. Genomic networks of hybrid sterility. *PLOS Genetics* **10**, e1004162 (2014).

89. Davis, B. W. *et al.* Mechanisms underlying mammalian hybrid sterility in two feline interspecies models. *Mol Biol Evol* **32**, 2534–2546 (2015).

90. Li, Y. *et al.* Regulatory divergences in dosage compensation cause hybrid male inviability in *Caenorhabditis*. 2024.02.06.577000 Preprint at https://doi.org/10.1101/2024.02.06.577000 (2024).

91. Xiong, T. *et al.* A polygenic explanation for Haldane's rule in butterflies. *Proceedings of the National Academy of Sciences* **120**, e2300959120 (2023).

92. Arévalo, L., Gardner, S. & Campbell, P. Haldane's rule in the placenta: Sex-biased misregulation of the Kcnq1 imprinting cluster in hybrid mice. *Evolution* **75**, 86–100 (2021).

93. Viswanath, A. & Cutter, A. D. Regulatory Divergence as a Mechanism for X-Autosome Incompatibilities in Caenorhabditis Nematodes. *Genome Biol Evol* **15**, evad055 (2023).

94. Manna, F., Martin, G. & Lenormand, T. Fitness landscape: an alternative theory for the dominance of mutations. *Genetics* **189**, 923–937 (2011).

95. Turelli, M. & Orr, H. A. The dominance theory of Haldane's Rule. *Genetics* **140**, 389–402 (1995).

96. Presgraves, D. C. Patterns of postzygotic isolation in *Lepidoptera*. *Evolution* **56**, 1168–1183 (2002).

97. Muller, H. J. Isolating mechanisms, evolution and temperature. Chap 6:71-125 (1942).

98. Straub, T. & Becker, P. B. Dosage compensation: the beginning and end of generalization. *Nat Rev Genet* **8**, 47–57 (2007).

99. Zhang, Y. & Oliver, B. Dosage compensation goes global. *Current Opinion in Genetics & Development* **17**, 113–120 (2007).

100. Pessia, E., Makino, T., Bailly-Bechet, M., McLysaght, A. & Marais, G. A. B. Mammalian X chromosome inactivation evolved as a dosage-compensation mechanism for dosage-sensitive genes on the X chromosome. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 5346–5351 (2012).

101. Slavney, A., Arbiza, L., Clark, A. G. & Keinan, A. Strong constraint on human genes escaping X-inactivation is modulated by their expression level and breadth in both sexes. *Molecular Biology and Evolution* **33**, 384–393 (2016).

102. Sidorenko, J. *et al.* The effect of X-linked dosage compensation on complex trait variation. *Nat Commun* **10**, 3009 (2019).

103. Navarro-Cobos, M. J., Balaton, B. P. & Brown, C. J. Genes that escape from X-chromosome
inactivation: Potential contributors to Klinefelter syndrome. *Am J Med Genet C Semin Med*
Genet **184**, 226–238 (2020).

104. San Roman, A. K. *et al.* The human inactive X chromosome modulates expression of the active X
chromosome. *Cell Genomics* **3**, 100259 (2023).

105. Coyne, J. A. The genetic basis of Haldane's rule. *Nature* **314**, 736–738 (1985).

106. Charlesworth, B., Coyne, J. A. & Barton, N. H. The relative rates of evolution of sex
chromosomes and autosomes. *The American Naturalist* **130**, 113–146 (1987).

107. Nozawa, M., Onizuka, K., Fujimi, M., Ikeo, K. & Gojobori, T. Accelerated pseudogenization on
the neo-X chromosome in *Drosophila miranda*. *Nat Commun* **7**, 13659 (2016).

108. Hughes, J. F. *et al.* A gene deriving from the ancestral sex chromosomes was lost from the X and
retained on the Y chromosome in eutherian mammals. *BMC Biology* **20**, 133 (2022).

109. Amrein, H. & Axel, R. Genes Expressed in neurons of adult male *Drosophila*. *Cell* **88**, 459–469
(1997).

110. Lercher, M. J., Urrutia, A. O. & Hurst, L. D. Evidence that the human X chromosome is enriched
for male-specific but not female-specific genes. *Molecular Biology and Evolution* **20**, 1113–1116
(2003).

111. Wang, P. J., McCarrey, J. R., Yang, F. & Page, D. C. An abundance of X-linked genes expressed in
spermatogonia. *Nat Genet* **27**, 422–426 (2001).

112. Lamnissou, K., Loukas, M. & Zouros, E. Incompatibilities between Y chromosome and
autosomes are responsible for male hybrid sterility in crosses between *Drosophila virilis* and
Drosophila texana. *Heredity (Edinb)* **76 (Pt 6)**, 603–609 (1996).

113. Pantazidis, A. C. & Zouros, E. Location of an autosomal factor causing sterility in *Drosophila*
mojavensis males carrying the *Drosophila arizonensis* Y chromosome. *Heredity (Edinb)* **60 (Pt**
2), 299–304 (1988).

114. Araripe, L. O., Tao, Y. & Lemos, B. Interspecific Y chromosome variation is sufficient to rescue
hybrid male sterility and is influenced by the grandparental origin of the chromosomes.
Heredity **116**, 516–522 (2016).

115. Sackton, T. B., Montenegro, H., Hartl, D. L. & Lemos, B. Interspecific Y chromosome
introgressions disrupt testis-specific gene expression and male reproductive phenotypes in
Drosophila. *Proceedings of the National Academy of Sciences* **108**, 17046–17051 (2011).

116. Orr, H. A. Genetics of sterility in hybrids between two subspecies of *Drosophila*. *Evolution;*
international journal of organic evolution **43**, 180–189 (1989).

117. Nozawa, M. *et al.* Shared evolutionary trajectories of three independent neo-sex chromosomes
in *Drosophila*. *Genome Res* **31**, 2069–2079 (2021).

118. Murphy, P. J. & Berger, F. The chromatin source-sink hypothesis: a shared mode of chromatin-
mediated regulations. *Development* **150**, dev201989 (2023).

119. Francisco, F. O. & Lemos, B. How do Y-chromosomes modulate genome-wide epigenetic states:
genome folding, chromatin sinks, and gene expression. *Journal of Genomics* **2**, 94–103 (2014).

120. Wei, K. H.-C., Gibilisco, L. & Bachtrog, D. Epigenetic conflict on a degenerating Y chromosome
increases mutational burden in *Drosophila* males. *Nat Commun* **11**, 5537 (2020).

121. Muyle, A., Bachtrog, D., Marais, G. A. B. & Turner, J. M. A. Epigenetics drive the evolution of sex
chromosomes in animals and plants. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **376**, (2021).

122. Brown, E. J., Nguyen, A. H. & Bachtrog, D. The *Drosophila* Y chromosome affects
heterochromatin integrity genome-wide. *Molecular Biology and Evolution* **37**, 2808–2824
(2020).

123. Nozawa, M., Ikeo, K. & Gojobori, T. Gene-by-gene or localized dosage compensation on the
neo-X chromosome in *Drosophila miranda*. *Genome Biology and Evolution* **10**, 1875–1881
(2018).

124. Walters, J. R. & Hardcastle, T. J. Getting a full dose? Reconsidering sex chromosome dosage
compensation in the silkworm, *Bombyx mori*. *Genome Biology and Evolution* **3**, 491–504 (2011).

125. Veneti, Z., Bentley, J. K., Koana, T., Braig, H. R. & Hurst, G. D. D. A functional dosage
compensation complex required for male killing in *Drosophila*. *Science* **307**, 1461–1463 (2005).
126. Fukui, T. *et al.* The endosymbiotic bacterium *Wolbachia* selectively kills male hosts by targeting
the masculinizing gene. *PLOS Pathogens* **11**, e1005048 (2015).
127. Cheng, B., Kuppanda, N., Aldrich, J. C., Akbari, O. S. & Ferree, P. M. Male-killing spiroplasma
alters behavior of the dosage compensation complex during *Drosophila melanogaster*
embryogenesis. *Current Biology* **26**, 1339–1345 (2016).
128. Jaffe, E. & Laird, C. Dosage compensation in *Drosophila*. *Trends in Genetics* **2**, 316–321 (1986).
129. Pal Bhadra, M., Bhadra, U. & Birchler, J. A. Misregulation of sex-lethal and disruption of male-
specific lethal complex localization in *Drosophila* species hybrids. *Genetics* **174**, 1151–1159
(2006).
130. Chatterjee, R. N., Chatterjee, P., Pal, A. & Pal-Bhadra, M. *Drosophila simulans* Lethal hybrid
rescue mutation (Lhr) rescues inviable hybrids by restoring X chromosomal dosage
compensation and causes fluctuating asymmetry of development. *J Genet* **86**, 203–215 (2007).
131. Barbash, D. A. Genetic testing of the hypothesis that hybrid male lethality results from a failure
in dosage compensation. *Genetics* **184**, 313–316 (2010).
132. Brideau, N. J. *et al.* Two Dobzhansky-Muller genes interact to cause hybrid lethality in
Drosophila. *Science* **314**, 1292–1295 (2006).
133. Wit, E. de, Greil, F. & Steensel, B. van. Genome-wide HP1 binding in *Drosophila*: Developmental
plasticity and genomic targeting signals. *Genome Res.* **15**, 1265–1273 (2005).
134. Presgraves, D. C., Balagopalan, L., Abmayr, S. M. & Orr, H. A. Adaptive evolution drives
divergence of a hybrid inviability gene between two species of *Drosophila*. *Nature* **423**, 715–
719 (2003).
135. Tang, S. & Presgraves, D. C. Evolution of the *Drosophila* Nuclear Pore Complex results in
multiple hybrid Incompatibilities. *Science* **323**, 779–782 (2009).
136. Mendjan, S. *et al.* Nuclear pore components are involved in the transcriptional regulation of
dosage compensation in *Drosophila*. *Molecular Cell* **21**, 811–823 (2006).
137. Chen, J., Wang, M., He, X., Yang, J.-R. & Chen, X. The evolution of sex chromosome dosage
compensation in animals. *Journal of Genetics and Genomics* **47**, 681–693 (2020).
649

Acknowledgments: We thank D Presgraves, C. Haag, M. Kirkpatrick, S. Otto, M Raymond, J Welch and
anonymous reviewers for discussions and/or comments and K. McKean for editing. We thank CNRS
ABiMs cluster. **Funding**: This work was supported by CNRS. **Author contributions**: Original ideas TL,
DR; Model conception TL, DR; Code DR, TL; Simulations TL; Data analyses TL; Interpretation TL, DR;
First draft, editing and revisions TL, DR; Project management and funding TL. **Competing interests**: the
authors declare no conflict of interest. **Data and materials availability**: simulation code is available on
Zenodo https://zenodo.org/doi/10.5281/zenodo.12584015.

Fig. 1. Hybrid fitness in crosses between Independently evolving species. The figure shows the fitness of homogametic (x-axis) and heterogametic F1 (y-axis), between species that have evolved independently for 2.5×10^5, 10^6 or 4×10^6 generations (light, medium, and dark color on each panel, respectively). The data are obtained in each case using 15 independently evolving species and the 105 possible hybrid crosses (averaged in the two directions FxM and MxF for each of them). The fitness of hybrids is computed relative to the average fitness of intraspecific crosses for male and female offspring, respectively. Dots (or crosses for panel A) are mean values for all replicates, and small dots individual values. Contours represent the areas containing the individual values (the envelope is computed with a smoothed Gaussian kernel to facilitate vizualisation). The x=y line is added to show equal fitness reduction in male and female F1. HR corresponds to cases where points fall below this line (the fitness of the heterogametic sex is lower than that of the homogametic sex in F1 hybrids. LX corresponds to a larger effect in panels B-E than the one shown on panel A for autosomes. (**A**) Autosomal case (see method). (**B**) Neo-Y case (see methods). In gray, the control where inversions do not occur, so that the XY remains fully recombining throughout. In orange/red/darker red, inversions (and reversions) occur and recombination progressively stops between the X and the Y in each species independently (Fig S3 shows the progression of recombination arrest in the different replicates). After 2.5×10^5 generations, only two species evolved a non-recombining stratum on the Y, and all hybrids involving those species had a lower fitness in the heterogametic sex (those points form two clouds of points, whose contours are indicated by the orange arrows). (**C**) Half degenerate case (see methods). Note that regulatory divergence can accumulate initially for a larger number of dosage-compensated genes (initially 250), explaining why the fitness reduction on hybrids is larger for the same time period than in (B), where the simulations start with no dosage compensated gene. In gray, the control where inversions do not occur and where all the effect is therefore only driven by the shared ancestral stratum (the difference between the curves shows the contribution of newly evolving strata). (**D**) XO case (see methods). In this case, there is no control simulation where inversions evolve since the Y is

initially already fully non-recombining. Here too, regulatory divergence can accumulate initially for a larger number of dosage-compensated genes (500) than in (B) and (C), explaining the larger fitness reduction in hybrids. (**E**) Sex-limited case (see methods). The figure shows the result of simulations with only female-limited genes (x-axis) or only male-limited genes (y-axis), starting from a fully non-recombining Y. These simulations are performed independently and are randomly paired between male-limited and female-limited simulations to be represented on the same graph for better comparison with other cases (see methods). In the "unsorted" case, each male-limited gene become independently degenerate and silenced on either the X or Y during the course of the simulation (see Fig S11), while in the "sorted" case, it is initially decided, and the same for all replicates, which is lost on the Y and which is lost on the X (see methods). The fitness of the heterogametic F1 drops very quickly toward a value close to zero in the unsorted case.

	Dominance theory	Faster male theory	Meiotic drive theory	Regulatory theory
HR in XY species	✔	✔	✔	✔
HR in ZW species	✔	✗	✔	✔
Large X/Z effect	✔	✗	✔	✔
Importance of HR through sterility in several groups	✗	✔	✔	✔
Importance of HR through inviability in several groups	✔	✗	✗	✔
Darwin's corollary to HR	✔	✗	✗	✔
HR through inviability/sterility in species without/with global DC	✗	✗	✗	✔
HR in species with X inactivation in females	(1)	✔	✔	✔
HR on sterility sometimes caused by Y or W-autosomes interactions	(2)	(3)	✔	✔
HR on sterility in XY species with homomorphic sex chromosomes	(4)	✔	(4)	(4)
Theory also explains evolution of sex chromosomes	✗	✗	✗	✔

Table. 1. Summary of the comparison of main predictions of the different theories for Haldane's rule. Regulatory theory refers to the theory presented in this paper. In green/orange/red the theory adequately/partially/inadequately predicts the pattern indicated in the first column. The cases where the theories fail or succeed are explained in the text. There are four "unsettled" cases (in orange and numbered): (1) HR in mammal species with X inactivation in females could be explained by the dominance theory if HR concerns male sterility, but not male inviability. (2) Y-autosomes interactions are not predicted in the usual dominance theory, as the argument is usually based on male hemizygosity resulting from Y degeneration by selective interference. However, as we show, male hemizygosity from X degeneration could evolve in a regulatory model for some male-limited genes. (3) The faster male theory, even if it does not specifically predict rare Y-autosome interactions, could accommodate this observation. (4) The observation of HR in species lacking a hemizygous X challenge all theories based on sex-chromosomes, but these species might actually have a small non-recombining region, sufficient to generate HR (see text).

Supplementary material

1 Additionnal methods and results

1.1 Graphical illustration of the model

The model considers X and Y sex chromosomes with a sex determining locus (SDL) located at one end of the chromosome as shown on Fig S1a(a). The Y can evolve non-recombining strata (the figure S1a(a) illustrates a Y with two strata like on Figs S10a, S7-S9, in green and blue). 500 coding genes, each with its own cis-regulator are present on the Y and X.

As shown on Fig S1a(b), the expression of each gene is symmetrically controlled by two autosomal trans-regulators, one expressed in males and the other in females, allowing for free evolution of dosage compensation.

Fitness effect of deleterious mutations: each allele of the coding gene expresses a given amount of protein determined by the trait values corresponding to its cis and trans-regulators (see methods). Fig S1a(c) shows how the relative proportion of protein produced from the two alleles modulates the dominance level of the deleterious alleles that could occur in the coding sequence as shown by the curve.

Fig S1a(d) shows the fitness effect of expression level. The total amount of protein produced for a gene (as determined by trait values corresponding to its cis and trans-regulators) is under stabilizing selection around an optimal dosage, for each gene. Zero expression has a fitness effect of s_{max}, corresponding to the maximal fitness effect of deleterious mutations that can accumulate on that gene (such that knocking out a gene by accumulating deleterious mutations has the same effect as reducing its expression to zero). This s_{max} value is equal to 0.3 in all simulations.

754   **Fig. S1a. Graphical illustration of the model.** (a) Chromosomes and genes, (b) Cis

755   and trans regulators, (c) fitness effect of deleterious mutations, (d) fitness effect of

756   expression levels (see methods).

Parameter	Description	Value
N_g	Number of genes	500
N	Number of individuals	**10000**, 5000, 20000
$\bar{s}$	Average fitness effect of deleterious mutations in coding genes	0.05
h	Baseline dominance of deleterious mutations in coding genes	0.25
s_{max}	Maximum effect of cumulated effect of deleterious mutations on a given coding gene	0.3
I	Intensity of stabilizing selection on dosage	**1/10**, 1/30, 1/3
σ	Standard deviation of the phenotypic effect of regulatory mutations	0.2
Q_{opt}	Optimal expression level for coding genes	2
U_g	Mutation rate per coding gene per generation	$2.\,10^{-4}$
U_c	Mutation rate per cis-regulator per generation	$2.\,10^{-4}$
U_t	Mutation rate per trans-regulator per generation	10^{-4}
U_{inv}	Rate of inversion per chromosome per generation	10^{-5}
U_{rev}	Reversion rate of inversions per generation	10^{-6}
R_g	Recombination distance between consecutive coding genes	$5.\,10^{-4}$
R_c	Recombination distance between a coding gene and its cis-regulator	$5.\,10^{-5}$

Table S1b. Parameter values. Model's parameters, their notation, description and values (in bold their default

value when several values were considered).

1.2 Measure of regulatory divergence

The regulatory system for a given species and a given gene can be characterized by the vector $\mathbf{r} =$

(t_m, t_f, c_X), where t_m, t_f, c_X are the trait values for the male trans-regulator, the female trans-

regulator and the cis-regulator on the X. On the Y, the cis-regulatory trait tends towards zero after

recombination suppression and thus do not contribute to regulatory variation among independently

evolving species (before recombination suppression, c_Y equals c_X and therefore does not bring

additinnal information on regulatory divergence). At a given time and for a given gene, we can

therefore measure the regulatory divergence among independently evolving species as the mean over

replicates i of $(\mathbf{r}_i - \bar{\mathbf{r}}) . (\mathbf{r}_i - \bar{\mathbf{r}})$. This divergence can then be averaged across genes for a specific region

of the sex chromosome. Fig S4 shows this measure of divergence every 40 000 for windows of 100
consecutive genes on the se chromosomes for the simulations represented on Fig. 1.

Fig. S2. Divergence of regulatory traits through time. The x-axis is time (number of generations).
The y-axis is a measure of the divergence of regulatory traits through time in independently
evolving species. To compute it, the chromosome is divided in five equally sized regions: region
1 (brown) includes genes 1 to 100, region 2 (red) genes 101 to 200, region 3 (orange) genes 201
to 300, region 4 (light orange) genes 301 to 400, region 5 (yellow) genes 401 to 500. The
divergence measure is given every 40 000 generations averaged for the 100 genes in these five
regions, from the same simulations than those reported on Fig. 1: **(a)** Simulations with an
autosome, **(b)** starting with a neo-Y, **(c)** starting with a half-degenerate Y, **(d)** with a fully
degenerate Y. When recombination suppression evolves in some replicates, but not all, it causes
a slight acceleration of regulatory divergence, as expected. This effect can be seen on simulations
starting with a neo-Y (panel b). Recombination suppression occurs first in some replicates in the
first region (brown), then in the second (red) etc. It can also be seen on simulations starting with
a half-degenerated Y (panel c). Recombination suppression occurs first in some replicates in half
of the third region for genes 250 to 300 (orange), then in the fourth (light orange) and the fifth
(yellow).

1.3 Evolution of new non-recombining strata

In neo-Y (Fig 1B) and half-degenerated Y scenarios (Fig 1C), with new inversions and reversions,
recombination suppression evolves independently in the different replicates. Some inversions fix in
the population and are not reversed : they are "stabilized" (see ref. 34 for details about the underlying
processes leading to these fixations and stabilisation of these new strata). The size of newly occuring
inversions is drawn from a uniform distribution over the recombining fraction of the Y chromosome.
However, the process of fixation and stabilization favors inversions with intermediate but relatively
small size. The size of each new stabilized stratum can be followed independently in each replicate
through time. This is illustrated on Fig S3.

Fig S3. Evolution of recombination suppression in neo-Y and half-degenerated Y scenarios. (a) Evolution of recombination suppression in the neo-Y simulations corresponding to Fig 1B. The x-axis is time in number of generations, in log-scale, with the three time points illustrated on Fig 1 shown by the vertical dashed curves. The y-axis is the fraction of the Y with suppressed recombination corresponding to the accumulation of several strata. The gray lines represent individual replicates (F1 hybrids are generated by crosses among these replicates) and the black curve the mean across replicates. **(b)** Same as in (a), but for the simulations starting with a 'half-degenerate' Y, corresponding to Fig 1C.

1.4 Fitness effect per gene in derived and ancestral strata

Consider two species evolving in a neo-Y scenario (their common ancestor has a fully recombining Y).
Suppose that after t generations of divergence, species 1 evolves a non-recombining region up to
position z_1 on the Y chromosome, and species 2 up to z_2. We can define the region up to Min(z_1, z_2) as
a "parallel" stratum. This region is non-recombining in both species but has evolved independently in
each. The region between Min(z_1, z_2) and Max(z_1, z_2) can then be defined as a "private" stratum (i.e.
this region is non-recombining in only one of the two species). We can refer to these parallel and
private strata as "derived" strata. The mean effect of a gene (on the fitness of F1 individuals) from
these derived strata can be computed by averaging over genes in parallel or private regions in all
species pairs. Similarly, we can consider two species evolving in an XO scenario. Their common
ancestor has a fully non-recombining, degenerate and dosage-compensated Y (with a given dosage
compensation t_m). After t generations of divergence, the mean effect of one gene (on the fitness of F1
individuals) within this "ancestral" stratum (of 500 genes) can be computed and averaged over all pairs
of species. All these measures can be made independently for females and males (i.e. for the homo-
and heterogametic sex, respectively).

Figure S4(a) shows these mean fitness effects per gene in parallel, private and ancestral non-
recombining regions. The effect of genes is of the same order of magnitude among derived strata
(private or parallel), with a large difference between their effects in male and female F1. The measures
show some noise, especially initially when few derived strata have evolved (Fig S4(b) shows the
average number of genes present in parallel and private strata over time). The effect of genes in the
ancestral strata is also greater in males than in females, but the male effect is smaller (compared to
the derived strata). This observation confirms equations 7-9 where the expected effect on ancestral
strata per gene is expected to be weaker than on derived strata. Note that the mean effect per locus
is small here. However HR may not necessarily be caused by many small effects. Few dosage sensitive
loci may contribute strongly to HR, and in some case, a single gene could even cause sterility in hybrids
(see Sup. Mat. 2.4 about the effect of sex-limited genes).

Fig S4. Effect per gene and number of genes in different non-
recombining regions. (a) Mean effect of genes on F1 fitness in
different non-recombining regions (parallel, private, ancestral,
see above) after a given time of divergence between species (in
number of generations, x-axis). (b) Mean number of genes in
derived strata (parallel or private, see above) in neo-Y
simulations. The data are from the neo-Y simulations shown on
Fig 1B (for derived strata) and Fig 1D (for ancestral stratum).

1.5 Effect of population size and intensity of stabilizing selection on hybrid fitness

We investigated the effect of varying population size and the intensity of stabilizing selection on dosage
on the fitness of F1 hybrids. In the main scenarios reported on Fig. 1, population size was set to 10000
individuals and I = 1/10. We performed additional simulations with 5000 or 20000 individuals in
scenario 1 (neo-Y) and 3 (XO) to investigate the effect of increasing or decreasing population size. We
also performed simulations with I = 1/30 or 1/3 in scenario 1 (neo-Y) and 3 (XO) to investigate the
effect of increasing or decreasing the intensity of stabilizing selection (Fig S5). Other parameters have
predictable effect on the outcome and were not investigated. Notably, increasing the number of loci
or the effect of a gene knock-out (s_{max}) will directly increase the total fitness effect seen in hybrids (see
Eq. 7) and varying mutations rates will change the timescale. The effect of the parameters on the
process of Y evolution was also investigated in previous studies, and will influence the process in a
predictable way in the scenarios involving the formation of new sex-chromosome strata[34–36]. The effect
of varying I or N is less straightforward as it can be expected to have contradicting effects on the
outcome. Indeed, increasing both N and I speeds up the formation of new strata[34], but may
subsequently limit regulatory divergence by cis-trans coevolution (since this coevolution involves
crossing small "fitness valleys"). We find that the former effect prevails. Larger N and I leads indeed to
faster evolution of recombination suppression in neo-Y scenario, but it does not suppress cis trans
coevolution. In both Neo-Y and XO scenario, the fitness asymmetry between homo and heterogametic
F1 hybrids tends to be larger for larger N and I. In all cases, however, decreasing N and I also tends to
globally decrease the fitness of all F1, consistent with the expectation that stronger drift / lower
stabilizing selection on dosage allows faster cis-trans coevolution.

Fig S5. Hybrid fitness in crosses between Independently evolving species for varying population sizes and intensity of stabilizing selection on dosage. The figure is built like Fig 1 in the main text and shows the relative fitnesses of homo- and heterogametic F1 hybrids in neo-Y (panels A and C) and XO scenarios (panels B and D) after 4 million generations. Panels A and B show results for varying population sizes (values indicated on the plots, with I = 1/10). Panels C and D show results for varying intensity of stabilizing selection (values indicated on the plots, with N = 10000). Each dot represent F1 hybrids from a specific species pair. Larger dots represent mean values and contours around dots (computed with a smoothed Gaussian kernel) are added to facilitate visualization.

1.6 Darwin's corollary

The fitness of F1 hybrids often differs significantly between reciprocal crosses. For instance, Turelli and Orr [95] estimated that 15% of cases of Haldane's rule in *Drosophila* involved a strong asymmetry (the male F1 being sterile or inviable in only one direction of the cross between the two hybridizing species). A similar pattern is seen in *Anopheles* [3] and could be even more prevalent in *Lepidoptera* [96] or *Silene* [11]. Turelli and Moyle termed this pattern "Darwin's corollary", complementing Haldane's rule and the large X effect in the rules of speciation [11]. In our model, fitness asymmetries between reciprocal crosses often arise. The general reason for this asymmetry is that the heterogametic F1 will typically suffer from over-expression in one direction of the cross and from under-expression in the other. This pattern occurs in all cases (see detailed examples in Fig S7-S9). Everything being equal, under-expression is more deleterious than overexpression in our model, as is likely in most biologically plausible situations. In the extreme case, e.g. when a male-limited gene is entirely missing in one direction of the cross, male fitness is much more reduced than when its expression is doubled in the other direction of the cross (Fig S9b). The asymmetry in reciprocal crosses is particularly strong when a discrete event of large effect occurs in one species but not in the other (similarly, Muller noted that asymmetry between reciprocal hybrids must involve loci of large effect [97]). This occurs for instance when a species acquires a new non-recombining stratum and not the other. In this case, the F1 male inheriting the more degenerate Y will suffer more than the F1 male in the reciprocal cross (Fig S7a, S7b). This effect will occur mostly at intermediate steps of Y chromosome degeneration, i.e. when species have different strata, and before F1 male fitness is not too strongly reduced in all crosses (Fig S6a). Another case particularly conducive to strong fitness asymmetries in reciprocal crosses is when a male-limited gene

is missing in one direction, as explained above. However, if there are many male-limited genes, then
different genes will be lost on the X or Y in the different species, and the fitness of all F1 males will be
reduced. Darwin's corollary should be strongest when the number of male-limited genes on sex
chromosomes is not too large, maximizing the sampling variance of those genes and therefore the
fitness asymmetries in the reciprocal crosses (compare Fig S6b(a) and S6b(c), with 500 and 50 male-
limited genes on the chromosome respectively). When the fitness reduction in F1 males results from
many small effects, fitness asymmetries in reciprocal crosses tend to be weaker. This is the case for
instance when regulatory divergence occurs on a chromosome that is already fully dosage-
compensated (see Fig S6b for XO simulations). Overall, our model predicts substantial fitness
asymmetries in reciprocal crosses, and at intermediate times of species divergence, which is in line
with the available observations.

Fig S6a. Darwin's corollary in the neo-Y scenario. (**a**) Crosses asymmetries in neo-Y
simulations (same simulations as the one illustrated in Fig1B). The axes show the
absolute value of the difference in the fitness of males produced in reciprocal crosses
(y-axis) against time (x-axis, in millions of generations, in log-scale). A larger value
means that the male fitness reduction is larger in one direction of the cross. Gray dots
are the individual values obtained from all hybrids among independently evolving
populations; red dots indicate mean values. (**b**) The graph shows the slope of the
regression of the (signed) fitness difference between males produced in reciprocal
crosses against the (signed) difference in the size of the non-recombining region (*z*)
between the Ys in the two populations (y-axis). A positive slope indicates that males
suffer more in the direction of the cross where the father carries a Y with a larger non-
recombing region (i.e. a larger z value).

Fig S6b. Darwin's corollary in other scenarios. (a) Crosses asymmetries in XO
simulations (same simulations as the one illustrated in Fig1D). The graphs
show the absolute value of the difference in the fitness of males produced in
reciprocal crosses (y-axes) against time (x-axis, in millions of generations, in
log-scale). A larger value means that the male fitness reduction is larger in
one direction of the cross. Gray dots are the individual values obtained from
all hybrids among independently evolving populations; red dots indicate
mean values. **(b)** Same as in (a), but for male-limited simulations with 500
genes (same simulations as the unsorted case illustrated in Fig1E). (**c**): Same
as in (a), but for male-limited simulations with 50 genes. The x-axis does not
represent the exact same range of values in the three panels to better
emphasize the period during which asymmetries are maximized.

1.7 Numerical examples for the different situations discussed in the main text.

Fig S7a. Example of the effect of a gene located in a derived stratum present in only one species, with a drosophila-like DC mechanism. The figure shows species A and B, where the Y has a non-recombining stratum (in green) inherited from the common ancestor, and a newly evolved non-recombining stratum only in species B (in blue), as in Fig S10a. The example focuses on a gene in the derived (blue) stratum. In species A, DC does not evolve (since the gene is located in a recombining portion of the Y), while in species B, DC is achieved for that gene, since it is located in the non-recombining region. In this example, DC in species B is achieved by having male overexpression of the X, like in *Drosophila* (i.e. by evolving a male-specific trans-acting factor increasing X expression in males, t_m = 2 in species B). Male and female expression levels is indicated by the value of Q. Q = 2 is the optimal expression level (Q is computed as $(c_{X,i} + c_{Y,i})\bar{t}_{m,i}$ and $(c_{X1,i} + c_{X2,i})\bar{t}_{f,i}$ in males and females, respectively, see methods). F1 males show a departure from optimal expression, while F1 females maintain optimal expression, in both directions of the cross. Note that the reduction in F1 male fitness is potentially asymmetric in the two directions of the cross, resulting from either over or underexpression.

Fig S7b. Example of the effect of a gene located in a derived stratum present in only one species, with a mammal-like DC mechanism. The figure shows species A and B, where the Y has a non-recombining stratum (in green) inherited from the common ancestor, and a newly evolved non-recombining stratum only in species B (in blue), as in Fig S10a. The example focuses on a gene in the derived (blue) stratum (like in Fig S7a). In species A, DC does not evolve (since the gene is located in a recombining portion of the Y), while in species B, DC is achieved for that gene, since it is located in the non-recombining region. The example shows a mammal-like pattern of DC for species B: DC in species B is achieved by having overall X overexpression (by evolving stronger X cis-regulatory elements, c_{xB} = 2 in species B) which is corrected in females to avoid overshooting (by evolving a female-specific trans-acting factor decreasing X expression in females, t_f = 0.5 in species B). Male and female expression levels is indicated by the value of Q. Q = 2 is the optimal expression level (Q is computed as $(c_{X,i} + c_{Y,i})\bar{t}_{m,i}$ and $(c_{X1,i} + c_{X2,i})\bar{t}_{f,i}$ in males and females, respectively, see methods). F1 males show a greater departure from optimal expression than F1 females. Note that the reduction in F1 male fitness is potentially asymmetric in the two directions of the cross, resulting from either over or underexpression.

Fig S8. Example of the effect of a gene located in an ancestral stratum, with different DC mechanisms in the diverging species. The figure shows species A and B, where the Y has a non-recombining stratum (in green) inherited from the common ancestor, and a newly evolved non-recombining stratum only in species B (in blue), as in Fig S10a. The example focuses on a gene in the ancestral (green) stratum. In species A, DC is achieved, for the focal gene, by having overall X overexpression (by evolving a stronger X cis-regulatory element, c_{xA} = 2 in species A) which is corrected in females to avoid overshooting (by evolving a female-specific trans-acting factor decreasing X expression in females, t_f = 0.5 in species A). In species B, DC is achieved, for the same gene, by having male overexpression of the X (i.e. by evolving a male-specific trans-acting factor increasing X expression in males, t_m = 2 in species B). Male and female expression levels is indicated by the value of Q. Q = 2 is the optimal expression level (Q is computed as $(c_{X,i} + c_{Y,i})\bar{t}_{m,i}$ and $(c_{X1,i} + c_{X2,i})\bar{t}_{f,i}$ in males and females, respectively, see methods). Male F1 show a greater departure from optimal expression than female F1, in both directions of the cross. Note that the reduction in F1 male fitness is potentially asymmetric in the two directions of the cross, resulting from either over or underexpression.

Fig S9a. Example of the effect of a male-limited gene located in an ancestral stratum, with different DC mechanisms in the diverging species. The figure shows species A and B, where the Y has a non-recombining stratum (in green) inherited from the common ancestor, and a newly evolved non-recombining stratum only in species B (in blue), as in Fig S10a. The example focuses on a male-limited gene in the ancestral (green) stratum. The gene is not expressed in females. In species A, DC is achieved, for the focal gene, by having overall X overexpression (by evolving stronger X cis-regulatory elements, c_{xA} = 2 in species A) which is corrected in females to avoid overshooting (by evolving a female-specific trans-acting factor decreasing X expression in females, t_f = 0.5 in species A). In species B, DC is achieved, for the same gene, by having male overexpression of the X (i.e. by evolving a male-specific trans-acting factor increasing X expression in males, t_m = 2 in species B). Male and female expression levels is indicated by the value of Q. Q = 2 is the optimal expression level (Q is computed as $(c_{X,i} + c_{Y,i})\bar{t}_{m,i}$ and $(c_{X1,i} + c_{X2,i})\bar{t}_{f,i}$ in males and females, respectively, see methods). Male F1 show a departure from optimal expression in both directions of the cross. Note that the reduction in F1 male fitness is potentially asymmetric in the two directions of the cross, resulting from either over or underexpression. Note also that this situation is identical to Fig S7, ignoring females.

1005

1006

1007

Fig S9b. Example of the effect of a male-limited gene located in an ancestral stratum, with alternate X or Y
degeneration in the diverging species. The figure shows species A and B, where the Y has a non-recombining
stratum (in green) inherited from the common ancestor, and a newly evolved non-recombining stratum only in
species B (in blue), as in Fig S10a. The example focuses on a male-limited gene in the ancestral (green) stratum
which is silenced on the Y in species A ($c_{yA} = 0$), but silenced on the X in species B ($c_{xB} = 0$). Indeed, male-limited
genes can degenerate both ways, which differs from non-sex-limited cases shown in Fig S7 and S8 where
degeneration can only occur on the Y. The gene is not expressed in females. In both species, DC is achieved by
having male overexpression of either the X or Y, through the evolution of stronger cis-regulators ($c_{xA} = c_{yB} = 2$).
Male F1 show a strong departure from optimal expression in both directions of the cross, notably in one direction
where the gene is completely silenced. Note that the reduction in F1 male fitness is potentially asymmetric in
the two directions of the cross, resulting from either over or underexpression.

1019

1020

2 Computing hybrid fitness

In males, the total expression level Q_i for a gene i is equal to $(c_{X,i} + c_{Y,i})\bar{t}_{m,i}$, where $\bar{t}_{m,i}$ is the average strength of the *trans*-regulators expressed in males, while $c_{X,i}$ and $c_{Y,i}$ are the strength of cis regulators associated with the X and Y-linked copies of gene i, respectively. Symmetrically, it is $(c_{X1,i} + c_{X2,i})\bar{t}_{f,i}$ in females. The decrease in fitness resulting from the departure from optimal dosage W_i^Q can be expressed as a function of the intensity of stabilizing selection on the expression level, I :

$$W_i^Q = 1 - s_{max}\left(1 - e^{-I\left(\ln Q_i - \ln Q_{opt}\right)^2}\right) \qquad (1)$$

This expression ensures that complete loss of expression (at Q = 0) has the same fitness effect as the fitness effect of completely disruting the coding sequence (s_{max}, see methods). For the purpose of this analysis, we assume that Q_{opt}= 2 without loss of generality. We also define $\theta = I\,s_{max}$, which represents the overall intensity of stabilizing selection on dosage. We drop below the indices i referring to a specific gene, to simplify the notation.

A variety of of dosage compensation mechanisms have been described across a range of organisms, which may be represented by specific instance of our general model. Indeed, once the Y is fully silenced ($c_Y = 0$), and assuming that the population remains at Q_{opt} in both males and females (which is a good approximation unless stabilizing selection is particularly weak), we have

$$Q_{opt} = c_X\bar{t}_m = 2c_X\bar{t}_f \qquad (2)$$

We therefore have $\bar{t}_m = 2\bar{t}_f$, which defines dosage compensation. Hence the manner in which DC operates (i.e. the triplet $\bar{t}_m, \bar{t}_f, c_X$) can be delineated by a single parameter, given that there is only one degree of freedom in the way DC occurs. For the purpose of this description, we may, for instance choose to use $\bar{t}_m$. In comparison to the initial system in which $\bar{t}_m = \bar{t}_f = c_X = c_Y = 1$, a final compensation characterized by $\bar{t}_m$ =1 (i.e. $\bar{t}_m = 1, \bar{t}_f = 0.5, c_X = 2, c_Y = 0$) would correspond to the *Caenorhabditis elegans* case, where the X is inherently expressed twice as much ($c_X = 2$) to achieve optimal expression in males, while an hermaphrodite-limited *trans*-regulator halves expression ($\bar{t}_f = 0.5$) to recover optimal expression in hermaphrodites. This is also analogous to the mammal case where a female-limited *trans*-regulator halves expression by randomly silencing one X (instead of halving the expression of each X like in *C. elegans*). The case $\bar{t}_m$ = 2 (i.e. $\bar{t}_m = 2, \bar{t}_f = 1, c_X = 1, c_Y = 0$) would correspond to the *Drosophila* case, where a male-limited *trans*-acting factor doubles X expression to achieve optimal expression in males (with no change in females, $\bar{t}_f = 1, c_X = 1$).

2.1 Contribution of autosomes to the fitness reduction of F1 hybrids

On autosomes, cis and trans regulators can coevolve provided that the overall expression level stays close to its optimum value. This condition implies that, for each gene, t_m and t_f stay both close to $1/c$. Hence, autosomes can contribute to a reduction in hybrid fitness, but it will be symmetrical for males and females. Noting Δ the difference in t values between the two species for a given gene, and assuming that this difference is weak, we find that the reduction in fitness of F1 hybrids caused by misregulation of this gene is, to leading order in Δ:

1061

$$W_{hybrid} = 1 - \theta(\Delta/2t_m)^4 + O(\Delta)^5 \qquad (3)$$

1063

1064 This effect is modest (note the quartic exponent), but can lead to substantial fitness loss if cumulated
1065 over many genes.

2.2 Contribution of ancestral and derived Y strata to the fitness reduction of F1 hybrids

We can compute the male/female fitness ratio in hybrids, accounting
for two types of loci (Fig S10a). We consider first the Y-linked loci that
are degenerate and silenced in both species (assume there are k_1 such
loci). These loci are for instance located in a non-recombining region
that is common to both species (green stratum 1 in Fig S10a, where
the sex-detrmining locus, SDL, is located). Second, we consider loci
that are degenerate and silenced in only one species (assume there are
k_2 such loci). These loci are for instance located in a non-recombining
region that only evolved in one species (blue stratum 2 of species B in
Fig S10a). We compute the contribution to Haldane's rule of these two
types of loci in turn.

2.2.1 Contribution of an ancestral stratum (stratum 1)

1079 For loci that are degenerate in both species, the contribution of a given gene to the male and female
1080 fitness can be approximated assuming that the gene is fully silenced on the Y and dosage-compensated
1081 in both species, that is, without loss of generality $c_Y = 0$, $c_X = 2/t_m$, $t_f = t_m/2$. Noting Δ the
1082 difference in t_m values between the two species for a given gene, and assuming that this difference is
1083 weak, we find that the contribution to Haldane's rule of that gene is, to leading order in Δ:

1084

$$W_{XY} = 1 - \theta(\Delta/2t_m)^2 + O(\Delta)^3 \qquad (4a)$$

$$W_{XX} = 1 - \theta(\Delta/2t_m)^4 + O(\Delta)^5 \qquad (4b)$$

1087

1088 The fitness reduction in males is much larger than in females (which equals the fitness reduction
1089 expected for an autosomal locus). The male to female fitness ratio ρ is therefore

1090

$$\begin{aligned}\rho &= 1 - \theta(\Delta/2t_m)^2 + O(\Delta)^3 \\ &= 1 - \theta\, \text{var}(\Delta)C_1 + O(\Delta)^3\end{aligned} \qquad (4c)$$

1092

1093 where var(Δ) is a measure of the divergence in the DC mechanism between the hybridizing species,
1094 equal to $E(\Delta^2)$, and C_1 a positive constant equal to the inverse of $4t_m^2$.

2.2.2 Contribution of a derived stratum (stratum 2)

The male/female ratio contributed by the loci in stratum 2 can be approximated assuming that in
species A the Y genes are not degenerate and remain fully expressed (i.e. $\bar{t}_m = \bar{t}_f = c_X = c_Y = 1$),

while in species B the genes are degenerate, silenced, and dosage compensated (with a given t_m value,
i.e. without loss of generality $c_Y = 0$, $c_X = 2/t_m$, $t_f = t_m/2$). The reciprocal crosses are asymmetrical
for males, depending on which Y they inherit. In cross 1 (male B x female A), the Y in hybrid male
offspring is carrying degenerated genes, but in cross 2 (male A x female B), it does not.

Assuming weak stabilizing selection, we find, for the effect of one gene:

1103

1104
$$\rho = 1 - \theta \bar{C}_2 + O(\theta)^2 \tag{5}$$

1105

1106 Where C_2 is a constant equal to C_{2a} in cross 1, and to C_{2b} in cross 2 and to $\bar{C}_2 = (C_{2a} + C_{2b})/2$ on
1107 average over the two types of crosses, with:

1108

1109
$$C_{2a} = \text{Log}\left[\frac{1+t_m}{4}\right]^2 - \text{Log}\left[\frac{(2+t_m)^2}{8t_m}\right]^2$$
$$C_{2b} = \text{Log}[\frac{(1+t_m)(2+t_m)}{4t_m}]^2 - \text{Log}[\frac{(2+t_m)^2}{8t_m}]^2 \tag{6}$$

1110

1111 C_{2a} and C_{2b} are positive (except in a very small region around t_m=3 where C_{2a} is very close to zero,
1112 Fig S10b). C_{2a} could also be negative for very low values of t_m ($t_m < 0.17$), but these situations are
1113 probably not relevant as they imply varying cis regulator strength by an order of magnitude, and are
1114 far outside values consistent with extent DC systems (ie. t_m between 1 and 2). Fig S10b shows the
1115 value of these constants in the plausible range $0.5 < t_m < 4$.

1116

1117

1118 ### 2.2.3 The overall effect of ancestral and derived strata

1119 Hence with weak stabilizing selection and at small divergence, the male/female fitness ratio among
1120 F1 offspring, overall, is to leading order close to

1121

1122
$$\rho = 1 - \theta[k_1 C_1 \text{var}(\Delta) + k_2 \bar{C}_2] \tag{7}$$

1123

1124 With k_1 genes on the shared ancestral stratum and k_2 genes in the recently evolved stratum in one of
1125 the two species, where C_1 and $\bar{C}_2$ are positive constants that depend on the DC mechanism, and var(Δ)
1126 is a measure of the divergence in the DC mechanism between the hybridizing species. Hence, HR due

to disruption of DC can be caused by the portion of the Y that is degenerate and compensated in both species (provided they exhibit some divergence in DC), and by the portion of the Y that is degenerate and compensated in only one of them. Importantly, the fitness effect in hybrids directly scales with the size (i.e. the number of genes) of the non-recombining strata (shared or recently evolved), everything else being equal. However, this is not strongly discriminating with other theories. Such scaling is also expected, for instance, for the dominance or the meiotic drive theory.

In a drosophila-like situation where dosage compensation is achieved by overexpressing the X in males (t_m around 2), we have:

$$\text{Cross 1:} \frac{W_{male}}{W_{female}} = 1 - \theta[0.06k_1 var(\Delta) + 0.08k_2] \tag{8a}$$

$$\text{Cross 2:} \frac{W_{male}}{W_{female}} = 1 - \theta[0.06k_1 var(\Delta) + 0.16k_2] \tag{8b}$$

while in a *C. elegans* or mammal case (t_m around 1):

$$\text{Cross1:} \frac{W_{male}}{W_{female}} = 1 - \theta[0.25k_1 var(\Delta) + 0.47k_2] \tag{9a}$$

$$\text{Cross2:} \frac{W_{male}}{W_{female}} = 1 - \theta[0.25k_1 var(\Delta) + 0.15k_2] \tag{9b}$$

In all cases, Haldane's Rule occurs and combines a regulatory effect of the genes that are degenerate in one species and not the other, and a regulatory effect of the genes that are degenerate in both species (but not compensated exactly in the same way in the two species). The two effects combine and contribute to HR.

Loci that are degenerate in one species but not in the other generate an expression mismatch in hybrids, even if trans-acting factors are codominant. In the range of known DC mechanisms, i.e. with t_m values between 1 and 2, hybrid females will tend to show slight overexpression, males from cross 1 will show relatively severe underexpression, and males from cross 2 relatively severe overexpression (Fig S7-S9).

For loci that are degenerate in both species, the effect comes from the fact that genes may exhibit a quantitative difference in the way they are dosage compensated (since achieving optimal DC can be obtained by a different combination of cis and trans effects, i.e. different t_m values). This difference will be however more buffered in females (where cis effects are averaged over the two X) than in males (where cis effects are not averaged and only expressed from a single X, Fig S8). In both males and females, (autosomal) trans-effects are equally averaged. In species pairs sharing an identical and chromosomal-level DC mechanism, this effect is expected to be relatively minor, since genes should not exhibit quantitative differences in the DC mechanism. However, in many species, many genes escape this global DC mechanism and exhibit gene-level expression control [98–104]. These genes, if sufficiently numerous on the X could therefore also contribute to HR. Current evidence indicates that chromosome-wide DC is the exception, rather than the norm [58–60].

Finally, in systems where DC occurs around a Drosophila-like situation (t_m around 2), hybrid males
inheriting the most degenerated Y (from cross 1) suffer more than males inheriting the less
degenerated Y (from cross 2). The reverse occurs in systems where DC is close to a C. elegans/mammal
type (t_m around 1). However, the difference will manifest only in species pairs showing Y with different
degrees of degeneration (i.e. only if $k_2 \gg 1$ and larger than k_1).

2.3 Case of unbalanced females

In several Drosophila experiments, the fitness of F1 females carrying 2 attached X from the same
species (XXY females) was investigated to test the dominance theory of Haldane's rule. In these cases,
it was expected that homozygous females (carrying two X from the same parental species) would show
a large fitness reduction, like hemizygous males (but unlike standard heterozygous XX females). These
crosses have revealed that when HR was about F1 fertility, unbalanced females remained fertile (unlike
F1 males), while when HR was about F1 viability, unbalanced females were inviable (like F1 males).
These results generated considerable discussion [1,2,4,16,17,95,105,106].

Using the same approach as above, and assuming perfect dosage compensation in the parental
species, we can first compute the expected fitnesses of the different F1s for genes located on stratum
1 (shared between the two hybridizing species), assuming a small divergence in their regulatory traits
(small Δ) :

1181

1182
$$W_{XY} = W_{XXY} = 1 - \theta(\Delta/2t_m)^2 + O(\Delta)^3 \tag{10a}$$

1183
$$W_{XX} = 1 - \theta(\Delta/2t_m)^4 + O(\Delta)^5 \tag{10b}$$

1184

1185

For genes in that stratum, we find that
unbalanced females present the same
fitness reduction as males, much larger than
the one seen in regular F1 females. For
genes located on stratum 2, degenerate and
compensated only in species 2, and noting
X^d the X from species 2, we find that the
fitness reduction is proportional to θK,
where K is a constant that depends on how
X^d is dosage compensated (i.e. it depends on

the value of t_m). The different values of K are straightforward to compute and are illustrated in Fig.
S10c for the different F1 in the reciprocal crosses. Unbalanced females have a lower fitness than
standard females except near t_m =2 where the difference is small. Overall, combining effects in stratum
1 and 2, we can therefore conclude that this model predicts that unbalanced F1 females show a greater
fitness reduction compared to standard females. The comparison to males depends on whether genes
in stratum 1 or 2 contribute. For genes on stratum 1, the fitness reduction is the same as in males. For
genes on stratum 2, it depends on the direction of the cross and t_m values (see Fig. S10c). Overall, these
results indicate that unbalanced F1 females should often present a large reduction in fitness, much
more similar to that of F1 males than that of F1 females. This result holds for viability. For sterility, the
fitness of males and females (balanced or unbalanced) will differ, if genes involved in sterility are
expressed in a sex-specific manner (see next section for the effect of such genes). As shown below,

male-limited genes will cause a fitness reduction in F1 males, but female-limited genes will not cause
a fitness reduction in F1 females. This would therefore explain the contradictory results observed for
the *Drosophila* experiments mentioned above where unbalanced females are fertile when HR involves
sterility but are inviable when HR involves viability [1,105].

2.4 Case of sex-limited genes

We can consider the case of fertility genes. We consider that these genes are expressed in only one
sex (they are e.g. expressed in the germline), contrary to viability genes, which in the vast majority of
cases, will be expressed in both sexes). We will take male as the heterogametic sex as in XX/XY species,
but the same argument can be made by switching sexes for ZZ/ZW species.

Female-limited genes can obviously be lost from the Y, as they are not expressed in males. Apart from
this, their regulators can diverge between species, like in the case of autosomes, generating a
consistent but small decrease in fitness in F1 (as in Eq. 3). Given the lower effective population size on
the X (compared to autosomes), and the weaker selection pressure overall (everything else being
equal, sex-limited genes are only selected in one sex), we might nevertheless expect that the
regulatory divergence could somehow be faster on the X compared to autosomes (as seen in the
results, compare the female fitness drop illustrated on Fig 1E to that on Fig 1A).

Male limited genes can be lost from the Y (like genes expressed in both sexes), but they can also be
lost from the X (see empirical example in [107,108]). Indeed, there is no requirement that they remain
functional in XX females (contrary to genes expressed in both sexes that can only evolve Y silencing [35]).
Diploid expression is unstable after recombination arrest, but silencing can occur both ways.
Simulation results confirm this finding. However, despite this apparent symmetry, X and Y silencing is
not occurring at an equal rate. Male-limited genes are more likely to be silenced on the Y than on the
X (and indeed such cases are documented[109–111], note that it does not contradict the observation that
most genes remaining functional on the Y are male limited). The bias is for instance close to 6:4 in a
non-recombining stratum with 50 male-limited genes, but it is 9:1 in a stratum with 500 male-limited
genes (in both cases the simulation had a population size of 10 000 individuals, and the X or Y silencing
occurred relatively quickly). Presumably, the bias increases with the relative ease at which deleterious
mutations initially accumulate on the Y versus X once recombination has stopped (due to selective
interference), preferentially pushing the regulatory feedback loop in that direction. In all cases,
because one copy is silenced, these male-limited genes will evolve dosage compensation to stay
expressed at the right level in males (i.e. their cis-regulators on the X or Y will coevolve with the trans-
regulator expressed in males). In an F1 hybrid, when the focal gene has been silenced on the same
chromosome in both species, they will therefore cause a reduction in fitness that is similar to the case
of a gene expressed in both sexes (i.e. Eq 7 applies). If p is the chance that a gene is silenced on the Y,
and $1 - p$ on the X, this situation has $p^2 + (1 - p)^2$ chances to occur. If the focal gene is silenced on
the Y in one species, but on the X in the other, male F1 will however suffer from a larger fitness
reduction. In one direction of the cross, the F1 male will receive two silenced copies, and its fitness will
therefore be reduced by a factor $1 - s_{max}$. This has $p(1 - p)$ chances to occur. For an essential gene
involved in male fertility, a single case like this would be sufficient to cause complete male sterility. In
the other direction of the cross, the F1 male will receive two non-silenced copies, which will correspond
to severe overexpression. Its fitness will be reduced by a factor $(1 - \theta \log(2)^2)$, which is close to $(1 - \theta/2)$.
$\theta/2)$. This has also $p(1 - p)$ chances to occur. Simulation starting with an XY non recombing pair with
500 fully expressed male-limited genes leads to a dramatic reduction of F1 male fitness, due to this
sorting effect (Fig 1E, unsorted). Some genes become silenced on the X and some on the Y, but they
are not the same in the two diverging species, resulting in many mismatches. Note that genes are not
equally likely to degenerate on the Y and on the X. The former is much more likely (Fig S11), consistent

with the idea that the lower effective population size of the Y and the absence of recombination and
selective interference on the Y bias silencing and degeneration of the Y copy in this, otherwise,
symmetrical situation. A simulation starting with 500 male-limited genes that are already sorted leads
to an F1 male fitness reduction closer to that observed for the XO simulation with genes expressed in
both cases. The effect is stronger, however (compare figures 1D and 1E), as male-limited genes
silenced on the X (and maintained on the Y) will exhibit faster regulatory divergence, due to their lower
effective population size (3 times lower than that of genes silenced on the Y and maintained on the X).

Overall, we thus predict a stronger effect on HR of male-limited genes than of genes expressed in both
sexes. The argument was made for males but would apply to females in ZZ/ZW species. This finding
indicates that our regulatory theory could account for the importance of cases of sterility in HR,
without having to suppose a 'faster male' theory. It would also better explain why HR is not weaker in
Lepidoptera where males are homogametic [a difficulty of the faster male theory noted by Presgraves
[96]].

1266

Fig S11. (a) Number of male-limited dosage compensated genes evolving on the
X and Y chromosomes. In the "unsorted" simulations shown on Fig 1E, some
male-limited genes degenerate on the Y (and stay on the X and become dosage-
compensated, gray dots), and some on the X (and stay on the Y and become
dosage compensated, black dots). The figure shows how many genes (y-axis in
log-scale) end up in each category through time (x-axis in generations, in log-
scale). Despite the symmetry of the regulatory instability, many more genes
degenerate on the Y, probably due to selective interference on the non-
recombining Y and lower effective population size of the Y. (b) Corresponding
fitness of the heterogametic F1 (y-axis, log scale) through time (x-axis in
generations, in log-scale). The fitness decrease is fast due to the independent
degeneration of genes on the X or Y in different species.

2.5 X-autosomes versus Y-autosomes hybrid incompatibilities

The issue of a possible role of Y-autosome or X-Y interaction for Haldane rule has been proposed by
Muller[20] and often discussed. Several experiments showed that these interactions occur sometimes
and contribute to HR on sterility[112–115]. However, they are far from being the norm[116], and not an
explanation for Haldane rule on viability (since Y chromosomes have usually very limited effect on
viability) or Haldane rule in XO species lacking a Y. Most work on HR has therefore mainly focused on
X-autosomes interactions.

Most of our results rely on such X-autosomes interactions. Specifically, they concern the misregulation
of genes on the X, in crosses between species where at least one evolved silencing and degeneration
of those gene copy on the Y. However, it is useful to point out that in some F1 hybrids, the
misregulation may involve the non-silenced and non-degenerate Y copy of the gene inherited from the
species in which this gene is still on a recombining region of the Y (see e.g. Fig 7a). In such case,
overexpression in F1 males results from the presence of two fully functional copies of the gene (on
both the X and Y), in presence of partial dosage compensation (inherited from the species where the
Y copy is silenced). In this case, the X and Y gene copies play symmetrical role, and arguably, this is not
best described as a simple "X-autosome" interaction. The fitness reduction involves the autosomal
trans-acting factors, and both X and Y copies of the gene. However, the misregulation emerges because
in one species, the Y copy of the gene was silenced and the remaining X copy evolved DC. In this sense
only can it be presented as a problem of X-linked dosage compensated genes.

However, as simulations involving genes with sex-limited expression show, sometimes genes can also
degenerate on the X (or Z in ZW species). Example of genes lost on the X and with a local DC have been
reported[117]. In this case, the dosage compensated (functional) genes may be located on the Y (or W),
and their misregulation could also contribute to explain HR on sterility (see previous section). In the
simulations illustrated on Fig 1E, unsorted case, 10% of genes stay functional on the Y and degenerate
on the X (in a ZW system, similarly, a handful of female-limited genes would stay on the W and
degenerate on the Z). These genes would contribute to HR through Y-autosomes interactions. The
reason these genes contribute to HR is the same than for other sex-linked genes in our model: the
heterogametic sex will average autosomal trans-regulatory effects but not cis Y or W regulatory effects
(since a single cis functional sequence is inherited in the heterogametic sex). Hence, our model may
explain why Y / W linked genes could contribute to HR on sterility, and why such cases are likely to be
rare.

The argument about these Y effects could be somehow generalized. Our simulations concern genes
present on the ancestral autosomal pair and degenerating either on the X or on the Y. Yet, the
regulatory averaging imbalance between the homo and heterogametic sex in F1 hybrids could also, in
principle, apply to genes evolving *de novo* on the Y or the X. As far as the regulation of expression is
determined by the interaction between a sex-linked (X or Y) cis-acting factor and an autosomal trans-
acting factor, then misregulation is likely to occur more strongly in the heterogametic F1, due to the
difference in averaging between cis and trans effects in hetero and homogametic sex. A particular and
noteworthy case could involve sequences evolving on the Y (and absent on the X) and acting as
'chromatin sinks' (as reported in several studies [118]) for an autosomal transcription factor or any
autosomal factor that affects chromatin state. Such sequences could have evolved due to an
accumulation of transposable elements on the Y, or to regulate specific male-limited genes as often
reported[119–122]. The strength of a such a chromatin sink could be modeled in the same way as a cis-
regulatory strength. Consider, for instance, a species without this chromatin sink on the Y. In that
species, the autosomal factor is free to interact with its normal target elsewhere in the genome.
Consider now another species in which a sink sequence has evolved on the Y. Because this sink
sequence "absorbs" some of the autosomal factor, it is less available for its normal function, which
presumably would be detrimental. Hence, we would expect that, in this second species, this autosomal
factor has evolved to become overexpressed to compensate for this sink effect (and reach the proper
amount of factor not captured by the sink sequence). F1 males, however, will inherit either the Y with
or without the sink sequence (depending on the cross direction), but always an "averaged" production
of the autosomal factor. It will always result in over or underexpression of that factor in F1 males. In
F1 females, however, no such effect will occur (since in this example the sink effect is specific to males

having a Y with the sink sequence). Hence, it is theoretically and empirically possible that the effects
we describe on genes extends to any cis-regulatory element on the X or Y, not just coding genes and
their cis-regulatory sequences that were inherited from the ancestral autosomal pair. In particular, all
Y/W specific sequences interacting with autosomal factors could possibly play a role in HR in our
theory, including sequences acting as chromatin sinks.

3 Comparison to other theories

3.1 Comparison of the regulatory theory with the dominance theory

The dominance theory for Haldane's rule proposes that recessive incompatibilities occur, such that, in
an hybrid F1 autosomal background, F1 females are fit while F1 males are unfit. The reason why F1
females are fit is that incompatibilities carried on the X are "recessive", meaning that if the X were
homozygous (in an F1 hybrid background), these incompatibilities would manifest themselves. If such
females could be produced in a backcross or through genetic manipulations (such as the F1 females
with attached X in *Drosophila*), they would therefore have low fitness. The theory further considers
that hemizygous males express the incompatibilities because, having a single X, these incompatibilities
carried on the X cannot be masked. Different models proposed different ways to achieve this pattern.
For instance, different beneficial alleles can fix in different populations and some can turn out to be
incompatible in the sense just explained (those that turn out to be incompatible and dominant reduce
fitness in all F1 hybrids and thus do not create a sex bias) [22,95].

In the model of stabilizing selection proposed by Barton [24], combinations of phenotypically dominant
mutations can contribute to bring a population closer to an optimum. When different dominant
mutations (on autosomes and on the X) coming from two parental species, who adapted
independently, combine in an F1 female, they all add-up to result in a near optimal phenotype (since
all the mutations coming from each parental species independently add-up, as they are all dominant
and have additive effects between loci). In contrast, F1 males miss the dominant mutations that
occurred on the X of one parental species, causing a departure from the optimal phenotype (this model
assumes no mutation on the degenerate Y). "Homozygous" F1 females would also miss all the
mutations from the X of one of the two parental species (while having the dominant mutations on
autosomes from both species), resulting in low fitness. Note that this model does not require
specifically a different number of dominant mutations on the X of the two species.

Another model of stabilizing selection has been proposed that includes additive mutations (within and
between loci), but assumes that mutations in males have twice the phenotypic effect when
hemizygous (although not explicitly mentioning expression levels, this assumption might be viewed as
a way to account for the occurrence of dosage compensation)[25,26]. As in Barton's model, F1 females
remain close to the phenotypic optimum because all mutations are additive. F1 males, in contrast,
depart from the optimal phenotype because the mutation on their X mismatch with half the additive
autosomal allele (coming from the same parental species as the X), while the other half of additive
autosomal alleles do not add up with the missing X (from the other parental species). Here too, the
"homozygous" females exhibit the same phenotype, and low fitness, as the F1 males.

All these models do not propose the same processes to explain HR. In particular, they consider
different mutational input (the occurrence of recessive incompatibilities, the occurrence of
phenotypically dominant mutations or the occurrence of additive mutations that have twice their
phenotypic effect when hemizygous, respectively). They have one pattern in common, besides
predicting HR: they all predict that a female that would be homozygous for a parental X in an otherwise
F1 background has a lower fitness than a regular F1 female. This implies a dominance effect on the X,
i.e., heterozygotes have a higher fitness than the average of the two homozygotes. In all cases the

fitness of hemizygous F1 males is also found to be the same as the fitness of females made homozygous
for the X. For this reason, and despite being based on different types of mutations, they can all be
viewed as specific cases of a "general" dominance theory.

Our theory is based on another kind of mutational input (additive mutations on cis and trans-
regulators). Besides offering a mechanistic explanation of HR (specifically based on the misregulation
of dosage compensated genes) and an evolutionary explanation of the pattern (the incompatibilities
emerge from cis-trans coevolution and the degeneration and dosage compensation of sex-linked
genes), our theory does not fit well in the "general" dominance theory mentioned above. In our
regulatory model, F1 males show a reduced fitness compared to F1 females (Haldane's rule), but the
reason is not always a "recessive" effect, in the sense described above. Unlike in the dominance theory,
the effect of the F1 genetic background differs between males and females: they express different
trans-acting factors, both in the parental species and in the male and female hybrids (Fig S12). Hence,
the fitness of hemizygous males is not necessarily the same as the fitness of "homozygous" females.
Hemizygous males present the same fitness reduction as "homozygous" females only for genes located
in ancestral strata (Eq 10), but not in derived strata (Sup. mat. 2.5, Fig S10c). This is particularly clear
in the example shown in Figure S7a, where F1 females that would be made homozygous for the X
would retain optimal expression, contrarily to F1 males. Generally, for derived strata, the disruption of
dosage compensation always impacts more the heterogametic than the homogametic F1, although
the effect varies with the type of dosage compensation and the position of the loci involved (on an
ancestral or derived stratum, see Eq. 7). The difference between the fitness of F1 males and
"homozygous" females also varies with type of dosage compensation and the position of loci, although
"homozygous" females tend to have a lower fitness than F1 females (this is the case in the examples
shown on Figures S7b and S8). Note too that in the case of derived strata, incompatibilities are, strictly
speaking, not simple X-autosomes incompatibilities (see Sup. Mat. 2.5), as corresponding genes on the
Y are not degenerate in one of the two parental species.

1402

1403

Fig. S12. **Simplified fitness patterns in the dominance versus regulatory theory.** The
chromosome with a hook represents the Y. A key difference is that the fitness of
"homozygous" females differs between the two theories (indicated by "BC" as these
females can be obtained by backcrossing or by creating "attached X" F1 females in

Drosophila). In the dominance theory, "homozygous" females exhibit the same
fitness reduction as F1 males. In the regulatory theory, male and female fitness can
be uncoupled since they express different trans-regulators. The sex asymmetry
results from lack of averaging of cis-regulator effects in males (see above).

3.2 Criticisms of the dosage compensation hypothesis for Haldane's rule

The list below presents past arguments against a theory of HR based on the disruption of dosage
compensation. It has to be taken with precaution for three main reasons.

First, a firm theory of Haldane's rule based on the disruption of DC has never been made, so most of
these past arguments were made against a verbal and relatively vague theory. For instance, the relative
level of expression disruption of autosomal vs. sex-linked genes in hybrids is sometimes mentioned as
a decisive observation. Yet, this comparison is mostly uninformative. Autosomal misexpression
reduces the fitness of both sexes in F1, and is therefore irrelevant to HR. It does not indicate whether
the cause of HR is related to misexpression of sex-linked genes. If for instance DC is controlled by a
global mechanism, such as in mammals or *Drosophila*, it might be constrained and slow evolving
compared to autosomes. In this case, only few genes escaping this global DC might contribute to HR
(including sex-limited genes in the germline not regulated by this somatic global DC). As a result,
misregulation may appear larger overall on autosomes than on sex-linked genes, but this observation
would not rule out that the cause of HR is misregulation of sex-linked genes.

Second, the idea that HR may be caused by DC disruption, made in the past, is much narrower than
our regulatory theory. The idea usually considered is that the mechanism of dosage compensation is
disrupted in hybrids, due to a divergence in the genes *directly* involved in the DC pathway. This is much
narrower than in our regulatory theory, where the reduction of hybrid fitness is caused by the
misregulation of gene expression *for all genes that are dosage compensated*, not only for genes directly
involved in DC regulation. Our theory potentially includes most genes on the non-recombining part of
the X or Z, not just the genes directly identified in the pathways controlling DC (the latter can
nevertheless play a role in the misexpression of dosage compensated genes). If for instance, again, DC
is controlled by a global mechanism, such as in mammals or *Drosophila* (although local DC probably
precedes global DC, as seen on neo sex chromosomes[123]), only genes escaping this global DC might
contribute to HR, including sex-limited genes in the germline not regulated by this somatic global DC.
Their misexpression would play a key role in the fitness reduction of heterogametic F1, but they would
be entirely unaffected by the global DC mechanism, and entirely unrelated to the control of this DC
regulation. Note also that defining dosage compensation for sex-limited genes can only be made here
relative to the ancestral expression level (before the evolution of recombination suppression). Since
such genes are only expressed in one sex, male-female comparisons cannot be made, as usually done,
to assess DC. In these cases, the comparison of expression levels in parental species and the hybrids
would be the important comparison to make.

Third, the genetics of post-zygotic isolation is a broader topic than the genetic cause of HR. Many
incompatibilities can impact hybrid fitness, in addition to those causing HR (i.e. not specifically reducing
the fitness of the heterogametic sex). Importantly, such incompatibilites can be caused by many other
mechanisms than expression disruption. Even in the context of HR, exceptions to the rule are known
and relatively well understood (in particular nucleo-cytoplasmic incompatibilities[74]). Hence, assessing
the merit of the different theories of HR based on a specific gene disrupting hybrid fitness in a specific
cross must be done carefully. Documenting broad patterns such as those mentioned on Table 1 could
be a more informative and more robust way of comparing the different theories.

With these caveats in mind, here are 13 arguments made in the literature against a (relatively narrow)
theory based on the idea that the cause of HR is due to the direct disruption of the dosage
compensation pathway in hybrids.

The 1st major critique of the DC hypothesis is that the rule was observed in groups in which DC was
allegedly absent (such as in birds and lepidopterans [4,6]). However, DC has since been documented in
these groups[58,60], although global DC is perhaps less frequent in ZW species[124].

The 2nd argument is based on an experiment with *Drosophila* [51]. However, this experiment was based
on crosses with mutants in which DC was supposedly fully functional, but without showing that this
was the case. Hence, the results presented could not really discard DC disruption as a cause of HR.
Specifically, it could not discard the possibility that hybrid male sterility was caused by a failure in DC
downstream of *Sxl* regulation (which is the sex-determining switch in Drosophila).

The 3rd argument is that DC evolves very slowly, being an essential function under strong constraints
[52]. However, as our results show, divergence in DC can readily occur, even with substantial stabilizing
selection on expression levels. Note that DC evolution could be triggered by different phenomena,
including coevolution with cytoplasmic bacteria targeting DC pathways to achieve male killing [125–127].

The 4th argument proposes that if Haldane's rule was caused by DC disruption, cis and trans regulators
involved in DC should exhibit signs of divergence between species exhibiting Haldane's rule. Yet, Jaffe
and Laird [128] reported unpublished data where the X-linked *D. pseudoobscura Hsp82* gene remained
dosage compensated when transformed into various autosomal sites in *D. melanogaster,* suggesting
conservation of the cis-regulatory elements involved in DC for ~20 million years. Indeed, if cis-
regulators involved in DC are highly conserved, they could not cause Haldane's rule [4,51]. However,
further evidence showed that this conclusion, besides applying only to *Drosophila*, was premature.
More recent investigations revealed that both cis and trans regulators involved in DC were actually
fast evolving in *Drosophila melanogaster*. This is the case with *msl*, *mof,* and *mle* trans-acting genes
[52,83] as well as cis-acting binding sequences on the X [53]. The case of male lethality in *D. melanogaster* x
D. simulans hybrids is not clear-cut. Some studies support the role of DC in male lethality [129,130] while
others challenge this interpretation [131]. In this cross, both males and females show a reduction in
viability, but to a different degree depending on temperature. However, the effect of mutants is not
always evaluated in this GxE context. Barbash shows that deletion of mel-specific DC genes—ie, forcing
Xmel/Ysim hybrid males to develop using only *simulans* DC complex components—does not
exacerbate F1 male lethality when this lethality was first rescued by lhr mutation. The test is based on
the premise that the lhr rescue occurs and that playing with DC genes should show an independent
effect when manipulated. However, lhr might be acting by interacting with these DC genes, so there
may be no clear 'independent' manipulation of the DC phenotype in the first place in the experiment.
Barbash also argues that DC disruption is unlikely because a lower-than-expected number of genes
downregulated in lethal F1 males are located on the X chromosome. However, only the misregulation
associated with sex chromosomes will have a differential fitness effect on males vs females. HR
depends on the presence of the latter, not on the proportion of misregulation of X versus autosomes.
Whether there is more or less than "expected" misregulation on the X says nothing about the cause of
HR. What can be noticed, however, although it could be a coincidence, is that major genes involved in
these hybrid incompatibilities interact with the dosage compensation complex. For instance, *Lhr*
interacts with HP1, a chromodomain-containing protein that   izes to heterochromatic regions of
chromosomes [132] and is also involved in DC [133]. Nuclear pore complex proteins also cause hybrid male
lethality [134,135], and are also involved in DC [136].

The 5[th] argument is a refinement of the fourth. After the accumulating evidence demonstrating fast DC evolution in *D. melanogaster*, Tang and Presgraves [135] suggested that this phenomenon was not general and limited to that species. They concluded that the limited evolution of the MSL complex in *D. simulans* could not well explain the rapid evolution of Nup160 and the other Nup107 subcomplex genes in that species (these autosomal simulans alleles being involved in male hybrid lethality through incompatibility with the melanogaster X). This may be the correct interpretation, and this specific case may be regarded as mere exception to the general case. However, this argument can be nuanced by several points, that might warrant further investigation. First, the *mof* gene involved in the DC complex does show evidence of rapid adaptive evolution in *simulans* [83]. Second, Nup160 and the other Nup107 subcomplex genes are known to be involved in DC in *Drosophila* [136]. Third, DC disruption is not necessarily limited to the coevolution of the MSL complex and its cis-binding sites. For instance, MSL cis-binding sites could change location on the X, thereby changing the pattern of DC, as shown by the rapid and extensive turnover of individual binding sites of roX lncRNAs, which are essential for *Drosophila* DC [56]. Fourth, some genes are dosage-compensated but non-MSL-binding, suggesting that MSL is not the only mechanism for achieving DC [99]. This is also particularly the case for genes expressed in the germline, impacting male fertility in *Drosophila* [65].

The 6[th] argument, also by Tang and Presgraves [135] and also concerning Nup160 in *D. melanogaster*, note that Nup160sim kills both Xmel/Ysim hybrid males and Xmel ·Xmel/Ysim hybrid females, and in a similar way, suggesting a common cause. This would exclude the role of DC, on the premise that DC only concerns males in *Drosophila*. Like above, this may be the correct interpretation, and this case may well be an exception (there are several other known exceptions to HR in other groups, involving specific Y effects or incompatibilities with mitochondria, etc.). However, interestingly, our results show that, especially for ancient Y, unbalanced females should show an equal decrease in fitness to males, especially for viability (Eq. 10). (For hybrid sterility, it is easy to uncouple the fitness effect seen in males and unbalanced females: it is sufficient to have male-limited genes, as discussed above). This result relies on a quantitative DC trait divergence between species. In our model, the *Drosophila* DC system corresponds to a particular case where the strength of X cis-regulators does not change and where male trans regulators double X expression in males (case $t_m = 2$ in our notation). Any quantitative departure from this point (where the strength of X cis regulator changes) involves a correction in females. In a quantitative model, DC is a phenomenon that always involves both sexes, as long as t_m is not exactly 2. Hence, anytime males will show DC disruption (because of t_m divergence), so will unbalanced females.

The 7[th] argument has not directly been made against the DC hypothesis but against the dominance theory. However, this issue also concerns the DC disruption hypothesis. Haldane's rule is observed in species lacking heteromorphic sex chromosomes. In particular, in the genus *Aedes*, Haldane's rule applies to many interspecific crosses [3]. Yet, *Aedes* have homomorphic sex chromosomes with a sex locus. A priori this observation would rule out theories based on hemizygosity (dominance theory) or DC. However, recent genomic data have revealed that in *Ae. aegyptii*, the sex "locus" is a 1.5 Mb region with 30 genes in a 100 Mb non-recombining region encompassing the centromere of chromosome 1, that diverged between males and females. Hence, the premise that these species have a sex chromosome with just a sex-determining locus seems erroneous. More work is required to evaluate the extent of hemizygosity and the occurrence of (local) DC in these species, but the conclusion favoring the "faster male" hypothesis based on these *Aedes* data certainly requires re-evaluation.

The 8[th] argument is that one might expect a breakdown in dosage compensation to affect the homogametic more than the heterogametic sex in cases where DC involves an active mechanism in the homogametic sex, such as in mammals or *C. elegans* [6]. This argument does not hold up

theoretically. The larger impact of DC disruption in the heterogametic sex holds up irrespectively of the type of DC (Eq. 7). Despite showing extensive misregulation in *Caenorhabditis* hybrids, especially involving males, and involving X-autosome and cis-trans coevolution, a hallmark of DC disruption, Sánchez-Ramirez et al. [46] excluded DC as an explanation based on the observation that expression levels were not strongly disrupted in females. This reasoning is based on the idea that DC is a female phenomenon, as it works by halving X expression in females. This argument (similar to that of Laurie) does not hold, since it is in fact both a male and a female phenomenon (the halving in females corrects X overexpression evolving in both males and females). Even with a *C. elegans*-like system, DC disruption will be larger in males than in females (Eq. 7).

The 9th argument is that DC could play a role, but only in the case of species with fast and ongoing Y degeneration, dramatically reducing the scope for the general application of the DC hypothesis [50]. Indeed, the effect of DC disruption has been understood as resulting solely from the mismatch of having non-functional (but dosage-compensated) genes on the Y in one species but not in the other [50]. However, this is an incomplete picture, and overly restrictive, as DC disruption may also occur for genes that are degenerate in both species (Eq. 4, Fig 11).

The 10th argument is that the DC hypothesis is supposed to entail that "*any* anomaly (not just sterility and inviability) appearing in hybrids results from a disturbance in dosage compensation" [4]. Since morphological anomalies in hybrids do not seem to be more severe in *Drosophila* hybrid males, it would then indicate that DC disturbance is not occurring [4]. This argument seems greatly exaggerated, in the sense that there is no reason to suppose that all hybrid problems are DC-related. For many traits, the genes involved may not be sex-linked, so there is no reason to expect that they all show an HR pattern.

The 11th argument is that DC disruption could hardly predict partial hybrid sterility or inviability [4]. This would be true if DC was an all-or-nothing phenomenon, but, as our results show without ambiguity, a partial fitness reduction is very easy to obtain after short divergence. As is now better understood, DC occurs often on a gene-by-gene basis, even when a global DC system is in place [58,137]. This observation was not available at the time this critique was formulated, and is thus, now, less convincing than it was.

The 12th argument is that "it is difficult to envision how the failure of dosage compensation could explain the sterility of hybrids that are viable and morphologically normal" [4]. In this view, disruption of dosage compensation should affect hybrid viability more than fertility [17]. This argument is close to the previous one, in the sense that DC is viewed as a global chromosomal level process. However, it would be fairly easy to observe sterile but viable hybrids: it only requires that DC disruption only involves a few sterility genes in recently derived species. The same explanation applies (and was applied) to the dominance theory: it would only involve the occurrence of recessive incompatibilities on sterility genes in recently derived species [95]. We also directly show how genes only expressed in the heterogametic sex (like genes involved in fertility) can produce a stronger fitness reduction in heterogametic F1, compared to genes expressed in both sexes.

The 13th argument is that, in *Drosophila*, the absence of dosage compensation in the male germline excludes its disruption as a cause of hybrid male sterility [62]. This argument does not take into account that some form of dosage compensation may be occurring in the germline, even if does not involve the global somatic DC mechanism. Some regulation seems to take place on both the X and Y in the germline, beyond the effect of MSCI [65,72]. If this regulation is local (on a gene-by-gene basis) instead of following the global somatic mechanism, it would rather facilitate the evolution of regulatory divergence between species on those genes. We showed that male-limited genes could play a

1587　disproportionate effect on HR. This observation would tend to reinforce this conclusion. If DC
1588　divergence of viability genes is limited (because of the global somatic DC mechanism that remains
1589　constant), it would exacerbate the role of HR of genes expressed in the germline that escape this global
1590　regulation.