Title

Allele Sails: launching traits and fates into wild populations with DNA sequence modifiers

Authors

Michelle L. Johnson[1] (ORCID: 0000-0003-3646-8062), Bruce A. Hay[1]* (ORCID: 0000-0002-5486-0482), and Maciej Maselko[2]* (ORCID: 0000-0001-7831-848X).

Affiliations

[1] California Institute of Technology. Division of Biology and Biological Engineering.

1200 East California Boulevard, MC156-29, Pasadena, CA 91125

[2]Applied BioSciences, Macquarie University, North Ryde, NSW 2109, Australia

Contact info

*Correspondence: haybruce@caltech.edu, maciej.maselko@mq.edu.au

Abstract

Population-scale genome editing can be used to alter the composition or fate of wild populations. One approach to achieving these aims utilizes a synthetic gene drive element—a multi-gene cassette—to bring about an increase in the frequency of an existing allele. However, the use of gene drives is complicated by the multiple scientific, regulatory, and social issues associated with transgene persistence and gene flow. Alternatives in which transgenes are not driven could potentially avoid some of these issues. Here we propose an approach to population scale gene editing using a system we refer to as an Allele Sail. An Allele Sail consists of a genome editor (the Wind) that introduces DNA sequence edits (the Sail) at one or more sites, resulting in progeny that are viable and fertile. The editor, such as a sequence-specific nuclease, or a prime- or base-editor, is inherited in a Mendelian fashion. Meanwhile, the edits it creates experience a Super-Mendelian increase in frequency. We explore this system using agent-based modeling, and identify contexts in which a single, low frequency release of an editor brings edits to a very high frequency. We also identify conditions in which manipulation of sex determination can be used to bring about population suppression. Current regulatory frameworks often distinguish between transgenics as genetically modified organisms (GMOs), and their edited non-transgenic progeny as non-GMO. In this context an Allele Sail provides a path to alter traits and fates of wild populations in ways that may be considered more acceptable.

Keywords

gene drive, evolutionary rescue, Cas9, base editing, prime editing, retrotransposon, sex determination

Introduction

The ability to modify or build resilience into ecosystems is increasingly desirable to confront a range of challenges including vector-borne diseases, agricultural loss, the spread of invasive species, and climate change. One important tool to address these are population-scale genetic alterations. These changes can introduce beneficial traits into a population (population modification) or eliminate a harmful population (population suppression). For example, population modification could promote resilience of an endangered or threatened species by bringing currently beneficial genomic modifications to high frequency. It could also be used to introduce 'anticipatory' sequence changes designed to provide a benefit in a likely future environment altered by climate change or the introduction of an invasive disease vector. These possibilities are suggested by the growing number of contexts in which one or a modest number of alleles of large effect can produce significant phenotypic benefits for a population. Examples include a single locus that can confer heat resistance in mussels[1] and cattle[2], fungi resistance in certain plants[3], or varroa mite resistance in bees[4,5]. There are also collections of loci that have been suggested to contribute to coral resistance to heat[6,7], bird resistance to malaria[1,8], or frog resistance to chytrid fungi[9]. Allele modifications can also be used to reduce harm, for example by limiting the ability of insects to vector disease[10,11], or reducing the toxicity (poison production) of an invasive species such as cane toads. Finally, genetic alterations can be used to bring about population suppression, either directly[12] or by sensitizing pests to an outside stimulus.

One way to introgress important alleles into a population is via the release of individuals carrying the desired allele. A new allele under positive selection will spread, but may take many generations to reach high frequency depending on the size of the benefit and whether it is dominant, additive or recessive. Alternatively, if the allele is not under positive selection or if the release introduces transgenes meant to suppress the population, large and/or repeated introductions will be needed. These are resource intensive and impractical for many species. Additionally, controlled breeding and the large-scale introduction of these individuals into the target population could increase the frequency of other, unintended and possibly harmful alleles. It could also threaten the existing genetic diversity of the target population or subspecies.

These issues can be overcome via the use of synthetic gene-drives (SGD). SGDs are selfish genetic elements that spread to high frequency by biasing their own inheritance; they can include 'cargo' in addition to transgenes essential for the SGD's function. A wide variety of SGDs have been devised (reviewed in ([13,14])). However, the very features of a gene drive that makes it attractive—that it can rapidly bring to high frequency transgenes that may persist for extended periods in (and in some cases outside) the target region—create regulatory and social hurdles to their implementation.

Population-scale genetic alterations with phenotypic consequences can also involve more subtle modifications such as single base changes, or small insertions or deletions. Importantly, in some regulatory environments (Australia provides one example[15]) genome edits present in non-transgenic progeny of a transgenic for a DNA

sequence modifier such as a Cas9 nuclease are regulated as non-transgenic. Designation of a population carrying a high frequency of such edits as non-transgenic may facilitate regulatory approval and social acceptance. Here we explore a system that we refer to as an Allele Sail, which can, in the absence of gene drive, bring edits but not transgenes to high frequency for population modification, and suppression in certain contexts.

An Allele Sail consists of a chromosomally located genome editor (e.g. a site specific nuclease such as Cas9, or a base- or prime-editor. We call this editor the Wind, as it is responsible for pushing edits into the population. The editor is expressed in the germline (though the expression need not be germline specific) and introduces sequence modifications at one or more target sites located anywhere in the genome. The homozygous edited individuals are viable and fertile. While the editor is transmitted in a Mendelian fashion, the edits it creates (we call these edits the Sail) increase in frequency at a super-Mendelian rate as the editor encounters new unedited alleles in the germline each generation (Figure 1). We first consider the use of an Allele Sail in population modification, where we believe there are a variety of applications for conservation or infectious disease prevention. We then explore potential uses for population suppression in organisms that have sex determination systems in which the activity of a single gene is required for femaleness[16,17].

Figure 1. Graphical Representation of Allele Sail Components and Behavior. The Wind (an editor, in Blue) is inherited in a Mendelian fashion (left panel). When the editor is present in a germline that carries an unedited target locus located elsewhere in the genome, conversion to the edited state occurs at some frequency (middle panel). This pushes the Sail (the edited locus, in Pink) to higher frequency in the population (left and right panels).

Results

Population Modification

We consider the dynamics of a genome editor that introduces genomic edits at one or more sites, resulting in progeny that are viable and fertile. The editor is transmitted in a Mendelian manner, while the edits change in frequency as a function of

frequency of the editor, frequency of wild type alleles, fitness costs and editing efficiency. To explore the use of Allele Sails for population modification, we characterize behavior of the components using a discrete-time and generation stochastic model with a panmictic population (see methods for details). This type of model is often used to gain insight into population genetic processes and provides a format that allows comparison of methods for genetically altering populations.

We first consider ideal conditions: the editor alters a target sequence with 100% efficiency, and editing occurs in the germline and progeny of a female carrier as a result of maternal carryover of editing activity. The power of the Allele Sail system can be seen by comparing the frequency of edits over time when introduced in the presence or absence of an editor. These comparisons are shown in Figure 2A for an editor with no fitness cost introduced at a frequency of 10%, with the edit conferring either no fitness change, or an additive benefit or cost of 5% per allele. Recessive and dominant fitness costs/benefits are considered in Supplement Figure 1 and show largely similar dynamics. In the presence of an editor, edits with no cost or a benefit spread to allele fixation. In contrast, edits conferring a cost rise to high frequency (~75%) and then decline. In all editing scenarios the peak frequencies of the edits are much higher than those of the corresponding Mendelian allele alone.

The dynamics of edit frequency can be understood by considering the fate of the editor (Figure 2B). When the presence of edits has no effect on fitness the editor remains at its introduction frequency, continually generating new edits until fixation is reached. When the presence of the edit confers a benefit the editor increases in

frequency. This occurs because the editor spends more time in the presence of the higher fitness edited genotype (which it creates) than does its counterpart wildtype allele. The frequency of the editor plateaus when the edits are ubiquitous (allele fixation) because at this point all individuals have equal fitness. Conversely, when the edit results in a cost to carriers the frequency of the editor declines continuously since it now spends increased time in lower fitness edited individuals (so long as these never reach fixation) than does its wildtype allele counterpart, leading to its loss through natural selection.

The general relationship between introduction frequency and edit fitness costs/benefits on the frequency of edits is shown in heat maps which plot the frequency of edit homozygotes (Figure 2C) and carriers (homozygotes and heterozygotes) (Figure 2D) at the 50 generation time point. There is a large region of parameter space in which edits are pushed to very high frequency, and increasing the introduction frequency has the general effect of increasing the rate of spread, as well as the time spent at high frequency for those edits that do not reach allele fixation.

The presence of the editor may result in some fitness cost to carriers[18–20]. As such, we explored the context in which the editor but not the edits result in fitness costs. This scenario is illustrated in supplemental Fig. 2, which shows edit frequency at generation 50 as a function of introduction frequency and editor fitness costs, up to 10%. Costs on the editor cause its eventual loss from the population and thus reduce the parameter space in which a single introduction can push edits to high frequency, though increased introduction frequencies and/or multiple releases can compensate. In

some cases a guarantee of eventual loss may be desirable, such as when regulatory approval requires that transgenes do not persist in the population. In other contexts, in which the spread of edits to high frequency at minimal cost is the dominant consideration, it may be possible to take advantage of next generation editors that have increased specificity and reduced toxicity[20–23] to reduce any editor-associated fitness costs.

Figure 2. Behavior of a Neutral Editor & Non-Neutral Edit. A) The average allele frequency of edits after being introduced at 10% frequency. Introduced individuals

are either homozygous for the edit with no editor present, or homozygous for both edit and editor when the editor is present. The fitness benefit shown here is an additional 5% chance of survival for each copy of the edit present (additive benefits), where fitness cost is a 5% decrease in survival for each copy of the edit present. **B)** The average allele frequency of the editor, from the simulations shown in panel **A. C)** Average percent of the population that is homozygous for an edit after 50 generations, for various editor introduction frequencies and fitness costs associated with the edit. Pink tiles represent an average allele frequency of more than 99.9% of the population, across 20 simulations. Negative fitness costs represent fitness benefits, and boxed areas highlight specific allele frequencies to provide guidance for interpretation of the heat map colors. **D)** Average percent of the population that carries at least one copy of the edit, for various editor introduction frequencies and fitness costs associated with the edit.

Consequences of Altering Editing Efficiency

We have thus-far assumed that our editor has 100% efficiency in editing. CRISPR nucleases such as Cas9 can cleave and create loss of function (LOF) alleles (especially when using multiple gRNAs) in the Drosophila germline or the plant Arabidopsis thaliana at frequencies near 100%[24–31]. While base and prime editors also have significant levels of editing activity in Drosophila, they are not 100%; instead closer to 36% for a prime editor[32], and >90% for a base editor[20]. These encouraging rates notwithstanding, it is important to note that the editing enzymes used—nuclease,

reverse transcriptase, deaminase, and uracil-DNA glycosylase—are often derived from organisms that live in temperature ranges very different from those in which their use is intended, resulting in significantly reduced activity[20,33,34]. To explore these less-than-ideal scenarios, we model several representative examples in which the editor is introduced at a frequency of 10% into the wildtype population and has no associated fitness costs, for various editor efficiencies (between 15-100%), both with and without maternal carryover. These results are compared with a scenario in which edits are introduced directly into the population in the absence of the editor.

Figure 3 shows that decreasing the rate of editing from 100% to 50% or 15% still results in the rapid spread of a beneficial or neutral edit to high frequency by generation 50, as compared with the spread of a Mendelian allele on its own. A deleterious allele also undergoes a significant increase in frequency, though there is ultimately a decline (as when edits are created 100% of the time) when the frequency of the editor is so low it no longer generates edits faster than they are lost through natural selection. Even so, the peak frequency and time to decline can always be improved by increasing the editor introduction frequency (Supplement Figure 3).

Figure 3. Consequences of reducing editing efficiency for Population Modification. Comparison of edit frequencies over time when using editors with different efficiencies (15%, 50%, and 100%), for various fitnesses and presence or absence of maternal carryover (MC). Each line represents the average of 20 simulations. Fitness benefits and costs are additive and reflect a 5% increase or decrease in the chance of survival for an individual, respectively. Costs are associated with the Edit, and the Editor has neutral fitness.

Consequences of Genetic Linkage Between Editor and Edit

Thus far we have considered scenarios in which the editor and edit site are unlinked. In some cases, particularly if multiple changes are desired, some degree of linkage between the editor and one of the edits may be present. To explore the consequences of linkage we consider the extreme scenarios in which the edit and editor are either tightly linked (are always co-inherited) or unlinked (have equal chance of

being co-inherited or not), and the editor has a 50% probability of germline editing and there is no maternal carryover. We utilize a lower rate of cleavage because when editing rates are 100% there is no difference between the linked and unlinked scenarios; all progeny inherit an edit from the carrier parent regardless of linkage. As shown in the heat maps in Fig. 4, for a neutral editor and deleterious edit an absence of linkage is beneficial for spread of the edit (Figure 4). This is because an unlinked editor encounters more non-edited alleles than a linked editor (existing edits spend increased time in the presence of the editor). When fitness costs are associated with the editor, the unlinked version still performs slightly better than the linked (Supplement Figure 4). However, the differences are much smaller and more difficult to see, and beneficial edits spread rapidly regardless of linkage status. In summary, linkage can decrease rates of editing for neutral and deleterious alleles, but the effects are only significant when linkage is tight and editing rates are substantially below 100%.

Figure 4. Effects of Linkage between editor and edit on population modification. The frequency of editing is set to 50% and fitness costs are associated only with the edit. **A, B)** The percentage of the population that has at least one edited allele at generation 50 averaged over 20 simulations. For higher fitness costs, tight linkage between editor & edit **(A)** results in reduced ability to push edits to high frequency as compared with the case in which editor & edit are unlinked **(B)**.

Population Suppression

Transgene-based population suppression strategies take several forms. In one, a self-sustaining gene drive utilizes high frequency homing to spread a multi-gene cassette into (thereby inactivating) a haplosufficient gene required in somatic cells for female sexual identity, fertility, or viability. This drives the population towards a homozygous genotype that is fit in males but unfit in females, leading to a population crash[12,35,36]. There are also suppression strategies that do not rely on drive; Non-drive transgene-based approaches utilize periodic inundation of males to, by one mechanism or another, reduce the frequency of female progeny, of progeny generally, or fertile females[37–41].

Here we explore how an Allele Sail could be used for population suppression by causing sex ratio distortion. Our focus is on species in which expression of a single gene is needed for femaleness, and whose loss results in conversion of these individuals into fertile males. The Tra gene in medfly Ceratitis capitata provides one example (reviewed in [42]). Aromatase, encoded by the cyp19a1a gene, plays a similar

role in a number of vertebrates. Aromatase converts androgens to estrogens and its loss through chemical inhibition or mutation converts genetic females to fertile males[17,43,44]. This, combined with the fact that estrogen agonists can promote femaleness in genetic males[44] argues that aromatase activity is both necessary and sufficient for femaleness. Here we model the composition and fate of a population in which an editor such as Cas9 is introduced, creating LOF alleles (the edit) in aromatase. Because single locus sex specification occurs in species with diverse chromosomal systems, we model suppression in species with XY and ZW sex chromosomes. We also remove maternal carryover from these simulations, to allow the editor to persist in the female line for longer. We first consider the consequences of single releases and then multiple releases.

Dynamics of Alleles and Chromosomes following a Single Release

Results of a single release of the editor at a frequency of 10% carrying capacity are shown in Figure 5A. We compare these to results of a single release of males carrying transgenes that implement female sterile Repressible Inducible Dominant Lethal (fsRIDL), a system in which an autosomal transgene, transmitted in a Mendelian manner, causes death or sterility of female progeny carriers. The suppression effects of fsRIDL are immediate, but the total population also increases back to carrying capacity very rapidly. In contrast, release of an Allele Sail editor results in a relatively prolonged reduction in population size, which is particularly prominent in the ZW system (Figure 5A). To understand why a single release of an Allele Sail with no associated fitness costs only modestly reduced population size rather than collapsing it entirely, we first

examined the XY system and the frequency of the editor, the edit, and the Y chromosome over time. The initial release is of XX males homozygous for the editor and edit. This results in a transient spike in population numbers because when these mate with XX females only female progeny are produced, since the editor is germline specific. This increase in females leads to an increase in population, as their progeny will include both males and females because the edit is heterozygous in these first generation females. Interestingly, the editor undergoes a rapid decrease in frequency, even as the frequency of the edit increases dramatically and the population slowly returns to its carrying capacity (Figure 5B-F). These dynamics (also observed when XY males are released; Figure 5D) can be explained by considering how the frequency of males in the population changes over time. As edits accumulate, males (many now XX) constitute an increasing fraction of the population (Fig. 5A, dashed lines). The editor also spends more time in males than in females as a result of its germline editing activity (Supplement Figure 5). Since each female only mates once, males are in excess and the probability that an editor-bearing male will participate in reproduction is reduced. In consequence, the frequency of the editor declines while the edit stabilizes at an intermediate frequency, thereby preventing further population suppression. The Y chromosome (Fig. 5B) is lost from the population for similar reasons. As edits accumulate, XX males make up a larger fraction of the (increased) male population, resulting in a corresponding decrease in the probability that Y-bearing males participate in reproduction.

An important consequence of a single modest release of an editor in an XY system is that the Y is completely lost from the population even as the ratio of males to females approaches 1:1 (Figure 5B). This result implies that sex is now determined through a different mechanism. The aromatase loss-of-function edit plateaus at a frequency of 75%, indicating that the aromatase locus is now the primary sex determining locus, with males being edit+/edit+ and females edit+/edit-. This behavior is predicted by earlier modeling of switching of the heterogametic sex through intermediate states in which multiple systems are active[45]. It can be understood intuitively by considering the plot in Figure 6, which shows outcomes when populations are seeded with various frequencies of aromatase edits and then followed for 500 generations. Each point represents the simulation endpoint after 500 generations, by which time the sex ratio has approached the Fisherian 1:1[46]. The frequency of the original sex chromosome (either Y for XY, or Z for ZW) is plotted as the X-coordinate, and the frequency of the new sex chromosome, (the edit) is plotted as the Y-coordinate. The trajectory composed of these points represents a path by which sex can transition from male heterogametic to female heterogametic without a change in sex ratio. For a complete sex determination system turnover, the Y allele must only drop from 25% frequency to 0%, and cleaved aromatase must reach 75%. With sufficient editor, this turnover is easily achieved and sex determination becomes solely dependent on the status of the aromatase locus. The consequences of such transitions during attempts (particularly failed attempts) to suppress a population warrant further study.

We now consider a ZW system in which the W chromosome is required but not sufficient for femaleness/female viability (e.g. certain birds[47], crustaceans[48], and amphibians[49,50]) (Figure 5A, C). Release of ZZ males brings about a gradual but transient decrease in population size (Figure 5A, C), coupled with a transient increase in the male:female ratio (Figure 5A). These effects are due to a rise and subsequent fall in the frequency of edits (Figure 5C). The editor is ultimately lost for the same reasons as in the XY system: it is present more often in males than females (Supplement Figure 5), reducing its likelihood of participating in mating. When the WW genotype is inviable, the edits also ultimately decrease in frequency because they find themselves in inviable WW individuals more often than do the non-edited alleles. Placing the editor on the Z-chromosome can mitigate this effect, leading to increased suppression (Supplement Figure 6). In the XY system there is no equivalent lethal genotype since the creation of XX males drives the population towards an all XX genotype in which sex is now determined by the presence or absence of a functional aromatase allele. Also unlike the XY system, the W sex chromosome (in contrast to the Y) is not lost from the population because it is required (but not sufficient) for femaleness. Because of this, and because the Z is required for viability, neither can be lost and the heterogametic sex chromosome flipping cannot occur. In such a system the editor is quickly lost, edits decrease over time, and the W allele returns to a frequency of 25%.

Figure 5. Single Release of Allele Sail for Population Suppression. Values are averaged over 20 stochastic simulations, after a single release of transgenic individuals, with introductions being 10% of the carrying capacity. The counting of population size occurs before this additional release, so those individuals are not counted here. Additionally, there is no maternal carryover occurring in these simulations **A)** An overview of the total population over time, for XY and ZW systems using Allele Sail for suppression, compared to fsRIDL. **B-F)** A breakdown of allele frequency overtime, graphed as a proportion of the total population and also compared to the carrying capacity. **B)** For an XY system, releasing Editor/Editor, Edit/Edit, XX males. The introduction of additional X chromosomes leads to an initial spike in population, further explored in Supplemental Figure 7. **C)** For a ZW system, releasing Editor/Editor, Edit/Edit, ZZ males. The increase in Z alleles only increases the effects of our male-skewing editor. WW individuals are considered non-viable, and die off. **D)** Same as B, but releasing XY males, removing the spike. **E)** The same as C, but WW individuals are viable, leading to an overall increase in W alleles in the population. **F)** Same as E, but releasing ZW males instead of ZZ.

In the case where WW homozygotes are viable[48,50,51] (as might be the case with a newly derived sex chromosome) heterogametic sex flipping can occur, but is harder to achieve than in the XY case. The dynamics that support this conclusion are shown in Figure 5E, F and Figure 6, focusing on the behavior of the W chromosome. Figure 5E and Figure 5F show the consequences of a single release of ZZ or ZW individuals,

respectively. In both cases the transient drop in population size is coupled with a substantial and persistent rise in the frequency of the edits and the W chromosome. Both plateau as the editor is lost from the population through the mechanisms discussed above.

The reason the population returns to its carrying capacity even while edits remain at high frequency is due to the fact that WW individuals are viable and can be either male or female depending on the status of the aromatase gene. This point is illustrated in [Figure 6](#), which shows that there is a large region of parameter space in which ZZ males and ZW females coexist at a near 1:1 sex ratio with some fraction of ZW males and WW females. The presence of edits drives the population towards a WW state (with aromatase status again determining sex), but a much higher frequency of edits is required than with the XY system, since the Z allele must decrease in frequency from 75% to 0% (as opposed to a drop of the Y from 25% to 0%) in order for the transition to be complete. Thus, a modest single release is insufficient and leaves the population in a state with mixed sex determination systems. These results correspond to previous findings that modeled a scenario termed mildly male-determining XY to ZW turnover[45,52].

Figure 6. Paths to sex chromosome turnover. A population of the given system, either XY or ZW, was simulated with a single addition of various amounts of Edit (cleaved aromatase). Each point represents the allele frequencies and percent males of each simulation after 500 generations. The frequency of the old sex chromosome (either Y for XY, or Z for ZW) is plotted as the X-coordinate, and the frequency of the new sex chromosome, (the edit) is plotted as the Y-coordinate. Here, WW individuals are viable, allowing for complete turnover from a ZW system in which edits are absent, to a WW system in which the presence or absence of the edit determines sex. Note that all populations end near a ratio of 1:1 males:females.

Dynamics of Alleles and Chromosomes following Multiple Releases

Here we model the effects of repeated releases - once every generation. As illustrated in Figure 7, repeated releases of an aromatase editor at a frequency of 10% eventually causes population collapse (Figure 7A). This is in contrast to repeated releases of fsRIDL, which at the same introduction frequencies only decrease the total population size (Figure 7A). To explore multiple release scenarios in more depth we investigated the time to collapse. Plots of the average generation to collapse versus introduction frequency are shown in Figure 7B. An editor is much more effective than fsRIDL for low-frequency releases, causing collapse for releases between 10% and 20%, while fsRIDL only results in collapse when the release frequency is 25%, and even then only after roughly an additional 10 generations. For higher introduction frequencies, the time to collapse is comparable (Figure 7B). Reduced editing efficiencies can still lead to population collapse under some scenarios. An XY system can be collapsed faster than fsRIDL (assuming 100% female killing) when the editing efficiency is 90%, but not when the editing efficiency is 80%. Meanwhile, in a ZW system in which WW individuals are non-viable, an editing efficiency of 80% still promotes population collapse faster than with fsRIDL. (Supplement Figure 8).

Finally, we asked what the effects on population suppression were if the editor, in addition to cleaving a gene required for femaleness, also created LOF alleles in a haplosufficient gene required in somatic cells for female viability or fertility - a strategy sometimes referred to as sterilizing sex conversion[53]. As illustrated in Supplement Fig. 9, sterilizing sex conversion leads to faster collapse, bringing about elimination of a

population using repeated releases of only 5% of the starting population ([Supplement Figure 9](#)).

Figure 7. Comparison of population suppression using multiple releases of an Allele Sail or fsRIDL. A) Total population over time, averaged over 20 simulations. Transgenic individuals are being introduced at 10% of carrying capacity, at the start of every generation. The total population count shown here does not include this additional

population, only counting the number of surviving offspring. The XY system shown here introduces XX males, and the ZW introduces ZZ releases when WW offspring are non-viable or viable. **B)** The average generation of collapse for various introduction frequencies. Points plotted here are the average of 20 simulations, where all 20 simulations went to collapse within 50 generations. If all 20 simulations did not collapse within 50 generations, the corresponding point is not plotted. Transgenic individuals were introduced at the indicated introduction frequency, once every generation, until collapse. For introduction frequencies below 25%, an editor can collapse a population where fsRIDL cannot. At higher frequencies, the ability to collapse a population and time to collapse are comparable for strategies using fsRIDL and an editor.

Discussion

We set out to investigate the potential of a novel population-level genome modifier, which we call an Allele Sail. An Allele Sail consists of a genome modifier (the Wind) transmitted in a Mendelian manner, that introduces edits into a target locus (the Sail). A single release of an editor is capable of pushing a target allele to fixation, even in the presence of fitness costs, on either the editor or the edit. These observations suggest an Allele Sail can be a promising tool for population modification. The Allele Sail is a relatively simple system that is applicable for many species. It will also often be self-limiting when the editor or edits have costs. Even in populations where a beneficial allele is already present at low frequencies, an Allele Sail can push that trait into the population at much faster rates than those due to natural selection alone, depending on

how many transgenic individuals are released and the efficiency of the Wind/Editor. One concern may be that simple editors can only make small changes to the genome. However, small variations such as point mutations or small indels at one or a modest number of loci can have a large effect. As examples, point mutations have been found that contribute to plant disease resistance[3], animal heat tolerance[1,2], and honeybee sensitivity to Varroa mites[4,5]. As another point of reference, as of 2019, half of the pathogenic variants in the human ClinVar database were point-mutations, and almost 90% of clinically relevant insertions and deletions were less than 30bp[54]. Finally, recent work shows that larger fragments of DNA (which would albeit be considered transgenes) can be copy-pasted from one location to another using engineered retrotransposons[55,56]. These observations suggest it may soon be possible to push larger fragments of DNA into a population using an editing locus transmitted in a Mendelian manner

Use Case - when beneficial alleles already exist

One exciting application of Allele Sail is evolutionary rescue, the process by which introduction of new alleles into a threatened population allows it to adapt rapidly enough to survive a current or anticipated stress. In species where beneficial alleles exist in some populations but not others, two common methods of introducing favorable alleles into the threatened population are selective breeding and targeted gene flow. Examples include breeding toad-smart quolls[57], and introducing heat-tolerant corals into sensitive populations[58]. However, selective breeding is time and labor intensive, and can reduce genetic variation. Translocation of individuals can also lead to outbreeding

depression, while the rapid spread of the beneficial trait is not guaranteed. Genomic editing utilizing an Allele Sail may prove useful because the introduction frequencies can be low, bringing about an increase in trait frequency without dramatically modifying the genetics of the population at other loci.

Use Case - when beneficial alleles *don't* already exist

In populations in which the desired beneficial allele is not present, and a future threat is clear, an Allele Sail could be used for anticipatory conservation: pushing into a population an allele that is not currently strongly beneficial but that is projected to be so in the future. In these scenarios, simply introducing or re-introducing a beneficial allele into a threatened population may not be enough to prevent population collapse. If the population drops below its stochastic threshold before the allele reaches high frequencies, the presence of a beneficial allele at a low frequency may not save the population from extinction, or allow the population to persist in the face of rapidly changing conditions[59]. An Allele Sail can dramatically increase the spread of desired alleles before populations begin to experience serious pressure, slowing or preventing population decline to critical levels. An example of this challenge is highlighted by research into avian malaria in Hawaii; small releases of disease-resistant individuals into a population would likely face genetic dilution if the disease pressures are not yet severe, which might prevent the creation of a large, robustly disease-resistant population[60]. Increased transmission of resistance via an Allele Sail could combat dilution and greatly increase the likelihood of success.

Use Case - non-rescue

An Allele Sail can also spread alleles that are not beneficial to carriers, but are beneficial to others. For example, in invasive species such as Cane Toads, an Allele Sail could reduce the negative effects of the spreading population; if Cane Toads were engineered to produce less toxin or be completely non-toxic, their invasion would be much more survivable for local fauna. An Allele Sail could also spread an anticipatory maladaptation, seeding an increased susceptibility to certain toxicants before introducing said toxicants into a population, thereby increasing its effectiveness.

Suppression

In the context of population suppression we focused on using repeated introductions of an Allele Sail to bring about sex ratio distortion. An Allele Sail is (when modification rates are high) generally more effective than the non-drive fsRIDL introduced under similar conditions. One important finding is that Allele Sails are self-limiting, even when their presence does not result in a fitness cost, as they remove themselves from the population over time. Even if small numbers of Allele Sail individuals spread through migration into a secondary population, they would not seriously impact total population size. A chemical sex-distortion scheme, with some similarities to the one discussed here, is being tested for Brook trout in Idaho; YY males are created following supplementation with estrogens that create XY females. YY males are bred and released, leading to all XY-male progeny[61]; modeling suggests that regular releases, when combined with yearly 50% suppression via conventional means, have an 80% chance of eradicating brook trout within 9 years[62]. Using an Allele Sail could

drastically reduce the number of fish that would need to be released, and decrease the eradication time. Finally, if the editor creates a second set of edits that induce female sterility, the power of the system to bring about population elimination at low introduction frequencies is enhanced. An added benefit of this strategy is that sterilizing sex conversion likely has a higher tolerance to evolution of resistance than non-sterilizing sex conversion[53].

Altering Sex Determination

In our investigation of population suppression we encountered an interesting result concerning sex determination. By introducing a modest frequency of an Allele Sail that cleaves aromatase and creates fertile XX males in an XX female; XY male species, we were able to switch a population from having heterogametic males to heterogametic females, creating a new sex determination system. The sail we introduced, cleaved aromatase, is considered a mildly male determining mutant, matching up with Table 2B in Bull & Charnov[45]. The sex determination system turnover happens rapidly in our simulations, as the Y allele is nearly eliminated by generation 50. Introduction of the same frequency of editors into a ZW system did not yield a switch, but could (when WW is viable) do so under multiple release scenarios in which population elimination failed. The ZW to Edit/No Edit seems to be a much slower process, which is supported by the neutral equilibrium path length of these two systems.

Summary/Conclusions

The differences in behavior of different sex determination systems means that special consideration should be taken for different species. A system where WW

individuals are non-viable could be highly efficient, which could be the case in many bird species due to degeneration of the W chromosome[47,63]. Birds are a common agricultural pest, and many attempts have been made to reduce or eliminate invasive birds. Examples include Monk Parakeets and Sacred Ibis in the US[64], and the 20 exotic species of birds currently in Australia which are known to be agricultural pests[65]. In contrast, there are many species for whom WW individuals are viable, including crustaceans[48], amphibians[49,50], and reptiles[51]. When designing for these systems, sterilizing sex conversion may be necessary to achieve the desired outcomes. In cases where an XY system is involved, sex determination system turnover is a legitimate concern and the impacts of turnover would need to be seriously considered.

Despite these limitations, Allele Sails could be useful for suppression in all these cases, and across many species. Unlike homing-based drives, the editors in Allele Sail systems would not need to rely on homology directed repair mechanisms, and therefore have a much broader range of use. When used for suppression, Allele sails are also non-lethal, addressing concerns of humaneness[66]. An additional benefit of Allele Sails is that, in theory, the number of organisms bearing Cas9 or some other editor should never significantly increase in the population. In countries like Australia, offspring that only inherit an edit, but not an editor transgene, could be considered non-GMO, reducing regulatory hurdles[15]. Because of its broad applicability, both in terms of Suppression and Modification, we believe Allele Sails are a potentially useful tool for population control, both in reducing and eliminating invasive or harmful species, as well as saving or conserving endangered native species.

Methods

All modeling was done using a stochastic discrete-generation simulation, for populations with carrying capacity of 10,000 individuals, and individuals that are expected to bear 100 offspring each. The actual number of offspring is pulled from a poisson distribution.

Introduction frequencies represent a release of homozygous males, equal to that percent of the carrying capacity; they are an additional release of individuals on top of the carrying capacity. For example, a 10% introduction frequency of transgenic individuals will have a starting breeding population of 10,000 wildtype individuals, and 1000 transgenic individuals for a total population of 11,000. However, our measure of Total Population, as shown in the figures, does not consider the individuals artificially added to the population and does not count the added individuals when performing population-dependent growth.

Density-dependent growth was implemented as a survival modifier on all offspring; when the population size is equal to carrying capacity, each individual in a litter of size N has a 2/N chance of survival. This should lead to perfect population replacement when there are no fitness costs, as each mating produces 2 offspring. This value is modified by a density-dependence factor (eq. 1), where P is the current population size, K is the carrying capacity, and g is some arbitrary growth factor. We used $g = 10$, which means that if population size is near zero, each offspring has roughly a 20/N chance of surviving. Use of this equations leads to logistic growth.

$$Density\ dependence\ =\ \frac{g}{1+(g-1)*\frac{P}{K}},\ (1)$$

Fitness costs further modify the 2/N chance of survival, such that a 5% fitness cost leads to an individual having a 2/N * 0.95 chance of survival, and a 5% fitness benefit leads to an individual having a 2/N * 1.05 chance of survival.

Some simulations included maternal carryover modification and some did not - generally, all modification simulations used maternal carryover of the editor, while suppression simulations did not. These differences are noted in the text and figure legends. Maternal carryover was assumed to be 100%, meaning that if the mother carried an editor, both target alleles in the offspring would be edited regardless of whether the father carried an editor.

The Simulation can be found at https://github.com/HayLab/AlleleSail.

Data Availability

All data is available in the main text, the supplementary information files, or in the github: https://github.com/HayLab/AlleleSail

References

1. Tan, Y., Ma, C.-Y., Li, X.-X., Han, G.-D. & Dong, Y.-W. Genome-wide sequencing identifies a thermal-tolerance related synonymous mutation in the mussel, Mytilisepta virgata. *Commun Biol* **6**, 1–11 (2023).

2. Sosa, F. *et al.* Effects of the SLICK1 mutation in *PRLR* on regulation of core body temperature and global gene expression in liver in cattle. *animal* **16**, 100523 (2022).

3. Li, S. *et al.* Genome-edited powdery mildew resistance in wheat without growth penalties. *Nature* **602**, 455–460 (2022).

4. Sainsbury, J. *et al.* Marker assisted selection for Varroa destructor resistance in New Zealand honey bees. *PLOS ONE* **17**, e0273289 (2022).

5. Tsuruda, J. M., Harris, J. W., Bourgeois, L., Danka, R. G. & Hunt, G. J. High-Resolution Linkage Analyses to Identify Genes That Influence Varroa Sensitive Hygiene Behavior in Honey Bees. *PLOS ONE* **7**, e48276 (2012).

6. Selmoni, O., Bay, L. K., Exposito-Alonso, M. & Cleves, P. A. Finding genes and pathways that underlie coral adaptation. *Trends in Genetics* **40**, 213–227 (2024).

7. Researchers embrace a radical idea: engineering coral to cope with climate change. https://www.science.org/content/article/researchers-embrace-radical-idea-engineering-coral-cope-climate-change.

8. Foster, J. T. *et al.* Genetic structure and evolved malaria resistance in Hawaiian honeycreepers. *Molecular Ecology* **16**, 4738–4746 (2007).

9. Savage, A. E. & Zamudio, K. R. MHC genotypes associate with resistance to a frog-killing fungus. *Proceedings of the National Academy of Sciences* **108**, 16705–16710 (2011).

10. Franz, A. W. E. *et al.* Engineering RNA interference-based resistance to dengue virus type 2 in genetically modified Aedes aegypti. *Proceedings of the National Academy of Sciences* **103**, 4198–4203 (2006).

11. Ito, J., Ghosh, A., Moreira, L. A., Wimmer, E. A. & Jacobs-Lorena, M. Transgenic anopheline mosquitoes impaired in transmission of a malaria parasite. *Nature* **417**,

452–455 (2002).

12. Kyrou, K. *et al.* A CRISPR–Cas9 gene drive targeting doublesex causes complete population suppression in caged Anopheles gambiae mosquitoes. *Nat Biotechnol* **36**, 1062–1066 (2018).

13. Bier, E. Gene drives gaining speed. *Nat Rev Genet* **23**, 5–22 (2022).

14. Melesse Vergara, M., Labbé, J. & Tannous, J. Reflection on the Challenges, Accomplishments, and New Frontiers of Gene Drives. *Biodes Res* **2022**, 9853416.

15. Regulator, O. of the G. T. Overview of the status of organisms modified using gene editing and other new technologies. *The OGTR* https://www.ogtr.gov.au/resources/publications/overview-status-organisms-modified-using-gene-editing-and-other-new-technologies (2021).

16. Pane, A., Salvemini, M., Delli Bovi, P., Polito, C. & Saccone, G. The transformer gene in Ceratitis capitata provides a genetic basis for selecting and remembering the sexual fate. *Development* **129**, 3715–3725 (2002).

17. Lau, E. S.-W., Zhang, Z., Qin, M. & Ge, W. Knockout of Zebrafish Ovarian Aromatase Gene (cyp19a1a) by TALEN and CRISPR/Cas9 Leads to All-male Offspring Due to Failed Ovarian Differentiation. *Sci Rep* **6**, 37357 (2016).

18. Langmüller, A. M. *et al.* Fitness effects of CRISPR endonucleases in Drosophila melanogaster populations. *eLife* **11**, e71809.

19. Friskes, A. *et al.* Double-strand break toxicity is chromatin context independent. *Nucleic Acids Research* **50**, 9930–9947 (2022).

20. Doll, R. M., Boutros, M. & Port, F. A temperature-tolerant CRISPR base editor

mediates highly efficient and precise gene editing in *Drosophila*. *Sci. Adv.* **9**, eadj1568 (2023).

21. Bestas, B. *et al.* A Type II-B Cas9 nuclease with minimized off-targets and reduced chromosomal translocations in vivo. *Nat Commun* **14**, 5474 (2023).

22. Weber, Y. *et al.* Enhancing prime editor activity by directed protein evolution in yeast. *Nat Commun* **15**, 2092 (2024).

23. Skeens, E. *et al.* High-fidelity, hyper-accurate, and evolved mutants rewire atomic-level communication in CRISPR-Cas9. *Sci. Adv.* **10**, eadl1045 (2024).

24. Oberhofer, G., Johnson, M. L., Ivy, T., Antoshechkin, I. & Hay, B. A. Cleave and Rescue gamete killers create conditions for gene drive in plants. *bioRxiv* 2023.10.13.562303 (2024) doi:10.1101/2023.10.13.562303.

25. Oberhofer, G., Ivy, T. & Hay, B. A. Gene drive that results in addiction to a temperature-sensitive version of an essential gene triggers population collapse in Drosophila. *Proc Natl Acad Sci U S A* **118**, e2107413118 (2021).

26. Oberhofer, G., Ivy, T. & Hay, B. A. Split versions of Cleave and Rescue selfish genetic elements for measured self limiting gene drive. *PLoS Genet* **17**, e1009385 (2021).

27. Oberhofer, G., Ivy, T. & Hay, B. A. Gene drive and resilience through renewal with next generation Cleave and Rescue selfish genetic elements. *Proc Natl Acad Sci U S A* **117**, 9013–9021 (2020).

28. Oberhofer, G., Ivy, T. & Hay, B. A. Cleave and Rescue, a novel selfish genetic element and general strategy for gene drive. *Proc Natl Acad Sci U S A* **116**,

6250–6259 (2019).

29. Behavior of homing endonuclease gene drives targeting genes required for viability

    or female fertility with multiplexed guide RNAs.

    https://www.pnas.org/doi/10.1073/pnas.1805278115 doi:10.1073/pnas.1805278115.

30. Metzloff, M. *et al.* Experimental demonstration of tethered gene drive systems for

    confined population modification or suppression. *BMC Biol* **20**, 119 (2022).

31. Xie, K., Minkenberg, B. & Yang, Y. Boosting CRISPR/Cas9 multiplex editing

    capability with the endogenous tRNA-processing system. *Proc Natl Acad Sci U S A*

    112, 3570–3575 (2015).

32. Bosch, J. A., Birchak, G. & Perrimon, N. Precise genome engineering in Drosophila

    using prime editing. *Proceedings of the National Academy of Sciences* **118**,

    e2021996118 (2021).

33. Moreno-Mateos, M. A. *et al.* CRISPR-Cpf1 mediates efficient homology-directed

    repair and temperature-controlled genome editing. *Nat Commun* **8**, 2024 (2017).

34. Xiang, G., Zhang, X., An, C., Cheng, C. & Wang, H. Temperature effect on

    CRISPR-Cas9 mediated genome editing. *Journal of Genetics and Genomics* **44**,

    199–205 (2017).

35. Hammond, A. M. *et al.* The creation and selection of mutations resistant to a gene

    drive over multiple generations in the malaria mosquito. *PLoS Genet* **13**, e1007039

    (2017).

36. Hammond, A. *et al.* A CRISPR-Cas9 gene drive system targeting female

    reproduction in the malaria mosquito vector Anopheles gambiae. *Nat Biotechnol* **34**,

78–83 (2016).

37. Kandul, N. P. *et al.* Transforming insect population control with precision guided sterile males with demonstration in flies. *Nat Commun* **10**, 84 (2019).

38. Smidler, A. L. *et al.* A confinable female-lethal population suppression system in the malaria vector, *Anopheles gambiae*. *Sci. Adv.* **9**, eade8903 (2023).

39. Dobson, S. L. When More is Less: Mosquito Population Suppression Using Sterile, Incompatible and Genetically Modified Male Mosquitoes. *Journal of Medical Entomology* **58**, 1980–1986 (2021).

40. Burt, A. Gene drive designs for efficient and localisable population suppression using Y-linked editors.

41. Upadhyay, A. *et al.* Genetically engineered insects with sex-selection and genetic incompatibility enable population suppression. *eLife* **11**, e71230 (2022).

42. Makki, R. & Meller, V. H. A factor to control Medfly sex. *Science* **365**, 1380–1381 (2019).

43. Olmstead, A. W. *et al.* Sex reversal of the amphibian, Xenopus tropicalis, following larval exposure to an aromatase inhibitor. *Aquat Toxicol* **91**, 143–150 (2009).

44. Guiguen, Y., Fostier, A., Piferrer, F. & Chang, C.-F. Ovarian aromatase and estrogens: A pivotal role for gonadal sex differentiation and sex change in fish. *General and Comparative Endocrinology* **165**, 352–366 (2010).

45. Bull, J. J. & Charnov, E. L. Changes in the heterogametic mechanism of sex determination. *Heredity* **39**, 1–14 (1977).

46. Sexual Reproduction And Sexual Selection. in *Natural selection: A Complete*

Variorum Edition (eds. Fisher, R. A. & Fisher, R. A.) 0 (Oxford University Press, 1999). doi:10.1093/oso/9780198504405.003.0006.

47. Kuroiwa, A. Sex-Determining Mechanism in Avians. in *Avian Reproduction: From Behavior to Molecules* (ed. Sasanami, T.) 19–31 (Springer, Singapore, 2017). doi:10.1007/978-981-10-3975-1_2.

48. Molcho, J. *et al.* Three generations of prawns without the Z chromosome: Viable WW Macrobrachium rosenbergii all-female populations in polyculture with Oreochromis niloticus. *Aquaculture* **515**, 734531 (2020).

49. Mikamo, K. & Witschi, E. Masculinization and breeding of the WW Xenopus. *Experientia* **20**, 622–623 (1964).

50. Colombelli, B., Thiébaud, Ch. H. & Müller, W. P. Production of WW super females by diploid gynogenesis in Xenopus laevis. *Molec. Gen. Genet.* **194**, 57–59 (1984).

51. Booth, W., Johnson, D. H., Moore, S., Schal, C. & Vargo, E. L. Evidence for viable, non-clonal but fatherless Boa constrictors. *Biology Letters* **7**, 253–256 (2010).

52. Saunders, P. A., Neuenschwander, S. & Perrin, N. Sex chromosome turnovers and genetic drift: a simulation study. *J Evol Biol* **31**, 1413–1419 (2018).

53. Meccariello, A. *et al. Gene Drive and Genetic Sex Conversion in the Global Agricultural Pest Ceratitis Capitata - PREPRINT.* http://biorxiv.org/lookup/doi/10.1101/2023.08.16.553191 (2023) doi:10.1101/2023.08.16.553191.

54. Anzalone, A. V. *et al.* Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature* **576**, 149–157 (2019).

55. Zhang, X. *et al.* Harnessing eukaryotic retroelement proteins for transgene insertion into human safe-harbor loci. *Nat Biotechnol* (2024) doi:10.1038/s41587-024-02137-y.

56. Wang, Y. *et al.* CRISPR-Enabled Autonomous Transposable Element (CREATE) for RNA-based gene editing and delivery. 2024.01.29.577809 Preprint at https://doi.org/10.1101/2024.01.29.577809 (2024).

57. Kelly, E. & Phillips, B. L. Targeted gene flow and rapid adaptation in an endangered marsupial. *Conservation Biology* **33**, 112–121 (2019).

58. Anthony, K. *et al.* New interventions are needed to save coral reefs. *Nat Ecol Evol* **1**, 1420–1422 (2017).

59. Carlson, S. M., Cunningham, C. J. & Westley, P. A. H. Evolutionary rescue in a changing world. *Trends in Ecology & Evolution* **29**, 521–530 (2014).

60. Samuel, M. D., Liao, W., Atkinson, C. T. & LaPointe, D. A. Facilitated adaptation for conservation – Can gene editing save Hawaii's endangered birds from climate driven avian malaria? *Biological Conservation* **241**, 108390 (2020).

61. Kennedy, P. A., Meyer, K. A., Schill, D. J., Campbell, M. R. & Vu, N. V. Survival and Reproductive Success of Hatchery YY Male Brook Trout Stocked in Idaho Streams. *Transactions of the American Fisheries Society* **147**, 419–430 (2018).

62. Poirier, J., Davis, B. & Harris, J. Feasibility Assessment of Stocking YY Males to Eradicate Nonnative Brook Trout from Tyee Springs.

63. Zhou, Q. *et al.* Complex evolutionary trajectories of sex chromosomes across bird taxa. *Science* **346**, 1246338 (2014).

64. Avery, M. L. & Feare, C. J. Control or eradication: problems in the management of invasive birds. in *Invasive birds: global trends and impacts* (eds. Downs, C. T. & Hart, L. A.) 350–362 (CABI, UK, 2020). doi:10.1079/9781789242065.0350.

65. Tracey, J., Bomford, M., Hart, Q., Saunders, G. & Sinclair, R. Managing Bird Damage to Fruit and Other Horticultural Crops.

66. Sharp, T. M., Cope, H. & Saunders, G. New South Wales Code of Practice and Standard Operating Procedures for the Effective and Humane Management of Pest Birds. *NSW Department of Primary Industries, Orange, NSW* (2022).

Acknowledgements

Special thanks to Tobin Ivy for laying the groundwork for this simulation

Funding

This work was supported by a grant to MM from the Centre for Invasive Species Solutions (P01-B-005) Australian Department of Agriculture, Fisheries, and Forestry (4-FY94ZNX) and Grants to BAH from the Caltech Center for Evolutionary Sciences.

Contributions

Conceptualization: M.L.J, B.A.H. and M.M. Experimental design: M.L.J, B.A.H. and M.M. Coding: M.L.J. Writing: M.L.J, B.A.H. and M.M.

Corresponding authors

Correspondence to haybruce@caltech.edu and maciej.maselko@mq.edu.au

Ethics declaration

The authors declare no conflicts of interest.

Extended Data

Supplemental Figure 1. Allele Sail in response to different types of fitness costs: dominant, additive or recessive. Fitness costs and benefits are associated with the edit. For the dominant and recessive types, fitness costs/benefits are 10% (either positive or negative), and in the additive case are '5%', such that a heterozygous

individual has a cost/benefit of 5%, and a homozygous individual has a cost/benefit of 10%. **A)** The Average frequency of our edit, both with and without an editor, when transgenic (bearing edit, or edit and editor) individuals are introduced at 10% of carrying capacity and the edit has a 10% (dominant/recessive) fitness benefit. **B)** The same as A, but for a 10% fitness cost. **C)** A heatmap showing the relationship between introduction frequency and fitness costs, when the cost/benefits are dominant and apply to the edit. The percent carriers being plotted here is taken from Generation 50. **D)** The same as C, but for recessive cost/benefits.

Supplemental Figure 2. Behavior of a Non-neutral editor & a Neutral edit. A)
The average allele frequency of edits after being introduced at 10% frequency.
Introduced individuals are homozygous for both edit and editor. The non-neutral editor
includes an additive 5% fitness cost. **B)** Average percent of the population that is
homozygous for our edit after 50 generations, for various editor introduction frequencies
and fitness costs associated with the Sail. **C)** Average percent of the population that
carries at least one copy of the edit, for various editor introduction frequencies and
fitness costs.

Supplemental Figure 3. Increasing Introduction Frequency can Overcome Fitness Costs when Editor Efficiency is low. The Average Edited Allele Frequency over time when each copy of the edited allele confers a 5% fitness cost. Releases consist of all males homozygous for both the editor and the edit.

Supplemental Figure 4. Effects of Linkage on Population Modification, with Cost on the Editor. Rates of editing are 50%. No costs are associated with the edit. **A, B)** The percentage of the population that has at least one edited allele / sail present in their genome, averaged over 20 simulations. For moderate costs linkage between editor and edits results in lower rates of editing **(A)** than does independent assortment. **(B)**.

Supplemental Figure 5. Amounts of Edit and Editor by Sex. A) Average total population graphed with the average number of edit-bearing, editor-bearing, and Y-bearing individuals. The editor is introduced in males, ends up in only females for the first generation, and then drops quickly in females. The low level of editor in females is because editor offspring are skewed male. In the males, the editor is present at a higher frequency in males versus females and is also lost over time. The editor is removed due to dilution: as sex distortion increases, the editor is segregated into males and fewer editor males are chosen to participate in mating with the Non-editor females. **B)** The same as **A**, for a ZW system where WW is nonviable. As with the XY system in **A**, the editor has a higher frequency in males than females and decreases over time.

Supplemental Figure 6: Suppression Behavior of a Z-linked Editor.

A) The average population size over time, following a single release of transgenic individuals at 10% of the carrying capacity. Since the released individuals are males, and population numbers are determined by the number of fertile females, these additional males are not counted at generation 1. **B)** The average population size and the number of Editor alleles over time. Individuals are released every generation at either 5% or 10% of the carrying capacity. Those released individuals do not count towards the number of alleles in the population, but all of their offspring do.

Supplemental Figure 7. Amount of Males and Females under Population Suppression. A) XX males carrying both Editor and Edit are released into the population at a frequency of 20%, at Generation 0. Notice that after this release, the amount of females initially spikes due to increased X frequency. In the second generation, the amount of males and females are equal, and higher than carrying capacity due to an increased number of females. After this, the frequency of the edit increases in the population and the sex ratio becomes skewed towards male. **B)** Under the same conditions as A, XY males are released. Notice that there is no chromosomal imbalance between X's and Y's, leading to almost immediate sex skew and no spike in population.

Average generation of collapse, for each type and introduction frequency

Supplemental Figure 8: Average time to Population Collapse, for various editor efficiencies. Each dot represents the last non-zero generation of a simulation using the given editor efficiency. As such, simulations that ran to generation 50 but did not collapse are also shown here, but are represented by triangles instead of circles. Twenty simulations were run for each scenario, i.e., for each editor efficiency and system. In the fsRIDL case, changing the editor efficiency did not affect the simulation, as there is no editor present. Even so, there are 80 points for each introduction frequency.

Average generation of collapse, for each type and introduction frequency

Supplemental Figure 9. Average time to Population Collapse, including addition of an edit that sterilizes females when homozygous. The same as Figure 6B from the main text, but for different sex determination systems and varying modifications. Points plotted here are the average of 20 simulations, where all 20 simulations went to collapse within 50 generations. If all 20 simulations did not go to fixation within 50 generations, the corresponding point is not plotted here. Transgenic individuals were introduced at the corresponding introduction frequency, once every generation, until collapse. Cleaving a fertility gene, as well as a gene required for female development, significantly increases the efficiency of an Allele Sail as a suppressive mechanism.