1 **Disruption of Circadian Clock Induces Abnormal Mammary Morphology and Aggressive**
2 **Basal Tumorigenesis by Enhancing LILRB4 Signaling**
3
4 **Short title:** Circadian disruption causes tumorigenesis via LILRB4
5

6 **Authors**
7 Olajumoke Ogunlusi,[1][†] Mrinmoy Sarkar,[1][†] Arhit Chakrabarti,[2] Devon J Boland,[3] Tristan
8 Nguyen,[1] James Sampson[4], Christian Nguyen,[1] Danielle Fails,[5] Yava Jones-Hall,[6] Loning
9 Fu,[7] Bani Mallick,[2] Alex Keene,[1] Jeff Jones,[1] Tapasree Roy Sarkar[*1]
10

Affiliations
[1]Department of Biology, Texas A&M University, College Station, TX, USA.

[2]Department of Statistics, Texas A&M University, College Station, TX, USA.

[3]Texas A&M Institute of Genome Sciences & Society (TIGSS), College Station, TX, USA.

[4] Department of Biomedical Engineering, Texas A&M University, College Station, TX,

USA.

[5]Fortis Life Sciences, Montgomery, TX 77356, USA.

[6]Veterinary Pathobiology, Texas A&M University, College Station, TX, USA.

[7]Baylor College of Medicine, Houston, TX, USA.

20
21 [*]**Corresponding author:** tsarkar@bio.tamu.edu.
22 [†] These authors contributed equally to this work.

Abstract

Epidemiological studies have shown that circadian rhythm disruption (CRD) is associated with the risk of breast cancer. However, the role of CRD in mammary gland morphology and aggressive basal mammary tumorigenesis and the molecular mechanisms underlying CRD and cancer risk remain unknown. To investigate the effect of CRD on aggressive tumorigenesis, a genetically engineered mouse model that recapitulates the human basal type of breast cancer was used for this study. The effect of CRD on mammary gland morphology was investigated using wild-type mice model. The impact of CRD on the tumor microenvironment was investigated using the tumors from LD12:12 and CRD mice via scRNA seq. ScRNA seq was substantiated by multiplexing immunostaining, flow cytometry, and realtime PCR. The effect of LILRB4 immunotherapy on CRD-induced tumorigenesis was also investigated. Here we identified the impact of CRD on basal tumorigenesis and mammary gland morphology and identified the role of LILRB4 on CRD-induced lung metastasis. We found that chronic CRD disrupted mouse mammary gland morphology and increased tumor burden, and lung metastasis and induced an immunosuppressive tumor microenvironment by enhancing LILRB4a expression. Moreover, CRD increased the M2-macrophage and regulatory T-cell populations but decreased the M1-macrophage, and dendritic cell populations. Furthermore, targeted immunotherapy against LILRB4 reduced CRD-induced immunosuppressive microenvironment and lung metastasis. These findings identify and implicate LILRB4a as a link between CRD and aggressive mammary tumorigenesis. This study also establishes the potential role of the targeted LILRB4a immunotherapy as an inhibitor of CRD-induced lung metastasis.

Keywords

Circadian rhythm disruption, breast cancer, LILRB4, tumor microenvironment, immunotherapy.

Introduction

The circadian clock regulates the expression of several genes that impact physiology, metabolic

processes, and health outcomes (Panda, Antoch et al. 2002, Storch, Lipan et al. 2002, Kelleher,

Rao et al. 2014). Epidemiological studies have reported that circadian rhythm disruption (CRD),

such as that which occurs during shift work or travel across time zones, affects human health and

increases the risk of developing cancer, metabolic disorders, and cardiovascular disease (Hemmer,

Mareschal et al. 2021). According to the National Health Interview Survey, 12–35% of the US

population works irregular schedules, including night and rotating shifts (CDC 2015). CRD

increases the risk of different types of cancer, including lung (Papagiannakopoulos, Bauer et al.

2016, Pariollaud, Ibrahim et al. 2022), colon (Chun, Fortin et al. 2022), and breast (Van Dycke,

Rodenburg et al. 2015, Hadadi 2020) cancers.

Shiftwork was found to increase the incidence of breast cancer in nurses by approximately 50%,

suggesting a critical role for the circadian clock in breast cancer pathogenesis (Schernhammer,

Laden et al. 2001). Circadian clocks regulate the rhythmic expression of numerous genes in breast

tissues (Blakeman, Williams et al. 2016). Triple-negative breast cancer (TNBC) or basal phenotype

(Alluri and Newman 2014) encompasses a breast tumor subtype that is clinically negative for the

expression of the estrogen (ER) and progesterone (PR) receptors and lacks overexpression of the

Human Epidermal Growth Factor Receptor 2 (HER2) protein, with unique prognostic and

therapeutic implications (Anders and Carey 2008). TNBC is responsible for more than 15–20% of

all breast cancers, with a mortality rate of 40% (Almansour 2022). TNBC is relatively aggressive

and shows more resistance to chemotherapy than other subtypes of breast cancer (Almansour

2022). However, the effect of CRD on aggressive TNBC is poorly understood, and the precise

mechanism underlying CRD-induced tumorigenesis is not yet known.

The mammalian circadian machinery consists of an autoregulatory transcription-translation

feedback loop, where the "positive elements" circadian locomotor output cycle kaput (CLOCK)

and brain and muscle aryl hydrocarbon receptor nuclear translocator (ARNT)-like protein 1

(BMAL1) heterodimerize through their PAS domains and activate the transcription of the

"negative elements," i.e., the period genes (*Per1* and *Per2)* and the cryptochrome genes (*Cry1* and

Cry2). The PER/CRY heterodimers inhibit the transcription of their gene(s) by blocking

CLOCK/BMAL1-dependent transactivation (Kelleher, Rao et al. 2014). This feedback loop

regulates the transcription of an estimated 30% of the genome (Vitaterna, Shimomura et al. 2019).

The expression of core circadian clock genes is frequently dysregulated in human tumors,

indicating the tumor-suppressive role of the molecular clock (Battaglin, Chan et al. 2021,

Stephenson, Usselmann et al. 2021, Chun, Fortin et al. 2022). Epidemiological studies have also

emphasized that the risk of developing cancer increases with increasing years of shift work (Yuan,

Zhu et al. 2018), indicating that longitudinal experiments are better suited for characterizing how

CRD may affect tumorigenesis.

Here, we investigated how the loss of circadian function impacts mammary glands, tumorigenesis,

and the tumor microenvironment (TME). We used the jet lag schedule, which mimics the circadian

disturbance humans encounter during rotating shift work or frequent jet lag, and the genetically

engineered mouse model (GEMM) of TNBC. We specifically examined the role of leukocyte

immunoglobulin-like receptor 4a (LILRB4a or LILRB4 or gp49B), which is known to suppress

immunity in acute myeloid leukemia (AML) and solid tumors (Deng, Gui et al. 2018, Sharma

2021). Additionally, we investigated whether the elevation of LILRB4 expression could be a key

molecular event in CRD-induced mammary tumorigenesis and immunosuppressive TME. Overall,

this study shows how circadian desynchronization affects mammary gland morphology, aggressive

mammary tumorigenesis, and identifies a compelling target for immunotherapy of CRD-induced

tumorigenesis.

Methods

A. Animal husbandry:

The mice were housed under 12 h light and 12 h dark (LD 12:12) and circadian rhythm

disruption (CRD) conditions at room temperature (25°C), and food and water were

provided *ad libitum*. The genetically engineered mice model (GEMM), i.e., FVB-Tg(C3-

1-Tag)cJeg/JegJ mice (Jax stock #013591), were caged until they were 27 weeks old.

Normal FVB mice (Jax stock #001800) and BALB/cJ mice (Jax stock #000651) were also

used in this study. All animal care and treatments were per the Institutional Animal Care

and Use Committee of Texas A&M University under protocol # 2022-0094.

Mice were grouped randomly for exposure to standard light conditions (LD 12:12) or CRD

light conditions (consisting of an 8-hour light phase advance repeated every two days)

(Casiraghi, Oda et al. 2012, Papagiannakopoulos, Bauer et al. 2016, Pariollaud, Ibrahim et

al. 2022). Zeitgeber time (ZT) 0 corresponded to the onset of light, while ZT12

corresponded to the onset of dark. The LD 12:12 and CRD animal groups were provided

access to running wheels (Columbus Instruments) for several weeks. ActogramJ (ImageJ)

was used to assess voluntary running activities (Schmid, Helfrich-Förster et al. 2011).

B. 4T1 cell mouse TNBC model:

The 4T1 cells were maintained RPMI-1640 (ATCC, 30-2001) supplemented with in 10%

FBS and 1% penicillin and streptomycin in 5% CO_2. Briefly, 10,000 cells suspended in

sterile PBS were injected subcutaneously into the 4th mammary gland fat pad. The mice

were housed in LD or CRD conditions for 3-4 weeks before they were sacrificed to harvest

the tumors. To investigate the effect of immunotherapy on tumor progression and

metastasis, BALB/cJ mice were housed under LD 12:12 conditions for 4-5 days following

injection of 4T1 cells until palpable tumors developed. They were segregated and assigned

randomly either in LD or CRD conditions. On days 6, 9, and 12 (after the palpable tumors

developed), an anti-LILRB4 antibody (ABclonal, #A7073) (polyclonal, 50μg) was injected

intratumorally. The animals were checked regularly to observe their body weight and tumor

sizes.

C. Histology and tumor burden analysis:

The tumors and the mammary glands were harvested using aseptic techniques. Tumor

volumes were calculated for each breast tumor using caliper measurements after every 3

days. $Tumor\ volume\ = (length\ \times width^2\ )/2$ was used to measure the volume of the

tumors (Kersemans, Cornelissen et al. 2013), whereas the equation

$Tumor\ burden\ (in\ percentage)\ = (Tumor\ weight\ /mice\ body\ weight) \times 100\ \%$

was considered to determine the tumor burden (Hadadi 2020). The tumor, or the lung

samples were fixed in 10% neutral buffered formalin (NBF) for 48 hours before being

transferred to the Histology core facility at Texas A&M University. The tissue sections

were stained with hematoxylin and eosin (H&E) for pathological examinations.

D. Immunohistochemistry:

The formalin-fixed paraffin-embedded (FFPE) mouse mammary gland tissues were deparaffinized by incubating at 65°C for 2 hours, followed by dipping in xylene for 5 minutes each time for two changes. The endogenous peroxidase activity was inhibited by treating the tissue with hydrogen peroxide. Antigen retrieval was carried out by incubating the slides in pre-heated citrate antigen retrieval buffer (10 mM citrate, 0.05% Tween 20, pH 6.0) at 95°C for 20-25 minutes in a water bath. After blocking, the tissues were incubated overnight at 4°C with anti-ki67 polyclonal antibody (ABclonal, A16919), dissolved in the kit blocking solution. Following incubation with the secondary antibody availed with the IHC kit, the expression was visualized with DAB peroxidase substrate (Abcam, ab64264). The slides were counterstained with Mayer's Hematoxylin Solution (VitroVivo Biotech, VB-3000-1) before being mounted with Optic Mount I™ xylene mounting medium (Mercedes Scientific) and imaged under the Echo Revolve R4 fluorescent microscope (Echo).

E. Whole mount staining of mouse mammary gland:

The mammary gland tissues were spread on glass slides immediately after harvesting. The mammary glands were then fixed using Carnoy's fixative solution (VitroVivo Biotech, VB-3001-1) for three hours, followed by washing in 70% ethanol for 30 minutes. The tissues were incubated overnight at room temperature (RT) with Carmine Alum Staining Solution (VitroVivo Biotech, VB-3001-2). Stained tissues were dehydrated with gradients of ethanol concentrations and finally with three changes of xylene bath of 10 minutes each. The slides were then mounted with Optic Mount I™ xylene mounting medium (Mercedes Scientific) and observed under the microscope.

F. Single-cell RNA sequencing (scRNA Seq):

After brief washing in chilled sterile PBS, the tumors were kept in DMEM (ATCC, 30-2002), supplemented with 10% FBS (Sigma-Aldrich, F2442) and 1% antibiotic solution (Sigma Aldrich, P4333) on ice until minced and subsequently blended with gentleMACS Dissociator (Miltenyi Biotec, 130-093-235). MACS Miltenyi Tumor Dissociation Kit for mice (Miltenyi Biotech, 130-096-730) was used for further enzymatic digestion according to the manufacturer's protocol. The suspension of cells was then transferred to the Texas A&M Institute for Genome Sciences and Society (TIGSS) for scRNA-seq. Single-cell sequencing libraries were prepared on the Chromium platform (10x Genomics) using the Single Cell 5' v2 Full Kit (PN-1000263). Cell type assignment was performed manually based on unique marker genes identified in each cell cluster from the UMAP reduction. Markers were cross-referenced with published literature specific to mouse mammary gland tissues. Differentially expressed genes were identified using the MAST (Finak, McDavid et al. 2015) R package, and GO pathway enrichment analysis was performed with the clusterProfiler (Wu, Hu et al. 2021) R package. The CellChat R package analyzed cellular communication using default parameters. The most significant pathways were identified using the "rankNet" function and visualized with the "netVisual_diffInteraction" function.

G. *In vitro* **tumorosphere formation assay:**

The single cells (from tumors) were cultured in ultra-low attachment 96-well plates (ThermoFisher Scientific, 174927) in EpiCult™-B (mouse) medium supplemented with human recombinant EGF (hrEGF) (Stem Cell Technologies, 78006), human recombinant bFGF (hrbFGF) (Stem Cell Technologies, 78003), and heparin (Stem Cell Technologies, 07980). 500 cells per well from LD 12:12 and CRD group tumors were added and

incubated with standard cell culture conditions for 5 days in the CO_2 incubator before being

observed under the microscope.

H. **Organoid assay:**

Singlet cells from tumors (LD and CRD) were resuspended in Geltrex™ (Gibco,

ThermoFisher, A1413202) and plated on a 24-well plate. After the gel was solidified to

form the matrix, EpiCult™-B (mouse) medium supplemented with hrEGF, hrbFGF, and

heparin was added to assess the growth of the mouse mammary tumor cells from LD 12:12

and CRD conditions. After 7 days of incubation, the organoids formed were imaged under

the microscope.

I. **Flow cytometry:**

The dissociated tumor cells were counted and resuspended in 1 ml of cell staining buffer

(CSB) (BioLegend, 420201). Zombie NIR™ Fixable Viability Kit (BioLegend, 423105)

was applied to cells on ice for 20 min. After spinning down at 400 x g for 4 minutes, cells

were resuspended in 50 µl of the CSB and incubated with fluorochrome-conjugated

antibodies (Alexa Fluor® 700 anti-mouse CD4, Pacific Blue™ anti-mouse CD8a, Alexa

Fluor® 488 anti-mouse CD86, PE anti-mouse CD163, Alexa Fluor® 594 anti-mouse

CD45, FITC anti-mouse Ly-6G/Ly-6C, PE anti-mouse/human CD11b, Alexa Fluor® 647

anti-mouse I-A/I-E, FITC anti-mouse CD11c) in the dark for 20 minutes, following a

systematic approach via designing compatible multicolor flow cytometry panels. Cells

were fixed and permeabilized using 2% paraformaldehyde and 0.1% Triton X-100 in PBS,

respectively, and stained with anti-FoxP3 antibody (Alexa Fluor® 488 anti-

mouse/rat/human FOXP3) for 20 minutes in the dark at 4°C, in case of tumor infiltrated

lymphocyte-specific panel. Afterward, stained cells were rinsed with PBS and kept in PBS

215	at 4°C in the dark until analyzed using Cytek Aurora spectral flow cytometer at TAMU
216	Flow Cytometry Facility.

J. Quantitative PCR for detecting mRNA levels:

218	Total RNA was extracted from mammary glands or tumors using Quick-RNA™ MiniPrep
219	(Zymo Research, R1054), and the purity was analyzed by DeNovix DS-11 Series nanodrop
220	Spectrophotometer (DeNovix). RNA from each sample was reverse transcribed to cDNA
221	by High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems™, 4368814).
222	Analysis for gene expression by quantitative real-time PCR was performed using
223	SsoAdvanced Universal SYBR Green Supermix (BioRad, 1725271).

K. Human CRD analysis for breast cancer patients:

225	All statistical analyses were performed using the R software package (version 4.3.0.).
226	scRNA-seq data corresponding to the triple-negative breast cancer (TNBC) patients from
227	Wu *et al.*, 2021 (Wu, Al-Eryani et al. 2021) were pre-processed using standard steps
228	outlined in Hao *et al.,* (Hao, Hao et al. 2021). The uniform manifold approximation and
229	projection (UMAP) was performed on the processed data. The UMAP plot with cells
230	clustered according to the inferCNV cell type. The cells which had no infer-CNV call are
231	denoted as "unknown." To investigate the level of circadian rhythm disruption in each cell,
232	we calculated the CRD scores for the 358 circadian-related genes (CRGs) following He *et*
233	*al.,* 2022 (He, Fan et al. 2022) separately for malignant and non-malignant cells. A cutoff
234	of 75% based on quartiles was used as the threshold for the CRD level to segregate the
235	scores as high (denoted as CRD^{hi}) and low (denoted as CRD^{low}). The percentage of cells
236	that show CRD^{hi} scores by cell type, along with the p-value of the corresponding Binomial
237	proportion test, is used for this study. The mean normalized expression of the core

circadian genes by the cell type (malignant and non-malignant) is presented in this study. Bulk RNA-sequencing data from the publicly available TCGA BRCA dataset was used further to explore the expression pattern of the core circadian genes. The patients were segregated according to the analyzed tissue type, i.e., "Tumor" and "Surrounding" tissue. The mean expression of the genes (+/- SEM) over patients by the type of tissue analyzed. The Wilcoxon signed-rank test was used to test for the difference in expression between the tumor and surrounding tissues for each gene. The genes showing a significant difference are labeled in "red", while the insignificant genes are marked in "black".

L. Blood chemistry and cytokine array:

Pre-mortem terminal blood was collected from the FVB-Tg(C3-1-Tag)cJeg/JegJ mice. The blood samples were transferred to the Texas A&M TVMDL to analyze the blood chemistry parameters. The cytokine array was performed using the Proteome Profiler Mouse XL Cytokine Array kit (R&D Systems, ARY028). The blood samples were kept undisturbed for 30 minutes at room temperature before being spun at 2000 x g for 10 minutes at 4°C. The serum was separated for the cytokine array profiling according to the manufacturer's protocol and was scanned using ChemiDoc Imaging System (BioRad, 12003153).

M. mIF Staining of Mouse Tissues: The panels were developed and optimized for use in mouse tissue. The following primary antibodies from Fortis Life Sciences, Cell Signaling Technology, and ABclonal were utilized for both immunohistochemical and immunofluorescence staining: rabbit anti-mouse ARG1 [BLR161J], rabbit anti-human/mouse LILRB4 [A7073], rabbit anti-human/mouse CD163 [BLR087G], rabbit anti-mouse CD8a [BLR173J], and rabbit anti-mouse FOXP3 [D6O8R], and rabbit anti-mouse CD4 [BLR167J]. Multiplex immunofluorescence staining was performed with the

Akoya Opal™ Polaris 7-color IHC kit fluorophores (Akoya Biosciences

[NEL861001KT]). FFPE tissue sections were baked for 30 minutes, deparaffinized in

xylene, and rehydrated by serial passage through graded concentrations of ethanol.

Endogenous peroxidase in tissues was blocked with 0.9% H_2O_2/methanol for 40 min.

Multiple (7) sequential HIER treatments were performed for 20 min each at 92-96°C in

Citrate pH 6.0 (first cycle) or Tris EDTA pH9 (remaining cycles) buffer. After each HIER

cycle, sections were rinsed with DI water and cooled at RT for 20 minutes. Tissue sections

were blocked with 20% normal goat serum for 20 minutes before incubation with primary

antibody for 20 min. Then, sections were rinsed with TBST for 10 min and incubated with

HRP-conjugated secondary (A120-501P) for 20 min, followed by another 10 min rinse in

TBST. Incubation with an Opal fluorophore (Opal480, Opal520, Opal570, Opal620,

Opal690) was then done for 10 min, followed by a 10 min rinse in DI water. Bound primary

and secondary antibodies were then subject to the next HIER treatment (as aforementioned)

for 20 min. After washing in DI water and cooling at RT for 20 min, the process of staining

and antibody removal was repeated for the desired number of targets. Finally, the tissue

specimens were stained with 4′,6-diamidino-2-phenylindole (DAPI) for 10 min and

mounted in VECTASHIELD Vibrance Antifade Mounting Medium (ThermoFisher

Scientific). Akoya Biosciences' PhenoImager HT was used for multispectral imaging at

40× magnification.

N. Statistical analysis:

GraphPad Prism8 was used as the Statistical analysis tool except for the assessment of the

rhythmicity that was carried out using the JTK cycle. The significance was assessed using

either an Unpaired t-test or analysis of variance (ANOVA), with a false positive threshold

of 0.05 considered acceptable ($p<0.05$). Unless otherwise mentioned, the data were

presented as mean ± standard error of mean (SEM).

Results

Chronic CRD results in abnormal mammary gland morphology

The jet-lag paradigm mimics the CRD that humans face during shift work (Lee, Donehower et al.

2010) or travels across time zones. Mice were exposed to an 8 h light advance repeated every 2–3

days (Fig. 1A). Female Friend LeukemiaVirus B (FVB) mice (4 weeks old) were housed under

normal light and dark (LD 12:12), or chronic CRD conditions for 8 weeks before the mammary

glands were collected every 4 h for 24 h. We confirmed that this jet-lag protocol disrupted

circadian activity in our mouse model by assessing the locomotor activity (Fig. 1B). CRD did not

significantly impact the weight of mice (Fig. S1A). Real-time PCR analysis showed that CRD led

to the loss of rhythmicity (*PJTKcycle* < 0.05) for most genes, including *Cry1*, *Cry2*, *Bmal1*, *Clock,*

and *Per1* (Fig. 1C). *Per2* expression remained rhythmic in mice exposed to CRD (Fig. 1C),

suggesting the desynchronization of the core clock components. To understand the effect of CRD

on the mammary glands, we examined whole-mount preparations of inguinal mammary glands

from LD and CRD mice at 12 weeks. Chronic CRD decreased branching (bifurcations) (Fig. 1D,

upper panel, 1E, S1B) and the number of terminal end buds (TEBs) (Fig. 1D, lower panel, 1F,

S1B) in the mammary glands. Compared with the mammary glands of LD mice (control), CRD

mice showed a high frequency of ductal hyperplasia as found by using hematoxylin and eosin

(H&E) (Fig. 1G) and by using Ki67 (Fig. 1H, S1C) staining of the mammary glands.

Chronic CRD accelerates aggressive mammary tumorigenesis and lung metastasis

To investigate the effect of CRD on aggressive basal tumorigenesis, we used preclinical GEMM

of human basal tumors (FVB-Tg(C3-1-Tag)cJeg/JegJ (Maroulakou, Anver et al. 1994, Usary

2017) that provides a model for examining how CRD impacts aggressive mammary tumorigenesis

and metastasis. Female hemizygous mice develop mammary gland adenocarcinomas at

approximately 24 weeks of age (Maroulakou, Anver et al. 1994, Usary 2017). To investigate the

effect of CRD on tumorigenesis, mice were placed under LD or CRD conditions when they were

8 weeks old and maintained under CRD conditions through the end of the study at 27 weeks (Fig.

2A). The growth of primary tumors was measured every 3 days throughout this period. Because

of the highly aggressive tumors in CRD mice, we euthanized the animals at 27 weeks. All mice

were euthanized, and the tumor tissues were processed for other analyses. CRD did not

significantly impact the weight of mice (Fig. S2A). Peripheral blood cell counts showed no

significant differences between those mice (Fig. S2B, S2C). No significant changes were noted in

the blood glucose, cholesterol, or amylase (Fig. S2D-F) levels in CRD mice, although the albumin-

to-globulin ratio (AGR) was decreased (Fig. S2G). A high AGR shows a good prognosis for

survival in different solid-tumor patients (He, Pan et al. 2017). Therefore, CRD negatively affects

cancer prognosis by decreasing the AGR. scRNAseq (Fig. S3A) and real-time PCR (Fig. S3B) of

tumor samples showed that CRD did not substantially alter the expression of Per genes (*Per1*, and

Per2) (Fig. S3A, B) or other clock genes (*Clock*, *Cry2*) (Fig. S3A, B). *Cry1* expression

significantly increased in CRD-induced tumors (Fig. S3B).

Animals placed in CRD conditions had a substantially increased mammary tumor burden (Fig. 2B)

compared to the LD animals. CRD also accelerated tumor initiation at 18 weeks. In contrast, the

average time of tumor initiation for LD mice was 22 weeks (Fig. 2C). However, no difference was

observed in the spectrum of tumor grades assessed using histopathology (Elston and Ellis 2002)

between the two groups, with most tumors being grade 2 (Fig. S4A, B). Several technical differences, such as in the mouse genetic background (MMTV:PyMT C57BL/6J versus FVB-Tg(C3-1-TAg)cJeg/JegJ) and conditions of the animal facilities (such as light exposure) may explain why Hadadi *et al.* (Hadadi 2020) observed an effect of CRD on tumor grade, but the same was not found in our study. The number of metastatic foci in the lungs was significantly higher in the CRD mice (Fig. 2D, E). To investigate the impact of CRD on tumor progression, 4T1 (mice triple-negative breast cancer cell line) cells were introduced into the female BALB/cJ mice via orthotopic transplantation into the 4th mammary glands. When mice developed palpable tumors (~4-5 days after injection), they were placed into LD or CRD conditions for 3 weeks until tumors became aggressive. CRD led to a modest but non-significant increase in tumor burden (Fig. S4C). This data suggests that CRD does not affect mammary tumor progression. However, we did observe a significant increase in the formation of lung metastasis foci (Fig 2F, G). Flow cytometry was performed to investigate whether a higher metastatic potential was related to a higher cancer stem cell (CSC) population in CRD tumors. We observed a significant increase in the percentage of CSCs expressing CD49f markers (Fig. 2H, S4D) in CRD tumors compared with that in LD tumors. To investigate whether this enrichment was associated with an increase in the stemness potential of primary tumors, a mammosphere assay was performed using CRD and LD GEMM tumors. CRD tumor cells had a significantly higher (P<0.0001) mammosphere formation efficiency than LD tumor cells (Fig. 2I, S4E). Recent studies have used three-dimensional (3D) organoids as *in vitro* organs, with numerous applications ranging from disease modeling to drug screening (Meyer, Howden et al. 2011). Using small intestinal crypt-derived organoids, a recent study showed that genetic disruption of the circadian clock accelerates intestinal organoid

352 proliferation (Chun, Fortin et al. 2022). We established 3D organoids (tumoroids) from LD and

353 CRD tumors (Fig. 2J). Overall, CRD tumors exhibited more aggressive organoids than LD tumors.

Circadian desynchronization alters the TME

Breast TME is essential for the survival and immunosuppression of tumor cells (Binnewies,

Roberts et al. 2018). We performed scRNA analysis on cells isolated from GEMM basal tumor

tissues. Single cells were isolated from the tumors (Rodriguez de la Fuente, Law et al. 2021)

without surface marker selection. Using proper QC and filtration, we obtained single-cell

transcriptomic data for the studied tumors (N= 2/set). To analyze different cell populations in the

TME, 5,804 QC-positive single cells (LD and CRD tumors) were clustered according to their

expression profiles (Fig. 3A), and 20,764 genes were identified. The Seurat package was used for

preliminary clustering. Cells were clustered based on specific cell markers. Using the established

expression markers, such as for endothelial cells (*KLF2*)(Sangwung, Zhou et al. 2017), M2

macrophages (*SLC2A1*)(Wang, Wang et al. 2023), TNBC cancer cells (*TRPS1*)(Ai, Yao et al.

2021), proliferating cells (*TOP2A)*(Neubauer, Wirtz et al. 2016), myeloid cells (*FTH1*)(Moreira,

Silva et al. 2022), progenitor cells (*NES*)(Bernal and Arranz 2018), adipocytes

(*CLEC11A*)(Merrick, Sakers et al. 2019), mesenchymal cells (*S100A4*)(Fei, Qu et al. 2017),

angiogenic cells (*RPL35A*)(Wu, Al-Eryani et al. 2021), breast cancer stem cells *(FXYD3*)(Li,

Nishimura et al. 2023), dendritic cells or DCs (*CD68*) (Strobl, Scheinecker et al. 1998), cancer-

associated fibroblast or CAFs (*COL1A2*)(Han, Liu et al. 2020), epithelial cells (*EPCAM*)(Trzpis,

McLaughlin et al. 2007), cycling cells (*CDK4*)(Baker, Poulikakos et al. 2022), and B-cells (*CD83*)

(Grosche, Knippertz et al. 2020), we found that CRD altered different cell populations in the TME

(Fig. 3B). This showed that CRD induced a higher number of M2-macrophage, TNBC cancer cells,

cycling cells, adipocytes, CAFs and proliferating cells but a lower number of B-cells, and DCs in

the tumor (Fig. 3B). We next analyzed putative ligand–receptor (L-R) interactions in LD and CRD

tumors and modeled cell–cell interactions between different cell types that play vital roles in

maintaining cellular homeostasis. Cellular communication was investigated using CellChat, which

shows the differences in the total number of L-R interactions between different cell types in the

tumors (Fig. 3C). For example, the interaction between CAF with proliferating cells, TNBC cells,

and M2 macrophages increased under CRD conditions compared to LD conditions (Fig. 3C). We

identified 3429 and 3992 critical ligand–receptor pairings in LD and CRD tumors, respectively

(Fig. S5A). Ligand–receptor interactions with increased and decreased strength were also

identified (Fig. S5B). CellChat also identified signaling pathways linked to ligand–receptor

interactions (Fig. 3D). Specific signaling pathways, including those involving fibroblast growth

factor (FGF), WNT, and platelet-derived growth factor (PDGF), were exclusively detected in CRD

tumors. Oncostatin M, Tumor necrosis factor (TNF), and NOTCH, were detected solely in LD

tumors, and signaling pathways, including transforming growth factor β (TGFβ), epidermal

growth factor (EGF) and fibronectin 1, were detected in both LD and CRD tumors (Fig. 3D). Gene

Ontology (GO) analysis (Fig. 3E) indicated that genes associated with the acetyl CoA metabolic

process, antigen processing (and presentation), nuclear division, regulation of cellular pH,

carboxylic acid catabolic process, chromosome segregation were upregulated in CRD.

CRD induces immunosuppressive TME and turns tumors "cold"

The TME is critical for tumor progression and metastasis. Tumors are "hot" when they show signs

of inflammation characterized by an infiltration of T cells mobilizing to fight the cancerous cells,

whereas nonimmunogenic "cold" tumors lack infiltrating T cells, which makes it challenging to

provoke an immune response with immunotherapy drugs (Ren, Guo et al. 2022). To investigate

whether the increased prevalence of metastasis in CRD mice resulted from changes in the TME,

we characterized different immune cell populations in the TME of LD and CRD tumors using

scRNA-seq, flow cytometry, and immunostaining (MxIF). We determined whether CRD could

influence the proportion of various immune cells in the TME and, in turn, promote tumor

progression and metastasis. scRNA-seq with GEMM tumors showed that CRD reduced cytotoxic

T-cell infiltration (Fig. S6A) and enhanced regulatory T-cell (T_{reg}) infiltration (Fig. S6B). CRD-

induced M2-macrophage populations while decreasing M1-macrophage populations (Fig. S6C).

Flow cytometry analysis with CRD-induced 4T1 tumors supported the scRNA seq data. Our study

showed a significant inhibition of the number of infiltrating $CD8^+$ T cells (cytotoxic T-cells) (Fig.

4A) and an increase in the CD4/CD8 ratio (Fig. 4B) in CRD-induced tumors. The increased

CD4/CD8 ratio in CRD-induced tumors also indicates an immunosuppressive microenvironment.

FACS showed significant enrichment of the immunosuppressive T_{reg} ($CD4^+FoxP3^+$) population

(Fig. 4C) in the CRD-induced tumors. We identified anti-tumor (M1) and pro-tumor (M2)

macrophage populations in the LD and CRD-induced tumors. CRD induced M2-macrophage

infiltration while reducing M1-macrophage infiltration (Fig. 4D). Tumors from mice under CRD

conditions exhibited a ratio (M1:M2) that favored M2-mediated immunosuppression (Fig. 4E) and

facilitated tumor growth. CRD significantly enhanced the $CD45^+$ immune cell population in

tumors (Fig. 4F). CRD-induced tumors showed a lower dendritic cell population compared to LD-

induced tumors which support the anti-cancer properties of dendritic cells (CD11c) (Fig. 4G). Flow

cytometry study also showed a significant increase in the MDSC population (CD11b) (Fig. 4H) in

CRD-induced tumors, which further supports an immunosuppressive microenvironment. A donut

chart shows the differences in different immune cell populations in LD vs CRD-induced tumors

(Fig. 4I). Our immunofluorescence assay showed that CD8+ T-cell infiltration was decreased, but

M2 macrophages (CD163) and FOXP3 (T_{reg}) expression was enhanced in CRD-induced GEMM tumors compared with those in LD GEMM tumors (Fig. 4J), making the tumor "cold." CRD increased EMT gene expressions, such as fibronectin 1 (FN1) and Snail (SNAIL 1) (Fig. 4K). There was an increase in $S100A4^+$ and α-smooth muscle actin ($α-SMA^+$) CAF populations in CRD-induced tumors compared to those in LD tumors (Fig. 4L).

Our data suggest that CRD enhances tumor progression and lung metastasis by enhancing the immunosuppressive microenvironment and increasing the CSC population. Our scRNA-seq, and immunostaining results confirmed the "cold" characteristics of CRD tumors. The gating strategy used for flow cytometry is shown in Fig. S7. Chemokines play a pivotal role in the migration pattern of immune cells into the tumor and facilitate the immunosuppressive TME by enhancing the differentiation and infiltration of immunosuppressive cells, such as T_{reg} cells, myeloid-derived suppressor cells (MDSCs), and tumor-associated macrophages (TAMs). To investigate whether CRD altered the chemokine-cytokine network in the studied mice, we quantified the levels of 40 circulating chemokines in the plasma of LD and CRD mice using a Proteome Profiler. Marked increases in circulating IFN-γ, IL-1β, G-CSF, and IL-16 levels were observed in CRD mice (Fig. S8A). As plasma cytokine/chemokine levels do not represent the TME, we used the data from our scRNA-seq and real-time PCR to assess chemokine/cytokine network expression levels in LD and CRD tumors. Levels of specific cytokines/chemokines known to favor an immunosuppressive microenvironment (Vilgelm and Richmond 2019), such as CCL12 (Fig. 4M) and CXCL5 (Fig. 4N), were upregulated in CRD-induced GEMM tumors. Levels of CCL28, known to induce T_{reg} infiltration and create an immunosuppressive environment (Yang 2017), were upregulated in CRD tumors, as shown using scRNA-seq (Fig. S8B), but not significantly, as shown using real-time

PCR (Fig. S8C). Levels of IL-17β and IL10, were upregulated modestly (but not significantly) in

CRD-induced 4T1 tumors (Fig. S8C).

Malignant human breast cancer cells have high CRD

Earlier studies (Hadadi 2020) and the present study used GEMMs to investigate the effect of CRD

on tumorigenesis. However, more insights are needed into the molecular mechanisms underlying

CRD at the single-cell level and in human tumors. To investigate the process underlying CRD

during mammary tumorigenesis, we used a scRNA sequence dataset from a previously published

study (Wu, Al-Eryani et al. 2021). The dataset included TNBC, HER2-positive, and luminal

tumors. Using large-scale copy number variation (CNV) from the transcriptomics data (Tirosh,

Izar et al. 2016), we separated definitive malignant cells from potential nonmalignant cell

populations in TNBC (Fig. 5A), luminal (ER$^+$/PR$^+$) (Fig. S9A), and HER2 tumors (Fig. S9B).

Based on a previous study (He, Fan et al. 2022), the CRD score was calculated using the expression

of 358 circadian-related genes (He, Fan et al. 2022). Next, we quantified the CRD scores of

malignant (and nonmalignant) cells in different types of human breast tumors. Aggressive TNBC

comprised significantly (P < 0.01) more malignant cells with high CRD scores (Fig. 5B) than

luminal tumors (Fig. S9C). Malignant cells in HER2 tumors also showed a high CRD score (Fig

S9D). *Cry1*, *Per1*, and *Per2* expression decreased in TNBC malignant cells compared to non-

malignant cells, as shown using the scRNA-seq dataset (Fig. 5C). To define the role of CRD in

large patient populations, we used publicly available RNA-seq data from tumors (N=1104 patients)

and surrounding tissues (N = 113) of patients with breast cancer in The Cancer Genome Atlas

(TCGA). The expression of core clock genes (*Per1*, *Per2*, *Per3*, *Cry2*, *Rorc*, *Rorb*, *Nr1d1*, and

Arnt2) was significantly ($P < 0.05$) lower in the tumors than in the surrounding tissues (Fig. 5D).

These data suggest that the core clock is disrupted in human breast tumors.

CRD creates "cold" tumors by inducing LILRB4 expression

Next, to identify the molecular mechanism that regulates the CRD-induced immunosuppressive

TME, we searched for transcripts differentially expressed between the samples (mammary tumors

and glands) collected from mice exposed to ordinary light and dark conditions (LD 12:12) and

mice subjected to CRD. Differential gene expression (DGE) analysis of our scRNA-seq data

revealed that LILRB4 was upregulated in the tumor samples from CRD-induced GEMMs (Fig.

6A, S10A). LILRB4, an immunoreceptor tyrosine-based inhibitory motif-containing receptor,

suppresses T-cell activity via a signaling pathway involving apolipoprotein E (APOE), SH2-

domain-containing protein tyrosine phosphatase-2 (SHP-2), urokinase-like plasminogen activator

(uPAR or Plaur), and arginase 1 (ARG-1) (Sharma 2021). Recent studies have shown that LILRB4

creates an immunosuppressive microenvironment in AML (Deng, Gui et al. 2018) and solid

cancers (Sharma 2021) by inhibiting CD8$^+$ T-cell infiltration and inducing T$_{reg}$. Real-time PCR

showed that CRD enhanced LILRB4 levels significantly in the CRD-induced GEMM tumors (Fig.

6B), and CRD-induced 4T1 tumors (Fig. 6C). The immunofluorescence analysis showed an

elevation of ARG1 expression in CRD-induced GEMM tumors (Fig. 6D). The transcript level of

ARG1 was significantly increased in CRD-induced tumors (Fig. 6E). Increased LILRB4

expression enhanced the immunosuppressive TME as CD8+ T cell infiltration decreased, whereas

M2- macrophage (CD163) and T$_{reg}$ (FOXP3) populations increased in CRD-induced mammary

tumors from GEMMs, as shown using immunofluorescence assay (Fig. 6F). Furthermore, LILRB4

transcript levels were significantly increased in CRD-induced mammary glands (Fig. 6G).

Mammary gland LILRB4 transcript levels exhibited robust diurnal oscillations, increasing during

the light phase under normal and dark conditions. Under CRD, the amplitude of this rhythm was

dampened, but the phase was reserved (or only slightly shifted), and the transcript level peaked

during the dark phase (Fig. S10B). The transcript level of ARG1 increased modestly but non-

significantly in the CRD-induced mammary glands (Fig. S10C). The transcript level of the

chemokine CCL12 increased significantly in CRD-induced mammary glands (Fig. S10D). The

transcript level of CXCL5 is raised modestly but not significantly in CRD-induced mammary

glands (Fig. S10E). Overall, these changes in the immune milieus of the CRD-induced mammary

microenvironment can decrease adaptive immunity and enhance susceptibility to cancer and other

diseases.

To investigate whether LILRB4 expression was responsible for creating an immunosuppressive

TME, we explored the effects of LILRB4-immunotherapy on tumor progression in CRD mice by

treating them with an LILRB4 antibody (Sharma 2021). Briefly, 4T1 cells were orthotopically

transplanted into BALB/cJ mice to develop TNBC tumors. Four to five days (~4-5) days after the

transplantation, once palpable tumors developed, mice were placed into LD or CRD condition.

The LILRB4 antibody was administered on days 6, 9, and 12 after developing palpable tumors.

Mice were sacrificed 3 weeks after the development of palpable tumors (day 21) (Fig. 6H), and

lung metastasis and the TME were examined. We observed a significant decrease (P<0.0001) in

the prevalence of lung metastasis in LILRB4-antibody treated mice under CRD condition (Fig.

6I). Flow cytometric analysis showed an inhibition of the T_{reg} cell population in LILRB4-antibody

treated tumors under CRD condition, which alleviate the immunosuppressive microenvironment

(Fig. 6J). LILRB4-targeted therapy inhibited ARG1 transcript level (Fig. 6K) in CRD-induced

tumors. We also observed a modest but non-significant decrease in CCL12 (Fig. S10F) and CXCL5

(Fig. S10G) transcript levels in CRD-induced tumors. No significant difference was observed in

the number of lung metastasis foci in LD mice treated with the LILRB4-antibody (Fig S10H).

LILRB4-antibody treatment under LD conditions does not decrease the T_{reg} population (Fig. S10I),

ARG1 transcript level (Fig. S10J), and the chemokines (CCL12, CXCL5) (Fig. S10K, L)

significantly. These findings fortify the notion that elevated LILRB4 contributes to CRD-induced

aggressive tumorigenesis.

Discussion

Following the chronic jet lag protocol, here, we showed that CRD interrupted the biological clock,

affected the mammary gland morphology, and accelerated aggressive basal mammary

tumorigenesis and lung metastasis by altering the TME. CRD induced an immunosuppressive

TME and turned the tumors "cold." Malignant human breast cancer cells were shown to have high

CRD scores. We also identified the molecular mechanisms underlying CRD-induced aggressive

mammary tumorigenesis: CRD created "cold" tumors by inducing LILRB4 expression.

CRD-induced mammary glands exhibited underdeveloped (and abnormal) TEBs and

reduced branching morphology. CRD-induced mice showed mammary gland ductal hyperplasia,

which can be the early sign of ductal cancer (Behbod, Gomes et al. 2018), as seen by H&E and

Ki67 staining. This indicates the role of the circadian clock in mammary gland development and

the potential role of CRD in breast cancer initiation. Our study showed that CRD accelerated

spontaneous mammary tumor initiation and lung metastasis but not tumor progression, indicating

that CRD affects the early events of tumor initiation, rather than the growth of the established

tumors. LILRB4 expression was increased in mammary glands and CRD-induced mammary

tumors, revealing an unidentified mechanism responsible for CRD-induced tumorigenesis, lung

metastasis, and mammary gland development. The jet lag protocol used here mimics the effects of

rotating shift work or frequent eastbound trans-meridian flights and has been previously shown to

cause severe perturbations of circadian rhythmicity (Papagiannakopoulos, Bauer et al. 2016,

Hadadi, Taylor et al. 2020, Pariollaud, Ibrahim et al. 2022). Consistent with the results of published

studies (Pariollaud, Ibrahim et al. 2022), we found that CRD affected the rhythmicity of many

clock genes (e.g., *Clock, Bmal1, Per1*, and *Cry1*) in the mammary glands; however, *Per2*

expression remained rhythmic after CRD, indicating that *Per2* is more sensitive to phase

entrainment signals of molecular oscillators in peripheral tissues.

Using scRNA-seq and TCGA data, we showed that the expression of certain circadian

genes was decreased in aggressive human TNBCs compared with that in the surrounding

mammary tissues. Our analysis also revealed that the malignant cells in aggressive human TNBCs

had higher CRD scores and downregulated *PER1*, *PER2*, and *CRY1* significantly than

nonmalignant cells.

Our study showed that CRD induced an immunosuppressive tumor microenvironment,

which could be involved in increased tumor burden in mice maintained under the CRD schedule.

The TME consists of natural killer cells, $CD8^+$ T cells, $CD4^+$ helper T (T_h) cells, proinflammatory

macrophages (M1), and DCs, which elicit antitumor immune responses, whereas the presence of

MDSCs and $FOXP3^+$ T_{reg} counteracts tumor immunity (Labani-Motlagh, Ashja-Mahdavi et al.

2020). Our scRNA-seq study, and FACS showed that CRD decreased cytotoxic T-cell infiltration,

and the DC population. This supports the recent finding that DC and T-cell-autonomous circadian

clocks are responsible for time-of-the-day-dependent antitumor effects (Wang, Barnoud et al.

2023). CRD-induced $S100a4^+$ CAF, and α-SMA^+ CAF populations are known to promote tumor

progression and metastasis and contribute to an immunosuppressive TME (Costa, Kieffer et al.

2018). We found an increase in the M2 macrophage, and MDSC population and a decrease in the

M1 macrophage population in CRD tumors. M2 macrophages play a significant role in creating

an immunosuppressive microenvironment (Hao, Lu et al. 2012), whereas M1 macrophages induce

inflammation and kill tumor cells (Hao, Lu et al. 2012, Pan, Yu et al. 2020). MDSCs are also

known to create an immunosuppressive environment by suppressing T-cell responses and $CD8^+$ T-

cell infiltration (Li, Shi et al. 2021). Our FACS results corroborated the scRNA-seq results and

confirmed that CRD enhanced the T_{reg} population. CRD increased the leukocyte ($CD45^+$)

population, corroborating that the jet lag schedule enhances leukocyte levels in melanoma (Aiello,

Fedele et al. 2020). Our scRNA-seq and real-time PCR studies showed that CRD induced a pro-

tumorigenic and immunosuppressive microenvironment by increasing the expression of CCL12

and CXCL5. The upregulation of CCL12 (human orthologs of CCL2) (Yang J 2020) and CXCL5

reportedly lead to the accumulation of MDSCs (Li, Shi et al. 2021). CRD also enhances CCL28

production, which leads to the recruitment of T_{reg} cells in the TME (Ren, Yu et al. 2016).

We further demonstrated that CRD enhanced the expression of LILRB4 *in vivo* in healthy

mammary glands and mammary tumors. Recent studies have shown that LILRB4 creates an

immunosuppressive microenvironment in AML (Deng, Gui et al. 2018) and solid cancers (Sharma

2021) by inhibiting $CD8^+$ T-cell infiltration and inducing T_{reg}. LILRB4 inhibited the differentiation

of $CD8^+$ and $CD4^+$ T cells (Deng, Gui et al. 2018). By activating the JAK/STAT signaling pathway,

LILRB4 controls the expression of cytokines in macrophages (Truong, Hong et al. 2019). In

different malignant cancers, LILRB4 expression in MDSCs is associated with decreased survival

in patients (de Goeje, Bezemer et al. 2015). A recent study showed that LILRB4 blockade

increased the proportions of tumor immune infiltrates, effector (T_{eff}) levels, and altered the TME

toward reduced immunosuppression in solid tumors (Sharma 2021). Together, these findings

reveal that LILRB4 creates an immunosuppressive microenvironment, thereby decreasing its

antitumor efficacy. Our study showed that CRD activated LILRB4 signaling in healthy mammary

glands and aggressive TNBCs. Using real-time PCR and immunofluorescence analysis, we

observed increased *ARG1* expression in CRD-induced tumors. Analysis of the TME using

multiplexing immunostaining showed an immunosuppressive microenvironment as the cytotoxic

T-cell and M1 macrophage populations decreased. In contrast, T_{reg}, and M2 macrophage

populations were increased in CRD tumors. We found that the disruption of circadian rhythm

upregulated LILRB4 expression, which correlated with mammary tumor progression and

abnormal morphology in the mammary glands.

CRD upregulated the expression of immunosuppressive chemokines (CCL12) in mouse

mammary glands. CCL2 (an orthologous of mouse CCL12) increases stromal density and elevates

cancer risk (Sun, Glynn et al. 2017). An immunosuppressive environment can increase

susceptibility to opportunistic infections (Shu Kurizky, Dos Santos Neto et al. 2020) in CRD-

induced patients. This study demonstrated how an inhibitory immune receptor altered the TME

and influenced aggressive mammary tumorigenesis in response to CRD. We showed that the

targeted LILRB4 immunotherapy reduced CRD-induced lung metastasis by inhibiting

immunosuppressive TME.

Our findings suggest that LILRB4 is a potential therapeutic target for mitigating CRD-

induced cancer risk in populations exposed to chronic CRD, such as shift workers. LILRB4

inhibition was also found to improve the immunosuppressive microenvironment. However, how

the disruption of the cell-autonomous molecular clock by CRD could enhance LILRB4 expression

remains unknown. Further investigations are needed to determine why and how CRD enhances

LILRB4 expression. Additional investigations are required to determine whether the chronic

elevation of LILRB4 levels in the mammary glands in response to CRD occurs early in the disease

process and whether LILRB4 is present in other anatomical locations.

In conclusion, our study provides the link between CRD and aggressive basal mammary

tumorigenesis, metastasis, and abnormal mammary gland morphology. Based on our data, we

propose a signaling cascade (**Fig. 7**) as a possible mechanism behind CRD-induced mammary

gland disruption, enhanced aggressive tumorigenesis, lung metastasis, and immunosuppressive

TME.

609
List of abbreviations:

LD   Light and Dark

CRD   Circadian Rhythm Disruption

LILRB4   leukocyte immunoglobulin-like receptor 4

TNBC   Triple-negative Breast Cancer

ER   Estrogen receptor

PR   Progesterone receptor

HER2   Human Epidermal Growth Factor Receptor 2

SCN   Suprachiasmatic nucleus

CLOCK   Circadian locomotor output cycles kaput

BMAL-1   Brain and muscle Arnt-like protein-1

PAS   Per-Arnt-Sim domain

PER   Period

CRY   Cryptochrome

TME   Tumor microenvironment

GEMM   Genetically engineered mouse model

FVB   Friend LeukemiaVirus B

TEB   Terminal end bud

H&E   Hematoxylin and Eosin

AGR   Albumin-to-globulin ratio

scRNA-seq   Single-cell RNA sequence

631	CD	Cluster of differentiation
632	KLF2	Krüppel-like factors2
633	SLC2A1	Solute Carrier Family 2 Member 1
634	TRPS1	Transcriptional Repressor GATA Binding 1
635	TOP2A	Topoisomerase II Alpha
636	FTH1	Ferritin heavy chain 1
637	NES	Nestin
638	CLEC11A	C-Type Lectin Domain Containing 11A
639	S100A4	S100 Calcium Binding Protein A4
640	RPL35A	Ribosomal Protein L35a
641	FXYD3	FXYD Domain Containing Ion Transport Regulator 3
642	COL1A2	Collagen Type I Alpha 2 Chain
643	EPCAM	Epithelial cellular adhesion molecule
644	CDK	Cyclin-dependent kinase
645	M1	Type1 macrophage
646	M2	Type2 macrophage
647	CAFs	Cancer-associated fibroblasts
648	L-R	Ligand-receptor
649	FGF	Fibroblast growth factor
650	PDGF	Platelet-derived growth factor
651	TNF	Tumor necrosis factor
652	NOTCH	Neurogenic locus notch homolog protein
653	TGFβ	transforming growth factor-beta
654	CoA	Coenzyme A
655	MxIF	Multiplex immunofluorescence
656	Treg	Regulatory T-cell
657	α-SMA	Alpha-smooth muscle actin
658	FN1	fibronectin 1
659	FACS	Fluorescence-activated cell sorting
660	FoxP3	forkhead box P3
661	CSC	Cancer stem cell

662	MDSCs	Myeloid-derived suppressor cells
663	TAMs	Tumor-associated macrophages
664	IFN	Interferon
665	IL	Interleukin
666	G-CSF	Granulocyte colony-stimulating factor
667	CCL2/12	Chemokine (C-C motif) ligand 2/12
668	CXCL5	C-X-C motif chemokine 5
669	PCR	Polymerase chain reaction
670	TCGA	The Cancer Genome Atlas
671	DGE	Differential gene expression
672	APOE	apolipoprotein E
673	SHP-2	SH2 domain-containing protein tyrosine phosphatase-2
674	uPAR	Urokinase-like plasminogen activator
675	ARG1	Arginase 1
676	AML	Acute Myeloid Leukemia
677	mIHC	multiplex immunofluorescence histochemistry
678	ARNT2	Aryl hydrocarbon receptor nuclear translocator 2
679	NR1D1	Nuclear Receptor Subfamily 1 Group D Member 1
680	RORB & C	RAR Related Orphan Receptor B and C
681	ITIMs	Immunoreceptor tyrosine-based inhibitory motifs
682	ITAMs	Immunoreceptor tyrosine-based activating motifs
683	APCs	Antigen-presenting cells
684	JAK/STAT	Janus kinase/signal transducers and activators of transcription

685 **Availability of data and materials**

686 All data needed to evaluate the conclusions in the paper are present in the paper and/or the
687 Supplementary Materials.
688

689 **Competing interests:** Authors declare that they have no competing interests.
690

691 **Funding:** This work was funded by the Texas A&M University Faculty Development Grant (to
692 T.R.S) and Strategic Transformative Research Funds (STRP to TRS), NIH grant NIH 5
693 P01HL160471 (to A.K) and NIH R01NS131628 (to A.K), NIH R35GM151020 (to J.J), and NSF
694 CCF-1934904 (to B.M).

Author contributions: Conceptualization: T.R.S. Methodology: O.O., M.S., A.C, D.J.B., T.N., D.F., Y.J-H., J.J and T.R.S. Investigation: O.O., M.S., A.C, D.J.B., T.N.,C.,N., T.R.S. Visualization: O.O, M.S., D.F., YJ-H., T.R.S. Supervision: T.R.S. Analysis of data: T.R.S., B.M., J.J., A.K. Writing—original draft: O.O., M.S., T.R.S. Review and editing: A.K., B.M., L.F and J.J.

Acknowledgments
We thank Dr. Mahul Chakraborty for helping with real-time PCR, Dr. Gus Wright for helping with flow cytometry experiments and analysis, histology core facility (Texas A&M University) for histology work, Ashley Nguyen, David Arreola, Subiksha Sankar, and Aaryan Amin for helping with circadian cabinets.

Reference:

Ai, D., et al. (2021). "TRPS1: a highly sensitive and specific marker for breast carcinoma, especially for triple-negative breast cancer." Modern Pathology **34**(4): 710-719.

Aiello, I., et al. (2020). "Circadian disruption promotes tumor-immune microenvironment remodeling favoring tumor cell proliferation." Sci Adv **6**(42).

Alluri, P. and L. A. Newman (2014). "Basal-Like and Triple-Negative Breast Cancers: Searching for Positives Among Many Negatives." Surgical Oncology Clinics of North America **23**(3): 567-577.

Almansour, N. M. (2022). "Triple-Negative Breast Cancer: A Brief Review About Epidemiology, Risk Factors, Signaling Pathways, Treatment and Role of Artificial Intelligence." Front Mol. Biosci **9**.

Anders, C. and L. A. Carey (2008). "Understanding and treating triple-negative breast cancer." Oncology (Williston Park) **22**(11): 1233-1239; discussion 1239-1240, 1243.

Baker, S. J., et al. (2022). "CDK4: a master regulator of the cell cycle and its role in cancer." Genes Cancer **13**: 21-45.

Battaglin, F., et al. (2021). "Clocking cancer: the circadian clock as a target in cancer therapy." Oncogene **40**(18): 3187-3200.

Behbod, F., et al. (2018). "Modeling Human Ductal Carcinoma In Situ in the Mouse." <u>J Mammary</u>
<u>Gland Biol Neoplasia</u> **23**(4): 269-278.

Bernal, A. and L. Arranz (2018). "Nestin-expressing progenitor cells: function, identity and
therapeutic implications." <u>Cell Mol Life Sci</u> **75**(12): 2177-2195.

Binnewies, M., et al. (2018). "Understanding the tumor immune microenvironment (TIME) for
effective therapy." <u>Nat Med</u> **24**(5): 541-550.

Blakeman, V., et al. (2016). "Circadian clocks and breast cancer." <u>Breast Cancer Res</u> **18**(1): 89.

Casiraghi, L. P., et al. (2012). "Forced Desynchronization of Activity Rhythms in a Model of
Chronic Jet Lag in Mice." <u>Journal of Biological Rhythms</u> **27**(1): 59-69.

CDC (2015). "Work Organization Characteristics (NHIS 2015) Charts; National Health Interview
Survey (NHIS 2015)." [https://wwwn.cdc.gov/Niosh-whc/chart/OHS-](https://wwwn.cdc.gov/Niosh-whc/chart/OHS-WORKORG?OU=*&T=OU&V=R)
[WORKORG?OU=*&T=OU&V=R](https://wwwn.cdc.gov/Niosh-whc/chart/OHS-WORKORG?OU=*&T=OU&V=R).

Chun, S. K., et al. (2022). "Disruption of the circadian clock drives Apc loss of heterozygosity to
accelerate colorectal cancer." <u>Science Advances</u> **8**(32): eabo2389.

Costa, A., et al. (2018). "Fibroblast Heterogeneity and Immunosuppressive Environment in
Human Breast Cancer." <u>Cancer Cell</u> **33**(3): 463-479.e410.

de Goeje, P. L., et al. (2015). "Immunoglobulin-like transcript 3 is expressed by myeloid-derived
suppressor cells and correlates with survival in patients with non-small cell lung cancer."
<u>OncoImmunology</u> **4**(7): e1014242.

Deng, M., et al. (2018). "LILRB4 signalling in leukaemia cells mediates T cell suppression and
tumour infiltration." <u>Nature</u> **562**(7728): 605-609.

Elston, C. W. and I. O. Ellis (2002). "Pathological prognostic factors in breast cancer. I. The value
of histological grade in breast cancer: experience from a large study with long-term follow-up."
<u>Histopathology</u> **41**(3A): 154-161.

Fei, F., et al. (2017). "Role of metastasis-induced protein S100A4 in human non-tumor
pathophysiologies." <u>Cell Biosci</u> **7**: 64.

Finak, G., et al. (2015). "MAST: a flexible statistical framework for assessing transcriptional
changes and characterizing heterogeneity in single-cell RNA sequencing data." <u>Genome Biology</u>
16(1): 278.

Grosche, L., et al. (2020). "The CD83 Molecule - An Important Immune Checkpoint." <u>Front</u>
<u>Immunol</u> **11**: 721.

Hadadi, E., et al. (2020). "Chronic circadian disruption modulates breast cancer stemness and
immune microenvironment to drive metastasis in mice." <u>Nature Communications</u> **11**(1): 3193.

Hadadi, E., Taylor, W., Li, X. M., Aslan, Y., Villote, M., Rivière, J., Duvallet, G., Auriau, C., Dulong,
S., Raymond-Letron, I., Provot, S., Bennaceur-Griscelli, A., & Acloque, H. (2020). "Chronic
circadian disruption modulates breast cancer stemness and immune microenvironment to drive
metastasis in mice." <u>Nature communications</u> **11**(1).

Han, C., et al. (2020). "Biomarkers for cancer-associated fibroblasts." <u>Biomark Res</u> **8**(1): 64.

Hao, N. B., et al. (2012). "Macrophages in tumor microenvironments and the progression of
tumors." <u>Clin Dev Immunol</u> **2012**: 948098.

Hao, Y., et al. (2021). "Integrated analysis of multimodal single-cell data." <u>Cell</u> **184**(13): 3573-
3587.e3529.

He, J., et al. (2017). "Prognostic Effect of Albumin-to-Globulin Ratio in Patients with solid
tumors: A Systematic Review and Meta-analysis." <u>J Cancer</u> **8**(19): 4002-4010.

He, L., et al. (2022). "Single-cell transcriptomic analysis reveals circadian rhythm disruption
associated with poor prognosis and drug-resistance in lung adenocarcinoma." <u>J Pineal Res</u> **73**(1):
e12803.

Hemmer, A., et al. (2021). "The Effects of Shift Work on Cardio-Metabolic Diseases and Eating
Patterns." <u>Nutrients</u> **13**(11).

Kelleher, F. C., et al. (2014). "Circadian molecular clocks and cancer." <u>Cancer Lett</u> **342**(1): 9-18.

Kersemans, V., et al. (2013). "Subcutaneous tumor volume measurement in the awake,
manually restrained mouse using MRI." <u>Journal of Magnetic Resonance Imaging</u> **37**(6): 1499-
1504.

Labani-Motlagh, A., et al. (2020). "The Tumor Microenvironment: A Milieu Hindering and
Obstructing Antitumor Immune Responses." <u>Front Immunol</u> **11**: 940.

Lee, S., et al. (2010). "Disrupting circadian homeostasis of sympathetic signaling promotes
tumor development in mice." <u>PLoS One</u> **5**(6): e10995.

Li, K., et al. (2021). "Myeloid-derived suppressor cells as immunosuppressive regulators and
therapeutic targets in cancer." <u>Signal Transduction and Targeted Therapy</u> **6**(1): 362.

Li, M., et al. (2023). "FXYD3 functionally demarcates an ancestral breast cancer stem cell
subpopulation with features of drug-tolerant persisters." <u>J Clin Invest</u> **133**(22).

Maroulakou, I. G., et al. (1994). "Prostate and mammary adenocarcinoma in transgenic mice
carrying a rat C3(1) simian virus 40 large tumor antigen fusion gene." <u>Proc Natl Acad Sci U S A</u>
91(23): 11236-11240.

Merrick, D., et al. (2019). "Identification of a mesenchymal progenitor cell hierarchy in adipose
tissue." <u>Science</u> **364**(6438).

Meyer, J. S., et al. (2011). "Optic vesicle-like structures derived from human pluripotent stem
cells facilitate a customized approach to retinal disease treatment." <u>Stem Cells</u> **29**(8): 1206-
1218.

Moreira, A. C., et al. (2022) H-Ferritin Produced by Myeloid Cells Is Released to the Circulation
and Plays a Major Role in Liver Iron Distribution during Infection. <u>International Journal of</u>
<u>Molecular Sciences</u> **23**, DOI: 10.3390/ijms23010269

Neubauer, E., et al. (2016). "Comparative evaluation of three proliferation markers, Ki-67,
TOP2A, and RacGAP1, in bronchopulmonary neuroendocrine neoplasms: Issues and prospects."
<u>Oncotarget</u> **7**(27): 41959-41973.

Pan, Y., et al. (2020). "Tumor-Associated Macrophages in Tumor Immunity." <u>Front Immunol</u> **11**:
583084.

Panda, S., et al. (2002). "Coordinated transcription of key pathways in the mouse by the
circadian clock." Cell **109**(3): 307-320.

Papagiannakopoulos, T., et al. (2016). "Circadian Rhythm Disruption Promotes Lung
Tumorigenesis." Cell Metabolism **24**(2): 324-331.

Pariollaud, M., et al. (2022). "Circadian disruption enhances HSF1 signaling and tumorigenesis in
Kras-driven lung cancer." Science Advances **8**(39): eabo1123.

Ren, L., et al. (2016). "Hypoxia-induced CCL28 promotes recruitment of regulatory T cells and
tumor growth in liver cancer." Oncotarget **7**(46): 75763-75773.

Ren, X., et al. (2022). "Immunological Classification of Tumor Types and Advances in Precision
Combination Immunotherapy." Front Immunol **13**: 790113.

Rodriguez de la Fuente, L., et al. (2021). "Tumor dissociation of highly viable cell suspensions for
single-cell omic analyses in mouse models of breast cancer." STAR Protoc **2**(4): 100841.

Sangwung, P., et al. (2017). "KLF2 and KLF4 control endothelial identity and vascular integrity."
JCI Insight **2**(4): e91700.

Schernhammer, E. S., et al. (2001). "Rotating night shifts and risk of breast cancer in women
participating in the nurses' health study." J Natl Cancer Inst **93**(20): 1563-1568.

Schmid, B., et al. (2011). "A New ImageJ Plug-in "ActogramJ" for Chronobiological Analyses."
Journal of Biological Rhythms **26**(5): 464-467.

Sharma, N., Atolagbe, O., Ge, Z., Allison, J (2021). "LILRB4 suppresses immunity in solid tumors
and is a potential target for immunotherapy." J of Exp Medicine **218**(7).

Shu Kurizky, P., et al. (2020). "Opportunistic tropical infections in immunosuppressed patients."
Best Pract Res Clin Rheumatol **34**(4): 101509.

Stephenson, E. M., et al. (2021). "Cancer clocks in tumourigenesis: the p53 pathway and
beyond." Endocrine-Related Cancer **28**(4): R95-R110.

Storch, K. F., et al. (2002). "Extensive and divergent circadian gene expression in liver and heart."
Nature **417**(6884): 78-83.

Strobl, H., et al. (1998). "Identification of CD68+lin− Peripheral Blood Cells with Dendritic
Precursor Characteristics1." The Journal of Immunology **161**(2): 740-748.

Sun, X., et al. (2017). "CCL2-driven inflammation increases mammary gland stromal density and
cancer susceptibility in a transgenic mouse model." Breast Cancer Res **19**(1): 4.

Tirosh, I., et al. (2016). "Dissecting the multicellular ecosystem of metastatic melanoma by
single-cell RNA-seq." Science **352**(6282): 189-196.

Truong, A. D., et al. (2019). "Characterization and functional analyses of novel chicken leukocyte
immunoglobulin-like receptor subfamily B members 4 and 5." Poult Sci **98**(12): 6989-7002.

Trzpis, M., et al. (2007). "Epithelial cell adhesion molecule: more than a carcinoma marker and
adhesion molecule." Am J Pathol **171**(2): 386-395.

Usary, J., Darr, UB., Pfefferle, A, Perou, C. (2017). "Genetically Engineered Mouse Models of
Distinct Breast Cancer Subtypes." Curr Protoc Pharmacol **72**(14): 38.31-38.11.

Van Dycke, K. C. G., et al. (2015). "Chronically Alternating Light Cycles Increase Breast Cancer
Risk in Mice." Current biology : CB **25**(14): 1932-1937.

Vilgelm, A. E. and A. Richmond (2019). "Chemokines Modulate Immune Surveillance in
Tumorigenesis, Metastasis, and Response to Immunotherapy." Frontiers in Immunology **10**.

Vitaterna, M. H., et al. (2019). "Genetics of Circadian Rhythms." Neurol Clin **37**(3): 487-504.

Wang, C., et al. (2023). "Dendritic cells direct circadian anti-tumour immune responses." Nature
614(7946): 136-143.

Wang, Y., et al. (2023). "Cell death-related biomarker SLC2A1 has a significant role in prognosis
prediction and immunotherapy efficacy evaluation in pan-cancer." Frontiers in Genetics **13**.

Wu, S. Z., et al. (2021). "A single-cell and spatially resolved atlas of human breast cancers."
Nature genetics **53**(9): 1334-1347.

Wu, T., et al. (2021). "clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data." <u>The Innovation</u> **2**(3): 100141.

Yang J, A. M., Ling S, Teitz-Tennenbaum S, Zemans RL, Osterholzer JJ, Sisson TH, Kim KK. (2020).
"Diverse Injury Pathways Induce Alveolar Epithelial Cell CCL2/12, Which Promotes Lung
Fibrosis." <u>Am J Respir Cell Mol Biol.</u> **62**(5): 622-632.

Yang, Z., L., Liu, K.Y., Lin, F.J., Shi, H.M., Ou, Z.L. (2017). "CCL28 promotes breast cancer growth
and metastasis through MAPK-mediated cellular anti-apoptosis and pro-metastasis." <u>Oncology</u>
<u>Reports</u> **38**(3): 1393-1401.

Yuan, X., et al. (2018). "Night Shift Work Increases the Risks of Multiple Primary Cancers in
Women: A Systematic Review and Meta-analysis of 61 Articles." <u>Cancer Epidemiol Biomarkers</u>
<u>Prev</u> **27**(1): 25-40.

936
937

Figures

Fig. 1. CRD severely impairs rhythmicity and disrupts mammary gland morphogenesis. (A) Schematic representation of the CRD protocol. **(B)** FVB female mice were housed under LD 12:12 or CRD for 8 weeks. Representative actograms of LD and CRD mice. LD: 12 h light and 12 h dark; CRD: circadian rhythm disruption using jet lag protocol, represented by a shortening of the dark period by 8 h every third day. **(C)** Mammary glands were collected at the indicated times. Gene expression normalized to GAPDH expression was measured using quantitative real-time PCR data represented as mean ± SEM for n=3 females per time point and light condition. Rhythmicity was determined using JTK_Cycle analyses; *PJTKcycle < 0.05, **PJTKcycle < 0.01, ***PJTKcycle < 0.001, and ****PJTKcycle < 0.0001. CRD disrupts branching morphogenesis in WT virgin FVB mice. **(D, E, F)** Representative images of branching and development of terminal ducts in LD and CRD animals are shown in LD and CRD-induced mammary glands **(D, E, F)**, scale bar: 350 μm, (*P < 0.05, error bars indicate SEM, n = 3). **(G)** The effect of CRD on mammary ductal hyperplasia is shown by hematoxylin and eosin (H&E) staining of the mammary gland **(G)** and by using Ki67 **(H)** of the mammary glands. Scale bar: 100 μm.

956
Fig. 2. CRD increases aggressive mammary tumorigenesis and lung metastasis. Experimental
timeline for evaluation of the effect of CRD on spontaneous TNBC in (FVB-Tg(C3-1-TAg)cJeg
(C3-TAg) mice **(A)**. Dashed lines highlight the start and end points of the experiment. **(B)** Tumor
burden (tumor to body weight ratio) as % in LD (n = 9) or CRD (n = 10) conditions. Column data
represent the mean ± SEM values for individual animals. The effect of CRD on tumor initiation is
shown in **(C)**. Number of metastatic foci in the lungs of LD (n=6) and CRD (n =7) mice at the
time of their sacrifice is shown in **(D)**. P-value obtained from a binomial two-sided test. Indicated
(n) represents the number of independent experiments as biological replicates. Metastatic foci in
the lungs are shown using H&E staining, scale bar 150 μm **(E)**. CRD-induced lung metastasis in
4T1 tumor model in BALB/cJ mice as shown by the graphical representation **(F)** and H&E staining
(G), scale bar 100 μm. **(H)** The percentage of CD49f-breast cancer stem cell (CSC) subpopulations
in LD and CRD tumors is shown as a graph. Results represent the mean ± SEM of 3–4 independent
experiments. P-value obtained from an unpaired two-sided *t*-test. Mammosphere formation
efficiency (MFE%) of LD and CRD tumor cells (n =3) is shown in Figure. **(I)** with lines indicating
the mean ± SEM. P-value obtained from an unpaired two-sided *t*-test. The formation of organoids
(tumoroids) from LD and CRD tumors is shown in the **J**. Scale bar, 250 μm.
973

974
Fig. 3. CRD alters the tumor microenvironment. Cell clusters from 10x Genomics scRNA-seq
analysis visualized using Uniform Manifold Approximation (UMAP). Colors indicate clusters of
various cell types **(A)**. Fraction of cells from tumors (LD and CRD) in each cluster **(B)**. Clusters
were annotated for their cell types as predicted using canonical markers and signature-based
annotation with Garnett. Expression markers for endothelial cells (*KLF2*), M2 macrophages
(*SLC2A1*), TNBC cancer cells (*TRPS1*), proliferating cells (*TOP2A)*, myeloid cells (*FTH1)*,
progenitor cells (*NES*), adipocytes (*CLEC11A*), mesenchymal cells (*S100A4*), angiogenic cells
(*RPL35A*), breast cancer stem cells *(FXYD3)*, dendritic cells (*CD68*), cancer-associated fibroblast
(*COL1A2*), epithelial cells (*EPCAM*), cycling cells (*CDK4*), and B-cells (*CD83*). The comparison
of the total number of interactions among different cell populations between LD and CRD tumors

is shown in **C**. Edge width is proportional to the number of interactions, which assesses the number
of ligand–receptor pairs contributing to the communication between two interacting cell
populations. CellChat analysis quantitatively analyzes intercellular communication networks **(D)**.
The analysis shows differential ligand–receptor interactions between LD and CRD tumors.
Significant signaling pathways were ranked based on differences in the information flow within
the inferred networks between LD and CRD tumors. Dot plot profiling of the Gene Ontology (GO)
analyses in each cluster **(E)**. The dot plot represents the average expression of GOs per cluster. The
color gradient of dots represents the expression level, whereas the size represents the percentage
of cells expressing any genes per cluster. The number indicates the total number of genes identified
belonging to GO pathways significantly enriched in the cluster.
995

Fig. 4. CRD turns tumors "cold." Flow cytometry showed a significant decrease in cytotoxic CD8 T cell population (A). An overall ratio of CD4/CD8 is shown in **(B)**. Flow cytometry revealed a significant increase in the FoxP3$^+$ T$_{reg}$ population in the tumors of CRD mice (n = 4) **(C).** The percentage of antitumor M1 macrophage population and pro-tumor M2 macrophage population (n = 4) in LD and CRD-induced 4T1 tumors is shown in **(D),** and an overall ratio of M1/M2 is shown in **(E)**. The percentage of total leukocytes (CD45$^+$ cells) in LD and CRD-induced tumors is shown in **(F)**. The percentage of DC cells **(G)** and MDSC **(H)** is shown in LD and CRD-induced tumors via flow cytometry (n=4). P-value obtained from an unpaired two-sided *t*-test. The relative distribution of key immune cell types in LD and CRD-induced tumors is represented in the donut chart **(I).** Representative immunostaining image of cytotoxic CD8 T-cell (CD8, green), M2

macrophages (CD186, Cyan), and T_{reg} (FOXP3, White) in the LD and CRD-induced GEMM
tumors (n = 4) Scale bar= 100μm **(J)**. Single-cell RNA seq (scRNA Seq) study showed epithelial
to mesenchymal (EMT) markers (Fibronectin and Snail1) **(K),** and cancer-associated fibroblast
(CAF) population based on S100a4, and α-SMA markers **(L),** in LD and CRD-induced tumors.
The expression of chemokines *Ccl12* **(M)** and *Cxcl5* **(N),** is shown using real-time PCR in LD and
CRD-induced tumors (n = 3 - 4).
1048

1049
Fig. 5. The human circadian clock is disrupted in breast cancer. UMAP of malignant and
nonmalignant cells (total cells = 42,512) captured across patients with (n = 9) triple-negative breast
tumors. Moving from the plot in **A**, 30,719 cells had an unknown inferCNV call and 11,793 had a
known inferCNV call. These 11,793 single cells were used for downstream analysis to detect
"malignant" and "nonmalignant" cells. The percentage of malignant and nonmalignant cells with
CRD^{high} is shown in **B**. The mean normalized expression values of the core circadian genes by the
cell type (malignant and nonmalignant) are presented in **C**. Red represents significant changes in
expression between malignant and nonmalignant cells. **(D)** The mean expression of the clock genes
(+/- standard error of mean) over patients by the type of tissue analyzed in TNBC tumors (n =
1104) and surrounding tissue (n = 113) using TCGA.
1060

1061
Fig. 6. CRD enhances LILRB4a expression to create an immunosuppressive tumor
microenvironment. (A) Heatmaps of differentially expressed genes between LD 12:12 and CRD
tumors identified using DESeq2 analyses of scRNA-seq. The transcript level of LILRB4 was
analyzed in LD and CRD-induced GEMM tumors **(B)** and 4T1 tumors **(C)** using real-time PCR.
(D) The expression of arginase 1 (*Arg1*, a downstream target of LILRB4a) was analyzed using
immunofluorescence. Scale bar: 300μm **(E)** The transcript level of *Arg1* in LD and CRD-induced
GEMM tumors was analyzed using real-time PCR. **(F)** Representative MxIF showing the
expression of LILRB4, CD8, CD163, and FOXP3 markers in LD and CRD-induced GEMM
tumors. Scale bar: 50μm. **(G)** The transcript level of *Lilrb4* in LD and CRD-induced mammary
glands was analyzed using real-time PCR. **(H)** 4T1 cells were injected into the mammary fat pads

of BALB/cJ mice for the LILRB4-targeted immunotherapy. The number of metastatic foci in the lungs of CRD (control) and CRD (LILRB4-antibody) mice at the time of their sacrifice is shown in **I.** The percentage of T_{reg} cells in control and LILRB4-antibody-treated CRD-induced tumors is shown in **J.** The transcript level of *Arg1* in control and LILRB4-antibody-treated CRD-induced tumors is shown by real-time PCR **(K).**

Fig 7. Diagrammatic representation of the molecular mechanism of CRD-induced immunosuppressive microenvironment in breast cancer. CRD enhances aggressive TNBC and lung metastasis by creating an immunosuppressive tumor microenvironment. CRD also disrupts the mammary gland morphology. Our study revealed that CRD enhanced LILRB4 expression, which induces an immunosuppressive "cold" TME by increasing the M2 macrophage and T_{reg} populations and decreasing M1 macrophage infiltration. Inhibition of LILRB4 via LILRB4-targeted antibody alleviates CRD-induced immunosuppressive TME and inhibits lung metastasis.