Forage conservation is a neglected nitrous oxide source

Seongmin Yang[a], Maheen Mahmood[a,1], Rudra Baral[b,2], Hui Wu[c], Marc Almloff[a,3], Lauren E. Stanton[a], Doohong Min[b], Brenda K. Smiley[d], J. Chris Iiams[d], Jisang Yu[e], and Jeongdae Im[a,4]

This manuscript was compiled on April 22, 2024

Agricultural activities are the major anthropogenic source of nitrous oxide (N_2O), an important greenhouse gas and ozone-depleting substance. However, the role of forage conservation as a potential source of N_2O has rarely been studied. We investigated N_2O production from the simulated silage of the three major crops—maize, alfalfa, and sorghum—used for silage in the US, which comprises over 90% of the total silage production. Our findings revealed a substantial N_2O could be generated, which potentially places forage conservation as the third largest N_2O source in the agricultural sector. Notably, the application of chlorate as an additive significantly reduced N_2O production, but neither acetylene nor intermittent exposure to oxygen showed any impact. Overall, the results highlight that denitrifiers, rather than nitrifiers, are responsible for N_2O production from silage, which was confirmed by molecular analyses. Our study reveals a previously unexplored source of N_2O and provides a crucial mechanistic understanding for effective mitigation strategies.

greenhouse gas | forage conservation | nitrous oxide | denitrification | sustainable agriculture

Nitrous oxide (N_2O) is the third most important greenhouse gas (GHG), following carbon dioxide (CO_2) and methane. On a molar basis, N_2O has a warming potential 300 times greater than CO_2 and remains in the atmosphere for an extended period, estimated at 100–150 years (1). Recent studies have also emphasized N_2O as the primary ozone-depleting substance in the stratosphere (2, 3). Atmospheric N_2O concentrations have increased by over 20% compared to preindustrial levels, with the fastest increase occurring in the last five decades (4, 5). While natural processes and human activities contribute to N_2O production, agricultural activities are one of the dominant sources, responsible for over two-thirds of global anthropogenic N_2O emissions (5–7). Numerous countries compile GHG emission inventories following the technical guidelines provided by the Intergovernmental Panel on Climate Change. For example, the United States Environmental Protection Agency (US EPA) annually publishes a report titled "Inventory of U.S. Greenhouse Gas Emissions and Sinks," which accounts for N_2O emissions from agricultural activities through three sources: agricultural land management, manure management, and the burning of agricultural residues.

Forages, the plant materials consumed by herbivores, are conserved to sustain livestock during periods of limited pasture growth or inadequate grazing conditions (8, 9). Globally significant for productive and efficient livestock production, forage conservation methods mainly involve hay and silage. For long-term storage, hay is dried to below 20% moisture (12%–20%, w/w) to curtail microbial growth and stored under aerobic conditions (8). Conversely, silage is produced at higher moisture levels (40%–70%, w/w) and stored strictly under anaerobic conditions to facilitate fermentation. Indigenous or exogenous lactic acid bacteria (LAB) convert soluble carbohydrates into organic acids, predominantly lactic acid, which acts as a natural preservative that inhibits unwanted microorganisms (10, 11). Commercial silage inoculants containing LAB, such as *Lentilactobacillus buchneri* and other facultative anaerobic bacteria, are used as additives to enhance silage fermentation (12). The market for global silage inoculants reached USD 503 million in 2021 and is projected to grow to USD 630 million by 2028 at a compound annual growth rate of 3.8% (13). Silage is a significant segment of the global livestock industry, with 162.3 million metric tons (MMT) harvested in the US in 2022 (14, 15).

Despite its high production volume and extensive use, forage conservation has been limitedly studied as a potential source of GHG emissions (15). While gas production during forage conservation has been investigated, previous studies have primarily focused on odorous chemicals, such as volatile organic compounds and ammonia (NH_3) (16–18). At best, the detection of N_2O has been reported in

Significance Statement

N_2O is the third most important greenhouse gas (GHG) and agriculture contributes 80% of the total anthropogenic emissions in the US. The major sources of N_2O in the agricultural sector identified by the USEPA include agricultural land management, manure management, and the field burning of agricultural residues. Here, we show that forage conservation could be a significant unaccounted source of N_2O, surpassing the field burning by 30. Our study provides a mechanistic understanding of N_2O production and a simple and effective remedy for reducing N_2O emissions. The findings have substantial implications for mitigating climate change, informing policymakers, and guiding future research on reducing greenhouse gas emissions from livestock production.

Author affiliations: [a]Department of Civil Engineering, Kansas State University, Manhattan, KS 66506; [b]Department of Agronomy, Kansas State University, Manhattan, KS 66506; [c]Department of Statistics, Kansas State University, Manhattan, KS 66506; [d]Corteva Agriscience, Forage Additive Research, Johnston, IA 50131; [e]Department of Agricultural Economics, Kansas State University, Manhattan, KS 66506

J.I. and S.Y designed research; S.Y., M.M., R.B., M.A., and L.E.S. performed research; H.W., D.M., B.K.S., J.C.I, J.Y, and J.I. analyzed data, and J.Y., and J.I. wrote the manuscript

The authors declare no competing interest.

[4]To whom correspondence should be addressed. E-mail: jeongdaei@ksu.edu

previous studies (19–22), but a quantitative assessment is still lacking. That is, our study, to our knowledge, is the first to provide comprehensive quantitative estimates on a per-crop basis, which can be scaled to national- and sector-level estimates for N_2O emission from forage conservation.

The precise mechanism underlying N_2O production remains partially understood, but microbial processes are widely considered the main contributors to N_2O emissions (5, 23–26). While nearly all microorganisms involved in the biogeochemical nitrogen cycle can potentially produce N_2O, specific microbial pathways including heterotrophic denitrification, NH_3 oxidation, and nitrifier denitrification, are pivotal to N_2O production (26, 27). Heterotrophic denitrification is a multistep respiration process that involves the reduction of oxidized mineral forms of nitrogen (i.e., nitrate (NO_3^-) and nitrite (NO_2^-)) to gaseous nitric oxide (NO), N_2O, and dinitrogen (N_2). This process typically occurs under anaerobic conditions, although recent discoveries have identified a new group of aerobic denitrifying bacteria (28). Conversely, NH_3 oxidation and nitrifier denitrification occur under aerobic conditions. Nitrification involves a two-step process: the oxidation of NH_3 to NO_2^- by ammonia-oxidizing bacteria (AOB) and archaea (AOA), followed by further oxidation to NO_3^- by NO_2^--oxidizing bacteria. N_2O is indirectly produced through the chemical decomposition of intermediate or end products of NH_3 oxidation (hydroxylamine, nitroxyl hydride, or NO_2^-) (29). Additionally, certain AOB can directly produce N_2O through nitrifier denitrification by oxidizing NH_3 to NO_2^- and subsequently reducing it to NO and N_2O (30–32).

In this study, we conducted a comprehensive quantitative assessment of N_2O emissions during forage conservation, especially in silage form. Simulated silages derived from three major silage crops in the US—maize, alfalfa, and sorghum—were monitored for N_2O emissions over a 4-week period. Our findings showed significant N_2O release from silages, making forage conservation the third-largest source of N_2O emissions within the agricultural sector. Further experiments confirmed the significant role of denitrification in N_2O production in conserved forages, as validated by molecular analyses.

Results

Chemical properties of silage materials. The freshly chopped, non-inoculated plant materials showed characteristic nutritional properties for each crop (SI Fig. S1). Among them, alfalfa, a leguminous crop, showed notably higher concentrations of proteins and amino acids than maize and sorghum. Conversely, cereal crops, such as maize and sorghum, exhibited a higher starch content than alfalfa. The total protein and amino acid contents in alfalfa decreased as it matured, accompanied by an increase in fiber (acid detergent fiber (ADF) and neutral detergent fiber (aNDF)) and lignin contents. Notably, the alfalfa variety HybriForce 3400 consistently showed higher protein and amino acid contents than HVX MegaTron, regardless of maturity stage.

N_2O emissions from simulated silage. The total amount of N_2O produced in the maize, alfalfa (HybriForce 3400, harvested at mid-bud stage, Cv_2 Hv_1), and sorghum silage over 28 d of incubation was 6.7 ($\pm$0.7), 62.3 ($\pm$4.0), and

Fig. 1. Cumulative N_2O production of (a) maize, alfalfa, and sorghum and (b) two distinct alfalfa varieties harvested at two different stages of maturity. The labels Cv_1 and Cv_2 denote the two alfalfa varieties, HVX MegaTron and HybriForce 3400, respectively. Similarly, the Hv_1 and Hv_2 denote alfalfa samples harvested at the mid-bud and early flowering stages, respectively. The error bars represent the standard deviations derived from the triple incubations. Error bars may not be visible if their magnitude is smaller than the symbols.

1.8 ($\pm$0.1) mL, which corresponded to 18.2 ($\pm$1.9), 169.7 ($\pm$10.9), and 4.8 ($\pm$0.2) g CO_2 equivalent (eq.) per kg_{DM}, respectively (Fig. 1a). N_2O emissions began immediately after incubation commenced, with the majority ($> 90\%$) being produced within 24 h for maize and sorghum and 5 d for alfalfa. For alfalfa harvested at the same growth stages, no significant difference in N_2O production was observed between the varieties ($p > 0.05$) (Fig. 1b). However, incubations with alfalfa harvested at the later maturity stages (i.e., early flowering stage, Hv_2) produced significantly lower N_2O emissions ($p < 0.05$) regardless of the varieties (Fig. 1b). The statistical analysis is summarized in SI Fig. S2.

Effects of different treatments on N_2O production. The effects of various treatments on N_2O production were examined within the same sample group, revealing consistent trends regardless of the crops or alfalfa varieties harvested at different growth stages (Fig. 2a). Compared to the controls (I^-), the addition of inoculants (I^+) had no significant effect on N_2O production ($p > 0.05$), except for sorghum, where a significant difference was observed ($p < 0.05$). Notably, applying chlorate treatment and the inoculant (I^+ Ch^+) significantly reduced N_2O production by up to 99%. The addition of acetate as an external carbon source alongside the inoculant (I^+ Ac^+) resulted in a statistically insignificant but numerically lower N_2O production than I^+ ($p > 0.05$). However, the effect of acetate on N_2O reduction was significant in alfalfa Cv_2 Hv_1 and sorghum ($p < 0.05$). Conversely, neither acetylene (C_2H_2) addition nor intermittent O_2 exposure at different

Fig. 2. Effects of various treatments on N_2O production. (a) No inoculant (I^-), crop-specific commercial silage inoculant (I^+), inoculant and chlorate ($I^+ Ch^+$), and inoculant and acetate ($I^+ Ac^+$). (b) Acetylene (C_2H_2) addition and intermittent exposure to oxygen at different periods. (c) Different chlorate concentrations. Oxygen was added to separate bottles on each injection date. The N_2O production was normalized by control, with the shaded area representing the standard deviation of the control. The labels Cv_1 and Cv_2 denote two alfalfa varieties, HVX MegaTron and HybriForce 3400, respectively. Similarly, the labels Hv_1 and Hv_2 denote alfalfa samples harvested at the mid-bud and early flowering stages, respectively. The error bars denote the standard deviations derived from the triple incubations.

periods (Days 3, 5, and 10) impacted N_2O production (Fig. 2b). Lower chlorate concentrations, as low as 0.01% (w/w), still achieved 92% N_2O reduction (Fig. 2c). Residual chlorate could not be quantified due to technical limitations in ion chromatography. The statistical analysis is summarized in SI Table S1.

Analysis of the correlation between N_2O production and fresh matter nutrient parameters. The relationship between various nutrient parameters and N_2O production in the controls (I^-) was measured using a Pearson correlation coefficient (Fig. 3). Parameters related to protein and amino acids, including crude protein (r = 0.98, p = 0.031), total amino acids (r = 0.99, p = 0.001), NO_3^--N (r = 0.94, p = 0.017), NH_3-N (r = 0.96, p = 0.010), and NDICP (r = 0.92, p = 0.029), exhibited strong correlations with N_2O production (r > 0.8 and p < 0.05). Conversely, parameters related to carbohydrates and fats, including ADF, aNDF, lignin, starch, ethanol-soluble carbohydrates, and total fatty acids, exhibited no significant correlation with N_2O production (r < 0.4 or p > 0.05).

Gene and transcript abundance dynamics. The abundance of the genes and transcripts in incubations with alfalfa harvested at the early flowering stage (Hv_2) under various treatments is summarized in Fig. 4. Notably, the abundance of *narG*, the gene encoding membrane-bound nitrate reductase, exhibited a marked two-order-of-magnitude decrease with the addition of chlorate ($I^+ Ch^+$), in which N_2O production was reduced by up to 99%, compared to I^+. The abundance of other denitrification genes showed no trend over time across the various treatments. Notably, the abundances of bacterial and archaeal *amoA* gene encoding ammonia monooxygenase were lower than those of denitrification genes and no trend was observed over time across the various treatments. Transcript analysis revealed the expression of *narG* was completely suppressed by the addition of chlorate ($I^+ Ch^+$), similar to the gene abundance. Transcripts of archaeal and bacterial *amoA*, both *nir* genes encoding nitrite reductase, and *nosZ* gene encoding nitrous oxide reductase clade II were not detected. The expression level of *napA* gene encoding periplasmic nitrate reductase, *norB* gene encoding nitric oxide reductase,

and *nosZ* gene encoding nitrous oxide reductase clade I were not affected by the treatments used.

Discussion

N_2O production mechanism. The majority of N_2O production in this study occurred within the first week of incubation (Fig. 1a), which is consistent with previous studies ([21], [22]). During this initial phase, transient aerobic or microaerobic conditions can be expected due to residual oxygen, followed by anaerobic conditions ([33]). Under these conditions, both ammonia oxidizers and denitrifiers are potential contributors to N_2O production ([34], [35]). Ammonia oxidizers are recognized for producing N_2O through direct (i.e., nitrifier denitrification) and indirect (i.e., abiotic processes) mechanisms. Furthermore, ammonia oxidizers can contribute to N_2O production by converting NH_3 to NO_3^-, subsequently fueling denitrification. However, N_2O production did not increase when oxygen was added at different time points (p > 0.05, Fig. 3b), indicating that the contribution of nitrification to N_2O production may not be significant. This finding is further supported by the observation that C_2H_2 (10 Pa), a potent inhibitor of bacterial and archaeal NH_3 oxidation ([36], [37]), did not affect N_2O production (p > 0.05, Fig. 3b). The absence of bacterial and archaeal *amoA* transcripts also confirmed the non-involvement of ammonia oxidizers in N_2O production (Fig. 4). At high concentrations (1–20 kPa), C_2H_2 has been shown to inhibit N_2O reductase activity of denitrifying microorganisms ([38], [39]). However, such inhibition (i.e., increased N_2O production) was not observed in this study (Fig. 2b). The expression of Clade I *nosZ* remained unaffected in the samples supplemented with C_2H_2 (SI Fig. S3), suggesting that C_2H_2 did not inhibit denitrifying bacteria at the concentration used in this study (i.e., 10 Pa).

Due to the chemical similarities between NO_3^- and chlorate, chlorate has been used as an inhibitor for dissimilatory NO_3^- reduction, the first step in denitrification ([40], [41]). The significant decrease in N_2O production, by up to 99% upon the addition of chlorate (Fig. 2a and SI Table S1), indicates that denitrifiers are the main contributors to N_2O production. This finding was further confirmed by the

Fig. 3. Correlation analysis of N_2O production and fresh matter nutrient parameters. DM: dry matter; CP: crude protein; ADICP: acid detergent insoluble CP; NDICP: neutral detergent insoluble CP; ADF: acid detergent treated fiber; aNDF: amylase treated neutral detergent fiber; ESC: ethanol soluble carbohydrates.

Yang *et al.*

Fig. 4. Abundance of functional genes (bar) and their transcripts (black circle) in incubation with alfalfa harvested at the early flowering stage (Hv₂) under various treatments. The error bars represent the standard deviations of the triplicate incubation. Error bars may not be visible if their magnitude is smaller than the symbols.

diminished abundance of the *narG* genes and transcripts in the chlorate-amended samples (Fig. 4). In another study, the same concentration of chlorate (0.1% w/w) was used as a ruminant supplement to reduce *E. coli* O157:H7 population (42), suggesting that chlorate has the potential to be used as a silage additive to reduce N_2O emissions. Further studies are warranted to assess the potential hazards such as the ultimate fate of the added chlorate and its impact on animal health if it remains in the silage.

Denitrification is a microbial process wherein NO_3^- is reduced to N_2 via intermediates including N_2O. Factors such as a low C/N ratio have been reported to lead to N_2O accumulation during denitrification (43, 44). The addition of acetate as an external carbon source, effectively increasing the C/N ratio, resulted in a slight reduction in N_2O production (Fig. 2a and SI Table S1). This suggests that N_2O production in conserved forage may be influenced, at least in part, by a low C/N ratio. Moreover, the findings also indicate that denitrification inhibitors, such as chlorate, can be combined

with an external carbon source, such as acetate, as an effective additive to mitigate N_2O emissions from the forage conservation process.

N_2O production and its relationship with nutritional parameters. N_2O production was closely correlated with most parameters related to protein and amino acids, including NO_3^--N (Fig. 3), presumably the main source of N_2O production. The NO_3^--N content in forages varies with the stage of plant maturity (45), and both alfalfa varieties harvested at later growth stages, which produced significantly less N_2O ($p < 0.05$) (Fig. 1b), contained lower NO_3^--N levels (SI Fig. S1). Reports have also demonstrated that nitrogen fertilization directly impact NO_3^--N content (45), implying that nitrogen fertilization, especially preceding harvest, may contribute to higher N_2O production at ensiling. Further studies are needed to investigate the effects of nitrogen fertilization schedule on N_2O production. Simple changes in agricultural practice may reduce N_2O emissions.

Gas/Source	2005	2019	2020	2021	2022
Agricultural Soil Management	291.5	309.3	290.5	294.0	_^{b}
Manure Management	14.5	17.4	17.5	17.4	_^{b}
Field Burning of Agricultural Residues	0.2	0.2	0.2	0.2	_^{b}
Forage Conservation^c	**5.6**	**5.6**	**5.7**	**5.8**	**5.3**

Note: Totals may not sum due to independent rounding

^{a} MMT: million metric tons

^{b} data not available

^{c} hypothetical estimation based on this study

A source of organic carbon is an important component of denitrification, serving as an electron donor. Many studies have shown that external carbon sources, such as methanol, ethanol, and acetate, stimulate denitrification and usually reduce N$_2$O production (46). Consistent with these findings, our study showed that the addition of acetate reduced N$_2$O production (Fig. 2a). However, carbohydrate-related parameters, such as ADF, aNDF, lignin, starch, and ethanol-soluble carbohydrates, did not correlate with N$_2$O production (Fig. 3), which could be due to the recalcitrance of these carbon sources. Similarly, denitrification was promoted by plant-based carbon substrates, such as rice straw, but there was a significant lag before denitrification became active (47).

Environmental implications. What gets measured gets managed. The first step in reducing GHG emissions is to measure them. The US EPA publishes an annual report titled "Inventory of U.S. Greenhouse Gas Emissions and Sinks," which estimates total GHG emissions by source across all sectors of the economy at the national level (1). Notably, agriculture is the largest contributor to N$_2$O emissions in the US, accounting for 80% in 2021. The EPA monitors major sources in the agricultural sector, including agricultural land management, manure management, and the field burning of agricultural residues (Table 1) (1). In our study, 18.2 ($\pm$1.9), 169.7 ($\pm$10.9), and 4.8 ($\pm$0.2) g CO$_2$ eq. per kg$_{DM-forage}$ of N$_2$O were produced from maize, alfalfa, and sorghum, respectively (Fig. 1a). According to the Crop Production 2022 Summary of the United States Department of Agriculture National Agricultural Statistics Service, the total production volumes of maize, alfalfa, and sorghum for silage in 2022 were 128.6, 17.4, and 5.6 MMT, respectively, comprising 93% of the total silage production combined (14). Assuming a similar amount of N$_2$O can be produced from each crop, the total N$_2$O emission potential amounts to 5.3 MMT CO$_2$ eq.. This makes forage conservation the third largest N$_2$O emitter in the agricultural sector, surpassing the field burning of agricultural residues by a factor of 30 (Table 1). Notably, N$_2$O emissions from silage of uncategorized crops (total production volume: 10.7 MMT in 2022, comprising 7% of the total silage production) were not considered in this study (14). Again, the first step to reducing GHG emissions is to measure them, as policymakers and decision-makers use GHG inventories to develop strategies and track progress in GHG emission reduction efforts (48, 49).

Limitations and Perspectives. There are, admittedly, limitations to our assessment, especially in extrapolating our data to the national scale. The microbial N$_2$O production, like any other microbial processes, is sensitive to various environmental factors such as temperature and nutritional parameters, which could result in underestimation or overestimation of the outcomes. There are additional uncertainties associated with the heterogeneity of farmers' practices, such as harvest time, inoculant use, moisture content (i.e., wilting), and oxygen exposure. Therefore, N$_2$O emission measurements from actual silage fermentation systems spanning the range of environmental and management variation across the US are warranted to achieve more accurate estimation results. With that, this work aims to draw attention to the silage process as an overlooked but abundant source of N$_2$O in the agricultural sector.

In this study, the use of chlorate was proposed as a simple and effective remedy to reduce N$_2$O emissions significantly (Fig. 2a), and additional experiments demonstrated that lower concentrations as low as 0.01% (w/w) still achieved 92% N$_2$O reduction (Fig. 2c). The chlorate (as sodium salt) price at the end of 2023 in the US was 795 USD per ton (50). The estimated cost to add 0.01% (w/w) chlorate as an additional silage additive is 0.08 USD per ton forage (dry weight), which is only approximately 5-8% compared to the silage inoculant cost (i.e., 1-1.5 USD per ton). Assessing the social cost of GHG emissions has become a common yardstick for estimating the benefits of formulating and implementing abatement policies (51). The social cost estimates of N$_2$O shown in recent studies range from 16 to 174 USD per kg-N$_2$O-N (52–54). Assuming that chlorate (0.01%, w/w) is added to achieve 90% N$_2$O reduction (Fig. 2c), the social cost saving could be 81-882 million USD with the expected total cost of 12 million USD for applying chlorate as an additive.

Materials and Methods

Forage sample preparation. Maize (Dyna Gro D53VC55RIB) and sorghum (Dyna Gro Super Sile 20) were harvested from a private farm near Manhattan, KS (39°38'N, 96°88'W) at their optimal maturity stages (i.e., 2/3 milk line and soft dough stages, respectively). The plants were chopped to a theoretical length of 2 cm using a standard forage harvester without inoculation. Two distinct alfalfa varieties, HVX MegaTron (WinField United L.L.C., Arden Hills, MN, USA) and HybriForce 3400 (Dairyland Seed Co.), were grown in experimental plots at the Kansas State University Agronomy Research Farm in Manhattan, Kansas (39°20´ N, 96°59´W). Each variety was harvested at two different stages of maturity (mid-bud and early flowering) using a sickle bar mower. The harvested alfalfa was subsequently chopped to a length of 2 cm using a stationary forage chopper.

Experimental design. We used a simulated silage model known as mini silos. These mini-silos comprised a 1-L glass jar (55) connected to a 3-L Tedlar bag (SI Fig. S4. Each crop was ensiled with four treatments: 1) no inoculant (control, I⁻), 2) crop-specific commercial silage inoculant (brand names omitted for confidentiality) following the manufacturer's instructions (I⁺), 3) inoculant + chlorate (0.1%, w/w, potassium salt) (I⁺ Ch⁺), and 4) inoculant + acetate (0.1%, w/w, sodium salt) (I⁺ Ac⁺). Chlorate was added to inhibit denitrifiers (40, 41), and acetate was added to increase the initial C/N ratio (43, 44). Separate inoculant, chlorate, and acetate solutions were prepared and sprayed onto the plants. The initial moisture content was measured using the conventional microwave oven method (56) and adjusted to 70% (w/w, wet weight basis) by adding deionized water to the treatment solutions. All crops were packed in mini-silos with a bulk density of 650 kg/m^3 (41 lb/ft3) (57), and each silo contained 650 g of the crops (wet weight), which was equivalent to 195 g$_{DM}$. The mini-silos were incubated at 30°C in the dark, and N$_2$O production was monitored regularly for up to 4 weeks. Gas bags were removed

on Days 1, 2, 3, 5, and 20 to characterize temporal variations in the volume and composition of gas production and replaced with new bags at each sampling. Fresh feed samples were collected and stored at -80°C for chemical analysis. Additionally, four mini-silos were prepared for each treatment and sacrificed on Days 1, 3, 5, and 20 for molecular analysis. A subset of the samples (5 g) was preserved in 5 mL RNA preservation solution (RNAprotect, QIAGEN, Hilden, Germany) and stored at -80°C. Each treatment comprised three replicates of the mini-silos.

Contribution of nitrification to N_2O emissions. To investigate the relative contribution of nitrification to N_2O production, mini-silos were prepared with alfalfa (HVX Megatron) harvested at the early flowering stage, and acetylene (10 Pa) [36] was added at the beginning of incubation. In addition, the mini-silos were intermittently exposed to oxygen on Days 3, 5, and 10, achieved by supplying 20 mL of air (4 mL of oxygen) through a stainless-steel tube that reached the center of the mini-silos (Supplementary Fig. 3). Oxygen was added to separate bottles on each injection date.

Analytical methods. The total gas production volume was measured using the water displacement method. N_2O was quantified using an Agilent 7890 gas chromatograph with an electron capture detector and an HP-PLOT/Q column (30 m × 0.53 mm × 40 m). Fresh forage samples were sent to Rock River Laboratory, Inc., Watertown, WI, for nutritional analysis. The following parameters were examined: crude protein (CP), total amino acid, NH3-N content, acid detergent fiber (ADF), amylase-treated neutral detergent fiber (aNDF), lignin, starch, and ethanol-soluble carbohydrate (ESC). NO_3^--N content (% of DM) was measured at the Kansas State University Soil Testing Laboratory.

Nucleic acid extraction and quantitative PCR. Microbial nucleic acid extraction from plant material, particularly from the epiphytic phyllosphere, poses challenges due to plant-derived biomolecules such as proteins and nucleic acids [58, 59]. In this study, we used a microbial DNA extraction method optimized for silage samples, as detailed in our previous study [60]. Briefly, for DNA extraction, five grams of the sample was suspended in 45 mL of sterile 0.85% NaCl solution, shaken on a rotary shaker at 120 rpm for 2 h at room temperature, filtered through two layers of gauze cloth to remove large plant debris, and centrifuged at 12,000 g for 15 min at 4°C [61]. The supernatant was discarded, and the pellet was stored at -80°C until subsequent DNA extraction. The RNA preservation solution containing the sample was shaken on a rotary shaker at 120 rpm for 2 h at room temperature. Four milliliters of the supernatant were collected, pelleted by centrifugation at 12,000 g for 15 min at 4°C, and immediately subjected to RNA

extraction. DNA and RNA extractions from the pellets were performed using the DNeasy and RNeasy PowerSoil kits (Qiagen, Hilden, Germany), following the manufacturer's instructions, with slight modifications. Cells were lysed by bead beating at 20°C for 5 min. RNA samples were subjected to DNase treatment (ezDNase™, Invitrogen, CA, USA) and reverse transcribed into complementary DNA (cDNA) using SuperScriptIV Reverse Transcriptase (Thermo Fisher Scientific, Waltham, MA, USA). PCR was conducted with universal 16S rRNA gene primers on DNase-treated RNA samples to confirm the absence of DNA contamination. Target genes and transcripts were quantified by qPCR using a CFX Opus 96 Real-Time PCR System (Bio-Rad, Hercules, CA, USA) in 20 μL reaction mixtures containing 10 μL of SsoAdvanced™ Universal SYBR Green Supermix (Bio-Rad, Hercules, CA, USA), 300 nM of each primer, and 2 μL of template DNA. Quantification was performed using standard curves prepared from serial 10-fold dilutions of cloned plasmids or double-stranded synthetic DNA fragments (gBlocks®, Integrated DNA Technologies, Coralville, IA, USA) in triplicate. Detailed information on primer sequences, standard sequences, and detection limits can be found in SI Table S2.

Statistical analysis . Statistical analyses were conducted using the MIXED procedures of SAS/STAT software, Version 9.4 (SAS Institute Inc., Cary, NC, USA). A two-way ANOVA was applied to the N_2O emission data, crop, treatment, and their interaction term as fixed effects. A Bonferroni multiplier adjustment was performed when comparing N_2O emissions between crops in each

treatment group or treatments in each crop group (conditional pairwise comparison). Probability values of $p < 0.05$ (2-tailed) were considered statistically significant for all comparisons. The correlation coefficients between N_2O production and nutrient variables were calculated as Pearson correlation coefficients. A Pearson correlation coefficient > 0.8 or < -0.8 with $p < 0.05$ (two-tailed) was deemed indicative of a strong correlation between the two variables.

Data Availability. All data are included in the article and Supplementary material.

ACKNOWLEDGMENTS. Please include your acknowledgments here, set in a single paragraph. Please do not include any acknowledgments in the Supporting Information, or anywhere else in the manuscript. Support for this research was provided by the Kansas NSF EPSCoR (OIA-1656006) and NSF CAREER Program (2144189).

1. USEPA, Inventory of u.s. greenhouse gas emissions and sinks: 1990-2021 – main report (2023).
2. AR Ravishankara, JS Daniel, RW Portmann, Nitrous oxide (n2o): The dominant ozone-depleting substance emitted in the 21st century. *Science* **326**, 123–125 (2009).
3. X Fang, et al., Challenges for the recovery of the ozone layer. *Nat. Geosci.* **12**, 592–596 (2019).
4. BD Hall, GS Dutton, JW Elkins, The noaa nitrous oxide standard scale for atmospheric observations. *J. Geophys. Res. Atmospheres* **112** (2007).
5. H Tian, et al., A comprehensive quantification of global nitrous oxide sources and sinks. *Nature* **586**, 248–256 (2020).
6. EA Davidson, D Kanter, Inventories and scenarios of nitrous oxide emissions. *Environ. Res. Lett.* **9** (2014).
7. DS Reay, et al., Global agriculture and nitrous oxide emissions. *Nat. Clim. Chang.* **2**, 410–416 (2012).
8. M Owens V N., *Preservation of Forage as Hay and Silage*, eds. RF Barnes, CJ Nelson, M Collins, KJ Moore. (Wiley-Blackwell) Vol. 56, 6th edition edition, p. 673 (2003).
9. H Finch, A Samuel, G Lane, *Conservation of grass and forage crops*. (Elsevier), pp. 513–526 (2014).
10. CE Müller, D von Rosen, P Udén, Effect of forage conservation method on microbial flora and fermentation pattern in forage and in equine colon and faeces. *Livest. Sci.* **119**, 116–128 (2008).
11. I Soundharrajan, et al., Application of customised bacterial inoculants for grass haylage production and its effectiveness on nutrient composition and fermentation quality of haylage. *3 Biotech* **7**, 1–9 (2017).
12. RE Muck, et al., Silage review: Recent advances and future uses of silage additives. *J. Dairy Sci.* **101**, 3980–4000 (2018).
13. Facts, Factors, Silage inoculants and enzymes market size, share, growth analysis report 2022 – 2028 (2023).
14. USDA, Crop production 2022 summary (2023).
15. G Grossi, P Goglio, A Vitali, AG Williams, Livestock and climate change: Impact of livestock on climate and mitigation strategies. *Animal Front.* **9**, 69–76 (2019).
16. SE Pusede, RC Cohen, On the observed response of ozone to nox and voc reactivity reductions in san joaquin valley california 1995–present. *Atmospheric Chem. Phys.* **12**, 8323–8339 (2012).
17. SD Hafner, et al., Emission of volatile organic compounds from silage: Compounds, sources, and implications. *Atmospheric Environ.* **77**, 827–839 (2013).
18. ME Morgan, RL Pereira, Volatile constituents of grass and corn silage. ii. gas-entrained aroma. *J. Dairy Sci.* **45**, 467–471 (1962).
19. AVI Bueno, et al., Natamycin as a potential silage additive: A lab trial using sugarcane to assess greenhouse gas emissions. *Revista Brasileira de Zootecnia* **49** (2020).
20. LC Wang, RH Burris, Mass spectrometric study of nitrogenous gases produced by silage. *J. Agric. Food Chem.* **8**, 239–242 (1960).
21. MS Krommweh, AJ Schmithausen, HF Deeken, W Büscher, GC Maack, A new experimental setup for measuring greenhouse gas and volatile organic compound emissions of silage during the aerobic storage period in a special silage respiration chamber. *Environ. Pollut.* **267**, 115513 (2020).
22. AJ Schmithausen, et al., Greenhouse gas formation during the ensiling process of grass and lucerne silage. *J. Environ. Manag.* **304** (2022).
23. AJ Thomson, G Giannopoulos, J Pretty, EM Baggs, DJ Richardson, Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philos. Transactions Royal Soc. B: Biol. Sci.* **367**, 1157–1168 (2012).
24. HW Hu, D Chen, JZ He, Microbial regulation of terrestrial nitrous oxide formation: Understanding the biological pathways for prediction of emission rates. *FEMS Microbiol.*

869 *Rev.* **39**, 729–749 (2015).

25. IM Klimasmith, AD Kent, Micromanaging the nitrogen cycle in agroecosystems. *Trends Microbiol.* **30**, 1045–1055 (2022).

26. MMM Kuypers, HK Marchant, B Kartal, The microbial nitrogen-cycling network. *Nat. Rev. Microbiol.* **16**, 263–276 (2018).

27. K Butterbach-Bahl, EM Baggs, M Dannenmann, R Kiese, S Zechmeister-Boltenstern, Nitrous oxide emissions from soils: How well do we understand the processes and their controls? *Philos. Transactions Royal Soc. B: Biol. Sci.* **368** (2013).

28. J Yang, et al., A critical review of aerobic denitrification: Insights into the intracellular electron transfer. *Sci. The Total. Environ.* **731**, 139080 (2020).

29. S Liu, et al., Abiotic conversion of extracellular nh2oh contributes to n2o emission during ammonia oxidation. *Environ. Sci. Technol.* **51**, 13122–13132 (2017).

30. N Wrage-Mönnig, et al., The role of nitrifier denitrification in the production of nitrous oxide revisited. *Soil Biol. Biochem.* **123**, A3–A16 (2018).

31. N Wrage, G Velthof, M van Beusichem, O Oenema, Role of nitrifier denitrification in the production of nitrous oxide. *Soil Biol. Biochem.* **33**, 1723–1732 (2001).

32. JA Kozlowski, KD Kits, LY Stein, Comparison of nitrogen oxide metabolism among diverse ammonia-oxidizing bacteria. *Front. Microbiol.* **7** (2016).

33. CL Ávila, BF Carvalho, Silage fermentation—updates focusing on the performance of micro-organisms. *J. Appl. Microbiol.* **128**, 966–984 (2020).

34. F Schinner, R Öhlinger, E Kandeler, R Margesin, *Nitrification and Denitrification.* (Springer Berlin Heidelberg), pp. 144–161 (1996).

35. R Barnard, PW Leadley, BA Hungate, Global change, nitrification, and denitrification: A review. *Glob. Biogeochem. Cycles* **19**, 1–13 (2005).

36. Z Jia, R Conrad, Bacteria rather than archaea dominate microbial ammonia oxidation in an agricultural soil. *Environ. Microbiol.* **11**, 1658–1671 (2009).

37. P Berg, L Klemedtsson, T Rosswall, Inhibitory effect of low partial pressures of acetylene on nitrification. *Soil Biol. Biochem.* **14**, 301–303 (1982).

38. T Yoshinari, R Hynes, R Knowles, Acetylene inhibition of nitrous oxide reduction and measurement of denitrification and nitrogen fixation in soil. *Soil Biol. Biochem.* **9**, 177–183 (1977).

39. R Knowles, Denitrification. *Microbiol. Rev.* **46**, 43–70 (1982).

40. I Kučera, Interference of chlorate and chlorite with nitrate reduction in resting cells of paracoccus denitrificans. *Microbiology* **152**, 3529–3534 (2006).

41. O R.S., D Capone, *Use of "specific" inhibitors in biogeochemistry and microbial ecology*, ed. KC Marshall. (Springer US) Vol. 10, 1 edition, (1988).

42. TR Callaway, et al., Sodium chlorate supplementation reduces e. coli o157:h7 populations in cattle. *J. Animal Sci.* **80**, 1683–1689 (2002).

43. MJ Kampschreur, H Temmink, R Kleerebezem, MS Jetten, MC van Loosdrecht, Nitrous oxide emission during wastewater treatment. *Water Res.* **43**, 4093–4103 (2009).

44. H Itokawa, K Hanaki, T Matsuo, Nitrous oxide production in high-loading biological nitrogen removal process under low cod/n ratio condition. *Water Res.* **35**, 657–664 (2001).

45. LS Murphy, GE Smith, Nitrate accumulations in forage crops. *Agron. J.* **59**, 171–174 (1967).

46. X Fu, et al., Application of external carbon source in heterotrophic denitrification of domestic sewage: A review. *Sci. The Total. Environ.* **817**, 153061 (2022).

47. B Zhou, J Duan, L Xue, J Zhang, L Yang, Effect of plant-based carbon source supplements on denitrification of synthetic wastewater: focus on the microbiology. *Environ. Sci. Pollut. Res.* **26**, 24683–24694 (2019).

48. G Iyer, et al., Measuring progress from nationally determined contributions to mid-century strategies. *Nat. Clim. Chang.* **7**, 871–874 (2017).

49. MJ Gidden, et al., Aligning climate scenarios to emissions inventories shifts global benchmarks. *Nature* **624**, 102–108 (2023).

50. ChemAnalyst, Sodium chlorate price trend and forecast (2024).

51. JE Aldy, MJ Kotchen, RN Stavins, JH Stock, Keep climate policy focused on the social cost of carbon. *Science* **373**, 850–852 (2021).

52. K Rennert, et al., Comprehensive evidence implies a higher social cost of co2. *Nature* **610**, 687–692 (2022).

53. T Wang, F Teng, Damage function uncertainty increases the social cost of methane and nitrous oxide. *Nat. Clim. Chang.* **13**, 1258–1265 (2023).

54. DR Kanter, et al., Improving the social cost of nitrous oxide. *Nat. Clim. Chang.* **11**, 1008–1010 (2021).

55. FE Contreras-Govea, RE Muck, DR Mertens, PJ Weimer, Microbial inoculant effects on silage and in vitro ruminal fermentation, and microbial biomass estimation for alfalfa, bmr corn, and corn silages. *Animal Feed. Sci. Technol.* **163**, 2–10 (2011).

56. KRdS J. W. Lacerda, M. J. R. ; Freitas, Determining forage dry matter using microwave oven and conventional method. *Biosci. J.* **35**, 185–190 (2009).

57. D Darby, J Jofriet, Density of silage in horizontal silos. *Can. Agric. Eng.* **35**, 275–280 (1993).

58. KC Dent, JR Stephen, WE Finch-Savage, Molecular profiling of microbial communities associated with seeds of beta vulgaris subsp. vulgaris (sugar beet). *J. Microbiol. Methods* **56**, 17–26 (2004).

59. S Ikeda, et al., Development of a bacterial cell enrichment method and its application to the community analysis in soybean stems. *Microb. Ecol.* **58**, 703–714 (2009).

60. S Yang, J Lee, M Mahmood, D Min, J Im, Evaluation and optimization of lysis method for microbial dna extraction from epiphytic phyllosphere samples. *J. Microbiol. Methods* **206**, 106677 (2023).

61. A Saito, S Ikeda, H Ezura, K Minamisawa, Microbial community analysis of the phytosphere using culture-independent methodologies. *Microbes environments* **22**, 93–105 (2007).