Research article

Effect of C-to-T transition at CpG sites on tumor suppressor genes in tumor

development in cattle evaluated by somatic mutation analysis in enzootic bovine

leukosis

Running Title: Mode of EBL development based on somatic mutation

Asami Nishimori,[a#] Kiyohiko Andoh,[a] Yuichi Matsuura,[a] Tomohiro Okagawa,[b] Satoru

Konnai[b,c]

[a] National Institute of Animal Health, National Agriculture and Food Research

Organization, Tsukuba, Ibaraki, Japan

[b] Department of Advanced Pharmaceutics, Faculty of Veterinary Medicine, Hokkaido

University, Sapporo, Hokkaido, Japan.

[c] Department of Disease Control, Faculty of Veterinary Medicine, Hokkaido University,

Sapporo, Hokkaido, Japan

Address correspondence to Asami Nishimori, nishimoria466@affrc.go.jp

Abstract: 223 words

Text excluding the references and figure legends: 4,910 words

Abstract

Oncogenic transformation of normal cells is caused by mutations and

chromosomal abnormalities in cancer-related genes. Enzootic bovine leukosis (EBL) is

a malignant B-cell lymphoma caused by bovine leukemia virus (BLV) infection in cattle.

Although a small fraction of BLV-infected cattle develops EBL after a long latent period,

the mechanisms for oncogenesis in EBL cattle remain largely unknown. In this study,

we analyzed the types and patterns of somatic mutations in cancer cells from 36 EBL

cases, targeting 21 cancer-related genes. Various somatic mutations were identified in 8

genes, *TP53*, *NOTCH1*, *KMT2D*, *CREBBP*, *KRAS*, *PTEN*, *CARD11*, and *MYD88*. In

addition, *TP53* gene was found to be mutated in 69.4% of EBL cases, with most being

biallelic mutations. In some cases, associations were observed between the ages at

which cattle had developed EBL and somatic mutation patterns; young onset of EBL

possibly occurs due to congenital mutations, high impact mutations affecting protein

translation, and biallelic mutations. Furthermore, nucleotide substitution patterns

indicated that cytosine at CpG sites tended to be converted to thymine in many EBL

cases, which was considered to be the result of spontaneous deamination of

5-methylctosine. These results demonstrate how somatic mutations have occurred in

cancer cells leading to EBL development, thereby explaining its pathogenic mechanism.

These findings will contribute to a better understanding and future elucidation of

disease progression in BLV infection.

Importance

Enzootic bovine leukosis (EBL) is a malignant and lethal disease in cattle.

Currently, there are no effective vaccines or therapeutic methods against bovine

leukemia virus (BLV) infection, resulting in severe economic losses in livestock

industry. This study provides a renewed hypothesis to explain the general mechanisms

of EBL onset by combining the previous finding that several integration sites of BLV

provirus can affect the increase in survival and proliferation of infected cells. We

demonstrate that two additional random events are necessary for oncogenic

transformation in infected cell clones, elucidating the reason why only few infected

cattle develop EBL. Further exploration of somatic mutation and BLV integration sites

could support this hypothesis more firmly, potentially contributing to the development

of novel control methods for EBL onset.

Keywords: somatic mutation, enzootic bovine leukosis, *TP53*, mutation signature

Introduction

Cancer is a disease of cell proliferation caused by chromosomal aberrations and somatic mutations in genes related to proliferation, the cell cycle, and DNA damage repair. Currently, at least 100 oncogenes and at least 30 tumor suppressor genes have been identified [1]. Oncogenes are considered to enhance cell proliferation by gain-of-function mutations, whereas tumor suppressor genes such as *RB1* and *TP53* cause loss of control over normal cell growth and differentiation by loss-of-function mutations or deletion [1, 2]. Generally, both copies of tumor suppressor genes on homologous chromosomes need to be inactivated to achieve oncogenic transformation, which is a concept known as Knudson's "two-hit" theory [3]. The first "hit" is not necessarily a somatic mutation; Li-Fraumeni syndrome is a hereditary and familial cancer caused by a germline mutation in *TP53* gene. Patients with Li-Fraumeni syndrome have only one active *TP53* gene, leading to the development of multiple types of tumors at a young age due to a somatic mutation as the second "hit" in the normal allele [4, 5]. Thus, analyzing somatic mutations in cancer cells is important for understanding the history and process of oncogenesis.

Enzootic bovine leukosis (EBL) is a malignant B-cell lymphoma in cattle caused by bovine leukemia virus (BLV) infection, which belongs to the family *Retroviridae*, genus *Deltaretrovirus* [6]. BLV is closely related to human T-cell leukemia virus type I (HTLV-1) that infects human T cells and causes adult T-cell leukemia/lymphoma (ATL). BLV and HTLV-1 exhibit similar life cycles and disease progression. Their viral genomes are integrated into the host genome as a provirus, and only a small fraction (1–5%) of infected individuals develops lymphoma after a long latent period [6]. Experimental infection with BLV in cattle has shown two types of

viral propagation during the infection *in vivo*. BLV actively replicates and propagates by

infecting new lymphocytes in the early phase of infection (infectious cycle), but it

expresses few viral antigens after the development of host acquired immunity and

propagates via clonal expansion of infected cells in the chronic stage (mitotic cycle) [7].

Because infecting new target cells results in generating a new clone of infected cells

with a different provirus integration site, the infectious cycle is important to increase the

clonal variation of infected cells *in vivo*. A recent study reported that clonal variation

and provirus integration sites were involved in the mechanisms of EBL and ATL

development [8]. Rosewick et al. reported that BLV/HTLV-1 proviruses in tumor

genomes of EBL animals or ATL patients were preferentially integrated near cancer

driver genes, and that the proviruses could disturb mRNA expression of host genes

located in their vicinity by gene interruption, depending on viral 5′ long terminal repeat

poly-(A) signal or expression of 3′ antisense RNA-dependent chimeric transcript.

However, the provirus integration sites did not fully explain the reasons why only a

small fraction of infected individuals developed EBL or ATL, because the same pattern

of integration sites was observed in animals at a polyclonal non-malignant stage [8].

The authors discussed in their report that provirus integration sites would contribute to

initial selection of viral-infected cell clones at an early aleukemic stage, and the

accumulation of further somatic mutation in the host genome was needed for the onset

of lymphoma. Therefore, mutation analysis of cancer-related genes will enable the more

detailed mechanisms of EBL development to be clarified.

To date, many studies of the mutation analyses of bovine *TP53* gene have been

reported using clinical samples or cell lines derived from EBL cattle [9–13]. However,

unlike the research on ATL [14], there is no information on the mutation status of other

cancer-related genes involved in EBL development. Moreover, although the analysis of

mutation signatures, which is an *in silico* method to evaluate the mutation process of

oncogenesis from the pattern of single nucleotide variants (SNVs), is used in the field of

human cancer research [15, 16], neither mutational patterns nor evaluation of allelic

variation based on the "two-hit" theory has been fully elucidated in cancer cells of EBL

cattle. What is more, recently, EBL onset in young cattle (less than 3 years old) has been

frequently reported in Japan, despite the common ages of EBL development for

BLV-infected cattle being approximately 5–8 years [17, 18]. Examining the relation

between cattle ages when EBL developed and patterns of somatic mutations in cancer

cells will generate much interest. Therefore, in this study, a somatic mutation analysis

targeting the selected 21 genes was performed to clarify genomic changes related to

oncogenesis in EBL cattle. The pattern of identified somatic mutations was then

evaluated based on the ages of EBL development, and the mutation process of

oncogenesis was evaluated using mutation signature analysis.

Results

Cancer-related genes in which somatic mutations were detected in EBL cases

According to the procedures shown in Fig. 1, somatic mutation analyses were

conducted for 36 EBL cases. The results showed that somatic mutations were detected

in coding sequence (CDS) regions of only 8 of 21 targeted cancer-related genes, *TP53*,

NOTCH1, *KMT2D*, *CREBBP*, *KRAS*, *PTEN*, *CARD11*, and *MYD88* (Tables 1, 2 and S3;

summarized in Fig. 2A). There were identical mutations in different EBL cattle in *TP53*

(p.Arg167His, p.Arg241Trp, p.Arg242Trp p.Arg242Gln, and p.Arg275His) and

NOTCH1 (p.Cys468fs) genes, implying the presence of mutational hotspots in these

genes (Table 2). *TP53* was the most frequently mutated gene, with 69.4% (25/36) of

EBL cases possessing at least one *TP53* mutation in the cancer cells (Fig. 2B).

Moreover, 88.0% (22/25) of *TP53* mutations were detected as biallelic mutations,

namely multiple mutations or mutations associated with loss of heterozygosity (LOH)

(Fig. 2A). Although the mutation frequencies of *NOTCH1* and *CREBBP* genes were not

high, they were the only two genes in which biallelic mutations were observed in EBL

cases with wildtype *TP53* gene, such as cases #5, #29, and #34 (Fig. 2A,B). It should be

noted that *TP53* mutations with LOH in several cases were validated by comparisons

between variant allele frequencies (VAFs) of blood and tumor samples for germline

mutations located on chromosome 19 (Table S4).

Age-associated characteristics of somatic mutations

One remarkable point in the associations between the ages at which cattle had

developed EBL and somatic mutation patterns is evident in cases #29 and #31, which

involved young cattle less than 36 months old (Table S1). Both normal and cancer cells

in these cases possessed a frameshift deletion in *NOTCH1* gene that was not registered

as a known Variant ID (Fig. 2A and Table 2), suggesting a possible congenital variation.

The possibility of significant contamination of cancer cells in blood samples could be

ruled out because, in cases #29 and #31, there were different somatic mutations in

CREBBP and *TP53* genes, respectively (Fig. 2A), and they were not detected in blood

samples from these cases (Table S3). Interestingly, case #34 showed the same frameshift

mutation in *NOTCH1* gene as a somatic mutation (Table 2). As for considerably older

cattle, the mutation frequency in *TP53* gene was clearly decreased in EBL cases more

than 96 months old (Fig. 2A). However, there was no significant difference between

158 ages of EBL cases with *TP53* wildtype gene and those with a mutated gene, because

159 several young EBL cattle were also free of *TP53* mutation (Fig. 2C). However, ages at

160 development of EBL were significantly younger in the cases with high-impact

161 mutations, which exert a large influence on protein translation, such as stop-gained and

162 frameshift mutation, or with biallelic mutations (Fig. 2D,E).

164 **Analysis of the mutation pattern in SNVs**

165 In all somatic mutations, an advanced analysis focusing on the SNVs was

166 performed. Given the small number of SNVs per EBL case in this study, the 36 EBL

167 cases were grouped by their ages into two categories: an old group over 36 months old

168 ($\geq$ 36-m-old) and a young group less than 36 months old (< 36-m-old). Mutation

169 patterns were then compared among those groups and all cases. The frequency of base

170 substitution from the reference allele to the alternative allele among detected SNVs

171 indicated a large number of mutations involving G-to-A and its complementary

172 sequence C-to-T in EBL cattle (Fig. 3A). That tendency was particularly more

173 pronounced in the $\geq$ 36-m-old group (G-to-A, 0.38; C-to-T, 0.44) than in the < 36-m-old

174 group (G-to-A, 0.33; C-to-T, 0.25).

176 **Evaluation of single base substitution (SBS) signatures**

177 In the SBS signature, SNVs can be classified into 96 different contexts by

178 considering 6 types of base substitutions (C>A, C>G, C>T, T>A, T>C and T>G,

179 including complementary strands), as well as 4 types of base immediately 5´ and 4 types

180 of base immediately 3´. Mutation spectra were compared based on SBS signatures

181 among all cases, the $\geq$ 36-m-old group, and the < 36-m-old group (Fig. 3B). As a result,

3 mutation contexts, A[C>T]G, C[C>T]G and G[C>T]G, were found to predominate in

all cases and the ≥ 36-m-old group, which indicated that cytosine with guanine on the 3´

side, i.e., cytosine at CpG sites, tended to be converted to thymine. Interestingly, hotspot

mutations in *TP53* gene, observed across multiple EBL cases and/or reported in

previous studies, corresponded to CpG sites with only one exception (p.Phe229Leu,

T[C>A]A) (Table 2). In contrast, of 12 SNVs in *TP53* gene first reported in this study,

eleven were inconsistent with CpG sites. The mutation spectrum in the < 36-m-old

group showed that the context C[T>C]G, in addition to C[C>T]G and G[C>T]G, was

slightly predominant (Fig. 3B), although the small number of SNVs in the young group

should be considered.

At this point, the mutation spectrum obtained from SNVs in the ≥ 36-m-old

group was defined as Signature A, and that obtained from SNVs in the < 36-m-old

group was defined as Signature B. The similarities of mutation context patterns were

compared between Signature A or B and the reference signatures, classically known

COSMIC signatures 1–21 (Fig. S2A) [15, 16]. Both Signature A and B were similar to

COSMIC signatures 1 and 6, and Signature A in particular showed the highest cosine

similarity with COSMIC signature 1 (Fig. 3C). We then attempted to evaluate the

mutation process of oncogenesis by mutation spectra for all cases, the ≥36-m-old group,

and the <36-m-old group using MutSignatures R package [19]. The COSMIC signatures

used for the evaluation were selected based on validated evidence and their likelihood

of involvement in EBL development (Fig. S2A); for example, COSMIC signature 4

(tobacco smoking) and signature 11 (temozolomide chemotherapy) were excluded

because they were less likely to be associated with cattle diseases. As a result, the main

components of the mutation process in EBL cases were COSMIC signatures 1, 6, 7, and

18 (Fig. S2B), and above all, COSMIC signature 1 accounted for the majority in all

cases and the $\geq$ 36-m-old group (Fig. 3D). However, not only signature 1, but also

COSMIC signature 6 seemed to be largely involved in EBL development in the

<36-m-old group. According to the mutation profiles in the COSMIC database, the

mutational processes of COSMIC signatures 1 and 6 are "spontaneous deamination of

5-methylcytosine" and "mismatch repair (MMR) deficiency", respectively (Fig. S2A).

Discussion

In this study, somatic mutations of 21 cancer-related genes were analyzed in

cancer cells from EBL cattle, clearly distinguishing between somatic and germline

mutations. The results uncovered several key findings: 1) somatic mutation based on the

"two-hit" theory, such as multiple mutations on the same genes or LOH-associated

mutations, were frequently observed at EBL onset (Fig. 2A); 2) many EBL cases relied

on somatic mutations in *TP53* gene for oncogenesis (Fig. 2B); and 3) CpG sites were

the main targets of these somatic mutations (Fig. 3B and Table 2). It is well-known that

CpG sites are major targets for methylation induced by cell division, and that

5-methylcytosine is prone to deamination to thymine, leading to a G·T mismatch in

double-stranded DNA [20–23]. Thus, the C-to-T mutations at CpG sites are related to

aging of the animal as a mutation process, which is supported by the observation that

the number of mutations correlates with patients' ages in COSMIC signature 1 [16].

COSMIC signature 1 is hence considered to be a cell division clock. Besides, an

increase in the rate of LOH can result from mitotic recombination [24]. Consequently,

the present study suggests that physiological and accidental mutations or chromosomal

abnormalities caused by repeated cell division in infected cells for a long period are the

factors contributing to EBL onset, particularly in cattle over 36 months old. However, it

is important to note that the ages of cattle when EBL developed may not correctly

reflect the period of disease progression from viral infection to lymphoma onset, since

the time of BLV infection varied among EBL cases.

234   Regarding the associations between EBL onset in young cattle and somatic

mutation patterns, the presence of congenital variations, high-impact mutations

significantly affecting protein translation, and biallelic mutations primarily in *TP53*,

NOTCH1, and *CREBBP* genes may elevate the probability of EBL development in a

short period (Fig. 2A,D,E). *TP53* gene is a representative tumor suppressor gene coding

p53, which acts as a transcription factor upregulated in response to DNA damage,

activating downstream genes involved in cell-cycle control, apoptosis, and DNA repair

[25]. NOTCH1 is a receptor protein expressed on the cell membrane that regulates cell

differentiation and self-renewal by interaction with ligands on adjacent cells [26].

CREBBP is a lysine acetyltransferase modulating chromatin accessibility through

histone acetylation [27]. Although *NOTCH1* and *CREBBP* genes can act as both

oncogenes and tumor suppressor genes, they probably functioned as tumor suppressor

genes at EBL onset, because most of their mutations were inactivating forms like

stop-gained and frameshift mutations (Fig. 2A). For these reasons, somatic mutations in

TP53, *NOTCH1*, and *CREBBP* genes would result in accelerating the accumulation of

secondary mutations due to deficiencies in DNA damage repair and cell division

regulation.

251   Further, the evaluation of the mutation process based on the known signatures

showed that COSMIC signature 6, not only signature 1, was mainly involved in

oncogenesis in the < 36-m-old group (Fig. 3D). It should be considered that the number

of SNVs from the < 36-m-old group was quite small, and hence the accuracy of the

estimation by mutation signature analysis might be insufficient; therefore, further

analyses are needed to confirm if COSMIC signature 6 is involved in young onset of

EBL. Nevertheless, the involvement of COSMIC signature 6, representing MMR

deficiency, is worth considering as a possible factor causing young EBL onset, because

the evaluation results for SNVs from all cases ($n = 44$) included COSMIC signature 6,

whereas that from the $\geq$ 36-m-old group ($n = 32$) did not (Fig. 3D).

Although the mechanism of EBL development had not been fully elucidated,

Rosewick et al. reported that cancer cells from animals developing EBL had proviruses

integrated near cancer driver genes, and that these proviruses could increase survival

and proliferation in infected cells by disturbing host gene expression [8]. They

concluded in their report that it will result in extending the half-life of infected cells and

promote the accumulation of further secondary mutations in the genome of oncogenic

candidate clones. The present results strongly support this model for EBL development

by clarifying the detail of how secondary somatic mutations accumulated in cancer cells.

Now, we can propose a renewed hypothesis to explain the general mechanisms of EBL

onset; three events need to be completed for tumor development in BLV-infected cattle

(Fig. 4). The first event is provirus integration near cancer driver genes, as shown in the

previous study [8], which is randomly decided during the infectious cycle in the early

phase of BLV infection. The second event is an accidental single mutation (the first

"hit") at CpG sites in cancer-related genes, particularly *TP53*, *NOTCH1*, and *CREBBP*

genes, caused by cell division repeated over long periods. The third event is an

additional mutation on another allele or chromosomal aberration like LOH as the

second "hit". The oncogenic transformation of BLV-infected cells requires the

occurrence of all three random events in the same cell clone, and thus the rate of EBL

onset in infected cattle becomes very small.

Based on our hypothesis, we can discuss host or viral factors previously

reported to be involved in EBL development. The associations between EBL onset and

types of cattle major histocompatibility complex BoLA-DRB3 allele [28], viral

genotypes [29], or provirus loads during mitotic cycle [30] have been reported in

previous studies. At this point, we consider that these factors are related to expanding

the diversity of provirus integration sites. The diversity of provirus integration sites,

namely clonal variation of infected cells, is increased during the infectious cycle when

BLV infects new target cells, followed by a large decrease in clone numbers due to

negative selection by the host immune response [7]. In addition, the provirus loads in

the chronic phase of BLV infection tend to reflect the number of infected cell clones in

the blood [7]. Therefore, these host or viral factors may contribute to a weak immune

response against BLV or efficient viral replication in an early phase, resulting in an

increase in initial candidate clones with the potential for transformation.

In the research field of ATL caused by HTLV-1 infection, somatic mutation

analysis by whole exome sequencing (WES) was conducted using clinical samples from

ATL patients. Similar to the present results, age-related C-to-T transitions at CpG sites

were predominantly observed on mutation signature analysis [14]. In contrast, the

mutation frequency in *TP53* genes of ATL samples was around 20% on WES analysis,

suggesting that the oncogenic process in ATL development was less dependent on *TP53*

mutations than in EBL cases. In addition, the rate of detecting somatic mutations was

reported as 2.3 mutation/Mbp/samples according to the WES in ATL patients. Since the

total length of the CDS region in our target resequencing is approximately 73 kbp

(Table 1), focusing on selected cancer-related genes worked well for efficiently

detecting somatic mutations in the present study. As a limitation of the present study,

which is also relevant to WES analysis, a portion of EBL cases (6/36) did not show any

somatic mutation in the targeted genes (Fig. 2A), probably because the targeted-gene

panel was selected based on information from various human B-cell lymphomas, rather

than actual data from WES analysis in cattle. Moreover, the experimental approach

cannot identify somatic mutations in promoter regions or epigenetic inactivation. For

these reasons, further efforts will be required to improve the methods or study

approaches for mutation analysis in bovine B cells.

In this study, several characteristic EBL cases were identified. As for cases #29

and #31, which showed a possible congenital variation in *NOTCH1* genes, they both

harbored biallelic and high-impact somatic mutations on *CREBBP* or *TP53* genes,

making it unclear whether that congenital variation directly caused the young onset of

EBL (Fig. 2A). Moreover, these cases were inconsistent with the characteristics of

hereditary cancer, such as Li-Fraumeni syndrome, because there was no second "hit" on

NOTCH1 genes in their genome. However, it is known that *NOTCH1*, as well as

CREBBP, can be affected by a single mutation due to haploinsufficiency [26, 27, 31].

Based on the observation that case #34 also exhibited the same frameshift mutation as a

somatic mutation, the identified congenital variation is likely a functional driver

mutation that plays a role in oncogenesis. Unfortunately, there is no additional

information on cases #29 and #31 regarding their history, and thus, we cannot discuss

their blood relationship. Let us consider two possibilities for the congenital variant; it

may be a variant inherited from the parents, or it may have incidentally occurred during

the period from fertilized egg to early embryonic development.

For the three cases that did not exhibit biallelic mutations in *TP53* gene, only a

single *TP53* mutation was observed in each case (#7, #18, and #28) (Fig. 2A). The

somatic mutations from cases #7 and #18, p.Arg167His (R167H) and p.Arg241Trp

(R241W), respectively, were mutational hotspots in *TP53* gene, which were also

detected in other EBL cases reported in previous studies (Table 2) [9–12]. According to

the previous study examining the functions of mutated *TP53*, R241W was not only a

loss-of-function mutation impairing the transcriptional activity of p53 protein, but it

also acted as a dominant negative variant inhibiting wildtype p53 in the presence of

both variants [12]. Conversely, R167H is known as an orthologous mutation of R175H

in human *TP53* gene, which is one of the major hotspot mutations observed in various

human cancers, showing a dominant negative effect [9, 32–34]. Altogether, it is

suggested that the introduction of mutations with dominant negative effects as the first

"hit" resulted in oncogenic transformation by inhibiting the normal allele in cases #7

and #18. However, all other cases with R167H or R241W mutations (cases #2, #12, #13,

#16, and #19) showed the biallelic *TP53* mutations (Table S3). This suggests that the

dominant negative effects of these mutations may not completely suppress the function

of the normal allele, and that biallelic inactivity is more advantageous for oncogenesis.

Lastly, regarding the association of age with EBL development, it is interesting

that EBL in cattle more than 96 months old showed less dependence on *TP53* mutations

(Fig. 2A). This finding may be linked to immunosenescence [35]; the decline in the

functions of the immune response due to aging causes a weakened elimination of

precancerous abnormal cells, creating a tumor-prone condition. Considering that

mutation frequencies of *TP53* gene in diffuse large B-cell lymphoma, one of the

nonviral and sporadic human cancers, range from 17.6% to 23.2% [36], the

development of EBL in very old cattle may resemble that of spontaneous tumors rather

than virus-associated tumors.

In conclusion, this study provides novel insights into the pathogenic

mechanism of somatic mutations in cancer cells for EBL development. Furthermore, the

present results suggest the possibility that the mutation patterns in young cattle

developing EBL are different from those in EBL cattle at common ages. To strengthen

the hypothesis inferred from our results, it is necessary to accumulate data using a larger

number of clinical samples in the future.

Materials and methods

Study design

Fig. 1 summarizes the study strategy. Two samples each were collected from 36

cattle with EBL; one was a blood sample mainly containing normal cells, and the other

was a tumor sample mainly containing cancer cells. Because it is known that several

EBL cases do not show visible lymphocytosis when a tumor develops and thus, cancer

cells are hardly detected in the peripheral blood [17, 37], such EBL cases not showing

leukemic signs were selectively used. DNA was extracted from blood and tumor

samples and used for PCR amplification of 21 targeted cancer-related genes, followed

by amplicon sequencing and variant calling using next-generation sequencing

technology. Two variant lists of blood and tumor samples from the same EBL case were

compared, and the variants that existed in only tumor samples were identified as

somatic mutations. All identified somatic mutations were used for mutation pattern

analysis.

Clinical samples from EBL cattle

The information on EBL cases used in this study are listed in Table S1. All

cases were diagnosed as EBL in a domestic livestock hygiene service center and meat

inspection center in Japan, and they were kindly provided for research use in the

previous studies [37, 38]. To select EBL cases not showing leukemic signs, it was

confirmed that lymphocyte counts or clonality values measured by the

RAISING-CLOVA method [38] were low in the blood (Table S1). In cases #1–32,

peripheral blood mononuclear cells (PBMCs) were isolated from blood samples using

Percoll (GE Healthcare Life Sciences, Little Chalfont, UK) density gradient

centrifugation for DNA extraction. In cases #33–36, DNA was extracted directly from

whole blood samples. Cancer cells were isolated from tumor samples, as described

previously [37]; briefly, tumor samples were cut into small pieces, suspended in

phosphate-buffered saline, and passed through a 40-μm cell strainer (Corning Inc.,

Corning, NY, USA) for removal of cell debris. For DNA extraction, the Dneasy Blood

and Tissue kit (QIAGEN, Hilden, Germany), Wizard Genomic DNA purification kits

(Promega, Madison, WI, USA), or Quick-DNA Miniprep Kits (Zymo Research, Irvine,

CA, USA) were used.

Amplicon sequencing of cancer-related genes

To determine the targeted cancer-related genes, the COSMIC (Catalogue Of

Somatic　　　Mutations　　　In　　　Cancer)　　　[39]　　　Cancer　　　Browser　　　database

(https://cancer.sanger.ac.uk/cosmic/browse/tissue) was used, and 21 genes that showed

high mutation frequencies in various human B-cell lymphomas were selected (Table 1).

Primer sets that included all CDS of the 21 targeted genes were designed for long PCR,

and 3–4 sets each were mixed for amplification as multiplex PCR (multiplex set A–U,

Table S2). The targeted genes in the DNA sample from EBL cattle were amplified using

the multiplex primers in a reaction mixture containing 12.5 μl of KOD One PCR Master

Mix (TOYOBO, Osaka, Japan), 3.0 μl of primer mix (each 2.5 μM), 0.5 μl of dimethyl

sulfoxide (FUJIFILM Wako, Osaka, Japan), 0.2–1.0 μl of DNA template, and extra

double-distilled water up to 25 μl. The PCR conditions were as follows: a two-step

procedure consisting of 10 sec at 98 °C and 2 min at 68 °C for 25–35 cycles decided

depending on the concentration of the template DNA. Following DNA purification

using the QIAquick PCR Purification Kit (QIAGEN), the amplicons were pooled and

submitted to Macrogen Japan Co., Ltd. (https://www.macrogen-japan.co.jp/) for the

DNA-seq analysis. Briefly, a sequence library was prepared using the TruSeq Nano

DNA Kit (Illumina, San Diego, CA, USA), and next-generation sequencing analysis

was conducted using a NovaSeq 6000 (150-bp pair-end reads, approximately 5.7-Gbp

total read bases per sample).

Data analysis

The identification of somatic mutations was performed using "Variant Calling

Pipeline using GATK4" in accordance with online instructions

(https://github.com/gencorefacility/variant-calling-pipeline-gatk4) with some

modifications. The FASTQ data obtained in DNA-seq of blood and tumor samples from

36 EBL cattle are deposited in the DDBJ database as submission no. DRR531829–

DRR531900. The reads were trimmed using Trimmomatic version 0.39 and mapped

against the reference genome (*Bos taurus*: ARS-UCD1.2 also known as

GCF_002263795.1) using the MEM algorithm in BWA version 0.7.17. The output

BAM files were indexed using samtools version 1.6. The variant calling was then

conducted for each BAM file using HaplotypeCaller in GATK version 4.3.0.0, and base

quality scores were adjusted by GATK's BQSR algorithm. The variant filtering was

performed using GATK's VariantFiltration with the following criteria: $QD < 2$, $FS > 60$,

$MQ < 40$, $QUAL < 30$, $SOR > 4$, $MQRankSum < -12.5$, $ReadPosRankSum < -8$ for

SNVs, and $QD < 2$, $FS > 200$, $QUAL < 30$, $SOR > 10$, $ReadPosRankSum < -20$ for

insertions or deletions (INDELs). Gene annotations of called variants were added in

VCF files using SnpEff version 5.1 with the reference genome, ARS-UCD1.2.99. The

variants called only in tumor samples were sorted using bcftools version 1.9.

Of the somatic mutations called by the above procedure, the variants whose

read depth was less than 20 or which agreed with known Variant ID in the Ensembl

database were excluded from the following analysis. When somatic mutations were also

called in blood samples, both VAFs of blood and tumor samples were compared

statistically, and it was found that VAF was significantly higher in tumor samples than

in blood samples. The definition of chromosome zygosity at the detected somatic

mutations was as follows: VAF > 0.7 was considered homozygosity, VAF 0.3–0.7

(SNVs) or VAF 0.2–0.7 (INDELs) was considered heterozygosity, and VAF < 0.3

(SNVs) or VAF < 0.2 (INDELs) was considered a minor mutation. Fig. S1 indicates the

interpretation and classification of the patterns of somatic mutations. In comparison

with a single heterozygous mutation, multiple heterozygous mutations on a certain

target gene were interpreted as meaning that one mutation existed on one allele, and a

second mutation existed on a different allele of a homologous chromosome. In addition,

a single homozygous mutation was interpreted as a mutation with LOH that resulted in

loss of one copy of the target gene by chromosomal abnormalities such as

recombination and deficiency. All somatic mutations identified in this study were

visually checked in the BAM files using IGV software [40].

Mutation signature analysis

R package MutSignatures version 2.1.5 [19] with R version 4.3.0 was used for

mutation signature analysis of EBL cattle using VCF files generated above in

accordance   with   online   instructions

(https://cran.r-project.org/web/packages/mutSignatures/vignettes/get_sarted_with_mutS

ignatures.html). Bovine whole genome sequences, BSgenome.Btaurus.UCSC.bosTau9,

were used as a reference genome. The reference signature datasets listed on the

COSMIC website [39] were obtained by the getCosmicSignatures command.

Statistical analysis

Fisher's exact test was used to evaluate differences between VAFs of blood and

tumor samples when somatic mutations were called in both samples. Associations

between cattle ages when EBL developed and patterns of somatic mutations in cancer

cells were explored using Mann-Whitney's U test. A p-value less than 0.05 was defined

as significant.

Appendix

The FASTQ data in this study were deposited in the DDBJ database as submission no. DRR531829–DRR531900. All supplemental information is shown in the supplemental material.

Acknowledgements

The authors are grateful to the staff at the livestock hygiene service centers and meat hygiene inspection centers in Hokkaido, Iwate, Ibaraki, and Hyogo prefectures in Japan for kindly providing clinical EBL samples. Xiao Bo provided excellent technical support.

This study was supported by JSPS KAKENHI Grant-in-Aid for Early-Career Scientists (19K16008 and 23K14101) (A.N.). The funder had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

A.N. designed the research. A.N., K.A., and T.O. performed data collection and analysis. K.A., Y.M., T.O., and S.K. provided intellectual input, laboratory materials, and/or analytic tools. A.N. drafted the manuscript. All authors read and approved the manuscript and contributed to manuscript revisions.

Conflict of interest statement

The authors declare no conflict of interest.

References

References

1.   Soto AM, Sonnenschein C. 2004. The somatic mutation theory of cancer: growing

problems   with   the   paradigm?   Bioessays   26(10):1097–1107.   doi:

10.1002/bies.20087.

2.   Kontomanolis EN, Koutras A, Syllaios A, Schizas D, Mastoraki A, Garmpis N,

Diakosavvas M, Angelou K, Tsatsaris G, Pagkalos A, Ntounis T, Fasoulakis Z.

2020. Role of Oncogenes and Tumor-suppressor Genes in Carcinogenesis: A

Review. Anticancer Res. 40(11):6009–6015. doi: 10.21873/anticanres.14622.

3.   Knudson AG Jr. 1971. Mutation and cancer: statistical study of retinoblastoma.

Proc Natl Acad Sci U S A 68(4):820–823. doi: 10.1073/pnas.68.4.820.

4.   Srivastava S, Zou Z, Pirollo K, Blattner W, Chang EH. 1990. Germ-line

transmission of a mutated p53 gene in a cancer-prone family with Li–Fraumeni

syndrome. Nature 348(6303):747–749. doi: 10.1038/348747a0

5.   Malkin D. 1993. p53 and the Li-Fraumeni syndrome. Cancer Genet Cytogenet

66(2):83–92. doi: 10.1016/0165-4608(93)90233-c

6.   Gillet N, Florins A, Boxus M, Burteau C, Nigro A, Vandermeers F, Balon H,

Bouzar AB, Defoiche J, Burny A, Reichert M, Kettmann R, Willems L. 2007.

Mechanisms of leukemogenesis induced by bovine leukemia virus: prospects for

novel   anti-retroviral   therapies   in   human.   Retrovirology   4:18.   doi:

10.1186/1742-4690-4-18

7.   Gillet NA, Gutiérrez G, Rodriguez SM, de Brogniez A, Renotte N, Alvarez I, Trono

K, Willems L. 2013. Massive depletion of bovine leukemia virus proviral clones

located in genomic transcriptionally active sites during primary infection. PLoS

Pathog 9(10):e1003687. doi: 10.1371/journal.ppat.1003687.

8.   Rosewick N, Durkin K, Artesi M, Marcais A, Hahaut V, Griebel P, Arsic N,

Avettand-Fenoel V, Burny A, Charlier C, Hermine O, Georges M, Van den Broeke

A. 2017. Cis-perturbation of cancer drivers by the HTLV-1/BLV proviruses is an

early determinant of leukemogenesis. Nature Communications 8:15264 doi:

10.1038/ncomms15264.

9.   Konishi M, Kameyama K, Yamamoto T. 2021. Mutations in the tumor suppressor

gene p53 in cattle are associated with enzootic bovine leukosis. Vet Microbiol

263:109269. doi: 10.1016/j.vetmic.2021.109269.

10. Dequiedt F, Kettmann R, Burny A, Willems L. 1995. Mutations in the p53

Tumor-Suppressor Gene Are Frequently Associated with Bovine Leukemia

Virus-Induced Leukemogenesis in Cattle but Not in Sheep. Virology 209 (2):676–

683. doi: 10.1006/viro.1995.1303.

11. Ishiguro N, Furuoka H, Matsui T, Horiuchi M, Shinagawa M, Asahina M, Okada K.

1997. P53 Mutation As a Potential Cellular Factor for Tumor Development in

Enzootic Bovine Leukosis. Vet Immunol Immunopathol 55(4):351–358. doi:

10.1016/s0165-2427(96)05721-2.

12. Tajima S, Zhuang WZ, Kato MV., Okada K, Ikawa Y, Aida Y. 1998. Function and

conformation of wild-type p53 protein are influenced by mutations in bovine

leukemia virus-induced B-cell lymphosarcoma. Virology 243 (1):235–246.

13. Komori H, Ishiguro N, Horiuchi M, Shinagawa M, Aida Y. 1996. Predominant p53

mutations in enzootic bovine leukemic cell lines. Vet Immunol Immunopathol 52

(1-2):53–63. doi: 10.1016/0165-2427(95)05538-x.

14. Kataoka K, Nagata Y, Kitanaka A, Shiraishi Y, Shimamura T, Yasunaga J, Totoki Y,

Chiba K, Sato-Otsubo A, Nagae G, Ishii R, Muto S, Kotani S, Watatani Y, Takeda J,

Sanada M, Tanaka H, Suzuki H, Sato Y, Shiozawa Y, Yoshizato T, Yoshida K, Makishima H, Iwanaga M, Ma G, Nosaka K, Hishizawa M, Itonaga H, Imaizumi Y, Munakata W, Ogasawara H, Sato T, Sasai K, Muramoto K, Penova M, Kawaguchi T, Nakamura H, Hama N, Shide K, Kubuki Y, Hidaka T, Kameda T, Nakamaki T, Ishiyama K, Miyawaki S, Yoon SS, Tobinai K, Miyazaki Y, Takaori-Kondo A, Matsuda F, Takeuchi K, Nureki O, Aburatani H, Watanabe T, Shibata T, Matsuoka M, Miyano S, Shimoda K, Ogawa S. 2015. Integrated molecular analysis of adult T cell leukemia/lymphoma. Nat Genet 47(11) :1304–1315.

15. Nik-Zainal S, Alexandrov LB, Wedge DC, Van Loo P, Greenman CD., Raine K, Jones D, Hinton J, Marshall J, Stebbings LA, Menzies A, Martin S, Leung K, Chen L, Leroy C, Ramakrishna M, Rance R, Lau KW, Mudie LJ, Varela I, McBride DJ, Bignell GR, Cooke SL, Shlien A, Gamble J, Whitmore I, Maddison M, Tarpey PS, Davies HR, Papaemmanuil E, Stephens PJ, McLaren S, Butler AP, Teague JW, Jönsson G, Garber JE, Silver D, Miron P, Fatima A, Boyault S, Langerød A, Tutt A, Martens JWM, Aparicio SAJR, Borg Å, Salomon AV, Thomas G, Børresen-Dale AL, Richardson AL, Neuberger MS, Futreal PA, Campbell PJ, Stratton MR. 2012. Mutational Processes Molding the Genomes of 21 Breast Cancers. Cell 149(5):979–993. doi: 10.1016/j.cell.2012.04.024.

16. Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, Bignell GR, Bolli N, Borg A, Børresen-Dale AL, Boyault S, Burkhardt B, Butler AP, Caldas C, Davies HR, Desmedt C, Eils R, Eyjörd JE, Foekens JA, Greaves M, Hosoda F, Hutter B, Ilicic T, Imbeaud S, Imielinski M, Jäger N, Jones DTW, Jones D, Knappskog S, Kool M, Lakhani SR, López-Otín C, Martin S, Munshi NC, Nakamura H, Northcott PA, Pajic M, Papaemmanuil E, Paradiso A, Pearson JV,

Puente XS, Raine K, Ramakrishna M, Richardson AL, Richter J, Rosenstiel P,

Schlesner M, Schumacher TN, Span PN, Teague JW, Totoki Y, Tutt ANJ,

Valdés-Mas R, van Buuren'MM, van 't Veer L, Vincent-Salomon A, Waddell N,

Yates LR, Australian Pancreatic Cancer Genome Initiative, ICGC Breast Cancer

Consortium, ICGC MMML-Seq Consortium, ICGC PedBrain, Zucman-Rossi J,

Futreal PA, McDermott U, Lichter P, Meyerson M, Grimmond SM, Siebert R,

Campo E, Shibata T, Pfister SM, Campbell PJ, Stratton MR. 2013. Signatures of

mutational processes in human cancer. Nature 500(7463):415–421. doi:

10.1038/nature12477.

17. Nishimori A, Konnai S, Okagawa T, Maekawa N, Goto S, Ikebuchi R, Nakahara A,

Chiba Y, Ikeda M, Murata S, Ohashi K. 2017. Identification of an Atypical

Enzootic Bovine Leukosis in Japan by Using a Novel Classification of Bovine

Leukemia Based on Immunophenotypic Analysis. Clin Vaccine Immunol

24(9):e00067-17. doi: 10.1128/CVI.00067-17.

18. Oguma K, Suzuki M, Sentsui H. 2017. Enzootic bovine leukosis in a

two-month-old calf. Virus Res 233:120–124. doi: 10.1016/j.virusres.2017.03.016

19. Fantini D, Vidimar V, Yu Y, Condello S, Meeks JJ. 2020. MutSignatures: an R

package for extraction and analysis of cancer mutational signatures. Sci Rep

10(1):18217. doi: 10.1038/s41598-020-75062-0.

20. Jones M, Wagner R, Radman M. 1987. Mismatch repair of deaminated

5-methyl-cytosine. J Mol Biol 194(1):155–159. doi:

10.1016/0022-2836(87)90724-8.

21. Bird A. 2002. DNA methylation patterns and epigenetic memory. Genes Dev

16(1):6–21. doi: 10.1101/gad.947102.

22. Velicescu M, Weisenberger DJ, Gonzales FA, Tsai YC, Nguyen CT, Jones PA. 2002.

Cell Division Is Required for de Novo Methylation of CpG Islands in Bladder

Cancer Cells. Cancer Res. 62(8):2378–2384.

23. Kim JY, Siegmund KD, Tavaré S, Shibata D. 2005. Age-related human small

intestine methylation: evidence for stem cell niches. BMC Med 3:10. doi:

10.1186/1741-7015-3-10.

24. Luo G, Santoro IM, McDaniel LD, Nishijima I, Mills M, Youssoufian H, Vogel H,

Schultz RA, Bradley A. 2000. Cancer predisposition caused by elevated mitotic

recombination in Bloom mice. Nat Genet 26(4):424–429. doi: 10.1038/82548.

25. Rufini A, Tucci P, Celardo I, Melino G. 2013. Senescence and aging: the critical

roles of p53. Oncogene 32(43):5129–5143. doi: 10.1038/onc.2012.640.

26. Lobry C, Oh P, Mansour MR, Look AT, Aifantis I. 2014. Notch signaling:

switching an oncogene to a tumor suppressor. Blood 123(16):2451–2459. doi:

10.1182/blood-2013-08-355818.

27. Zhang J, Vlasevska S, Wells VA, Nataraj S, Holmes AB, Duval R, Meyer SN, Mo T,

Basso K, Brindle PK, Hussein S, Dalla-Favera R, Pasqualucci L. 2017. The

CREBBP Acetyltransferase Is a Haploinsufficient Tumor Suppressor in B-cell

Lymphoma. Cancer Discov 7(3):322–337. doi: 10.1158/2159-8290.CD-16-1417.

28. Lo CW, Borjigin L, Saito S, Fukunaga K, Saitou E, Okazaki K, Mizutani T, Wada S,

Takeshima SN, Aida Y. 2020. BoLA-DRB3 Polymorphism is Associated with

Differential Susceptibility to Bovine Leukemia Virus-Induced Lymphoma and

Proviral Load. Viruses 12(3):352. doi: 10.3390/v12030352.

29. Murakami H, Uchiyama J, Suzuki C, Nikaido S, Shibuya K, Sato R, Maeda Y,

Tomioka M, Takeshima SN, Kato H, Sakaguchi M, Sentsui H, Aida Y, Tsukamoto

K. 2018. Variations in the viral genome and biological properties of bovine

leukemia virus wild-type strains. Virus Res 253:103–111. doi:

10.1016/j.virusres.2018.06.005.

30. Kobayashi T, Inagaki Y, Ohnuki N, Sato R, Murakami S, Imakawa K. 2020.

Increasing Bovine leukemia virus (BLV) proviral load is a risk factor for

progression of Enzootic bovine leucosis: A prospective study in Japan. Prev Vet

Med 178:104680 doi: 10.1016/j.prevetmed.2019.04.009.

31. Southgate L, Sukalo M, Karountzos ASV, Taylor EJ, Collinson CS, Ruddy D,

Snape KM, Dallapiccola B, Tolmie JL, Joss S, Brancati F, Digilio MC,

Graul-Neumann LM, Salviati L, Coerdt W, Jacquemin E, Wuyts W, Zenker M,

Machado RD, Trembath RC. 2015. Haploinsufficiency of the NOTCH1 Receptor

as a Cause of Adams-Oliver Syndrome With Variable Cardiac Anomalies. Circ

Cardiovasc Genet 8(4):572–581. doi: 10.1161/CIRCGENETICS.115.001086.

32. Chène P. 1998. In vitro analysis of the dominant negative effect of p53 mutants. J

Mol Biol 281(2):205–209. doi: 10.1006/jmbi.1998.1897.

33. Petitjean A, Mathe E, Kato S, Ishioka C, Tavtigian SV, Hainaut P, Olivier M. 2007.

Impact of mutant p53 functional properties on TP53 mutation patterns and tumor

phenotype: lessons from recent developments in the IARC TP53 database. Hum

Mutat 28(6):622–629. doi: 10.1002/humu.20495.

34. Olivier M, Hollstein M, Hainaut P. 2010. TP53 mutations in human cancers: origins,

consequences, and clinical use. Cold Spring Harb Perspect Biol 2(1):a001008. doi:

10.1101/cshperspect.a001008.

35. Liu Z, Liang Q, Ren Y Guo C, Ge X, Wang L, Cheng Q, Luo P, Zhang Y, Han X.

2023. Immunosenescence: molecular mechanisms and diseases. Sig Transduct

Target Ther 8:200 doi: 10.1038/s41392-023-01451-2.

36. Young KH, Leroy K, Møller MB, Colleoni GWB, Sánchez-Beato M, Kerbauy FR,

Haioun C, Eickhoff JC, Young AH, Gaulard P, Piris MA, Oberley TD, Rehrauer

WM, Kahl BS, Malter JS, Campo E, Delabie J, Gascoyne RD, Rosenwald A,

Rimsza L, Huang J, Braziel RM, Jaffe ES, Wilson WH, Staudt LM, Vose JM, Chan

WC, Weisenburger DD, Greiner TC. 2008. Structural profiles of TP53 gene

mutations predict clinical outcome in diffuse large B-cell lymphoma: an

international collaborative study. Blood 112(8):3088–3098. doi:

10.1182/blood-2008-01-129783.

37. Nishimori A, Andoh K, Matsuura Y, Kumagai A, Hatama S. 2021. Establishment of

a simplified inverse polymerase chain reaction method for diagnosis of enzootic

bovine leukosis. Arch Virol 166(3):841–851. doi: 10.1007/s00705-020-04945-4.

38. Okagawa T, Shimakura H, Konnai S, Saito M, Matsudaira T, Nao N, Yamada S,

Murakami K, Maekawa N, Murata S, Ohashi K. 2022. Diagnosis and Early

Prediction of Lymphoma Using High-Throughput Clonality Analysis of Bovine

Leukemia Virus-Infected Cells. Microbiol Spectr 10(6):e0259522. doi:

10.1128/spectrum.02595-22.

39. Forbes SA, Beare D, Boutselakis H, Bamford S, Bindal N, Tate J, Cole CG, Ward S,

Dawson E, Ponting L, Stefancsik R, Harsha B, Kok CY, Jia M, Jubb H, Sondka Z,

Thompson S, De T, Campbell PJ. 2017. COSMIC: somatic cancer genetics at

high-resolution. Nucleic Acids Res 45(D1):D777–D783. doi:

10.1093/nar/gkw1121.

40. Robinson JT., Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G,

Mesirov JP. 2011. Integrative Genomics Viewer. Nat Biotechnol 29(1):24–26. doi:

10.1038/nbt.1754.

 Figure legends

 Figure 1

A scheme for somatic mutation analysis in cattle with EBL development. Thirty-six

EBL cattle collected in previous studies [37, 38] were used for the mutation analysis.

Blood and tumor samples derived from the same EBL cases were examined to identify

variants in their genome using amplicon sequencing that targeted 21 cancer-related

genes. Following variant calling, variant lists were compared between blood and tumor

samples, and the variants existing only in tumor samples were identified as somatic

mutations. All identified somatic mutation were used for mutation pattern analysis.

Figure 2

Somatic mutation analysis in 36 EBL cases at various ages in months. (A) Identified

somatic mutations of 8 of 21 targeted cancer-related genes are shown in ascending order

of age for EBL cases. Genes in which a somatic mutation are not identified at all are

omitted. The top panel indicates mutation type, and the middle panel indicates detailed

classification of multiple mutations and presence or absence of loss of heterozygosity

(LOH) and germline mutations. The bottom panel indicates breed, sex, and age in

months of each case. Information on identified somatic mutations based on each

targeted gene or each EBL case is listed in Table 2 or Table S3, respectively. (B)

Number of cases and frequencies and types of somatic mutations are shown. (C–E)

Ages of cattle are compared among 36 EBL cases based on (C) presence of *TP53*

mutation, (D) intensities of mutation impact, and (E) presence of biallelic mutation.

TP53 wt, wildtype *TP53*; *TP53* mut, mutated *TP53*. High mutation impact means

stop-gained or frameshift mutation. Biallelic mutation includes multiple mutations or

683 mutation with LOH. Significant differences between the two groups were evaluated by

684 Mann-Whitney's U test (*P*-value < 0.05).

686 **Figure 3**

687 Analysis of mutation patterns in identified single nucleotide variants (SNVs). (A) Rates

688 of nucleotide substitutions in SNVs from 36 EBL cases are shown with a relative color

689 scale. Identified SNVs are put together in all cases or in two groups divided on the basis

690 of cattle age in months. The identical somatic mutation observed in different cases were

691 counted independently. ≥ 36-m-old, group of EBL cases over 36 months old; <

692 36-m-old, group of EBL cases less than 36 months old. (B) Mutation spectra based on

693 single base substitution (SBS) signatures are drawn using the MutSignatures R package.

694 Identified SNVs are divided into 96 different contexts and their mutation frequencies

695 calculated. The top 3 tri-nucleotide mutations in each group are marked in the spectra.

696 Mutation spectra in the ≥ 36-m-old group and the < 36-m-old group are defined as

697 Signatures A and B, respectively. (C) Heatmap shows similarity between Signature A/B

698 and known COSMIC signature 1–21. Cosine distances between each pair of signatures

699 were calculated and are shown by color intensity. (D) Evaluation of the mutation

700 process using known COSMIC signatures was performed using MutSignatures. The

701 vertical axis indicates the number of mutations imputed for each signature. COSMIC

702 signatures used for estimation are sorted according to validated evidence and reasonable

703 involvement in tumor development.

705 **Figure 4**

706 Hypothesis of the general mechanisms of EBL onset. There are 3 events for

BLV-infected cells to be transformed into neoplasms. First, infected cell clones whose

proviruses are integrated near cancer driver genes acquire long life and reproductive

activity [8]. Other clones in which the proviruses do not integrate near cancer driver

genes cannot overcome this first obstacle, and thus they are not able to be candidate

future cancer cells. Second, single somatic mutation occurs accidentally mainly at CpG

sites in cancer-related genes, e.g., *TP53*, *NOTCH1,* and *CREBBP*, as a first "hit".

Finally, an additional mutation or loss of heterozygosity (LOH) occurs on the left

normal allele as a second "hit". These 3 random events need to occur in the same

infected clone and be completed during the lives of infected animals, which is

considered to be the reason why only a small fraction of BLV-infected cattle develops

EBL after a long latent period.

Supplemental figure 1

Diagram of data interpretation of somatic mutation analysis in this study. Images of the

cording sequence (CDS) of targeted genes, read coverage in the IGV view, and

homologous chromosomes are shown. (A) In normal cells, there are only germline

mutations indicated in black on the targeted gene. (B) Single heterozygous somatic

mutation indicated in red is observed in the CDS region of cancer cells. It is interpreted

to be located on 1 allele of the homologous chromosome (pink). (C) Double somatic

mutations are heterozygously observed in the same targeted gene. This result is

interpreted to mean that the first mutation is located on one allele (pink), with the

second mutation on another allele (sky blue). (D) Single homozygous somatic mutation

is observed, which is interpreted as meaning that, following single heterozygous

mutation on one allele, loss of heterozygosity (LOH) occurs by chromosomal

recombination or deficiency.

Supplemental figure 2

Information on known COSMIC signatures obtained from the database. (A) In

COSMIC single base substitution (SBS) signatures 1–21, mutational process, supported

evidence, and presence of validation for real signatures are listed based on the COSMIC

website. (B) Mutation spectra of COSMIC SBS signatures 1, 6, 7 and 18 are drawn

using MutSignatures R package and reference signatures datasets. Tri-nucleotide

mutations at notable peaks in each signature are marked.

Table 1. Targeted cancer-related genes in the bovine genome

Table 2. Somatic mutations of 21 cancer-related genes identified from 36 EBL cases

Supplemental Table 1. Basic information on clinical EBL cases evaluated in this study

Supplemental Table 2. Primers used for amplicon sequencing

Supplemental Table 3. Details of somatic mutations identified in each EBL case

Supplemental Table 4. Loss of heterozygosity of germline mutations in chromosome

19

Figure 1

Figure 2

Figure 3

Figure 4

Random event 1)
Provirus integration near cancer driver genes (Rosewick *et al.*, 2017)

Random event 2)
Single somatic mutation mainly at CpG sites in cancer-related genes (<u>first "hit"</u>)

e.g., *TP53*, *NOTCH1*, *CREBBP*

Random event 3)
Additional mutation of another allele or LOH (<u>second "hit"</u>)

EBL onset (1–5%)

Aging

Cell division

Methylation at CpG sites

Life of infected cattle

BLV infection

BLV-infected clones in which the proviruses <u>do not</u> integrate near cancer driver genes

BLV-infected clones in which the proviruses integrate near cancer driver genes

Table 1. Targeted cancer-related genes in the bovine genome

Gene	Chr	Location	Gene ID	Transcript ID	mRNA length	CDS length	Protein length	Exon
TBL1XR1	1	90,030,514-90,202,770	ENSBTAG00000021910	ENSBTAT00000068590.1	2233	1,575	524	15
NRAS	3	28,614,965-28,624,861	ENSBTAG00000046797	ENSBTAT00000012897.6	4214	570	189	7
EZH2	4	112,024,592-112,069,782	ENSBTAG00000009426	ENSBTAT00000084115.1	2274	2,274	757	20
KMT2D	5	30,764,350-30,796,346	ENSBTAG00000014429	ENSBTAT00000083957.1	17350	17,049	5,682	50
KRAS	5	84,749,225-84,782,460	ENSBTAG00000009778	ENSBTAT00000043161.4	1007	570	189	5
EP300	5	112,282,183-112,344,853	ENSBTAG00000016198	ENSBTAT00000021556.6	7236	7,236	2,411	31
TET2	6	19,898,416-19,995,309	ENSBTAG00000017682	ENSBTAT00000023517.6	7056	6,024	2,007	10
TNFAIP3	9	75,740,934-75,756,753	ENSBTAG00000000436	ENSBTAT00000069028.1	5253	2,352	783	8
B2M	10	103,095,506-103,103,769	ENSBTAG00000048782	ENSBTAT00000086688.1	1038	360	119	4
NOTCH1	11	103,935,364-103,980,160	ENSBTAG00000022799	ENSBTAT00000017746.6	8190	7,602	2,533	34
ATM	15	17,853,846-17,996,481	ENSBTAG00000003111	ENSBTAT00000040104.4	9290	9,165	3,054	62
CD79A	18	51,344,675-51,348,682	ENSBTAG00000001882	ENSBTAT00000002451.4	1361	672	223	5
TP53	19	27,376,072-27,388,416	ENSBTAG00000001069	ENSBTAT00000001420.4	2160	1,161	386	11
CD79B	19	48,132,030-48,135,645	ENSBTAG00000044204	ENSBTAT00000061488.3	1401	690	229	6
MYD88	22	11,609,349-11,613,789	ENSBTAG00000000563	ENSBTAT00000000735.4	2786	1155	384	5
PIM1	23	11,066,555-11,071,948	ENSBTAG00000000396	ENSBTAT00000000511.6	2758	942	313	6
BCL2	24	61,396,513-61,588,241	ENSBTAG00000019302	ENSBTAT00000025701.6	1075	690	229	2
CREBBP	25	3,054,344-3,173,309	ENSBTAG00000026403	ENSBTAT00000005092.6	8515	7,308	2,435	31
SOCS1	25	9,913,007-9,913,678	ENSBTAG00000004386	ENSBTAT00000005753.5	672	672	223	1
CARD11	25	40,416,268-40,522,996	ENSBTAG00000014698	ENSBTAT00000037106.5	4515	3,465	1,154	25
PTEN	26	9,466,271-9,564,180	ENSBTAG00000009498	ENSBTAT00000012499.6	1722	1,722	573	9

Table 2. Somatic mutations of 21 cancer-related genes identified from 36 EBL cases

Gene	Chr	Mutation type	Genome location	Exon	CDS	Amino acid substitution	Mutation contexts	No. of EBL cases	Previous reports
TP53	19	stop gained	27380089	4/11	c.175G>T	p.Glu59*	T[C>A]A	33	-
		stop gained	27380015	4/11	c.249G>A	p.Trp83*	G[C>T]C	4	-
		missense	27379974	4/11	c.290G>A	p.Gly97Asp	G[C>T]C	36	-
		stop gained	27379449	5/11	c.382C>T	p.Gln128*	C[C>T]A	30	-
		missense	27379400	5/11	c.431C>T	p.Pro144Leu	C[C>T]G	33	[9]
		missense	27379338	5/11	c.493G>C	p.Val165Leu	A[C>G]A	17	-
		missense	27379331	5/11	c.500G>A	p.Arg167His	A[C>T]G	2, 7, 13,16, 19	[9–11]
		missense	27379319	5/11	c.512A>G	p.His171Arg	A[T>C]G	19	-
		missense	27379193	6/11	c.558C>G	p.His186Gln	A[C>G]C	22	-
		missense	27378842	7/11	c.687C>A	p.Phe229Leu	T[C>A]A	14	[9]
		missense	27378819	7/11	c.710G>A	p.Gly237Glu	C[C>T]C	6	-
		missense	27378808	7/11	c.721C>T	p.Arg241Trp	C[C>T]G	12, 18	[9, 11, 12]
		missense	27378805	7/11	c.724C>T	p.Arg242Trp	G[C>T]G	11, 21	[9, 12]
		missense	27378804	7/11	c.725G>A	p.Arg242Gln	C[C>T]G	9, 15	[9, 10]
		missense	27378780	7/11	c.749T>C	p.Leu250Pro	C[T>C]G	28	-
		stop gained	27378778	7/11	c.751G>T	p.Glu251*	T[C>A]C	10	-
		frameshift	27378769	7/11	c.758_759delCT	p.Ser253fs	-	9	-
		missense	27378376	8/11	c.824G>A	p.Arg275His	G[C>T]G	3, 22, 25	[9]
		missense	27378365	8/11	c.835G>A	p.Glu279Lys	T[C>T]C	27	-
		missense	27377694	10/11	c.988C>T	p.Arg330Cys	A[C>T]G	17	[9]
		stop gained	27377679	10/11	c.1003C>T	p.Arg335*	C[C>T]G	31	-
NOTCH1	11	frameshift	103956330	8/34	c.1403delG	p.Cys468fs	-	29[†], 31[†], 34	-
		missense	103956322	8/34	c.1412A>G	p.Gln471Arg	C[T>C]G	34	-
		missense	103946432	20/34	c.3248G>A	p.Arg1083His	G[C>T]G	6	-
KMT2D	5	missense	30777496	25/50	c.6344G>A	p.Arg2115Gln	C[C>T]G	22	-
		missense	30792244	45/50	c.16073G>T	p.Gly5358Val	T[C>A]C	4	-
		missense	30793539	46/50	c.16408G>A	p.Val5470Met	A[C>T]G	2	-
		missense	30796297	50/50	c.17000C>T	p.Ser5667Phe	T[C>T]C	26	
CREBBP	25	stop gained	3147752	2/31	c.733C>T	p.Gln245*	C[C>T]A	5	-
		frameshift	3085919	14/31	c.2650dupC	p.Gln884fs	-	29	-
		stop gained	3078462	17/31	c.3283C>T	p.Gln1095*	C[C>T]A	21	-
		stop gained	3061108	27/31	c.4480C>T	p.Gln1494*	T[C>T]A	5	-
KRAS	5	missense	84753447	2/5	c.34G>A	p.Gly12Ser	C[C>T]A	5	-
		missense	84753447	2/5	c.34G>T	p.Gly12Cys	C[C>T]A	30	-
PTEN	26	missense	9466850	1/9	c.580G>A	p.Asp194Asn	T[C>T]T	21	-
		stop gained	9538357	5/9	c.898C>T	p.Arg300*	A[C>T]G	30	-
CARD11	11	missense	40519575	22/25	c.2845G>A	p.Asp949Asn	T[C>T]C	23	-
MYD88	22	missense	11611445	3/5	c.881C>G	p.Ser294Cys	T[C>G]C	26	-

The symbol (†) indicates that the variant was called a germline mutation.