The autophagy protein, ATG14 safeguards against unscheduled pyroptosis activation

to enable embryo transport during early pregnancy

Pooja Popli[1], Arin K. Oestreich[2,3], Vineet K. Maurya[4], Marina N. Rowen[2,3], Yong Zhang[5],

Michael J. Holtzman[5], Ramya Masand[1], John Lydon[4], Shizuo Akira[6,7], Kelle H. Moley[2,3], and

Ramakrishna Kommagani[1,8*]

[1]Department of Pathology and Immunology, Baylor College of Medicine, One Baylor Plaza,

Houston, TX 77030, USA

[2]Department Obstetrics and Gynecology,

[3]Center for Reproductive Health Sciences, Washington University School of Medicine, St.

Louis, MO, 63110, USA

[4]Department of Molecular and Cellular Biology, Baylor College of Medicine, One Baylor Plaza,

Houston, TX 77030, USA

[5]Department of Medicine and Department of Cell Biology, Washington University School of

Medicine, St. Louis, MO, 63110, USA

[6]Department of Host Defense, Research Institute for Microbial Diseases (RIMD),

[7]Laboratory of Host Defense, World Premier Institute Immunology Frontier Research Center

(WPI-IFReC), Osaka University, Osaka 565-0871, Japan

[8]Department of Molecular Virology and Microbiology, Baylor College of Medicine, One Baylor

Plaza, Houston, TX 77030, USA.

*Correspondence to:

Ramakrishna Kommagani
Associate Professor
Department of Pathology and Immunology
Department of Molecular Virology and Microbiology
Alkek Center for Metagenomics and Microbiome Research
T228A, One Baylor Plaza
Baylor College of Medicine
Houston, TX, 77030
Email: Rama.Kommagani@bcm.edu
Phone: (713)-798-5085
Fax: 713-798-5838

Short title: Cellular self-digestion for female reproductive tract functions

The authors have declared that no conflict of interest exists.

Non-standard Abbreviations

WT: Wild Type

HESCs: Human Endometrial stromal cells

dpc: Days post coitum

E2: Estrogen

P4: Progesterone

VPS 15: Vacuolar sorting protein 15

cKO: Conditional knockout

FRT: Female reproductive tract

GSDMD: Gasdermin D

Keywords: fallopian tube, oviduct, embryo transport, uterus, pregnancy failure, infertility

Abstract

Recurrent pregnancy loss (RPL), characterized by two or more failed clinical pregnancies, poses a significant challenge to reproductive health. In addition to embryo quality and endometrial function, proper oviduct function is also essential for successful pregnancy establishment. Therefore, structural abnormalities or inflammation resulting from infection in the oviduct may impede the transport of embryos to the endometrium, thereby increasing the risk of miscarriage. However, our understanding of the biological processes that preserve the oviductal cellular structure and functional integrity is limited. Here, we report that Atg14-dependent autophagy plays a crucial role in maintaining the cellular integrity of the oviduct by controlling inflammatory responses, thereby supporting efficient embryo transport. Specifically, the conditional depletion of the autophagy-related gene, *Atg14* in the oviduct causes severe structural abnormalities compromising its cellular integrity leading to the abnormal retention of embryos. Interestingly, the selective loss of *Atg14* in oviduct ciliary epithelial cells did not impact female fertility, highlighting the specificity of ATG14 function in distinct cell types within the oviduct. Mechanistically, loss of *Atg14* triggered unscheduled pyroptosis via altering the mitochondrial integrity leading to inappropriate embryo retention and impeded embryo transport in the oviduct. Finally, pharmacological activation of pyroptosis in pregnant mice phenocopied the genetically induced defect and caused impairment in embryo transport. Together, we found that ATG14 safeguards against unscheduled pyroptosis activation to enable embryo transport from the oviduct to uterus for the successful implantation. Of clinical significance, these findings provide possible insights into the underlying mechanism(s) of early pregnancy loss and might aid in developing novel prevention strategies using autophagy modulators.

Introduction

A successful pregnancy is orchestrated by the sequential and coordinated events happening in the FRT [1]. Each of these events is crucial to advance to the next step in pregnancy. For example, as a first step, the ovary undergoes ovulation to release a mature ovum which is collected by oviduct fimbriae and funnelled through the infundibulum into the ampulla where they get fertilized by sperm. The embryos then pass through the ampulla-isthmus junction (AIJ) to enter the isthmus of the oviduct before exiting via the distal utero-tubal junction (UTJ) to implant on the wall of the uterus. Recently, the intricate structural and cellular diversity of the oviduct has drawn significant attention, emphasizing its vital role in facilitating embryo development and transport [2-4]. Morphologically, the oviduct is divided into four evolutionarily conserved regions: the infundibulum (nearest the ovary), ampulla (the site of fertilization), isthmus (serving as a sperm reservoir and site for early embryonic development), and utero-tubal junction (connected to the uterus). Each of these segments comprises epithelial, stromal, and smooth muscle cells with lumen lined up with an epithelium containing secretory (PAX8[+]) and multi-ciliated (FOXJ1[+]) cells. While the infundibulum and ampulla are composed of more ciliated than secretory cells, the isthmus has more secretory than ciliated cells [5, 6]. Successful pregnancy requires that the embryo transit the entire oviduct orchestrated by the following: (1) secretory cells producing oviduct fluid, (2) beating cilia to ensure unidirectional flow of the fluid, and (3) periodic contractions of surrounding muscle to propel fluid flow. Therefore, a combination of ciliary and muscular activity contributes to the overall success of oviduct embryo transport. This combination of ciliary and muscular activity plays a crucial role in the effective transport of the embryo through the oviduct. Any perturbations in these early pregnancy events can lead to adverse ripple effects that can compromise pregnancy outcomes. Previous studies have reported that impaired oviductal transport of embryos can lead to pregnancy failure and cause infertility in mice [7-10]. However, the underlying mechanism is not yet completely understood.

Autophagy is a cellular process, evolutionarily conserved from yeast to mammals, that

recycles long-lived proteins and organelles to maintain cell energy homeostasis. Autophagy

is classically activated through the nutrient sensor or mammalian target of the rapamycin

complex. Genetic screens for autophagy-defective mutants in yeast and other fungi have

currently identified 41 autophagy-related (ATG) genes that play a primary role in autophagy

[11, 12]. Approximately half of these genes have homologs in higher organisms, and *Atg14*

(also known as Barkor for Beclin 1 (*Becn1*)-associated autophagy-related key regulator) is

one of them [13, 14]. ATG14 is part of a protein complex that is composed of Beclin 1, vacuolar

sorting protein 15 (VPS15), and VPS34 (also named as Pik3c3, the catalytic subunit of the

class III phosphatidylinositol 3-kinase), and this ATG14-containing complex plays an important

role in the initiation process of autophagy.

Pyroptosis is a type of inflammatory cell death mediated by gasdermin (GSDM) and is

a product of continuous cell expansion until the cytomembrane ruptures, resulting in the

release of cellular contents that can activate strong inflammatory and immune responses.

Gasdermin family proteins are the primary executioners of pyroptosis. Cytotoxic N-terminal of

gasdermins generated from caspases mediated cleavage of gasdermin proteins oligomerizes

and forms pores across the cell membrane, leading to the release of proinflammatory

cytokines such as interleukin-1β (IL-1β) and interleukin-18 (IL-18). Aberrant activation of

pyroptosis has been implicated in the progression of many diseases including cancer and

autoimmune, cardiovascular, and infectious diseases [15-17]. Recent studies have shown that

the interactions between pyroptosis and autophagy play an important role in various

physiological and pathological processes [15-20]. For example, autophagy inhibition

upregulates galangin-induced pyroptosis in human glioblastoma multiforme cells and

promotes pneumococcus-induced pyroptosis [21, 22]. However, the studies defining the

molecular interactions between the pyroptosis, and autophagy pathways are very limited.

With a better understanding of the fundamental process of autophagy, its

pathophysiological functions have begun to be appreciated in the female reproductive tract.

For example, mice deficient in key autophagy genes such as genetic knockout of *Atg7* or

Becn1 result in primary ovarian insufficiency and reduced progesterone production [23, 24].

Similarly, recent studies from our group established the roles of three different autophagy-

specific genes: ATG16L, FIP200/RB1CC1, or BECN1 in endometrial physiological processes,

including receptivity and decidualization [25-27]. However, none of these proteins exhibited

any discernible impact on oviduct function. Surprisingly, in this study, we revealed a critical

role for Atg14 in maintaining proper oviduct function, specifically enabling the transport of

embryos to the uterus- a function distinct from that of other autophagy-related proteins. Loss

of ATG14 in the oviduct resulted in severe structural abnormalities, compromising its cellular

integrity, ultimately leading to embryo retention and infertility.

Materials and Methods

Animal care and use:

All animal studies were approved by the Institutional Animal Care and Use Committee of Washington University School of Medicine, Saint Louis, MO, USA and Use Committee of Baylor College of Medicine, Houston, TX, USA. *Atg14* f/f mice were provided by gift from Dr. Shizuo Akira at the Department of Host Defense, Research Institute for Microbial Diseases (RIMD), Osaka University, and previously described [28]. Wild type Pr-cre mice were provided by Dr. John Lydon at Baylor College of Medicine, Houston and previously described [29]. Atg14$^{flox/flox}$ mice, in which exon 4 was flanked by loxp sites, were bred to progesterone receptor cre (PR$^{cre/+}$) mice to generate (Atg14$^{flox/flox}$; PR$^{cre/+}$ mice), hereafter referred to as *Atg14* cKO mice. Both control and conditional knockout females were generated by crossing females carrying homozygous *Atg14*$^{flox/flox}$ alleles with *Atg14* cKO males. *Foxj1-cre* mice were a generous gift from Dr. Michael J. Holtzman at Washington University St. Louis and previously described [30]. *Foxj1/Atg14* cKO mice were generated by crossing females carrying homozygous *Atg14* $^{f/f}$/Foxj1$^{f/f}$ females with *Atg14*$^{f/f}$/Foxj1-cre males. All mice were age-matched and on a C57BL/6 genetic background (The Jackson Laboratory, Bar Harbor, ME). Mice were genotyped by PCR analysis of genomic DNA isolated from tail clippings using the gene-specific primers listed in **Supplementary Table 1.**

Fertility analysis and timed mating

Female fertility was determined by mating cohorts of *Atg14* cKO experimental (n=6) and control *Atg14* f/f (n=4) females individually starting at 8 weeks of age with sexually mature males of proven fertility. Similarly, breeding trials were set up for *Foxj1/Atg14*-cre females (n=6) and *Foxj1*/*Atg14* control 8-week-old f/f females (n=6). The numbers of litters and pups were tracked over 6 months for each female. Pups per litter for each genotype are reported as mean ± SEM. For timed mating, the morning on which the copulatory plug was first observed was considered 1 dpc. To visualize implantation sites, mice received a tail vein

injection of 50 µL of 1% Chicago Sky Blue dye (Sigma-Aldrich, St. Louis, MO, USA) at 5 dpc

just before sacrifice.

Steroid hormone treatments

The hormonal profile of pregnancy at the time of implantation was done using a previously

described experimental scheme [26]. Briefly, *Atg14* cKO and control females (8 weeks old)

were bilaterally ovariectomized under ketamine anesthesia with buprenorphine-SR as an

analgesic. Mice were allowed to rest for two weeks to dissipate all endogenous ovarian

hormones. After the resting period, mice were injected with 100 ng of estrogen (E2; Sigma-

Aldrich) dissolved in 100 µL of sesame oil on two consecutive days and then allowed to rest

for two days. At this point, mice were randomly divided into three groups of five: Vehicle-

treated (E2 priming) mice received four consecutive days of sesame oil injections; E2 group

mice received three days of sesame oil injections followed by a single injection of 50 ng of E2

on the fourth day; The E2/P4 mice received 1 mg of progesterone (P4; Sigma-Aldrich) for

three consecutive days followed by a single injection of 1 mg P4 plus 50 ng E2 on the fourth

day. All hormones were delivered by subcutaneous injection in a 90:10 ratio of sesame oil:

ethanol. Mice were euthanized 16 hours after the final hormone injection to collect the uteri. A

small piece of tissue from one uterine horn was processed in 4% neutral buffered

paraformaldehyde for histology, and the remaining tissue was snap-frozen and stored at -80

°C.

Hormone Analysis

For serum hormone levels measurement, blood was collected from D4 pregnant mice before

mice were sacrificed. Serum was separated from the blood by centrifugation and stored at -

80°C before hormone analysis. Serum P4 and E2 levels were measured by using ELISA kits

(Enzo life Sciences) according to the manufacturer's instructions.

Hematoxylin and eosin staining

Tissues were fixed in 4% paraformaldehyde, embedded in paraffin, and then sectioned (5 μm)

with a microtome (Leica Biosystem, Wetzlar, Germany). Tissue sections were deparaffinized,

rehydrated, and stained with Hematoxylin and Eosin (H&E) as described previously [31]. All

the histology was performed on three sections from each tissue of individual mice, and one

representative section image is shown in the respective figures.

Histological analysis

For histological analysis, the collected tissues (oviduct or uteri) were fixed in 4%

paraformaldehyde, embedded in paraffin, and sectioned. Sections (5 μm) were

immunostained (n=5 per group) as described previously [31]. Briefly, after deparaffinization,

sections were rehydrated in an ethanol gradient and then boiled for 20 min. in citrate buffer

(Vector Laboratories Inc., Newark, CA, USA) for antigen retrieval. Endogenous peroxidase

activity was quenched with Bloxall (Vector Laboratories Inc.), and tissues were blocked with

2.5% goat serum in PBS for 1 hr (Vector Laboratories Inc.). After washing in PBS three times,

tissue sections were incubated overnight at 4 °C in 2.5% goat serum containing the primary

antibodies listed in **Supplementary Table 2.** Sections were incubated for 1 hr with biotinylated

secondary antibody, washed, and incubated for 45 min with ABC reagent (Vector Laboratories

Inc.). Color was developed with 3, 3'-diaminobenzidine (DAB) peroxidase substrate (Vector

Laboratories Inc.), and sections were counter-stained with hematoxylin. Finally, sections were

dehydrated and mounted in Permount histological mounting medium (Thermo Fisher

Scientific, Waltham, MA, USA).

Transmission electron microscopy (TEM)

For ultrastructural analysis, oviducts were fixed in 2% paraformaldehyde/2.5% glutaraldehyde

(Ted Pella Inc., Redding, CA) in 100 mM cacodylate buffer, pH 7.2 for 1 hr at room temperature

and then overnight at 4°C. Samples were washed in cacodylate buffer and postfixed in 1%

osmium tetroxide (Ted Pella Inc.) for 1 hr. Samples were then rinsed extensively in dH20 prior

to en bloc staining with 1% aqueous uranyl acetate (Ted Pella Inc.) for 1 hr. Following several

rinses in dH20, samples were dehydrated in a graded series of ethanol and embedded in

Eponate 12 resin (Ted Pella Inc.). For initial evaluation semithin sections (0.5 μm) were cut

with a Leica Ultracut UCT7 ultramicrotome (Leica Microsystems Inc., Bannockburn, IL) and

stained with toluidine blue. Sections of 95 nm were then cut and stained with uranyl acetate

and lead citrate and viewed on a JEOL 1200 EX II transmission electron microscope (JEOL

USA Inc., Peabody, MA). Images at magnifications of 3,000X to 30,000X were taken with an

AMT 8-megapixel digital camera (Advanced Microscopy Techniques, Woburn, MA).

Immunofluorescence analysis

Formalin-fixed and paraffin-embedded sections were deparaffinized in xylene, rehydrated in

an ethanol gradient, and boiled in a citrate buffer (Vector Laboratories Inc.) for antigen

retrieval. After blocking with 2.5% goat serum in PBS (Vector laboratories) for 1 hr at room

temperature, sections were incubated overnight at 4°C with primary antibodies

(**Supplementary Table 2**) diluted in 2.5% normal goat serum. After washing with PBS,

sections were incubated with Alexa Fluor 488-conjugated secondary antibodies (Life

Technologies, Carlsbad, CA, USA) for 1 hr at room temperature, washed, and mounted with

ProLong Gold Antifade Mountant with DAPI (Thermo Fisher Scientific). All

Immunofluorescence images were obtained using a Zeiss LSM 880 confocal microscope (10x

and 40x objective lens).

Fresh oviduct tissues from 8-week-old mice from control (n=3) and cKO (n=3) were

collected and fixed in 4 % PFA at 4ºC overnight. After that, oviducts were washed with PBS

and cryoprotected using gradually increased sucrose concentrations in a row (15 and 30%

w/v). 24 h later, oviducts were embedded in an OCT medium, frozen (−80 °C), and cryo-

sectioned in the CM3050S Cryostat (Leica Biosystems, Germany). The cryosections were

mounted on Superfrost plus glass slides (Fisher Scientific, Pittsburgh, PA) and stored at −80°C

until used.

Cryo-sections were processed for immunofluorescence staining as described before

[32]. Briefly, cryosections were blocked in 2.5% normal goat serum. After washing with PBS,

sections were blocked with Mouse on Mouse (M.O.M.) IgG blocking reagent (M.O.M.

Fluorescein Kit, Vector Laboratories, #FMK-2201) diluted in 1% BSA per the manufacturer's

instructions. After blocking, sections were incubated overnight at 4°C with primary antibodies

(**Supplementary Table 2**) diluted in 2.5% normal goat serum. Next day, following washing

with PBS, sections were incubated with Alexa Fluor 488-conjugated secondary antibodies

(Life Technologies, Carlsbad, CA, USA) for 1 hr at room temperature, washed, and mounted

with ProLong Gold Antifade Mountant with DAPI (Thermo Fisher Scientific). All

Immunofluorescence images were imaged using a Nikon Fluorescent microscope.

Western blotting

Protein lysates (40 µg per lane) from uteri or oviducts were loaded on a 4-15% SDS-PAGE

gel (Bio-Rad, Hercules, CA, USA), separated in 1X Tris-Glycine Buffer (Bio-Rad), and

transferred to PVDF membranes via a wet electro-blotting system (Bio-Rad), all according to

the manufacturer's directions [33]. PVDF membranes were blocked for 1 hour in 5% non-fat

milk in Tris-buffered saline containing 0.1% Tween-20 (TBS-T, Bio-Rad), then incubated

overnight at 4 °C with antibodies listed in **Supplementary Table 2** in 5% BSA in TBS-T. Blots

were then probed with anti-Rabbit IgG conjugated with horseradish peroxidase (1:5000, Cell

Signaling Technology, Danvers, MA, USA) in 5% BSA in TBS-T for 1 hr at room temperature.

Signal was detected with the Pierce™ ECL Western Blotting Substrate (Millipore, Billerica,

MA, USA), and blot images were collected with a Bio-Rad ChemiDoc imaging system.

RNA Isolation and Quantitative Real-Time RT-PCR Analysis

Tissues/cells were lysed in RNA lysis buffer, and total RNA was extracted with the Purelink

RNA mini kit (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions.

RNA was quantified with a Nano-Drop 2000 (Thermo Fisher Scientific). Then, 1 µg of RNA

was reverse transcribed with the High-Capacity cDNA Reverse Transcription Kit (Thermo

Fisher Scientific). The amplified cDNA was diluted to 10 ng/µL, and qRT-PCR was performed

with primers listed in the **Supplementary Table 1** and TaqMan 2X master mix (Applied

Biosystems/Life Technologies, Grand Island, NY, USA) on a 7500 Fast Real-time PCR system

(Applied Biosystems/Life Technologies). The delta-delta cycle threshold method was used to

normalize expression to the reference gene 18S.

Treatment of mice with Polyphyllin VI

Polyphyllin VI (Selleck chemicals, Houston, TX, USA), a pharmacological agent that induces

caspase-1-mediated pyroptosis, was used to study its effects on embryo transport in the

oviduct [34]. Eight-week-old C57BL/6 mice were injected for three consecutive days starting

from 1 dpc with Polyphyllin VI activator dissolved in 40% PEG-300 (15mg/kg body weight).

Dimethyl sulfoxide with 40% PEG-300 was administered as a vehicle.

Statistics

A two-tailed paired student t-test was used to analyze data from experiments with two

experimental groups and one-way ANOVA followed by Tukey's post hoc multiple range test

was used for multiple comparisons. All data are presented as mean ±SEM. GraphPad Prism

9 software was used for all statistical analyses. Statistical tests, including p values, are

reported in the corresponding figure legends or, when possible, directly on the data image. To

ensure the reproducibility of our findings, experiments were replicated in a minimum of three

independent samples, to demonstrate biological significance, and at least three independent

times to ensure technical and experimental rigor and reproducibility.

Results

Conditional deletion of *Atg14* in the FRT results in infertility despite the normal ovarian function

To explore the role of *Atg14* in uterine function, we first determined its expression levels in uterus during early pregnancy (Day 1-7) in mice. We found distinct expression of ATG14 in all the uterine compartments (luminal epithelium, glands, and stroma) by day 1 of pregnancy, which disappeared by day 2 of pregnancy (**Fig. 1A**). However, the ATG14 expression in uterus reappeared by day 3 and persisted through the day 7 of pregnancy. The period from day 3 to day 7 is critical, as it is during this time when uterus begins to prepare for embryo implantation and undergoes decidualization process. Consistent with protein expression, similar *Atg14* expression at mRNA levels was noted in uteri from early pregnant mice (**Fig. 1B**). This analysis suggests a potential role for ATG14 protein in the uterine physiologic adaptations during early pregnancy. Thus, to study the role of ATG14 in uterine function, we generated a conditional knockout (PR^cre/+ /*Atg14*^flox/flox) mouse model by crossing *Atg14* flox/flox mice with mice expressing Cre recombinase under the control of progesterone receptor promoter (PR^cre/+). Histological examination of the uterus from adult females showed no gross morphological differences between *Atg14* cKO and control mice (**Fig. S1A**). Further, we did not find any overt defects in ovary as cKO mice had normal follicles and corpus luteum as like their corresponding controls (**Fig. S1B**). Analysis of transcript levels from the uterus, ovary, and liver showed that while *Atg14* levels were efficiently depleted in uteri from cKO mice the levels were unaltered in the ovary and liver samples in control and cKO mice (**Fig. 1C**). Immunofluorescence analysis further confirmed the efficient deletion of ATG14 in all uterine compartments (epithelium and stroma) of cKO mice uteri compared to corresponding control groups (**Fig. 1D**).

Considering the effective ablation, a 6-month breeding study was performed mating virile wild-type male mice with adult *Atg14* cKO and control female mice. We found that *Atg14* cKO females did not deliver any litter during the 6-months trial period. However, the control female (*Atg14* flox/flox) mice delivered an average of ~7-8 pups per litter every month (**Table**

1) (**Fig. 1E).** These findings suggest an indispensable role of *Atg14* in female fertility with intact ovarian function.

Loss of *Atg14* results in impaired embryo implantation and uterine receptivity in mice

Based on the normal ovarian morphology, we posited that the infertility observed in females with *Atg14* cKO status could be attributed to compromised uterine functioning. Analysis of the implanting embryos at Day 5 of pregnancy showed no implantation sites in the uteri of *Atg14* cKO females, whereas ~8 to 9 implantation sites were seen in the uteri of control mice at 5 dpc (**Fig. 2A, left panel**). Whilst there was no embryo present in the uterine lumen of *Atg14* cKO mice, a fully attached embryo encapsulated by the luminal uterine epithelium, was seen in control mice uteri (**Fig. 2A, middle panel**). Additionally, MUC1, a receptive marker expression persisted in the luminal epithelium of cKO mice uteri at 5 dpc as well as 4 dpc (**Fig. 2A, right panel, Fig. 2C**). The process of successful embryo attachment and implantation within the uterus necessitates a transition from a non-receptive to a receptive state, a transformation orchestrated under the regulated influence of steroid hormones. Thus, the effects of *Atg14* loss on uterine responsiveness to steroid hormones E2 and P4 were characterized. We performed an established hormone-induced uterine receptivity experiment, which involved ordered and co-stimulatory actions of E2 and P4 leading to the initiation of a receptive phase [35]. In response to E2 treatment, uteri from both *Atg14* control and cKO mice showed similar proliferation as evidenced by Ki-67 staining (**Fig. 2B, middle panel**). Following the co-stimulation with E2+P4 treatment, epithelial proliferation was inhibited with concomitant induction of stromal cell proliferation in control mice uteri indicating fully receptive uteri (**Fig. 2B**). Interestingly, cKO mice uteri failed to elicit sub-epithelial stromal cell proliferation and showed intact P4-driven inhibition of epithelial proliferation. Consistently, uteri from *Atg14* cKO mice at 4 dpc showed a reduced number of proliferating stromal cells (**Fig. 2D**). The absence of significant changes in E2-induced targets, such as *Lif*, *Mcm2*, *Ccnd1*, and *Fgf18*, at 4 dpc (**Fig. 2E**) supports our conclusion that ATG14 is required for P4-mediated but not for E2-mediated actions during uterine receptivity. Moreover, the normal serum levels of E2 and P4

337   at D4 of pregnancy rule out any hormonal imbalances, strongly suggesting that the observed

338   phenotype is primarily due to the uterine-specific loss of *Atg14* (**Fig. 2F**).

***Atg14* is required for maintaining oviductal cell structural integrity and embryo transport**

Given that *Atg14* cKO mice had impaired embryo implantation, we wondered whether

embryos are reaching the uterus timely through the oviduct. To determine this, we flushed

embryos from both the control and cKO mice uteri on day 4 of pregnancy. Interestingly, in cKO

mice, we could retrieve only 1-2% of embryos from their uteri, whereas in control mice, 100%

of well-developed blastocysts were retrieved from their uteri (**Fig. 3A & B**). To ensure the

timely transport of all embryos from the oviducts to the uteri, we also flushed oviducts from

both control and cKO mice. As expected, in the control mice oviduct flushing, we could not

recover any embryos or blastocysts, indicating their precise and timely transport to the uterus.

Unexpectedly, oviduct flushing from cKO mice resulted in the retrieval of approximately 80-

90% of blastocysts, suggesting their potential entrapment within the oviducts, impeding their

transit to the uterus (**Fig. 3A & B**). At 4 dpc, there was no significant difference in the average

number of blastocysts, morula, or non-viable or retrieved from *Atg14* control uteri and cKO

oviducts (**Fig. S4A).** However, we noted that the percentage of developmentally delayed

embryos appeared to be higher in *Atg14* cKO oviducts compared to the embryos retrieved

from control uteri (**Fig. 3C**). The histological analysis further confirmed an entrapped embryo

in the ampulla of *Atg14* cKO at 4 dpc as shown in **Fig. 3D**. Given the embryo retention

phenotype in oviduct, we sought to determine if ATG14 is expressed in this region. Consistent

with previous studies reporting PR-cre activity in the isthmus [8, 29], we found a significant

depletion of ATG14 in the isthmus compared to the ampulla (**Fig. S2A**). Further, the increased

p62 and LC3B expression in cKO oviducts suggests that the observed embryo retention

phenotype might be attributed to the loss of ATG14-dependent autophagy in cKO oviducts

(**Fig. S2B**).

Similarly, when we super-ovulated the mice and looked for the embryos on day 4 of

pregnancy, we could recover ~20-25 embryos from the cKO mice oviducts compared to only

4-5 embryos that were able to reach the uterus. In comparison, in control mice, 90% of

embryos were able to complete their pre-destined and timely transport to the uterus (**Fig. 3E**),

except 1-2% of unfertilized embryos, which remained in their oviducts.

To understand the underlying cause for retained embryos in cKO mice oviducts, we

performed histology analysis and determined the structural morphology of their oviducts. The

cKO mice oviduct lining shows marked eosinophilic cytoplasmic change akin to

decidualization in human oviducts. Some of the cells showed degenerative changes with

cytoplasmic vacuolization and nuclear pyknosis, loss of nuclear polarity, and loss of distinct

cell borders giving an appearance of fusion of cells (**Fig. 3F**). The marked cytologic

enlargement appears to cause luminal obliteration or narrowing resulting in a completely

unorganized, obstructed, and narrow lumen, thereby, hampering the path of embryos to reach

the uterus as evident from an entrapped embryo shown in **Fig. 3D**. Further, epithelial

(cytokeratin KRT8-positive) and myosalpinx (α-smooth muscle active [SMA]-positive,) marker

analysis revealed completely distorted epithelial structures (in terms of loss of epithelial cell

integrity) with no overt defects in the muscle organization in oviducts of cKO mice (**Fig. 3F,**

lower panel). Notably, the ampullary region from cKO oviducts seems to be normal with intact

epithelial and smooth muscle structures as evident from KRT8 and SMA staining (**Fig. 3F,**

upper panel).

Selective loss of *Atg14* in oviduct cilia is dispensable for female fertility

The oviduct epithelium primarily consists of two types of cells: ciliated and secretory

(non-ciliated) cells. Ciliated cells play a role in embryo and oocyte transport by means of ciliary

beat, and non-ciliated/secretory cells produce an oviductal fluid that is rich in amino acids and

various molecules, thereby providing an optimal micro-environment for sperm capacitation,

fertilization, embryonic survival, and development [36]. Therefore, we determined whether

oviducts from cKO mice possess normal ciliated and secretory cell composition. To do so, we

examined the expression of various markers, including acetylated-α-tubulin (cilia marker),

FOXJ1 (ciliogenesis markers), and PAX8 (non-ciliated/ secretory cell marker) in control and

cKO oviducts. As shown in **Fig. 4A,** while we observed normal ciliary structures in the ampulla

of both control and cKO oviducts, there was a substantial loss of the ciliary epithelial cells

(indicated by fewer α-tubulin and FOXJ1-positive cells) **(Fig. 4B, left panel and Fig. S3)** as

well as secretory cells (indicated by fewer PAX-8 positive cells) in the isthmus of cKO oviducts

(**Fig. 4B, right panel)**.

Given the importance of cilia in embryo transport, we wondered whether the loss of

Atg14 in oviduct cilia has any impact on embryo transport. To address this, we generated a

Foxj1-cre/+; *Atg14* f/f mouse model, wherein *Atg14* will be ablated only in ciliary epithelial

cells. We observed that ciliated epithelial cells that were positive for acetylated α-tubulin

staining did not appear to be different in Foxj1$^{Cre/+;}$ *Atg14* f/f mice oviduct compared to controls

suggesting normal ciliogenesis in Foxj1$^{Cre/+}$; *Atg14* f/f mice (**Fig. 4C**).

The 6-month breeding trial revealed that loss of *Atg14* in oviductal cilia had no impact

on fertility (**Table 2**, **Fig. 4D & E**). Consistently, D5 implantation study analysis showed ~7-8

visible embryo implantation sites in *Foxj1/Atg14* cKO mice like their corresponding controls

suggesting that embryos were able to make their way to the uterus in a timely manner and

undergo implantation despite the ablation of *Atg14* in the oviduct ciliary epithelial cells (**Fig.**

4F). These findings suggest that ciliary expression of *Atg14* is dispensable for embryo

transport.

ATG14 maintains mitochondria integrity and prevents unscheduled pyroptosis
activation in the oviduct to enable embryo transport

412   To gain more insights into structural defects, we performed TEM analysis on oviducts

413   collected from both control and cKO females on day 4 of pregnancy. Interestingly, oviducts

414   from cKO females had numerous altered mitochondrial structures with abnormally enlarged

415   mitochondria and a less dense matrix compared to control oviducts that had small and

416   compact mitochondria with tight cristae and a dense matrix (**Fig. 5A**). Corroborating and

extending those findings, we observed marked reductions of mitochondria network (as defined

by TOMM20 staining) in the sub-nuclear region of oviducts from *Atg14* cKO mice compared

to more densely packed mitochondrial network in the sub-nuclear region of control ones (**Fig.**

5B). Further co-localization study analysis revealed a uniform co-localization of Cytochrome

C with the TOM20 positive mitochondrial network in control oviducts reflecting the intact

mitochondrial integrity (**Fig. 5B**). In contrast, in *Atg14* cKO oviducts, cells with disrupted

mitochondrial networks exhibited increased cytosolic leakage of Cytochrome C (**Fig. 5B**).

Additionally, analysis of various mitochondrial functional and architecture markers (*Cox4i2*,

Pink1, *Opa1,* and *Drp1* showed reduced expression in *Atg14* cKO compared to controls (**Fig.**

5C).

Having identified the cellular mechanisms of Atg14, we aimed to delineate the

underlying molecular mechanisms associated with Atg14 function in the oviduct. Since we

found mitochondrial defects with ATG14 loss and altered mitochondrial structures have been

linked to the activation of pyroptosis, we determined if ATG14 regulates pyroptosis in oviduct

[37-39]. First, we assessed the key primary executors of the pyroptosis pathway, Gasdermin

D (GSDMD), and caspase-1. Immunofluorescence analysis revealed a remarkable

upregulation in GSDMD and caspase-1 expression in the cKO oviducts compared to controls

(**Fig. 5D & E**). However, histological analysis of uterus and ovary showed no induction in

GSDMD expression compared to their corresponding controls (**Fig. 5F**). Western blot analysis

further confirmed elevated expression of caspase-1 and GSDMD in cKO oviducts in

comparison to control oviducts as shown in **Fig. 5G.** Additionally, the qPCR analysis

demonstrated elevated levels of inflammatory markers, such as *Tnf-α* and *Cxcr3* in cKO

oviducts compared to control ones (**Fig. 5H**). Based on these findings, we posit that Atg14

plays a crucial role in regulating the pyroptotic pathway by preserving the mitochondrial

structural and functional integrity, and activation of pyroptosis owing to loss of ATG14 within

the oviduct.

To substantiate the notion, we also evaluated the impact of unscheduled activation of

pyroptosis on embryo transport. To test this, we employed a pyroptosis inducer, Polyphyllin

VI, and chose the optimal dose based on the established studies [34]. Wild-type females were

mated with virile males and following the plug detection treated with Polyphyllin VI for three

consecutive days from 1 to 4 dpc (**Fig. 5E**). We chose to treat to 1 to 4 dpc for treatment to

activate unwarranted pyroptosis in oviduct during the embryo transport. Following the

treatments, the oviducts, and uteri from both vehicle-treated and Polyphyllin VI-treated were

flushed. We found that pregnant females treated with Polyphyllin VI showed ~50% embryo

retention in the oviduct, whereas in the vehicle-treated group, no embryos were retained in

the oviduct (**Fig. 6A-C**). Further analysis of embryos retrieved from Polyphyllin-treated

oviducts showed more percentage of developmentally delayed and non-viable embryos

compared to embryos recovered from control uteri (**Fig. 6D**). The average number of embryos

recovered from polyphyllin- and vehicle-treated mice was not significantly different (**Fig. S4B**).

Histological analysis showed a marked induction in GSDMD expression compared to vehicle-

treated mice oviducts (**Fig. 6E**). However, precisely activating the unscheduled pyroptosis

during the critical period of embryo transport is technically challenging. Nonetheless, our

findings provide evidence that unscheduled pyroptosis adversely affects embryo transport

through the oviduct. Taken together, these results demonstrate that *Atg14* safeguards

pyroptosis activation in the oviduct and allows the smooth transport of embryos to the uterus

(**Fig. 6F**).

Discussion

In this study, we delineated the essential role of ATG14 in maintaining the structural integrity of the oviduct by preventing pyroptosis, which enables smooth embryo transit during early pregnancy. Specifically, we dissected the tissue-specific functions of ATG14 in the FRT using Cre driver mice targeting the female reproductive tract. Given that PR-Cre expresses postnatally in different tissues of FRT such as corpus luteum, oviducts, and different cellular compartments (epithelial, stromal, and myometrium) of uteri [29], we found that conditional ablation of ATG14-mediated functions in FRT resulted in infertility owing to hampered transport of embryos from the oviduct. Additionally, *Atg14* loss in uteri causes impaired embryo implantation and receptivity. Its ablation in the oviduct might lead to the activation of pyroptosis, an inflammatory-associated apoptosis pathway that causes the retention of embryos in the oviduct and prevents their timely transport to the uterus. However, the ovarian-specific functions were found to be unaffected despite the loss of *Atg14* in the corpus luteum.

The human endometrium is a complex dynamic tissue that undergoes sequential phases of proliferation and differentiation to support embryo implantation during the conceptive. However, in the absence of an implanting embryo, the endometrium sheds (menstruation) and initiates the regeneration process [40]. Previous reports indicated that autophagy is modulated in the human endometrium during the menstrual cycle. For example, autophagy is altered during the proliferative and secretory phases of the menstrual cycle with its highest activity occurring during the secretory phase when the stroma is decidualized [41]. Similarly, the level of autophagy was higher in postmenopausal human uterine epithelial cells compared to premenopausal uterine epithelial cells indicating the onset of autophagy upon estrogen deprivation [42]. Although all these studies reported that autophagy is hormonally regulated, however, the role of autophagy-specific proteins in the endometrium was not established till our group's recent reports. Specifically, studies from our group reported the role of three different autophagy-associated proteins: FIP200 [25], ATG16L [27], and BECLIN-1 in uterine functions [26]. Conditional ablation of *Fip200* and *Atg16l* in the uterus displayed fertility defects owing to impaired implantation, uterine receptivity, and decidualization defects.

On the contrary, loss of *Beclin1* in the uterus caused progressive loss of endometrial progenitor stem cells resulting in severe uterine developmental defects and rendering the mice infertile [26]. Although the autophagy-related proteins we studied so far influenced uterine functions, [25-27], we found a distinct role for ATG14 in both the uterine and oviduct-specific functions. It is intriguing to note that the absence of ATG14 did not affect the tissue integrity of the uterus. However, the severe structural abnormalities in the oviduct due to *Atg14* ablation unearthed a unique function of Atg14 in maintaining oviduct homeostasis. This distinctive role of ATG14, unlike other autophagy proteins, might be due to its pivotal role in the assembly of PtdIns3K complexes which is not the case for either FIP200 or ATG16L [40, 43, 44]. Nevertheless, understanding the specific contributions of each core autophagy protein in reproductive tract functions is necessary which requires substantial efforts.

The retention of preimplantation embryos in the oviduct has been established as a significant contributor to implantation failures, presenting challenges to the overall reproductive health of women [8-10]. Embryo transport in the oviduct is known to be controlled primarily by two major physiological responses: ciliary activity and muscle contractility. In our study, a dramatic decrease in the number of Foxj1^{+ve} ciliary epithelial cells and Pax-8^{+ve} secretory cells in *Atg14* cKO mice implied cell-type specific actions for ATG14 in the oviduct. Interestingly, the specific ablation of *Atg14* in Foxj1+ve ciliary epithelial cells of the oviduct does not appear to impact fertility. This suggests that the absence of ATG14 within ciliary cells does not impact the process of embryo transport. Although unexpected, this is consistent with other reports that have demonstrated the dispensability of cilia for embryo transport in the oviduct [8, 9].

Pyroptosis is a highly inflammatory form of programmed cell death, characterized by cell swelling, flattening of the cytoplasm, and large bubbles-like protrusions on the plasma membrane [45-47]. Several recent studies found a link between autophagy and pyroptosis [48]. *In vivo* studies also found that mice lacking the autophagy genes *Atg14*, *Fip200*, *Atg5*, or *Atg7* in myeloid cells had more pronounced lung inflammation [49]. In our study, cellular swelling, and fused membranous structures (a unique feature of activation of pyroptosis)

observed in the oviductal epithelial folds from *Atg14* cKO mice. This aberrant activation of the

pyroptosis pathway due to loss of *Atg14* appears to adversely affect the oviduct cellular

structural integrity. These findings have implications beyond the oviduct, as autophagy can

modulate the inflammatory signaling pathway through different avenues. For example, first,

autophagy prevents mitochondrial reactive oxygen species release that can activate the

inflammasome by inhibiting IL1-β and IL18 production through the digestion of dysfunctional

mitochondria [50]. Second, autophagy is capable of targeting inflammasome complexes for

degradation, which prevents the cleavage of pro-IL-1β and pro-IL-18 into biologically active

forms. Finally, autophagy machinery further regulates IL-1β levels by engulfing and degrading

pro-IL-1β proteins [50]. In our study, accumulation of abnormally enlarged and dysfunctional

mitochondria in the absence of ATG14 in the oviduct clearly suggests the induction of severe

inflammatory conditions hampering the embryo transit through the oviduct. These findings are

further substantiated by the upregulation of pivotal mediators of the pyroptosis program,

including GSDMD and caspase-1, in *Atg14* cKO mice. Notably, the induction of pyroptosis in

the absence of ATG14 is specifically localized to the oviducts, with no parallel activation

detected in either the ovary or uterus. One potential reason for this observed phenomenon

could be the oviduct's unique molecular signatures that get activated in response to the

presence of gametes. A recent study by Finnerty et al. (2024) reported that sperm induce pro-

inflammatory conditions in the oviduct, which are preceded by an anti-inflammatory response

triggered by the presence of embryos [51]. The persistent activation of inflammatory or

pyroptotic conditions following the loss of Atg14 suggests that Atg14 may be a critical

autophagy protein responsible for suppressing pyroptosis in the oviduct. Moreover,

understanding the complex rheostat between the ATG14 and inflammation regulatory axis is

highly intriguing and will necessitate future studies employing sophisticated models, such as

combined knockout mice where ATG14 is deleted alongside key inflammatory regulators (e.g.,

NLRP3, GSDMD, or CASPASE-1).

In this report, we revealed tissue-specific roles of ATG14 in governing oviductal

transport, and uterine receptivity which are necessary for embryo implantation and pregnancy

establishment. Of mechanistic insights, we delineated a novel function of ATG14 as a negative

regulator of pyroptosis, which is vital for proper embryo transport in the oviduct (**Fig. 6F**). Thus,

this study not only sheds light on the involvement of autophagy in oviductal embryo transport

but also underscores the importance of autophagy in preserving the oviduct tissue integrity.

Such insights hold promise for advancing our understanding of gynecological pathologies

associated with the oviduct including tubal pregnancies.

Acknowledgments

We thank Goutham Davuluri, and Ashirbad Guria from our group for the technical assistance. We thank Dr. Robert Lawrence (senior editor at Baylor College of Medicine) for assistance with manuscript editing. We thank WashU Molecular Microbiology Imaging Facility for assisting us with transmission electron microscopy. This work was funded, in part, by the National Institutes of Health/National Institute of Child Health and Human Development (grants R01HD102680, R01HD104813, and R01HD065435) to RK, R01 HD-042311 to JPL, and in part by National Institutes of Health/National Heart, Lung, and Blood Institute (grant R35HL145242) to MJH.

Authors' roles

PP and RK designed experiments, conducted most of the studies, analyzed the data, and wrote the manuscript. AKO, VKM, and MNR, conducted some of the experiments. MJH, JL, SA, YZ, and KHM, provided critical reagents for the study. RM helped us with oviduct pathology phenotype analysis. RK conceived the project, supervised the work, and wrote the manuscript. All authors critically reviewed the manuscript.

Conflict of Interest

The authors have declared that no conflict of interest exists.

570   **Table 1**: **Six-month breeding trial of *Atg14* control and cKO females with wild-type males**

571   **Table 2**: **Six-month breeding trial of *Foxj1*/*Atg14* control and cKO females with wild-type**

572   **males**

 Figure legends

 Figure 1: *Atg14* loss in the FRT results in infertility. (**A**) Representative immunofluorescent

 images of uteri from pregnant mice (n=3) at the indicated days of pregnancy stained with an

 ATG14-specific antibody (green). LE: luminal epithelium, G: glands, S: stroma, Scale bar: 100

 μm. Rabbit IgG was used as an isotype control for staining. (**B**) Relative transcript levels of

 Atg14 mRNA in uteri from pregnant mice (n=3-5) at indicated days of pregnancy. mRNA levels

 are normalized to levels of 18S m-RNA. Data are presented as mean ±SEM; **P<0.01,

 P>0.05, ns=non-significant. (**C**) Relative mRNA levels of *Atg14* in 8-week-old virgin control

 and cKO mice uteri, ovary, and liver (n=5). mRNA levels are normalized to levels of 18S

 mRNA. Data are presented as mean ±SEM; ***P<0.001, P>0.05, ns=non-significant. (**D**)

 Representative immunofluorescent images of ATG14 expression in different uterine

 compartments in control (n=5) and *Atg14* cKO mice (n=5). LE: luminal epithelium, G: glands,

 S: stroma. (**E**) (*left panel*) Relative number of pups/female/litter and (*right panel*) and relative

 number of total pups/months of *Atg14* control (n=4) and cKO mice (n=5) sacrificed after the

 breeding trial. Data are presented as mean ±SEM; ***P<0.001; P>0.05, ns=non-significant.

 Figure 2: *Atg14* is critical for embryo implantation, and uterine receptivity. (**A**) Gross

 images of 5.0 dpc uteri of control (n=5) and *Atg14* cKO mice (n=5) injected with Chicago Sky

 Blue dye to visualize implantation sites (denoted by black arrows) (*left panel*). H&E-stained

 cross-sections (4X & 40X) of 5.0 dpc uteri of control (n=5) and *Atg14* cKO (n=5) mice to

 visualize embryo implantation (*Middle panel*). The asterisk denotes the embryo.

 Immunofluorescence analysis of uterine tissues from control (n=5) and *Atg14* cKO mice (n=5),

 stained with MUC1 and KRT8 (*right panel*). LE: luminal epithelium, G: glands, S: stroma. (**B**)

 Representative immunofluorescence images of uteri from control (n=5) and *Atg14* cKO mice

 (n=5) stained for Ki-67 following Oil or E2 or E2+P4 treatment (n=5 mice/group); scale bar:

 100 μm. LE: luminal epithelium, G: glands, S: stroma. (**C**) Immunofluorescence analysis of

 KRT8 (green), MUC1 (Red), and (**D**) Ki-67 (red) in the uteri of 4 dpc control and *Atg14* cKO

 mice; scale bars, 100 μm. (**E**) Relative transcript levels of *Lif*, *Mcm2*, *Ccnd1,* and *Fgf18* in

control and cKO uteri at 4dpc. mRNA levels are normalized to levels of 18S m-RNA. Data are

presented as mean ±SEM; ns=non-significant. (**F**) Levels of steroid hormones estradiol and

progesterone from serum collected during euthanasia of 4 dpc control or cKO mice.

Figure 3: *Atg14* is critical for embryo transport in oviduct. (**A**) Representative images

(*upper panel*) and (**B**) Percentage of embryos (*lower panel*) collected at 4 dpc from the uteri

or the oviducts of control (n=6) or *Atg14* cKO mice (n=6-8). (**C**) Percentage of blastocysts,

morulae, developmentally delayed or nonviable embryos collected from *Atg14* control mice

uteri and Atg14 cKO mice oviducts at 4 dpc. (**D**) Histological analysis using H&E staining of

the ampullary and isthmic region of the oviduct from control and *Atg14* cKO female mice at 4

dpc (n=3 mice/genotype). (**E**) Embryos retrieved from the oviduct and uterus of super-ovulated

Atg14 control or cKO mice at 4 dpc (n=3 mice/genotype). (**F**) Immunofluorescence analysis of

KRT8 (green), and α-SMA (Red), in the oviduct of 4 dpc control (n=5) and *Atg14* cKO mice

(n=5) ampulla (*upper panel*) and isthmus (*lower panel*).

Figure 4: *Atg14* loss in oviduct cilia is dispensable for embryo transport. (A)

Immunofluorescence analysis of acetylated α-tubulin (green), and DAPI (blue) in oviduct of 4

dpc control (n=5) and *Atg14* cKO mice (n=5). (**B**) Immunohistochemical analysis of FOXJ1

and PAX8 at 4 dpc (n=5). (**C**) Immunohistochemical analysis of acetylated α-tubulin in 8-week-

old control (n=5) and *Foxj1*/*Atg14* cKO mice (n=5). Images are taken at 6X and 40X. Scale

bar: 100μm and 10 μm. (**D & E**) Relative number of pups/females/litter and relative number

of total pups/months of control (n=5) and *Foxj1*/*Atg14* cKO mice (n=5) sacrificed after the

breeding trial. Data are presented as mean ± SEM; P>0.05, ns=non-significant. (**F**) Gross

images of 5.0 dpc uteri of control (n=3) and *Foxj1/Atg14 cKO mice* (n=3) injected with Chicago

Sky Blue dye to visualize implantation sites (denoted by black arrows).

Figure 5: ATG14 facilitates embryo transport in the oviduct by preserving mitochondrial

integrity and inhibiting the activation of pyroptosis. (**A**) Transmission electron microscopy

of oviducts at 4 dpc from control (n=3) and *Atg14* cKO (n=3)**.** Black arrowheads in control

oviducts show small, compact mitochondria with tight cristae. Red arrowheads in cKO oviducts

show abnormally enlarged mitochondria with loose cristae. (**B**) Immunofluorescence analysis

of TOM20 (green) and Cytochrome C (red) in *Atg14* control and cKO oviducts cryo-sections.

The inset shows the zoomed image of the selected area from control and cKO oviducts,

Arrowheads in cKO inset oviduct section show the leaked cytochrome C in the cytoplasm.

Tissues were counterstained with DAPI (blue) to visualize nuclei; scale bars, 10 μm. (**C**)

Relative transcript levels of *Opa1*, *Cox4i2*, *Drp1,* and Pink1 in oviduct tissues. Data are

presented as mean ±SEM. *P<0.05; **P<0.01 compared with controls. 18S was used as an

internal control (**D**) Immunofluorescence analysis of GSDMD (red) + KRT8 (green), Caspase-

1 (red) + KRT8 (green) in oviducts of adult control (n=5), and *Atg14* cKO mice (n=5). Tissues

were counterstained with DAPI (blue) to visualize nuclei; scale bars, 100 μm. (**E**)

Immunohistochemical analysis of GSDMD expression in adult oviduct tissues (*left panel*). The

middle and -right panels show the zoom-in images to show the relative GSDMD expression in

the ampulla and isthmus section from control and cKO oviducts. (**F**) Immunohistochemical

analysis of GSDMD expression in adult uterus and ovary tissues (**G**) Western blotting to show

protein levels of Caspase-1, and GSDMD in oviduct tissues. β-actin is used as a loading

control. (**H**) Relative transcript levels of *Tnf-α* and *Cxcr3* in oviduct tissues. Data are presented

as mean ±SEM. *P<0.05; **P<0.01; ***P<0.001 compared with controls. 18S was used as an

internal control.

Figure 6: Pharmacological activation of pyroptosis in the oviduct inhibits embryo

transport. (**A**) Experimental strategy for pyroptosis activation in pregnant female mice (**B**)

Embryos flushed from the vehicle (n=3) or polyphyllin VI treated (n=3) D-4 pregnant females.

(**C**) Percentage of embryos recovered from oviducts or uteri. (**D**) Percentage of blastocysts,

morulae, developmentally delayed or nonviable embryos collected from vehicle or Polyphyllin

IV oviducts at 4 dpc. (**E**) Immunohistochemistry to show GSDMD expression in the isthmus

section of polyphyllin IV-treated and vehicle-treated mice. 40X objective, Scale bar: 5 μm. (**F**)

Graphical illustration to show embryo transport and pyroptosis regulation in the oviduct.

References

References

1.   Wang, H. and S.K. Dey, *Roadmap to embryo implantation: clues from mouse models.* Nat Rev
Genet, 2006. **7**(3): p. 185-99.
2.   Ford, M.J., et al., *Oviduct epithelial cells constitute two developmentally distinct lineages that*
are spatially separated along the distal-proximal axis. Cell Rep, 2021. **36**(10): p. 109677.
3.   McKey, J., et al., *Integration of mouse ovary morphogenesis with developmental dynamics of*
the oviduct, ovarian ligaments, and rete ovarii. Elife, 2022. **11**.
4.   Harwalkar, K., et al., *Anatomical and cellular heterogeneity in the mouse oviduct-its potential*
roles in reproduction and preimplantation developmentdagger. Biol Reprod, 2021. **104**(6): p.
1249-1261.
5.   Dirksen, E.R. and P. Satir, *Ciliary activity in the mouse oviduct as studied by transmission and*
scanning electron microscopy. Tissue Cell, 1972. **4**(3): p. 389-403.
6.   Li, S., et al., *Estrogen receptor alpha is required for oviductal transport of embryos.* FASEB J,
2017. **31**(4): p. 1595-1607.
7.   Jiang, R., et al., *CDC42 governs normal oviduct multiciliogenesis through activating AKT to*
ensure timely embryo transport. Cell Death Dis, 2022. **13**(9): p. 757.
8.   Herrera, G.G.B., et al., *Oviductal Retention of Embryos in Female Mice Lacking Estrogen*
Receptor alpha in the Isthmus and the Uterus. Endocrinology, 2020. **161**(2).
9.   Wang, H., et al., *Aberrant cannabinoid signaling impairs oviductal transport of embryos.* Nat
Med, 2004. **10**(10): p. 1074-80.
10.   Qian, J., et al., *Caffeine consumption during early pregnancy impairs oviductal embryo*
transport, embryonic development and uterine receptivity in mice. Biol Reprod, 2018. **99**(6): p.
1266-1275.
11.   Klionsky, D.J., *Autophagy: from phenomenology to molecular understanding in less than a*
decade. Nat Rev Mol Cell Biol, 2007. **8**(11): p. 931-7.
12.   Wen, X. and D.J. Klionsky, *An overview of macroautophagy in yeast.* J Mol Biol, 2016. **428**(9 Pt
A): p. 1681-99.
13.   Itakura, E., et al., *Beclin 1 forms two distinct phosphatidylinositol 3-kinase complexes with*
mammalian Atg14 and UVRAG. Mol Biol Cell, 2008. **19**(12): p. 5360-72.
14.   Sun, Q., et al., *Identification of Barkor as a mammalian autophagy-specific factor for Beclin 1*
and class III phosphatidylinositol 3-kinase. Proc Natl Acad Sci U S A, 2008. **105**(49): p. 19211-
6.
15.   Wang, Y.Y., X.L. Liu, and R. Zhao, *Induction of Pyroptosis and Its Implications in Cancer*
Management. Front Oncol, 2019. **9**: p. 971.
16.   Man, S.M., R. Karki, and T.D. Kanneganti, *Molecular mechanisms and functions of pyroptosis,*
inflammatory caspases and inflammasomes in infectious diseases. Immunol Rev, 2017. **277**(1):
p. 61-75.
17.   Wei, X., et al., *Role of pyroptosis in inflammation and cancer.* Cell Mol Immunol, 2022. **19**(9):
p. 971-992.
18.   Lin, L., et al., *Autophagy, Pyroptosis, and Ferroptosis: New Regulatory Mechanisms for*
Atherosclerosis. Front Cell Dev Biol, 2021. **9**: p. 809955.
19.   Li, X., et al., *Autophagy, Pyroptosis and Ferroptosis are Rising Stars in the Pathogenesis of*
Diabetic Nephropathy. Diabetes Metab Syndr Obes, 2024. **17**: p. 1289-1299.
20.   Zhao, H., et al., *The Role of Autophagy and Pyroptosis in Liver Disorders.* Int J Mol Sci, 2022.
23(11).
21.   Kong, Y., et al., *The Natural Flavonoid Galangin Elicits Apoptosis, Pyroptosis, and Autophagy*
in Glioblastoma. Front Oncol, 2019. **9**: p. 942.
22.   Kim, J.Y., et al., *Streptococcus pneumoniae induces pyroptosis through the regulation of*
autophagy in murine microglia. Oncotarget, 2015. **6**(42): p. 44161-78.

23. Song, Z.H., et al., *Germ cell-specific Atg7 knockout results in primary ovarian insufficiency in*
female mice. Cell Death Dis, 2015. **6**(1): p. e1589.
24. Gawriluk, T.R., et al., *Beclin-1 deficiency in the murine ovary results in the reduction of*
progesterone production to promote preterm labor. Proc Natl Acad Sci U S A, 2014. **111**(40):
p. E4194-203.
25. Oestreich, A.K., et al., *The autophagy protein, FIP200 (RB1CC1) mediates progesterone*
responses governing uterine receptivity and decidualizationdagger. Biol Reprod, 2020. **102**(4):
p. 843-851.
26. Popli, P., et al., *Beclin-1-dependent autophagy, but not apoptosis, is critical for stem-cell-*
mediated endometrial programming and the establishment of pregnancy. Dev Cell, 2023.
58(10): p. 885-897 e4.
27. Oestreich, A.K., et al., *The Autophagy Gene Atg16L1 is Necessary for Endometrial*
Decidualization. Endocrinology, 2020. **161**(1).
28. Matsunaga, K., et al., *Two Beclin 1-binding proteins, Atg14L and Rubicon, reciprocally regulate*
autophagy at different stages. Nat Cell Biol, 2009. **11**(4): p. 385-96.
29. Soyal, S.M., et al., *Cre-mediated recombination in cell lineages that express the progesterone*
receptor. Genesis, 2005. **41**(2): p. 58-66.
30. Zhang, Y., et al., *A transgenic FOXJ1-Cre system for gene inactivation in ciliated epithelial cells.*
Am J Respir Cell Mol Biol, 2007. **36**(5): p. 515-9.
31. Chadchan, S.B., et al., *Antibiotic therapy with metronidazole reduces endometriosis disease*
progression in mice: a potential role for gut microbiota. Hum Reprod, 2019. **34**(6): p. 1106-
1116.
32. Chojnacki, A.K., et al., *Tissue imaging reveals disruption of epithelial mitochondrial networks*
and loss of mitochondria-associated cytochrome-C in inflamed human and murine colon.
Mitochondrion, 2023. **68**: p. 44-59.
33. Kommagani, R., et al., *The Promyelocytic Leukemia Zinc Finger Transcription Factor Is Critical*
for Human Endometrial Stromal Cell Decidualization. PLoS Genet, 2016. **12**(4): p. e1005937.
34. Teng, J.F., et al., *Polyphyllin VI Induces Caspase-1-Mediated Pyroptosis via the Induction of*
ROS/NF-kappaB/NLRP3/GSDMD Signal Axis in Non-Small Cell Lung Cancer. Cancers (Basel),
2020. **12**(1).
35. Tong, W. and J.W. Pollard, *Progesterone inhibits estrogen-induced cyclin D1 and cdk4 nuclear*
translocation, cyclin E- and cyclin A-cdk2 kinase activation, and cell proliferation in uterine
epithelial cells in mice. Mol Cell Biol, 1999. **19**(3): p. 2251-64.
36. Ito, S., Y. Yamamoto, and K. Kimura, *Analysis of ciliogenesis process in the bovine oviduct based*
on immunohistochemical classification. Mol Biol Rep, 2020. **47**(2): p. 1003-1012.
37. Liu, Q., et al., *The role of mitochondria in NLRP3 inflammasome activation.* Mol Immunol,
2018. **103**: p. 115-124.
38. Shi, Y., et al., *Echinacoside Induces Mitochondria-Mediated Pyroptosis through Raf/MEK/ERK*
Signaling in Non-Small Cell Lung Cancer Cells. J Immunol Res, 2022. **2022**: p. 3351268.
39. Wang, Y., et al., *Mitochondrial ROS promote macrophage pyroptosis by inducing GSDMD*
oxidation. J Mol Cell Biol, 2019. **11**(12): p. 1069-1082.
40. Popli, P., A.J. Sun, and R. Kommagani, *The Multifaceted Role of Autophagy in Endometrium*
Homeostasis and Disease. Reprod Sci, 2022. **29**(4): p. 1054-1067.
41. Choi, J., et al., *Differential induction of autophagy by mTOR is associated with abnormal*
apoptosis in ovarian endometriotic cysts. Mol Hum Reprod, 2014. **20**(4): p. 309-17.
42. Zhou, S., et al., *Author Correction: Menopause-induced uterine epithelium atrophy results from*
arachidonic acid/prostaglandin E2 axis inhibition-mediated autophagic cell death. Sci Rep,
2020. **10**(1): p. 6915.
43. Fan, W., A. Nassiri, and Q. Zhong, *Autophagosome targeting and membrane curvature sensing*
by Barkor/Atg14(L). Proc Natl Acad Sci U S A, 2011. **108**(19): p. 7769-74.

44.   Diao, J., et al., *ATG14 promotes membrane tethering and fusion of autophagosomes to*
755   *endolysosomes.* Nature, 2015. **520**(7548): p. 563-6.
756   45.   Ma, X., et al., *LPS Mediates Bovine Endometrial Epithelial Cell Pyroptosis Directly Through Both*
757   *NLRP3 Classical and Non-Classical Inflammasome Pathways.* Front Immunol, 2021. **12**: p.
758   676088.
759   46.   Cookson, B.T. and M.A. Brennan, *Pro-inflammatory programmed cell death.* Trends Microbiol,
760   2001. **9**(3): p. 113-4.
761   47.   Jorgensen, I. and E.A. Miao, *Pyroptotic cell death defends against intracellular pathogens.*
762   Immunol Rev, 2015. **265**(1): p. 130-42.
763   48.   Liu, J., et al., *Autophagy blockage promotes the pyroptosis of ox-LDL-treated macrophages by*
764   *modulating the p62/Nrf2/ARE axis.* J Physiol Biochem, 2021. **77**(3): p. 419-429.
765   49.   Lu, Q., et al., *Homeostatic Control of Innate Lung Inflammation by Vici Syndrome Gene Epg5*
766   *and Additional Autophagy Genes Promotes Influenza Pathogenesis.* Cell Host Microbe, 2016.
767   **19**(1): p. 102-13.
768   50.   Netea-Maier, R.T., et al., *Modulation of inflammation by autophagy: Consequences for human*
769   *disease.* Autophagy, 2016. **12**(2): p. 245-60.
770   51.   Finnerty, R.M., et al., *Multi-omics analyses and machine learning prediction of oviductal*
771   *responses in the presence of gametes and embryos.* bioRxiv, 2024.

A

D-1 D-2 D-3 D-4

D-5 D-6 D-7

ATG14/DAPI IgG Isotype control

B

mRNA levels of *Atg14*/18S

Days of pregnancy

C

Control *Atg14* cKO

Uterus *Ovary* *Liver*

Atg14 mRNA levels/18s

**** ns ns

D

Control *Atg14* cKO

ATG14/DAPI

E

Average no. of pups/female/litter

Control *Atg14* cKO

****

Average pups/month

Control

Atg14 cKO

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6