Gut microbiota gate host exposure to metabolites from dietary Solanums

Catherine S. Liou[1], Mikhail Iakiviak[2], Jakub Rajniak[3], Pallavi P. Murugkar[2], Steven K. Higginbottom[4], Marissa Jasper[2], Michael A. Fischbach[2,3], Elizabeth S. Sattely[1,5]*

[1] Department of Chemical Engineering, Stanford University, Stanford, CA, USA
[2] Sarafan ChEM-H Institute, Stanford University, Stanford, CA, USA
[3] Department of Bioengineering, Stanford University, Stanford, CA, USA
[4] Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA, USA
[5] Howard Hughes Medical Institute, Stanford, CA, USA

*Corresponding author: sattely@stanford.edu

ABSTRACT

Dietary plants are molecularly rich but the fates of these compounds post-ingestion and their implications for human health are largely unknown. Here, we systematically characterized the major chemical contributions of widely consumed Solanum species (nightshades) to the human metabolome and established their mechanisms of production. Using untargeted metabolomics, we found a series of steroidal alkaloids resulting from tomatine, solanine, and chaconine that are dominant diet-derived compounds in systemic circulation following ingestion of potato and tomato. By comparing serum and tissue metabolomics of colonized and microbiome-depleted mice, we found that the gut microbiota metabolizes these compounds extensively, altering their absorption and gating host exposure to the diverse resulting metabolites. By screening metabolism with human stool samples, we established that steroidal glycoalkaloid metabolism varies inter-individually in a population. Further, using a collection of human commensal type strains, we found that a specific subset of strains in a community is responsible for steroidal glycoalkaloid metabolism, and chemical output of a community is determined by its strain-level composition. Microbial metabolism alters Solanum metabolite bioavailability and bioactivity in the context of acetylcholinesterase inhibition. The combination of dietary input and specific gut colonists dictate the composition of a variable component of the human metabolome. Our study provides insights into the molecular mechanisms of a diet-microbiome interaction and its effects on host physiology.

INTRODUCTION

Plants are sources of vast molecular diversity in diet, contributing not only carbohydrates,

proteins, fats, and vitamins but also specialized small molecules. These secondary metabolites,

produced by plants for defense, environmental stress tolerance, and nutrient acquisition, are

estimated to be ingested in significant quantities of at least 50 milligrams per day[1,2] and have been

associated with numerous pharmacological effects[3,4]. The immense chemical complexity of these

plants outmatches our understanding of the major molecular inputs from diet, which obscures

connections between dietary plant consumption and human health and disease states. Further

exacerbating this challenge is metabolite transformation after ingestion, which vastly expands the

array of diet-derived metabolites in the human metabolome.

In addition to host digestive processes, dietary compounds can be transformed by the

commensal bacteria that reside in the gut. Gut microbiota-specific metabolism has an outsized

impact on access to nutrients from diet[5] and we are only beginning to explore the specific

molecular contributions by bacterial metabolism to the host metabolome[6,7]. Biotransformation by

the gut microbiota can alter compound bioactivity and bioavailability; furthermore, microbiota-

specific metabolism can vary inter-individually, resulting in divergent molecular exposures and

health outcomes from a common dietary input[8,9]. Elucidation of bioactive compounds from diet,

their transformation by gut microbiota, and their ultimate impacts on human health will not only

expand our ability to take advantage of the benefits of dietary plants, but also provide valuable

insight into the molecular interactions that exist at the interface of the plant, bacterial, and animal

kingdoms.

Nightshades from the family Solanaceae, including potatoes, tomatoes, and eggplants,

are of particular interest among dietary plants. Solanums are some of the most abundantly

consumed plants in the United States[10] and are rich in specialized compounds studied in

numerous physiological contexts, including carotenoids, steroidal glycoalkaloids, and

flavonoids[11,12]. Diets rich in tomatoes and potatoes have been associated with different

pharmacological effects; tomato ingestion, for example, has been shown to reduce serum LDL

oxidation[13] and decrease the incidence of cardiovascular disease in a cohort of women[14].

However, the specific compounds that drive these pharmacological effects are still under active

investigation.

In this study, we use dietary Solanums as a model to explore the major molecular inputs

from a commonly consumed food, the role of gut microbiota in gating host exposure to these

inputs, and the impacts of commensal-specific metabolism on compound bioavailability and

bioactivity. We used untargeted metabolomics in animal models and identified steroidal alkaloids

as the major diet-derived compounds in host circulation following Solanum ingestion. We mapped

the metabolic fates of dietary steroidal glycoalkaloids and reconstituted the production of these

metabolites *in vitro* and *ex vivo* using a combination of gut commensal strains and human liver

enzymes. These results reveal a paradigm where specific bacterial strains are responsible for key

steps in compound metabolism, yielding metabolite and bioactivity profiles unique to individual

community compositions. Biotransformation by the gut microbiota is a layer of metabolism that

can profoundly impact interactions between diet and the host and must be characterized to fully

understand the physiological effects of a dietary molecule.

76   **RESULTS**

Major molecular inputs from dietary Solanums

We focused on Solanums, including potatoes and tomatoes, as abundantly consumed

plants known to be rich in specialized compounds and began by exploring the major molecular

inputs derived from ingestion of these plants. We reasoned that an untargeted metabolomics

approach would enable unbiased identification of the compounds encountered post-ingestion that

may mediate the pharmacological effects associated with these plants. We used liquid

chromatography coupled to mass spectrometry (LC-MS) to profile serum from Swiss Webster

mice provided with *ad libitum* access to ripe Microtom tomatoes in addition to a standard diet for

15 days. To capture a broad range of molecule classes in serum, analyses were performed using

both hydrophilic interaction liquid chromatography (HILIC) for the retention of more polar

compounds and reverse phase chromatography for the retention of nonpolar compounds.

Untargeted metabolomics was used to identify tomato-associated compounds in systemic

circulation; mass features were filtered to identify those that significantly accumulated at least

three-fold higher in samples from mice receiving tomatoes than in mice that did not receive

tomatoes (**Fig S1A**). Many of these molecules were absent in the tomato supplement and are

likely derived from the metabolism of tomato compounds or endogenous mouse compounds

produced in response to tomato consumption (**Fig 1A**). We identified one such metabolite to be

trimethylamine N-oxide (TMAO), known to be produced by intestinal bacteria from dietary

trimethylamine containing precursors like choline, betaine, and carnitine[15]. Serum TMAO levels

were increased 2.5-fold by tomato consumption (**Fig S1B**), an unexpected result considering the

association of circulating TMAO with red meat consumption and increased risk for cardiovascular

disease[16]. These data highlight the strengths of an untargeted approach in identifying unexpected
compounds resulting from a dietary intervention.

When subjected to tandem mass spectrometry (MS/MS), fourteen of the 28 tomato-
associated compounds found in serum had mass fragmentation patterns characteristic of
steroidal alkaloids (SAs) (**Fig S1C**). Steroidal alkaloids are nitrogen containing compounds
derived from cholesterol and accumulate highly in Solanums as glycosides called steroidal
glycoalkaloids (SGAs). Solanums are known to biosynthesize species-specific SGAs with
structural variations in both the steroidal aglycone and the tetraose or triose glycone. Commercial
tomato cultivars accumulate the SGA tomatine, comprised of a tomatidine aglycone conjugated
to a lycotetraose moiety, which is converted *in planta* to esculeoside A during fruit ripening (**Fig
1B**). We found major SA compounds in serum following tomato consumption to include tomatidine
and esculeogenin A, the aglycones derived from tomatine and esculeoside A, respectively, as
well as other oxidized versions of SA aglycones (**Fig 1A**). Interestingly, despite the abundance of
SGA and the lack of detectable SA aglycones in the provided tomato, we observed primarily
aglycones accumulating in the serum. These results suggest that SGAs undergo significant
metabolism following consumption.

Tomato steroidal glycoalkaloids are deglycosylated and further modified after ingestion

To track the metabolic fates of SGAs independently of the other dietary components in
tomatoes, we next provided Swiss Webster mice with pure tomatine in drinking water. Reasoning
that metabolites may accumulate more highly in certain tissues than in systemic circulation, we
used untargeted metabolomics to characterize the major tomatine metabolites that are present in
animal tissues and therefore have the potential to interact with tissue-specific physiology. We
surveyed a panel of tissues for steroidal alkaloid content, including two tissues along the digestive
tract (colonic tissue and luminal contents from the cecum), another two that are first-pass tissues
for nutrient absorption (mesenteric lymph nodes (MLNs) for molecules transported by the
lymphatic system and liver for molecules transported by the portal vein), and kidney, based on
previous studies that have tracked tissue accumulation of SGAs from potato[17].

In addition to some of the SA metabolites previously detected in systemic circulation
following tomato ingestion, we were able to identify several aglycone metabolites that
accumulated highly in specific tissues (**Fig 1C**). Selected compounds were structurally
characterized through retention time and MS/MS spectrum matching to synthesized standards,

including tomatidine oxidized at the C3 alcohol to yield the ketone 3-oxotomatidine (**Fig S2A**) and
sulfated to form tomatidine 3-sulfate (**Fig S2B**). We also noted the accumulation of metabolites
putatively identified through MS/MS analyses to be C3-OH epimerized, acetylated, and doubly
hydroxylated analogues of tomatidine. These metabolites reveal that extensive structural
modifications of SGAs beyond deglycosylation occur following ingestion.

Because only SGAs were present in the dietary intervention, we next wanted to probe how
the abundant aglycones in serum and tissues were formed. Tomatine and many of the SA
aglycone metabolites were found primarily in the cecal contents and the colon (**Fig 1C**). A few
aglycones, however, including 3-oxotomatidine, hydroxytomatidine, and tomatidine sulfate were
also abundant in the liver. We hypothesized that these compounds could be formed by hepatic
metabolism. To test this hypothesis, we incubated human liver protein extracts with tomatine and
tomatidine. While human liver enzymes did not modify the parent SGA tomatine *in vitro*, we
observed formation of both tomatidine sulfate (**Fig S2C**) and hydroxytomatidine (**Fig S2D**) after
incubation with the aglycone tomatidine. While we could not detect any hepatic oxidation of
tomatidine to form 3-oxotomatidine, human liver enzymes were able to perform the reverse
reaction and non-selectively reduce synthesized 3-oxotomatidine to tomatidine and a putative
3αOH epimer of tomatidine (**Fig S2E**). Using human liver enzymes, we were able to recapitulate
the production of a few SA metabolites from tomatidine. Many of the major metabolic
transformations present in serum metabolites, however, including tomatine deglycosylation, were
not catalyzed by human liver enzymes *in vitro*. These data suggest that human liver enzymes
have a limited role in generating systemically circulating metabolites from SGAs.

Potato steroidal glycoalkaloids from potato undergo similar metabolism as tomato SGAs

To explore the major molecular inputs of other dietary Solanums and track their metabolic
fates, we provided Swiss Webster mice with a single dose of the potato SGA solanine, followed
by diet supplemented with 10% potato peel. Using a similar approach as for the tomato dietary
supplement, we performed untargeted metabolomics to identify the major diet-derived
metabolites that accumulated in systemic circulation (**Fig S3A**). We found that dietary intervention
with potatoes significantly increased serum levels of several compounds, including hippuric acid
(**Fig S3B**), potentially derived from metabolism of phenolic compounds abundant in potato[18]. The
majority of the compounds that accumulated in serum with potato consumption, however, had
MS/MS spectra characteristic of solanidine-based steroidal alkaloids (**Fig S3C**). These

metabolites included the major SGAs solanine and chaconine present in potato (**Fig 1B**), as well
as the corresponding aglycone solanidine. Reminiscent of the metabolic fates of tomatine, we
observed compounds produced by extensive structural modifications to the aglycone solanidine,
including hydroxylation, sulfation, and oxidation. We additionally observed hydrogenation of the
Δ5,6 double bond of solanidine to form 5α and 5β isomers of dihydrosolanidine, confirmed by
MS/MS spectrum and retention time matching to a synthetic standard (**Fig S4A**).

To investigate the biosynthetic origins of these serum metabolites, we next incubated
solanine, chaconine, and solanidine with human liver proteins *in vitro*. As was the case with the
tomato SGA, human liver proteins did not metabolize glycosides solanine and chaconine. We
recapitulated the formation of a subset of the modified aglycone metabolites observed in serum
by incubating liver proteins with solanidine, namely sulfated (**Fig S4B**) and hydroxylated (**Fig
S4C**) solanidine. However, other modified aglycone metabolites, including keto- and saturated
analogues of solanidine, were not produced. These data show that, like tomato SGAs, potato
SGAs undergo extensive modifications following deglycosylation, many of which are produced
independently of hepatic metabolism.

Taken together, we found that steroidal alkaloid aglycones represent a substantial portion
of the detectable diet-derived metabolites in systemic circulation following a meal of tomatoes and
potatoes. These data support other studies that have noted steroidal alkaloids and their
metabolites to be small molecule biomarkers characteristic of tomato consumption[19–21]. We further
characterized the SA metabolites observed *in vivo* to uncover previously uncharacterized
modifications that occur to SGAs following ingestion, including oxidations and hydrogenations.

Gut microbiota gate the formation of steroidal alkaloid aglycones *in vivo*

As we could not detect aglycone formation from SGAs by human metabolic enzymes *in
vitro*, we next wanted to explore the mechanisms through which SGAs are deglycosylated *in vivo*.
Considering the relative abundance of SA aglycones within the digestive tract (**Fig 1C**) and the
established capabilities of gut commensals in polysaccharide hydrolysis[22], we hypothesized that
gut microbiota could play a role in mediating SGA metabolism. Fecal pellets from conventional
mice and mice treated with broad spectrum antibiotics[23] were incubated with exogenously added
tomatine. We observed formation of the aglycone tomatidine only in the feces of mice that had
not received antibiotics (**Fig S5A**), indicating that native murine commensals are capable of
tomatine deglycosylation.

In addition to gut microbiota, host digestive processes including acid hydrolysis in the
stomach and glycosidases in the intestine may also contribute to SGA deglycosylation. To assess
the contributions of gut microbiota towards SGA deglycosylation *in vivo*, we provided C57BL/6J
mice with diet supplemented with 10% tomato concurrently with antibiotics treatment. The
aglycone tomatidine was absent from serum (**Fig 2A**) and tissues (**Fig S5B**) in mice receiving
antibiotics treatment. Other modified aglycone metabolites, including 3-oxotomatidine and
tomatidine sulfate, were similarly depleted by antibiotics treatment (**Fig 2B**). Because the potato
SGAs are structurally distinct to tomatine, we also explored whether the gut microbiota is similarly
necessary for solanine and chaconine deglycosylation. C57BL/6J mice concurrently receiving
antibiotics and a diet containing potato peels were deficient in the aglycone solanidine, both in
systemic circulation (**Fig 2C**) and tissues that highly accumulated SGAs (**Fig S5C**). These
animals were also deficient in modified analogues of solanidine (**Fig S5D**). These data show that
SGA deglycosylation is performed primarily by gut microbiota and not host digestion. Additionally,
we observe that SGA deglycosylation by gut microbiota precede and gate the formation of the
extensive modified aglycone metabolites characterized *in vivo*.

To explore the relevance of this metabolism to humans, we analyzed urine from human
colectomy patients lacking an intact colon and therefore deficient in colonic microbiota[24]. Even
without a specified dietary intervention, we were able to detect potato SGAs solanine and
chaconine co-occurring in 8 out of 12 colectomy donors and 9 out of the 17 donors with intact
colons, indicative of recent potato consumption (**Fig S5E**). In donor samples that also contained
potato SGAs, the aglycone solanidine (**Fig 2D**) was largely depleted in colectomy patients
compared to donors with intact colons. We also detected solanidine metabolites modified by
hydroxylation, sulfation, and hydrogenation, transformations that we had observed in previous
mouse studies; these molecules were similarly depleted in colectomy patients relative to patients
with intact colons (**Fig S5F**). These data show not only that SA metabolites are found in systemic
circulation with presumed dietary exposure to Solanums, but that metabolism by gut microbiota
dictate SGA metabolism and host exposure to steroidal alkaloid aglycones.

Inter-individual variation in *ex vivo* SGA metabolism by human stool samples

A key implication of SGAs being metabolized exclusively by gut microbiota is the potential
variation in metabolism between individuals, with unique bacterial communities generating distinct
metabolite profiles from a common molecular input. Having established the necessity of gut

microbiota in metabolizing SGAs, we next explored how these compounds might be metabolized

differently between individuals. We began by identifying the major compounds produced from

SGAs by complex human gut microbial communities. Thirty human stool samples[25] were

incubated *ex vivo* with the major SGAs from dietary Solanums and assayed for product formation

by LC-MS. In addition to potato and tomato SGAs, we also included SGAs from eggplant,

solamargine and solasodine, which share a solasodine aglycone but differ in their glycone

moieties. MS fragmentation was used to identify mass features containing characteristic steroidal

alkaloid substructures (**Fig S6A**). Using this approach, we found compounds that had not been

previously observed in our analyses of mouse samples, including putative phosphoethanolamine

conjugation to the aglycones tomatidine and solasodine (**Fig S6B**).

Comparing glycoalkaloid conversion by different human donor samples, we observed

significant inter-individual differences in SGA deglycosylation, with some samples extensively

metabolizing SGAs and others inactive (**Fig 3A**). SGA metabolism also appeared to correlate with

identity of the glycone. Solanine and the eggplant SGA solasonine share the same solatriose

glycone and were metabolized by a similar subset of samples. In the same manner, another

subset of donors was able to metabolize chaconine and the eggplant SGA solamargine, which

share a chacotriose sugar.

We further tested the ability of donor samples to metabolize steroidal alkaloid aglycones.

In addition to variation in SGA deglycosylation, we also observed inter-individual variation in the

major modifications made to exogenously added aglycone. Hydrogenation of Δ5,6 aglycones

solanidine and solasodine was previously observed in mice following potato consumption (**Fig**

S5D) and occurred only in two out of the 30 human stool samples surveyed (**Fig 3B**). More

common modifications included phosphoethanolamine conjugation to tomatidine and solasodine,

as well as oxidation and sulfation of all three aglycones surveyed. The variation in the abundance

and identity of metabolites produced *ex vivo* by different human stool samples indicate that, from

a common input of dietary Solanums, individuals may be exposed to different compounds

depending on compositions of gut microbial communities.

We noted that some samples that did not metabolize SGAs were nonetheless active on

the corresponding aglycone. For example, sample 26 did not significantly metabolize either

solanine or solasonine, but still produced abundant sulfonated aglycone when incubated directly

with solanidine and solasodine (**Fig S7**), respectively. These results suggest that SGAs may be

metabolized cooperatively by the gut microbiota, with different strains responsible for

deglycosylation and aglycone modification.

Cooperative metabolism by human commensal type strains

Having noted significant variation in SGA metabolism by human stool samples, we sought to identify specific human commensal strains involved in these metabolic steps. We individually cultured 116 human commensal type strains, selected to represent strains that are abundant and prevalent in a Western microbiome[26], in rich medium containing potato and tomato SGAs and assayed for compound metabolism using LC-MS. Following incubation with tomatine, aglycones were produced by only a few strains out of those surveyed, including *Roseburia intestinalis* L1-82, *Bacteroides uniformis* ATCC 8492, and strains from the genus *Parabacteroides* (**Fig 4**). Among these strains, *Parabacteroides distasonis* ATCC 8503 was notable for being able to metabolize all three SGAs surveyed. Generally, however, the capacity of a strain to deglycosylate one SGA did not guarantee the metabolism of other SGAs. *Ruminococcus gnavus* ATCC 29149, for example, was able to hydrolyze chaconine but not solanine, despite the identical aglycone moiety. These data show that SGA deglycosylation is a substrate-specific transformation performed by only select strains in a community.

To circumvent the necessity for SGA deglycosylation and directly assay for aglycone modification, we additionally incubated the commensal strains directly with aglycones tomatidine and solanidine. Bacteroidetes were found to be broadly active on these compounds to generate the same sulfated and phosphoethanolamine-conjugated metabolites previously produced by human stool. Some of the strains that were active on SGAs were also capable of modifying the resulting aglycone; *Bacteroides uniformis* ATCC 8492 produced sulfonated and phosphoethanolamine-conjugated compounds following incubation with both tomatine and tomatidine. However, SGA deglycosylation appeared to be generally independent of aglycone modification, with different subsets of strains active on the aglycone and the corresponding glycoside.

Since SGA deglycosylation and aglycone modification are generally carried out by different strains, we hypothesized that SGAs may be metabolized cooperatively in a community context, reliant on a subset of strains to liberate the aglycone before further modification by other members of the community. We explored this possibility by incubating cocultures of defined composition with tomatine in rich medium. Six *Bacteroides* strains were selected for their ability to modify tomatidine but not deglycosylate tomatine (**Fig S8**). Inclusion of either *Roseburia intestinalis* LI-82 or *Roseburia inulinivorans* DSM 16841, strains capable of deglycosylation but

not aglycone modification, in the coculture enabled production of sulfonated,
phosphoethanolamine-conjugated, and acetylated tomatidine. From these results, we show that,
despite their relative hydrophobicity, steroidal alkaloid aglycones are exchanged as intermediates
between bacterial strains in a community. Furthermore, we see that the major compounds derived
from SGAs *in vitro* can be manipulated by the strains present in a community. These results
support our findings that that inter-individual differences in microbiome composition (**Fig 3**) drive
divergent host exposure to SA metabolites.

SGA metabolism by gut microbiota dictates compound bioavailability and bioactivity

Having established that SGAs are major dietary inputs from dietary Solanums and that gut
microbiota dictate the metabolic fates of dietary SGAs, we next explored the consequences of
this metabolism on host physiology. As hydrolysis of the glycone by gut microbiota significantly
impact physiochemical properties like solubility and molecular size, we began by investigating the
impact of SGA deglycosylation on compound bioavailability. We used a Caco-2 human colonic
cell monolayer as a model for compound transport across the intestinal enterocyte barrier[27].
Potato SAs solanine, chaconine, and solanidine were added to the apical side of the monolayer
representing the intestinal lumen. Following incubation, compound transport across the
monolayer was determined using LC-MS. While the SGAs solanine and chaconine primarily
remained on the apical side, the majority of the aglycone solanidine was observed to accumulate
on the basolateral side, indicating that solanidine is better transported across the epithelium than
SGAs (**Fig S9A**). We next reproduced these differences in bioavailability *in vivo* in mice fed
potatoes concurrently with antibiotics treatment. The total quantity of SGA-derived compounds
absorbed into systemic circulation was estimated as the sum of serum SGAs and tracked
agylcones.   We found that animals with intact gut microbiota had significantly higher
concentrations of total SGA-derived metabolites in serum relative to mice that had been treated
with antibiotics, with aglycones representing the majority of these compounds (**Fig 5A**). Taken
together, these data show that metabolism of SGAs by the gut microbiota convert SGAs to a
better absorbed form and influence circulating concentrations of these specialized steroidal
alkaloid compounds in the host.

Having noted differences in bioavailability resulting from SGA metabolism, we next
explored the impact of gut microbiota-mediated SGA metabolism on compound biological activity.
SAs have been shown to have numerous effects on host physiology, including anti-inflammation[28]

and chemoprevention through cytotoxicity in cancer cell lines[29–31]. We began by exploring the impact of deglycosylation on SGA cytotoxicity *in vitro*. We treated the human colon cancer cell line Caco-2 with potato SGAs solanine and chaconine and observed concentration-dependent loss of cell viability. In contrast, the aglycone solanidine had a negligible impact on cell viability (**Fig S9B**). A similar connection between glycosylation and *in vitro* toxicity was observed for the tomato SGA tomatine and the aglycone tomatidine (**Fig S9C**)[31]. These data suggest that SGA deglycosylation by gut microbiota inactivates the toxic potential of these compounds.

In addition to cytotoxicity and uniquely among the dietary SGAs, the potato SGAs solanine and chaconine are established cholinergic toxins in humans, capable of inhibiting the two cholinesterases that terminate acetylcholine (ACh) neurotransmission[32,33] and implicated in poisonings caused by the ingestion of improperly stored potatoes[34]. Considering the established interactions between potato SGAs and these physiologically essential cholinesterases, we next explored the effects of gut microbiota-mediated SGA metabolism on ACh hydrolysis. Recombinant human acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) were treated with solanine, chaconine, and solanidine, with enzyme activity measured by tracking hydrolysis of substrate mimics acetylthiocholine and butyrylthiocholine, respectively[35]. The SGAs solanine and chaconine inhibited acetylthiocholine hydrolysis relative to a vehicle treated control in a dose-dependent manner while the aglycone solanidine did not significantly impact AChE activity at concentrations up to 100 μM (**Fig 5B)**. While AChE activity was not impacted at the concentrations tested, solanidine did inhibit BChE activity to similar extents as did the SGAs (**Fig S9D**). These results show that deglycosylation of potato SGAs solanine and chaconine to the aglycone solanidine alleviates AChE inhibition but not BChE inhibition by these compounds. Furthermore, we showed with physiologically relevant dietary interventions that SGAs can accumulate to micromolar quantities gastrointestinal tissue. In animals concurrently receiving 5% potato peel in diet with antibiotics, colonic levels of SGAs accumulated to 2.98 ± 0.38 pmol/mg fresh weight tissue (**Fig S5C**), concentrations which corresponded to about 40% inhibition of AChE activity *in vitro*.

In addition to its role in the central nervous system, ACh is a key neurotransmitter in the enteric nervous system[36]. Because AChE and BChE can both contribute to ACh hydrolysis in the gastrointestinal tract, we aimed to investigate the effects of SGA metabolites on a physiological mix of cholinesterases. We chose to focus on the colon as the primary tissue of interest, as SGAs are present in high concentrations in the gastrointestinal tract and extensively metabolized by colonic microbiota. Protein homogenates were prepared from human colonic tissue samples and

Caco-2 colonic epithelial cells. We incubated these samples with equal concentrations of
solanine, chaconine, and solanidine, and assayed for acetylthiocholine hydrolysis. Treatment of
both Caco-2 and colon tissue homogenates with SGAs strongly inhibited ACh hydrolysis (**Fig 5C**).
In contrast, treatment of tissue homogenates with an equal concentration of the aglycone
solanidine only slightly inhibited ACh hydrolysis; in Caco-2 lysates, ACh hydrolysis was
completely unaffected by solanidine. The observed differences in the effect of solanidine on ACh
hydrolysis by Caco-2 lysates and colonic tissue may result from differences in relative BChE
quantities. Indeed, treatment of these samples with the BChE selective inhibitor iso-OMPA
showed more significant contributions of BChE towards ACh hydrolysis in colonic tissue than in
Caco-2 (**Fig S9E**). The modest inhibition of ACh hydrolysis in colonic tissue by solanidine was
rescued by concurrent treatment with iso-OMPA, while ACh inhibition by the glycosides solanidine
and chaconine remained unaffected (**Fig S9F**). These data show that potato SGAs inhibit overall
ACh hydrolysis in relevant host tissues and deglycosylation of these compounds alleviates this
inhibition.

To directly connect microbial SGA metabolism to the activity of host AChE, we used spent
media from individual human stool samples incubated with solanine to modulate AChE activity.
These donor samples were selected to have a range of solanine metabolism (**Fig 3A**), including
samples inactive on solanine and samples that deglycosylated solanine. Solanine metabolism by
different donor samples resulted in significant differences in ACh hydrolysis. Media from inactive
stool samples inhibited AChE while stool samples with extensive aglycone production showing
only slight AChE inhibition (**Fig 5C**). These data show that the inter-individual differences that
exist in microbiota-mediated SGA metabolism can translate to varying levels of host AChE
inhibition *ex vivo*. Taken together, our mapping of the metabolic fates of dietary SGAs revealed
the crucial role of gut microbiota in controlling host exposure to and host enzyme interactions with
these compounds.

382   **DISCUSSION**

Beyond steroidal glycoalkaloids from Solanums, other saponins of plant origin have been
shown to undergo gut microbiome-mediated deglycosylation. Ginsenosides from the root of
Panax ginseng were found to be deglycosylated to the protopanaxadiol aglycone, mediated by
Bacteroides, *Bifidobacterium*, *Fusobacterium*, and *Prevotella* strains[37]. In contrast to our
observations with SGA metabolism, however, ginsenosides do not appear to be solely dependent

on the gut microbiome for metabolism and are also hydrolyzed by host digestive processes[38].
Beyond saponins, other dietary plant glycosides are also known to be metabolized by the gut
microbiota. A notable example are flavonoids, which undergo microbiota-mediated
deglycosylation and extensive metabolism to form phloroglucinol, catechol, and equol, among
other products[39]. The ability of broad specificity host glycosidases to metabolize flavonoids varies
with the identity and linkage of the glycone moiety[40]. In contrast to many plant glycosides
encountered in the diet and potentially resulting from the complexity of the glycone moieties, we
find SGAs from Solanums to be unusually resistant to hydrolysis by host enzymes and digestion.
These results support previous observations that tomatine is stable to incubation in acidic
conditions at $37°C$[41], and is reminiscent of nondigestible dietary fibers that are fermented by gut
commensals[42].

In this work, we have established a direct link between human gut commensals and SGA
metabolism, both in the context of complex communities from human stool and with individual
type strains in culture. Under the conditions used to culture the commensal type strains, we found
that SGA metabolism is performed by a small subset of the strains in a representative community.
While Solanum plant pathogens are known to deglycosylate SGAs (**Supplemental Discussion**)
and clearly benefit from deactivation of these plant defense compounds, the motivation of gut
commensals to deglycosylate SGAs is less apparent. One possible motive for commensal
metabolism of SGAs is access to the liberated sugars as a competitive advantage in a community
context; gut commensals are known to harvest sugars from plant glycosides for use as carbon
sources[43]. Regardless, identification of the strains involved in SGA deglycosylation enables
further investigation into the ecological and genetic bases of this metabolism, understanding of
the prevalence of this metabolism across human populations, and the rational design of synthetic
communities that produce a desired metabolite output[44].

Though SA aglycones are formed from gut microbiota-mediated deglycosylation of dietary
SGAs, we have found that they are transient intermediates both *in vivo* and in the context of a
synthetic bacterial community. Our efforts to characterize the metabolite fates of SGAs have led
to the identification of aglycones that are modified further by hydroxylation, sulfation, 3OH
oxidation, Δ5,6 saturation, and phosphoethanolamine conjugation. There is rich biology
associated with each of these transformations promoted by subsets of gut microbial communities
(**Supplemental Discussion**). A major transformation that we observed in both mouse systemic
circulation and in human urine involved hydrogenation of the Δ5,6 double bond of the potato
aglycone solanidine. A previous report of such a compound noted its formation in bovine rumen

fluid from a solanine input[45]. More broadly, however, this reduction is reminiscent of gut

microbiota-mediated hydrogenation of cholesterol to coprostanol. Bacteria identified to be

coprostanol producers include *Bacteroides* sp. D8[46] and strains from the genus *Eubacterium*[47,48].

In human gut communities, the cholesterol dehydrogenase ismA from uncultured Clostridia has

been implicated in coprostanol formation[49]. Despite the analogous transformations observed for

solanidine and cholesterol, however, we observed solanidine hydrogenation to be performed by

Bacteroidetes, including *Alistipes finegoldii* DSM 17242 (**Fig 4**). A BLAST search revealed no

close homologs for *ismA* homologs in the solanidine reducing strains. As such, we hypothesize

that solanidine hydrogenation proceeds by a route distinct to the one identified for coprostanol

formation.

Crucially, metabolism of dietary SGAs by gut microbiota can transform the bioavailability

and bioactivity of these compounds in the host. Potato SGAs are AChE inhibitors, a class of

compound associated with drastic pharmacological and toxicological impacts. Systemic AChE

inhibition, including irreversible enzyme inhibition by sarin and organophosphates, can be lethal

due to the local accumulation of ACh and continued stimulation of vital nervous processes.

However, reversible and tissue specific AChE inhibition is a clinical strategy for the treatment of

Alzheimer's disease[50], glaucoma[51], and post-operative ileus[52]. The evidence for the toxicological

impacts of potato SGAs is extensive; improper storage of potatoes can result in overaccumulation

of SGAs and has led to numerous reports of SGA-induced poisonings[34], with an estimated toxic

dose of 2-5 mg/kg body weight in humans[53]. Acute symptoms of SGA toxicity commonly involve

nausea, vomiting and diarrhea[54], indicative of effects local to the gastrointestinal tract. Moreover,

treatment with the acetylcholine receptor inhibitor atropine was found to alleviate toxicity following

of solanine administration in mice[55], suggestive of a role for cholinesterase inhibition in mediating

toxicity. We show in this study that the gut microbiota alleviates the AChE inhibitory properties of

potato SGAs solanine and chaconine by deglycosylating the compounds to a non-inhibitory form

(**Fig 5C**). We hypothesize that gut microbiota can significantly diminish the toxicity of these

compounds *in vivo*.

In the colon where we found SGAs to significantly accumulate, acetylcholine is additionally

a key regulator of multiple processes including chloride secretion[56], epithelial proliferation[57], fluid

secretion[58], and intestinal propulsion[59]. Considering the role of ACh in these diverse physiological

processes, we suspect that AChE inhibition by SGAs may impact colonic physiology through a

myriad of mechanisms beyond overt toxicity. While there is still much to be investigated regarding

the contributions of the gut microbiota to mitigating the symptoms of SGA exposure, we show that

gut microbial metabolism can significantly deplete SGA levels in the colon *in vivo* following a

potato dietary intervention (**Fig S5C**). Furthermore, variation in the abilities of individual microbiota

to deglycosylate potato SGAs (**Fig 3A**) can lead to interindividual differences in host exposure to

these bioactive SGAs. Finally, these results emphasize the need for considering gut microbiota

in future studies investigating the physiological impacts of SGAs.

In addition to the depletion of bioactive SGAs by microbiota-mediated deglycosylation, the

products of this metabolism may have distinct physiological effects. For example, dietary

intervention with the aglycone tomatidine was found to increase mTORC1 signaling in mice,

resulting in increased skeletal muscle hypertrophy and decreased adiposity[60]. Further modified

versions of steroidal aglycones produced by host or bacterial metabolism may also result in

differential activity relative to the unmodified aglycone. For example, modification of the bile acid

lithocholic acid by oxidation at the C3 alcohol, a transformation that we also observed with

tomatidine and solanidine, enabled binding of the compound to host RORγt receptors and

resulted in inhibition of T_H17 cell differentiation[61].

Taken together, our characterization of the metabolic fates of SGAs and the role of

microbiota in determining host exposure lays a foundation towards a more complete

understanding of the effects that this abundant class of dietary plant metabolites may have on

host physiology.

MATERIALS AND METHODS

Chemicals

Steroidal alkaloid compounds were acquired from commercial sources: tomatine (Tokyo

Chemical Industry), tomatidine (Cayman Chemical Company), solanine (Sigma Aldrich),

chaconine (Sigma Aldrich), solanidine (Carbosynth), solamargine (MedChemExpress),

solasonine (MedChemExpress), solasodine (Santa Cruz Biotechnology). All steroidal alkaloid

compounds were prepared as 10 mM stocks in vehicle. Steroidal glycoalkaloids were dissolved

in dimethylsulfoxide (DMSO). Tomatidine and solasodine were dissolved in methanol and

solanidine was dissolved in ethanol. Other chemical standards were acquired as follows:

trimethylamine N-oxide (Cayman Chemical Company), trigonelline chloride (Cayman Chemical

Company), and hippuric acid (Sigma Aldrich).

Mouse Studies

Animal studies were performed under a protocol approved by the Stanford University Institutional Animal Care and Use Committee. For all studies, groups of five 6-8 week old female mice were used and allowed to acclimate for a week before any treatment.

Ad libitum fresh tomato feeding

Swiss Webster mice (Taconic) were fed *ad libitum* with AIN93G diet (Dyets, Inc). Three days following the start of AIN93G diet, 40 g of red ripe Microtom tomatoes were provided daily alongside diet for each cage of five mice for 15 days, with uneaten tomatoes removed daily before replenishing with fresh tomatoes. The quantities of tomatine and esculeoside A in the provided tomatoes were determined by liquid chromatography-mass spectrometry (LC-MS) as described below to be about 0.72 μmol/g fresh weight 16sculeoside A and 0.074 μmol/g fresh weight tomatine. Molar quantities of both compounds were estimated using a commercial tomatine standard (Tokyo Chemical Industry). Mice were sacrificed by CO_2 asphyxiation 24 hours following the final provision of tomatoes. Blood was collected by cardiac puncture into Microtainer SST tubes (BD) and serum harvested. All samples were frozen in liquid nitrogen and stored at -80°C until analysis.

Pure tomatine feeding

Swiss Webster mice (Taconic) were fed *ad libitum* with AIN93G diet (Dyets, Inc). Three days following the start of AIN93G diet, mice were provided with 15 μM pure tomatine in drinking water. Drinking water was sterile filtered and refreshed every 2 to 4 days for 15 days, after which mice were sacrificed by CO_2 asphyxiation and serum, cecal contents, colon, liver, kidneys, and mesenteric lymph nodes (MLNs) were harvested. Samples were frozen in liquid nitrogen and stored at -80°C until analysis.

Potato feeding

Swiss Webster mice (Taconic) were fed *ad libitum* with AIN93G diet (Dyets, Inc). One week following the start of AIN93G diet, mice were dosed with 0.5 μmol solanine per mouse by oral gavage. Two days following this single dose, mice were provided with 10% w/w potato peels incorporated into AIN93G diet. Potato-supplemented diet was prepared by autoclaving peels from Yukon Gold potatoes at 121°C for 30 minutes, lyophilizing to dryness, and homogenizing using a blender to generate dry powder. Potato peel powder was mixed into powdered AIN93G diet (Dyets, Inc) at 10% w/w, moistened with water to pellet, and then lyophilized to dryness. The quantity of solanine and chaconine in the provided potato-supplemented diet was

determined by LC-MS as described below to be about 0.4 µmol/g diet chaconine and 0.11
µmol/g diet solanine. Mice that did not receive potato diet instead received similarly pelleted
AIN93G. After one day of potato-supplemented diet, mice were sacrificed by CO_2 asphyxiation
and serum was collected. Samples were frozen in liquid nitrogen and stored at -80°C until
analysis.

Antibiotics treatment for *ex vivo* feces metabolism
Swiss Webster mice (Taconic) were fed ad libitum with AIN93G diet (Dyets, Inc). Three days
following the start of AIN93G diet, mice received five consecutive daily doses of 10 mg each
ampicillin, vancomycin, metronidazole, and neomycin by oral gavage[62]. Each mouse received
0.25 mL of a combined stock of 40 g/L of each antibiotic prepared in phosphate buffered saline
(PBS, pH 7.2). Mice not treated with antibiotics instead received 0.25 mL of PBS by oral
gavage. Following five daily doses of antibiotics by oral gavage, antibiotics were provided *ad*
libitum in drinking water for three days: ampicillin (1 g/L), vancomycin (0.5 g/L), neomycin (1
g/L), and metronidazole (1 g/L)[23]. Feces were collected into sterile tubes and processed as
described below.

Concurrent antibiotics treatment and dietary intervention
C57BL/6J mice (Jackson Laboratory) were fed *ad libitum* with AIN93G diet (Dyets, Inc). One
week following the start of AIN93G diet, antibiotics were provided *ad libitum* in drinking water for
the duration of the study: ampicillin (1 g/L), vancomycin (0.5 g/L), neomycin (1 g/L), and
metronidazole (1 g/L). Antibiotics were refreshed every 3-4 days. Six days following the start of
antibiotics treatment, mice were provided potato-supplemented diet (5% dry weight), prepared
as described above. Another set of mice was provided tomato-supplemented diet (10% dry
weight) seventeen days following the start of antibiotics treatment. Sugar plum tomatoes were
incorporated into AIN93G diet as described for potato supplemented diet above, but surface-
sterilized with 70% ethanol instead of autoclaved. Animals received potato supplemented diet
for 8 days or tomato supplemented diet for 7 days, and were then sacrificed by CO_2
asphyxiation. Serum, colon, livers, cecal contents, kidneys, and MLNs were collected and
frozen in liquid nitrogen and stored at -80°C until analysis.

Sample preparation for LC-MS
Harvested tissues were lyophilized to dryness and then homogenized using a ball mill
homogenizer (Retsch). Homogenized tissues were extracted with a fixed volume of 2:1:1
acetonitrile (MeCN):methanol(MeOH):water(H_2O) per dry mass tissue, vortexed to mix, and then
centrifuged at 18,000 *g* for 10 minutes. The resulting supernatant was optionally diluted (for

cecal content samples) and then filtered through a 0.45 μm PTFE for analysis by LC-MS, as

described below.

Serum samples were thawed and proteins precipitated by the addition of three volumes of ice

chilled 3:1 MeCN:MeOH. Samples were incubated on ice for 10 minutes, and then centrifuged

at 18,000 g for 10 minutes at 4°C. To concentrate the sample, supernatant was transferred to a

new microfuge tube, flash-frozen in liquid nitrogen, and lyophilized to dryness. Dried samples

were resuspended in 2:1:1 MeCN:MeOH:H_2O to the original volume of serum and filtered

through a 0.45 μm PTFE for analysis by LC-MS.

Steroidal alkaloid metabolism by human liver enzymes

To assay for metabolism of steroidal alkaloids by human liver enzymes, 1 mg/mL of liver S9

proteins from pooled human donors (Sekisui XenoTech) was incubated in PBS pH 7.4,

supplemented with 25 μM tomatine, tomatidine, solanine, chaconine, or solanidine. Samples

were incubated at 37°C for 5 minutes. Metabolism was initiated by the addition of cofactors

required for metabolic reactions and supplied as two separate mixtures: 1) 1 mM glutathione, 2

mM NADPH, and 0.5 mM acetyl-CoA, and 2) 200 μM 3'-phosphoadenosine-5'-phosphosulfate

(PAPS), 500 μM uridine diphosphate glucuronic acid (UDPGA), 25 μg/L alamethicin, 2 mM

magnesium chloride, and 2 mM NADPH. Each reaction was prepared in duplicate, with one set

immediately quenched by the addition of three volumes of cold 3:1 MeCN:MeOH. The other set

of reactions was incubated at 37°C for 60-90 minutes and then quenched as described above.

Quenched samples were stored at -80°C overnight, thawed, and centrifuged at 18,000 g for 10

minutes. The resulting supernatant was collected and filtered through 0.45 μm PTFE for

analysis by LC-MS.

Synthesis of standards for steroidal alkaloid metabolites

3-oxotomatidine

Synthesis of 3-oxotomatidine was adapted from previous reports[63]. Tomatidine hydrochloride

was dissolved in dichloromethane (DCM). Three molar equivalents of Dess-Martin periodinane

(DMP, Alfa Aesar) was added and the mixture was stirred at room temperature overnight. The

reaction was quenched with one volume of 0.8 M sodium thiosulfate and saturated sodium

bicarbonate, stirred at room temperature for 1 hour, and then extracted with three subsequent

volumes of DCM. The organic layers were combined and dried on anhydrous calcium sulfate.

LC-MS analysis of the reaction products at this stage generally revealed a conversion of about

50%. To improve substrate conversion, three equivalents of DMP were again added to this

mixture and the work repeated as described above. LC-MS analysis of the crude reaction after a

second round of reaction generally showed little detectable remaining unreacted tomatidine.

To isolate the produced 3-oxotomatidine, crude reaction products in DCM were dried down,

resuspended in a small volume of N,N-dimethylformamide (DMF), and separated by reversed

phase chromatography using a 100Å, 30 µm, 12 g Sfär C18 D Duo column (Biotage). Water

with 0.1% (v/v) formic acid and acetonitrile with 0.1% (v/v) formic acid were used as mobile

phase solvents with a flow rate of 12 mL/min. Chromatographic separation was achieved using

the following linear gradient (with percentages indicating levels of acetonitrile with formic acid):

10%, 3 column volumes (CV); 10% to 20%, 3 CV; 20% to 50%, 25 CV; 50% to 95%, 5 CV; 95%,

10 CV. Fractions were assayed by LC-MS; fractions containing pure 3-oxotomatidine were

combined and evaporated to dryness.

<u>Tomatidine sulfate</u>

Tomatidine sulfate synthesis was adapted from previous reports for the sulfation of dietary

phenolics[64]. Tomatidine hydrochloride was dissolved in pyridine and combined with one molar

equivalent of sulfur trioxide pyridine complex. The mixture was sparged with argon and stirred at

65°C for 24 hours. The reaction was quenched with one volume of water and the organic

solvent evaporated. The crude reaction mixture was analyzed by LC-MS for retention time

match and MS/MS match to the mass feature in biological samples.

<u>Dihydrosolanidine</u>

Dihydrosolanidine was synthesized from solanidine (Carbosynth) by hydrogenation mediated by

a palladium on carbon catalyst. Solanidine was dissolved in ethanol and one mass equivalent of

palladium of carbon was added with stirring. Hydrogen gas was bubbled through the mixture for

6 hours and the reaction was filtered to remove catalyst. The crude reaction mixture was

analyzed by LC-MS for retention time match and MS/MS match to the sample mass feature.

Human urine from colectomy patients

Urine samples were collected from 12 patients with total colectomies (6 male and 6 female) and

17 patients with intact colons with no known gastrointestinal disease and no antibiotic use for

one month pre-collection (11 male and 6 female)[24]. Donors did not receive specified dietary

regimens or supplements.

Urine samples were centrifuged at 12,000 g for 10 minutes and then analyzed by LC-MS as

described below. Sample creatinine levels were measured using a colorimetric creatinine assay

kit (Abcam) and normalization factors were calculated using the Cockcroft-Gault equation[65]. Ion

counts for metabolites of interest were multiplied by the normalization factor to enable

comparison in compound abundance between samples.

Anaerobic culturing

Work with fecal samples and individual commensal strains was performed under anaerobic

conditions using an anaerobic chamber (Coy, 10% CO_2, 5% H_2). To maintain a reducing

environment for fecal bacteria during sample collection and handling, mouse and human feces

were suspended in pre-reduced transport medium[26]. Transport medium contained, per L: 0.2 g

KCl, 0.1 g $CaCl_2$, 0.1 g $MgCl_2$, 0.2 g KH_2PO_4, 1.15 g Na_2HPO_4, 3 g NaCl, 0.1 g uric acid, 0.1 g

glutathione. Human fecal samples were cultured in standard amino acid complete minimal

medium (SAAC), prepared as described by Dodd *et al*[66]. Human commensal type strains were

grown on either Brain Heart Infusion (Teknova) agar plates supplemented with 5 µg/mL hemin

and 0.5 µg/mL menadione (BHIS) or Columbia agar plates supplemented with 10% sheep blood

(Col-B). Liquid cultures were grown in Mega medium, a rich medium adapted from Han *et al*[67];

for our use, Mega medium was modified to exclude soluble starch and Tween 80 for combability

with downstream LC-MS analysis.

Ex vivo **steroidal alkaloid metabolism by mouse feces**

To assay for steroidal glycoalkaloid metabolism by murine fecal bacteria, freshly collected fecal

pellets were suspended at 2% w/v in pre-reduced transport medium and then vortexed until

homogenized. Larger particulates were allowed to settle, and the slurry was transferred to a

new tube. An equal volume of transport medium containing 40 µM tomatine was added to the

2% w/v fecal slurry, resulting in a final incubation of 1% w/v feces with 20 µM tomatine. Samples

were incubated anaerobically for 36 hours at 37°C. Following incubation, two volumes of 3:1

MeCN:MeOH were added to extract metabolites and quench metabolism. Samples were

vortexed, centrifuged at 12,000 *g* for 10 minutes, and then supernatant was diluted and filtered

through a 0.45 µm PTFE filter for analysis by LC-MS, as described below.

***Ex vivo* human stool metabolism**

De-identified stool samples from human donors were collected as part of the RAMP study

(Rejuvenating the Aging Microbiota with Prebiotics, clinical trial ID: NCT03690999)[25]. This

cohort was comprised of 89 individuals (49 female and 40 male) ranging from 60 years to 83.9

years. A subset of feces samples from thirty donors prior to any dietary intervention were

randomly selected and assayed for SGA metabolism. Samples were stored at -20°C in donor's

home freezers until they were transferred to the research laboratory and stored at -80°C until

use. Frozen samples were transferred into an anaerobic chamber, suspended at 10% w/v in

pre-reduced transport medium with 20% glycerol, vortexed for 2 minutes to homogenize, and

then allowed to settle. The stool slurry was aliquoted into cryotubes, flash-frozen in liquid

nitrogen, and stored at -80°C until use. For use in metabolism assays, frozen stool

resuspensions were thawed, diluted to 1% v/w in pre-reduced transport medium, and combined

in 96 well plate format with an equal volume of 2x SAAC containing 40 µM of steroidal alkaloid

substrate. Each reaction contained final concentrations of 0.5% w/v stool sample, 1x SAAC, and

20 µM steroidal alkaloid substrate. Each combination of stool sample and substrate was

prepared in two sets of triplicates; one set was incubated anaerobically at 37°C for 48 hours

while another set was immediately frozen in liquid nitrogen as an unmetabolized control and

stored at -80°C. Following incubation for 48 hours, the other set of reactions was also frozen in

liquid nitrogen and stored at -80°C.

To mitigate effects from differential evaporation between wells, samples were lyophilized to

dryness in 96 well plates and then resuspended in three volumes of 2:1:1 MeCN:MeOH:H_2O.

Plates were vortexed and sonicated for 5 minutes to resuspend, and then stored at -80°C

overnight to precipitate any proteins. Samples were centrifuged at 3,000 *g* for 10 minutes,

diluted in resuspension solvent, and then filtered through a 0.45 µm PTFE filter for analysis by

LC-MS, as described below.

Human commensal type strain culturing and metabolism

Human commensal type strains were streaked from glycerol stocks onto Columbia blood agar plates and incubated anaerobically at 37°C for about 2 days. Single colonies were inoculated into Mega medium and grown anaerobically at 37°C. After 2 days growth, strains were subcultured into fresh Mega medium and further incubated for 48 hours. Individual cultures were pelleted by centrifugation at 6000 g for 10 minutes, and then resuspended in pre-reduced PBS. Resuspended cultures were used to inoculate modified Mega medium (without Tween and soluble starch) supplemented with 25 µM steroidal alkaloid substrate and grown anaerobically at 37°C for 2 days. After incubation, cultures were flash-frozen in liquid nitrogen and lyophilized to dryness to mitigate the effects of evaporation. Dried reactions were resuspended in three times the original reaction volume with 2:1:1 MeCN:MeOH:H_2O, vortexed and sonicated to resuspend, and then frozen at -80°C. Prior to analysis by LC-MS, samples were centrifuged at 4000 g for 15 minutes to remove particulates, diluted, and then filtered through a 0.45 µm PTFE filter.

Phylogenetic analysis of human commensal type strains

Taxonomic classifications of the human commensal type strains were determined using the GTDBtk v.1.6.0 classify workflow which uses 120 marker genes from bacterial whole genome sequences to place strains of interest in a GTDBtk reference tree (release202)[68]. Bacterial genomes were obtained from the NIH Human Microbiome Project[69]. Trees were visualized using Geneious.

Metabolism of steroidal alkaloids by commensal cocultures

Bacteroides and *Roseburia* strains were streaked from glycerol stocks onto BHIS plates and incubated anaerobically at 37°C for one day. Single colonies were inoculated into pre-reduced Mega medium and grown anaerobically at 37°C for about one day. Liquid cultures were subcultured once more into Mega medium and allowed to grow anaerobically for a day. An equal volume of each culture was combined, pelleted by centrifugation at 6,000 g for 10 minutes, and resuspended in one volume of fresh Mega medium supplemented with 20 µM tomatine substrate. Cocultures were incubated anaerobically at 37°C for three days. Samples were frozen at -80°C and lyophilized to dryness. Dried samples were then extracted with three

volumes of 2:1:1 MeCN:MeOH:H_2O, vortexed and sonicated to resuspend, and centrifuged at
4000 g for 15 minutes and filtered through a 0.45 μm PTFE filter prior to LC-MS analysis.

In vitro cell viability assays

The human intestinal epithelial cell line Caco-2 (ATCC, provided by Sarah Heilshorn, Stanford)
and used between passages 5 to 15. Cells were grown and maintained in DMEM (ATCC)
supplemented with 10% fetal bovine serum, 1% nonessential amino acids, 100 U/mL penicillin,
and 100 μg/mL streptomycin at 37°C in 5% CO_2. Cells were seeded into 96 well plates at a
density of 10,000 cells per well and incubated for 17 hours for cells to adhere. Cells were then
treated with varying concentrations of steroidal alkaloid compounds dissolved in solvent vehicle.
The total amount of vehicle added to each well was normalized to 0.4% v/v. As a nontreated
control, cells were treated with vehicle alone. Cells were incubated with compound at 37°C for
24 hours, after which cell viability was determined using the CellTiter-Glo Luminescent Cell
Viability Assay (Promega). All treatments were performed in triplicate and repeated in three
separate experiments.

Caco-2 permeability assay

Transport of steroidal alkaloid compounds _in vitro_ were determined using differentiated Caco-2
cells as a model monolayer[27]. Cells were seeded into 12 well plates fitted with 0.4 μm
polycarbonate Transwell inserts (Corning) at a density of 300,000 cells per insert. The apical
compartment above the insert was filled with 0.5 mL of medium and the basolateral
compartment below the insert was filled with 1.5 mL of medium. Medium was refreshed on both
the apical and basolateral sides every 2 to 3 days for 21 days for monolayer formation. After 21
days, cells were rinsed with serum free DMEM supplemented with 1% nonessential amino
acids, 100 U/mL penicillin, and 100 μg/mL streptomycin. The apical compartment was filled with
serum-free DMEM additionally supplemented with nontoxic concentrations of steroidal alkaloid
compounds, as determined by viability assays described above (solanine: 10 μM, chaconine: 3
μM, solanidine: 10 μM). A subset of wells was treated with vehicle only as a control. The
basolateral compartment was filled with serum-free DMEM supplemented with 4% bovine serum
albumin (BSA) to mimic physiological protein concentrations on either side of the intestinal
epithelium. After 17 hours of incubation with steroidal alkaloid compounds, medium was

collected from both the apical and basolateral components. An aliquot of medium was assayed
for extracellular LDH using the LDH-Glo Cytotoxicity Assay (Promega) to ensure that cell
monolayers were not compromised by compound treatment. All compound-treated wells
showed similar LDH levels to vehicle only treatment, indicating that compound treatments at the
selected concentrations did not impact cell viability. The remainder of the medium collected was
assayed for steroidal alkaloid content using LC-MS. To prepare medium for analysis, samples
were diluted with nine volumes of 3:1 MeCN:MeOH, kept at -20°C overnight to precipitate
proteins, centrifuged at 10,000 g for 10 minutes, and filtered through a 0.45 μm PTFE filter.

Cholinesterase activity assay

Acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) activities were measured using
Ellman's method, adapted to a 96-well format[35]. For acetylcholinesterase activity, 5 ng of
recombinant human AChE (Sigma) or 30 μg of tissue lysate was incubated with steroidal
alkaloid compound or at room temperature for 30 minutes in 50 mM Tris pH 8.0 in a total
volume of 90 μL. To initiate reactions, 5 μL of 10 mM acetylthiocholine iodide and 5 μL of 10
mM 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB) were added to each well and absorbance at 405
nm was measured over the course of 30 minutes. BChE activity was analogously measured
using 5 ng of recombinant human BChE (R&D Systems) and butyrylthiocholine iodide. Enzyme
activity was quantified as the linear slope of absorbance over time. Relative inhibition by
steroidal alkaloid compounds was quantified by normalization to enzyme activity in the presence
of vehicle.

Colon lysate preparation

Normal human colon tissue from three de-identified human donors was obtained from the
Stanford Tissue Bank. About 100 mg of tissue was homogenized in 2 mL of tissue lysis buffer
(50 mM Tris pH 8, 150 mM sodium chloride, 0.5% Triton X-100, 10 μg/mL leupeptin, 10 μg/mL
aprotinin, 20 μM pepstatin, and 2.5 mM benzamidine hydrochloride) using a gentleMACs Tissue
Dissociator (Miltenyi). Homogenized samples were centrifuged at 2,000 g for 5 minutes at 4°C
to collect lysate, transferred to new centrifuge tubes, and centrifuged at 9,000 g for 20 minutes
at 4°C to pellet cell debris.

To prepare protein lysates from Caco-2 intestinal epithelial cells, cells were maintained as

described above. A confluent 10 cm plate was rinsed three times with cold PBS, and then lysed

by adding tissue lysis buffer and incubating with rocking at 4°C for 20 minutes. Lysate was

collected and centrifuged at 9,000 g at 4°C for 20 minutes to remove cell debris.

To the resulting supernatants from human tissue and Caco-2 cells, glycerol was added to the

resulting supernatant for a final concentration of 20%. Protein concentrations in lysates were

estimated using BSA standards and a detergent-compatible Bradford assay (Abcam). Lysates

were aliquoted, flash-frozen in liquid nitrogen, and stored at -80°C until analysis by Ellman's

assay as described above. To assay the contributions of AChE to acetylthiocholine hydrolysis in

these homogenates, BChE-specific inhibitor iso-OMPA was added to homogenates at a

concentration of 500 µM and incubated at room temperature for 30 minutes before the assay

was initiated with acetylthiocholine and DTNB.

Acetylcholinesterase inhibition by human stool samples

A subset of eleven of the previously assayed de-identified human stool samples was selected

for their differences in solanine metabolism, including samples that did not appear to metabolize

solanine (02, 17, 22, 28), samples that deglycosylated solanine (07, 13, 23, 30), and samples

that appeared to partially deglycosylate solanine by removing a glucose moiety from the glycone

(08, 15, 16). *Ex vivo* incubations with 20 µM solanine or DMSO vehicle were prepared as

described above for each of the donor samples. For compatibility with Ellman's assay, we

minimized the presence of free thiols in the assay medium; these *ex vivo* stool incubations were

performed in pre-reduced SAAC minimal medium formulated without cysteine. Samples were

incubated anaerobically at 37°C for 48 hours, after which an aliquot was removed for LC-MS

analysis. This aliquot was diluted in 3:1 MeCN:MeOH to precipitate proteins, centrifuged at

12,000 *g* for 10 minutes, diluted in 2:1:1 MeCN:MeOH:H_2O, and then filtered through a 0.45 µm

PTFE filter. The remainder of the samples was spun down at 12,000 *g* for 20 minutes and the

resulting supernatant used to inhibit recombinant human AChE as described above, with a few

adaptations. These AChE assays were performed in spent supernatant from human stool

cultures instead of 50 mM Tris pH 8. Metabolism by stool samples appeared to lower media pH;

to adjust the assay pH to levels amenable to AChE activity and DTNB-thiol conjugation, equal

volumes of sodium hydroxide was added to samples at a final concentration of 6 mM. Sodium

hydroxide was not added to media-only controls (i.e. sterile).

LC-MS

Ad libitum fresh tomato feeding and pure tomatine feeding

Mouse serum and tissues resulting from fresh tomato supplementation and pure tomatine feeding were analyzed on an Agilent 1260 Infinity II HPLC using both reversed-phase liquid chromatography (LC) and normal phase LC. Reversed-phase LC was conducted using a Gemini NX-C18 2x100 mm, 5 µm column (Phenomenex) at 40°C. Mobile phase solvents were water with 0.1% (v/v) formic acid and acetonitrile with 0.1% (v/v) formic acid at a flow rate of 0.4 mL/min. A volume of 3 µL of sample was injected. Chromatographic separation was achieved using the following linear gradient (with percentages indicating the levels of acetonitrile with formic acid): 3% to 97%, 20 minutes; 97%, 5 minutes; 97% to 3%, 1 minute; 3%, 5 minutes. Normal phase liquid chromatography was conducted using a Poroshell 120 HILIC-Z, 2.1x100 mm, 2.7 µm column (Agilent) at 40°C. Mobile phase solvents were 10 mM ammonium formate supplemented with 0.1% formic acid and 90% acetonitrile with 10 mM ammonium formate supplemented with 0.1% formic acid at a flow rate of 0.25 mL/ min. A volume of 2 µL of sample was injected. Chromatographic separation was achieved using the following linear gradient (with percentages indicating the levels of acetonitrile-containing mobile phase): 100% to 60%, 15 minutes; 60%, 2 minutes; 60% to 100%, 1 minute; 100%, 8 minutes. For both types of chromatography, coupled mass spectrometry (MS) data was collected with an Agilent 6545 quadrupole time-of-flight (qTOF) ESI mass spectrometer in positive polarity. Parameters for the 6545 qTOF MS were as follows: mass range: 50-1700 m/z, gas temperature: 250°C, nebulizer: 10 psi, drying gas flow rate: 12 L/min, sheath gas temperature: 300°C, sheath gas flow rate: 12 L/min, fragmentor: 100 V, skimmer: 50 V.

Human urine

Urine from human colectomy patients and donors with intact colons were chromatographically separated on an Agilent 1260 Infinity II HPLC using both reversed-phase LC. LC was conducted using a SB-C18 3.0x100 mm, 1.8 µm column (Agilent) at 40°C. Mobile phase solvents were water with 0.1% (v/v) formic acid and acetonitrile with 0.1% (v/v) formic acid at a flow rate of 0.4 mL/min. A volume of 5 µL of undiluted sample was injected. Chromatographic separation was achieved using the following linear gradient (with percentages indicating the levels of acetonitrile with formic acid): 3% to 97%, 20 minutes; 97%, 4 minutes; 97% to 3%, 1 minute; 3%, 4 minutes. Coupled mass spectrometry (MS) data was collected with an Agilent 6545 quadrupole time-of-flight (qTOF) ESI mass spectrometer in positive polarity. Parameters for the 6545 qTOF MS were as follows: mass range: 50-1700 m/z, gas temperature: 250°C, nebulizer: 60 psi, drying

gas flow rate: 6 L/min, sheath gas temperature: 250°C, sheath gas flow rate: 11 L/min,

fragmentor: 140 V, skimmer: 65 V.

All other studies

Samples from all other studies were separated on an Agilent 1290 Infinity II HPLC using

reversed phase (LC). Chromatography for mouse serum and tissue samples and human liver

enzyme incubations was performed using a Zorbax RRHT Eclipse Plus C18, 2.1x100 mm, 1.8

µm column (Agilent) at 40°C. Mobile phase solvents were water with 0.1% (v/v) formic acid and

acetonitrile with 0.1% (v/v) formic acid at a flow rate of 0.4 mL/min. Chromatographic separation

was achieved using the following linear gradient (with percentages indicating the levels of

acetonitrile with formic acid): 3% to 97%, 20 minutes; 97%, 5 minutes; 97% to 3%, 1 minute;

3%, 5 minutes. Chromatography for human stool incubations, commensal type strain cultures,

bacterial cocultures, and Caco-2 monolayer transport studies was performed using a Zorbax

RRHT Eclipse Plus C18, 2.1x50 mm, 1.8 µm column (Agilent) at 40°C. Mobile phase solvents

were water with 0.1% (v/v) formic acid and acetonitrile with 0.1% (v/v) formic acid at a flow rate

of 0.6 mL/min. Chromatographic separation was achieved using the following linear gradient

(with percentages indicating the levels of acetonitrile with formic acid): 3% to 60%: 6 minutes,

60% to 95%: 0.1 minutes, 95%: 1.5 minutes, 95% to 3%: 0.1 minutes, 3%: 1 minute. For all

samples, a volume of 1 µL of sample was injected. MS data was collected using an Agilent 6546

qTOF ESI mass spectrometer in positive polarity. Parameters for the 6546 qTOF MS were as

follows: mass range: 50-1700 m/z, gas temperature: 325°C, nebulizer: 35 psi, drying gas flow

rate: 10 L/min, sheath gas temperature: 400°C, sheath gas flow rate: 12 L/min, fragmentor: 140

V, skimmer: 45 V.

Data analysis: untargeted metabolomics

Untargeted metabolomics was performed using XCMS Online[70]. Mass features were detected

using the centWave algorithm, with the following parameters:

Maximal tolerated m/z deviation: 30 ppm

Chromatographic peak width [min, max]: [10 seconds, 60 seconds]

Signal to noise threshold: 10

Minimum m/z difference in features with overlapping retention times: 0.01

Retention time correction was performed using the obiwarp method, with a step size of m/z 0.5.

Peak alignment was performed with bandwidth of 3 seconds and minimum fraction (minfrac) of

samples necessary for a valid group of 0.5.

From the table of mass features generated by XCMS Online, filtering criteria were applied to

identify features present or differentially accumulating in certain treatment groups. The first

filtering criteria was retention time; all mass features that eluted before 1 minute were assumed

to not be retained on the column and therefore discarded. Feature peak areas that were

statistically significantly different between treatment groups were then identified by Student's t-

test ($p < 0.1$ for normal phase analysis and $p < 0.05$ for reverse phase analysis). Features that

were higher in the treatment group of interest (e.g. higher in the serum of tomato-fed mice than

in the serum of un-supplemented mice) were identified by fold change difference (ratio of

average peak areas between groups). Mass features above a threshold signal (peak height >

10^4) were then retained. Finally, putative parent ions were identified by retention time and peak

area clustering to remove adducts, potential fragment ions, dimers, and isotopologues.

<u>Data analysis: semi-targeted analysis for steroidal alkaloids in complex samples</u>

To search for compounds in human stool incubations containing steroidal alkaloid

substructures, one set of replicates was analyzed with a constant collision energy of 40V (for

tomatidine-derived compounds) or 60V (for solanidine- and solasodine-derived compounds).

Ions characteristic of tomatidine, solanidine, and solasodine fragmentation at these collision

energies were extracted; tomatidine-derived: $[M+H]^+$ 161.1325, solanidine-derived: $[M+H]^+$

98.0964, and solasodine-derived: $[M+H]^+$ 157.1012. To find the unfragmented mass features

that yielded these characteristic fragments, the same sample was run with a collision energy of

0V and mass features that accumulated at the retention time of interest after stool sample

incubation were identified for further characterization by MS/MS.

<u>Data analysis: targeted analysis</u>

Extracted ion chromatograms (EIC) were extracted from raw data files using MassHunter

Qualitative Data Analysis software and integrated using MassHunter Quantitative Data Analysis

software. EICs were extracted using a 20-40 ppm window centered on the exact m/z value

(**Supplemental Table 1**).

Statistics

Statistical analyses were performed using Prism 9 (GraphPad). Pairs of sample groups were

analyzed using two-sided Welch's t test to account for unequal variance. Multiple groups of

samples were analyzed using a one-way ANOVA with multiplicity-adjusted P values calculated

using Tukey's multiple comparison test. Gaussian distribution was assumed.

DATA AVAILABILITY

All data in this manuscript are available upon request. Any materials generated within this manuscript will be made available, as possible.

ACKNOWLEDGEMENTS

This work was supported by a USDA NIFA grant 2022-67017-36242 (to E.S.S.). C.S.L. was supported by a Bio-X Bowes Fellowship. We thank J.L. Sonnenburg (Stanford) and T.W. Meyer (Stanford) for generously providing human samples for analysis; M. Carter for assistance accessing human stool samples; J.Z. Long and J.T. Kim (Stanford) for helpful discussions and preliminary experiments; E.A. Schulman (Stanford) for assistance with preliminary experiments; A.G. Cheng (Stanford) for helpful discussions and sharing protocols; J. Arreola (Stanford) for assistance with strain growth; S. Jain (C.Z. Biohub) for assistance with phylogenetic analyses; D.L. Wengier (Stanford) for assistance with acquiring tomatoes; J.C.T. Liu (Stanford) for assistance with structural characterization and comments on this manuscript; J.E. Blum and S.P. Niehs for comments on this manuscript.

CONTRIBUTIONS

C.S.L. and E.S.S designed experiments and analyzed data. C.S.L performed all experiments in the manuscript. M.I. and P.P.M. helped prepare human commensal strains. S.K.H. and M.J. assisted with mouse studies. J.R. assisted with human urine metabolomics analysis. C.S.L. wrote the paper with edits from E.S.S., M.A.F., and J.R.

DECLARATION OF INTERESTS

M.A.F. is a cofounder and director of Kelonia, a cofounder of Revolution Medicines, and a member of the scientific advisory boards of the Chan Zuckerberg Initiative and NGM Bio. The remaining authors declare no competing interests.

FIGURES

Figure 1. Major molecular inputs from dietary Solanums

A) Major mass features in mouse serum resulting from tomato intervention (schematic above)
identified using untargeted metabolomics in positive ionization mode with reversed-phase (C:
C18) and normal phase (H: HILIC) chromatography. Abundances for each individual (n=5
mice per treatment) are shown in the heat map as extracted ion chromatogram (EIC) peak
areas normalized to the mean peak area for that mass feature across all individuals. Mass
features also observed in the tomato supplement are marked with a red circle. Putative
compound assignments are provided, with SA denoting putative steroidal alkaloid (SA)
compounds. Assignments with an asterisk (*) have been confirmed by retention time and
MS/MS matches to a chemical standard.

B) The major steroidal glycoalkaloids (SGAs) present in tomato and potato. Glycone monomers
are colored by identity (blue: glucose, yellow: galactose, green: xylose, pink: rhamnose)
C) Accumulation of SGA-derived metabolites in mouse tissue following dietary intervention with
purified tomatine. Metabolite abundances are represented by the mean EIC peak area for the
compound (n=5 mice), normalized to the maximum peak area of that mass feature across
different tissues. Peak areas were normalized by dry mass to enable comparison between
different tissue types. The maximum EIC peak area across tissues is shown in the bars on
the right, with molar quantities in pmol/mg dry weight shown for select compounds. Structures
that have been characterized by comparison to a chemical standard or reproduced by *in vitro*
enzyme reactions are named. Other structures are hypothesized based on *m/z* and MS/MS
spectra; the positions of hydroxylations are not known. For molecules with multiple detected
isomers, the most abundant isomer is shown.
D) Major mass features in mouse serum (n=5 mice per treatment) resulting from dietary potato
intervention for 1 day, identified using untargeted metabolomics in positive ionization mode
with C18 chromatography. Mass features also observed in the potato supplement are marked
with a brown circle.

Figure 2. Gut microbiota gate host exposure to steroidal alkaloid aglycones

A) Tomatine and tomatidine concentrations in serum (nM), cecal contents (pmol/mg fresh
weight), and liver (pmol/mg fresh weight) following a 10% tomato dietary intervention in
C57BL/6J mice receiving antibiotics (Abx). Values shown are the mean±SD for each group,
with individual replicates overlaid (n=5 mice per treatment). P values are determined using a
two-sided Welch's T test. ns = not significant (P > 0.05).
B) Tomatine-derived metabolites in tissues following dietary intervention with 10% tomato diet in
C57BL/6J mice receiving antibiotics. Abundances for each metabolite are normalized to the
maximum EIC peak area across tissues shown in bars. Peak areas were normalized by
sample dry mass to enable comparison between different tissue types. Values shown are the
mean for each treatment group (n=5 mice per treatment). Structures as drawn are
hypothesized based on *m/z* and MS/MS spectra; the position of hydroxylation is not known.
For molecules with multiple detected isomers, the most abundant isomer is shown.
C) Chaconine, solanine, and solanidine concentrations in serum (nM), cecal contents (pmol/mg
fresh weight), and liver (pmol/mg fresh weight) following a 5% potato dietary intervention in
C57BL/6J mice receiving antibiotics. Values shown are the mean±SD for each treatment
group, with individual replicates overlaid (n=5 mice per treatment). P values are determined
using a two-sided Welch's T test. ns = not significant (P > 0.05).

D) Solanidine and chaconine in human urine from colectomy patients (n=12 donors) and donors
with intact colons (n=17 donors). Compound abundance is shown as EIC peak area
normalized to the creatinine content in that sample. Values shown are for individual donor
samples, with the molar concentration of solanidine in one intact colon sample shown in pmol
solanidine per mg creatinine.

Figure 3. Inter-individual variation in *ex vivo* glycoalkaloid metabolism

A) Deglycosylation of tomato, potato, and eggplant SGAs by thirty human stool samples after incubation with substrate (25 µM) in SAAC minimal medium. Data shown are the change in mean EIC peak area of the tracked metabolite before and after 48 hours of incubation (n=3 replicate incubations per donor, substrate, and time point combination). Values are normalized to the maximum change registered across all donors. Total aglycone EIC peak areas were estimated as the sum of EIC peak areas of all tracked aglycones. A sterile sample without added human stool was included as a negative control.

B) Modification of tomato, potato, and eggplant steroidal alkaloid aglycones by human stool samples after incubation with substrate (25 µM) in SAAC minimal medium. Putative structures for modified aglycones are shown to the left. Donor samples are ordered according to deglycosylation activity shown in panel A. Data shown are the change in mean EIC peak area of the tracked metabolite before and after 48 hours of incubation (n=3 replicate incubations per donor, substrate, and time point combination).

Figure 4. SGA metabolism by human commensal type strains

Metabolism of tomato and potato steroidal alkaloids by human commensal type strains after 48 hours of growth in rich Mega medium supplemented with steroidal alkaloid substrate (25 µM). A

phylogenetic tree of the surveyed strains is shown on the left, with boxes colored according to
phylum. Metabolite abundances are represented as the EIC peak area following incubation scaled
to the maximum EIC peak area of any sample incubated with the same substrate, values of which
are shown in grey bars at the bottom of the figure. Data are colored according to strain phylum.
Metabolite modifications: -2H: oxidation of 3OH, +2H: Δ5,6 saturation, +PEA:
phosphoethanolamine conjugation. Data shown are representative of two experiments.

Figure 5. Microbiota-mediated SGA metabolism impacts compound bioavailability and bioactivity

A) Total steroidal alkaloid concentration in the serum of C57BL/6J mice fed potato-supplemented diet and treated with antibiotics. Aglycone concentrations were estimated using a solanidine standard curve. Data shown are the mean±SD (n=5 mice). P values were determined using a two-sided Welch's T test.

B) *In vitro* inhibition of recombinant human AChE activity by potato steroidal alkaloids. Data shown are normalized to AChE treated with vehicle only (DMSO for solanine, chaconine and

ethanol for solanidine) with a total vehicle concentration of 1% for all treatments. Values
shown are the mean±SD (n=3 assays per concentration). 5 ng of AChE was used in each
reaction.

C) *In vitro* inhibition of cholinesterase activity by 50 µM potato SAs in human protein
homogenates from normal colonic tissue (n=3 donors) and Caco-2 colonic epithelial cells.
Activity was normalized to the activity of the vehicle-treated sample. Values shown are the
mean±SD of three replicate assays per homogenate. 5 ng each of recombinant human AChE
and BChE were used for pure protein reactions, while 35 µg total protein was used for each
homogenate. Multiplicity-adjusted P values were calculated using Tukey's multiple
comparison tests.

D) Inhibition of recombinant human AChE activity by spent media from human stool (n=11
donors) incubated with 20 µM solanine in SAAC minimal medium. Activity was normalized to
the activity of AChE treated with spent media from the same donor sample incubated with
vehicle only. Values shown are the mean±SD of three AChE activity assays for each donor
and compound combination. The relative abundances of select metabolites following
incubation with solanine are shown below for each stool sample. Values shown are EIC peak
areas normalized to the maximum peak area across all samples. P values were determined
using a two-sided Welch's T test. Ns: not significant with P>0.05.

1026

REFERENCES

1. Murphy, M. M. *et al.* Phytonutrient Intake by Adults in the United States in Relation to Fruit and Vegetable Consumption. *J. Acad. Nutr. Diet.* **112**, 222–229 (2012).
2. Tennant, D. R., Davidson, J. & Day, A. J. Phytonutrient intakes in relation to European fruit and vegetable consumption patterns observed in different food surveys. *Br. J. Nutr.* **112**, 1214–1225 (2014).
3. Butelli, E. *et al.* Enrichment of tomato fruit with health-promoting anthocyanins by expression of select transcription factors. *Nat. Biotechnol.* **26**, 1301–1308 (2008).
4. Fahey, J. W., Zhang, Y. & Talalay, P. Broccoli sprouts: An exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 10367–10372 (1997).
5. Turnbaugh, P. J. *et al.* An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* **444**, 1027–1031 (2006).
6. Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. & Goodman, A. L. Mapping human microbiome drug metabolism by gut bacteria and their genes. *Nature* **570**, 462–467 (2019).
7. Marcobal, A. *et al.* A metabolomic view of how the human gut microbiota impacts the host metabolome using humanized and gnotobiotic mice. *ISME J.* **7**, 1933–1943 (2013).
8. Fuhrman, B. J. *et al.* Equol Status Modifies the Association of Soy Intake and Mammographic Density in a Sample of Postmenopausal Women. *Cancer Epidemiol. Biomarkers Prev.* **17**, 33–42 (2008).
9. Koeth, R. A. *et al.* Intestinal microbiota metabolism of l-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat. Med.* **19**, 576–585 (2013).
10. Economic Research Service (ERS), U.S. Department of Agriculture (USDA). Food Availability (Per Capita) Data System.
11. Chaudhary, P., Sharma, A., Singh, B. & Nagpal, A. K. Bioactivities of phytochemicals present in tomato. *J. Food Sci. Technol.* **55**, 2833–2849 (2018).
12. Schieber, A. & Saldaña, M. Potato Peels: A Source of Nutritionally and Pharmacologically Interesting Compounds – A Review. *Food* **3**, 23–29 (2009).
13. Agarwal, S. & Rao, A. V. Tomato lycopene and low density lipoprotein oxidation: A human dietary intervention study. *Lipids* **33**, 981–984 (1998).
14. Sesso, H. D., Liu, S., Gaziano, J. M. & Buring, J. E. Dietary Lycopene, Tomato-Based Food Products and Cardiovascular Disease in Women. *J. Nutr.* **133**, 2336–2341 (2003).
15. Simó, C. & García-Cañas, V. Dietary bioactive ingredients to modulate the gut microbiota-derived metabolite TMAO. New opportunities for functional food development. *Food Funct.* **11**, 6745–6776 (2020).
16. Zhu, W. *et al.* Gut Microbial Metabolite TMAO Enhances Platelet Hyperreactivity and Thrombosis Risk. *Cell* **165**, 111–124 (2016).
17. Norred, W., Nishie, K. & Osman, S. Excretion, distribution and metabolic fate of 3H-alpha-chaconine. *Res. Commun. Chem. Pathol. Pharmacol.* **13**, (1976).
18. Krupp, D., Doberstein, N., Shi, L. & Remer, T. Hippuric Acid in 24-Hour Urine Collections Is a Potential Biomarker for Fruit and Vegetable Consumption in Healthy Children and Adolescents,. *J. Nutr.* **142**, 1314–1320 (2012).
19. Hövelmann, Y., Lewin, L., Steinert, K., Hübner, F. & Humpf, H. Mass Spectrometry–Based Analysis of Urinary Biomarkers for Dietary Tomato Intake. *Mol. Nutr. Food Res.* **64**, 2000011 (2020).
20. Cichon, M. J. *et al.* Plasma Metabolomics Reveals Steroidal Alkaloids as Novel Biomarkers of Tomato Intake in Mice. *Mol. Nutr. Food Res.* **61**, 1–9 (2017).

21. Dzakovich, M. P. *et al.* Transcriptomics and Metabolomics Reveal Tomato Consumption Alters Hepatic Xenobiotic Metabolism and Induces Steroidal Alkaloid Metabolite Accumulation in Mice. *Mol. Nutr. Food Res.* **68**, 2300239 (2024).

22. Cockburn, D. W. & Koropatkin, N. M. Polysaccharide Degradation by the Intestinal Microbiota and Its Influence on Human Health and Disease. *J. Mol. Biol.* **428**, 3230–3252 (2016).

23. Rakoff-Nahoum, S., Paglino, J., Eslami-Varzaneh, F., Edberg, S. & Medzhitov, R. Recognition of Commensal Microflora by Toll-Like Receptors Is Required for Intestinal Homeostasis. *Cell* **118**, 229–241 (2004).

24. Mair, R. D., Sirich, T. L., Plummer, N. S. & Meyer, T. W. Characteristics of Colon-Derived Uremic Solutes. *Clin. J. Am. Soc. Nephrol.* **13**, 1398–1404 (2018).

25. Carter, M. M. *et al. A Human Milk Oligosaccharide Alters the Microbiome, Circulating Hormones, Cytokines and Metabolites in a Randomized Controlled Trial of Older Individuals*. http://medrxiv.org/lookup/doi/10.1101/2023.08.18.23294085 (2023) doi:10.1101/2023.08.18.23294085.

26. Cheng, A. G. *et al.* Design, construction, and in vivo augmentation of a complex gut microbiome. *Cell* **185**, 3617-3636.e19 (2022).

27. Hubatsch, I., Ragnarsson, E. G. E. & Artursson, P. Determination of drug permeability and prediction of drug absorption in Caco-2 monolayers. *Nat. Protoc.* **2**, 2111–2119 (2007).

28. Kuo, C. Y. *et al.* Tomatidine Attenuates Airway Hyperresponsiveness and Inflammation by Suppressing Th2 Cytokines in a Mouse Model of Asthma. *Mediators Inflamm.* **2017**, (2017).

29. Shieh, J. M. *et al.* α-Tomatine Suppresses Invasion and Migration of Human Non-Small Cell Lung Cancer NCI-H460 Cells Through Inactivating FAK/PI3K/Akt Signaling Pathway and Reducing Binding Activity of NF-κB. *Cell Biochem. Biophys.* **60**, 297–310 (2011).

30. Kúdelová, J. *et al.* Alpha-tomatine activates cell cycle checkpoints in the absence of DNA damage in human leukemic MOLT-4 cells. *J. Appl. Biomed.* **11**, 93–103 (2013).

31. Friedman, M. *et al.* Tomatine-containing green tomato extracts inhibit growth of human breast, colon, liver, and stomach cancer cells. *J. Agric. Food Chem.* **57**, 5727–5733 (2009).

32. Roddick, J. G. The acetylcholinesterase-inhibitory activity of steroidal glycoalkaloids and their aglycones. *Phytochemistry* **28**, 2631–2634 (1989).

33. Nigg, H. N. *et al.* Inhibition of Human Plasma and Serum Butyrylcholinesterase (EC 3.1.1.8) by *α* -Chaconine and *α* -Solanine. *Toxicol. Sci.* **33**, 272–281 (1996).

34. McMillan, M. & Thompson, J.C. An outbreak of suspected solanine poisoning in schoolboys: examination of criteria of solanine poisoning. *QJM Int. J. Med.* **48**, (1979).

35. Ellman, G. L., Courtney, K. D., Andres, V. & Featherstone, R. M. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **7**, 88–95 (1961).

36. Goyal, R. K. & Hirano, I. The Enteric Nervous System. *N. Engl. J. Med.* **334**, 1106–1115 (1996).

37. Kim, D. H. Gut microbiota-mediated pharmacokinetics of ginseng saponins. *J. Ginseng Res.* **42**, 255–263 (2018).

38. Han, B. H., Park, M. H. & Han, Y. N. Degradation of ginseng saponins under mild acidic conditions. *Planta Med.* **44**, 146–149 (1982).

39. Schneider, H., Schwiertz, A., Collins, M. D. & Blaut, M. Anaerobic transformation of quercetin-3-glucoside by bacteria from the human intestinal tract. *Arch. Microbiol.* **171**, 81–91 (1999).

40. Day, A. J. *et al.* Deglycosylation of flavonoid and isoflavonoid glycosides by human small intestine and liver β-glucosidase activity. *FEBS Lett.* **436**, 71–75 (1998).

41. Friedman, M., Kozukue, N. & Harden, L. A. Preparation and Characterization of Acid Hydrolysis Products of the Tomato Glycoalkaloid α-Tomatine. *J. Agric. Food Chem.* **46**, 2096–2101 (1998).

42. Zhao, L. *et al.* Gut bacteria selectively promoted by dietary fibers alleviate type 2 diabetes.
Science **359**, 1151–1156 (2018).
43. Rakoff-Nahoum, S., Coyne, M. J. & Comstock, L. E. An Ecological Network of
Polysaccharide Utilization among Human Intestinal Symbionts. *Curr. Biol.* **24**, 40–49 (2014).
44. Wang, M. *et al.* Strain dropouts reveal interactions that govern the metabolic output of the
gut microbiome. *Cell* **186**, 2839-2852.e21 (2023).
45. King, R. R. & McQueen, R. E. Transformations of Potato Glycoalkaloids by Rumen
Microorganisms. *J. Agric. Food Chem.* **29**, 1101–1103 (1981).
46. Gérard, P. *et al. Bacteroides* sp. Strain D8, the First Cholesterol-Reducing Bacterium
Isolated from Human Feces. *Appl. Environ. Microbiol.* **73**, 5742–5749 (2007).
47. Freier, T. A., Beitz, D. C., Li, L. & Hartman, P. A. Characterization of Eubacterium
coprostanoligenes sp. nov., a Cholesterol-Reducing Anaerobe. *Int. J. Syst. Bacteriol.* **44**,
137–142 (1994).
48. Eyssen, H. J., Parmentier, G. G., Compernolle, F. C., De Pauw, G. & Piessens-Denef, M.
Biohydrogenation of Sterols by *Eubacterium* ATCC 21,408— *Nova* Species. *Eur. J.*
Biochem. **36**, 411–421 (1973).
49. Kenny, D. J. *et al.* Cholesterol Metabolism by Uncultured Human Gut Bacteria Influences
Host Cholesterol Level. *Cell Host Microbe* **28**, 245-257.e6 (2020).
50. Ibach, B. & Haen, E. Acetylcholinesterase Inhibition in Alzheimers Disease. *Curr. Pharm.*
Des. **10**, 231–251 (2004).
51. Almasieh, M., Zhou, Y., Kelly, M. E., Casanova, C. & Di Polo, A. Structural and functional
neuroprotection in glaucoma: role of galantamine-mediated activation of muscarinic
acetylcholine receptors. *Cell Death Dis.* **1**, e27–e27 (2010).
52. Andrew Luckey, Edward Livingston, & Yvette Tache. Mechanisms and Treatment of
Postoperative Ileus. *Arch. Surg.* **138**, 206 (2003).
53. Slanina, P. Solanine (glycoalkaloids) in potatoes: Toxicological evaluation. *Food Chem.*
Toxicol. **28**, 759–761 (1990).
54. Mensinga, T. T. *et al.* Potato glycoalkaloids and adverse effects in humans: An ascending
dose study. *Regul. Toxicol. Pharmacol.* **41**, 66–72 (2005).
55. Patil, B. C., Sharma, R. P., Salunkhe, D. K. & Salunkhe, K. Evaluation of solanine toxicity.
Food Cosmet. Toxicol. **10**, 395–398 (1972).
56. Yajima, T., Inoue, R., Matsumoto, M. & Yajima, M. Non-neuronal release of ACh plays a key
role in secretory response to luminal propionate in rat colon. *J. Physiol.* **589**, 953–962
(2011).
57. Yu, H. *et al.* Acetylcholine acts through M3 muscarinic receptor to activate the EGFR
signaling and promotes gastric cancer cell proliferation. *Sci. Rep.* **7**, 40802 (2017).
58. Billipp, T. E. *et al. Tuft Cell-Derived Acetylcholine Regulates Epithelial Fluid Secretion.*
http://biorxiv.org/lookup/doi/10.1101/2023.03.17.533208 (2023)
doi:10.1101/2023.03.17.533208.
59. Vizi, S. E., Bertaccini, G., Impicciatore, M. & Knoll, J. Evidence that Acetylcholine Released
by Gastrin and Related Polypeptides Contributes to their Effect on Gastrointestinal Motility.
Gastroenterology **64**, 268–277 (1973).
60. Dyle, M. C. *et al.* Systems-based discovery of tomatidine as a natural small molecule
inhibitor of skeletal muscle atrophy. *J. Biol. Chem.* **289**, 14913–14924 (2014).
61. Hang, S. *et al.* Bile acid metabolites control TH17 and Treg cell differentiation. *Nature* **576**,
143–148 (2019).
62. Kuss, S. K. *et al.* Intestinal Microbiota Promote Enteric Virus Replication and Systemic
Pathogenesis. *Science* **334**, 249–252 (2011).
63. Chagnon, F. *et al.* Unraveling the structure-activity relationship of tomatidine, a steroid
alkaloid with unique antibiotic properties against persistent forms of Staphylococcus aureus.
Eur. J. Med. Chem. **80**, 605–620 (2014).

64. Almeida, A. F., Santos, C. N. & Ventura, M. R. Synthesis of New Sulfated and Glucuronated
Metabolites of Dietary Phenolic Compounds Identified in Human Biological Samples. *J.*
Agric. Food Chem. **65**, 6460–6466 (2017).
65. Cockcroft, D. W. & Gault, H. Prediction of Creatinine Clearance from Serum Creatinine.
Nephron **16**, 31–41 (1976).
66. Dodd, D. *et al.* A gut bacterial pathway metabolizes aromatic amino acids into nine
circulating metabolites. *Nature* **551**, 648–652 (2017).
67. Han, S. *et al.* A metabolomics pipeline for the mechanistic interrogation of the gut
microbiome. *Nature* **595**, (2021).
68. Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: A toolkit to classify
genomes with the genome taxonomy database. *Bioinformatics* **36**, 1925–1927 (2020).
69. The Human Microbiome Project Consortium. A framework for human microbiome research.
Nature **486**, 215–221 (2012).
70. Tautenhahn, R., Patti, G. J., Rinehart, D. & Siuzdak, G. XCMS Online: A Web-Based
Platform to Process Untargeted Metabolomic Data. *Anal. Chem.* **84**, 5035–5039 (2012).
71. Cahill, M. G., Caprioli, G., Vittori, S. & James, K. J. Elucidation of the mass fragmentation
pathways of potato glycoalkaloids and aglycons using Orbitrap mass spectrometry. *J. Mass*
Spectrom. **45**, 1019–1025 (2010).
72. Caprioli, G., Cahill, M., Logrippo, S. & James, K. Elucidation of the mass fragmentation
pathways of tomatidine and β1-hydroxytomatine using orbitrap mass spectrometry. *Nat.*
Prod. Commun. **10**, 575–576 (2015).
73. Falany, C. N., Wheeler, J., Tae Sung Oh & Falany, J. L. Steroid sulfation by expressed
human cytosolic sulfotransferases. *J. Steroid Biochem. Mol. Biol.* **48**, 369–375 (1994).
74. Yao, L. *et al.* A biosynthetic pathway for the selective sulfonation of steroidal metabolites by
human gut bacteria. *Nat. Microbiol.* **7**, 1404–1418 (2022).
75. Johnson, E. L. *et al.* Sphingolipids produced by gut bacteria enter host metabolic pathways
impacting ceramide levels. *Nat. Commun.* **11**, 1–11 (2020).
76. Panevska, A., Skočaj, M., Križaj, I., Maček, P. & Sepčić, K. Ceramide
phosphoethanolamine, an enigmatic cellular membrane sphingolipid. *Biochim. Biophys.*
Acta - Biomembr. **1861**, 1284–1292 (2019).
77. Osbourn, A., Bowyer, P., Lunness, P., Clarke, B. & Daniels, M. Fungal Pathogens of Oat
Detoxify Host Saponins. *Mpmi* **8**, (1995).
78. Kaup, O., Gräfen, I., Zellermann, E. M., Eichenlaub, R. & Gartemann, K. H. Identification of
a tomatinase in the tomato-pathogenic actinomycete Clavibacter michiganensis subsp.
michiganensis NCPPB382. *Mol. Plant. Microbe Interact.* **18**, 1090–1098 (2005).
79. Hennessy, R. C. *et al.* A Screening Method for the Isolation of Bacteria Capable of
Degrading Toxic Steroidal Glycoalkaloids Present in Potato. *Front. Microbiol.* **9**, 2648
(2018).
80. Wang, W. *et al.* Glutamicibacter halophytocola-mediated host fitness of potato tuber moth
on Solanaceae crops. *Pest Manag. Sci.* **78**, 3920–3930 (2022).
1217