Gamma-TuRC proteins contribute to the dynamic organization of MTOCs during *Drosophila* spermatogenesis

Elham Alzyoud[1#], Dóra Németh[1,2#], Viktor Vedelek[1], Titanilla Szögi[3], Mónika Krecsmarik[1], Zoltán Lipinszki[4*], Rita Sinka[1*]

[1]Department of Genetics, University of Szeged, Szeged, Hungary
[2]Faculty of Science and Informatics, Doctoral School of Biology, University of Szeged, Szeged, Hungary
[3]Department of Pathology, University of Szeged, Szeged, Hungary
[4]HUN-REN Biological Research Centre, Institute of Biochemistry, MTA SZBK Lendület Laboratory of Cell Cycle Regulation, Szeged, Hungary

\# These authors contributed equally to this work
* Authors for correspondence (rsinka@bio.u-szeged.hu, lipinszki.zoltan@brc.hu)

Abstract

The initiation of microtubule formation in eukaryotes is primarily facilitated by γ-tubulin and the highly conserved protein complex called γ-Tubulin Ring Complex (γ-TuRC) in various microtubule-organizing centers (MTOCs). Non-centrosomal MTOCs have a range of functions and contribute to diverse cellular activities in different cell types. While the heterogeneity of tissue-specific MTOCs and γ-TuRC in *Drosophila* testis has been described, their molecular composition and physiological significance are poorly understood. In this study, we investigate the testis-specific distribution of the canonical γ-TuRC by *in vivo* tagging of two complex members, Grip163 and Grip84. We find that both the testis-specific (t- γ-TuRC) and canonical form of γ-TuRC are simultaneously present and colocalize with each other at the basal body, apical nuclear tip, and near the elongating mitochondrial derivatives in the post-meiotic spermatids. We propose the possibility of the formation of MTOCs containing both t-γ-TuRC and γ-TuRC complexes. We also find some of the known γ-TuRC binding partners localizing to the apical nuclear tip. Additionally, we identify the Golgi localization of Grip84 in spermatocytes, alongside its centrosomal and diverse post-meiotic localization. These results highlight and prove the importance of the different γ-TuRCs in organizing the diverse MTOCs during the extensive rearrangement of cell organelles in *Drosophila* spermatogenesis.

Introduction

The microtubule (MT) network in eukaryotic cells plays a crucial role in cell differentiation and division, as well as movement through the cilia and flagella. The spatial re-organization of the MT cytoskeleton is a fast and dynamic process that requires distinct subcellular nucleation sites called microtubule organizing centers (MTOCs) that act as templates for the MT arrays to initiate. The main MTOC in dividing cells is the centrosome, which consists of a pair of centrioles (an older mother and a newer daughter centriole), and the pericentriolar material (PCM). When cells exit the cell cycle and start differentiation, their centrosomes become inactive, resulting in a gradual loss of their MTOC activity [1]. To support different functions and structures, the cells employ a variety of non-centrosomal MT organizing centers (ncMTOCs) [2]. The heterogeneity of ncMTOCs is reflected in the molecular composition and the cellular localization, and their function is unique to each cell type [3].

MTs are polar structures containing a plus and a minus end. The highly dynamic plus end is generally where the majority of MT polymerization and depolymerization occurs, while the minus end is often stabilized and capped [4]. The main factor known to specifically associate with microtubule minus ends is the γ-tubulin ring complex (γ-TuRC) [5,6]. PCM and ncMTOCs both recruit γTuRCs to the microtubule nucleation site [7,8]. γTuRC is a multiprotein complex consisting of γ-tubulin and its associated proteins (γ-tubulin ring proteins (Grips) in Drosophila, or γ-tubulin complex proteins (GCPs) in human) that are essential for γTuRC integrity and MT nucleation activity. The conserved and essential core of the γTuRC is the small complex (γTuSC) that consists of two γ-tubulins and one copy of Grip84 and Grip91 in Drosophila [9]. The large ring complex comprises multiple γ-TuSCs and additional Grips, including Grip75, Grip128, Grip163, and Grip71 [10,11] and is able to nucleate microtubule organization more efficiently than the small complex.

Similarly to mammals, Drosophila spermatogenesis involves a series of stage-dependent morphological changes when stem cells give rise to mature sperm. During this process, round spermatids of 10 μm in diameter elongate to become 1850 μm in length [12], which requires an extensive remodelling of existing cellular organelles, as well as the addition of new structures. After meiosis, each spermatid inherits a centriole that will serve a basal body around a ring-shaped PCM-like macromolecular structure called the centriole adjunct. The growth of the axoneme starts from the basal body and continues to elongate towards the spermatid tail [13]. Opposite to the basal body, on one side of the nucleus, the acroblast starts to develop, which is a Golgi-derived organelle. During the later stages of elongation, the acroblast will be converted

into the acrosome. Another important event that occurs after meiosis is the fusion of the mitochondria into
two giant mitochondrial derivatives, called the nebenkern. These mitochondrial derivatives elongate with
the axoneme during sperm maturation and accumulate paracrystalline material in one of them, which could
provide structural support for the growing tail [12,14].

Growing evidence suggests that testis-specific MTOC components play important roles in basal body and
ncMTOC formation and also in the positioning and shaping of the post-meiotic organelles during
Drosophila spermatogenesis [15–17]. For instance, testis-specific CnnT expression contributes to the
conversion of mitochondria into a ncMTOC, independent of core pericentriolar Cnn [16]. In addition to the
canonical γ-TuRC genes (Grip84, Grip91, Grip75, Grip128, Grip163, and Grip71), recently we have
identified three additional testis-specific paralogs of the γ-TuRC components, t-Grip84, t-Grip91, and t-
Grip128 [15,18]. We have shown, that from the round spermatid stage onward, these t-γ-TuRC proteins start
to localize to the centriole adjunct, then to the nuclear tip, and finally surrounding the mitochondria of the
elongating cyst. We also presented that they bind to γ-Tubulin and Mzt1 and interact with each other in
the post-meiotic stages [15]. We and others have shown that Mzt1 binds to the N-termini of the GCP proteins
(Zupa et al., 2021) and interacts with the N-terminal regions of Grip91, t-Grip91 and Grip128 in
Drosophila [15,17]. Our finding raised the question of whether the canonical γ-TuRC proteins are replaced
with their testis-specific counterparts, or whether they are able to cooperate to meet the needs of ncMTOCs
during the late stages of spermatogenesis. We also aimed to understand whether the testis-specific γ-TuRC
can interact with the canonical γ-TuRC or with any other proteins. In order to address these queries, we
created *in situ* tagged *Grip163* and *Grip84* transgenic lines to elucidate the adaptable localization and
identify interacting partners of the γ-TuRC during the development of germ cells.

Results

Phenotypic characterization of the *Grip163*[Δ11] mutant line

We have previously reported that t-γ-TuRC proteins are present only in the post-meiotic spermatids [15].
However, there is a lack of data regarding the relationship between the canonical and the testis-specific γ-
TuRCs, as well as the presence or localization of ubiquitously expressed γ-TuRC components in the late
stages of spermatogenesis.

Therefore, we tested the presence and possible function of the γ-TuRC component Grip163, which has no
testis-specific paralogs, in sperm development [19,10,15]. To achieve this, we induced mutation in the *Grip163*
gene, and performed an *in situ* tagging of the *Grip163* gene, using the CRISPR-Cas9 gene editing

technology. Using a specific guided RNA designed for *Grip163*, we deleted 11 base pairs from the coding

region of the gene (hereafter, *Grip163^{Δ11}*). This deletion caused a frameshift leading to a premature stop

codon in the coding region of *Grip163* (**S.Figure 1a**). Interestingly, we found that in contrast to the G*rip84*

and *Grip91* homozygotes, which are lethal [20,21], the homozygous *Grip163^{Δ11}* mutant animals are viable,

but male and female sterile (**S.Figure 1c).** The seminal vesicles of the *Grip163^{Δ11}* males were empty,

without mature sperms (**Figure 1 a-b**). A very similar phenotype was observed in *Grip128* and *Grip75*

(γ-TuRC components) mutants [22].

Based on this phenotype we investigated both the early and late stages of spermatogenesis in

*Grip163^{Δ11}.*flies. We found an abnormal nebenkern-nucleus ratio in the round spermatids, which is a

hallmark of abnormal mitotic division (**Figure 2 a-c**). Nevertheless, many of the cysts are able to elongate

in the *Grip163^{Δ11}* mutant, visualized by the β2tub-GFP transgene (**Figure 1 a, b**). To further investigate

the elongation of the cysts, we tested whether the individualization proceeds normally. To visualize every

elongated post-meiotic cyst, we stained the actin cones of the individualization complex (IC) in the

Grip163^{Δ11} testis and found the lack of IC in *Grip163^{Δ11}*, providing evidence for the failure of proper cyst

elongation, which is the prerequisite of individualisation **(Figure 1 c, d).** For further evidence, we used

anti-pan polyglycylated tubulin (Axo49) antibody to visualize mature axonemes of the spermatids, at the

onset of individualization [23]. We observed significantly shorter axonemes in the *Grip163^{Δ11}* mutant

compared to the WT, indicating abnormal spermatid elongation **(Figure 1 e-g).** Axoneme elongation

depends on the mitochondria, which provide a structural platform for MT reorganization to support the

robust elongation taking place at the tip of the sperm tail (Nouguchi 2011). Based on this we tested the

presence and appearance of late elongating mitochondria using dj-GFP as a marker. In accordance with

the defect observed in axoneme elongation, it was detected that the mitochondria in the *Grip163^{Δ11}*

mutants also exhibited incomplete elongation (**Figure 1 e, f**). Based on this general spermatid elongation

defect of *Grip163^{Δ11}* flies, the detailed morphology of post-meiotic cysts was further investigated using

transmission electron microscopy (TEM).

In wild-type testis, the 64 spermatids develop simultaneously and on the TEM image of elongated cysts

we observed the two mitochondrial derivatives attached to the axoneme, which has the classical 9+2

arrangement of microtubules in each spermatid (**Figure 2 d-h**). In *Grip163^{Δ11}* mutants, cysts became

disorganized, with fewer spermatids per cyst and with disturbed mitochondrial:axonemal ratio (**Figure 2**

b). Frequently, multiple mitochondrial derivatives were attached to a single axoneme, and we also

observed multiple damaged axonemes **(Figure 2 h)**, supporting that Grip163 is necessary for normal

axoneme formation and/or stability. Since the γ-TuRC is essential for centrosome formation [11,20,24,25], we

examined the centrosome/basal body distribution at different stages of spermatogenesis in the testes of

Grip163[A11] males. We tested the distribution of the centrosomal/basal body localized γ-Tubulin and the

PACT domain of DPlp fused with GFP (GFP-PACT) [26]. In the wild type animals both proteins localize to

the centrosome/basal body during different stages of spermatogenesis **(Figure 3 a, b)**. In the *Grip163[A11]*

mutant flies we found strong GFP-PACT positive centrosomal and basal body signals, but the γ-Tubulin

signal in spermatocytes and spermatids was weak compared to the wild type **(Figure 3 c, d)**. In wild type

males, GFP-PACT is located close to the nucleus and the nebenkern, as expected in round spermatids. In

contrast, multiple and scattered GFP-PACT signals were observed in the *Grip163[A11]* mutants' testis

(Figure 3 c-d). We could detect γ-Tubulin in the post-meiotic elongating spermatids, but those were

scattered along the cysts of *Grip163[A11]* mutant **(Figure 3 c, d)**. All these data suggest that Grip163 could

be essential for the formation of stable γ-TuRC prior to and at the onset of meiosis, however, the newly

organized testis-specific γ-TuRC could contribute to the γ-Tubulin binding after meiosis.

Interestingly, mutants of the canonically expressed γ-TuRC components, *Grip128* and *Grip75,* are also

viable and sterile with scattered basal body distribution [22]. Contrary to the essentiality of γ-TuSC genes,

the absence of the additional complex members contributing to the formation of γ-TuRC, is not critical

for viability, but they are required for normal fertility in both sexes [22].

Localization of γ-TuRC proteins during spermatogenesis

To test the subcellular localization of Grip163 at different stages of spermatogenesis, we generated

endogenously GFP-tagged Grip163 lines. We and others have shown that the N- or C-termini of the γ-

TuRC proteins are responsible for the binding of their interacting partners [15,17], therefore we tagged the

Grip163 close the N-terminus, without interfering with the functional GCP domains of the protein

(S.Figure 1 a). The fertility of the GFP-Grip163 males was comparable to the wild-type animals,

indicating that the inserted fluorophore does not alter the function of the protein **(S.Figure 1 c)**.

GFP-Grip163 signal was present on the centrosomes of spermatocytes and after meiosis, it appeared on

the basal body/centriole adjunct of the developing spermatids as well **(Figure 4 a-e)**. Similar to the testis-

specific γ-TuRC members, GFP-Grip163 was also detected at the apical end of the nuclei and

accumulation was observed at the mitochondrial surface **(Figure 4 d, e)**. These results strongly suggest

that the testis-specific γ-TuRC does not replace the canonical γ-TuRC, but both are present and form

similarly distributed versatile ncMTOCs in the post-meiotic spermatids.

The localization of GFP-Grip163 to the centriole adjunct has prompted an inquiry into the potential presence of other canonical γ-TuSC components in the subsequent developmental stages. To investigate this, we applied CRISPR technology to *in situ* tag Grip84 with GFP. We designed the incorporation in a way that all 5 potential isoforms of Grip84 will have the N-terminally inserted GFP tag, outside of the GCP domain (**S.Figure 1 b**). It was previously reported that the classical mutant of *Grip84* is lethal, therefore the normal viability and fertility of the *in situ* GFP-tagged Grip84 serves as evidence that the GFP tag does not alter the normal function of Grip84 (**S.Figure 1 c**) [27].

We found that GFP-Grip84 is present on the centrosomes of spermatocytes, similarly to the GFP-Grip163 protein (**Figure 5 a-e**).

This localization of GFP-Grip84 was also verified by its colocalization with the centrosomal component Asterless (Asl) (**Figure 6 a, S.Figure 2 a, b**) [28].

Following meiosis, GFP-Grip84 is present on the centriole adjunct and then appears at the apical nuclear tip tip and exhibits a strong overlap with the elongated mitochondria, akin to the GFP-Grip163. We wanted to know whether the post-meiotic localization of GFP-Grip84 overlaps with its testis-specific paralogue, t-Grip84. To do this we co-expressed GFP-Grip84 and t-Grip84-mCherry together and found complete colocalization after the meiotic stages (**Figure 5 f, g**) [15]. These results provide additional evidence that both the ubiquitously expressed γ-TuRC and the testis-specific γ-TuRC may together play a role in the formation of the diverse microtubule organizing centers in post-meiotic stages of spermatids.

Apart from the centrosomal localization of GFP-Grip84, we also noticed strong bright punctae in the cytoplasm of spermatocytes, whose distribution is reminiscent of t-ER-Golgi units. It is known that centrosomal proteins are associated with the Golgi apparatus in mammalian cells [29]. Golgi organization is special in Drosophila, as the scattered Golgi stacks are found in close proximity to the ER exit sites (t-ER-Golgi) [30]. To test the t-ER-Golgi localization of GFP-Grip84, we used GM130 as a *cis*-Golgi, as well as GCC88, as a *trans*-Golgi-specific antibody for co-immunostaining (**Figure 6 b-e, S.Figure 2 c-d**). We observed strong co-localization of GFP-Grip84 with the *cis*-Golgi, suggesting that GFP-Grip84 is connected to the t-ER-Golgi units of spermatocytes.

In a comprehensive analysis of the effector proteins of Rab GTPases, Grip84, Grip91 and γ-Tubulin were identified as interacting partners of the Golgi-localizing Rab2 and Rab4 [31]. Rab2A and Rab2B proteins were shown to be essential for acrosome formation and male fertility in mice [32], therefore it seemed reasonable to check whether the Golgi-localizing Rab2 might also be present at the apical end where the acrosome is located during nuclear elongation in Drosophila spermatids. We tested the germline-specific

Bam-Gal4 driven pUASP-Rab2-YFP localization in spermatocytes and spermatids. We detected Rab2-
YFP on the t-ER-Golgi units adjacent to the *trans*-Golgi-specific antibody GCC88 in spermatocytes and
on the acrosome containing apical nuclear tip of the elongated spermatids (**Figure 6 d, e**). Localization of
GFP-Grip84 to the Golgi apparatus led us to hypothesize that Drosophila γ-TuRC, similarly to the
mammalian γ-TuRC is necessary to the Golgi organization and also raised the possibility of a conserved
function of Rab2 in acrosome formation. Additional experiments can address these highly intriguing
prospects.
200
Post-meiotic ncMTOCs contain centrosomal components
The versatile localization pattern of GFP-Grip84 prompted us to investigate whether any of the already
known binding partners of the γ-TuRC could be present on the newly identified localization foci. It was
shown previously that γ-Tubulin is localized in the apical region of the spermatid nuclei, specifically at
the distal end of the MTs present within the dense complex [15,33]. We previously demonstrated the presence
of t-γ-TuRC and Mtz1 at the apical tip of the nucleus, suggesting the presence of a ncMTOC organized
opposite to the basal body during nuclear elongation [15]. Therefore, we carefully examined the precise
localization of the centriole component, Sas4, a known interacting partner of γ-TuRC proteins by
immunostaining using anti-Sas4 antibody [34,35]. We detected Sas4 on the centrosome of mitotic
spermatocytes and the basal body of the spermatids after meiosis (**Figure 7 a-c**). Intriguingly, we detected
endogenous Sas4 on the apical tip of the elongating spermatid nuclei as well (**Figure 6 c**). Apart from the
γ-TuRC proteins, Sas4 has multiple centriolar binding partners, including Sas6, Asl, Ana1, and Dplp [28,35–
38]. Therefore, we wondered whether they are also present on the ncMTOC at the apical tip of the spermatid
nuclei. To achieve this we performed immunostaings using gene-specific antibodies [28,34,39]. We detected
Sas6 and Asl on the centriole adjunct and also at the apical tip of the nuclei (**S.Figure 3 a, b**), however,
Ana1 and GFP-PACT colocalized with t-Grip91 exclusively at the centriole adjunct region of the
elongating leaf and canoe stage nuclei, in line with the reported localization pattern of these proteins [33]
(**Figure 7 d, S.Figure 3 c**). These results indicate that the ncMTOC localised to the apical nuclear tip
contains several well-defined binding partners of the γ-TuRC.
220
Composition of t-γ-TuRC
We next examined whether the partially overlapping localization pattern might be a consequence of the
physical interaction between the canonical γ-TuRC and the t-γ-TuRC and/or the t-γ-TuRC proteins and

the γ-TuRC interacting protein Sas4. To address this, we performed a yeast two-hybrid (Y2H) analysis,
where we tested t-γ-TuRC proteins with the canonical γ-TuSC protein Grip91 and Sas4. Due to the large
size of the proteins, we divided all three t-γ-TuRC proteins and Grip91 into an N-terminal (N) and a C-
terminal (C) part. We found a specific interaction between t-Grip84-C and Grip91-N, which suggests the
possibility of the presence of a testis-specific and non-testis-specific mixed γ-TuRC (**Figure 8 a, S.Figure**
4). We also detected interaction between Sas4 and t-Grip91-N, t-Grip91-C, t-Grip128-N and t-Grip128-C
however, we could not find an interaction between Sas4 and t-Grip84-N, t-Grip84-C (**Figure 8 a, S.Figure**
4). To support Y2H results and prove the direct binding between proteins, we performed *in vitro* binding
experiments. We used immobilised GST-tagged recombinant proteins (N- or C-terminal parts of the t-γ-
TuRC proteins, as indicated) as bait, which were mixed with [35]S-methionine-labelled Sas4 or Grip91,
respectively, produced by a coupled *in vitro* transcription-translation (IVTT) reaction followed by
autoradiography (**Figure 8 b-e**). The *in vitro* binding experiments confirmed the results of the Y2H assay
(**Figure 8 a-e**). Taken together the *in vivo* localization pattern and the *in vitro* interaction results suggest
a versatile γ-TuRC containing ncMTOC formation in Drosophila spermatids.

Discussion

Drosophila spermiogenesis and spermatogenesis represent an ideal model for studying mitotic and meiotic
cell division, centrosome and basal body formation and dynamic organelle reorganization after meiosis.
It was proven to be a good model for understanding centrosomal and non-centrosomal MT nucleation due
to the intensive morphological changes and organelle movement in the post-meiotic spermatids [15–17]. MT
nucleation was reported, among others, at the surface of the Golgi [40], mitochondria [16], nuclear envelope [41]
and plasma membrane [42] in mammalian cells. We previously identified and described the function and
versatile localization of three testis-specific γ-TuRC proteins in the post-meiotic spermatids [15](**Figure 9**).
The described centriole adjunct, nuclear tip and mitochondrial surface localization of the t-γ-TuRC
proteins and their interaction partner Mzt1 raised the question of whether the ubiquitously expressed γ-
TuRC is replaced after meiosis for the testis-specific one or both are present due to the multiple ncMTOC
formation during spermatid elongation and acrosome formation. γ-Tubulin was already reported to
localize to both tips of the elongating nuclei and to the surface of the giant mitochondria during elongation
[15,16,33]. The testis-specific splice variant of Centrosomin (Cnn), CnnT, was also reported to promote non-
centrosomal MT growth on the surface of the mitochondrial derivative of spermatids [16]. However, we
lacked the canonical γ-TuRC localization after the meiotic stages of spermatogenesis, despite both CnnT

and γ-Tubulin can bind to γ-TuRC proteins. Mutants for the small complex components Grip91 and
Grip84 are lethal and have defects in spindle assembly [20,21], which indicates their essentiality for the whole
fly development. It has been demonstrated that the small complex is the minimal unit needed for MT
nucleation [9]. However, RNAi depletion of the ubiquitously expressed Grip163 results in loss of Grip84
recruitment to the ncMTOC in tracheal cells [11].

Our experiments showed that, in contrast to the essentiality of the γ-TuSC, the γ-TuRC member Grip163
is only essential for fertility, similar to the already characterized γ-TuRC members Grip75 and Grip128
[22]. From the elongation and basal body stability defects of *Grip163* mutant spermatids, we inferred that
the presence of Grip163 protein is essential after meiosis. This was supported by the localization of GFP-
Grip163, as we detected its presence in the localization foci of t-γ-TuRC members after meiosis. These
results raised the possibility that t-γ-TuRC and the canonical γ-TuRC may be present simultaneously, even
as a mixed complex, in the aforementioned ncMTOCs (**Figure 9**). This hypothesis was further supported
by the fact that Grip84, a member of γ-TuSC, also exhibits an overlapping localization pattern in the late
stages of spermatogenesis and the physical interaction between t-Grip84 and the ubiquitously expressed
Grip91. Our results strongly suggest that both testis-specific and canonical γ-TuRC members are involved
in the formation of post-meiotic ncMTOCs.

In vivo tagging of Grip84 enabled us to describe the Golgi localization of the γ-TuSC component Grip84
in primary spermatocytes, in addition to the extensively studied centrosomal localization of the Grip84 [21].
Until now γ-Tubulin containing Golgi outpost-associated MT nucleation sites was only described to
regulate distal dendritic branching in *Drosophila* neurons [43]. Golgi non-centrosomal MT assembly is
mediated by the γ-Tubulin-interacting AKAP450 protein, which is anchored by the cis-Golgi-associated
GM130 protein. This provides evidence that the Golgi apparatus may become a major site of MT
nucleation by acting as a γ-TuRC docking site in mammalian cells by a mechanism similar to that in the
centrosome (Rivero et al., 2009). Our results suggest that a similar mechanism could be present in the
primary spermatocytes. Further experiments are needed to determine exactly which isoform of the Grip84
protein is associated with the Golgi (**Figure 9**).

All three proteins of γ-TuSC (Grip84, Grip91 and γ-Tubulin) interact with the constitutively active, GTP-
bound Rab2 protein [31], and Rab2 and Grip84 both localize to both the Golgi and the apical end of the
nucleus, suggesting a role for γ-TuSC and Rab2 in the formation of the Golgi and lysosome-associated
acrosome. Rab2A and Rab2B proteins were shown to be components of the proacrosomal membrane and
essential for acrosome formation and male fertility in mice [32].

Drosophila Rab2 is an essential gene and recruitment of Rab2 takes place on the membranes of late
endosomes, where it controls the fusion of LAMP-containing biosynthetic carriers and lysosomes to late
endosomes in somatic cells [44]. t-ER-Golgi units contribute to the formation of the acroblast, which forms
the acrosome by fusing lysosomal structures during nuclear elongation in Drosophila [45]. The molecular
determinants of acrosome formation are hardly known in Drosophila, however, several retrograde
tethering complexes were described in the formation of a functional Golgi ribbon in the acroblast (Fári et
al 2016). Acrosomal proteins such as Sneaky, Lamp1 and Rab2, as well as the ubiquitous and testis-
specific γ-TuRC, its partner proteins (Mzt1, Sas4) and other centrosomal proteins (Asl, Sas6), are all
localized to the apical tip of the nucleus. Further biochemical studies are needed to clarify the precise
interactions between the aforementioned proteins. Nevertheless, these results raise the possibility of a link
between the ncMTOC and acrosome organization, which is present at the dense complex forming
perinuclear MTs at the apical end of the nuclei(**Figure 9**).
298

299
Figure 1. Mitochondrial and axonemal abnormalities lead to abnormal spermatid elongation in
Grip163^{Δ11} mutant
(a, b) Microtubules are visualized with the β2tub-GFP transgene (green), while the nuclei stained with
DAPI (blue) in WT (a) and _Grip163^{Δ11}_ mutant spermatids (b) Elongated cysts are present both in WT and
Grip163^{Δ11}, however, the seminal vesicle (dashed lines) of _Grip163^{Δ11}_ mutant does not contain β2tub-
GFP positive mature sperms. (c, d) The individualization complexes (IC) were stained with Phalloidin
(red), ICs are formed normally (arrows) in WT, (c) and missing in _Grip163^{Δ11}_ mutant (d). (e, f) Fully
developed axonemes are visualized by Axo49 tubulin antibody (red) and elongated mitochondria are with
DJ-GFP transgene (green) both in the testis and seminal vesicle (dashed line). Elongated polyglycylated
axonemes and mitochondria were shorter in _Grip163^{Δ11}_ mutant and the seminal vesicle was empty (dashed
line) (f) compared to the WT (e). (g) Diagrams show the average length of Axo49 positive cysts in WT
and _Grip163^{Δ11}_ mutant. (WT n=12, _Grip163^{Δ11}_ n=22) Statistical significance was tested by one-way
ANOVA. (scale bars: 50 μm).

313
Figure 2: Abnormal mitochondrial and axonemal structures in the post-meiotic cysts of *Grip163^{Δ11}*
mutant

(a, b) Phase contrast images show a normal 1:1 (white outline), nucleus: nebenkern ratio in WT **(a)** and
an abnormal ratio (red outline) in *Grip163^{411}* mutant **(b)**. **(c)** Violin plot showing the nucleus: nebenkern
ratio in WT (n=186) and *Grip163^{411}* mutant round spermatids (n=112). **(d, h)** Transmission electron
micrograms of testis cross-sections, representing elongating cysts of WT **(d,f)** and *Grip163^{411}* mutant
(e,g,h). Disorganized cysts with axonemal and mitochondrial abnormalities are characteristic in
Grip163^{411} mutant **(f-h)** WT elongating cyst shows normal axoneme structure with 9+2 tubulin dimers
(highlighted in green) and two mitochondrial derivatives (yellow and orange), with paracrystalline
material formation in one of them (orange) **(f)**. In *Grip163^{411}* mutant the mitochondria and the axoneme
detached from each other, and multiple mitochondrial derivatives (highlighted in orange and yellow)
attached to one axoneme (green). Mitochondria fused, enlarged (blue arrowheads) and detached (arrow)
from the axoneme and disintegrated axonemes with microtubule bundles and the axial membrane opening
is characteristic of *Grip163^{411}* mutant cyst (black arrowheads) **(g, h)**. (scale bars: A-E 2 μm, F-H 500 nm)
328

Figure 3: Mislocalization of centrosomal markers in *Grip163^{Δ11}* mutant

(a-b) GFP-PACT (green) and γ-tubulin (red) are colocalizing on the centrosome (arrows) and basal body (arrowheads) in WT spermatocytes and spermatids **(a, b)**. γ-tubulin, but not GFP-PACT centrosomal localization is affected in *Grip163^{Δ11}* mutant spermatocytes. In the round spermatids, GFP-PACT and γ-tubulin are colocalised **(c)**, but the post-meiotic elongating cysts are disorganized with scattered basal bodies. **(d)**. (scale bars: 20 μm)

Figure 4: Localization of GFP-Grip163 during spermatogenesis.
(a-c) The endogenously tagged GFP-Grip163 colocalizes with γ-tubulin at the centrosome and the basal body throughout spermatogenesis (arrows). **(d, e)** GFP-Grip163 localizes additionally to the apical tip of the elongating nuclei **(d, inset)**, near the elongating mitochondria (arrowheads) **(e)** with overlapping localization with the t-γ-TuRC represented by HA-t-Grip91 **(e)**. (scale bars: 10 μm)

350
351
Figure 5: Localization of GFP-Grip84 during spermatogenesis.
(A-E) In vivo tagged GFP-Grip84 localizes both to the centrosome (**A**) and the basal body (**B**, **C**) (white
arrows) and in the post-meiotic elongating cyst GFP-Grip84 localizes also to the apical tip of the
elongating nuclei (**D**) (red arrows). Additionally, it is localized surrounding the elongating mitochondria
(**C**, **D**) (orange arrowheads). Golgi-like localization of GFP-Grip84 was observed in spermatocytes (**C**)
(white arrowheads). **(F, G)** GFP-Grip84 localizes to the centrosome before meiosis (blue arrows) and
shows overlapping localization with t-Grip84-mCh at the basal body (white arrows) and the apical tip of
the elongating nuclei (red arrows) in the post-meiotic spermatids (Scale bars: 20 μm).
360
361

Figure 6: γ-TuSC protein GFP-Grip84 is localizing to the Golgi in spermatocytes

(A) GFP-Grip84 colocalizes with centrosomal protein Asl (anti-Asl, white arrow) and also to the Golgi apparatus (white arrowhead). **(B, C)** GFP-Grip84 colocalizes with antibody-stained *cis*-Golgi marker GM130 in the spermatocytes (arrowheads). **(D, E)** Rab2-YFP localizes to the Golgi (arrow) in spermatocytes visualised by the *trans*-Golgi marker GCC88 and also to the apical end of the elongating nuclei (arrowhead). (Scale bars: A, C, D 10 μm, B, E 20 μm).

370
Figure 7: Sas4 is a component of the ncMTOC at the apical tip of the elongating nuclei
(A, B) Antibody-stained Sas4 (red) colocalizes with the endogenous tag GFP-Grip84 at the centrosome
(arrows) but not on the Golgi (arrowheads) in the spermatocytes. **(C)** Sas4 (red) is localizing to the basal
body of the elongating spermatids (arrows) and additionally to the apical tip of the nuclei (arrowheads).
(D) Antibody-stained Ana1 localizes only to the basal body (arrows) of the elongating spermatids, while
the t-γ-TuRC component HA-t-Grip91 localizes to the basal body(arrow) and the apical tip (arrowheads)
of the elongating spermatids. (Scale bars: 20 μm).
378
379

a

	t-Grip84-NT	t-Grip84-CT	t-Grip91-NT	t-Grip91-CT	t-Grip128-NT	t-Grip128-CT
Sas4	-	-	+	+	+	+
Grip91-NT	-	+	-	-	-	-
Grip91-CT	-	-	-	-	-	-

Figure 8: Interactions of t-γ-TuRC proteins with Sas4 and the canonical Grip91.
(A) Summary of Y2H analysis of the N- or C-terminal parts of t-γ-TuRC proteins (as indicated in the table) with Sas4, and the N- or C-terminal parts of Grip91, respectively. (B-E) Autoradiographic images showing the presence or absence of interaction between purified and immobilised GST-t-TuRC protein fragments and ^{35}S-labelled Sas4 or Grip91, synthesized by *in vitro* transcription-translation (IVTT) reaction.

389 **Figure 9: Summary of localization of γ-TuRC/t-γ-TuRC protein containing MTOCs during**

390 **different stages of spermatogenesis.**

Materials and Methods

Fly stocks, mutants, and fertility test

Flies were maintained on cornmeal agar medium at 25˚C in standard laboratory conditions. Fly stocks used in this study were obtained from Bloomington Drosophila Stock Center, except stated otherwise. w^{1118} used as wild-type control (3605, BDSC), *P{DJ-GFP.S}AS1/CyO*. The β2-tubulin-GFP line was provided by David M. Glover. GFP-PACT line was obtained from Jordan Raff's and [26]. HA-t-Grip91 and t-Grip84-mCh were described before (Alzyoud et al., 2021).

To establish the mutant allele *Grip163^Δ11^* the pCFD5-Grip163-gRNA line was crossed with vas-Cas9 source line to generate deletion in *Grip163*. The deletions were identified by *Grip163* specific PCR and further confirmed by direct sequencing. *Grip163^Δ11^* line was established and used for further analysis.

To establish the genomic tagged lines of *Grip84* and *Grip163* pCFD5-Grip84-gRNA and pCFD5-Grip163-gRNA lines were crossed with the vas-Cas9 source lines , w[1118]; PBac{y[+mDint2] GFP[E.3xP3]=vas-Cas9}VK00027 (BL51324), y[1] M{RFP[3xP3.PB] GFP[E.3xP3]=vas-Cas9}ZH-2A w[1118]/FM7c (BL51323) respectively, progeny of this cross were injected with the homology arm carrying pJET1.2 plasmid (Thermo Fisher Scientific). Transgenic candidates were identified by gene specific PCR and confirmed by sequencing of the insertion sites. For fertility tests, individual males were crossed with three w[1118] virgin females. After 3 days the parents were removed and 14 days after crossing, the hatched progeny was counted in every tube. Experiments were repeated 4 times with > 10 males per condition.

DNA constructs

To make the genomic GFP-Grip84, and GFP-Grip163 line, annealed oligos of the sgRNAs (ATCCTTGAGCGGCTACCGCT), (GTGCTTCTCCGGTCCAGGCT) for Grip84 and Grip163 respectively were cloned into pCFD5 vector (Addgene, 112645) at Bbs1 restriction site using HiFi DNA Assembly Master Mix (E2621S, NEB) [46]. To establish the homology arms 1534bp upstream, 1529bp downstream, 1515bp upstream and 1738bp downstream of the intended GFP insertion site for Grip84 and Grip163 respectively in addition to GFP were amplified using High fidelity DNA polymerase. The purified PCR products were cloned into pJET1.2 plasmid (Thermo Fisher Scientific) using HiFi DNA Assembly Master Mix (E2621S, NEB) [46]. GFP sequence was introduced 187bp upstream for Grip84, and 229bp upstream of the Grip163 sgRNA. Stable transgenic lines were established as described in Port et al. [47; 48].

For Y2H analysis t-Grip84 (1-399 aa N-term, 400-728 aa C-term), t-Grip128 (1-550 aa N-term, 551-975

aa C-term), full-length were cloned into pGAD424. full-lengths of Sas4, and t-Grip91s (1-966 aa N-term,

967-1932 aa C-term) were cloned into pGBT9, using HiFi DNA Assembly Master Mix (E2621S, NEB).

To generate GST-tagged bait proteins for the *in vitro* pull-down assay, full-length Grip84, N- and C-

terminal parts of t-Grip84, t-Grip91 and t-Grip128 were inserted into the pDEST15 (11802014, Thermo

Fisher Scientific) vector by Gateway cloning (11791-020, Thermo Fisher Scientific). Sas4 and Grip91

cDNAs were obtained from the *Drosophila* Gold collection (DGRC) and inserted into the pDEST17

plasmid (11803012, Thermo Fisher Scientific) by Gateway cloning. Oligonucleotide primers used in

cloning are listed in **(S. Table 1)**.

Staining and microscopy

Preparation and staining of tests with formaldehyde or methanol:acetone fixation were performed as

described by White-Cooper *et. al* [49]. Briefly, testes were dissected and fixed in 4% formaldehyde for 25

minutes or semi-squashed using a coverslip on poly-L-lysine coated slides, then frozen with liquid

nitrogen and fixed for 5 minutes in ice-cold methanol followed by 5 minutes in acetone. After fixation

samples were blocked and antibodies were added, the antibodies used in the study are listed in **(S.Table**

2). Stained samples were mounted in SlowFade Gold antifade reagent (S36967, Life Technologies).

Images were taken using Olympus BX51 fluorescent microscope or Olympus Fluoview Fv10i confocal

microscope.

Testes for transmission electron microscopy were dissected and fixed overnight at 4°C in 2%

paraformaldehyde and 2% glutaraldehyde. After rinsing in phosphate-buffered saline (PBS), the testes

were further fixed for 1 h in 2% OsO_4, followed by dehydration in increasing ethanol concentrations and

rinsing in uranyl acetate and acetone. The samples were embedded in Embed812 (Electron Microscopy

Sciences, Hatfield, PA, USA) then ultrathin (70 nm) sections were cut on an Ultracut S ultra-microtome

(Leica, Wetzlar, Germany), which were mounted on copper grids. The grids were counterstained with

uranyl acetate (Merck, Germany) and lead citrate (Merck, Germany) and were examined and

photographed with a JEOL JEM 1400 transmission electron microscope (JEOL, Tokyo, Japan). The

images were processed with the TEM Center software (JEOL, Tokyo, Japan).

Yeast two-hybrid assay

Y2H assays were carried out using the Matchmaker two-hybrid system (K1605, Clontech). One of the

baits pGBT9: N- and C- terminal parts of t-Grip91 and full-length γ-Tub23C was cotransformed with one

of the preys pGAD424: N- and C- terminal parts of t-Grip84, t-Grip91, t-Grip128, full-length Mzt1 and γ-Tub23C into PJ69 yeast cells [50]. Individual colonies were streaked out on yeast synthetic double drop-out plates that lacked tryptophan and leucine (Y0750, Merck), after growing, they were inoculated on quadruple dropout plates lacking (tryptophan, leucine, histidine, and adenine (Y2021, Merck) contains 10mM 3-amino-1, 2, 4-aminotrizole (3-AT) (A8056, Sigma).

Recombinant protein production

Recombinant GST-tagged bait proteins were expressed, purified and immobilized onto glutathione sepharose beads as described previously [51].

Coupled *in vitro* transcription-translation (IVTT) reaction and *in vitro* binding assay

[35]S-methionine-labelled prey proteins (Grip91, Sas4) were produced *in vitro* using the TnT T7 Quick Coupled IVTT kit (L1170, Promega). The detailed protocol of the GST-IVTT binding assay is described previously [52].

Quantification and statistical analysis

Data analysis and graph production was done using Origin 9.0 (OriginLab Corporation, Northampton, MA, USA). One-way ANOVA followed by Tukey's HSD test was used to comparisons between WT and mutants.

Acknowledgements
We are grateful to the members of RS group. We are indebted to David M. Glover, Helen White-Cooper, Sean Munro and Jordan Raff for fly stocks and reagents; for the Drosophila Genomics Resource Center (supported by NIH grant 2P40OD010949) for cDNA clones and to Ferenc Jankovics for critical reading of the manuscript.

Funding
This work was supported by Hungarian Science Foundation (K132155) to RS and by the Hungarian Academy of Sciences (Lendület Program Grant (LP2017-7/2017)) to ZL.

References

1. Muroyama, A., Seldin, L. & Lechler, T. Divergent regulation of functionally distinct γ-tubulin
complexes during differentiation. *J. Cell Biol.* **213**, 679–692 (2016).
2. Muroyama, A. & Lechler, T. Microtubule organization, dynamics and functions in
differentiated cells. *Dev. Camb. Engl.* **144**, 3012–3021 (2017).
3. Tillery, M., Blake-Hedges, C., Zheng, Y., Buchwalter, R. & Megraw, T. Centrosomal and
Non-Centrosomal Microtubule-Organizing Centers (MTOCs) in Drosophila melanogaster.
Cells **7**, 121 (2018).
4. Desai, A. & Mitchison, T. J. Microtubule polymerization dynamics. *Annu. Rev. Cell Dev.
Biol.* **13**, 83–117 (1997).
5. Moritz, M., Braunfeld, M. B., Guénebaut, V., Heuser, J. & Agard, D. A. Structure of the γ-
tubulin ring complex: A template for microtubule nucleation. *Nat. Cell Biol.* **2**, 365–370
(2000).
6. Zheng, Y., Wong, M. L., Alberts, B. & Mitchison, T. Nucleation of microtubule assembly by
a gamma-tubulin-containing ring complex. *Nature* **378**, 578–583 (1995).
7. Fry, A. M., Sampson, J., Shak, C. & Shackleton, S. Recent advances in pericentriolar
material organization: ordered layers and scaffolding gels. *F1000Research* **6**, 1622 (2017).
8. Kollman, J. M., Merdes, A., Mourey, L. & Agard, D. A. Microtubule nucleation by γ-tubulin
complexes. *Nat. Rev. Mol. Cell Biol.* **12**, 709–721 (2011).
9. Oegema, K. *et al.* Characterization of Two Related Drosophila γ-tubulin Complexes that
Differ in Their Ability to Nucleate Microtubules. *J. Cell Biol.* **144**, 721–733 (1999).
10. Gunawardane, R. N. *et al.* Characterization and reconstitution of Drosophila gamma-tubulin
ring complex subunits. *J. Cell Biol.* **151**, 1513–1524 (2000).
11. Vérollet, C. *et al.* Drosophila melanogaster gamma-TuRC is dispensable for targeting
gamma-tubulin to the centrosome and microtubule nucleation. *J. Cell Biol.* **172**, 517–528
(2006).
12. Tokuyasu, K. T. Dynamics of spermiogenesis in Drosophila melanogaster. VI. Significance
of 'onion' nebenkern formation. *J. Ultrastruct. Res.* **53**, 93–112 (1975).
13. Avasthi, P. & Marshall, W. F. Stages of ciliogenesis and regulation of ciliary length. *Differ.
Res. Biol. Divers.* **83**, S30-42 (2012).
14. Laurinyecz, B. *et al.* Sperm-Leucylaminopeptidases are required for male fertility as
structural components of mitochondrial paracrystalline material in Drosophila melanogaster
sperm. *PLoS Genet.* **15**, 1–24 (2019).
15. Alzyoud, E., Vedelek, V., Réthi-Nagy, Z., Lipinszki, Z. & Sinka, R. Microtubule Organizing
Centers Contain Testis-Specific γ-TuRC Proteins in Spermatids of Drosophila. *Front. Cell
Dev. Biol.* **9**, 727264 (2021).
16. Chen, J. V., Buchwalter, R. A., Kao, L.-R. & Megraw, T. L. A Splice Variant of Centrosomin
Converts Mitochondria to Microtubule-Organizing Centers. *Curr. Biol. CB* **27**, 1928-1940.e6
(2017).
17. Tovey, C. A. *et al.* γ-TuRC Heterogeneity Revealed by Analysis of Mozart1. *Curr. Biol. CB*
28, 2314-2323.e6 (2018).
18. Vedelek, V. *et al.* Analysis of Drosophila melanogaster testis transcriptome 06 Biological
Sciences 0604 Genetics. *BMC Genomics* **19**, 1–19 (2018).

19. Wasbrough, E. R. *et al.* The Drosophila melanogaster sperm proteome-II (DmSP-II). *J. Proteomics* **73**, 2171–2185 (2010).

20. Barbosa, V., Yamamoto, R. R., Henderson, D. S. & Glover, D. M. Mutation of a Drosophila gamma tubulin ring complex subunit encoded by discs degenerate-4 differentially disrupts centrosomal protein localization. *Genes Dev.* **14**, 3126–3139 (2000).

21. Colombié, N. *et al.* The Drosophila gamma-tubulin small complex subunit Dgrip84 is required for structural and functional integrity of the spindle apparatus. *Mol. Biol. Cell* **17**, 272–282 (2006).

22. Vogt, N., Koch, I., Schwarz, H., Schnorrer, F. & Nüsslein-Volhard, C. The gammaTuRC components Grip75 and Grip128 have an essential microtubule-anchoring function in the Drosophila germline. *Dev. Camb. Engl.* **133**, 3963–3972 (2006).

23. Bre, M. H. *et al.* Axonemal tubulin polyglycylation probed with two monoclonal antibodies: widespread evolutionary distribution, appearance during spermatozoan maturation and possible function in motility. *J. Cell Sci.* **109**, 727–738 (1996).

24. Bärenz, F., Mayilo, D. & Gruss, O. J. Centriolar satellites: Busy orbits around the centrosome. *Eur. J. Cell Biol.* **90**, 983–989 (2011).

25. Gupta, H. *et al.* SAS-6 Association with γ-Tubulin Ring Complex Is Required for Centriole Duplication in Human Cells. *Curr. Biol.* **30**, 2395-2403.e4 (2020).

26. Martinez-Campos, M., Basto, R., Baker, J., Kernan, M. & Raff, J. W. The Drosophila pericentrin-like protein is essential for cilia/flagella function, but appears to be dispensable for mitosis. *J. Cell Biol.* **165**, 673–683 (2004).

27. Bourbon, H.-M. *et al.* A P-insertion screen identifying novel X-linked essential genes in Drosophila. *Mech. Dev.* **110**, 71–83 (2002).

28. Dzhindzhev, N. S. *et al.* Asterless is a scaffold for the onset of centriole assembly. *Nature* **467**, 714–718 (2010).

29. Allan, V. J., Thompson, H. M. & McNiven, M. A. Motoring around the Golgi. *Nat. Cell Biol.* **4**, E236–E242 (2002).

30. Kondylis, V. & Rabouille, C. The Golgi apparatus: Lessons from Drosophila. *FEBS Lett.* **583**, 3827–3838 (2009).

31. Gillingham, A. K., Sinka, R., Torres, I. L., Lilley, K. S. & Munro, S. Toward a Comprehensive Map of the Effectors of Rab GTPases. *Dev. Cell* **31**, (2014).

32. Morohoshi, A. *et al.* FAM71F1 binds to RAB2A and RAB2B and is essential for acrosome formation and male fertility in mice. *Development* **148**, dev199644 (2021).

33. Riparbelli, M. G., Persico, V. & Callaini, G. The Microtubule Cytoskeleton during the Early Drosophila Spermiogenesis. *Cells* **9**, 2684 (2020).

34. Fu, J. *et al.* Conserved molecular interactions in centriole-to-centrosome conversion. *Nat. Cell Biol.* **18**, 87–99 (2016).

35. Gopalakrishnan, J. *et al.* Tubulin nucleotide status controls Sas-4-dependent pericentriolar material recruitment. *Nat. Cell Biol.* **14**, 865–873 (2012).

36. Galletta, B. J. *et al.* A centrosome interactome provides insight into organelle assembly and reveals a non-duplication role for Plk4. *Nat. Commun.* **7**, 12476 (2016).

37. Gopalakrishnan, J. *et al.* Sas-4 provides a scaffold for cytoplasmic complexes and tethers them in a centrosome. *Nat. Commun.* **2**, 359 (2011).

38. Guruharsha, K. G. *et al.* A Protein Complex Network of Drosophila melanogaster. *Cell* **147**, 690–703 (2011).

39. Dzhindzhev, N. S. *et al.* Plk4 phosphorylates Ana2 to trigger Sas6 recruitment and
procentriole formation. *Curr. Biol. CB* **24**, 2526–2532 (2014).
40. Efimov, A. *et al.* Asymmetric CLASP-dependent nucleation of non-centrosomal
microtubules at the trans-Golgi network. *Dev. Cell* **12**, 917–930 (2007).
41. Becker, R. *et al.* Myogenin controls via AKAP6 non-centrosomal microtubule-organizing
center formation at the nuclear envelope. *eLife* **10**, e65672 (2021).
42. Brodu, V., Baffet, A. D., Le Droguen, P.-M., Casanova, J. & Guichet, A. A developmentally
regulated two-step process generates a noncentrosomal microtubule network in Drosophila
tracheal cells. *Dev. Cell* **18**, 790–801 (2010).
43. Mukherjee, A., Brooks, P. S., Bernard, F., Guichet, A. & Conduit, P. T. Microtubules
originate asymmetrically at the somatic golgi and are guided via Kinesin2 to maintain
polarity within neurons. *eLife* **9**, e58943 (2020).
44. Lund, V. K., Madsen, K. L. & Kjaerulff, O. Drosophila Rab2 controls endosome-lysosome
fusion and LAMP delivery to late endosomes. *Autophagy* **14**, 1520–1542 (2018).
45. Fári, K., Takács, S., Ungár, D. & Sinka, R. The role of acroblast formation during
Drosophila spermatogenesis. *Biol. Open* **5**, 1102–1110 (2016).
46. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred
kilobases. *Nat. Methods* **6**, 343–345 (2009).
47. Port, F. & Bullock, S. L. Creating Heritable Mutations in Drosophila with CRISPR-Cas9. in
Methods in Molecular Biology vol. 1478 145–160 (2016).
48. Port, F., Chen, H. M., Lee, T. & Bullock, S. L. Optimized CRISPR/Cas tools for efficient
germline and somatic genome engineering in Drosophila. *Proc. Natl. Acad. Sci. U. S. A.*
111, (2014).
49. White-cooper, H. Analysis of Meiosis and Morphogenesis. *Drosoph. Cytogenet. Protoc.*
247, 468 (2004).
50. James, P., Halladay, J. & Craig, E. A. Genomic Libraries and a Host Strain Designed for
Highly Efficient Two-Hybrid Selection in Yeast. *Genetics* **144**, 1425–1436 (1996).
51. Karman, Z. *et al.* Novel perspectives of target-binding by the evolutionarily conserved PP4
phosphatase. *Open Biol.* **10**, 200343 (2020).
52. Réthi-Nagy, Z., Ábrahám, E. & Lipinszki, Z. GST-IVTT pull-down: a fast and versatile
in vitro method for validating and mapping protein-protein interactions. *FEBS Open Bio* **12**,
1988–1995 (2022).
53. Sinka, R., Gillingham, A. K., Kondylis, V. & Munro, S. Golgi coiled-coil proteins contain
multiple binding sites for Rab family G proteins. *J. Cell Biol.* **183**, 607–15 (2008).
603
604

605
Supplementary Figure 1: Structural arrangement of *Grip163* and *Grip84* genes and transcripts.
(A) Gene structure, transcript composition and protein domains (GCP-N, GCP-C) of Grip163. gRNA
position (11,997,349..11,997,371 [+]) and the position of the *in vivo* inserted GFP tag (11,997,601 [-]) is
labelled. *Grip163^{Δ11}* mutant has an 11bp deletion (11,997,369..11,997,359). The DNA and the protein
sequences surrounding the deletion are shown. **(B)** Gene structure of Grip84 with the sites of gRNA
(19,564,635..19,564,657 [-]) and the GFP tag (19,564,845 [-]). The inserted GFP tag labels all five Grip84

isoforms **(C)** Graph represents the fertility test of WT (n=14), *Grip163^{A11}* (n=6), *Grip163^{A11}/Cyo* (n= 12),
GFP-Grip84 (n=12) and GFP-Grip163 (n=15) males. Statistical significance was tested by one-way
ANOVA.

615
Supplementary Figure 2: Grip84 and Grip163 localization during spermatogenesis.

(A-D) The endogenous tag GFP-Grip84 colocalizes with antibody-stained Asl (red) at the **(A)** centrosome
(arrow) and the **(B)** basal body (arrow) of the spermatocytes and the spermatids, but not with the Golgi
localized GFP-Grip84 (arrowhead). **(C, D)** GFP-Grip84 localizes adjacent to antibody-stained *trans*-Golgi
marker GCC88 (arrowheads) and to the centrosome (arrow) in spermatocytes. **(E, F)** in vivo tagged GFP-
Grip163 colocalizes with the testis-specific HA-t-Grip91 (arrow) in the post-meiotic stages. (Scale bars:
20 μm).
623

624
Supplementary Figure 3: Localization of centrosomal and basal body component proteins in
spermatids.
(A, B) Asl, Sas6 and Sas4 localization to the basal body (arrow) and the apical tip (arrowhead) of
elongating spermatids are visualised by immunostaining using gene-specific antibodies. **(C)** GFP-PACT
localizes to the basal body (arrows) of the elongating spermatids, while the HA-t-Grip91 localizes to the
basal body (arrow) and the apical tip (arrowheads) of the elongating spermatids. (Scale bars: 20 μm)

Supplementary Figure 4: Interactions tested by Yeast Two-Hybrid assay
Mated yeast was plated as 10-fold serial dilutions (left to right) on -Leu, -Trp medium that is selective for the bait and prey plasmids and -Leu, -Trp, -His, -Ade medium that is selective for the bait, prey, and the interaction between the tested proteins. Negative control (-control) is the empty prey with the corresponding bait vector.

638   **Supplementary Table1 List of antibodies.**

Antibody /Dye	Dilution	Source	Identifier	Fixation method
Anti-HA (rat)	1:100	Sigma-Aldrich/Roche	Cat# 11867423001	Formaldehyde
Anti-Sas4 (rat)	1:500	[34]		Methanol-acetone
Anti-Sas6 (rat)	1:500	[39]		Methanol-acetone
GM130 (rabbit)	1:500	Abcam	ab30637	Formaldehyde
GCC88 (G. pig)	1:100	[53]		Formaldehyde
Anti-GFP (rabbit)	1:500	Thermo Fisher Scientific	Cat#A-11122	Methanol-acetone
anti-pan polyglycylated Tubulin Axo49 (mouse)	1:5000	Sigma-Aldrich/Merck	Cat# MABS276	Formaldehyde
anti-γ-Tubulin (mouse)	1:5000	Sigma-Aldrich/Merck	Cat# T6557	Formaldehyde
anti-Ana1 (rabbit)	1:5000	[34]	-	Formaldehyde
anti-Asl (rabbit)	1:1000	[28]	-	Methanol-acetone
G.pig Alexa Fluor 546 secondary	1:400	ThermoFisher	Cat# A-11074	
Mouse Alexa Fluor 488 secondary	1:400	Thermo Fisher Scientific	Cat#A-11029	
Mouse Alexa Fluor 546 secondary	1:400	Thermo Fisher Scientific	Cat#A-11030	
Mouse Alexa Fluor 633 secondary	1:400	Thermo Fisher Scientific	Cat#A-21052	
Rat Alexa Fluor 488 secondary	1:400	Thermo Fisher Scientific	Cat#A-21208	
Rat Alexa Fluor 568 secondary	1:400	Thermo Fisher Scientific	Cat#A-11081	

Rabbit Alexa Fluor 488 secondary	1:400	Thermo Fisher Scientific	Cat#A-11008	
Rabbit Alexa Fluor 546 secondary	1:400	Thermo Fisher Scientific	Cat#A-11010	
4',6-Diamidino-2-Phenylindole, Dihydrochloride (DAPI)	1 μg/ml	Thermo Fisher Scientific	Cat#D1306	

640 **Supplementary Table2 List and sequence of primers**

Primer Name	Primer Sequence
Grip163-gRNA-fwd	tgcagtgcttctccggtccaggct
Grip163-gRNA-rev	aaacagcctggaccggagaagcac
Grip163-1.5 kb upstream_fwd	gtgccacctgacgtctaagagatacatcccataattaccatacttacccag
Grip163-1.5 kb upstream_rev	tgctcaccatcagggggtttcagggcgcg
Grip163-GFP_fwd	gaaacccctgatggtgagcaagggcgag
Grip163-GFP_rev	gctgtagttgcttgtacagctcgtccatg
Grip163-before PAM_fwd	gctgtacaagcaactacagccatactccg
Grip163-before PAM_rev	ccaggctgtgccacaaaaggtcgttcatc
Grip163-Mutant PAM_fwd	ccttttgtggcacagcctggaccggagaag
Grip163-Mutant PAM_rev	gagttagctcactcattaggggtcaggactggacgacag
pJET1_fwd	cctaatgagtgagctaactcac
pJET1_rev	tcttagacgtcaggtggcac
Grip84-gRNA-fwd	tgcaatccttgagcggctaccgct
Grip84-gRNA-rev	aaacagcggtagccgctcaaggat
Grip84-1.5 kb upstream-fwd	ccacctgacgtctaagatgacccgctgaccatggc
Grip84-1.5 kb upstream-rev	ctcaccattacgggtgtagtgtcggattc
Grip84-GFP_fwd	cacccgtaatggtgagcaagggcgag
Grip84-GFP_rev	acaaagaccttgtacagctcgtccatg
Grip84-before PAM_fwd	gctgtacaaggtctttgtcaggcgcggc
Grip84-before PAM_rev	acagagcggtagccgctcaaggattg
Grip84-mutant PAM_fwd	cggctaccgctctgtgaccaacaaaacg
Grip84-mutant PAM_rev	ttagctcactcattagggctcgcagtgcagatcgtc
YTH Sas4_fwd	atcgccggaattcccaatgcaggaggctggcgaaag
YTH Sas4_rev	ggtcgacggatccccctaatacttggcatagtctgtgtccattatg
YTH t-Grip91-N_fw	atcgccggaattcccaatggagacgcagccatttaaaattg
YTH t-Grip91-N_rev	ggtcgacggatccccttaaactaacatagggggtatcaatttcaac
YTH t-Grip91-C_fw	atcgccggaattcccaatggaaagtcttcgtgaag
YTH t-Grip91-C_rev	ggtcgacggatccccttacagtgatcttttcctttcc

YTH t-Grip84-N_fw	aaagagatcgaattcccaatgaactccgacaaatcg
YTH t-Grip84-N_rev	ggtcgacggatccccttaattgtgctcctccacc
YTH t-Grip84-C_fw	aaagagatcgaattcccaatgttgcctttacacattcg
YTH t-Grip84-C_rev	ggtcgacggatccccttaaatattggaagctatttgatcc
YTH t-Grip128-N_fw	aaagagatcgaattcccaatgcaagaggacgagag
YTH t-Grip128-N_rev	ggtcgacggatccccttatcgttcatcggtcacaatc
YTH t-Grip128-C_fw	aaagagatcgaattcccaatgacagagtacgtcttaagaaaatc
YTH t-Grip128-C_rev	ggtcgacggatccccttaaatatggccgagcaaac
GST t-Grip91-N_fw	cgacggtaccatggagacgcagccatttaaaat
GST t-Grip91-N_rev	gcggccgcttaaactaacataggggtatcaatttcaac
GST t-Grip128-N_fw	cgacggtaccatgcaagaggacgagagaca
GST t-Grip128-N_rev	gcggccgcttatcgttcatcggtcacaatc
GST t-Grip128-C_fw	cgacggtaccatgacagagtacgtcttaagaaaatc
GST t-Grip128-C_rev	gcggccgcttaaatatggccgagcaaaca
GST t-Grip84-N_fw	gaattcaaatgaactccgacaaatcggaaagtg
GST t-Grip84-N_rev	gcggccgcttaattgtgctcctccacc
GST t-Grip84-C_fw	gaattcaaatgttgcctttacacattcg
GST t-Grip84-C_rev	gcggccgcttaaatattggaagctatttgatcc

642