GABA-ergic inhibition in human MT predicts visuo-spatial intelligence mediated by reverberation with frontal cortex

Yuan Gao[1#], Yong-Chun Cai[2#], Dong-Yu Liu[1,3], Juan Yu[1,3], Jue Wang[1], Ming li[4], Bin Xu[1], Teng-Fei Wang[2], Gang Chen[1,3,6], Georg Northoff[5,6*], Ruiliang Bai[1,3,7*], Xue Mei Song[1,3,5,7*]

[1]Department of Neurosurgery of the Second Affiliated Hospital and Interdisciplinary Institute of Neuroscience and Technology, Zhejiang University School of Medicine, Hangzhou 310029, China;

[2]Department of Psychology and Behavioral Sciences, Zhejiang University, Hangzhou 310028, China;

[3]Key Laboratory of Biomedical Engineering of Ministry of Education, Qiushi Academy for Advanced Studies, College of Biomedical Engineering and Instrument Science, Zhejiang University, Hangzhou 310027, China;

[4] College of Intelligence Science and Technology, National University of Defense Technology, Changsha 410073, China;

[5]Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, Zhejiang University School of Medicine, Hangzhou 310013, China;

[6]University of Ottawa Institute of Mental Health Research, University of Ottawa; Ottawa, ON, K1Z 7K4, Canada;

[7]MOE Frontier Science Center for Brain Science & Brain- Machine Integration, Zhejiang University，Hangzhou 311121, China;

[#]Co-first author

23 *Correspondence: songxuemei@zju.edu.cn, ruiliangbai@zju.edu.cn,

24   georg.northoff@theroyal.ca

Abstract

The canonical theory emphasizes fronto-parietal network (FPN) is key in mediating

general fluid intelligence (gF). Meanwhile, recent studies show that multiple sensory

regions in occipito-temporal border also play a key role in gF. However, the

underlying mechanism is not yet clear. To investigate this issue, this study selects

human MT complex (MT+), a region locates at the occipito-temporal border

representing multiple sensory flows as a target brain area. Using ultra-high field

magnetic resonance spectroscopy (MRS) to measure GABA/glutamate concentrations

in MT+ combining resting-state fMRI functional connectivity (FC), behavioral

examinations including MT+ perception suppression test and gF subtest in visuo-

spatial component, we reveal that MT+ GABA and frontal-MT+ FC significantly

correlate with the performance of visuo-spatial intelligence. Further, serial mediation

model demonstrates that MT+ GABA predicting visuo-spatial gF fully mediated by

reverberation effect between frontal and MT+ network. Our finding highlights that

sensory cortex could integrate into complex cognition system as an intellectual hub.

Introduction

General fluid intelligence (gF) is a current problem-solving ability, which shows high

inter-individual differences in humans(Cattell & Raymond, 1963). At the beginning of

the last century, Spearman(Spearman, 1904) proposed that some general or g factor

contributes to our cognition. Visuo-spatial intelligence, usually tested by visual

materials, is considered to have high g-loading(Colom et al., 2006; Deary et al., 2010;

Jung & Haier, 2007). Recently, Duncan et al. show coding of gF in distributed

regions including a new multiple-demand (MD) core, the multiple sensory regions at

the temporo-occipital border(Assem et al., 2020; Duncan et al., 2020). However, more

evidence is needed to uncover how the sensory cortex is a genuine cognitive core. The

human extrastriate cortical region, middle temporal complex (MT+), locates at the

temporo-occipital border(Dumoulin et al., 2000), and is a key region in the multiple

representations of sensory flows (including optic, tactile, and auditory flows)(Bedny

et al., 2010; Ricciardi et al., 2007); this ideally suits it to be a new MD core. To

further investigate this issue, this work conducted multi-level examination including

biochemical (glutamatergic - gabaergic in MT+), regional-systemic (brain

connectivity with MT+ - based), and behavioral (visual motion function in MT+)

levels to reveal if the function of MT+ contributes to the visuo-spatial component of

gF.

Using a well-known visual motion paradigm (center-surround antagonism)(Liu et

al., 2016; Tadin et al., 2003), Melnick et al found a strong link between suppression

index (SI) of motion perception and the scores of the block design test (BDT, a

subtest of the Wechsler Adult Intelligence Scale (WASI), which measures the visuo-

spatial component (3D domain) of gF (Melnick et al., 2013). Based on these findings,

MT+ is likely to play a key role in MD system of gF. Strong connectivity between the

core regions suggests a mechanism for large-scale information exchange and

integration in the brain(Barbey, 2018; Cole et al., 2012). This potential conjunctive

coding may overlap with the inhibition and/or excitation mechanism of MT+. Taken

together, we hypothesized that 3D visuo-spatial intelligence (the performance of

BDT) might be predicted by the inhibitory and/or excitation mechanisms in MT+ and

the integrative functions connecting MT+ with frontal cortex (Figure 1a). To verify

the specificity of MT+, we used primary visual cortex (V1) - based GABA/Glu as

control as it mediates the 2D rather than 3D visual domain(Born & Bradley, 2005).

We employ ultra-high field (7T) magnetic resonance spectroscopy (MRS)

technology to reliably resolve GABA and Glu concentrations(Ende, 2015; Liu et al.,

2022; Song et al., 2021), and first demonstrate GABAergic inhibition mechanisms

(but not excitatory Glu) in MT+ region involve in the 3D visuo-spatial ability. Further,

analysis of functional brain connectivity at rest reveals that the network (between

MT+ and frontal cortex) relating to MT+ GABA and perceptual suppression

contribute the visuo-spatial intelligence. Our results provide direct evidence that

inhibitory mechanisms centered on GABA levels in MT+ region (a sensory cortex)

mediate multi-level visuo-spatial component (3D domain) of gF thus drawing a direct

connection of biochemistry, brain connectivity, and behavioral levels.

Figure 1. A priori hypothesis and experimental design. (**a**) Schematic of *a priori* hypothesis. The

inhibition mechanism centered on MT+ GABA, including molecular level - the GABAergic

inhibition in MT+ (green circle), brain connectivity of MT+ - frontal relating to MT+ GABA (blue

circle), and perceptual suppression of visual motion (the behavioral presentation of MT+ function,

red circle), contributes to the visuo-spatial component of gF (3D domain, yellow circle). (**b**)

Schematic of experimental design. Session 1 (rectangle box of short line) was the functional MRI

and MRS scanning at resting state. Session 2 (rectangle box of solid line) was another region of

MRS acquisition. In the two sessions, the order of MRS scanning regions (MT+ and V1) was

counter balanced across participants. There was a structure MRI scanning before each MRS data

acquisition. Interval between the two sessions was used for behavioral measurement (rectangle

box of dotted line): block design task (BDT) and psychophysical task - motion discrimination.

Sold lines indicate the experiment sequence.

Results

To determine whether the function of MT+ cortex contributes to visuo-spatial component (3D domain) of gF, we adopted the experimental design depicted in Figure 1b. Participants underwent two MRI sessions: the first encompassing resting-state fMRI and magnetic resonance spectra (MRS), and the second solely involving MRS. A 30-minute interval separated these sessions, during which participants performed motion discrimination tasks (using center-surround antagonism stimuli)(Tadin et al., 2003) and the block design test (BDT), which assesses the visuo-spatial ability (3D domain) of gF(Fangmeier et al., 2006). Both the volume-of-interests (VOIs) of MRS scanning in the left MT+ (targeted brain area) and the left primary visual cortex (V1, control brain area) had dimensions of $2\times2\times2$ cm^3, and the MRS scanning sequences were randomized across the two sessions. The MT+ MRS VOIs were demarcated using an anatomical landmark(Dumoulin et al., 2000). For 14 subjects, we also utilized fMRI to functionally pinpoint the MT+ to validate the placement of the VOI (Figure 2a, b). The V1 MRS VOIs were anatomically defined (Methods). Here, MRS data after extensive quality control (31/36 in MT+, and 28/36 in V1) were taken for further analysis (Methods).

Figure 2. MT+ localizer scans and MT+ MRS VOI placement. **(a)** Single task block designs.

First: a cross fixation on the center of the screen (10s). Second: a moving grating (2^{o}) toward left

last 10s. Third: the grating keeps static for 10s. Fourth: the grating moves toward right last 10s.

The localizer scans consist of 8 blocks. (**b**) MT+ location and MRS VOI placement. The upper

template is the horizontal view. The lower templates from left to right are coronal and sagittal

views. The warm color indicates the overlap of fMRI activation of MT+ across 14 subjects, the

cold color bar indicates the overlap of MRS VOIs across all subjects.

GABA and Glu concentrations in MT+ and V1 and their relation to SI and BDT

An example of MRS from voxel located in MT+ is shown in Figure 3a. LCModel

fittings for GABA spectra from all subjects in MT+ ($n = 31$) and V1 ($n = 28$) are

illustrated in Figure 3b (color scale presents the BDT scores). We discerned a

significant association between the inter-subjects' BDT scores and the GABA levels

in MT+ voxels, but not in V1 voxels. Quantitative analysis displayed that BDT

significantly correlates with GABA concentrations in MT+ voxels ($r = 0.39$, $P = 0.03$,

$n = 31$, Figure 3c). After using partial correlation to control for the effect of age, the

relationship remains significant ($r_{partial} = 0.43$ $P = 0.02$, 1 participant excluded due to

the age greater than mean + 2.5SD) In contrast, there was no obvious correlation

between BDT and GABA levels in V1 voxels (figure supplement 1a). We show that

SI significantly correlates with GABA levels in MT+ voxels ($r = 0.44$, $P = 0.01$, $n =$

31, Figure 3d). In contrast, no significant correlation between SI and GABA

concentrations in V1 voxels was observed (figure supplement 1b). This finding

supports the hypothesis that motion perception is associated with neural activity in

MT+ area, but not in V1(Schallmo et al., 2018). LCModel fittings for Glu spectra

from all subjects in MT+ ($n = 31$) and V1 ($n = 28$) voxels are presented in figure

supplement 2a.

   Unlike in the case of GABA, no significant correlations between BDT and Glu

levels were found in both MT+ and V1 voxels (figure supplement 2b, c). While, as

expected(Song et al., 2021), we observed significant positive correlations between

GABA and Glu concentrations in both MT+ ($r = 0.62$, $P < 0.001$, $n = 31$) and V1

voxels (r = 0.56, P = 0.002, n = 28) (figure supplement 3a, b). Additionally, a

significant correlation between SI and BDT was discerned (r = 0.51, P = 0.002, n =

34, figure supplement 4a, $r_{partial}$ = 0.67, P < 0.001, 1 participant excluded due to the

age greater than mean + 2.5SD), corroborating previous conclusions(Melnick et al.,

2013). Two outliers evident in Figure 3d were excluded, with consistent results

depicted in figure supplement 4b.

Figure 3. MRS spectra and the relationships between GABA levels and SI / BDT. (**a**) Example

spectrum from the MT+ voxel of one participant. The first line is the LCModel fitting result of all

metabolites, and the following lines show the Glu and GABA spectra fitting with LCModel, and

then the baseline. (**b**) Individual participants fitted GABA MRS spectra from the MT+ (top) and

V1 (bottom) voxels from baseline measurement. The colors of the GABA spectra represent the

individual differences of BDT. The color bar represents the scores of BDT. (**c) and (d**) Pearson's

correlations showing significant positive correlations between MT+ GABA and BDT scores (**c**),

between MT+ GABA and SI (**d**). The ribbon between dotted lines represents the 95% confidence

interval, and the black regression line represents the Pearson's correlation coefficient (r). GABA

and Glu concentrations (Conc.) are absolute, with units of mmol per kg wet weight (Methods).

168 **Figure supplement 1.** Relationships between GABA concentration in V1 and BDT/SI. **(a)** There

169 is no significant relationship between BDT and GABA concentrations in V1 region. **(b)** There is

170 also no obvious correlation between SI and GABA concentrations in V1 region. GABA

171 concentration (Conc.) is absolute, with units of mmol per kg wet weight (Methods).

Figure supplement 2. Individual Glu MRS spectra from MT+ / V1 regions and relationships

between BDT and Glu concentrations in MT+ / V1 regions. **(a)** Individual participant fitted

Glutamate MRS spectra from the MT+ (top, $n = 31$) and V1 (bottom, $n = 28$) voxels from baseline

measurement. The colors of the Glutamate spectra represent the individual differences of BDT.

The color bar represents the scores of BDT. **(b)** There is no significant relationship between BDT

and Glu concentrations in MT+ region. **c** There is also no significant correlation between BDT and

Glu concentrations in V1 region. Glutamate concentration (Conc.) is absolute, with units of mmol

per kg wet weight (Methods).

Figure supplement 3. Correlations between GABA and Glu concentrations in MT+ and V1

regions. **(a)** There is significant correlation between GABA and Glu concentrations in MT+

region. **(b)** The levels of GABA also significantly correlate with Glu in V1 region. GABA and Glu

concentrations (Conc.) are absolute, with units of mmol per kg wet weight (Methods).

Figure supplement 4. Two linear correlations. **(a)** Significant positive correlation between

suppression index and BDT scores (took out two outlines). **(b)** BDT also significantly correlates

with GABA concentration in MT+ region (without two outliers, having the similar result shown in

Figure 2d).

MT - frontal FC relates to SI and BDT

We next took the left MT+ as the seed region and separately measured interregional FCs between the seed region and each voxel in the frontal regions (a priori search space). These measurements were correlated with performance in 3D visuospatial ability (BDT) to identify FCs with significant correlations. Results from connectivity-BDT analysis are summarized in Table 1 and shown in Figure 4a. We found that brain regions with FC strength to the seed region (left MT+) significantly correlated with BDT scores were situated within the canonical cognitive cores of MD system or FPN (Brodmann areas (BAs) 6, 9, 10, 46, 47)(Assem et al., 2020; Deary et al., 2010; Duncan et al., 2020; Duncan et al., 2000; Gray et al., 2003; Jung & Haier, 2007). Across the whole brain search, the similar FCs (between MT+ and frontal cognitive cores) still showed significant correlations with BDT scores (Table supplement 1) (also shown in figure supplement 5a). Additionally, we identified certain parietal regions (BAs 7, 39, 40) with significant correlations between their connectivity to the left MT+ and the BDT scores (Table supplement 1) (also shown in figure supplement 5a). These significant connections between MT+ and FPN/MD system suggest that left MT+ is involved in the efficient information integration network, serving as a maker of intellectual hub.

Then, we correlated MT+ - based global FCs with SI. Though spatial suppression during motion perception (quantified by SI) is considered to be the function of area MT+(Melnick et al., 2013), the top-down modulation from the frontal cortex can increase surround suppression(Liu et al., 2016). Our connectivity-SI analysis in the

frontal regions (a priori search space) displayed 3 brain regions in which FCs strength

significantly correlated with SI: right BA4/6, left BA6, and right BA46 (summarized

in Table supplement 2, and shown in Figure 4b). Across the whole brain search, we

identified total 7 brain regions in which FCs strength significantly correlated with SI,

and 3 of these were in the frontal cortex. This is consistent with the results obtained

by connectivity-SI analysis in a priori search space (frontal cortex) (Table supplement

3 and figure supplement 5b).

226 **Table 1 FC of voxels showing significant correlation with BDT scores across subjects in**
227 **frontal cortex.**

FC number	Connected regions	BA	Size	Peak coordinate MNI (x, y, z)	r	P
1	Frontal_Sup_Orb_R	11	33	(12,63, -19.5)	-0.57	0.0011
2	Frontal_Inf_Orb_L	47	24	(-34.5,28.5, -13.5)	-0.63	0.0003
3	Frontal_Med_Orb_R	11	41	(3,43.5, -12)	-0.58	0.0009
4	Frontal_Inf_Orb_R	47	48	(-31.5,24, -12)	0.59	0.0008
5	Frontal_Inf_Orb_R	47	29	(25.5,30, -13.5)	0.67	0.0001
6	Insula_L	\	26	(-28.5,27,0)	0.67	0.0001
7	Frontal_Inf_Oper_R	45	41	(43.5,16.5,6)	0.64	0.0002
8	Frontal_Sup_R	10	25	(31.5,57,9)	0.59	0.0008
9	Frontal_Mid_L	10	82	(-33,48,12)	0.62	0.0003
10	Frontal_Inf_Oper_R	44	49	(51,7.5,21)	0.59	0.0007
11	Frontal_Inf_Oper_R	46	96	(49.5,16.5,28.5)	-0.62	0.0003
12	Frontal_Mid_L	10	32	(-31.5,49.5,24)	0.57	0.0012
13	Frontal_Mid_R	10	102	(31.5,36,30)	0.59	0.0009
14	Precentral_L	6	46	(-49.5, -1.5,34.5)	0.59	0.0007
15	Frontal_Mid_R	9	107	(51,19.5,40.5)	-0.67	0.0001
16	Frontal_Sup_L	9	35	(-9,60,37.5)	-0.69	0.0001
17	Frontal_Sup_Medial_R	9	74	(4.5,52.5,43.5)	-0.57	0.0011
18	Frontal_Sup_R	6	136	(28.5, -7.5,63)	-0.64	0.0002
19	Supp_Motor_Area_L	6	48	(-10.5,6,54)	0.63	0.0003
20	Frontal_Mid_L	6	119	(-24,4.5,55.5)	0.63	0.0003
21	Precentral_L	6	229	(-24, -18,66)	0.60	0.0005
22	Frontal_Sup_R	6	32	(16.5, -18,67.5)	0.68	0.0001
23	Precentral_R	6	80	(30, -24,70.5)	0.70	0.0001
24	Precentral_R	6	23	(16.5, -25.5,76.5)	0.70	0.0001

228 Single voxel threshold $P < 0.005$ (t > 3.057 or t < -3.057), adjacent size ≥ 23 voxels (AlphaSim

229 corrected).

230

231 **Table supplementary 1. FC of voxels showing significant correlation with BDT scores across**
232 **subjects in whole brain.**

FC number	Connected regions	BA	Size	Peak coordinate MNI (x, y, z)	r	P
1	Frontal_Med_OrbR	11	44	(2,43.5, -12)	-0.58	0.0009
2	Frontal_Inf_Oper_R	45	49	(43.5,16.5,6)	0.64	0.0002
3	Precentral_L	6	46	(-49.5, -1.5,34.5)	0.59	0.0007
4	Precentral_L	6	237	(-24, -18,66)	0.68	0.0001
5	Precentral_R	6	80	(31, -25,72)	0.67	0.0001
6	Frontal_Mid_L	10	82	(-33,48,12)	0.62	0.0003
7	Insula_L	47	124	(-33,15, -9)	0.64	0.0002
8	Insula_L	13	107	(-31.5,9,10.5)	0.63	0.0002
9	Insula_L	13	44	(-42, -10.5,7.5)	0.64	0.0002
10	Frontal_Inf_Oper_R	44	49	(51,7.5,21)	0.59	0.0007
11	Frontal_Inf_Oper_R	46	96	(49.5,16.5,28.5)	-0.62	0.0003
12	Frontal_Mid_R	10	102	(31.5,36,30)	0.59	0.0009
13	Paracentral_Lobule_L	6	46	(-15,21,51)	-0.64	0.0002
14	Supp_Motor_Area_L	6	48	(-10.5,6,54)	0.63	0.0003
15	Frontal_Mid_L	6	119	(-24,4.5,55.5)	-0.67	0.0001
16	Frontal_Sup_R	6	136	(29, -9,65)	0.57	0.0014
17	Frontal_Sup_MedialR	9	90	(8,51,43)	-0.59	0.0009
18	Frontal_Mid_R	9	108	(50,19,41)	-0.67	0.0001
19	Occipital_Mid_L	19	47	(-43, -83,9)	-0.64	0.0002
20	Occipital_Mid_L	37	45	(-40.5, -63,4.5)	-0.62	0.0004
21	Temporal_Mid_R	39	56	(45, -57,4.5)	-0.71	0.0000
22	Temporal_Mid_L	39	54	(-45, -48,12)	-0.55	0.0021
23	Temporal_Mid_L	40	102	(-48, -55.5,16.5)	-0.65	0.0001
24	Temporal_Mid_R	21	105	(64, -2, -19)	-0.61	0.0004
25	Temporal_Sup_R	22	228	(65, -42,11)	-0.6	0.0006
26	Precuneus_L	30	155	(1.5, -51,16.5)	-0.64	0.0002
27	Cingulum_Ant_R	32	41	(9,15,27)	0.68	0.0000
28	Cingulum_Mid_R	31	73	(15, -46.5,36)	-0.63	0.0002
29	Cingulum_Mid_L	23	80	(-3, -15,30)	0.63	0.0003
30	Lingual_R	18	46	(21, -93, -16)	0.58	0.0009
31	Parietal_Sup_L	7	54	(-19.5, -63,55.5)	0.66	0.0001
32	Parietal_Sup_L	7	48	(-23, -71,58)	0.63	0.0003
33	Postcentral_L	40	41	(-31.5, -39,58.5)	0.68	0.0001
34	Postcentral_L	40	40	(-48, -36,58)	0.71	0.0000
35	Thalamus_L	Wm	108	(-24, -31.5,12)	0.64	0.0002
36	Parietal_Sup_L	5	37	(-37, -48,63)	0.59	0.0008
37	Vermis_10	-	37	(-3, -43, -37)	-0.68	0.0001
38	-	-	40	(12, -19.5, -42)	-0.68	0.0000

233 Single voxel threshold $P < 0.01$ (t > 2.771 or t < -2.771), adjacent size ≥ 37 voxels (AlphaSim corrected).

235 **Table supplement 2. FCs of voxels showing significant correlation with SI across subjects in**

FC number	Connected regions	BA	Size	Peak coordinate	r	P
				MNI (x, y, z)		
1	Frontal_Inf_Oper_R	46	80	(48,15,28.5)	-0.65	0.0001***
2	Precentral_R	4/6	106	(33, -25.5,63)	0.72	0.0000***
3	Precentral_L	6	26	(-30, -24,72)	0.66	0.0001***

236 **frontal cortex.**

237 Single voxel threshold $P < 0.005$ (t > 3.057 or t < -3.057), adjacent size $\geq$ 22 voxels (AlphaSim
238 corrected).

240 **Table supplement 3. FCs of voxels showing significant correlation with SI across subjects in**
241 **whole Brain.**

FC number	Connected regions	BA	Size	Peak coordinate	r	P
				MNI (x, y, z)		
1	Cerebelum_Crus1_L	18	70	(-29, -85, -24)	0.51	0.0052
2	Cerebelum_6_R	18	39	(12, -85, -17)	0.65	0.0002
3	Calcarine_R	18	70	(15, -91.5,12)	0.63	0.0003
4	Frontal_Inf_Oper_R	46	127	(48,15,28.5)	-0.65	0.0001
5	Precentral_R	4/6	179	(32, -24,70)	0.71	0.0001
6	Precentral_L	6	59	(-30, -23,72)	0.66	0.0001
7	-	-	37	(26, -37, -11)	-0.67	0.0001

Single voxel threshold $P < 0.01$ (t > 2.771 or t < -2.771), adjacent size $\geq$ 37 voxels (AlphaSim
corrected).

Figure 4. Significant FCs from connectivity-behavior analyses in *a priori* search

space. The seed region is the left MT+. The significant FCs are obtained from *a priori* space

(frontal cortex). (**a**) The significant FCs obtained from connectivity-BDT analysis. Single voxel

threshold *P* < 0.005, adjacent size ⩾ 23 (*AlphaSim* correcting, Methods). (**b**) The significant FCs

obtained from connectivity-SI analysis. Single voxel threshold *P* < 0.005, adjacent size ⩾ 22

(*AlphaSim* correcting, Methods). Positive correlations are shown in warm colors, while, negative

correlations are shown in cold colors.

Figure supplement 5. Significant FCs searched from connectivity-behavior (BDT/SI) analyses

in the whole brain. The seed region is the left MT+. The significant FCs are obtained from the

entire brain search, single voxel threshold $P < 0.01$, adjacent size ≥ 37 voxels (AlphaSim

correcting, Methods). Positive correlations are shown in warm colors, while, negative correlations

are shown in cold colors. **(a)** the significant FCs obtained from connectivity-BDT analysis. **(b)** the

significant FCs obtained from connectivity-SI analysis.

263 **Table 2. Correlations between FC in Table 1 and GABA/Glu concentrations in MT+.**

FC number	MT+ GABA concentrations			MT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	-0.07	0.72	0.75	-0.11	0.58	0.85
2	-0.28	0.14	0.36	-0.27	0.15	0.81
3	-0.13	0.52	0.59	-0.07	0.71	0.85
4	0.10	0.59	0.64	0.12	0.54	0.85
5	0.14	0.48	0.58	0.24	0.21	0.81
6	0.31	0.11	0.33	0.15	0.43	0.85
7	0.20	0.30	0.48	0.07	0.74	0.85
8	0.14	0.45	0.58	0.05	0.79	0.85
9	0.20	0.30	0.48	0.10	0.60	0.85
10	-0.13	0.49	0.58	-0.16	0.41	0.85
11	**-0.56**	**0.0018**	**0.02***	-0.22	0.25	0.81
12	0.18	0.34	0.51	0.15	0.43	0.85
13	0.20	0.30	0.48	0.05	0.81	0.85
14	0.39	0.04	0.14	0.22	0.24	0.81
15	-0.40	0.03	0.12	-0.21	0.27	0.81
16	-0.27	0.15	0.36	-0.12	0.53	0.85
17	0.17	0.37	0.52	0.06	0.74	0.85
18	0.26	0.18	0.39	0.16	0.40	0.85
19	0.39	0.03	0.12	0.31	0.10	0.81
20	0.01	0.98	0.98	0.14	0.46	0.85
21	0.40	0.03	0.12	0.24	0.21	0.81
22	0.22	0.25	0.48	0.06	0.76	0.85
23	**0.69**	**0.0001**	**0.002****	0.47	0.01	0.24
24	0.41	0.03	0.12	0.001	0.97	0.97

264 Bold font indicates the significant correlations survived from multi correlation correction.

266 **Table supplement 4. Correlations between FC in Table supplement 1 and GABA/Glu**

267 **concentrations in MT+**

FC number	MT+ GABA concentrations			MT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	-0.13	0.49	0.55	-0.08	0.68	0.84
2	0.17	0.38	0.49	0.05	0.82	0.88
3	0.39	0.04	0.22	0.22	0.24	0.69
4	0.41	0.03	0.19	0.24	0.21	0.69
5	**0.69**	**0.0001**	**0.0038****	0.47	0.01	0.38
6	0.2	0.3	0.41	0.1	0.6	0.81
7	0.26	0.17	0.34	0.21	0.28	0.69
8	0.22	0.24	0.35	0.25	0.19	0.69
9	0.23	0.23	0.35	0.02	0.9	0.92
10	-0.13	0.49	0.55	-0.16	0.41	0.70
11	**-0.56**	**0.0018**	**0.03***	-0.22	0.25	0.69
12	0.2	0.3	0.41	0.05	0.81	0.88
13	-0.35	0.06	0.25	0.07	0.73	0.84
14	0.39	0.03	0.19	0.31	0.1	0.69
15	0.006	0.98	0.98	0.14	0.46	0.70
16	0.26	0.18	0.34	0.16	0.4	0.70
17	0.15	0.42	0.51	0.04	0.83	0.88
18	-0.4	0.03	0.19	-0.21	0.27	0.69
19	-0.37	0.05	0.24	-0.15	0.44	0.70
20	-0.16	0.39	0.49	-0.19	0.33	0.70
21	-0.31	0.1	0.28	-0.29	0.13	0.69
22	-0.22	0.24	0.35	-0.07	0.72	0.84
23	-0.1	0.6	0.65	-0.12	0.53	0.77
24	-0.09	0.64	0.68	-0.09	0.63	0.83
25	-0.3	0.11	0.28	-0.14	0.46	0.70
26	-0.08	0.66	0.68	-0.08	0.69	0.84
27	0.28	0.13	0.29	0.1	0.59	0.81
28	-0.27	0.16	0.34	-0.2	0.29	0.69
29	0.33	0.08	0.28	0.25	0.19	0.69
30	0.24	0.21	0.35	0.19	0.32	0.70
31	0.24	0.22	0.35	0.36	0.05	0.69
32	0.23	0.24	0.35	0.23	0.23	0.69
33	0.44	0.02	0.19	0.18	0.35	0.70
34	0.32	0.09	0.28	0.002	0.99	0.99
35	0.32	0.09	0.28	0.31	0.1	0.69
36	0.30	0.11	0.28	0.22	0.25	0.69
37	-0.29	0.13	0.29	-0.23	0.23	0.69
38	-0.13	0.48	0.55	-0,15	0.42	0.70

268 Bold font indicates the significant correlations survived from multi correlation correction.

269 23

270 **Table supplement 5. Correlations between FCs in Table supplement 2 and GABA/Glu**
271 **concentrations in MT+.**

FC number	MT+ GABA concentrations			MT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	**-0.56**	**0.0017**	**0.0017****	-0.21	0.27	0.27
2	**0.70**	**0.0001**	**0.0003*****	**0.49**	**0.007**	**0.021***
3	**0.65**	**0.0001**	**0.0002****	**0.40**	**0.03**	**0.045***

272 *: $P_{FDR} < 0.05$; **: $P_{FDR} < 0.01$; ***: $P_{FDR} < 0.001$; Bold font indicates the significant correlations
273 survived from multi correlation correction.

275 **Table supplement 6. Correlations between FCs in Table supplement 3 and GABA/Glu**
276 **concentrations in MT+.**

FC number	MT+ GABA concentrations			MT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	**0.37**	**0.049**	**0.049***	0.32	0.09	>0.05
2	**0.37**	**0.049**	**0.049***	0.41	0.025	>0.05
3	**0.48**	**0.008**	**0.01***	0.21	0.28	>0.05
4	**-0.58**	**0.001**	**0.002****	-0.22	0.26	>0.05
5	**0.69**	**0.0001**	**0.0007**	0.46	0.01	>0.05
6	**0.66**	**0.0001**	**0.0004*****	0,39	0.04	>0.05
7	**-0.48**	**0.0077**	**0.01***	-0.35	0.067	>0.05

277 *: $P_{FDR} < 0.05$; **: $P_{FDR} < 0.01$; ***: $P_{FDR} < 0.001$; Bold font indicates the significant correlations
278 survived from multi correlation correction.

Local MT+ GABA acts on SI and BDT via global MT-frontal connectivity

To determine whether local neurotransmitter levels (such as GABA and Glu) in the MT+ region mediate the broader 3D visuo-spatial ability of BDT, which as a component of gF, is linked to the frontal cortex(Fangmeier et al., 2006), we correlated the significant FCs of MT – frontal in Figure 4a (also shown in Table 1) with the GABA and Glu levels in MT+ region. The results revealed that only two FCs significantly correlated with inhibitory GABA levels in MT+: 1) the FC of left MT+ - right BA 46 (significantly negative correlation, $r = - 0.56$, $P = 0.02$, $n = 29$, FDR correction, Figure 5a left); 2) the FC of left MT+ - right BA 6 (significantly positive correlation, $r = 0.69$, $P = 0.002$, $n = 29$, FDR correction, Figure 5b left) (also shown in Table 2). There were no significant correlations between these FCs and the excitatory Glu levels in MT+ (Table 2). Across the whole brain search, we obtained the same two MT+ - frontal FCs significantly correlating with both MT+ GABA levels and BDT (Table supplement 4), this is consistent with the results in a priori search space (frontal cortex) (Table 2). We then correlated the significant FCs in Figure 5b (also in Table supplement 2) with GABA and Glu concentrations in MT+ and found that almost all the correlations are significant except one (between the FC of left MT+ - right BA46 and the Glu levels in MT+) (Table supplement 5). Among the three FCs, the clusters of two FCs have substantial voxel overlap with the FCs we found by the connectivity-BDT analysis (Figure 5a, b). Across the whole brain search, there were total 7 brain regions in which FCs strength were significantly correlated with SI, all the 7 FCs significantly correlated the MT+ GABA levels, while, no FC

had significant correlation with the MT+ Glu levels (Table supplement 6). Taken

together, our results displayed that the overlap FCs from the analyses of connectivity -

behavior (BDT and SI) -GABA are the MT+ - BA 46 and MT+ - BA 6 (Figure 5a, b).

These results suggest that the FCs of MT+ - frontal regions (BA 46 and BA 6)

coupling with local MT+ GABA underlie the neural basis for both the simple motion

perception (quantified by SI) and the complex 3D visuo-spatial ability (quantified by

BDT).

In order to fully investigate the potential roles of multivariable contributing to BDT

scores, serial mediation analyses(Hayes, 2013) were applied to the MR and behavioral

data. Following our hypothesis, the independent variable (X) is MT+ GABA, the

dependent variable (Y) is BDT scores, the covariate is the age, and the mediators are

FC (M1) and SI (M2). We used the overlap clusters from the analyses of connectivity-

BDT-GABA and connectivity-SI-GABA to compute the FC of MT+ - BA46, 1

participant was excluded due to his age greater than mean + 2.5SD. The serial

mediation model is shown in Figure 5c. GABA levels in MT+ significantly negatively

correlated with the FC of MT+ - BA46 (β = - 0.32, P < 0.001), which in turn

significantly negatively correlated with SI (β = - 0.19, P = 0.035), and consequently,

significantly positively correlated with BDT (β = 38.5, P = 0.009). Critically,

bootstrapped analyses revealed that our hypothesized indirect effect (i.e., MT+ GABA

$\rightarrow$ FC of MT+ - BA46 $\rightarrow$ SI $\rightarrow$ BDT) was significant (β = 2.28, SE = 1.54, 95% CI =

[0.03, 5.94]). The model accounted for 34% of the variance in BDT. However, when

considering the MT-BA6 FC as the mediator M1, the serial model does not show a

significant indirect effect. Consequently, we explored a mediation model, which

revealed that the MT-BA6 FC totally mediates the relationship between GABA and

BDS. (Figure 5d). For sensitivity purposes, we tested the alternative models, in which

the order of the mediators was reversed. The pathway that MT+ GABA was predicted

to be associated with SI, followed by the FC of MT+ - BA 46, and then BDT, did not

yield the chained mediation effects on BDT (figure supplement 6).

Figure 5. Local MT+ GABA acts on SI and BDT via global MT-frontal connectivity. **(a)** Significant negative correlation between the FC of left MT+ - right DLPFC (BA46) and MT+ GABA (*FDR* correction). **(b)** Significant positive correlation between the FC of left MT+ - right (pre) motor cortex (BA4/6) and MT+ GABA (*FDR* correction). In **(a)** and **(b)**, left: the significant FCs obtained from connectivity-BDT analysis; right: the significant FCs obtained from connectivity-SI analysis. **(c)** Significant pathways: MT+ GABA→ FC (left MT+ - right BA46, negative correlation) → SI (negative correlation) → BDT (positive correlation). This pathway can explain 34% of the variance in BDT. **(d)** Significant pathways: MT+ GABA → FC (left MT+ - right BA6, positive correlation) → BDT (positive correlation).The bolded lines represent the hypothesized mediation effect. The dotted lines represent alternative pathways. [*]: P < 0.05; [**]: P < 0.01; [***]: P < 0.001.

Figure supplement 6. The alternative serial mediation models from local MT+ GABA to global

performance of BDT. The pathway that MT+ GABA was predicted to be associated with SI,

followed by the FC of MT+ - BA 46, and then BDT, did not yield the chained mediation effects

on BDT. *: P < 0.05; **: < 0.01; ***: P < 0.001.

To summarize (shown in Figure 6), the results from the serial mediation analyses

are consistent with our hypothesis. That is, higher GABAergic inhibition in MT+

relates to stronger negative FC between MT+ and BA46, leading to enhanced ability

for surround suppression (filtering out irrelevant information(Tadin, 2015), ultimately

resulting in more efficient visual 3D processing of gF (the higher BDT scores).

Figure 6. Sketch depicting the multi-level inhibitory mechanisms centered on MT+ GABA

contributing to visuo-spatial intelligence. Inhibitory GABA in MT+ (a sensory cortex, shown in

green circle), coupling with the conjunctive coding between MT+ and BA46 (cognitive control

core, shown in purple circle), and mediated by motion surround suppression (shown in red circle),

contributes to visuo-spatial intelligence (BDT, 3D domain, shown in red and white building

blocks). In this sketch, the two-colored parallel arrows show the negative FC between MT+ and

BA46, the colored arrows below the green circle display the inhibition mechanisms centered on

MT+ GABA (2), filtered the irrelevant information in (1) and focused on the efficient visuo-

spatial information (3). Black long arrows display the direction of information flow: from input

information (1) to visuo-spatial intelligence (4).

Discussion

Here, we provide evidence that MT+ inhibitory mechanisms mediate processing in the visuo-spatial component (3D domain) of gF on multiple levels, that is, from molecular over brain connectivity to behavior. First, this study found that higher MT+ inhibitory GABA levels (but not excitatory Glu) relate to FC between MT+ and BA 46 that contribute to both SI and BDT. Our serial mediation analyses indicate that the inhibitory mechanisms linking to MT+, including GABA levels in MT+ (but not Glu), FCs of MT+ - BA46 coupling with MT+ inhibitory GABA (but not excitatory Glu), and behavior (SI indexing perceptual suppression in MT+) predict the inter-subject variance in the 3D gF task (BDT) (Figure 5c). Second, we demonstrate discrete GABAergic inhibition mechanisms in MT+ that mediate the strong FCs between MT+ - frontal regions (BA46 and BA6): significant negative correlation with the FC of MT+ - BA 46 (Figure 5a), whereas significant positive correlation with the FC of MT+ - BA 6 (Figure 5b). This indicates that different frontal regions, DLPFC (BA 46) and premotor cortex (BA6), contribute uniquely to gF through MT+ - based inhibitory mechanisms.

The goal of our research is to reveal that the inhibitory (not excitatory) mechanism in MT+ contributes to multi-level processing in 3D visuo-spatial ability (BDT). Monkey electrophysiological experiments revealed that selective attention gates the visual cortex, including area MT, effectively suppressing the irrelevant information(Everling et al., 2002; Treue & Maunsell, 1996). These findings aligns with the "neural efficiency" hypothesis of intelligence(Haier et al., 1988), which puts

forward the human brain's ability to suppress the repetition of information. Neural

suppression is associated with the balance between excitation and inhibition (EIB),

usually represented by covariation between Glutamate and GABA(Ozeki et al., 2009).

Here, this study exploited the high spectral resolution afforded by ultrahigh field (7T)

MRS to reliably resolve GABA measurement, adequately discriminate the glutamate

and glutamine signals, and resolve the high accuracy Glu measurement(Ende, 2015).

This work implemented the MRS scanning in MT+ (3D visual domain) and V1 (2D

visual domain) regions and found that MT+ inhibitory GABA (but not excitatory Glu)

significantly correlated with BDT, i.e., the higher GABA levels in MT+ (rather than

excitatory Glu) relate to higher visual 3D processing (BDT) (Figure 3c). We searched

the global MT+ - based FCs with the connectivity-BDT analyses (in priori search

space and whole brain search to valid), and then, correlated these significant FCs with

the GABA and Glu concentrations in MT+, and found two FCs (MT+ - BA46 and

MT+ - BA6) significantly correlating with MT+ inhibitory GABA (whereas no FC

significantly correlated with MT+ excitatory Glu). Accordingly, our results emphasize

the importance of MT+ inhibitory GABA (but not excitatory Glu) in processing the

3D visual-spatial intelligence (BDT). Crucially, a previous study has reported a strong

link between motion spatial suppression (SI) and the performance of BDT(Melnick et

al., 2013). Our recent human study(Song et al., 2021) and other study's animal

experiments(Ozeki et al., 2009; Sato et al., 2016) demonstrated that the conjoint

action of inhibition (GABA) and excitation (Glu) underlies visual spatial suppression.

In this work, our novel data show the chained mediation effects from local MT+

GABA to more global BDT: MT+ GABA → FC (MT+ and BA46) → SI → BDT.

Thereby, our data indicate that inhibitory mechanisms in MT+, from the biochemistry

over FC to the behavioral level, can predict the inter-subject variance in the 3D gF

task (BDT) (Figure 5c).

Another interesting finding reveal that GABAergic inhibition in MT+ coupling

with distinct brain connectivity mechanisms between BA 46-MT+ and BA6-MT+. A

previous human fMRI experiment found that the positive and negative correlations

between BDT and the activation of frontal regions appeared at different reasoning

phases (validation or integration phases during reasoning)(Fangmeier et al., 2006). On

the one hand, a monkey electrophysiological experiment reported the delayed

modulation from PFC (especially in DLPFC (BA 46))) to area MT during a visual

motion task(Zaksas & Pasternak, 2006). Computational models converged with

empirical data of awake monkey experiments present slow temporal modulation from

PFC to MT/MST(Donner et al., 2009; Siegel et al., 2015; Wang, 2002; Wimmer et al.,

2015). On the other hand, human MEG studies(Donner et al., 2009; Wilming et al.,

2020) reported that the gamma-band activity in the visual cortex (including area MT)

exhibited high coherence with the activity in (pre-) motor regions (BA 4/6). These

works suggest that the relation of long-range FC and local inhibition mechanism

(MT+ GABA) might indicate that inhibition in MT+ contributes to efficient long-

range integration and coordination in distant brain areas like the prefrontal and

premotor cortex.

How does MT+ assemble into the MD system as an intellectual hub rather than a

simple input module? The results in Figure 5a, c showed that the overlap brain

regions from the analyses of connectivity-BDT-GABA/connectivity-SI-GABA are the

MT+ - BA 46. This overlap couples with local visual suppression (SI) and

consequently plays an important role in intelligence (BDT). The direction

discrimination task in this work (the visual motion paradigm of center-surround

antagonism) was previously considered a mainly local function of MT+(Melnick et

al., 2013; Tadin, 2015; Tadin et al., 2003). However, our results with connectivity-SI

analyses revealed that both local (FC within BA 18) and global brain connectivity (FC

between MT+ and frontal regions) contribute to SI (Table supplement 3). In human

psychophysical experiments(Melnick et al., 2013; Tadin et al., 2003) the brief

stimulus duration (~100 ms) in motion discrimination precludes most top-down

attentional effects(Wang, 2002; Zaksas & Pasternak, 2006), while, attention, which

predicted the performance of the motion discrimination task, was sustained

throughout the stimulus intervals(Siegel et al., 2015). Furthermore, animal

experiments have revealed that the local circuits in the visual cortex combing with the

top-down modulation and intracortical horizontal connection mediate the visual-

spatial suppression(Angelucci et al., 2002; Keller et al., 2020; Li et al., 2019; Zhang et

al., 2014). Our results (shown in Figure 5a, b, right) present the intrinsic binding of

local GABAergic inhibition in MT+, which suppressing redundancy of visual motion

processing (SI), and the activity of brain connectivity between MT+ and frontal

regions. These individually inherent traits may contribute to the individual difference

in 3D visuo-spatial ability (Figure 5a, b, left). A candidate divisive normalization

model(Carandini & Heeger, 2011; Reynolds & Heeger, 2009) can explain how such

reverberation affects the process of suppressing the irrelevant information, from

perception to intelligence(Melnick et al., 2013; Tadin, 2015). We summarize a

framework (Figure 6) to indicate and visualize our findings.

Together, this study offers a comprehensive insight into how the information

exchange and integration between the sensory cortex (MT+) and cognition core of BA

46, coupling with the MT+ GABA, can predict the performance of 3D visuo-spatial

ability (BDT). Our results provide direct evidence that a sensory cortex (MT+), at

biochemistry, connectivity, and behavior levels, can assemble into complex cognition

as an intellectual hub.

Materials and Methods

Subjects

Thirty-six healthy subjects (18 female, mean age: 23.6 years$\pm$2.1, range: 20 to 29

years) participated in this study, they were recruited from Zhejiang University. All

subjects had normal or corrected-to-normal vision. In addition, they reported no

psychotropic medication use, no illicit drug use within the past month, no alcohol use

within 3 days prior to scanning, and right-handed. This experiment was approved by

the Ethics Review Committee of Zhejiang University and conducted in accordance

with the Helsinki Declaration. All participants signed informed consent forms prior to

the start of the study and were compensated for their time. All subjects participated in

the motion spatial suppression psychophysical, resting-state fMRI and MRS (MT+

and V1 regions, in random sequence) experiments, but only part of the MRS data

(31/36 in MT+ region and 28/36 in V1 region) survived quality control (see the part

of MRS date processing). The sample size is determined by the statistic requirement

(30 sample for Person correlation statistical analysis).

Motion surrounding suppression measurement

All stimuli were generated using Matlab (MathWorks, Natick, MA) with

Psychophysics Toolbox(Brainard, 1997), and were shown on a linearized monitor

(1920×1080 resolution, 100-Hz refresh rate, Cambridge Research System, Kent, UK).

The viewing distance was 72 cm from the screen, with the head stabilized by a

chinrest. Stimuli were drawn against a gray (56 cd per m^{-2}) background.

A schematic of the stimuli and trial sequences is shown in our recent study(Song et

al., 2021). The stimulus was a vertical drifting sinusoidal grating (contrast, 50%;

spatial frequency, 1 cycle/^{o}; speed, 4°/s) of either small (diameter of 2°) or large

(diameter of 10°) size. The edge of the grating was blurred with a raised cosine

function (width, 0.3°). A cross was presented in the center of the screen at the

beginning of each trial for 500ms, and participants were instructed to fixate at the

cross and to keep fixating at the cross throughout the trial. In each trial, a grating of

either large or small size was randomly presented at the center of the screen. The

grating drifted either leftward or rightward, and participants were asked to judge the

perceived moving direction by a key press. Response time was not limited. The

grating was ramped on and off with a Gaussian temporal envelope, and the grating

duration was defined as 1 SD of the Gaussian function. The duration was adaptively

adjusted in each trial, and duration thresholds were estimated by a staircase procedure.

Thresholds for large and small gratings were obtained from a 160-trial block that

contained four interleaved 3-down/1-up staircases. For each participant, we computed

the correct rate for different stimulus durations separately for each stimulus size.

These values were then fitted to a cumulative Gaussian function, and the duration

threshold corresponding to the 75% correct point on the psychometric function was

estimated for each stimulus size.

Stimulus demonstration and practice trials were presented before the first run.

Auditory feedback was provided for each wrong response. To quantify the spatial

suppression strength, we calculated the spatial suppression index (SI), defined as the

difference of log10 thresholds for large versus small stimuli(Schallmo et al., 2018;

Tadin et al., 2003):

$$SI = log_{10}(\text{large threshold}) - log_{10}(\text{small threshold}) \qquad (1)$$

Block design task measurement

The block design task was administered in accordance with the WAIS-IV

manual(Wechsler, 2008). Specifically, participants were asked to rebuild the figural

pattern within a specified time limit using a set of red and white blocks. The time

limits were set as 30 s to 120 s according to different levels of difficulty. The patters

were presented in ascending order of difficulty, and the test stopped if two

consecutive patterns were not constructed in the allotted time. The score was

determined by the accomplishment of the pattern and the time taken. A time bonus

was awarded for rapid performance in the last six patterns. The score ranges between

0 and 66 points, with higher scores indicating better perceptual reasoning.

MR experimental procedure

MR experiments were performed in a 7T whole body MR system (Siemens

Healthcare, Erlangen, Germany) with a Nova Medical 32 channel array head coil.

Sessions included resting-state functional MRI, fMRI localizer scan, structural image

scanning, and MRS scan. Resting-state scans were acquired with 1.5-mm isotropic

resolution (transverse orientation, TR/TE = 2000/20.6 ms, 160 volumes, slice number

= 90, flip angle = 70^o, eyes closed). Structural images were acquired using a

MP2RAGE sequence (TR/ TI1/ TI2 = 5000/901/3200 ms) with 0.7-mm isotropic

resolution. MRS data were collected within two regions (MT+ and V1) for each

subject, and we divided them into two sessions to avoid discomfort caused by long

scanning. The order of MRS VOIs (MT+ and V1) in the two sessions was

counterbalanced across participants. Interval between two sessions was used for block

design and motion discrimination tasks. One session included fMRI localizer scan,

structural image scanning, and MRS scan for the MT+ region; the other session

included structural image scan, and MRS scan for the V1 region. Spectroscopy data

were acquired using a ^{1}H-MRS single-voxel short-TE STEAM (Stimulated Echo

Acquisition Mode) sequence(Frahm et al., 1989) (TE/TM/TR = 6/32/7100ms) with

4096 sampling points, 4-kHz bandwidth, 16 averages, 8 repetitions, 20×20×20 mm^3

VOI size, and VAPOR (variable power and optimized relaxation delays) water

suppression(Tkáč et al., 1999). Prior to acquisition, first- and second-order shims were

adjusted using FASTMAP (fast, automatic shimming technique by mapping along

projections(Gruetter, 1993). Two non-suppressed water spectra were also acquired:

one for phase and eddy current correction (only RF pulse, 4 averages) and another for

metabolite quantification (VAPOR none, 4 averages). Voxels were positioned based

on anatomical landmarks using a structural image scan collected in the same session,

while avoiding contamination by CSF, bone, and fat. The MT+ VOIs were placed in

the ventrolateral occipital lobe, which was based on anatomical landmarks(Dumoulin

et al., 2000; Schallmo et al., 2018). We did not distinguish between the middle

temporal (MT) and medial superior temporal (MST) areas in these MT+ VOIs(Huk et

al., 2002). For 14 subjects, we also functionally identified MT+ as a check on the

placement of the VOI. A protocol was used with a drifting grating (15% contrast)

alternated with a static grating across blocks (10 s block duration, 160 TRs total).

Using fMRI BOLD signals, these localizer data were processed online to identify the

MT+ voxels in the lateral occipital cortex, which responded more strongly to moving

vs. static gratings. In addition, we only used the left MT+ as the target region to scan,

which was motivated by studies showing that left MT+ was more effective at causing

perceptual effects(Tadin et al., 2011). For V1 region, the VOI was positioned on each

subject's calcarine sulcus on the left side(Tadin et al., 2011) based on anatomical

landmarks(Boucard et al., 2007; Dumoulin et al., 2000)

MRS data processing

Spectroscopy data were preprocessed and quantified using magnetic resonance signal

processing and analysis, https://www.cmrr.umn.edu/ downloads/mrspa/), which runs

under MATLAB and invokes the interface of the LCModel (Version 6.3-1L)(Chen et

al., 2019). First, we used the non-suppressed water spectra to perform eddy current

correction and frequency/phase correction. Second, we checked the quality of each

FID (16 averages) visually and removed those with obviously poor quality. Third, the

absolute concentrations of each metabolite were quantitatively estimated via the

Water-Scaling method. For partial-volume correction, the tissue water content was

computed as follows(Ernst et al., 1993):

575   *Tissue water content=fgm*0.78 + fwm*0.65 + fcsf*0.97*   (2)

where *fgm*, *fwm*, and *fcsf* were the GM/WM/CSF volume fraction in MRS VOI and

we used FAST (fMRI's automated segmentation tool, part of the FSL toolbox)(Zhang

et al., 2001) to segment the three tissue compartments from the T1-weighted structural

brain images. For water T2 correction, we set water T2 as 47ms (Marjańska et al.,

2012). Our concentrations were mM per kg wet weight. Furthermore, LCModel

analysis was performed on all spectra within the chemical shift range of 0.2 to 4.0

ppm.

Poor spectral quality was established by a Cramer-Rao Lower Bound (CRLB) of

more than 20%(Cavassila et al., 2001), and some data were excluded from further

analysis. The details were described in our recently paper(Song et al., 2021).

Rs-fMRI data processing and analysis

Resting-state functional image was analyzed in the Data Processing and Analysis for

Brain Imaging DPABI toolbox(Yan et al., 2016) based on SPM 12

(http://www.fil.ion.ucl.ac.uk/spm/). The preprocessing steps included discard of the

first five volumes, slice timing, realignment to the 90th slice, coregistration of each

subject's T1-weighted anatomical and functional images, segmentation of the

anatomical images into six types of tissues using DARTEL, linear detrend, regressing

nuisance variables (including realignment Friston 24-parameter, global signal, white

matter and CSF signal) (Friston et al., 1996), normalization to the standard Montreal

Neurological Institute (MNI) space with the voxel size of $1.5 \times 1.5 \times 1.5$ mm^3 using

DARTEL, spatial smoothing with a Gaussian kernel of 3 mm full-width-half-

maximum (FWHM), and band-pass filtering with Standard frequency band (SFB,

0.01–0.1 Hz). Spherical ROI with a radius of 6mm was placed in left MT. The

coordinate for left MT (-46, -72, -4, in MNI space) was obtained by our localizer

fMRI experiment. We calculated the seed-to-voxel whole brain FC map for each

subject. All the FC values were Fisher-Z-transformed.

We did a similar connectivity-behavior analysis to a previous study (Song et al.,

2008). First, we computed the Pearson's correlation coefficient between BDT scores

and the FC values across subjects in a voxel-based way. Then, to evaluate the

significance, we transformed the *r*-value into *t*-value ($t = \frac{\sqrt{df} \cdot r}{\sqrt{1-r^2}}$), where *df* denotes to

the degrees of freedom, and *r* is the Pearson's correlation coefficient between BDT

scores and the FC values. Here, *df* was equal to 27. The brain regions in which the FC

values to the seed region was significantly correlated with the BDT scores were

obtained with a threshold of *P* < 0.005 for regions of *a priori* ($|t_{(27)}| \geq 3.057$, and

adjacent cluster size ≥ 23 voxels (AlphaSim corrected)), and *P* < 0.01 for whole-brain

analyses ($|t(27)| \geq 2.771$ and adjacent cluster size ≥ 37 voxels (AlphaSim corrected).

Statistical Analysis

PROCESS version 3.4, a toolbox in SPSS, was used to examine the mediation model.

There are some prerequisites for mediation analysis: the independent variable should

be a significant predictor of the mediator, and the mediator should be a significant

predictor of the dependent variable.

SPSS 20 (IBM, USA) was used to conduct all the remaining statistical analysis in

the study. We evaluated the correlation of variables (GABA, Glu, SI, BDT) using

Pearson's correlation analysis. Differences or correlations were considered

statistically significant if *P* < 0.05. Significances with multiple comparisons were

tested with false discovery rate (*FDR*) correction. The effect of age on intelligence

was controlled for by using partial correlation in the correlation analysis and was

taken as a covariate in the serial mediation model analysis.

Data availability

628 All data needed to evaluate the conclusions in the paper are present in the paper

629 and/or the Supplementary information. Source data are provided with this paper and

630 have been archived at GitHub. They could be downloaded with reasonable request.

631 .

Code availability

633 This paper does not report original code.

References

Angelucci, A., Levitt, J. B., Walton, E. J., Hupe, J.-M., Bullier, J., & Lund, J. S. (2002). Circuits for local and
global signal integration in primary visual cortex. *Journal of Neuroscience*, *22*(19), 8633-8646.
Assem, M., Glasser, M. F., Van Essen, D. C., & Duncan, J. (2020). A domain-general cognitive core
defined in multimodally parcellated human cortex. *Cerebral Cortex*, *30*(8), 4361-4380.
Barbey, A. K. (2018). Network neuroscience theory of human intelligence. *Trends in cognitive sciences*,
22(1), 8-20.
Bedny, M., Konkle, T., Pelphrey, K., Saxe, R., & Pascual-Leone, A. (2010). Sensitive period for a
multimodal response in human visual motion area MT/MST. *Current Biology*, *20*(21), 1900-
1906.
Born, R. T., & Bradley, D. C. (2005). Structure and function of visual area MT. *Annu Rev Neurosci*, *28*,
157-189. https://doi.org/10.1146/annurev.neuro.26.041002.131052
Boucard, C. C., Hoogduin, J. M., van der Grond, J., & Cornelissen, F. W. (2007). Occipital proton
magnetic resonance spectroscopy (1H-MRS) reveals normal metabolite concentrations in
retinal visual field defects. *PLoS One*, *2*(2), e222.
Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis*, *10*(4), 433-436.
Carandini, M., & Heeger, D. J. (2011). Normalization as a canonical neural computation. *Nat Rev
Neurosci*, *13*(1), 51-62. https://doi.org/10.1038/nrn3136
Cattell, & Raymond, B. (1963). Theory of fluid and crystallized intelligence: A critical experiment.
Journal of Educational Psychology, *54*(1), 1-22.
Cavassila, S., Deval, S., Huegen, C., Van Ormondt, D., & Graveron‐Demilly, D. (2001). Cramér–Rao

bounds: an evaluation tool for quantitation. *NMR in Biomedicine: An International Journal*
Devoted to the Development and Application of Magnetic Resonance In Vivo, *14*(4), 278-283.

Chen, X., Fan, X., Hu, Y., Zuo, C., Whitfield-Gabrieli, S., Holt, D., Gong, Q., Yang, Y., Pizzagalli, D. A., &
Du, F. (2019). Regional GABA concentrations modulate inter-network resting-state functional
connectivity. *Cerebral Cortex*, *29*(4), 1607-1618.

Cole, M. W., Yarkoni, T., Repovs, G., Anticevic, A., & Braver, T. S. (2012). Global connectivity of
prefrontal cortex predicts cognitive control and intelligence. *J Neurosci*, *32*(26), 8988-8999.
https://doi.org/10.1523/JNEUROSCI.0536-12.2012

Colom, R., Jung, R. E., & Haier, R. J. (2006). Distributed brain sites for the g-factor of intelligence.
Neuroimage, *31*(3), 1359-1365. https://doi.org/10.1016/j.neuroimage.2006.01.006

Deary, I. J., Penke, L., & Johnson, W. (2010). The neuroscience of human intelligence differences.
Nature reviews neuroscience, *11*(3), 201-211.

Donner, T. H., Siegel, M., Fries, P., & Engel, A. K. (2009). Buildup of choice-predictive activity in human
motor cortex during perceptual decision making. *Curr Biol*, *19*(18), 1581-1585.
https://doi.org/10.1016/j.cub.2009.07.066

Dumoulin, S. O., Bittar, R. G., Kabani, N. J., Baker, C. L., Jr., Le Goualher, G., Bruce Pike, G., & Evans, A.
C. (2000). A new anatomical landmark for reliable identification of human area V5/MT: a
quantitative   analysis   of   sulcal   patterning.   *Cereb   Cortex*,   *10*(5),   454-463.
https://doi.org/10.1093/cercor/10.5.454

Duncan, J., Assem, M., & Shashidhara, S. (2020). Integrated Intelligence from Distributed Brain
Activity. *Trends Cogn Sci*, *24*(10), 838-852. https://doi.org/10.1016/j.tics.2020.06.012

Duncan, J., Seitz, R. J., Kolodny, J., Bor, D., Herzog, H., Ahmed, A., Newell, F. N., & Emslie, H. (2000). A
neural   basis   for   general   intelligence.   *Science*,   *289*(5478),   457-460.
https://doi.org/10.1126/science.289.5478.457

Ende, G. (2015). Proton Magnetic Resonance Spectroscopy: Relevance of Glutamate and GABA to
Neuropsychology. *Neuropsychol Rev*, *25*(3), 315-325. https://doi.org/10.1007/s11065-015-
9295-8

Ernst, T., Kreis, R., & Ross, B. (1993). Absolute quantitation of water and metabolites in the human
brain. I. Compartments and water. *Journal of magnetic resonance, Series B*, *102*(1), 1-8.

Everling, S., Tinsley, C. J., Gaffan, D., & Duncan, J. (2002). Filtering of neural signals by focused
attention in the monkey prefrontal cortex. *Nature neuroscience*, *5*(7), 671-676.

Fangmeier, T., Knauff, M., Ruff, C. C., & Sloutsky, V. (2006). fMRI evidence for a three-stage model of
deductive reasoning. *Journal of cognitive neuroscience*, *18*(3), 320-334.

Frahm, J. a., Bruhn, H., Gyngell, M., Merboldt, K., Hänicke, W., & Sauter, R. (1989). Localized high‐
resolution proton NMR spectroscopy using stimulated echoes: initial applications to human
brain in vivo. *Magnetic resonance in medicine*, *9*(1), 79-93.

Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S., & Turner, R. (1996). Movement‐related
effects in fMRI time‐series. *Magnetic resonance in medicine*, *35*(3), 346-355.

Gray, J. R., Chabris, C. F., & Braver, T. S. (2003). Neural mechanisms of general fluid intelligence. *Nat*
Neurosci, *6*(3), 316-322. https://doi.org/10.1038/nn1014

Gruetter, R. (1993). Automatic, localized in vivo adjustment of all first‐ and second‐order shim coils.
Magnetic resonance in medicine, 29(6), 804-811.

Haier, R. J., Siegel Jr, B. V., Nuechterlein, K. H., Hazlett, E., Wu, J. C., Paek, J., Browning, H. L., &

699   Buchsbaum, M. S. (1988). Cortical glucose metabolic rate correlates of abstract reasoning and
700    attention studied with positron emission tomography. *Intelligence*, *12*(2), 199-217.

701 Hayes, A. F. (2013). *Introduction to Mediation, Moderation, and Conditional Process Analysis: A*
702    *Regression‐Based Approach* (Vol. 51). The Guilford Press.

703 Huk, A. C., Dougherty, R. F., & Heeger, D. J. (2002). Retinotopy and functional subdivision of human
704    areas MT and MST. *Journal of Neuroscience*, *22*(16), 7195-7205.

705 Jung, R. E., & Haier, R. J. (2007). The Parieto-Frontal Integration Theory (P-FIT) of intelligence:
706    converging neuroimaging evidence. *Behav Brain Sci*, *30*(2), 135-154; discussion 154-187.
707    https://doi.org/10.1017/S0140525X07001185

708 Keller, A. J., Roth, M. M., & Scanziani, M. (2020). Feedback generates a second receptive field in
709    neurons of the visual cortex. *Nature*, *582*(7813), 545-549.

710 Li, M., Song, X. M., Xu, T., Hu, D., Roe, A. W., & Li, C.-Y. (2019). Subdomains within orientation columns
711    of primary visual cortex. *Science advances*, *5*(6), eaaw0807.

712 Liu, D. Y., Ju, X., Gao, Y., Han, J. F., Li, Z., Hu, X. W., Tan, Z. L., Northoff, G., & Song, X. M. (2022). From
713    Molecular to Behavior: Higher Order Occipital Cortex in Major Depressive Disorder. *Cereb*
714    *Cortex*, *32*(10), 2129-2139. https://doi.org/10.1093/cercor/bhab343

715 Liu, L. D., Haefner, R. M., & Pack, C. C. (2016). A neural basis for the spatial suppression of visual
716    motion perception. *Elife*, *5*, e16167.

717 Marjańska, M., Auerbach, E. J., Valabrègue, R., Van de Moortele, P. F., Adriany, G., & Garwood, M.
718    (2012). Localized 1H NMR spectroscopy in different regions of human brain in vivo at 7 T: T2
719    relaxation times and concentrations of cerebral metabolites. *NMR in Biomedicine*, *25*(2), 332-
720    339.

721 Melnick, M. D., Harrison, B. R., Park, S., Bennetto, L., & Tadin, D. (2013). A strong interactive link
722    between sensory discriminations and intelligence. *Curr Biol*, *23*(11), 1013-1017.
723    https://doi.org/10.1016/j.cub.2013.04.053

724 Ozeki, H., Finn, I. M., Schaffer, E. S., Miller, K. D., & Ferster, D. (2009). Inhibitory stabilization of the
725    cortical network underlies visual surround suppression. *Neuron*, *62*(4), 578-592.

726 Reynolds, J. H., & Heeger, D. J. (2009). The normalization model of attention. *Neuron*, *61*(2), 168-185.

727 Ricciardi, E., Vanello, N., Sani, L., Gentili, C., Scilingo, E. P., Landini, L., Guazzelli, M., Bicchi, A., Haxby, J.
728    V., & Pietrini, P. (2007). The effect of visual experience on the development of functional
729    architecture in hMT+. *Cerebral Cortex*, *17*(12), 2933-2939.

730 Sato, T. K., Haider, B., Häusser, M., & Carandini, M. (2016). An excitatory basis for divisive
731    normalization in visual cortex. *Nature neuroscience*, *19*(4), 568-570.

732 Schallmo, M. P., Kale, A. M., Millin, R., Flevaris, A. V., Brkanac, Z., Edden, R. A., Bernier, R. A., & Murray,
733    S. O. (2018). Suppression and facilitation of human neural responses. *Elife*, *7*.
734    https://doi.org/10.7554/eLife.30334

735 Siegel, M., Buschman, T. J., & Miller, E. K. (2015). Cortical information flow during flexible
736    sensorimotor decisions. *Science*, *348*(6241), 1352-1355.
737    https://doi.org/10.1126/science.aab0551

738 Song, M., Zhou, Y., Li, J., Liu, Y., Tian, L., Yu, C., & Jiang, T. (2008). Brain spontaneous functional
739    connectivity and intelligence. *Neuroimage*, *41*(3), 1168-1176.
740    https://doi.org/10.1016/j.neuroimage.2008.02.036

741 Song, X. M., Hu, X. W., Li, Z., Gao, Y., Ju, X., Liu, D. Y., Wang, Q. N., Xue, C., Cai, Y. C., Bai, R., Tan, Z. L., &

Northoff, G. (2021). Reduction of higher-order occipital GABA and impaired visual perception in acute major depressive disorder. *Mol Psychiatry*, *26*(11), 6747-6755. https://doi.org/10.1038/s41380-021-01090-5

Spearman, C. (1904). " General Intelligence" Objectively Determined and Measured. *American Journal of Psychology*, *15*, 201-293.

Tadin, D. (2015). Suppressive mechanisms in visual motion processing: From perception to intelligence. *Vision Res*, *115*(Pt A), 58-70. https://doi.org/10.1016/j.visres.2015.08.005

Tadin, D., Lappin, J. S., Gilroy, L. A., & Blake, R. (2003). Perceptual consequences of centre-surround antagonism in visual motion processing. *Nature*, *424*(6946), 312-315. https://doi.org/10.1038/nature01800

Tadin, D., Silvanto, J., Pascual-Leone, A., & Battelli, L. (2011). Improved motion perception and impaired spatial suppression following disruption of cortical area MT/V5. *Journal of Neuroscience*, *31*(4), 1279-1283.

Tkáč, I., Starčuk, Z., Choi, I. Y., & Gruetter, R. (1999). In vivo 1H NMR spectroscopy of rat brain at 1 ms echo time. *Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine*, *41*(4), 649-656.

Treue, S., & Maunsell, J. H. (1996). Attentional modulation of visual motion processing in cortical areas MT and MST. *Nature*, *382*(6591), 539-541.

Wang, X. J. (2002). Probabilistic decision making by slow reverberation in cortical circuits. *Neuron*, *36*(5), 955-968. https://doi.org/10.1016/s0896-6273(02)01092-9

Wechsler, D. (2008). Wechsler Memory Scale–Fourth Edition (WMS-IV) technical and interpretive manual.

Wilming, N., Murphy, P. R., Meyniel, F., & Donner, T. H. (2020). Large-scale dynamics of perceptual decision information across human cortex. *Nat Commun*, *11*(1), 5109. https://doi.org/10.1038/s41467-020-18826-6

Wimmer, K., Compte, A., Roxin, A., Peixoto, D., Renart, A., & de la Rocha, J. (2015). Sensory integration dynamics in a hierarchical network explains choice probabilities in cortical area MT. *Nat Commun*, *6*, 6177. https://doi.org/10.1038/ncomms7177

Yan, C.-G., Wang, X.-D., Zuo, X.-N., & Zang, Y.-F. (2016). DPABI: data processing & analysis for (resting-state) brain imaging. *Neuroinformatics*, *14*, 339-351.

Zaksas, D., & Pasternak, T. (2006). Directional signals in the prefrontal cortex and in area MT during a working memory for visual motion task. *J Neurosci*, *26*(45), 11726-11742. https://doi.org/10.1523/JNEUROSCI.3420-06.2006

Zhang, S., Xu, M., Kamigaki, T., Hoang Do, J. P., Chang, W.-C., Jenvay, S., Miyamichi, K., Luo, L., & Dan, Y. (2014). Long-range and local circuits for top-down modulation of visual cortex processing. *Science*, *345*(6197), 660-665.

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE transactions on medical imaging*, *20*(1), 45-57.

Acknowledgements

The authors thank Prof. Dost Ongur and Fei Du for guidance on the MRS data processing, and thank Zhejiang University 7T Brain Imaging Research Center. This work was supported by STI 2030—Major Projects (2021ZD0200401 to X.M.S., 2022ZD0206000 to R.B.), the National Natural Science Foundation of China Grants (61876222, 32000761, 82222032), Humanities and Social Sciences Ministry of Education (20YJC880095, 18YJA190001), the Key R&D Program of Zhejiang (2022C03096 to X.M.S.), the European Union's Horizon 2020 Framework Program for Research and Innovation under the Specific Grant Agreement No. 785907 (Human Brain Project SGA2 to G.N.), and the MOE Frontier Science Center for Brain Science & Brain- Machine Integration, Zhejiang University.

Author contributions

S.X.M. and Y.G. designed the experiment and G.N. guided the logic of the analysis. Y.G., Y.C. and D.L. conducted human experiments, analyzed data, and Y.G. created figures. R.B. guided the MRI experiments. J.Y., J.W. and B.X. were in charge of data collection and partial date analysis. T. W., M.L. and G.C. guided the data analysis. S.X.M., Y.G. and G.N. wrote the manuscript.

Competing interests

The authors declare no competing interests.

48