GABA-ergic inhibition in human hMT+ predicts visuo-spatial intelligence mediated through the frontal cortex

Yuan Gao[1#], Yong-Chun Cai[2#], Dong-Yu Liu[1,3], Juan Yu[1,3], Jue Wang[1], Ming li[4], Bin Xu[1], Teng-Fei Wang[2], Gang Chen[1,3], Georg Northoff[5,6*], Ruiliang Bai[1,3,7*], Xue Mei Song[1,3,5,7*]

[1]Department of Neurosurgery of the Second Affiliated Hospital and Interdisciplinary Institute of Neuroscience and Technology, Zhejiang University School of Medicine, Hangzhou 310029, China;

[2]Department of Psychology and Behavioral Sciences, Zhejiang University, Hangzhou 310028, China;

[3]Key Laboratory of Biomedical Engineering of Ministry of Education, Qiushi Academy for Advanced Studies, College of Biomedical Engineering and Instrument Science, Zhejiang University, Hangzhou 310027, China;

[4]College of Intelligence Science and Technology, National University of Defense Technology, Changsha 410073, China;

[5]Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, Zhejiang University School of Medicine, Hangzhou 310013, China;

[6]University of Ottawa Institute of Mental Health Research, University of Ottawa; Ottawa, ON, K1Z 7K4, Canada;

[7]MOE Frontier Science Center for Brain Science & Brain- Machine Integration, Zhejiang University，Hangzhou 311121, China;

[#]Co-first author

23   *Correspondence: songxuemei@zju.edu.cn, ruiliangbai@zju.edu.cn,

24   georg.northoff@theroyal.ca

Abstract

Abstract

The prevailing opinion emphasizes fronto-parietal network (FPN) is key in mediating

general fluid intelligence (gF). Meanwhile, recent studies show that human MT

complex (hMT+), located at the occipito-temporal border and involved in 3D

perception processing, also plays a key role in gF. However, the underlying

mechanism is not clear, yet. To investigate this issue, our study targets visuo-spatial

intelligence, which is considered to have high loading on gF. We use ultra-high field

magnetic resonance spectroscopy (MRS) to measure GABA/glutamate concentrations

in hMT+ combined with resting-state fMRI functional connectivity (FC), behavioral

examinations including hMT+ perception suppression test, and a gF subtest for its

visuo-spatial component. Our findings show that both GABA in hMT+ and frontal-

hMT+ functional connectivity significantly correlate with the performance of visuo-

spatial intelligence. Further, serial mediation model demonstrates that the effect of

hMT+ GABA on visuo-spatial gF is fully mediated by the hMT+ frontal FC.

Together, our findings highlight the importance in integrating sensory and frontal

cortices in mediating the visuospatial component of general fluid intelligence.

Introduction

General fluid intelligence (gF) is a current problem-solving ability, which shows high inter-individual differences in humans(Cattell & Raymond, 1963). At the beginning of the last century, Spearman(Spearman, 1904) proposed that some general or g factor contributes to our gF. One key component of gF is visuo-spatial intelligence that, usually tested by visual materials, shows high g-loading(Colom et al., 2006; Deary et al., 2010; Jung & Haier, 2007). The exact neural mechanisms of the interplay of visuo spatial intelligence with gF remain yet unclear, though.

The "neuro-efficiency" hypothesis is one explanation for individual differences in gF (Haier et al., 1988). This hypothesis puts forward that the human brain's ability to suppress irrelevant information leads to more efficient cognitive processing. Correspondingly, using a well-known visual motion paradigm (center-surround antagonism)(Liu et al., 2016; Tadin et al., 2003), Melnick et al found a strong link between suppression index (SI) of motion perception and the scores of the block design test (BDT, a subtest of the Wechsler Adult Intelligence Scale (WAIS), which measures the visuo-spatial component (3D domain) of gF (Melnick et al., 2013). Motion surround suppression (SI), a specific function of human extrastriate cortical region, middle temporal complex (hMT+), aligns closely with this region's activities (Gautama & Van Hulle, 2001). Furthermore, hMT+ is a sensory cortex involved in visual perception processing (3D domain) (Cumming & DeAngelis, 2001). These findings suggest that hMT+ potentially plays a significant role in 3D visuo-spatial gF by facilitating the efficient processing of 3D visual information and suppressing irrelevant information.

However, more evidence is needed to uncover how the hMT+ functions as a core region

for 3D visuo-spatial intelligence. We here hypothesize a key role of hMT+ GABA and its

functional connectivity with the frontal cortex in mediating visuospatial gF(Figure 1a).

To investigate our hypothesis, this work conducted multi-level examination including

biochemical (Glutamatergic - GABAergic in hMT+), regional-systemic (brain

connectivity with hMT+ - based), and behavioral (visual motion function in hMT+)

levels to reveal if hMT+ contributes to the 3D visuo-spatial component of gF. We

employ ultra-high field (7T) magnetic resonance spectroscopy (MRS) technology to

reliably resolve GABA and Glu concentrations(Ende, 2015; Liu et al., 2022; Song et

al., 2021). To verify the specificity of hMT+, we used primary visual cortex (V1) -

based GABA/Glu as control as it mediates the 2D rather than 3D visual domain

(Cumming & DeAngelis, 2001).

Our findings first demonstrate that GABAergic inhibition mechanisms (but not

excitatory Glu) in hMT+ region relate to 3D visuo-spatial ability. Further, analysis of

functional brain connectivity at rest reveals that the network (between hMT+ and

frontal cortex) relating to hMT+ GABA and perceptual suppression contribute to visuo-

spatial gF. Our results provide direct evidence that GABA-ergic inhibitory mechanisms

in hMT+ region (a sensory cortex) mediate the multi-level visuo-spatial component (3D

domain) of gF thus drawing a direct connection of biochemistry, brain connectivity,

and behavior.

Figure 1. Hypothesis and experimental design. (**a**) Schematic of hypothesis. The

inhibition mechanism centered on hMT+, including the molecular level: the

GABAergic inhibition in hMT+ (green circle), brain connectivity level: hMT+ - frontal

functional connectivity (blue circle), and behavior level: hMT+ specific surround

suppression of visual motion (red circle), contributes to the visuo-spatial component of

gF (3D domain, yellow circle). (**b**) Schematic of experimental design. Session 1

(rectangle box of short line) was the functional MRI and MRS scanning at resting state.

Session 2 (rectangle box of solid line) was another region of MRS acquisition. In the

two sessions, the order of MRS scanning regions (hMT+ and V1) was counter balanced

across participants. There was a structural MRI scanning before each MRS data

acquisition. The interval between the two sessions was used for behavioral

measurement (rectangle box of dotted line): block design task (BDT) and

psychophysical task - motion discrimination. Sold lines indicate the experiment

sequence.

Results

To determine whether the function of hMT+ cortex contributes to visuo-spatial component (3D domain) of gF, we adopted the experimental design depicted in Figure 1b. Thirty-six healthy subjects participated in this study. Participants underwent two MRI sessions: the first encompassing resting-state fMRI and magnetic resonance spectra (MRS), and the second solely involving MRS. A 30-minute interval separated these sessions, during which participants performed motion discrimination tasks (using center-surround antagonism stimuli)(Tadin et al., 2003) and the block design test (BDT), which assesses the visuo-spatial ability (3D domain) of gF (Fangmeier et al., 2006). In the motion discrimination tasks, a grating of either large or small size was randomly presented at the center of the screen. The grating drifted either leftward or rightward, and participants were asked to judge the perceived moving direction. While in the Block design test, participants were asked to rebuild the figural pattern within a specified time limit using a set of red and white blocks. Both the volume-of-interests (VOIs) of MRS scanning in the left hMT+ (targeted brain area) and the left primary visual cortex (V1, control brain area) had dimensions of $2\times2\times2$ cm^3, with the MRS scanning sequences being randomized across the two MRS sessions. The hMT+ MRS VOIs were demarcated using an anatomical landmark (Dumoulin et al., 2000). For 14 subjects, we also utilized fMRI to functionally pinpoint the hMT+ to validate the placement of the VOI (Figure 2a, b). The V1 MRS VOIs were anatomically defined (Methods). Here, MRS data after extensive quality control (31/36 in hMT+, and 28/36 in V1) were taken for further analysis (Methods).

Figure 2. hMT+ localizer scans and hMT+ MRS VOI placement. **(a)** Single task block

designs. First: a cross fixation on the center of the screen (10s). Second: a moving

grating (2°) toward left last 10s. Third: the grating keeps static for 10s. Fourth: the

grating moves toward right last 10s. The localizer scans consist of 8 blocks. **(b)** hMT+

location and MRS VOI placement. The upper template is the horizontal view. The lower

templates from left to right are coronal and sagittal views. The warm color indicates the

overlap of fMRI activation of hMT+ across 14 subjects, the cold color bar indicates the

overlap of MRS VOIs across all subjects.

Figure supplement 1. The left hemisphere V1 MRS scanning ROI. The color bar

indicates the overlap of the MRS VOIs across all subjects.

GABA and Glu concentrations in hMT+ and V1 and their relation to SI and BDT

An example of a MRS voxel located in hMT+ is shown in Figure 3a. LCModel fittings for GABA spectra from all subjects in hMT+ ($n = 31$) and V1 ($n = 28$) are illustrated in Figure 3b (color scale presents the BDT scores). We discerned a significant association between the inter-subjects' BDT scores and the GABA levels in hMT+ voxels, but not in V1 voxels. Quantitative analysis displayed that BDT significantly correlates with GABA concentrations in hMT+ voxels ($r = 0.39$, $P = 0.03$, $n = 31$, Figure 3c). After using partial correlation to control for the effect of age, the relationship remains significant ($r_{partial} = 0.43$ $P = 0.02$, 1 participant excluded due to the age greater than mean + 2.5SD). In contrast, there was no correlation between BDT and GABA levels in V1 voxels (figure supplement 1a). Further, we show that SI significantly correlates with GABA levels in hMT+ voxels ($r = 0.44$, $P = 0.01$, $n = 31$, Figure 3d). In contrast, no significant correlation between SI and GABA concentrations in V1 voxels was observed (figure supplement 1b). This finding is in line with prior results, which indicates that motion perception is associated with neural activity in hMT+ area, but not in V1 (Schallmo et al., 2018). LCModel fittings for Glu spectra from all subjects in hMT+ ($n = 31$) and V1 ($n = 28$) voxels are presented in figure supplement 2a.

Unlike in the case of GABA, no significant correlations between BDT and Glu levels were found in both hMT+ and V1 voxels (figure supplement 2b, c). While, as expected (Song et al., 2021), we observed significant positive correlations between GABA and Glu concentrations in both hMT+ ($r = 0.62$, $P < 0.001$, $n = 31$) and V1 voxels ($r = 0.56$, $P = 0.002$, $n = 28$) (figure supplement 3a, b). Additionally, significant correlations

between SI and BDT, small threshold and SI were discerned ($r = 0.59$, $P = 0.0002$, $n =$

34, figure 3e, $r_{\text{partial}} = 0.67$, $P < 0.001$, 1 participant excluded due to the age greater than

mean $+$ 2.5SD; $r = -0.43$, $P = 0.016$, $r_{partial} = 0.44$, $P = 0.014$). While there was no

significant correlation between large threshold and SI, corroborating previous

conclusions (Melnick et al., 2013). Two outliers evident in Figure 3e were excluded,

with consistent results depicted in figure supplement 4a. Further, two outliers evident

in Figure 3d were excluded, with consistent results depicted in figure supplement 4b.

Figure 3. MRS spectra and the relationships between GABA levels and SI / BDT. **(a)** Example spectrum from the hMT+ voxel of one participant. The first line is the LCModel fitting result of all metabolites, and the following lines show the Glu and GABA spectra fitting with LCModel, and then the baseline. **(b)** Individual participants fitted GABA MRS spectra from the hMT+ (top) and V1 (bottom) voxels from baseline measurement. The colors of the GABA spectra represent the individual differences of BDT. The color bar represents the scores of BDT. **(c) and (d)** Pearson's correlations showing significant positive correlations between hMT+ GABA and BDT scores **(c)**, between hMT+ GABA and SI **(d)**. GABA and Glu concentrations (Conc.) are absolute, with units of mmol per kg wet weight (Methods). **(e)** Peason's correlation showing significant positive correlations between SI and BDT. **(f)** Peason's correlation showing significant negative correlations between BDT and small threshold. **(g)** No correlation

between BDT and large threshold. The ribbon between dotted lines represents the 95%

confidence interval, and the black regression line represents the Pearson's correlation

coefficient (r).

Figure supplement 2. Relationships between GABA concentration in V1 and

BDT/SI. **(a)** There is no significant relationship between BDT and GABA

concentrations in V1. **(b)** There is also no correlation between SI and GABA

concentrations in V1. GABA concentration (Conc.) is absolute, with units of mmol

per kg wet weight (Methods).

Figure supplement 3. Individual Glu MRS spectra from hMT+ / V1 regions and

relationships between BDT and Glu concentrations in hMT+ / V1 regions. **(a)**

Individual participant fitted Glutamate MRS spectra from the hMT+ (top, *n* = 31) and

V1 (bottom, *n* = 28) voxels from baseline measurement. The colors of the Glutamate

spectra represent the individual differences of BDT. The color bar represents the scores

of BDT. **(b)** There is no significant relationship between BDT and Glu concentrations

in hMT+. **(c)** There is also no significant correlation between BDT and Glu

concentrations in V1. Glutamate concentration (Conc.) is absolute, with units of mmol

per kg wet weight (Methods).

Figure supplement 4. Correlations between GABA and Glu concentrations in hMT+

and V1 regions. **(a)** There is significant correlation between GABA and Glu

concentrations in hMT+ region. **(b)** The levels of GABA also significantly correlate

with Glu in V1 region. GABA and Glu concentrations (Conc.) are absolute, with units

of mmol per kg wet weight (Methods).

Figure supplement 5. Two linear correlations. **(a)** Significant positive correlation

between suppression index and BDT scores (took out two outlines, , having the similar

result shown in Figure 2e). **(b)** BDT also significantly correlates with GABA

concentration in hMT+ region (without two outliers, having the similar result shown in

Figure 2d).

MT - frontal FC relates to SI and BDT

We next took the left hMT+ as the seed region and separately measured interregional

FCs between the seed region and each voxel in the frontal regions (a priori search space).

These measurements were correlated with the performance in 3D visuospatial ability

(BDT) to identify FCs with significant correlations. Results from connectivity-BDT

analysis are summarized in Table 1 and shown in Figure 4a. We found that brain regions

with high FC strength to the seed region (left hMT+) significantly correlated with BDT

scores, these included mainly regions situated within the canonical cognitive cores of

FPN (Brodmann areas (BAs) 6, 9, 10, 46, 47)(Assem et al., 2020; Deary et al., 2010;

Duncan et al., 2020; Duncan et al., 2000; Gray et al., 2003; Jung & Haier, 2007). Across

the whole brain search, the similar FCs (between hMT+ and frontal cognitive cores)

still showed significant correlations with BDT scores (Table supplement 1) (also shown

in figure supplement 5a). Additionally, we identified certain parietal regions (BAs 7,

39, 40) with significant correlations of their functional connectivity to the left hMT+

and the BDT scores (Table supplement 1) (also shown in figure supplement 5a). These

significant connections between hMT+ and FPN system suggest that left hMT+ is

involved in the efficient information integration network mediating the visuo spatial

component of gF.

To address the question whether spatial suppression plays a role, we correlated

hMT+ - based global FCs with SI. Though spatial suppression during motion perception

(quantified by SI) is considered to be the function of area hMT (Gautama & Van Hulle,

2001), the top-down modulation from the frontal cortex can increase surround

suppression (Liu et al., 2016). Our functional connectivity-SI analysis in the frontal

regions (a priori search space) displayed 3 brain regions in which FCs strength

significantly correlated with SI: right BA4/6, left BA6, and right BA46 (summarized in

Table supplement 2, and shown in Figure 4b). Across the whole brain search, we

identified total 7 brain regions in which FCs strength significantly correlated with SI,

and 3 of these were in the frontal cortex. This is consistent with the results obtained by

the functional connectivity-SI analysis in a priori search space (frontal cortex) (Table

supplement 3 and figure supplement 5b).

Table 1 FC of voxels showing significant correlation with BDT scores across subjects in
frontal cortex.

FC number	Connected regions	BA	Size	Peak coordinate MNI (x, y, z)	r	P
1	Frontal_Sup_Orb_R	11	33	(12,63, -19.5)	-0.57	0.0011
2	Frontal_Inf_Orb_L	47	24	(-34.5,28.5, -13.5)	-0.63	0.0003
3	Frontal_Med_Orb_R	11	41	(3,43.5, -12)	-0.58	0.0009
4	Frontal_Inf_Orb_R	47	48	(-31.5,24, -12)	0.59	0.0008
5	Frontal_Inf_Orb_R	47	29	(25.5,30, -13.5)	0.67	0.0001
6	Insula_L	\	26	(-28.5,27,0)	0.67	0.0001
7	Frontal_Inf_Oper_R	45	41	(43.5,16.5,6)	0.64	0.0002
8	Frontal_Sup_R	10	25	(31.5,57,9)	0.59	0.0008
9	Frontal_Mid_L	10	82	(-33,48,12)	0.62	0.0003
10	Frontal_Inf_Oper_R	44	49	(51,7.5,21)	0.59	0.0007
11	Frontal_Inf_Oper_R	46	96	(49.5,16.5,28.5)	-0.62	0.0003
12	Frontal_Mid_L	10	32	(-31.5,49.5,24)	0.57	0.0012
13	Frontal_Mid_R	10	102	(31.5,36,30)	0.59	0.0009
14	Precentral_L	6	46	(-49.5, -1.5,34.5)	0.59	0.0007
15	Frontal_Mid_R	9	107	(51,19.5,40.5)	-0.67	0.0001
16	Frontal_Sup_L	9	35	(-9,60,37.5)	-0.69	0.0001
17	Frontal_Sup_Medial_R	9	74	(4.5,52.5,43.5)	-0.57	0.0011
18	Frontal_Sup_R	6	136	(28.5, -7.5,63)	-0.64	0.0002
19	Supp_Motor_Area_L	6	48	(-10.5,6,54)	0.63	0.0003
20	Frontal_Mid_L	6	119	(-24,4.5,55.5)	0.63	0.0003
21	Precentral_L	6	229	(-24, -18,66)	0.60	0.0005
22	Frontal_Sup_R	6	32	(16.5, -18,67.5)	0.68	0.0001
23	Precentral_R	6	80	(30, -24,70.5)	0.70	0.0001
24	Precentral_R	6	23	(16.5, -25.5,76.5)	0.70	0.0001

256 Single voxel threshold $P < 0.005$ (t > 3.057 or t < -3.057), adjacent size $\geq$ 23 voxels (AlphaSim

257 corrected).

259 **Table supplementary 1. FC of voxels showing significant correlation with BDT scores across**
260 **subjects in whole brain.**

FC number	Connected regions	BA	Size	Peak coordinate MNI (x, y, z)	r	P
1	Frontal_Med_OrbR	11	44	(2,43.5, -12)	-0.58	0.0009
2	Frontal_Inf_Oper_R	45	49	(43.5,16.5,6)	0.64	0.0002
3	Precentral_L	6	46	(-49.5, -1.5,34.5)	0.59	0.0007
4	Precentral_L	6	237	(-24, -18,66)	0.68	0.0001
5	Precentral_R	6	80	(31, -25,72)	0.67	0.0001
6	Frontal_Mid_L	10	82	(-33,48,12)	0.62	0.0003
7	Insula_L	47	124	(-33,15, -9)	0.64	0.0002
8	Insula_L	13	107	(-31.5,9,10.5)	0.63	0.0002
9	Insula_L	13	44	(-42, -10.5,7.5)	0.64	0.0002
10	Frontal_Inf_Oper_R	44	49	(51,7.5,21)	0.59	0.0007
11	Frontal_Inf_Oper_R	46	96	(49.5,16.5,28.5)	-0.62	0.0003
12	Frontal_Mid_R	10	102	(31.5,36,30)	0.59	0.0009
13	Paracentral_Lobule_L	6	46	(-15,21,51)	-0.64	0.0002
14	Supp_Motor_Area_L	6	48	(-10.5,6,54)	0.63	0.0003
15	Frontal_Mid_L	6	119	(-24,4.5,55.5)	-0.67	0.0001
16	Frontal_Sup_R	6	136	(29, -9,65)	0.57	0.0014
17	Frontal_Sup_MedialR	9	90	(8,51,43)	-0.59	0.0009
18	Frontal_Mid_R	9	108	(50,19,41)	-0.67	0.0001
19	Occipital_Mid_L	19	47	(-43, -83,9)	-0.64	0.0002
20	Occipital_Mid_L	37	45	(-40.5, -63,4.5)	-0.62	0.0004
21	Temporal_Mid_R	39	56	(45, -57,4.5)	-0.71	0.0000
22	Temporal_Mid_L	39	54	(-45, -48,12)	-0.55	0.0021
23	Temporal_Mid_L	40	102	(-48, -55.5,16.5)	-0.65	0.0001
24	Temporal_Mid_R	21	105	(64, -2, -19)	-0.61	0.0004
25	Temporal_Sup_R	22	228	(65, -42,11)	-0.6	0.0006
26	Precuneus_L	30	155	(1.5, -51,16.5)	-0.64	0.0002
27	Cingulum_Ant_R	32	41	(9,15,27)	0.68	0.0000
28	Cingulum_Mid_R	31	73	(15, -46.5,36)	-0.63	0.0002
29	Cingulum_Mid_L	23	80	(-3, -15,30)	0.63	0.0003
30	Lingual_R	18	46	(21, -93, -16)	0.58	0.0009
31	Parietal_Sup_L	7	54	(-19.5, -63,55.5)	0.66	0.0001
32	Parietal_Sup_L	7	48	(-23, -71,58)	0.63	0.0003
33	Postcentral_L	40	41	(-31.5, -39,58.5)	0.68	0.0001
34	Postcentral_L	40	40	(-48, -36,58)	0.71	0.0000
35	Thalamus_L	Wm	108	(-24, -31.5,12)	0.64	0.0002
36	Parietal_Sup_L	5	37	(-37, -48,63)	0.59	0.0008
37	Vermis_10	-	37	(-3, -43, -37)	-0.68	0.0001
38	-	-	40	(12, -19.5, -42)	-0.68	0.0000

261 Single voxel threshold $P < 0.01$ (t > 2.771 or t < -2.771), adjacent size ≥ 37 voxels (AlphaSim corrected).

263 **Table supplement 2. FCs of voxels showing significant correlation with SI across subjects in**
264 **frontal cortex.**

FC number	Connected regions	BA	Size	Peak coordinate MNI (x, y, z)	r	P
1	Frontal_Inf_Oper_R	46	80	(48,15,28.5)	-0.65	0.0001
2	Precentral_R	4/6	106	(33, -25.5,63)	0.72	0.0000
3	Precentral_L	6	26	(-30, -24,72)	0.66	0.0001

265 Single voxel threshold $P < 0.005$ (t > 3.057 or t < -3.057), adjacent size $\geq$ 22 voxels (AlphaSim
266 corrected).

269 **Table supplement 3. FCs of voxels showing significant correlation with SI across subjects in**
270 **whole Brain.**

FC number	Connected regions	BA	Size	Peak coordinate	r	P
				MNI (x, y, z)		
1	Cerebelum_Crus1_L	18	70	(-29, -85, -24)	0.51	0.0052
2	Cerebelum_6_R	18	39	(12, -85, -17)	0.65	0.0002
3	Calcarine_R	18	70	(15, -91.5,12)	0.63	0.0003
4	Frontal_Inf_Oper_R	46	127	(48,15,28.5)	-0.65	0.0001
5	Precentral_R	4/6	179	(32, -24,70)	0.71	0.0001
6	Precentral_L	6	59	(-30, -23,72)	0.66	0.0001
7	-	-	37	(26, -37, -11)	-0.67	0.0001

Single voxel threshold $P < 0.01$ (t > 2.771 or t < -2.771), adjacent size ≥ 37 voxels (AlphaSim
corrected).

Figure 4. Significant FCs from connectivity-behavior analyses in *a priori* search space.

The seed region is the left hMT+. The significant FCs are obtained from *a priori* space

(frontal cortex). (**a**) The significant FCs obtained from connectivity-BDT analysis.

Single voxel threshold *P* < 0.005, adjacent size ≥ 23 (*AlphaSim* correcting, Methods).

(**b**) The significant FCs obtained from connectivity-SI analysis. Single voxel threshold

P < 0.005, adjacent size ≥ 22 (*AlphaSim* correcting, Methods). Positive correlations are

shown in warm colors, while, negative correlations are shown in cold colors. The paired

deep green, deep pink, light green circles on (**a**) and (**b**) indicate the overlap regions in

left BA6, right BA6, right BA46(DLPFC) between Connectivity - BDT analysis and

Connectivity – SI analysis, respectively.

Figure supplement 6. Significant FCs searched from connectivity-behavior

(BDT/SI) analyses in the whole brain. The seed region is the left hMT+. The

significant FCs are obtained from the entire brain search, single voxel threshold *P* <

0.01, adjacent size ≥ 37 voxels (AlphaSim correcting, Methods). Positive correlations

are shown in warm colors, while, negative correlations are shown in cold colors. **(a)**

the significant FCs obtained from connectivity-BDT analysis. **(b)** the significant FCs

obtained from connectivity-SI analysis.

Figure supplement 7. Significant FCs searched from connectivity-behavior (BDT)

analyses in the frontal **(a)** and whole brain **(b)**. The seed region is the left V1. **(a)** The

significant FCs are obtained from the frontal search, single voxel threshold $P < 0.005$,

adjacent size ≥ 23 (AlphaSim correcting, same methods for hMT+). **(b)** The

significant FCs are obtained from the entire brain search, single voxel threshold $P <$

0.01, adjacent size ≥ 37 voxels (AlphaSim correcting, same methods for hMT+). Only

positive correlations were detected, shown in warm colors.

304
305

306 **Table 2. Correlations between FC in Table 1 and GABA/Glu concentrations in hMT+.**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	-0.07	0.72	0.75	-0.11	0.58	0.85
2	-0.28	0.14	0.36	-0.27	0.15	0.81
3	-0.13	0.52	0.59	-0.07	0.71	0.85
4	0.10	0.59	0.64	0.12	0.54	0.85
5	0.14	0.48	0.58	0.24	0.21	0.81
6	0.31	0.11	0.33	0.15	0.43	0.85
7	0.20	0.30	0.48	0.07	0.74	0.85
8	0.14	0.45	0.58	0.05	0.79	0.85
9	0.20	0.30	0.48	0.10	0.60	0.85
10	-0.13	0.49	0.58	-0.16	0.41	0.85
11	**-0.56**	**0.0018**	**0.02***	-0.22	0.25	0.81
12	0.18	0.34	0.51	0.15	0.43	0.85
13	0.20	0.30	0.48	0.05	0.81	0.85
14	0.39	0.04	0.14	0.22	0.24	0.81
15	-0.40	0.03	0.12	-0.21	0.27	0.81
16	-0.27	0.15	0.36	-0.12	0.53	0.85
17	0.17	0.37	0.52	0.06	0.74	0.85
18	0.26	0.18	0.39	0.16	0.40	0.85
19	0.39	0.03	0.12	0.31	0.10	0.81
20	0.01	0.98	0.98	0.14	0.46	0.85
21	0.40	0.03	0.12	0.24	0.21	0.81
22	0.22	0.25	0.48	0.06	0.76	0.85
23	**0.69**	**0.0001**	**0.002****	0.47	0.01	0.24
24	0.41	0.03	0.12	0.001	0.97	0.97

307 Bold font indicates the significant correlations survived from multi correlation correction.

309 **Table supplement 4. Correlations between FC in Table supplement 1 and GABA/Glu**

310 **concentrations in hMT+**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	-0.13	0.49	0.55	-0.08	0.68	0.84
2	0.17	0.38	0.49	0.05	0.82	0.88
3	0.39	0.04	0.22	0.22	0.24	0.69
4	0.41	0.03	0.19	0.24	0.21	0.69
5	**0.69**	**0.0001**	**0.0038**[**]	0.47	0.01	0.38
6	0.2	0.3	0.41	0.1	0.6	0.81
7	0.26	0.17	0.34	0.21	0.28	0.69
8	0.22	0.24	0.35	0.25	0.19	0.69
9	0.23	0.23	0.35	0.02	0.9	0.92
10	-0.13	0.49	0.55	-0.16	0.41	0.70
11	**-0.56**	**0.0018**	**0.03**[*]	-0.22	0.25	0.69
12	0.2	0.3	0.41	0.05	0.81	0.88
13	-0.35	0.06	0.25	0.07	0.73	0.84
14	0.39	0.03	0.19	0.31	0.1	0.69
15	0.006	0.98	0.98	0.14	0.46	0.70
16	0.26	0.18	0.34	0.16	0.4	0.70
17	0.15	0.42	0.51	0.04	0.83	0.88
18	-0.4	0.03	0.19	-0.21	0.27	0.69
19	-0.37	0.05	0.24	-0.15	0.44	0.70
20	-0.16	0.39	0.49	-0.19	0.33	0.70
21	-0.31	0.1	0.28	-0.29	0.13	0.69
22	-0.22	0.24	0.35	-0.07	0.72	0.84
23	-0.1	0.6	0.65	-0.12	0.53	0.77
24	-0.09	0.64	0.68	-0.09	0.63	0.83
25	-0.3	0.11	0.28	-0.14	0.46	0.70
26	-0.08	0.66	0.68	-0.08	0.69	0.84
27	0.28	0.13	0.29	0.1	0.59	0.81
28	-0.27	0.16	0.34	-0.2	0.29	0.69
29	0.33	0.08	0.28	0.25	0.19	0.69
30	0.24	0.21	0.35	0.19	0.32	0.70
31	0.24	0.22	0.35	0.36	0.05	0.69
32	0.23	0.24	0.35	0.23	0.23	0.69
33	0.44	0.02	0.19	0.18	0.35	0.70
34	0.32	0.09	0.28	0.002	0.99	0.99
35	0.32	0.09	0.28	0.31	0.1	0.69
36	0.30	0.11	0.28	0.22	0.25	0.69
37	-0.29	0.13	0.29	-0.23	0.23	0.69
38	-0.13	0.48	0.55	-0,15	0.42	0.70

311 Bold font indicates the significant correlations survived from multi correlation correction.

312 25

313 **Table supplement 5. Correlations between FCs in Table supplement 2 and GABA/Glu**

314 **concentrations in hMT+.**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	**-0.56**	**0.0017**	**0.0017****	-0.21	0.27	0.27
2	**0.70**	**0.0001**	**0.0003*****	**0.49**	**0.007**	**0.021***
3	**0.65**	**0.0001**	**0.0002****	**0.40**	**0.03**	**0.045***

315 **: P_{FDR} < 0.05; **: P_{FDR} < 0.01; ***: P_{FDR} < 0.001; Bold font indicates the significant correlations*

316 survived from multi correlation correction.

318 **Table supplement 6. Correlations between FCs in Table supplement 3 and GABA/Glu**

319 **concentrations in hMT+.**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	**0.37**	**0.049**	**0.049***	0.32	0.09	>0.05
2	**0.37**	**0.049**	**0.049***	0.41	0.025	>0.05
3	**0.48**	**0.008**	**0.01***	0.21	0.28	>0.05
4	**-0.58**	**0.001**	**0.002****	-0.22	0.26	>0.05
5	**0.69**	**0.0001**	**0.0007**	0.46	0.01	>0.05
6	**0.66**	**0.0001**	**0.0004*****	0,39	0.04	>0.05
7	**-0.48**	**0.0077**	**0.01***	-0.35	0.067	>0.05

320 **: P_{FDR} < 0.05; **: P_{FDR} < 0.01; ***: P_{FDR} < 0.001; Bold font indicates the significant correlations*

321 survived from multi correlation correction.

Local hMT+ GABA acts on SI and BDT via global MT-frontal connectivity

To determine whether local neurotransmitter levels (such as GABA and Glu) in the hMT+ region mediate the broader 3D visuo-spatial ability of BDT, which as a component of gF, is linked to the frontal cortex (Fangmeier et al., 2006), we correlated the significant FCs of hMT+ – frontal in Figure 4a (also shown in Table 1) with the GABA and Glu levels in hMT+ region. The results revealed that only two FCs significantly correlated with inhibitory GABA levels in hMT+: 1) the FC of left hMT+ - right BA 46 (significantly negative correlation, $r = - 0.56$, $P = 0.02$, $n = 29$, FDR correction, Figure 5a left); 2) the FC of left hMT+ - right BA 6 (significantly positive correlation, $r = 0.69$, $P = 0.002$, $n = 29$, FDR correction, Figure 5b left) (also shown in Table 2). There were no significant correlations between these FCs and the excitatory Glu levels in hMT+ (Table 2). Across the whole brain search, we obtained the same two hMT+ - frontal FCs significantly correlating with both hMT+ GABA levels and BDT (Table supplement 4), this is consistent with the results in the priori search space (frontal cortex) (Table 2). We then correlated the significant FCs in Figure 5b (also in Table supplement 2) with GABA and Glu concentrations in hMT+ and found that almost all the correlations are significant except one (between the FC of left hMT+ - right BA46 and the Glu levels in hMT+) (Table supplement 5). Among the three FCs, the clusters of two FCs have substantial voxel overlap with the FCs we found by the connectivity-BDT analysis (Figure 5a, b). Across the whole brain search, there were total 7 brain regions in which FCs strength were significantly correlated with SI, all the 7 FCs significantly correlated the hMT+ GABA levels, while, no FC had significant

correlation with the hMT+ Glu levels (Table supplement 6).

Taken together, our results displayed that the overlap FCs from the analyses of

connectivity - behavior (BDT and SI) -GABA are the hMT+ - BA 46 and hMT+ - BA

6 (Figure 5a, b). These results suggest that the coupling of FCs of hMT+ - frontal

regions (BA 46 and BA 6) with local hMT+ GABA provides the neural basis for both

the simple motion perception (quantified by SI) and the complex 3D visuo-spatial

ability (quantified by BDT).

In order to fully investigate the potential roles of the multiples variables contributing to

BDT scores, serial mediation analyses (Hayes, 2013) were applied to both MR and

behavioral data. Following our hypothesis, the independent variable (X) is hMT+

GABA, the dependent variable (Y) is BDT scores, the covariate is the age, and the

mediators are FC (M1) and SI (M2). We used the overlap clusters from the analyses of

connectivity-BDT-GABA and connectivity-SI-GABA to compute the FC of hMT+ -

BA46, 1 participant was excluded due to his age greater than mean + 2.5SD. The serial

mediation model is shown in Figure 5c. GABA levels in hMT+ significantly negatively

correlated with the FC of hMT+ - BA46 (β = - 0.32, P < 0.001), which in turn

significantly negatively correlated with SI (β = - 0.19, P = 0.035), and consequently,

significantly positively correlated with BDT (β = 38.5, P = 0.009). Critically,

bootstrapped analyses revealed that our hypothesized indirect effect (i.e., hMT+ GABA

$\rightarrow$ FC of hMT+ - BA46 $\rightarrow$ SI $\rightarrow$ BDT) was significant (β = 2.28, SE = 1.54, 95% CI =

[0.03, 5.94]). The model accounted for 34% of the variance in BDT. However, when

considering the MT-BA6 FC as the mediator M1, the serial model does not show a
28

significant indirect effect. Consequently, we explored a mediation model, which

revealed that the MT-BA6 FC totally mediates the relationship between GABA and

BDS. (Figure 5d). For sensitivity purposes, we tested the alternative models, in which

the order of the mediators was reversed. The pathway that hMT+ GABA was predicted

to be associated with SI, followed by the FC of hMT+ - BA 46, and then BDT, did not

yield the chained mediation effects on BDT (figure supplement 6).

Figure 5. Local hMT+ GABA acts on SI and BDT via global MT-frontal connectivity.

(**a**) Significant negative correlation between the FC of left hMT+ - right DLPFC (BA46)

and hMT+ GABA (*FDR* correction). (**b**) Significant positive correlation between the

FC of left hMT+ - right (pre) motor cortex (BA4/6) and hMT+ GABA (*FDR* correction).

In (**a**) and (**b**), left: the significant FCs obtained from connectivity-BDT analysis; right:

the significant FCs obtained from connectivity-SI analysis. (**c**) Significant pathways:

hMT+ GABA→ FC (left hMT+ - right BA46, negative correlation) → SI (negative

correlation) → BDT (positive correlation). This pathway can explain 34% of the

variance in BDT. **(d)** Significant pathways: hMT+ GABA → FC (left hMT+ - right

BA6, positive correlation) → BDT (positive correlation). The bolded lines represent

the hypothesized mediation effect. The dotted lines represent alternative pathways. *: P

< 0.05; **: P < 0.01; ***: P < 0.001.

Figure supplement 8. The alternative serial mediation models from local hMT+

GABA to global performance of BDT. The pathway that hMT+ GABA was predicted

to be associated with SI, followed by the FC of hMT+ - BA 46, and then BDT, did not

yield the chained mediation effects on BDT. *: P < 0.05; **: P < 0.01; ***: P < 0.001.

Figure Supplement 9. Correlation matrix shows correlations between all measured in

our biochemistry and behaviour level. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$.

To summarize (shown in Figure 6), the results from the serial mediation analyses are

consistent with our hypothesis. That is, higher GABAergic inhibition in hMT+ relates

to stronger negative FC between hMT+ and BA46, leading to enhanced ability for

surround suppression (filtering out irrelevant information(Tadin, 2015), which

ultimately results in more efficient visual 3D processing as key component of gF (the

higher BDT scores).

Figure 6. Sketch depicting the multi-level inhibitory mechanisms centered on hMT+

GABA contributing to visuo-spatial intelligence. Inhibitory GABA in hMT+ (a sensory

cortex, shown in green circle), coupling with the functional connectivity between

hMT+ and BA46 (cognitive control core, shown in purple circle), and mediated by

motion surround suppression (shown in red circle), contributes to visuo-spatial

intelligence (BDT, 3D domain, shown in red and white building blocks). In this sketch,

the two-colored parallel arrows show the negative FC between hMT+ and BA46, the

colored arrows below the green circle display the inhibition mechanisms centered on

hMT+ GABA (2), filtered the irrelevant information in (1) and focused on the efficient

visuo-spatial information (3). Black long arrows display the direction of information

flow: from input information (1) to visuo-spatial intelligence (4).

Discussion

Discussion

Here, we provide evidence that hMT+ inhibitory mechanisms mediate processing in the

visuo-spatial component (3D domain) of gF on multiple levels, that is, from molecular

over brain connectivity to behavior. First, this study found that higher hMT+ inhibitory

GABA levels (but not excitatory Glu) relate to FC between hMT+ and BA 46 that

contribute to both SI and 3D visuo-spatial intelligence (BDT). Our serial mediation

analyses indicate that the inhibitory mechanisms related to hMT+ and its GABA levels

(but not Glu), FCs of hMT+ - BA46 coupling with hMT+ inhibitory GABA (but not

excitatory Glu), and behavior (SI indexing perceptual suppression in hMT+) predict the

inter-subject variance in the 3D visuo-spatial intelligence (BDT) (Figure 5c). Second,

we demonstrate discrete GABAergic inhibition mechanisms in hMT+ that mediate the

strong FCs between hMT+ - frontal regions (BA46 and BA6): significant negative

correlation with the FC of hMT+ - BA 46 (Figure 5a), whereas there is significant

positive correlation with the FC of hMT+ - BA 6 (Figure 5b). This indicates that

different frontal regions, DLPFC (BA 46) and premotor cortex (BA6), contribute

uniquely to gF through hMT+ - based inhibitory mechanisms.

The goal of our research is to reveal that the inhibitory (not excitatory) mechanism

in hMT+ contributes to multi-level processing in 3D visuo-spatial ability (BDT).

Monkey electrophysiological experiments revealed that selective attention gates the

visual cortex, including area MT, effectively suppressing the irrelevant

information(Everling et al., 2002; Treue & Maunsell, 1996). These findings align with

the "neural efficiency" hypothesis of intelligence (Haier et al., 1988), which puts

forward the human brain's ability to suppress the repetition of information. Neural

suppression is associated with the balance between excitation and inhibition (EIB),

usually represented by covariation between Glutamate and GABA(Ozeki et al., 2009).

Here, this study exploited the high spectral resolution afforded by ultrahigh field (7T)

MRS to reliably resolve GABA measurement, to adequately discriminate the glutamate

and glutamine signals, and to resolve the high accuracy Glu measurement(Ende, 2015).

This work implemented the MRS scanning in hMT+ (3D visual domain) and V1 (2D

visual domain) regions and found that hMT+ inhibitory GABA (but not excitatory Glu)

significantly correlated with 3D visuo-spatial intelligence, i.e., the higher GABA levels

in hMT+ (rather than excitatory Glu) relate to higher visual 3D processing (Figure 3c).

We searched the global hMT+ - based FCs with the connectivity-BDT analyses (in

priori search space and whole brain search to valid), and then, correlated these

significant FCs with the GABA and Glu concentrations in hMT+ . We found two FCs

(MT+ - BA46 and hMT+ - BA6) significantly correlating with hMT+ inhibitory GABA

(whereas no FC significantly correlated with hMT+ excitatory Glu). Accordingly, our

results emphasize the importance of hMT+ inhibitory GABA (but not excitatory Glu)

in processing the 3D visual-spatial intelligence (BDT).

Our recent human study(Song et al., 2021) and other study's animal

experiments(Ozeki et al., 2009; Sato et al., 2016) demonstrated that the conjoint action

of inhibition (GABA) and excitation (Glu) underlies visual spatial suppression. In this

work, our novel data show the chained mediation effects from local hMT+ GABA to

more global 3D visuo-spatial intelligence: hMT+ GABA → FC (MT+ and BA46) →

SI → BDT. Thereby, our data indicate that inhibitory mechanisms in hMT+, from the

biochemical level of GABA over FC to the behavioral level, can predict the inter-

subject variance in the 3D visuo-spatial intelligence (BDT) (Figure 5c).

Another interesting finding reveal that GABAergic inhibition in hMT+ coupling with

distinct functional connectivity patterns between BA 46-MT+ and BA6-MT+. A

previous human fMRI experiment found that the positive and negative correlations

between 3D visuo-spatial intelligence and the activation of frontal regions appeared at

different reasoning phases (validation or integration phases during

reasoning)(Fangmeier et al., 2006). On the one hand, a monkey electrophysiological

experiment reported the delayed modulation from PFC (especially in DLPFC (BA 46)))

to area MT during a visual motion task(Zaksas & Pasternak, 2006). Computational

models converged with empirical data of awake monkey experiments slowing temporal

modulation from PFC to MT/MST(Donner et al., 2009; Siegel et al., 2015; Wang, 2002;

Wimmer et al., 2015). On the other hand, human MEG studies(Donner et al., 2009;

Wilming et al., 2020) reported that gamma-band activity in visual cortex (including

area MT) exhibits high coherence with the activity in (pre-) motor regions (BA 4/6).

These results suggest that the relation of long-range FC and local inhibitory mechanism

(MT+ GABA) support our findings that inhibitory GABA in hMT+ contributes to

efficient long-range integration and coordination in distant brain areas like the

prefrontal and premotor cortex.

How does hMT+ functions as a core region for 3D visuo-spatial intelligence. rather

than as simple input module? The results in Figure 5a, c showed that the overlap brain

regions from the analyses of connectivity-BDT-GABA/connectivity-SI-GABA are the

hMT+ - BA 46. This overlap couples with local visual suppression (SI) and

consequently plays an important role in 3D visuo-spatial intelligence. The direction

discrimination task in this work (the visual motion paradigm of center-surround

antagonism) was previously considered a mainly local function of hMT+(Melnick et

al., 2013; Tadin, 2015; Tadin et al., 2003). However, our results with connectivity-SI

analyses revealed that both local (FC within BA 18) and global brain connectivity (FC

between hMT+ and frontal regions) contribute to SI (Table supplement 3). In human

psychophysical experiments(Melnick et al., 2013; Tadin et al., 2003) the brief stimulus

duration (~100 ms) in motion discrimination precludes most top-down attentional

feedback effects(Wang, 2002; Zaksas & Pasternak, 2006), while, attention, which

predicted the performance of the motion discrimination task, was sustained throughout

the stimulus intervals(Siegel et al., 2015). Furthermore, animal experiments have

revealed that the local circuits in the visual cortex combining with top-down modulation

and intracortical horizontal connection mediate the visual-spatial suppression

(Angelucci et al., 2002; Keller et al., 2020; Li et al., 2019; Zhang et al., 2014).

Our results (shown in Figure 5a, b, right) present the intrinsic binding of local

GABAergic inhibition in hMT+, which suppressing redundancy of visual motion

processing (SI), and the activity of brain connectivity between hMT+ and frontal

regions. These individually inherent traits may contribute to the individual difference

in 3D visuo-spatial ability (Figure 5a, b, left). A candidate divisive normalization

model(Carandini & Heeger, 2011; Reynolds & Heeger, 2009) can explain how such

reverberation affects the process of suppressing the irrelevant information, from

perception to intelligence(Melnick et al., 2013; Tadin, 2015). We summarize a

framework (Figure 6) to indicate and visualize our findings.

Recently, Duncan et al. demonstrated coding of general fluid intelligence (gF) in

distributed regions, defining them as part of multi-demand (MD) systems (Assem et al.,

2020; Duncan et al., 2020). The MD system encompasses a range of cognitive domains,

including working memory, mathematics, language, and relational reasoning.

According to Melnick et al. (2013), motion surround suppression (SI) and time

thresholds for small and large gratings, which reflect hMT+ functionality, are correlated

with Verbal Comprehension, Perceptual Reasoning, Working Memory, and Processing

Speed indicators. Additionally, Fedorenko et al. identified multi-demand activation

regions around the occipito-temporal areas, potentially overlapping with hMT+

(Fedorenko et al., 2013). As a key region in the representation of sensory flows

(including optic, tactile, and auditory flows)(Fetsch et al., 2011; Gu et al., 2006), hMT+

shows potential to be central to the MD system. Future research could focus on multi-

task paradigms to further investigate the mechanisms of hMT+ and its relationship with

broader cognitive functions.

Together, this study offers a comprehensive insight into how the information

exchange and integration between the sensory cortex (hMT+) and cognition core of BA

46, coupling with the hMT+ GABA, can predict the performance of 3D visuo-spatial

ability (BDT). Our results provide direct evidence that a sensory cortex area (hMT+),

its GABA biochemistry, functional connectivity, and cognition behavior levels, can

assemble into complex cognition as an 3D visuo-spatial intellectual hub.

Materials and Methods

Subjects

Thirty-six healthy subjects (18 female, mean age: 23.6 years $\pm$ 2.1, range: 20 to 29 years)

participated in this study, they were recruited from Zhejiang University. All subjects

had normal or corrected-to-normal vision. In addition, they reported no psychotropic

medication use, no illicit drug use within the past month, no alcohol use within 3 days

prior to scanning, and right-handed. This experiment was approved by the Ethics

Review Committee of Zhejiang University and conducted in accordance with the

Helsinki Declaration. All participants signed informed consent forms prior to the start

of the study and were compensated for their time. All subjects participated in the motion

spatial suppression psychophysical, resting-state fMRI and MRS (MT+ and V1 regions,

in random sequence) experiments, but only part of the MRS data (31/36 in hMT+ region

and 28/36 in V1 region) survived quality control (see the part of MRS date processing).

The sample size is determined by the statistic requirement (30 sample for Person

correlation statistical analysis).

Motion surrounding suppression measurement

All stimuli were generated using Matlab (MathWorks, Natick, MA) with Psychophysics

Toolbox(Brainard, 1997), and were shown on a linearized monitor (1920×1080

resolution, 100-Hz refresh rate, Cambridge Research System, Kent, UK). The viewing

distance was 72 cm from the screen, with the head stabilized by a chinrest. Stimuli were

drawn against a gray (56 cd per m^{-2}) background.

A schematic of the stimuli and trial sequences is shown in our recent study(Song et

al., 2021). The stimulus was a vertical drifting sinusoidal grating (contrast, 50%; spatial

frequency, 1 cycle/°; speed, 4°/s) of either small (diameter of 2°) or large (diameter of

10°) size. The edge of the grating was blurred with a raised cosine function (width,

0.3°). A cross was presented in the center of the screen at the beginning of each trial for

500ms, and participants were instructed to fixate at the cross and to keep fixating at the

cross throughout the trial. In each trial, a grating of either large or small size was

randomly presented at the center of the screen. The grating drifted either leftward or

rightward, and participants were asked to judge the perceived moving direction by a

key press. Response time was not limited. The grating was ramped on and off with a

Gaussian temporal envelope, and the grating duration was defined as 1 SD of the

Gaussian function. The duration was adaptively adjusted in each trial, and duration

thresholds were estimated by a staircase procedure. Thresholds for large and small

gratings were obtained from a 160-trial block that contained four interleaved 3-down/1-

up staircases. For each participant, we computed the correct rate for different stimulus

durations separately for each stimulus size. These values were then fitted to a

cumulative Gaussian function, and the duration threshold corresponding to the 75%

correct point on the psychometric function was estimated for each stimulus size.

Stimulus demonstration and practice trials were presented before the first run.

Auditory feedback was provided for each wrong response. To quantify the spatial

suppression strength, we calculated the spatial suppression index (SI), defined as the

difference of log10 thresholds for large versus small stimuli(Schallmo et al., 2018;

Tadin et al., 2003):

$$SI = log_{10}(\ large\ threshold) - log_{10}(\ small\ threshold) \qquad (1)$$

Block design task measurement

The block design task was administered in accordance with the WAIS-IV

manual(Wechsler, 2008). Specifically, participants were asked to rebuild the figural

pattern within a specified time limit using a set of red and white blocks. The time limits

were set as 30 s to 120 s according to different levels of difficulty. The patters were

presented in ascending order of difficulty, and the test stopped if two consecutive

patterns were not constructed in the allotted time. The score was determined by the

accomplishment of the pattern and the time taken. A time bonus was awarded for rapid

performance in the last six patterns. The score ranges between 0 and 66 points, with

higher scores indicating better perceptual reasoning.

MR experimental procedure

MR experiments were performed in a 7T whole body MR system (Siemens Healthcare,

Erlangen, Germany) with a Nova Medical 32 channel array head coil. Sessions included

resting-state functional MRI, fMRI localizer scan, structural image scanning, and MRS

scan. Resting-state scans were acquired with 1.5-mm isotropic resolution (transverse

orientation, TR/TE = 2000/20.6 ms, 160 volumes, slice number = 90, flip angle = 70°,

eyes closed). Structural images were acquired using a MP2RAGE sequence (TR/ TI1/

TI2 = 5000/901/3200 ms) with 0.7-mm isotropic resolution. MRS data were collected

within two regions (MT+ and V1) for each subject, and we divided them into two

sessions to avoid discomfort caused by long scanning. The order of MRS VOIs (MT+

and V1) in the two sessions was counterbalanced across participants. Interval between

two sessions was used for block design and motion discrimination tasks. One session

included fMRI localizer scan, structural image scanning, and MRS scan for the hMT+

region; the other session included structural image scan, and MRS scan for the V1

region. Spectroscopy data were acquired using a ^{1}H-MRS single-voxel short-TE

STEAM (Stimulated Echo Acquisition Mode) sequence(Frahm et al., 1989)

(TE/TM/TR = 6/32/7100ms) with 4096 sampling points, 4-kHz bandwidth, 16 averages,

8 repetitions, 20×20×20 mm^3 VOI size, and VAPOR (variable power and optimized

relaxation delays) water suppression(Tkáč et al., 1999). Prior to acquisition, first- and

second-order shims were adjusted using FASTMAP (fast, automatic shimming

technique by mapping along projections(Gruetter, 1993). Two non-suppressed water

spectra were also acquired: one for phase and eddy current correction (only RF pulse,

4 averages) and another for metabolite quantification (VAPOR none, 4 averages).

Voxels were positioned based on anatomical landmarks using a structural image scan

collected in the same session, while avoiding contamination by CSF, bone, and fat. The

hMT+ VOIs were placed in the ventrolateral occipital lobe, which was based on anatomical landmarks(Dumoulin et al., 2000; Schallmo et al., 2018). We did not distinguish between the middle temporal (MT) and medial superior temporal (MST) areas in these hMT+ VOIs(Huk et al., 2002). For 14 subjects, we also functionally identified hMT+ as a check on the placement of the VOI. A protocol was used with a drifting grating (15% contrast) alternated with a static grating across blocks (10 s block duration, 160 TRs total). Using fMRI BOLD signals, these localizer data were processed online to identify the hMT+ voxels in the lateral occipital cortex, which responded more strongly to moving vs. static gratings. In addition, we only used the left hMT+ as the target region to scan, which was motivated by studies showing that left hMT+ was more effective at causing perceptual effects(Tadin et al., 2011). For V1 region, the VOI was positioned on each subject's calcarine sulcus on the left side(Tadin et al., 2011) based on anatomical landmarks(Boucard et al., 2007; Dumoulin et al., 2000)

MRS data processing

Spectroscopy data were preprocessed and quantified using magnetic resonance signal processing and analysis, https://www.cmrr.umn.edu/ downloads/mrspa/), which runs under MATLAB and invokes the interface of the LCModel (Version 6.3-1L) (Chen et al., 2019). First, we used the non-suppressed water spectra to perform eddy current correction and frequency/phase correction. Second, we checked the quality of each FID (16 averages) visually and removed those with obviously poor quality. Third, the absolute concentrations of each metabolite were quantitatively estimated via the Water-

Scaling method. For partial-volume correction, the tissue water content was computed

as follows(Ernst et al., 1993):

$Tissue\ water\ content = fgm*0.78 + fwm*0.65 + fcsf*0.97$ $\qquad$ (2)

where *fgm*, *fwm*, and *fcsf* were the GM/WM/CSF volume fraction in MRS VOI and

we used FAST (fMRI's automated segmentation tool, part of the FSL toolbox)(Zhang

et al., 2001) to segment the three tissue compartments from the T1-weighted structural

brain images. For water T2 correction, we set water T2 as 47ms (Marjańska et al., 2012).

Our concentrations were mM per kg wet weight. Furthermore, LCModel analysis was

performed on all spectra within the chemical shift range of 0.2 to 4.0 ppm.

Poor spectral quality was established by a Cramer-Rao Lower Bound (CRLB) of

more than 20%(Cavassila et al., 2001), and some data were excluded from further

analysis. The details were described in our recently paper(Song et al., 2021).

Rs-fMRI data processing and analysis

Resting-state functional image was analyzed in the Data Processing and Analysis for

Brain Imaging DPABI toolbox(Yan et al., 2016) based on SPM 12

(http://www.fil.ion.ucl.ac.uk/spm/). The preprocessing steps included discard of the

first five volumes, slice timing, realignment to the 90th slice, coregistration of each

subject's T1-weighted anatomical and functional images, segmentation of the

anatomical images into six types of tissues using DARTEL, linear detrend, regressing

nuisance variables (including realignment Friston 24-parameter, global signal, white

matter and CSF signal) (Friston et al., 1996), normalization to the standard Montreal

Neurological Institute (MNI) space with the voxel size of $1.5\times 1.5\times 1.5$ mm^3 using

DARTEL, spatial smoothing with a Gaussian kernel of 3 mm full-width-half-maximum

(FWHM), and band-pass filtering with Standard frequency band (SFB, 0.01–0.1 Hz).

Spherical ROI with a radius of 6mm was placed in left MT. The coordinate for left MT

(-46, -72, -4, in MNI space) was obtained by our localizer fMRI experiment. We

calculated the seed-to-voxel whole brain FC map for each subject. All the FC values

were Fisher-Z-transformed.

We did a similar connectivity-behavior analysis to a previous study (Song et al.,

2008). First, we computed the Pearson's correlation coefficient between BDT scores

and the FC values across subjects in a voxel-based way. Then, to evaluate the

significance, we transformed the r-value into t-value ($t = \frac{\sqrt{df}\cdot r}{\sqrt{1-r^2}}$), where df denotes to

the degrees of freedom, and r is the Pearson's correlation coefficient between BDT

scores and the FC values. Here, df was equal to 27. The brain regions in which the FC

values to the seed region was significantly correlated with the BDT scores were

obtained with a threshold of $P < 0.005$ for regions of *a priori* ($|t_{(27)}| \geq 3.057$, and

adjacent cluster size ≥ 23 voxels (AlphaSim corrected)), and $P < 0.01$ for whole-brain

analyses ($|t (27)| \geq 2.771$ and adjacent cluster size ≥ 37 voxels (AlphaSim corrected).

Statistical Analysis

PROCESS version 3.4, a toolbox in SPSS, was used to examine the mediation model.

There are some prerequisites for mediation analysis: the independent variable should be a significant predictor of the mediator, and the mediator should be a significant predictor of the dependent variable.

SPSS 20 (IBM, USA) was used to conduct all the remaining statistical analysis in the study. We evaluated the correlation of variables (GABA, Glu, SI, BDT) using Pearson's correlation analysis. Differences or correlations were considered statistically significant if $P < 0.05$. Significances with multiple comparisons were tested with false discovery rate (FDR) correction. The effect of age on intelligence was controlled for by using partial correlation in the correlation analysis and was taken as a covariate in the serial mediation model analysis.

Data availability

Source data are provided with this paper and have been archived at GitHub and could be downloaded with reasonable request. https://github.com/Yrehearsal/GABA-_hMT-_Intelligence

.

Code availability

This paper does not report original code.

References

Angelucci, A., Levitt, J. B., Walton, E. J., Hupe, J.-M., Bullier, J., & Lund, J. S. (2002). Circuits for local and global signal integration in primary visual cortex. *Journal of Neuroscience*, *22*(19), 8633-8646.

Assem, M., Glasser, M. F., Van Essen, D. C., & Duncan, J. (2020). A domain-general cognitive core defined
in multimodally parcellated human cortex. *Cerebral Cortex*, *30*(8), 4361-4380.

Boucard, C. C., Hoogduin, J. M., van der Grond, J., & Cornelissen, F. W. (2007). Occipital proton magnetic
resonance spectroscopy (1H-MRS) reveals normal metabolite concentrations in retinal visual
field defects. *PLoS One*, *2*(2), e222.

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis*, *10*(4), 433-436.

Carandini, M., & Heeger, D. J. (2011). Normalization as a canonical neural computation. *Nat Rev Neurosci*,
13(1), 51-62. https://doi.org/10.1038/nrn3136

Cattell, & Raymond, B. (1963). Theory of fluid and crystallized intelligence: A critical experiment. *Journal*
of Educational Psychology, *54*(1), 1-22.

Cavassila, S., Deval, S., Huegen, C., Van Ormondt, D., & Graveron‑Demilly, D. (2001). Cramér–Rao
bounds: an evaluation tool for quantitation. *NMR in Biomedicine: An International Journal*
Devoted to the Development and Application of Magnetic Resonance In Vivo, *14*(4), 278-283.

Chen, X., Fan, X., Hu, Y., Zuo, C., Whitfield-Gabrieli, S., Holt, D., Gong, Q., Yang, Y., Pizzagalli, D. A., & Du,
F. (2019). Regional GABA concentrations modulate inter-network resting-state functional
connectivity. *Cerebral Cortex*, *29*(4), 1607-1618.

Colom, R., Jung, R. E., & Haier, R. J. (2006). Distributed brain sites for the g-factor of intelligence.
Neuroimage, *31*(3), 1359-1365. https://doi.org/10.1016/j.neuroimage.2006.01.006

Cumming, B. G., & DeAngelis, G. C. (2001). The physiology of stereopsis. *Annu Rev Neurosci*, *24*, 203-
238. https://doi.org/10.1146/annurev.neuro.24.1.203

Deary, I. J., Penke, L., & Johnson, W. (2010). The neuroscience of human intelligence differences. *Nature*
reviews neuroscience, *11*(3), 201-211.

Donner, T. H., Siegel, M., Fries, P., & Engel, A. K. (2009). Buildup of choice-predictive activity in human
motor cortex during perceptual decision making. *Curr Biol*, *19*(18), 1581-1585.
https://doi.org/10.1016/j.cub.2009.07.066

Dumoulin, S. O., Bittar, R. G., Kabani, N. J., Baker, C. L., Jr., Le Goualher, G., Bruce Pike, G., & Evans, A. C.
(2000). A new anatomical landmark for reliable identification of human area V5/MT: a
quantitative analysis of sulcal patterning. *Cereb Cortex*, *10*(5), 454-463.
https://doi.org/10.1093/cercor/10.5.454

Duncan, J., Assem, M., & Shashidhara, S. (2020). Integrated Intelligence from Distributed Brain Activity.
Trends Cogn Sci, *24*(10), 838-852. https://doi.org/10.1016/j.tics.2020.06.012

Duncan, J., Seitz, R. J., Kolodny, J., Bor, D., Herzog, H., Ahmed, A., Newell, F. N., & Emslie, H. (2000). A
neural basis for general intelligence. *Science*, *289*(5478), 457-460.
https://doi.org/10.1126/science.289.5478.457

Ende, G. (2015). Proton Magnetic Resonance Spectroscopy: Relevance of Glutamate and GABA to
Neuropsychology. *Neuropsychol Rev*, *25*(3), 315-325. https://doi.org/10.1007/s11065-015-
9295-8

Ernst, T., Kreis, R., & Ross, B. (1993). Absolute quantitation of water and metabolites in the human brain.
I. Compartments and water. *Journal of magnetic resonance, Series B*, *102*(1), 1-8.

Everling, S., Tinsley, C. J., Gaffan, D., & Duncan, J. (2002). Filtering of neural signals by focused attention
in the monkey prefrontal cortex. *Nature neuroscience*, *5*(7), 671-676.

Fangmeier, T., Knauff, M., Ruff, C. C., & Sloutsky, V. (2006). fMRI evidence for a three-stage model of
deductive reasoning. *Journal of cognitive neuroscience*, *18*(3), 320-334.

Fetsch, C. R., Pouget, A., DeAngelis, G. C., & Angelaki, D. E. (2011). Neural correlates of reliability-based
cue weighting during multisensory integration. *Nat Neurosci*, *15*(1), 146-154.
https://doi.org/10.1038/nn.2983

Frahm, J. a., Bruhn, H., Gyngell, M., Merboldt, K., Hänicke, W., & Sauter, R. (1989). Localized high‐
resolution proton NMR spectroscopy using stimulated echoes: initial applications to human
brain in vivo. *Magnetic resonance in medicine*, *9*(1), 79-93.

Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S., & Turner, R. (1996). Movement‐related effects
in fMRI time‐series. *Magnetic resonance in medicine*, *35*(3), 346-355.

Gautama, T., & Van Hulle, M. M. (2001). Function of center-surround antagonism for motion in visual
area MT/V5: a modeling study. *Vision Res*, *41*(28), 3917-3930. https://doi.org/10.1016/s0042-
6989(01)00246-2

Gray, J. R., Chabris, C. F., & Braver, T. S. (2003). Neural mechanisms of general fluid intelligence. *Nat*
Neurosci, *6*(3), 316-322. https://doi.org/10.1038/nn1014

Gruetter, R. (1993). Automatic, localized in vivo adjustment of all first‐and second‐order shim coils.
Magnetic resonance in medicine, *29*(6), 804-811.

Gu, Y., Watkins, P. V., Angelaki, D. E., & DeAngelis, G. C. (2006). Visual and nonvisual contributions to
three-dimensional heading selectivity in the medial superior temporal area. *J Neurosci*, *26*(1),
73-85. https://doi.org/10.1523/jneurosci.2356-05.2006

Haier, R. J., Siegel Jr, B. V., Nuechterlein, K. H., Hazlett, E., Wu, J. C., Paek, J., Browning, H. L., &
Buchsbaum, M. S. (1988). Cortical glucose metabolic rate correlates of abstract reasoning and
attention studied with positron emission tomography. *Intelligence*, *12*(2), 199-217.

Hayes, A. F. (2013). *Introduction to Mediation, Moderation, and Conditional Process Analysis: A*
Regression‐Based Approach (Vol. 51). The Guilford Press.

Huk, A. C., Dougherty, R. F., & Heeger, D. J. (2002). Retinotopy and functional subdivision of human areas
MT and MST. *Journal of Neuroscience*, *22*(16), 7195-7205.

Jung, R. E., & Haier, R. J. (2007). The Parieto-Frontal Integration Theory (P-FIT) of intelligence: converging
neuroimaging evidence. *Behav Brain Sci*, *30*(2), 135-154; discussion 154-187.
https://doi.org/10.1017/S0140525X07001185

Keller, A. J., Roth, M. M., & Scanziani, M. (2020). Feedback generates a second receptive field in neurons
of the visual cortex. *Nature*, *582*(7813), 545-549.

Li, M., Song, X. M., Xu, T., Hu, D., Roe, A. W., & Li, C.-Y. (2019). Subdomains within orientation columns
of primary visual cortex. *Science advances*, *5*(6), eaaw0807.

Liu, D. Y., Ju, X., Gao, Y., Han, J. F., Li, Z., Hu, X. W., Tan, Z. L., Northoff, G., & Song, X. M. (2022). From
Molecular to Behavior: Higher Order Occipital Cortex in Major Depressive Disorder. *Cereb*
Cortex, *32*(10), 2129-2139. https://doi.org/10.1093/cercor/bhab343

Liu, L. D., Haefner, R. M., & Pack, C. C. (2016). A neural basis for the spatial suppression of visual motion
perception. *Elife*, *5*, e16167.

Marjańska, M., Auerbach, E. J., Valabrègue, R., Van de Moortele, P. F., Adriany, G., & Garwood, M. (2012).
Localized 1H NMR spectroscopy in different regions of human brain in vivo at 7 T: T2 relaxation
times and concentrations of cerebral metabolites. *NMR in Biomedicine*, *25*(2), 332-339.

Melnick, M. D., Harrison, B. R., Park, S., Bennetto, L., & Tadin, D. (2013). A strong interactive link between
sensory discriminations and intelligence. *Curr Biol*, *23*(11), 1013-1017.
https://doi.org/10.1016/j.cub.2013.04.053

Ozeki, H., Finn, I. M., Schaffer, E. S., Miller, K. D., & Ferster, D. (2009). Inhibitory stabilization of the
cortical network underlies visual surround suppression. *Neuron*, *62*(4), 578-592.

Reynolds, J. H., & Heeger, D. J. (2009). The normalization model of attention. *Neuron*, *61*(2), 168-185.

Sato, T. K., Haider, B., Häusser, M., & Carandini, M. (2016). An excitatory basis for divisive normalization
in visual cortex. *Nature neuroscience*, *19*(4), 568-570.

Schallmo, M. P., Kale, A. M., Millin, R., Flevaris, A. V., Brkanac, Z., Edden, R. A., Bernier, R. A., & Murray,
S. O. (2018). Suppression and facilitation of human neural responses. *Elife*, *7*.
https://doi.org/10.7554/eLife.30334

Siegel, M., Buschman, T. J., & Miller, E. K. (2015). Cortical information flow during flexible sensorimotor
decisions. *Science*, *348*(6241), 1352-1355. https://doi.org/10.1126/science.aab0551

Song, M., Zhou, Y., Li, J., Liu, Y., Tian, L., Yu, C., & Jiang, T. (2008). Brain spontaneous functional
connectivity and intelligence. *Neuroimage*, *41*(3), 1168-1176.
https://doi.org/10.1016/j.neuroimage.2008.02.036

Song, X. M., Hu, X. W., Li, Z., Gao, Y., Ju, X., Liu, D. Y., Wang, Q. N., Xue, C., Cai, Y. C., Bai, R., Tan, Z. L., &
Northoff, G. (2021). Reduction of higher-order occipital GABA and impaired visual perception
in acute major depressive disorder. *Mol Psychiatry*, *26*(11), 6747-6755.
https://doi.org/10.1038/s41380-021-01090-5

Spearman, C. (1904). " General Intelligence" Objectively Determined and Measured. *American Journal*
of Psychology, *15*, 201-293.

Tadin, D. (2015). Suppressive mechanisms in visual motion processing: From perception to intelligence.
Vision Res, *115*(Pt A), 58-70. https://doi.org/10.1016/j.visres.2015.08.005

Tadin, D., Lappin, J. S., Gilroy, L. A., & Blake, R. (2003). Perceptual consequences of centre-surround
antagonism in visual motion processing. *Nature*, *424*(6946), 312-315.
https://doi.org/10.1038/nature01800

Tadin, D., Silvanto, J., Pascual-Leone, A., & Battelli, L. (2011). Improved motion perception and impaired
spatial suppression following disruption of cortical area MT/V5. *Journal of Neuroscience*, *31*(4),
1279-1283.

Tkáč, I., Starčuk, Z., Choi, I. Y., & Gruetter, R. (1999). In vivo 1H NMR spectroscopy of rat brain at 1 ms
echo time. *Magnetic Resonance in Medicine: An Official Journal of the International Society for*
Magnetic Resonance in Medicine, *41*(4), 649-656.

Treue, S., & Maunsell, J. H. (1996). Attentional modulation of visual motion processing in cortical areas
MT and MST. *Nature*, *382*(6591), 539-541.

Wang, X. J. (2002). Probabilistic decision making by slow reverberation in cortical circuits. *Neuron*, *36*(5),
955-968. https://doi.org/10.1016/s0896-6273(02)01092-9

Wechsler, D. (2008). Wechsler Memory Scale–Fourth Edition (WMS-IV) technical and interpretive
manual.

Wilming, N., Murphy, P. R., Meyniel, F., & Donner, T. H. (2020). Large-scale dynamics of perceptual
decision information across human cortex. *Nat Commun*, *11*(1), 5109.
https://doi.org/10.1038/s41467-020-18826-6

Wimmer, K., Compte, A., Roxin, A., Peixoto, D., Renart, A., & de la Rocha, J. (2015). Sensory integration
dynamics in a hierarchical network explains choice probabilities in cortical area MT. *Nat*
Commun, *6*, 6177. https://doi.org/10.1038/ncomms7177

Yan, C.-G., Wang, X.-D., Zuo, X.-N., & Zang, Y.-F. (2016). DPABI: data processing & analysis for (resting-

state) brain imaging. *Neuroinformatics*, *14*, 339-351.

Zaksas, D., & Pasternak, T. (2006). Directional signals in the prefrontal cortex and in area MT during a

working memory for visual motion task. *J Neurosci*, *26*(45), 11726-11742.

https://doi.org/10.1523/JNEUROSCI.3420-06.2006

Zhang, S., Xu, M., Kamigaki, T., Hoang Do, J. P., Chang, W.-C., Jenvay, S., Miyamichi, K., Luo, L., & Dan, Y.

(2014). Long-range and local circuits for top-down modulation of visual cortex processing.

Science, *345*(6197), 660-665.

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov

random field model and the expectation-maximization algorithm. *IEEE transactions on medical*

imaging, *20*(1), 45-57.

Acknowledgements

The authors thank Prof. Dost Ongur and Fei Du for guidance on the MRS data processing, and thank Zhejiang University 7T Brain Imaging Research Center. Thank Prof. Xinyi Lai for supporting MRI data acquisition. Thank Fen Yang for technical assistance This work was supported by STI 2030—Major Projects (2021ZD0200401 to X.M.S., 2022ZD0206000 to R.B.), the National Natural Science Foundation of China Grants (U1909205, 61876222, 32000761, 82222032), Humanities and Social Sciences Ministry of Education (20YJC880095, 18YJA190001), the Key R&D Program of Zhejiang (2022C03096 to X.M.S., 2022ZJJH02-06 to G.C.), the European Union's Horizon 2020 Framework Program for Research and Innovation under the Specific Grant Agreement No. 785907 (Human Brain Project SGA2 to G.N.), and the MOE Frontier Science Center for Brain Science & Brain- Machine Integration, Zhejiang University.

Author contributions

856 S.X.M. and Y.G. designed the experiment and G.N. guided the logic of the analysis.

857 Y.G., Y.C. and D.L. conducted human experiments, analyzed data, and Y.G. created

858 figures. R.B. guided the MRI experiments. J.Y., J.W. and B.X. were in charge of data

859 collection and partial date analysis. T. W., M.L. and G.C. guided the data analysis.

860 S.X.M., Y.G. and G.N. wrote the manuscript.

Competing interests

863 The authors declare no competing interests.