GABA-ergic inhibition in human hMT+ predicts visuo-spatial intelligence mediated through the frontal cortex

Yuan Gao[1#], Yong-Chun Cai[2#], Dong-Yu Liu[1,3], Juan Yu[1,3], Jue Wang[1], Ming li[4], Bin Xu[1], Teng-Fei Wang[2], Gang Chen[5], Georg Northoff[6*], Ruiliang Bai[3,7*], Xue Mei Song[1,3*]

[1]Department of Neurosurgery of the Second Affiliated Hospital, Interdisciplinary Institute of Neuroscience and Technology, Zhejiang University School of Medicine, Hangzhou 310029, China;

[2]Department of Psychology and Behavioral Sciences, Zhejiang University, Hangzhou 310028, China;

[3]Key Laboratory of Biomedical Engineering of Ministry of Education, Qiushi Academy for Advanced Studies, College of Biomedical Engineering and Instrument Science, Zhejiang University, Hangzhou 310027, China;

[4]College of Intelligence Science and Technology, National University of Defense Technology, Changsha 410073, China;

[5]Oujiang Laboratory (Zhejiang Lab for Regenerative Medicine, Vision and Brain Health), Wenzhou, Zhejiang 325603, China;

[6]University of Ottawa Institute of Mental Health Research, University of Ottawa; Ottawa, ON, K1Z 7K4, Canada;

[7]MOE Frontier Science Center for Brain Science & Brain- Machine Integration, Zhejiang University，Hangzhou 311121, China;

[#]Co-first author

23 *Correspondence: songxuemei@zju.edu.cn, ruiliangbai@zju.edu.cn,

24   georg.northoff@theroyal.ca

Abstract

The prevailing opinion emphasizes fronto-parietal network (FPN) is key in mediating

general fluid intelligence (gF). Meanwhile, recent studies show that human MT

complex (hMT+), located at the occipito-temporal border and involved in 3D

perception processing, also plays a key role in gF. However, the underlying

mechanism is not clear, yet. To investigate this issue, our study targets visuo-spatial

intelligence, which is considered to have high loading on gF. We use ultra-high field

magnetic resonance spectroscopy (MRS) to measure GABA/glutamate concentrations

in hMT+ combining resting-state fMRI functional connectivity (FC), behavioral

examinations including hMT+ perception suppression test and gF subtest in visuo-

spatial component. Our findings show that both GABA in hMT+ and frontal-hMT+

functional connectivity significantly correlate with the performance of visuo-spatial

intelligence. Further, serial mediation model demonstrates that the effect of hMT+

GABA on visuo-spatial gF is fully mediated by the hMT+ frontal FC. Together our

findings highlight the importance in integrating sensory and frontal cortices in

mediating the visuospatial component of general fluid intelligence.

Introduction

General fluid intelligence (gF) is a current problem-solving ability, which shows high

inter-individual differences in humans(Cattell & Raymond, 1963). At the beginning of

the last century, Spearman(Spearman, 1904) proposed that some general or g factor

contributes to our gF. One key component of gF is visuo-spatial intelligence, usually

tested by visual materials, shows high g-loading(Colom et al., 2006; Deary et al.,

2010; Jung & Haier, 2007). The exact neural mechanisms of the interplay of visuo

spatial intelligence with gF remain yet unclear, though.

The"neuro-efficiency" hypothesis is one explanation for individual differences in

gF (Haier et al., 1988). This hypothesis puts forward that the human brain's ability to

suppress irrelevant information leads to more efficient cognitive processing.

Correspondingly, using a well-known visual motion paradigm (center-surround

antagonism)(Liu et al., 2016; Tadin et al., 2003), Melnick et al found a strong link

between suppression index (SI) of motion perception and the scores of the block

design test (BDT, a subtest of the Wechsler Adult Intelligence Scale (WAIS), which

measures the visuo-spatial component (3D domain) of gF (Melnick et al., 2013).

Motion surround suppression (SI), a specific function of human extrastriate cortical

region, middle temporal complex (hMT+), aligns closely with this region's activities

(Gautama & Van Hulle, 2001). Furthermore, hMT+ is a sensory cortex involved in

visual perception processing (3D domain) (Cumming & DeAngelis, 2001). These

findings suggest that hMT+ potentially plays a significant role in 3D visuo-spatial gF

by facilitating the efficient processing of 3D visual information and suppressing

irrelevant information. However, more evidence is needed to uncover how the hMT+

functions as a core region for 3D visuo-spatial intelligence.

Frontal cortex is usually recognized as the cognitive core region (Duncan et al.,

2000; Gray et al., 2003). Strong connectivity between the cognitive regions suggests a

mechanism for large-scale information exchange and integration in the brain(Barbey,

2018; Cole et al., 2012). Therefore, the potential conjunctive coding may overlap

with the inhibition and/or excitation mechanism of hMT+. Taken together, we

hypothesized that 3D visuo-spatial intelligence (as measured by BDT) might be

predicted by the inhibitory and/or excitation mechanisms in hMT+ and the integrative

functions connecting hMT+ with frontal cortex (Figure 1a).

To investigate our hypothesis, this work conducted multi-level examination

including biochemical (Glutamatergic - GABAergic in hMT+), regional-systemic

(brain connectivity with hMT+ - based), and behavioral (visual motion function in

hMT+) levels to reveal if hMT+ contributes to the 3D visuo-spatial component of gF.

We employ ultra-high field (7T) magnetic resonance spectroscopy (MRS) technology

to reliably resolve GABA and Glu concentrations(Ende, 2015; Liu et al., 2022; Song

et al., 2021). To verify the specificity of hMT+, we used early visual cortex (EVC,

primarily in V1) - based GABA/Glu as control as it mediates the 2D rather than 3D

visual domain(Cumming & DeAngelis, 2001).

Our findings first demonstrate that GABAergic inhibition mechanisms (but not

excitatory Glu) in hMT+ region relate to 3D visuo-spatial ability. Further, analysis of

functional brain connectivity at rest reveals that the network (between MT+ and

frontal cortex) relating to MT+ GABA and perceptual suppression contribute the

visuo-spatial intelligence. Our results provide direct evidence that inhibitory

mechanisms centered on GABA levels in MT+ region (a sensory cortex) mediate

multi-level visuo-spatial component (3D domain) of gF thus drawing a direct

connection of biochemistry, brain connectivity, and behavior.

Figure 1. Hypothesis and experimental design. (**a**) Schematic of hypothesis. The inhibition

mechanism centered on MT+ GABA, including the molecular level: the GABAergic inhibition in

MT+ (cyan circle), brain connectivity level: hMT+ - frontal functional connectivity (blue circle),

and behavior level: hMT+ specific surround suppression of visual motion (red circle), contributes

to the visuo-spatial component of gF (3D domain, yellow circle). (**b**) Schematic of experimental

design. Session 1 (rectangle box of short line) was the functional MRI and MRS scanning at

resting state. Session 2 (rectangle box of solid line) was another region of MRS acquisition. In the

two sessions, the order of MRS scanning regions (hMT+ and EVC (primarily in V1)) was counter

balanced across participants. There was a structure MRI scanning before each MRS data

acquisition. The interval between the two sessions was used for behavioral measurement

(rectangle box of dotted line): block design task (BDT) and psychophysical task - motion

discrimination. Sold lines indicate the experiment sequence.

Results

To determine whether the function of hMT+ cortex contributes to visuo-spatial

component (3D domain) of gF, we adopted the experimental design depicted in

Figure 1b. Thirty-six healthy subjects participated in this study. Participants

underwent two MRI sessions: the first encompassing resting-state fMRI and magnetic

resonance spectra (MRS), and the second solely involving MRS. A 30-minute interval

separated these sessions, during which participants performed motion discrimination

tasks (using center-surround antagonism stimuli)(Tadin et al., 2003) and the block

design test (BDT), which assesses the visuo-spatial ability (3D domain) of

gF(Fangmeier et al., 2006). In the motion discrimination tasks, a grating of either

large or small size was randomly presented at the center of the screen. The grating

drifted either leftward or rightward, and participants were asked to judge the

perceived moving direction. While in the Block design test, participants were asked to

rebuild the figural pattern within a specified time limit using a set of red and white

blocks. Both the volume-of-interests (VOIs) of MRS scanning in the left hMT+

(targeted brain area) and the left EVC (primarily in V1, control brain area) had

dimensions of $2\times2\times2$ cm^3, and the MRS scanning sequences were randomized across

the two sessions. The hMT+ MRS VOIs were demarcated using an anatomical

landmark(Dumoulin et al., 2000). For 14 subjects, we also utilized fMRI to

functionally pinpoint the hMT+ to validate the placement of the VOI (Figure 2a, b).

The EVC (primarily in V1) MRS VOIs (figure supplement 1) were anatomically

defined (Methods). Here, MRS data after extensive quality control (31/36 in hMT+,

and 28/36 in EVC (primarily in V1)) were taken for further analysis (Methods).

Figure 2. hMT+ localizer scans and hMT+ MRS VOI placement. (**a**) Single task block designs.

First: a cross fixation on the center of the screen (10s). Second: a moving grating (2^{o}) toward left

last 10s. Third: the grating keeps static for 10s. Fourth: the grating moves toward right last 10s.

The localizer scans consist of 8 blocks. (**b**) hMT+ location and MRS VOI placement. The upper

template is the horizontal view. The lower templates from left to right are coronal and sagittal

views. The warm color indicates the overlap of fMRI activation of hMT+ across 14 subjects, the

cold color bar indicates the overlap of MRS VOIs across all subjects.

Figure supplement 1. The left hemisphere EVC (primarily in V1) MRS scanning ROI. The color

bar indicates the overlap of the MRS VOIs across all subjects.

GABA and Glu concentrations in hMT+ and EVC (primarily in V1) and their

relation to SI and BDT

An example of a MRS voxel located in hMT+ is shown in Figure 3a. LCModel

fittings for GABA spectra from all subjects in hMT+ ($n = 31$) and EVC (primarily in

V1) ($n = 28$) are illustrated in Figure 3b (color scale presents the BDT scores). We

discerned a significant association between the inter-subjects' BDT scores and the

GABA levels in hMT+ voxels, but not in EVC (primarily in V1) voxels. Quantitative

analysis displayed that BDT significantly correlates with GABA concentrations in

hMT+ voxels ($r = 0.39$, $P = 0.03$, $n = 31$, Figure 3c). After using partial correlation to

control for the effect of age, the relationship remains significant ($r_{partial} = 0.426$ $P =$

0.02, 1 participant excluded due to the age greater than mean + 2.5SD) In contrast,

there was no obvious correlation between BDT and GABA levels in EVC (primarily

in V1) voxels (figure supplement 2a). We show that SI significantly correlates with

GABA levels in hMT+ voxels ($r = 0.44$, $P = 0.01$, $n = 31$, Figure 3d). In contrast, no

significant correlation between SI and GABA concentrations in EVC (primarily in

V1) voxels was observed (figure supplement 2b). These findings suggest that the

relationship between motion suppression and GABA+ is specific to hMT+, but not in

EVC (primarily in V1), which

is similar with prior observation (Schallmo et al., 2018). LCModel fittings for Glu

spectra from all subjects in hMT+ ($n = 31$) and EVC (primarily in V1) ($n = 28$) voxels

are presented in figure supplement 3a.

Unlike in the case of GABA, no significant correlations between BDT and Glu

levels were found in both hMT+ and EVC (primarily in V1) voxels (figure

supplement 3b, c). While, as expected(Song et al., 2021), we observed significant

positive correlations between GABA and Glu concentrations in both hMT+ ($r = 0.62$,

$P = 0.0002$, $n = 31$) and EVC (primarily in V1) voxels ($r = 0.56$, $P = 0.002$, $n = 28$)

(figure supplement 4a, b). Additionally, significant correlations between SI and BDT,

duration threshold of small grating and BDT was discerned ($r = 0.59$, $P = 0.0002$, $n =$

34, Figure 3e, $r_{partial} = 0.67$, $P < 0.001$, 1 participant excluded due to the age greater

than mean + 2.5SD; $r = -0.43$, $P = 0.016$, $r_{partial} = 0.44$, $P = 0.014$, Figure 3f). While

there was no significant correlation between duration threshold of large grating and

BDT (Figure 3g), corroborating previous conclusions(Melnick et al., 2013). Two

outliers evident in Figure 3e were excluded, with consistent results depicted in figure

supplement 5a. Further, two outliers evident in Figure 3d were excluded, with

consistent results depicted in figure supplement 5b.

Figure 3. MRS spectra and the relationships between GABA levels and SI / BDT. (**a**) Example spectrum from the hMT+ voxel of one participant. The first line is the LCModel fitting result of all metabolites, and the following lines show the Glu and GABA spectra fitting with LCModel, and then the baseline. (**b**) Individual participants fitted GABA MRS spectra from the hMT+ (top) and EVC (primarily in V1) (bottom) voxels from baseline measurement. The colors of the GABA spectra represent the individual differences of BDT. The color bar represents the scores of BDT. (**c**) **and (d)** Pearson's correlations showing significant positive correlations between hMT+ GABA and BDT scores (**c**), between hMT+ GABA and SI (**d**). (**e**) Peason's correlation showing significant positive correlations between SI and BDT. (**f**) Peason's correlation showing significant negative correlations between BDT and duration threshold of small grating. (**g**) No correlation between BDT and duration threshold of large grating. The ribbon between dotted lines represents the 95% confidence interval, and the black regression line represents the Pearson's correlation coefficient (*r*). GABA and Glu concentrations (Conc.) are absolute, with units of mmol per kg wet weight (Methods).

Figure supplement 2. Relationships between GABA concentration in EVC (primarily in V1) and

BDT/SI. (a) There is no significant relationship between BDT and GABA concentrations in EVC

(primarily in V1) region. (b) There is also no obvious correlation between SI and GABA

concentrations in EVC (primarily in V1) region. GABA concentration (Conc.) is absolute, with

units of mmol per kg wet weight (Methods).

Figure supplement 3. Individual Glu MRS spectra from hMT+ / EVC (primarily in V1) regions

and relationships between BDT and Glu concentrations in hMT+ / EVC (primarily in V1) regions.

(a) Individual participant fitted Glutamate MRS spectra from the hMT+ (top, $n = 31$) and EVC

(primarily in V1) (bottom, $n = 28$) voxels from baseline measurement. The colors of the Glutamate

spectra represent the individual differences of BDT. The color bar represents the scores of BDT. (b)

There is no significant relationship between BDT and Glu concentrations in hMT+ region. c There

is also no significant correlation between BDT and Glu concentrations in EVC (primarily in V1)

region. Glutamate concentration (Conc.) is absolute, with units of mmol per kg wet weight

(Methods).

Figure supplement 4. Correlations between GABA and Glu concentrations in hMT+ and EVC

(primarily in V1) regions. **(a)** There is significant correlation between GABA and Glu

concentrations in hMT+ region. **(b)** The levels of GABA also significantly correlate with Glu in

EVC (primarily in V1) region. GABA and Glu concentrations (Conc.) are absolute, with units of

mmol per kg wet weight (Methods).

Figure supplement 5. Two linear correlations. **(a)** Significant positive correlation between

suppression index and BDT scores (took out two outlines). **(b)** Suppression index also

significantly correlates with GABA concentration in hMT+ region (without two outliers, having

the similar result shown in Figure 3d).

MT - frontal FC relates to SI and BDT

We next took the left hMT+ as the seed region and separately measured interregional FCs between the seed region and each voxel in the frontal regions (a priori search space). These measurements were correlated with performance in 3D visuospatial ability (BDT) to identify FCs with significant correlations. Results from connectivity-BDT analysis are summarized in Table 1 and shown in Figure 4a. We found that brain regions with FC strength to the seed region (left hMT+) significantly correlated with BDT scores were situated within the canonical cognitive cores of FPN (Brodmann areas (BAs) 6, 9, 10, 46, 47)(Assem et al., 2020; Deary et al., 2010; Duncan et al., 2020; Duncan et al., 2000; Gray et al., 2003; Jung & Haier, 2007). Across the whole brain search, the similar FCs (between hMT+ and frontal cognitive cores) still showed significant correlations with BDT scores (Table supplement 1) (also shown in figure supplement 6a). Additionally, we identified certain parietal regions (BAs 7, 39, 40) with significant correlations between their connectivity to the left hMT+ and the BDT scores (Table supplement 1) (also shown in figure supplement 6a). These significant connections between hMT+ and FPN system suggest that left hMT+ is involved in the efficient information integration network mediating the visuo spatial component of gF.

To address the question whether spatial suppression plays a role, we correlated hMT+ - based global FCs with SI. Though spatial suppression during motion perception (quantified by SI) is considered to be the function of area hMT+(Gautama & Van Hulle, 2001; Tadin et al., 2011), the top-down modulation from the frontal

cortex can increase surround suppression(Liu et al., 2016). Our functional

connectivity-SI analysis in the frontal regions (a priori search space) displayed 3 brain

regions in which FCs strength significantly correlated with SI: right BA4/6, left BA6,

and right BA46 (summarized in Table supplement 2, and shown in Figure 4b). Across

the whole brain search, we identified total 7 brain regions in which FCs strength

significantly correlated with SI, and 3 of these were in the frontal cortex. This is

consistent with the results obtained by the functional connectivity-SI analysis in a

priori search space (frontal cortex) (Table supplement 3 and figure supplement 6b).

We also did the V1 functional connectivity-BDT correlations as control analysis

(figure supplement 7). Only positive correlations in the frontal area were detected,

suggesting that in the 3D visuo-spatial intelligence task, V1 plays a role in

feedforward information processing. However, hMT+, which showed specific

negative correlations in the frontal, be suggested involving in the inhibition

mechanism. These results further emphasize the de-redundancy function of hMT+ in

3D visuo-spatial intelligence.

Table 1 FC of voxels showing significant correlation with BDT scores across subjects in frontal cortex.

FC number	Connected regions	BA	Size	Peak coordinate MNI (*x, y, z*)	*r*	*P*
1	Frontal_Sup_Orb_R	11	33	(12,63, -19.5)	-0.57	0.0011
2	Frontal_Inf_Orb_L	47	24	(-34.5,28.5, -13.5)	-0.63	0.0003
3	Frontal_Med_Orb_R	11	41	(3,43.5, -12)	-0.58	0.0009
4	Frontal_Inf_Orb_R	47	48	(-31.5,24, -12)	0.59	0.0008
5	Frontal_Inf_Orb_R	47	29	(25.5,30, -13.5)	0.67	0.0001
6	Insula_L	\	26	(-28.5,27,0)	0.67	0.0001
7	Frontal_Inf_Oper_R	45	41	(43.5,16.5,6)	0.64	0.0002
8	Frontal_Sup_R	10	25	(31.5,57,9)	0.59	0.0008
9	Frontal_Mid_L	10	82	(-33,48,12)	0.62	0.0003
10	Frontal_Inf_Oper_R	44	49	(51,7.5,21)	0.59	0.0007
11	Frontal_Inf_Oper_R	46	96	(49.5,16.5,28.5)	-0.62	0.0003
12	Frontal_Mid_L	10	32	(-31.5,49.5,24)	0.57	0.0012
13	Frontal_Mid_R	10	102	(31.5,36,30)	0.59	0.0009
14	Precentral_L	6	46	(-49.5, -1.5,34.5)	0.59	0.0007
15	Frontal_Mid_R	9	107	(51,19.5,40.5)	-0.67	0.0001
16	Frontal_Sup_L	9	35	(-9,60,37.5)	-0.69	0.0001
17	Frontal_Sup_Medial_R	9	74	(4.5,52.5,43.5)	-0.57	0.0011
18	Frontal_Sup_R	6	136	(28.5, -7.5,63)	-0.64	0.0002
19	Supp_Motor_Area_L	6	48	(-10.5,6,54)	0.63	0.0003
20	Frontal_Mid_L	6	119	(-24,4.5,55.5)	0.63	0.0003
21	Precentral_L	6	229	(-24, -18,66)	0.60	0.0005
22	Frontal_Sup_R	6	32	(16.5, -18,67.5)	0.68	0.0001
23	Precentral_R	6	80	(30, -24,70.5)	0.70	0.0001
24	Precentral_R	6	23	(16.5, -25.5,76.5)	0.70	0.0001

Single voxel threshold $P < 0.005$ (t > 3.057 or t < -3.057), adjacent size ≥ 23 voxels (AlphaSim corrected).

Table supplementary 1. FC of voxels showing significant correlation with BDT scores across subjects in whole brain.

FC number	Connected regions	BA	Size	Peak coordinate MNI (*x, y, z*)	*r*	*P*
1	Frontal_Med_OrbR	11	44	(2,43.5, -12)	-0.58	0.0009
2	Frontal_Inf_Oper_R	45	49	(43.5,16.5,6)	0.64	0.0002
3	Precentral_L	6	46	(-49.5, -1.5,34.5)	0.59	0.0007
4	Precentral_L	6	237	(-24, -18,66)	0.68	0.0001
5	Precentral_R	6	80	(31, -25,72)	0.67	0.0001
6	Frontal_Mid_L	10	82	(-33,48,12)	0.62	0.0003
7	Insula_L	47	124	(-33,15, -9)	0.64	0.0002
8	Insula_L	13	107	(-31.5,9,10.5)	0.63	0.0002
9	Insula_L	13	44	(-42, -10.5,7.5)	0.64	0.0002
10	Frontal_Inf_Oper_R	44	49	(51,7.5,21)	0.59	0.0007
11	Frontal_Inf_Oper_R	46	96	(49.5,16.5,28.5)	-0.62	0.0003
12	Frontal_Mid_R	10	102	(31.5,36,30)	0.59	0.0009
13	Paracentral_Lobule_L	6	46	(-15,21,51)	-0.64	0.0002
14	Supp_Motor_Area_L	6	48	(-10.5,6,54)	0.63	0.0003
15	Frontal_Mid_L	6	119	(-24,4.5,55.5)	-0.67	0.0001
16	Frontal_Sup_R	6	136	(29, -9,65)	0.57	0.0014
17	Frontal_Sup_MedialR	9	90	(8,51,43)	-0.59	0.0009
18	Frontal_Mid_R	9	108	(50,19,41)	-0.67	0.0001
19	Occipital_Mid_L	19	47	(-43, -83,9)	-0.64	0.0002
20	Occipital_Mid_L	37	45	(-40.5, -63,4.5)	-0.62	0.0004
21	Temporal_Mid_R	39	56	(45, -57,4.5)	-0.71	0.0000
22	Temporal_Mid_L	39	54	(-45, -48,12)	-0.55	0.0021
23	Temporal_Mid_L	40	102	(-48, -55.5,16.5)	-0.65	0.0001
24	Temporal_Mid_R	21	105	(64, -2, -19)	-0.61	0.0004
25	Temporal_Sup_R	22	228	(65, -42,11)	-0.6	0.0006
26	Precuneus_L	30	155	(1.5, -51,16.5)	-0.64	0.0002
27	Cingulum_Ant_R	32	41	(9,15,27)	0.68	0.0000
28	Cingulum_Mid_R	31	73	(15, -46.5,36)	-0.63	0.0002
29	Cingulum_Mid_L	23	80	(-3, -15,30)	0.63	0.0003
30	Lingual_R	18	46	(21, -93, -16)	0.58	0.0009
31	Parietal_Sup_L	7	54	(-19.5, -63,55.5)	0.66	0.0001
32	Parietal_Sup_L	7	48	(-23, -71,58)	0.63	0.0003
33	Postcentral_L	40	41	(-31.5, -39,58.5)	0.68	0.0001
34	Postcentral_L	40	40	(-48, -36,58)	0.71	0.0000
35	Thalamus_L	Wm	108	(-24, -31.5,12)	0.64	0.0002
36	Parietal_Sup_L	5	37	(-37, -48,63)	0.59	0.0008
37	Vermis_10	-	37	(-3, -43, -37)	-0.68	0.0001
38	-	-	40	(12, -19.5, -42)	-0.68	0.0000

Single voxel threshold *P* < 0.01 (t > 2.771 or t < -2.771), adjacent size ≥ 37 voxels (AlphaSim corrected).

273 **Table supplement 2. FCs of voxels showing significant correlation with SI across subjects in**

FC number	Connected regions	BA	Size	Peak coordinate	r	P
				MNI (x, y, z)		
1	Frontal_Inf_Oper_R	46	80	(48,15,28.5)	-0.65	0.0001
2	Precentral_R	4/6	106	(33, -25.5,63)	0.72	0.0000
3	Precentral_L	6	26	(-30, -24,72)	0.66	0.0001

274 **frontal cortex.**

275 Single voxel threshold $P < 0.005$ (t > 3.057 or t < -3.057), adjacent size $\geq$ 22 voxels (AlphaSim

276 corrected).

278 **Table supplement 3. FCs of voxels showing significant correlation with SI across subjects in**
279 **whole Brain.**

FC number	Connected regions	BA	Size	Peak coordinate	r	P
				MNI (*x, y, z*)		
1	Cerebelum_Crus1_L	18	70	(-29, -85, -24)	0.51	0.0052
2	Cerebelum_6_R	18	39	(12, -85, -17)	0.65	0.0002
3	Calcarine_R	18	70	(15, -91.5,12)	0.63	0.0003
4	Frontal_Inf_Oper_R	46	127	(48,15,28.5)	-0.65	0.0001
5	Precentral_R	4/6	179	(32, -24,70)	0.71	0.0001
6	Precentral_L	6	59	(-30, -23,72)	0.66	0.0001
7	-	-	37	(26, -37, -11)	-0.67	0.0001

281 Single voxel threshold $P < 0.01$ (t > 2.771 or t < -2.771), adjacent size $\geq$ 37 voxels (AlphaSim
282 corrected).

Figure 4. Significant FCs from connectivity-behavior analyses in *a priori* search space. The seed region is the left hMT+. The significant FCs are obtained from *a priori* space (frontal cortex). (**a**) The significant FCs obtained from connectivity-BDT analysis. Single voxel threshold $P < 0.005$, adjacent size ≥ 23 (*AlphaSim* correcting, Methods). (**b**) The significant FCs obtained from connectivity-SI analysis. Single voxel threshold $P < 0.005$, adjacent size ≥ 22 (*AlphaSim* correcting, Methods). Positive correlations are shown in warm colors, while negative correlations are shown in cold colors. The paired deep blue, light blue, red circles on (**a**) and (**b**) indicate the overlap regions in left BA6, right BA6, right BA46(DLPFC) between Connectivity - BDT analysis and Connectivity – SI analysis, respectively.

Figure supplement 6. Significant FCs searched from connectivity-behavior (BDT/SI) analyses in

the whole brain. The seed region is the left hMT+. The significant FCs are obtained from the

entire brain search, single voxel threshold $P < 0.01$, adjacent size ≥ 37 voxels (AlphaSim

correcting, Methods). Positive correlations are shown in warm colors, while, negative correlations

are shown in cold colors. **(a)** the significant FCs obtained from connectivity-BDT analysis. **(b)** the

significant FCs obtained from connectivity-SI analysis.

Figure supplement 7. Significant FCs searched from connectivity-behavior (BDT) analyses in the

frontal **(a)** and whole brain **(b)**. The seed region is the left V1. **(a)** The significant FCs are obtained

from the frontal search, single voxel threshold $P < 0.005$, adjacent size ≥ 23 (AlphaSim correcting,

same methods for hMT+). **(b)** The significant FCs are obtained from the entire brain search, single

voxel threshold $P < 0.01$, adjacent size ≥ 37 voxels (AlphaSim correcting, same methods for

hMT+). Only positive correlations were detected, shown in warm colors.

323 **Table 2. Correlations between FC in Table 1 and GABA/Glu concentrations in hMT+.**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	-0.07	0.72	0.75	-0.11	0.58	0.85
2	-0.28	0.14	0.36	-0.27	0.15	0.81
3	-0.13	0.52	0.59	-0.07	0.71	0.85
4	0.10	0.59	0.64	0.12	0.54	0.85
5	0.14	0.48	0.58	0.24	0.21	0.81
6	0.31	0.11	0.33	0.15	0.43	0.85
7	0.20	0.30	0.48	0.07	0.74	0.85
8	0.14	0.45	0.58	0.05	0.79	0.85
9	0.20	0.30	0.48	0.10	0.60	0.85
10	-0.13	0.49	0.58	-0.16	0.41	0.85
11	**-0.56**	**0.0018**	**0.02***	-0.22	0.25	0.81
12	0.18	0.34	0.51	0.15	0.43	0.85
13	0.20	0.30	0.48	0.05	0.81	0.85
14	0.39	0.04	0.14	0.22	0.24	0.81
15	-0.40	0.03	0.12	-0.21	0.27	0.81
16	-0.27	0.15	0.36	-0.12	0.53	0.85
17	0.17	0.37	0.52	0.06	0.74	0.85
18	0.26	0.18	0.39	0.16	0.40	0.85
19	0.39	0.03	0.12	0.31	0.10	0.81
20	0.01	0.98	0.98	0.14	0.46	0.85
21	0.40	0.03	0.12	0.24	0.21	0.81
22	0.22	0.25	0.48	0.06	0.76	0.85
23	**0.69**	**0.0001**	**0.002****	0.47	0.01	0.24
24	0.41	0.03	0.12	0.001	0.97	0.97

324 Bold font indicates the significant correlations survived from multi correlation correction.

326 **Table supplement 4. Correlations between FC in Table supplement 1 and GABA/Glu**

327 **concentrations in hMT+**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	-0.13	0.49	0.55	-0.08	0.68	0.84
2	0.17	0.38	0.49	0.05	0.82	0.88
3	0.39	0.04	0.22	0.22	0.24	0.69
4	0.41	0.03	0.19	0.24	0.21	0.69
5	**0.69**	**0.0001**	**0.0038**[**]	0.47	0.01	0.38
6	0.2	0.3	0.41	0.1	0.6	0.81
7	0.26	0.17	0.34	0.21	0.28	0.69
8	0.22	0.24	0.35	0.25	0.19	0.69
9	0.23	0.23	0.35	0.02	0.9	0.92
10	-0.13	0.49	0.55	-0.16	0.41	0.70
11	**-0.56**	**0.0018**	**0.03**[*]	-0.22	0.25	0.69
12	0.2	0.3	0.41	0.05	0.81	0.88
13	-0.35	0.06	0.25	0.07	0.73	0.84
14	0.39	0.03	0.19	0.31	0.1	0.69
15	0.006	0.98	0.98	0.14	0.46	0.70
16	0.26	0.18	0.34	0.16	0.4	0.70
17	0.15	0.42	0.51	0.04	0.83	0.88
18	-0.4	0.03	0.19	-0.21	0.27	0.69
19	-0.37	0.05	0.24	-0.15	0.44	0.70
20	-0.16	0.39	0.49	-0.19	0.33	0.70
21	-0.31	0.1	0.28	-0.29	0.13	0.69
22	-0.22	0.24	0.35	-0.07	0.72	0.84
23	-0.1	0.6	0.65	-0.12	0.53	0.77
24	-0.09	0.64	0.68	-0.09	0.63	0.83
25	-0.3	0.11	0.28	-0.14	0.46	0.70
26	-0.08	0.66	0.68	-0.08	0.69	0.84
27	0.28	0.13	0.29	0.1	0.59	0.81
28	-0.27	0.16	0.34	-0.2	0.29	0.69
29	0.33	0.08	0.28	0.25	0.19	0.69
30	0.24	0.21	0.35	0.19	0.32	0.70
31	0.24	0.22	0.35	0.36	0.05	0.69
32	0.23	0.24	0.35	0.23	0.23	0.69
33	0.44	0.02	0.19	0.18	0.35	0.70
34	0.32	0.09	0.28	0.002	0.99	0.99
35	0.32	0.09	0.28	0.31	0.1	0.69
36	0.30	0.11	0.28	0.22	0.25	0.69
37	-0.29	0.13	0.29	-0.23	0.23	0.69
38	-0.13	0.48	0.55	-0,15	0.42	0.70

328 Bold font indicates the significant correlations survived from multi correlation correction.

329 23

330 **Table supplement 5. Correlations between FCs in Table supplement 2 and GABA/Glu**

331 **concentrations in hMT+.**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	**-0.56**	**0.0017**	**0.0017****	-0.21	0.27	0.27
2	**0.70**	**0.0001**	**0.0003*****	**0.49**	**0.007**	**0.021***
3	**0.65**	**0.0001**	**0.0002****	**0.40**	**0.03**	**0.045***

332 *: $P_{FDR} < 0.05$; **: $P_{FDR} < 0.01$; ***: $P_{FDR} < 0.001$; Bold font indicates the significant correlations

333 survived from multi correlation correction.

335 **Table supplement 6. Correlations between FCs in Table supplement 3 and GABA/Glu**

336 **concentrations in hMT+.**

FC number	hMT+ GABA concentrations			hMT+ Glu concentrations		
	r	*P*	*FDR*	*r*	*P*	*FDR*
1	**0.37**	**0.049**	**0.049**[*]	0.32	0.09	>0.05
2	**0.37**	**0.049**	**0.049***	0.41	0.025	>0.05
3	**0.48**	**0.008**	**0.01**[*]	0.21	0.28	>0.05
4	**-0.58**	**0.001**	**0.002**[**]	-0.22	0.26	>0.05
5	**0.69**	**0.0001**	**0.0007**	0.46	0.01	>0.05
6	**0.66**	**0.0001**	**0.0004**[***]	0,39	0.04	>0.05
7	**-0.48**	**0.0077**	**0.01***	-0.35	0.067	>0.05

337 *: $P_{FDR} < 0.05$; **: $P_{FDR} < 0.01$; ***: $P_{FDR} < 0.001$; Bold font indicates the significant correlations

338 survived from multi correlation correction.

Local hMT+ GABA acts on SI and BDT via global hMT-frontal connectivity

To determine whether local neurotransmitter levels (such as GABA and Glu) in the hMT+ region mediate the broader 3D visuo-spatial ability of BDT, which as a component of gF, is linked to the frontal cortex(Fangmeier et al., 2006), we correlated the significant FCs of hMT – frontal in Figure 4a (also shown in Table 1) with the GABA and Glu levels in hMT+ region. The results revealed that only two FCs significantly correlated with inhibitory GABA levels in hMT+: 1) the FC of left hMT+ - right BA 46 (significantly negative correlation, $r = - 0.56$, $P = 0.02$, $n = 29$, FDR correction, Figure 5a left); 2) the FC of left hMT+ - right BA 6 (significantly positive correlation, $r = 0.69$, $P = 0.002$, $n = 29$, FDR correction, Figure 5b left) (also shown in Table 2). There were no significant correlations between these FCs and the excitatory Glu levels in hMT+ (Table 2). Across the whole brain search, we obtained the same two hMT+ - frontal FCs significantly correlating with both hMT+ GABA levels and BDT (Table supplement 4), this is consistent with the results in the priori search space (frontal cortex) (Table 2). We then correlated the significant FCs in Figure 5b (also in Table supplement 2) with GABA and Glu concentrations in hMT+ and found that almost all the correlations are significant except one (between the FC of left hMT+ - right BA46 and the Glu levels in hMT+) (Table supplement 5). Among the three FCs, the clusters of two FCs have substantial voxel overlap with the FCs we found by the connectivity-BDT analysis (Figure 5a, b). Across the whole brain search, there were total 7 brain regions in which FCs strength were significantly correlated with SI, all the 7 FCs significantly correlated the hMT+ GABA levels, while, no FC

had significant correlation with the hMT+ Glu levels (Table supplement 6).

Taken together, our results displayed that the overlap FCs from the analyses of

connectivity - behavior (BDT and SI) -GABA are the hMT+ - BA 46 and hMT+ - BA

6 (Figure 5a, b). These results suggest that the FCs of hMT+ - frontal regions (BA 46

and BA 6) coupling with local hMT+ GABA provides the neural basis for both the

simple motion perception (quantified by SI) and the complex 3D visuo-spatial ability

(quantified by BDT).

In order to fully investigate the potential roles of the multiples variables

contributing to BDT scores, serial mediation analyses(Hayes, 2013) were applied to

both the MR and behavioral data. Following our hypothesis, the independent variable

(X) is hMT+ GABA, the dependent variable (Y) is BDT scores, the covariate is the

age, and the mediators are FC (M1) and SI (M2). We used the overlap clusters from

the analyses of connectivity-BDT-GABA and connectivity-SI-GABA to compute the

FC of hMT+ - BA46, 1 participant was excluded due to his age greater than mean +

2.5SD. The serial mediation model is shown in Figure 5c. GABA levels in hMT+

significantly negatively correlated with the FC of hMT+ - BA46 (β = - 0.32, P =

0.0009), which in turn significantly negatively correlated with SI (β = - 0.19, P =

0.035), and consequently, significantly positively correlated with BDT (β = 38.5, P =

0.009). Critically, bootstrapped analyses revealed that our hypothesized indirect effect

(i.e., hMT+ GABA $\rightarrow$ FC of hMT+ - BA46 $\rightarrow$ SI $\rightarrow$ BDT) was significant (β = 2.28,

SE = 1.54, 95% CI = [0.03, 5.94]). The model accounted for 34% of the variance in

BDT. However, when considering the hMT-BA6 FC as the mediator M1, the serial

model does not show a significant indirect effect. Consequently, we explored a

mediation model, which revealed that the hMT-BA6 FC totally mediates the

relationship between GABA and BDS. (Figure 5d). For sensitivity purposes, we tested

the alternative models, in which the order of the mediators was reversed. The pathway

that hMT+ GABA was predicted to be associated with SI, followed by the FC of

hMT+ - BA 46, and then BDT, did not yield the chained mediation effects on BDT

(figure supplement 8).

Figure 5. Local hMT+ GABA acts on SI and BDT via global hMT-frontal connectivity. (a) Significant negative correlation between the FC of left hMT+ - right DLPFC (BA46) and hMT+ GABA (*FDR* correction). (b) Significant positive correlation between the FC of left hMT+ - right (pre) motor cortex (BA4/6) and hMT+ GABA (*FDR* correction). In (a) and (b), left: the significant FCs obtained from connectivity-BDT analysis; right: the significant FCs obtained from connectivity-SI analysis. (c) Significant pathways: hMT+ GABA→ FC (left hMT+ - right BA46, negative correlation) → SI (negative correlation) → BDT (positive correlation). This pathway can explain 34% of the variance in BDT. (d) Significant pathways: hMT+ GABA → FC (left hMT+ - right BA6, positive correlation) → BDT (positive correlation). The bolded lines represent the hypothesized mediation effect. The dotted lines represent alternative pathways. *: $P < 0.05$; **: $P <$

0.01; ***: $P < 0.001$.

Figure supplement 8. The alternative serial mediation models from local hMT+ GABA to global

performance of BDT. The pathway that hMT+ GABA was predicted to be associated with SI,

followed by the FC of hMT+ - BA 46, and then BDT, did not yield the chained mediation effects

on BDT. *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$.

To summarize (shown in Figure 6), the results from the serial mediation analyses

are consistent with our hypothesis. That is, higher GABAergic inhibition in hMT+

relates to stronger negative FC between hMT+ and BA46, leading to enhanced ability

for surround suppression (filtering out irrelevant information(Tadin, 2015), which

ultimately resulting in more efficient visual 3D processing as key component of gF

(the higher BDT scores).

Figure 6. Sketch depicting the multi-level inhibitory mechanisms centered on hMT+ GABA contributing to visuo-spatial intelligence. Inhibitory GABA in hMT+ (a sensory cortex, shown in green circle), coupling with the functional connectivity between hMT+ and BA46 (cognitive control core, shown in purple circle), and mediated by motion surround suppression (shown in red circle), contributes to visuo-spatial intelligence (BDT, 3D domain, shown in red and white building blocks). In this sketch, the two-colored parallel arrows show the negative FC between hMT+ and BA46, the colored arrows below the green circle display the inhibition mechanisms centered on hMT+ GABA (2), filtered the irrelevant information in (1) and focused on the efficient visuo-spatial information (3). Black long arrows display the direction of information flow: from input information (1) to visuo-spatial intelligence (4).

Figure Supplement 9. Correlation matrix shows correlations between all measured in our

biochemistry and behaviour level. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$.

Discussion

Here, we provide evidence that hMT+ inhibitory mechanisms mediate processing in the visuo-spatial component (3D domain) of gF on multiple levels, that is, from molecular over brain connectivity to behavior. First, this study found that higher hMT+ inhibitory GABA levels (but not excitatory Glu) relate to FC between hMT+ and BA 46 that contribute to both SI and BDT. Our serial mediation analyses indicate that the inhibitory mechanisms related to hMT+ and its GABA levels in hMT+ (but not Glu), FCs of hMT+ - BA46 coupling with hMT+ inhibitory GABA (but not excitatory Glu), and behavior (SI indexing perceptual suppression in hMT+) predict the inter-subject variance in the 3D gF task (BDT) (Figure 5c). Second, we demonstrate discrete GABAergic inhibition mechanisms in hMT+ that mediate the strong FCs between hMT+ - frontal regions (BA46 and BA6): significant negative correlation with the FC of hMT+ - BA 46 (Figure 5a), whereas there is significant positive correlation with the FC of hMT+ - BA 6 (Figure 5b). This indicates that different frontal regions, DLPFC (BA 46) and premotor cortex (BA6), contribute uniquely to gF through hMT+ - based inhibitory mechanisms.

 The goal of our research is to reveal that the inhibitory (not excitatory) mechanism in hMT+ contributes to multi-level processing in 3D visuo-spatial ability (BDT). Monkey electrophysiological experiments revealed that selective attention gates the visual cortex, including area MT, effectively suppressing the irrelevant information(Everling et al., 2002; Treue & Maunsell, 1996). These findings align with the "neural efficiency" hypothesis of intelligence(Haier et al., 1988), which puts

forward the human brain's ability to suppress the repetition of information. Neural

suppression is associated with the balance between excitation and inhibition (EIB),

usually represented by covariation between Glutamate and GABA(Ozeki et al., 2009).

Here, this study exploited the high spectral resolution afforded by ultrahigh field (7T)

MRS to reliably resolve GABA measurement, to adequately discriminate the

glutamate and glutamine signals, and to resolve the high accuracy Glu

measurement(Ende, 2015).

This work implemented the MRS scanning in hMT+ (3D visual domain) and EVC

(primarily in V1) (2D visual domain) regions and found that hMT+ inhibitory GABA

(but not excitatory Glu) significantly correlated with BDT, i.e., the higher GABA

levels in hMT+ (rather than excitatory Glu) relate to higher visual 3D processing

(BDT) (Figure 3c). Basically, this study contains the data of SI, BDT, GABA in MT+

and EVC (primarily in V1), Glu in MT+ and EVC (primarily in V1)-all 6

measurements. we made a correlation matrix to reporting all values in figure

supplement 9.

We searched the global hMT+ - based FCs with the connectivity-BDT analyses (in

priori search space and whole brain search to valid), and then, correlated these

significant FCs with the GABA and Glu concentrations in hMT+. We found two FCs

(hMT+ - BA46 and hMT+ - BA6) significantly correlating with hMT+ inhibitory

GABA (whereas no FC significantly correlated with hMT+ excitatory Glu).

Accordingly, our results emphasize the importance of hMT+ inhibitory GABA (but

not excitatory Glu) in processing the 3D visual-spatial intelligence (BDT).

Our recent human study(Song et al., 2021) and other study's animal

experiments(Ozeki et al., 2009; Sato et al., 2016) demonstrated that the conjoint

action of inhibition (GABA) and excitation (Glu) underlies visual spatial suppression.

In this work, our novel data show the chained mediation effects from local hMT+

GABA to more global BDT: hMT+ GABA → FC (hMT+ and BA46) → SI → BDT.

Thereby, our data indicate that inhibitory mechanisms in hMT+, from the biochemical

level of GABA over FC to the behavioral level, can predict the inter-subject variance

in the 3D gF task (BDT) (Figure 5c).

Another interesting finding reveal that GABAergic inhibition in hMT+ coupling

with distinct functional connectivity patterns between BA 46-hMT+ and BA6-hMT+.

A previous human fMRI experiment found that the positive and negative correlations

between BDT and the activation of frontal regions appeared at different reasoning

phases (validation or integration phases during reasoning)(Fangmeier et al., 2006). On

the one hand, a monkey electrophysiological experiment reported the delayed

modulation from PFC (especially in DLPFC (BA 46))) to area MT during a visual

motion task(Zaksas & Pasternak, 2006). Computational models converged with

empirical data of awake monkey experiments slowing temporal modulation from PFC

to MT/MST(Donner et al., 2009; Siegel et al., 2015; Wang, 2002; Wimmer et al.,

2015). On the other hand, human MEG studies(Donner et al., 2009; Wilming et al.,

2020) reported that the gamma-band activity in the visual cortex (including area MT)

exhibited high coherence with the activity in (pre-) motor regions (BA 4/6). These

results suggest that the relation of long-range FC and local inhibitory mechanism

(hMT+ GABA) support our findings that inhibition in hMT+ contributes to efficient

long-range integration and coordination in distant brain areas like the prefrontal and

premotor cortex.

How does hMT+ assemble into the cognitive system as an intellectual hub rather

than a simple input module? The results in Figure 5a, b showed that the overlap brain

regions from the analyses of connectivity-BDT-GABA/connectivity-SI-GABA are the

hMT+ - BA 46. This overlap couples with local visual suppression (SI) and

consequently plays an important role in intelligence (BDT). The direction

discrimination task in this work (the visual motion paradigm of center-surround

antagonism) was previously considered a mainly local function of hMT+(Melnick et

al., 2013; Tadin, 2015; Tadin et al., 2003). However, our results with connectivity-SI

analyses revealed that both local (FC within BA 18) and global brain connectivity (FC

between hMT+ and frontal regions) contribute to SI (Table supplement 3). In human

psychophysical experiments(Melnick et al., 2013; Tadin et al., 2003) the brief

stimulus duration (~100 ms) in motion discrimination precludes most top-down

attentional effects(Wang, 2002; Zaksas & Pasternak, 2006), while, attention, which

predicted the performance of the motion discrimination task, was sustained

throughout the stimulus intervals(Siegel et al., 2015). Furthermore, animal

experiments have revealed that the local circuits in the visual cortex combining with

top-down modulation and intracortical horizontal connection mediate the visual-

spatial suppression(Angelucci et al., 2002; Keller et al., 2020; Li et al., 2019; Zhang et

al., 2014).

Our results (shown in Figure 5a, b, right) present the intrinsic binding of local

GABAergic inhibition in hMT+, which suppressing redundancy of visual motion

processing (SI), and the activity of brain connectivity between hMT+ and frontal

regions. These individually inherent traits may contribute to the individual difference

in 3D visuo-spatial ability (Figure 5a, b, left). A candidate divisive normalization

model(Carandini & Heeger, 2011; Reynolds & Heeger, 2009) can explain how such

reverberation affects the process of suppressing the irrelevant information, from

perception to intelligence(Melnick et al., 2013; Tadin, 2015). We summarize a

framework (Figure 6) to indicate and visualize our findings.

Recently, Duncan et al. demonstrated coding of general fluid intelligence (gF) in

distributed regions, defining them as part of multi-demand (MD) systems (Assem et

al., 2020; Duncan et al., 2020). The MD system encompasses a range of cognitive

domains, including working memory, mathematics, language, and relational reasoning.

According to Melnick et al. (2013), motion surround suppression (SI) and time

thresholds for small and large gratings, which reflect hMT+ functionality, are

correlated with Verbal Comprehension, Perceptual Reasoning, Working Memory, and

Processing Speed indicators. Additionally, Fedorenko et al. identified multi-demand

activation regions around the occipito-temporal areas, potentially overlapping with

hMT+ (Fedorenko et al., 2013). As a key region in the representation of sensory flows

(including optic, and auditory flows)(Fetsch et al., 2011; Gu et al., 2006), hMT+

shows potential to be central to the MD system. Future research could focus on multi-

task paradigms to further investigate the mechanisms of hMT+ and its relationship with broader cognitive functions.

Together, this study offers a comprehensive insight into how the information exchange and integration between the sensory cortex (hMT+) and cognition core of BA 46, coupling with the hMT+ GABA, can predict the performance of 3D visuo-spatial ability (BDT). Our results provide direct evidence that a sensory cortex area (hMT+), its GABA biochemistry, functional connectivity, and cognition behavior levels, can assemble into complex cognition as an intellectual hub.

Materials and Methods

Subjects

Thirty-six healthy subjects (18 female, mean age: 23.6 years±2.1, range: 20 to 29 years) participated in this study, they were recruited from Zhejiang University. All subjects had normal or corrected-to-normal vision. In addition, they reported no psychotropic medication use, no illicit drug use within the past month, no alcohol use within 3 days prior to scanning, and right-handed. This experiment was approved by the Ethics Review Committee of Zhejiang University and conducted in accordance with the Helsinki Declaration. All participants signed informed consent forms prior to the start of the study and were compensated for their time. All subjects participated in the motion spatial suppression psychophysical, resting-state fMRI and MRS (hMT+ and EVC (primarily in V1) regions, in random sequence) experiments, but only part

of the MRS data (31/36 in hMT+ region and 28/36 in EVC (primarily in V1) region)

survived quality control (see the part of MRS date processing). The sample size is

determined by the statistic requirement (30 sample for Person correlation statistical

analysis).

Motion surrounding suppression measurement

All stimuli were generated using Matlab (MathWorks, Natick, MA) with

Psychophysics Toolbox(Brainard, 1997), and were shown on a linearized monitor

(1920×1080 resolution, 100-Hz refresh rate, Cambridge Research System, Kent, UK).

The viewing distance was 72 cm from the screen, with the head stabilized by a

chinrest. Stimuli were drawn against a gray (56 cd per m^{-2}) background.

A schematic of the stimuli and trial sequences is shown in our recent study(Song et

al., 2021). The stimulus was a vertical drifting sinusoidal grating (contrast, 50%;

spatial frequency, 1 cycle/^{o}; speed, 4°/s) of either small (diameter of 2°) or large

(diameter of 10°) size. The edge of the grating was blurred with a raised cosine

function (width, 0.3°). A cross was presented in the center of the screen at the

beginning of each trial for 500ms, and participants were instructed to fixate at the

cross and to keep fixating at the cross throughout the trial. In each trial, a grating of

either large or small size was randomly presented at the center of the screen. The

grating drifted either leftward or rightward, and participants were asked to judge the

perceived moving direction by a key press. Response time was not limited. The

grating was ramped on and off with a Gaussian temporal envelope, and the grating

duration was defined as 1 SD of the Gaussian function. The duration was adaptively

adjusted in each trial, and duration thresholds were estimated by a staircase procedure.

Thresholds for large and small gratings were obtained from a 160-trial block that

contained four interleaved 3-down/1-up staircases. For each participant, we computed

the correct rate for different stimulus durations separately for each stimulus size.

These values were then fitted to a cumulative Gaussian function, and the duration

threshold corresponding to the 75% correct point on the psychometric function was

estimated for each stimulus size.

Stimulus demonstration and practice trials were presented before the first run.

Auditory feedback was provided for each wrong response. To quantify the spatial

suppression strength, we calculated the spatial suppression index (SI), defined as the

difference of log10 thresholds for large versus small stimuli(Schallmo et al., 2018;

Tadin et al., 2003):

599
$$SI = log_{10}(\ large\ threshold) - log_{10}(\ small\ threshold) \qquad (1)$$

Block design task measurement

The block design task was administered in accordance with the WAIS-IV

manual(Wechsler, 2008). Specifically, participants were asked to rebuild the figural

pattern within a specified time limit using a set of red and white blocks. The time

limits were set as 30 s to 120 s according to different levels of difficulty. The patters

were presented in ascending order of difficulty, and the test stopped if two

consecutive patterns were not constructed in the allotted time. The score was

determined by the accomplishment of the pattern and the time taken. A time bonus

was awarded for rapid performance in the last six patterns. The score ranges between

0 and 66 points, with higher scores indicating better perceptual reasoning.

MR experimental procedure

MR experiments were performed in a 7T whole body MR system (Siemens

Healthcare, Erlangen, Germany) with a Nova Medical 32 channel array head coil.

Sessions included resting-state functional MRI, fMRI localizer scan, structural image

scanning, and MRS scan. Resting-state scans were acquired with 1.5-mm isotropic

resolution (transverse orientation, TR/TE = 2000/20.6 ms, 160 volumes, slice number

= 90, flip angle = 70^{o}, eyes closed). Structural images were acquired using a

MP2RAGE sequence (TR/ TI1/ TI2 = 5000/901/3200 ms) with 0.7-mm isotropic

resolution. MRS data were collected within two regions (hMT+ and EVC (primarily

in V1)) for each subject, and we divided them into two sessions to avoid discomfort

caused by long scanning. The order of MRS VOIs (hMT+ and EVC (primarily in V1))

in the two sessions was counterbalanced across participants. Interval between two

sessions was used for block design and motion discrimination tasks. One session

included fMRI localizer scan, structural image scanning, and MRS scan for the hMT+

region; the other session included structural image scan, and MRS scan for the EVC

(primarily in V1) region. Spectroscopy data were acquired using a [1]H-MRS single-

voxel short-TE STEAM (Stimulated Echo Acquisition Mode) sequence(Frahm et al.,

1989) (TE/TM/TR = 6/32/7100ms) with 4096 sampling points, 4-kHz bandwidth, 16

averages, 8 repetitions, 20×20×20 mm^3 VOI size, and VAPOR (variable power and

optimized relaxation delays) water suppression(Tkáč et al., 1999). Prior to acquisition,

first- and second-order shims were adjusted using FASTMAP (fast, automatic

shimming technique by mapping along projections(Gruetter, 1993). Two non-

suppressed water spectra were also acquired: one for phase and eddy current

correction (only RF pulse, 4 averages) and another for metabolite quantification

(VAPOR none, 4 averages). Voxels were positioned based on anatomical landmarks

using a structural image scan collected in the same session, while avoiding

contamination by CSF, bone, and fat. The hMT+ VOIs were placed in the

ventrolateral occipital lobe, which was based on anatomical landmarks(Dumoulin et

al., 2000; Schallmo et al., 2018). We did not distinguish between the middle temporal

(MT) and medial superior temporal (MST) areas in these hMT+ VOIs(Huk et al.,

2002). For 14 subjects, we also functionally identified hMT+ as a check on the

placement of the VOI. A protocol was used with a drifting grating (15% contrast)

alternated with a static grating across blocks (10 s block duration, 160 TRs total).

Using fMRI BOLD signals, these localizer data were processed online to identify the

hMT+ voxels in the lateral occipital cortex, which responded more strongly to moving

vs. static gratings. In addition, we only used the left hMT+ as the target region to scan,

which was motivated by studies showing that left hMT+ was more effective at

causing perceptual effects(Tadin et al., 2011). For EVC (primarily in V1) region, the

VOI was positioned on each subject's calcarine sulcus on the left side(Tadin et al.,

2011) based on anatomical landmarks(Boucard et al., 2007; Dumoulin et al., 2000)

MRS data processing

Spectroscopy data were preprocessed and quantified using magnetic resonance signal

processing and analysis, https://www.cmrr.umn.edu/ downloads/mrspa/), which runs

under MATLAB and invokes the interface of the LCModel (Version 6.3-1L) (Chen et

al., 2019). First, we used the non-suppressed water spectra to perform eddy current

correction and frequency/phase correction. Second, we checked the quality of each

FID (16 averages) visually and removed those with obviously poor quality. Third, the

absolute concentrations of each metabolite were quantitatively estimated via the

Water-Scaling method. For partial-volume correction, the tissue water content was

computed as follows(Ernst et al., 1993):

663   *Tissue water content=fgm*0.78 + fwm*0.65 + fcsf*0.97*   (2)

where *fgm*, *fwm*, and *fcsf* were the GM/WM/CSF volume fraction in MRS VOI and

we used FAST (fMRI's automated segmentation tool, part of the FSL toolbox)(Zhang

et al., 2001) to segment the three tissue compartments from the T1-weighted structural

brain images. For water T2 correction, we set water T2 as 47ms (Marjańska et al.,

2012). Our concentrations were mM per kg wet weight. Furthermore, LCModel

analysis was performed on all spectra within the chemical shift range of 0.2 to 4.0

ppm.

Poor spectral quality was established by a Cramer-Rao Lower Bound (CRLB) of

more than 20%(Cavassila et al., 2001), and some data were excluded from further

analysis. The details were described in our recently paper(Song et al., 2021).

Rs-fMRI data processing and analysis

Resting-state functional image was analyzed in the Data Processing and Analysis for

Brain Imaging DPABI toolbox(Yan et al., 2016) based on SPM 12

(http://www.fil.ion.ucl.ac.uk/spm/). The preprocessing steps included discard of the

first five volumes, slice timing, realignment to the 90th slice, coregistration of each

subject's T1-weighted anatomical and functional images, segmentation of the

anatomical images into six types of tissues using DARTEL, linear detrend, regressing

nuisance variables (including realignment Friston 24-parameter, global signal, white

matter and CSF signal) (Friston et al., 1996), normalization to the standard Montreal

Neurological Institute (MNI) space with the voxel size of $1.5\times 1.5\times 1.5$ mm^3 using

DARTEL, spatial smoothing with a Gaussian kernel of 3 mm full-width-half-

maximum (FWHM), and band-pass filtering with Standard frequency band (SFB,

0.01–0.1 Hz). Spherical ROI with a radius of 6mm was placed in left MT. The

coordinate for left MT (-46, -72, -4, in MNI space) was obtained by our localizer

fMRI experiment. We calculated the seed-to-voxel whole brain FC map for each

subject. All the FC values were Fisher-Z-transformed.

We did a similar connectivity-behavior analysis to a previous study (Song et al.,

2008). First, we computed the Pearson's correlation coefficient between BDT scores

and the FC values across subjects in a voxel-based way. Then, to evaluate the

significance, we transformed the *r*-value into *t*-value ($t = \frac{\sqrt{df} \cdot r}{\sqrt{1-r^2}}$), where *df* denotes to

the degrees of freedom, and *r* is the Pearson's correlation coefficient between BDT

scores and the FC values. Here, *df* was equal to 27. The brain regions in which the FC

values to the seed region was significantly correlated with the BDT scores were

obtained with a threshold of *P* < 0.005 for regions of *a priori* ($|t_{(27)}| \geq 3.057$, and

adjacent cluster size ≥ 23 voxels (AlphaSim corrected)), and *P* < 0.01 for whole-brain

analyses ($|t_{(27)}| \geq 2.771$ and adjacent cluster size ≥ 37 voxels (AlphaSim corrected).

Statistical Analysis

PROCESS version 3.4, a toolbox in SPSS, was used to examine the mediation model.

There are some prerequisites for mediation analysis: the independent variable should

be a significant predictor of the mediator, and the mediator should be a significant

predictor of the dependent variable.

SPSS 20 (IBM, USA) was used to conduct all the remaining statistical analysis in

the study. We evaluated the correlation of variables (GABA, Glu, SI, BDT) using

Pearson's correlation analysis. Differences or correlations were considered

statistically significant if *P* < 0.05. Significances with multiple comparisons were

tested with false discovery rate (*FDR*) correction. The effect of age on intelligence

was controlled for by using partial correlation in the correlation analysis and was

taken as a covariate in the serial mediation model analysis.

Data availability

716   Source data are provided with this paper and have been archived at zenodo and could

717   be downloaded with reasonable request：

718   https://doi.org/10.5281/zenodo.12789781

Code availability

721   This code has been uploaded to the GitHub：

722   https://github.com/Yrehearsal/GABA-_hMT-_Intelligence

References

Angelucci, A., Levitt, J. B., Walton, E. J., Hupe, J.-M., Bullier, J., & Lund, J. S. (2002). Circuits for local and
global signal integration in primary visual cortex. *Journal of Neuroscience*, *22*(19), 8633-8646.
Assem, M., Glasser, M. F., Van Essen, D. C., & Duncan, J. (2020). A domain-general cognitive core
defined in multimodally parcellated human cortex. *Cerebral Cortex, 30*(8), 4361-4380.
Barbey, A. K. (2018). Network neuroscience theory of human intelligence. *Trends in cognitive sciences*,
22(1), 8-20.
Boucard, C. C., Hoogduin, J. M., van der Grond, J., & Cornelissen, F. W. (2007). Occipital proton
magnetic resonance spectroscopy (1H-MRS) reveals normal metabolite concentrations in
retinal visual field defects. *PLoS One, 2*(2), e222.
Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis, 10*(4), 433-436.
Carandini, M., & Heeger, D. J. (2011). Normalization as a canonical neural computation. *Nat Rev
Neurosci, 13*(1), 51-62. https://doi.org/10.1038/nrn3136
Cattell, & Raymond, B. (1963). Theory of fluid and crystallized intelligence: A critical experiment.
Journal of Educational Psychology, 54(1), 1-22.
Cavassila, S., Deval, S., Huegen, C., Van Ormondt, D., & Graveron-Demilly, D. (2001). Cramér–Rao
bounds: an evaluation tool for quantitation. *NMR in Biomedicine: An International Journal
Devoted to the Development and Application of Magnetic Resonance In Vivo, 14*(4), 278-283.

Chen, X., Fan, X., Hu, Y., Zuo, C., Whitfield-Gabrieli, S., Holt, D., Gong, Q., Yang, Y., Pizzagalli, D. A., & Du,
F. (2019). Regional GABA concentrations modulate inter-network resting-state functional
connectivity. *Cerebral Cortex, 29*(4), 1607-1618.

Cole, M. W., Yarkoni, T., Repovs, G., Anticevic, A., & Braver, T. S. (2012). Global connectivity of
prefrontal cortex predicts cognitive control and intelligence. *J Neurosci, 32*(26), 8988-8999.
https://doi.org/10.1523/JNEUROSCI.0536-12.2012

Colom, R., Jung, R. E., & Haier, R. J. (2006). Distributed brain sites for the g-factor of intelligence.
Neuroimage, 31(3), 1359-1365. https://doi.org/10.1016/j.neuroimage.2006.01.006

Cumming, B. G., & DeAngelis, G. C. (2001). The physiology of stereopsis. *Annu Rev Neurosci, 24*, 203-
238. https://doi.org/10.1146/annurev.neuro.24.1.203

Deary, I. J., Penke, L., & Johnson, W. (2010). The neuroscience of human intelligence differences.
Nature reviews neuroscience, 11(3), 201-211.

Donner, T. H., Siegel, M., Fries, P., & Engel, A. K. (2009). Buildup of choice-predictive activity in human
motor cortex during perceptual decision making. *Curr Biol, 19*(18), 1581-1585.
https://doi.org/10.1016/j.cub.2009.07.066

Dumoulin, S. O., Bittar, R. G., Kabani, N. J., Baker, C. L., Jr., Le Goualher, G., Bruce Pike, G., & Evans, A. C.
(2000). A new anatomical landmark for reliable identification of human area V5/MT: a
quantitative analysis of sulcal patterning. *Cereb Cortex, 10*(5), 454-463.
https://doi.org/10.1093/cercor/10.5.454

Duncan, J., Assem, M., & Shashidhara, S. (2020). Integrated Intelligence from Distributed Brain Activity.
Trends Cogn Sci, 24(10), 838-852. https://doi.org/10.1016/j.tics.2020.06.012

Duncan, J., Seitz, R. J., Kolodny, J., Bor, D., Herzog, H., Ahmed, A., Newell, F. N., & Emslie, H. (2000). A
neural basis for general intelligence. *Science, 289*(5478), 457-460.
https://doi.org/10.1126/science.289.5478.457

Ende, G. (2015). Proton Magnetic Resonance Spectroscopy: Relevance of Glutamate and GABA to
Neuropsychology. *Neuropsychol Rev, 25*(3), 315-325. https://doi.org/10.1007/s11065-015-
9295-8

Ernst, T., Kreis, R., & Ross, B. (1993). Absolute quantitation of water and metabolites in the human
brain. I. Compartments and water. *Journal of magnetic resonance, Series B, 102*(1), 1-8.

Everling, S., Tinsley, C. J., Gaffan, D., & Duncan, J. (2002). Filtering of neural signals by focused
attention in the monkey prefrontal cortex. *Nature neuroscience, 5*(7), 671-676.

Fangmeier, T., Knauff, M., Ruff, C. C., & Sloutsky, V. (2006). fMRI evidence for a three-stage model of
deductive reasoning. *Journal of cognitive neuroscience, 18*(3), 320-334.

Fetsch, C. R., Pouget, A., DeAngelis, G. C., & Angelaki, D. E. (2011). Neural correlates of reliability-
based cue weighting during multisensory integration. *Nat Neurosci, 15*(1), 146-154.
https://doi.org/10.1038/nn.2983

Frahm, J. a., Bruhn, H., Gyngell, M., Merboldt, K., Hänicke, W., & Sauter, R. (1989). Localized high-
resolution proton NMR spectroscopy using stimulated echoes: initial applications to human
brain in vivo. *Magnetic resonance in medicine, 9*(1), 79-93.

Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S., & Turner, R. (1996). Movement-related effects
in fMRI time-series. *Magnetic resonance in medicine, 35*(3), 346-355.

Gautama, T., & Van Hulle, M. M. (2001). Function of center-surround antagonism for motion in visual
area MT/V5: a modeling study. *Vision Res, 41*(28), 3917-3930. https://doi.org/10.1016/s0042-
6989(01)00246-2
Gray, J. R., Chabris, C. F., & Braver, T. S. (2003). Neural mechanisms of general fluid intelligence. *Nat
Neurosci, 6*(3), 316-322. https://doi.org/10.1038/nn1014

Gruetter, R. (1993). Automatic, localized in vivo adjustment of all first-and second-order shim coils.

Magnetic resonance in medicine, 29(6), 804-811.
Gu, Y., Watkins, P. V., Angelaki, D. E., & DeAngelis, G. C. (2006). Visual and nonvisual contributions to
three-dimensional heading selectivity in the medial superior temporal area. *J Neurosci, 26*(1),
73-85. https://doi.org/10.1523/jneurosci.2356-05.2006
Haier, R. J., Siegel Jr, B. V., Nuechterlein, K. H., Hazlett, E., Wu, J. C., Paek, J., Browning, H. L., &
Buchsbaum, M. S. (1988). Cortical glucose metabolic rate correlates of abstract reasoning and
attention studied with positron emission tomography. *Intelligence, 12*(2), 199-217.
Hayes, A. F. (2013). *Introduction to Mediation, Moderation, and Conditional Process Analysis: A

Regression-Based Approach* (Vol. 51). The Guilford Press.

Huk, A. C., Dougherty, R. F., & Heeger, D. J. (2002). Retinotopy and functional subdivision of human
areas MT and MST. *Journal of Neuroscience, 22*(16), 7195-7205.
Jung, R. E., & Haier, R. J. (2007). The Parieto-Frontal Integration Theory (P-FIT) of intelligence:
converging neuroimaging evidence. *Behav Brain Sci, 30*(2), 135-154; discussion 154-187.
https://doi.org/10.1017/S0140525X07001185
Keller, A. J., Roth, M. M., & Scanziani, M. (2020). Feedback generates a second receptive field in
neurons of the visual cortex. *Nature, 582*(7813), 545-549.
Li, M., Song, X. M., Xu, T., Hu, D., Roe, A. W., & Li, C.-Y. (2019). Subdomains within orientation columns
of primary visual cortex. *Science advances, 5*(6), eaaw0807.
Liu, D. Y., Ju, X., Gao, Y., Han, J. F., Li, Z., Hu, X. W., Tan, Z. L., Northoff, G., & Song, X. M. (2022). From
Molecular to Behavior: Higher Order Occipital Cortex in Major Depressive Disorder. *Cereb
Cortex, 32*(10), 2129-2139. https://doi.org/10.1093/cercor/bhab343
Liu, L. D., Haefner, R. M., & Pack, C. C. (2016). A neural basis for the spatial suppression of visual
motion perception. *Elife, 5*, e16167.
Marjańska, M., Auerbach, E. J., Valabrègue, R., Van de Moortele, P. F., Adriany, G., & Garwood, M.
(2012). Localized 1H NMR spectroscopy in different regions of human brain in vivo at 7 T: T2
relaxation times and concentrations of cerebral metabolites. *NMR in Biomedicine, 25*(2), 332-
339.
Melnick, M. D., Harrison, B. R., Park, S., Bennetto, L., & Tadin, D. (2013). A strong interactive link
between sensory discriminations and intelligence. *Curr Biol, 23*(11), 1013-1017.
https://doi.org/10.1016/j.cub.2013.04.053
Ozeki, H., Finn, I. M., Schaffer, E. S., Miller, K. D., & Ferster, D. (2009). Inhibitory stabilization of the
cortical network underlies visual surround suppression. *Neuron, 62*(4), 578-592.
Reynolds, J. H., & Heeger, D. J. (2009). The normalization model of attention. *Neuron, 61*(2), 168-185.
Sato, T. K., Haider, B., Häusser, M., & Carandini, M. (2016). An excitatory basis for divisive
normalization in visual cortex. *Nature neuroscience, 19*(4), 568-570.

Schallmo, M. P., Kale, A. M., Millin, R., Flevaris, A. V., Brkanac, Z., Edden, R. A., Bernier, R. A., & Murray,
825   S. O. (2018). Suppression and facilitation of human neural responses. *Elife*, *7*.
826   https://doi.org/10.7554/eLife.30334

827   Siegel, M., Buschman, T. J., & Miller, E. K. (2015). Cortical information flow during flexible
828   sensorimotor decisions. *Science*, *348*(6241), 1352-1355.
829   https://doi.org/10.1126/science.aab0551

830   Song, M., Zhou, Y., Li, J., Liu, Y., Tian, L., Yu, C., & Jiang, T. (2008). Brain spontaneous functional
831   connectivity and intelligence. *Neuroimage*, *41*(3), 1168-1176.
832   https://doi.org/10.1016/j.neuroimage.2008.02.036

833   Song, X. M., Hu, X. W., Li, Z., Gao, Y., Ju, X., Liu, D. Y., Wang, Q. N., Xue, C., Cai, Y. C., Bai, R., Tan, Z. L., &
834   Northoff, G. (2021). Reduction of higher-order occipital GABA and impaired visual perception
835   in acute major depressive disorder. *Mol Psychiatry*, *26*(11), 6747-6755.
836   https://doi.org/10.1038/s41380-021-01090-5

837   Spearman, C. (1904). " General Intelligence" Objectively Determined and Measured. *American Journal*
838   *of Psychology*, *15*, 201-293.

839   Tadin, D. (2015). Suppressive mechanisms in visual motion processing: From perception to intelligence.
840   *Vision Res*, *115*(Pt A), 58-70. https://doi.org/10.1016/j.visres.2015.08.005

841   Tadin, D., Lappin, J. S., Gilroy, L. A., & Blake, R. (2003). Perceptual consequences of centre-surround
842   antagonism in visual motion processing. *Nature*, *424*(6946), 312-315.
843   https://doi.org/10.1038/nature01800

844   Tadin, D., Silvanto, J., Pascual-Leone, A., & Battelli, L. (2011). Improved motion perception and
845   impaired spatial suppression following disruption of cortical area MT/V5. *Journal of*
846   *Neuroscience*, *31*(4), 1279-1283.

847   Tkáč, I., Starčuk, Z., Choi, I. Y., & Gruetter, R. (1999). In vivo 1H NMR spectroscopy of rat brain at 1 ms
848   echo time. *Magnetic Resonance in Medicine: An Official Journal of the International Society*
849   *for Magnetic Resonance in Medicine*, *41*(4), 649-656.

850   Treue, S., & Maunsell, J. H. (1996). Attentional modulation of visual motion processing in cortical areas
851   MT and MST. *Nature*, *382*(6591), 539-541.

852   Wang, X. J. (2002). Probabilistic decision making by slow reverberation in cortical circuits. *Neuron*,
853   *36*(5), 955-968. https://doi.org/10.1016/s0896-6273(02)01092-9

854   Wechsler, D. (2008). Wechsler Memory Scale–Fourth Edition (WMS-IV) technical and interpretive
855   manual.

856   Wilming, N., Murphy, P. R., Meyniel, F., & Donner, T. H. (2020). Large-scale dynamics of perceptual
857   decision information across human cortex. *Nat Commun*, *11*(1), 5109.
858   https://doi.org/10.1038/s41467-020-18826-6

859   Wimmer, K., Compte, A., Roxin, A., Peixoto, D., Renart, A., & de la Rocha, J. (2015). Sensory integration
860   dynamics in a hierarchical network explains choice probabilities in cortical area MT. *Nat*
861   *Commun*, *6*, 6177. https://doi.org/10.1038/ncomms7177

862   Yan, C.-G., Wang, X.-D., Zuo, X.-N., & Zang, Y.-F. (2016). DPABI: data processing & analysis for (resting-
863   state) brain imaging. *Neuroinformatics*, *14*, 339-351.

864   Zaksas, D., & Pasternak, T. (2006). Directional signals in the prefrontal cortex and in area MT during a
865   working memory for visual motion task. *J Neurosci*, *26*(45), 11726-11742.
866   https://doi.org/10.1523/JNEUROSCI.3420-06.2006

Zhang, S., Xu, M., Kamigaki, T., Hoang Do, J. P., Chang, W.-C., Jenvay, S., Miyamichi, K., Luo, L., & Dan, Y.
(2014). Long-range and local circuits for top-down modulation of visual cortex processing.
Science, 345(6197), 660-665.
Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov
random field model and the expectation-maximization algorithm. *IEEE transactions on
medical imaging*, *20*(1), 45-57.

Acknowledgements

The authors thank Prof. Dost Ongur and Fei Du for guidance on the MRS data processing. Thank Zhejiang University 7T Brain Imaging Research Center, and thank Guohua Xu and Fen Yang for technical assistance. Thank Prof. Xinyi Lai for supporting MRI data acquisition. This work was supported by STI 2030—Major Projects (2021ZD0200401 to X.M.S., 2022ZD0206000 to R.B.), the National Natural Science Foundation of China Grants (U1909205, 61876222, 32000761, 82222032), Humanities and Social Sciences Ministry of Education (20YJC880095, 18YJA190001), the Key R&D Program of Zhejiang (2022C03096 to X.M.S., 2022ZJJH02-06 to G.C.), the European Union's Horizon 2020 Framework Program for Research and Innovation under the Specific Grant Agreement No. 785907 (Human Brain Project SGA2 to G.N.), and the MOE Frontier Science Center for Brain Science & Brain- Machine Integration, Zhejiang University.

Author contributions

S.X.M. and Y.G. designed the experiment and G.N. guided the logic of the analysis. Y.G., Y.C. and D.L. conducted human experiments, analyzed data, and Y.G. created

891 figures. R.B. guided the MRI experiments. J.Y., J.W. and B.X. were in charge of data

892 collection and partial date analysis. T. W., M.L. and G.C. guided the data analysis.

893 S.X.M., Y.G. and G.N. wrote the manuscript.

Competing interests

896 The authors declare no competing interests.