Large diversity in the O-chain biosynthetic cluster within populations of

Pelagibacterales

Jose M. Haro-Moreno[1], Mario López-Pérez[1], Carmen Molina-Pardines[1], and

Francisco Rodriguez-Valera[1,*]

[1]Evolutionary Genomics Group, División de Microbiología, Universidad Miguel

Hernández, Apartado 18, San Juan 03550, Alicante, Spain.

*Corresponding author: frvalera@umh.es

Universidad Miguel Hernández, División de Microbiología, Apartado 18, San Juan de

Alicante, 03550 Alicante, Spain.

Phone +34-965919313, Fax +34-965 919457

Running title: O-chain diversity in SAR11

21 **Keywords**: SAR11 / Pelagibacterales / O-antigen / Lipopolysaccharide (LPS) /

22 Single-amplified genomes / Genomic Islands / PacBio metagenomics

ABSTRACT

We have used single-amplified genomes (SAGs) and long-read metagenomics to examine the diversity of the O-chain polysaccharide biosynthesis cluster (OBC) in marine bacteria of the Pelagibacterales order. OBCs are notorious for their diversity and have been used to type strains in pathogens and saprophytes, but their patterns of variation in free-living bacteria are little known. We found that, for these marine heterotrophic bacteria, the diversity is comparable to that of saprophytes, such as Enterobacteriales i.e. nearly each strain carries a different OBC. However, although OBC inheritance was largely vertical, the existence of some shared clusters allowed a comparative analysis. The OBCs diverge along with the genome, which is taken as indicative of old horizontal gene transfer (HGT) events. Only 14 cases of recent HGT were detected and they happened independently of taxonomy or location. Thus, although the O-chain is a major phage receptor in Gram-negative bacteria, the exchange of the complete cluster seems to play a minor role in the phage-bacterium arms-race. By long-read metagenomics, we could detect 380 different OBCs in a single sampling site in the Mediterranean. A single population (single species and sample) of the endemic Ia.3/VII genomospecies had a set of 158 OBCs of which 130 were different. This large diversity in clonal lineages might reflect the large amount of metabolic pathways required to deal with the enormous chemical diversity of dissolved organic matter in the ocean.

INTRODUCTION

In a recent review (Dittmar et al., 2021) it was calculated that in excess of 400,000 independent metabolic pathways were required to metabolize marine DOM and several hundreds of thousands of ABC transporters have been identified in marine metagenome assemblies (Zhao et al., 2024). Thus, it is to be expected that a single species of a dominant heterotrophic microbe such as Pelagibacter could contain a large diversity of genes in its local pangenome. However, determining the diversity of strains within populations of bacteria has been a major conundrum in microbiology (Viver et al., 2024). Some recent studies in the gut microbiome by culture (Huang et al., 2023), metagenomics (Costea et al., 2017; Sharon et al., 2013; Truong et al., 2017), and long-read amplicons of flagellins (Hu et al., 2022) have concluded that few strains of each species dominate the population within a single individual human microbiome and that they are relatively stable through time. However, studies from metagenomics and single-cell genomics indicated that diversity within a single drop of seawater might be very high (Coleman et al., 2006; Kashtan et al., 2014; Rusch et al., 2007). Nevertheless, even the orders of magnitude of such diversity were hard to establish due to the small size of metagenomic reads or the incompleteness of single-amplified genomes (SAGs). A recent study by culture and metagenomics in an extreme aquatic environment revealed a very high strain diversity in *Salinibacter ruber* with several hundreds of strains within a single pond (Viver et al., 2024).

A very common (if not universal) component of the flexible genome (variable from one strain to another within the same species) is the gene cluster coding for the synthesis of external polysaccharides (Raetz and Whitfield, 2002; Rodriguez-Valera et al., 2016), such as the one containing the genes synthesizing the O-chain of the

lipopolysaccharide (often referred to as O-antigen) (Kalynych et al., 2014; Liu et al.,

2019; Samuel and Reeves, 2003). They have been known for long in

pathogenic/saprophytic bacteria (Huszczynski et al., 2019; Kenyon et al., 2017; Liu

et al., 2014; Mostowy and Holt, 2018); their enormous variability has been used to

type strains in epidemic outbreaks, i.e. assuming that each strain has a unique

combination of exposed polysaccharides. In the order Enterobacteriales, a recent

study by Holt and colleagues (Holt et al., 2020) clearly illustrates the patterns of

variation found at these organisms' major glycotype (combinations of exposed

polysaccharides) gene clusters. After analyzing more than twenty-seven thousand

genomes, they shed light on the evolution of these loci. Their enormous diversity (ca.

18,000 different OBCs were described) and rare exchange between different strains

explains their discriminating power used in epidemiology to identify outbreaks

produced by the same strain. On the other hand, when HGT was detected the

divergence of the genes indicated old transfers (long-term preservation), rare events

that do not affect the association between OBC and strain (Holt et al., 2020). These

results also discredit the view in which diversity of OBCs is explained by phage-host

arms-race that would require fast exchange of the OBCs.

Here we have focused on a study of O-chain biosynthetic gene cluster (OBC)

diversity in the marine order Pelagibacterales, taking advantage of the abundance of

this oligotrophic microbe in SAG databases and the use of long-read (PacBio HiFi)

metagenomics. Their photoheterotrophic lifestyle is well known and makes them

among the most relevant microbes in nutrient fluxes in the oligotrophic ocean (Brown

et al., 2012; Giovannoni, 2017; Grote et al., 2012; Wilhelm et al., 2007). Besides,

their streamlined genome size of ~1.3 Mb (one of the smallest for planktonic free-

living microbes) (Giovannoni et al., 2005; Grote et al., 2012), simplifies the analysis,

as they contain a single gene cluster involved in O-chain biosynthesis (e.g. no

capsular envelope has ever been found and they have no flagella). Furthermore, the

cluster is bounded by the two parts of the ribosomal RNA operon 16S-23S on one

side and 5S on the other (Wilhelm et al., 2007), which provides good markers for

bioinformatic searches. In addition, a fine taxonomy derived from the ITS can be

used to classify the genome with high reliability (García-Martínez and Rodríguez-

Valera, 2000) and the rRNA genes can be used as hallmarks for identifying

Pelagibacterales reads in long-read metagenomes obtained from the same location

in off-shore Mediterranean waters (Haro-Moreno et al., 2021; Zaragoza-Solas et al.,

2022). This way we have been able to dissect the diversity of OBC gene clusters as

a whole and then also at the population (one single species in one single sample)

level. Our findings about the diversity of these OBCs indicate that indeed local strain

diversity can be very high even at the single population level.

RESULTS

OBCs variation across the order Pelagibacterales.

Our first objective has been the analysis of the evolutionary dynamics in the

available diversity of Pelagibacterales OBCs, to check if the pattern described

previously for saprophytic microbes (Enterobacteriales) could be extrapolated to

free-living ones that never interact with immune systems. We screened a dataset

comprising nearly 1,700 marine SAGs (Berube et al., 2018; Haro-Moreno et al.,

2020; Pachiadaki et al., 2019; Thompson et al., 2019; Thrash et al., 2014) and 20

marine isolate genomes from the whole order Pelagibacterales (clades Ia, Ib, Ic IIa,

IIb and IIIa) (**Figure S1**). We avoided clades IV and V, the latter also known as

HIMB59, since they have been recently classified as different orders (Haro-Moreno

et al., 2020; Molina-Pardines et al., 2023; Viklund et al., 2013). By using the 16S-

ITS-23S rRNA operon and the 5S rRNA gene at the left and right ends, respectively,

we could recover 806 OBCs >10 Kb long (**Figure S1A**). The Pelagibacterales are

characterized by unusually high synonymous replacements (López-Pérez et al.,

2020a) and thus the 95% average nucleotide similarity (ANI) used for bacterial

species definition (Konstantinidis and Tiedje, 2005) gives a very split taxonomy that

contrasts with their high synteny and coverage values (Haro-Moreno et al., 2020).

Thus, we have used a genomospecies classification based on phylogenomics and

environmental distribution described in (Haro-Moreno et al., 2020).

After manual curation, we found an exception of the location of the OBC in genomes

from the genomospecies Ib.4, where there was a genomic rearrangement, that in

this clade was found between the 16S-ITS-23S at one end (like in the other clades)

but at the other end was located near a cluster of genes involved in the

peptidoglycan biosynthesis (**Figure S2A**) and two tRNAs coding for valine and

methionine located ~330 Kb distant from their location in the other clades. A total of

27 SAGs, all in the Ib.4, contained the relocated island. Given that there is no

complete genome available for this genomospecies, we collected into a single contig

three almost identical (>99 % ANI) SAGs into a partially complete (1.06 Mb) and

admittedly chimeric construct (López-Pérez et al., 2020b; Roda-Garcia et al., 2023).

Figure S2B shows the reconstructed genomic fragment, indicating the position of

the 16S, 23S, and 5S rRNA genes. Metagenomic under-recruitment confirmed that

the region in **Figure S2A** corresponded indeed to the large OBC found in other

clades and that it has been likely translocated to one of the tRNAs, a frequent target

of site-directed non-homologous recombination. Still, it was used for the subsequent

diversity comparisons and showed very similar behavior (see below).

One hundred and sixty-three islands (one-fifth of the dataset) were categorized as

complete, including eight from the rearranged OBCs found in Ib.4. The remaining

were partial islands, from which we could recover only the left-hand side (~23 %),

right-hand side (~22 %), or both sides but in different contigs within the same SAG

(~35 %), (**Figure S1A and Table S1**). Regarding their origin, nearly half of the OBCs

(#373) were recovered from a single sample in the Bermuda Atlantic Time-series

Study (BATS) (sample SWC-09) collected from a 10 m deep water column sample

(**Figure S1B**) (Pachiadaki et al., 2019). Therefore, we can consider all these SAGs

as belonging to a single community, and those within a single species, as members

of a single population. The remaining OBCs came from a variety of samples from the

Atlantic and Pacific Oceans (#170 and #197, respectively) (Berube et al., 2018;

Pachiadaki et al., 2019), and from the Mediterranean (Haro-Moreno et al., 2020)

(#33) and the Red Seas (#15) (Thompson et al., 2019) (**Figure S1B**). We classified

the Pelagibacterales genomes containing OBCs as members of the subclades Ia

and Ib (~90 % of the OBCs), and in much smaller numbers to subgroups Ic, IIa, IIb,

and IIIa (**Figure S1C**). This was to be expected since most samples were collected

from surface waters, where members of subclades Ia and Ib are largely

predominant. Members of subclades Ic and II are mainly found in meso- and

bathypelagic waters (Giovannoni, 2017; Thrash et al., 2014; Tsementzi et al., 2016),

although some members of the subclade IIa have been detected in surface waters

(Bolaños et al., 2022).

The analysis of the 163 complete OBCs showed a large size range, from 10 to 90 Kb (average size 46 Kb) (**Figure S3 and Table S1**). The number of encoded genes varied from 8 to 101, with an average of 45.7 genes per OBC. Grouped by taxa, the average OBC size varied slightly for members of subclades Ia.1 (53 Kb), Ia.3 (51 Kb), and Ia.4 (45 Kb), while subclades Ib.1 (n=13, 57 Kb), and IIa (n=11, 27 Kb) had the largest and smallest glycosylation islands, respectively (**Figure S3A**). OBCs tend to have a GC content lower than the average of the genome in all Gram-negative bacteria (da Silva Filho et al., 2018). For the Pelagibacterales SAGs described here, regardless of the clade considered, the value for the complete OBCs GC content was around 23.8 %, statistically significant (paired t-test, p-value $1e^{-116}$), lower than the average GC content of the genomes (29.4 %) (**Figure S3B**). The median intergenic spacer (very small in streamlined genomes) was also statistically different between genomes and their OBCs (paired t-test, p-value 0.003), due to the high dispersion of their values (**Figure S3C**). Both differences might reflect the presence of genes of exogenous origin within the OBCs, although it seems difficult to find many microbes with lower GC content than the Pelagibacterales.

Similarity of shared OBC-types.

To evaluate the presence of similar OBC ORFs among the Pelagibacterales genomes, we performed an all-vs-all comparison among them (see methods) (**Table S2**). As shown in **Figure S4**, the majority of the comparisons showed no similarity, measured as the fraction of orthologous genes shared among two OBCs at 50 % AAI and 70 % coverage. For instance, 75.7 % of the OBC pairs shared less than 5 % of orthologs, while this number increases to 94.4 % if we consider those with less

than 15 % of orthologous genes **(Figure S4)**. These results illustrate the enormous diversity of OBC genes within the Pelagibacterales order, similar to Enterobacteriales, where 18,384 OBCs resulted in only 2,654 (14.4 %) coding for the same gene sets (Holt et al., 2020). We thus consider two OBCs to belong to the same locus type (hitherto OBC-type) similar to (Holt et al., 2020) if they share at least 90% of the genes. As a result, from the initial set of 806 OBCs (including incomplete ones), 208 genomes shared the OBC-type with another genome in our dataset, while 598 genomes had unique (singleton) OBCs. This is typical of the so-called replacement flexible genomic islands that, although coding for a similar function or (in this case) structure, have completely different gene make-up (López-Pérez et al., 2014; Rodriguez-Valera et al., 2016). An advantage of this kind of genomic island for comparative analysis is that incomplete clusters can be used in pairwise comparisons assuming that similarity throughout some genes can be taken as a sign of a common OBC and likely also a similar (or identical) polysaccharide or glycotype. This is important when working with SAGs, which tend to be fragmented and incomplete, and even more so for long metagenomic reads (see below).

As expected, either the clusters are syntenic and have >85 % similarity or they are completely different in gene content with nearly no orthologous genes detected **(Figure S4)**. For the OBCs belonging to the same OBC-type, in most cases, similarity was very high over the whole stretch, but there were exceptions. For example, **Figure S5** shows in one of the pairs similarity has decreased throughout the right-hand side end of the OBC. In other cases (**Figure S5**) variation seems to be concentrated at the left-hand-side end. In these two cases (and in most) variability seems to increase at the ends (one of them), as was the case in the

Enterobacteriales (Holt et al., 2020). This fact improves the reliability of partial

sequence comparisons as was done with the metagenomic long reads (see below).

Figure S6 shows an average amino acid identity (AAI) cladogram tree

(Konstantinidis and Tiedje, 2005) rather than the more common average nucleotide

identity (ANI). AAI relationships are helpful when comparing highly divergent genome

sequences (Barco et al., 2020), as those analyzed here for the whole

Pelagibacterales order. **Figure 1A** only shows the genomes sharing OBC-types**.** A

somewhat expected observation was that in most cases SAGs sharing the OBCs

were also >99 % AAI overall i.e., they belonged to the same clonal frame or

genomovar (Viver et al., 2024). This result was not surprising, since this is the

fundament of strain serotyping of Gram-negative bacteria using O-antigens

(Kauffmann, 1947; Liu et al., 2019). However, a significant number of associations

(~13 %) were found between individuals belonging to different genomospecies from

the same family, or even from different families (**Figure 1A**). There seems to be a

bias for certain groups regarding the number of common OBC-types. For instance,

genomospecies Ia.3/V (also named as gWID given that has a widespread oceanic

distribution by metagenomic fragment recruitment (Delmont et al., 2019; Haro-

Moreno et al., 2020)), is one of the genomospecies with the highest number of

retrieved genomes (#79, **Figure S6**) but barely showed any shared OBC-type (#7,

Figure 1A), six of them all corresponded to the same genomovar. Whereas Ib.1,

with 87 genomes, had 37 sharing events, five of them jumping across different

genomovars. Many shared OBC-types were detected only among members of the

same genomospecies, and often between genomes with >95 % AAI (**Figure S7**).

Only 49 OBC-types were found across two different genomospecies (31 of the pairs

having more than 95 % AAI), while only a single OBC-type was found in four

different genomospecies and one in three (**Figure S7**). Most shared OBCs had the

same gene content or only varied in one gene, regardless of the dissimilarity

between genomes sharing the OBC **(Figure 1B)**. Lastly, the rate of shared OBC-

types (**Figure 1C**) drops dramatically below 95 % AAI, regardless of the taxa, and

was negligible (24 OBCs) in genomes with less than 90 % AAI. The same pattern

applies even when considering different taxa individually i.e. members of the same

genomospecies with less than 90 % AAI very seldom share OBC-type.

The finding of O-antigen loci shared among distant families suggests horizontal gene

transfer (HGT). However, as seen in **Figure 2A**, for most shared OBCs (82 %), the

locus genetic distance expressed as ANI (data not shown) or AAI was similar to or

slightly higher than the genome distance. Actually, the dN/dS was statistically

significantly higher for shared OBC genes (average 0.14 versus 0.1 for the whole

genome, paired t-test, p-value $<1e^{-15}$) indicating a slightly higher rate of positive

selection (**Figure 2B**). In any case, the values detected for OBC genes indicate that,

if they have been exchanged by HGT, it must have been an old event and for the

most part they are vertically inherited (long-term preservation) as described for the

Enterobacteriales (Holt et al., 2020). The O-chain is a primary target for phages and

might be subject to arms-race rapid evolution i.e. change much more rapidly than the

rest of the genome to evade phage predation (Letarov, 2023; Mostowy and Holt,

2018). However, the data indicates that this is not achieved by swapping the whole

OBC or is very rare, as only fourteen (~7 % of the total shared OBCs) had an OBC

distance that was at least half of the genomic distance, indicative of recent events

(**Figure 2A**). These results reveal that rather than rapidly exchanging their OBCs as

some scenarios suggest (Mostowy and Holt, 2018; Rostøl and Marraffini, 2019),

most OBCs are vertically transmitted over long evolutionary periods. Arms-race is

more likely to act at the level of mutation or recombination affecting single genes.

Given that nearly half of the SAGs in the database came from a single sample of

water in the Sargasso Sea (Pachiadaki et al., 2019), we studied how many different

OBC-types could be detected at a single location and sample. To do that, we

rarefied (Chao et al., 2014; Hsieh et al., 2016) the genomes and the number of

different OBC-types detected globally and at BATS. The 373 BATS´ SAGs for the

whole Pelagibacterales order had a weak sign of saturation (**Figure 3A**). As

expected this coverage was even lower (~25 %) considering all the 806 genomes for

Pelagibacterales as a whole. Therefore, we are far from finding all possible

variations in the Pelagibactales OBCs either in the whole ocean or in a single

sample. At the genomospecies level and in the single BATS sample, the numbers

were too small to allow for a sensible estimation of the total numbers of OBC-types

in a single population.

Diversity of O-antigen loci within a single population by long-read

metagenomics

Metagenomic samples sequenced with PacBio Sequel II can be a good alternative to

SAGs to study intrapopulation diversity. Given the large size of the reads, typically

between 5 to 15 Kb long (Haro-Moreno et al., 2021), the flexible part of a genome,

including fragments presenting part of the OBC can be retrieved before assembly.

As discussed before, retrieving only a few genes allows inferring the OBC-type

represented by the PacBio read. Thus, we have sequenced and analyzed five

different PacBio CCS metagenomes collected from a single location at an off-shore

Western Mediterranean Sea off Alicante (see methods). We took samples during

winter when the water column is fully mixed (MedWinter-JAN2019 (Haro-Moreno et

al., 2021), MedWinter-FEB2022), and during the stratification season at three

different layers in the photic zone, the upper photic (Med-OCT2021-15m), the deep

chlorophyll maximum (Med-SEP2022-60m, DCM) and the lower photic (Med-

OCT2021-75m, below DCM). The prokaryotic community at this location has been

thoroughly studied by short and long-read metagenomics before (Haro-Moreno et al.,

2021, 2019, 2018; López-Pérez et al., 2017) and the three depths selected when the

water column is stratified represent different epipelagic assemblages, as the

prokaryotic community significantly differed at these depths (Haro-Moreno et al.,

2018). Then, we selected all the reads whose 23S rRNA affiliated within the order

Pelagibacterales, and hence the left-hand end of the OBCs could be analyzed to

study the variability of the OBC-types within these samples. Only PacBio CCS reads

with at least 5 Kb of OBC were used, resulting in a total of 2,780 OBCs from the five

metagenomic samples, and the threshold of at least 90 % of orthologous genes was

applied for their classification into the same OBC-type. Hence, we found 2,440 OBC-

types in all the samples combined (**Figure S8**). The resulting rarefaction,

extrapolating to 10,000 reads, showed several OBC-types, between 4,500 and

6,500, for the whole Pelagibacterales order at this single sampling site and over the

course of a couple of years with the corresponding seasonal variation (**Figure S8**).

This number is certainly in the order of the one found for the order Enterobacteriales

(15,730 in 27,000 genomes) (Holt et al., 2020).

To be able to classify the recovered OBCs into genomospecies, we have used the

internal transcribed spacer (ITS) between the 16S rRNA-23S rRNA genes. Ecotypes

of Pelagibacterales and *Prochlorococcus* have been reliably classified based on their

sequence for a long time (Brown and Fuhrman, 2005; García-Martínez and

Rodríguez-Valera, 2000; Ngugi and Stingl, 2012; Rocap et al., 2002; Shibl et al.,

2014). More recently, a clear association between genomospecies classified by

phylogenomics and the ITS cladogram was confirmed (Haro-Moreno et al., 2020). By

examining the correspondence between ITS and 23S rRNA gene sequences, we

also determined genomospecies from the latter. Therefore, the long-reads with

complete ITS (n=1,501) and 23S rRNA sequences (n=2,780) were used to analyze

the OBC diversity in single genomospecies (**Table 1**). As expected, the diversity

found within the Pelagibacterales CCS reads was astounding, with many sequences

grouping with the main phylogenetic groups (Ia, Ib, Ic, and IIa). In the case of the

most abundant genomospecies (by number of OBC reads), Ia.3/VII and Ic.2 were

detected at significant values (>5 %). Ia.3/VII had 18.4 % of the recovered OBC

reads. This group has been characterized as dominating (highest metagenomic

recruitment) in epipelagic Mediterranean waters and hence it was also named as

gMED (Haro-Moreno et al., 2020). By using only the sequences from a single

sample (mixed water column, MedWinter-FEB2022) that had the largest number of

OBC reads (158 gMED-affiliated sequences), 130 different OBCs were retrieved in

this population, and the rarefaction curve suggested the presence of around 400

different OBC types (294 - 480, 95 % CI) (**Figure 3B**). A similar approach was also

applied to another genomospecies, Ic.2 (deep epipelagic), which was found

abundant (112 out of 474 OBC identified reads) in the lower photic (75 m deep)

sample. The rarefaction curve extrapolates to 254 different OBCs (140 - 310, 95 %

CI). Although the numbers are smaller than the ones found for the species *E. coli* as

a whole (#800 in (Holt et al., 2020)), the *E. coli* OBCs, although coming from a single

well-defined species, were collected from several different populations (e.g. hosts or

environments) and the genomic sampling was much larger (ca. 10,000 complete

genomes).

DISCUSSION

Polysaccharides are extremely versatile macromolecules in which an infinity of

structures and properties can be obtained by small modifications in their sugars,

bonding order, branching and length. In this sense, they are like proteins but they

have to be synthesized by laborious enzymatic steps, exported to the outside of the

cell and then polymerized. We have not analyzed the functionality of the genes

detected, among other reasons because they lack in most cases reliable annotations

beyond the general description of function. But the whole subject of polysaccharide

biosynthesis even in model organisms requires more extensive work. Structural

predictions and comparisons could help to solve the conundrum but are beyond the

scope of this work.

The main objective of this work has been estimating the numbers of strains, i.e.

lineages of the same species but with different flexible genomes, present in a single

population (same sample). To that end, we have studied the diversity of OBCs within

the Pelagibacterales order, as the variation of this region have been traditionally

used to type strains (Huszczynski et al., 2019; Kenyon et al., 2017; Liu et al., 2014;

Mostowy and Holt, 2018). Unfortunately, we are far from answering this question

with the kind of data used here, but we can infer (Hu et al., 2022) that it should be

slightly less than the numbers of different OBCs, i.e. in the order of hundreds of

strains. This number fits well with the numbers found for *Prochlorococcus* by

Kashtan and collaborators (Kashtan et al., 2014) by comparing whole SAGs, and it is

again surprisingly high for individuals (cells) belonging to the same species and

inhabiting a relatively homogeneous environment. They would certainly allow for

dealing with large numbers of substrates and conditions by one single species. It has

been estimated that dissolved organic matter (DOM) in the ocean contains in the

order of 100,000 different chemical formulas half of which have a half-life time of less

than two weeks (Zark et al., 2017). Therefore, the number of genes (e.g.

transporters, degradative enzymes) required to cope with such chemical diversity is

expected to be very large (Dittmar et al., 2021). Thus, it is not surprising that major

consumers of DOM such as Pelagibacterales species have proportionally large local

gene pools or pangenomes.

The overall diversity and evolution of OBCs described here in a free-living marine

proteobacterial order is very similar to that described using cultures of a

heterogeneous group of saprophytic and free-living bacteria like the

Enterobacteriales. This proves beyond doubt that the classical view (Lerouge and

Vanderleyden, 2002; Reeves, 1995) that O-chain diversity is due to the need of the

microbes to change their antigenic specificity as protection from host immune

systems is unlikely, even if this is partially true under some special circumstances.

Furthermore, the term "serotype" should be replaced by "glycotype" which is more

realistic (López-Pérez and Rodriguez-Valera, 2016). The variation at the level of

exposed polysaccharides in microbes belonging to the same population has been

revealed by metagenomics, culture, and other approaches as a constant feature, at

least, for aquatic microbes (Layoun et al., 2024; Roda-Garcia et al., 2023;

Rodriguez-Valera et al., 2016), including Gram positives (López-Pérez et al., 2020b;

Neuenschwander et al., 2018) and even Archaea (Martin-Cuadrado et al., 2015).

The reasons for such extreme flexibility in structures often critical to the survival of

cells in nature are still not clear, but in free-living prokaryotes, the most obvious

reason to have a high diversity of these markers, within a single population, is avoid

predation by protists or phages. Although some O-chains have been considered

more refractory to protist predation than others (Sintes and del Giorgio, 2014), there

is little doubt that for tiny cells like those of the Pelagibacterales, the predatory

pressure of phages is likely more important. Something along the same lines has

been described for the most abundant picocyanobacterium *Prochlorococcus*, and its

phages (Avrani et al., 2011; Coleman et al., 2006; Schwartz and Lindell, 2017). The

O-chain is a major target for phage receptor-binding proteins and thus there are two

potential phage-related explanations for its diversity- arms-race and density-

dependent negative selection (Abedon, 2022; Rodriguez-Valera et al., 2009). In the

latter, increased predation on abundant OBC-types, maintains a large diversity of

receptors, distributing the predation pressure among many. The swapping of O-

chains is too slow to be effective in an arms-race scenario and, over long

timeframes, seems an unlikely explanation. However, at a shorter timeframe, it is

clear that mutation or individual gene or cassette swapping could have a role in the

phage-host interaction as seen in several laboratory experiments.

What is the relevance of arms race processes then? In the Enterobacteriales, the

gain or loss of a small number of genes (1-3) has been described within relatively

short (epidemiological) timescales (Holt et al., 2020). Experimental studies have

shown that isolated SNPs can make a strain resistant to one phage and also very

small variations in the phage receptor-binding protein can revert the resistance

(Schwartz and Lindell, 2017). However, it has been shown also by experimental

work in cyanophages that a decrease in sensitivity to one phage can increase the

sensitivity to another (Schwartz and Lindell, 2017). Furthermore, significant changes

in the O-chain structure or composition in a Gram-negative bacterium can alter their

antibiotic sensitivity (Pernitzsch et al., 2021) and also, likely, its nutritional

preferences. O-chain mutations are notoriously pleiotropic, as they have multiple

phenotypic effects (Martínez de Tejada et al., 1995; Pagnout et al., 2019). In

addition, the synthesis of an external polysaccharide requires many steps that are

connected (synthesis of sugars, transport and linking to the growing external

polymer. All these steps must be coordinated and thus submitted to a complexity

limitation to change (Perlovsky and Frank-Kamenetskii, 2002). Finally, we would like

to suggest that once reached an equilibrium, any change could lead to a major

disruption of the species' adaptation to its niche. In the evolutionary landscape

analogy (Gokhale et al., 2009), it would imply falling from the mountain peak to the

valley and is likely to be very infrequent.

METHODS

Recovery of Pelagibacterales genomes. To evaluate the presence and the

number of glycosylation islands from the whole Pelagibacterales order (NCBI

taxonomy ID 54526), a compendium of nearly 4,100 genome assemblies was

downloaded from the NCBI database. Prior to the analysis, due to the incomplete

nature of MAGs, only assemblies from SAGs and isolates were considered, and their

degree of completeness and contamination of SAGs were estimated using CheckM

v1.1.2 (Parks et al., 2015). SAGs with > 50% completeness and < 5% contamination

were kept. A fast taxonomic classification of genomes was performed using the

GTDB-Tk v2.1.0 tool (Chaumeil et al., 2019) using the Genome Taxonomy Database

(GTDB) release R207 (Parks et al., 2018). Genomes belonging to clades IV and V

(HIMB59) or misclassified were removed from the dataset.

Phylogenomic classification. Using Phylophlan (Segata et al., 2013) , a total of

104 genes (26,134 amino acid positions) were used to classify the 806

Pelagibacterales genomes phylogenomically. Genomes of the HIMB59 order and

Rickettsia *spp.* were used as an outgroup. The resulting tree was analyzed using

iTOL (Letunic and Bork, 2016). Following the well-established SAR11 nomenclature

within subclades, phylotypes, and genomospecies described in (Haro-Moreno et al.,

2020; López-Pérez et al., 2020a), we used the median distance between nodes and

cophenetic correlation coefficient (interval comprised between 0 and 2) to define

them.

Genome annotation and retrieval of the O-chain biosynthetic gene cluster

(OBC). Prodigal v2.6.3 (Hyatt et al., 2010) was used to predict genes from contigs

retrieved from the individual genomes in the curated Pelagibacterales dataset

containing 1,700 genomes. Predicted protein-encoded genes were taxonomically

and functionally annotated against the NCBI NR database using DIAMOND 0.9.15

(Buchfink et al., 2015) and against COG (Tatusov et al., 2001) and TIGRFAM (Haft

et al., 2001) using HMMscan v3.3 (Eddy, 2011). tRNA and rRNA genes were

predicted using tRNAscan-SE v2.0.5 (Lowe and Eddy, 1996) and barrnap v0.92

(https://github.com/tseemann/barrnap), respectively.

Contigs containing a signal for the 23S and 5S rRNA genes were selected as they

are characterized as markers for the start and ending points of the OBC in

Pelagibacterales. If both markers were found within the same contig the island was

classified as complete.

Genomic pairwise comparison. Average nucleotide and amino acid identities (ANI

and AAI) between a pair of genomes were calculated using the JSpecies with default

parameters (Richter and Rossello-Mora, 2009) and CompareM

(https://github.com/donovan-h-parks/CompareM) software packages respectively. A

cladogram of the genomes containing the OBC (complete or partial) was constructed

using an all-vs-all matrix of the AAI values among genomes, followed by hierarchical

clustering with the average Euclidean distance among values with the hclust function

in R (R Development Core Team and Team, 2011).

Determination of O-chain biosynthesis locus types. AAI values between pairs of

OBCs were calculated with CompareM, considering 50 % amino acid identity as the

threshold to establish similarity among genes. We only considered the pairwise

combinations of A (complete OBC) vs A, A vs B (partial OBC), B vs A, A vs C (only

left-hand side OBC), C vs A, A vs D (only right-hand side OBC), D vs A, B vs C, C vs

B, B vs D, D vs B, C vs C, and D vs D. Two or more OBCs belonged to the same

OBC-type if they shared at least 90 % of the genes (considering the smallest OBC).

dN/dS values from OBCs and genomes. Estimation of the numbers of

synonymous (dS), non-synonymous (dN) mutations, and the dN/dS ratio was

performed using orthologr (Drost et al., 2015) against the set of genomes sharing an

OBC and their OBCs. The OBC cluster was extracted from the genome sequences

before calculation. Ortholog genes were detected using a reciprocal best-hit

approach using DIAMOND and the dN/dS ratios were estimated using the Comeron

algorithm (Comeron, 1995).

PacBio CCS15 metagenomic reads. Four marine samples were collected from the

same sampling site in the epipelagic Mediterranean Sea at 20 nautical miles off the

coast of Alicante (Spain) (37.35361°N, 0.286194°W). MedWinter-FEB2022 (20 m

deep) was collected during winter, when the water column is fully mixed. Med-

OCT2021-15m, Med-SEP2022-60m and Med-OCT2021-75m were collected in

summer, during a strong stratification period. We added to the comparison a winter

sample collected in January 2019 and sequenced with PacBio Sequel II (MedWinter-

JAN2019 (Haro-Moreno et al., 2021)). For each depth, 200 L were collected and

filtered on board as described in the study of Haro-Moreno et al. (2018). Briefly,

seawater samples were sequentially filtered through 20-, 5-, and 0.22-µm pore filter

polycarbonate filters (Millipore). Water was directly pumped onto the series of filters

to minimize the bottle effect. Filters were immediately frozen on dry ice and stored at

−80°C until processing.

DNA extraction was performed from the 0.22-µm filter (free-living bacteria) following

the MagAttract Purification Kit protocol (QUIAGEN). Metagenomes were sequenced

using PacBio Sequel II (one 8M SMRT Cell Run, 30-h movie) (Novogene, South

Korea). To improve the quality of the PacBio raw reads, we generated Highly

Accurate Single-Molecule Consensus Reads (CCS reads) using the CCS v6

program of the SMRT-link package. The minimum number of full-length subreads

required to generate a CCS read was set to 15 (> 99.99 % base call accuracy).

ITS and 23S phylogenies of Pelagibacterales genomes and PacBio CCS reads.
PacBio CCS15 reads >5 Kb were screened to detect 16S and 23S rRNA genes using barrnap. Using SILVA (Quast et al., 2013), sequences affiliated with Pelagibacterales were kept and complete internal transcribed spacers (ITS) and 23S rRNA genes were extracted for further analysis. Phylotype classifications based on the ITS and 23S rRNA gene were inferred using the neighbour-joining approach in MEGA11 (Tamura et al., 2021), with 1000 bootstraps and the Jukes-Cantor model of substitution. Phylotype assignment followed existing ITS and 23S nomenclatures (Brown et al., 2012; García-Martínez and Rodríguez-Valera, 2000; Ngugi and Stingl, 2012).

Rarefaction curves among O-antigen loci. We quantified the number of locus types, aka locus diversity, by applying an ecological modelling approach using iNEXT package in R (Hsieh et al., 2016). In this approach, each SAG or CCS read was considered ecological "sites" and the OBC-types were considered the observed "species" in those sites. The rarefaction curves were calculated by extrapolating our data to 1,000 SAGs and CCS reads for measuring at the genomospecies level, and to 10,000 CCS reads for the whole Pelagibacterales order.

ACKNOWLEDGEMENTS
This work was supported by grant "FLEX3GEN" PID2020-118052GB-I00 (cofounded with FEDER funds) from the Spanish Ministerio de Economía, Industria y Competitividad to F.R-V and M.L.-P. J.M.H.-M. was supported with a PhD fellowship from Margarita Salas program, cofounded by the Spanish Ministerio de Universidades and the European Union—Next-Generation EU (2021/PER/00020). C.M.-P. was supported by a Ph.D. fellowship from the Spanish Ministerio de Ciencia

549 e Innovación (PRE2021-098122). We are grateful to Ramunas Stepanauskas for

550 suggestions related to SAGs and comments on a draft of the manuscript.

AUTHOR CONTRIBUTIONS

553 F.R-V conceived the study. J.M.H-M, C.M-P, and M.L-P collected and processed the

554 metagenomic samples. J.M.H-M, M.L-P, and F.R-V analyzed the data. J.M.H-M and

555 F.R-V contributed to write the manuscript.

CONFLICT OF INTERESTS

558 The authors declare that they have no competing interests.

DATA AVAILABILITY

561 Metagenomic datasets have been submitted to NCBI SRA and are available under

562 BioProject accession number PRJNA1088973 (PacBio CCS15 reads: MedWinter-

563 FEB2022-CCS [SAMN40517308], Med-OCT2021-15m-CCS [SAMN40517305],

564 Med-SEP2022-60m-CCS [SAMN40517307] and Med-OCT2021-75m-CCS

565 [SAMN40517306]).

REFERENCES

572 Abedon ST. 2022. Frequency-dependent selection in Light of Phage Exposure In: Abedon
573  ST, editor. Bacteriophages as Drivers of Evolution: An Evolutionary Ecological
574  Perspective. Cham: Springer International Publishing. pp. 275–292. doi:10.1007/978-
575  3-030-94309-7_24

Avrani S, Wurtzel O, Sharon I, Sorek R, Lindell D. 2011. Genomic island variability facilitates
Prochlorococcus–virus coexistence. *Nature* **474**:604–608. doi:10.1038/nature10172
Barco RA, Garrity GM, Scott JJ, Amend JP, Nealson KH, Emerson D. 2020. A Genus
Definition for Bacteria and Archaea Based on a Standard Genome Relatedness
Index. *mBio* **11**:10.1128/mbio.02475-19. doi:10.1128/mbio.02475-19
Berube PM, Biller SJ, Hackl T, Hogle SL, Satinsky BM, Becker JW, Braakman R, Collins SB,
Kelly L, Berta-Thompson J, Coe A, Bergauer K, Bouman HA, Browning TJ, De Corte
D, Hassler C, Hulata Y, Jacquot JE, Maas EW, Reinthaler T, Sintes E, Yokokawa T,
Lindell D, Stepanauskas R, Chisholm SW. 2018. Data descriptor: Single cell
genomes of Prochlorococcus, Synechococcus, and sympatric microbes from diverse
marine environments. *Sci Data* **5**:1–11. doi:10.1038/sdata.2018.154
Bolaños LM, Tait K, Somerfield PJ, Parsons RJ, Giovannoni SJ, Smyth T, Temperton B.
2022. Influence of short and long term processes on SAR11 communities in open
ocean and coastal systems. *ISME Commun* **2**:1–11. doi:10.1038/s43705-022-00198-
1
Brockhurst MA, Chapman T, King KC, Mank JE, Paterson S, Hurst GDD. 2014. Running
with the Red Queen: the role of biotic conflicts in evolution. *Proc R Soc B Biol Sci*
281:20141382. doi:10.1098/rspb.2014.1382
Brown MV, Fuhrman JA. 2005. Marine bacterial microdiversity as revealed by internal
transcribed spacer analysis. *Aquat Microb Ecol* **41**:15.
Brown MV, Lauro FM, DeMaere MZ, Muir L, Wilkins D, Thomas T, Riddle MJ, Fuhrman JA,
Andrews-Pfannkoch C, Hoffman JM, McQuaid JB, Allen A, Rintoul SR, Cavicchioli R.
2012. Global biogeography of SAR11 marine bacteria. *Mol Syst Biol* **8**:595.
doi:10.1038/msb.2012.28
Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND.
Nat Methods **12**:59–60. doi:10.1038/nmeth.3176
Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM. 2014. Rarefaction
and extrapolation with Hill numbers: a framework for sampling and estimation in
species diversity studies. *Ecol Monogr* **84**:45–67. doi:10.1890/13-0133.1
Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. *Bioinformatics* **36**:1925–1927.
doi:10.1093/bioinformatics/btz848
Coleman ML, Sullivan MB, Martiny AC, Steglich C, Barry K, Delong EF, Chisholm SW. 2006.
Genomic Islands and the Ecology and Evolution of Prochlorococcus. *Science*
311:1768–1770. doi:10.1126/science.1122050
Comeron JM. 1995. A method for estimating the numbers of synonymous and
nonsynonymous substitutions per site. *J Mol Evol* **41**:1152–1159.
doi:10.1007/BF00173196
Costea PI, Coelho LP, Sunagawa S, Munch R, Huerta-Cepas J, Forslund K, Hildebrand F,
Kushugulova A, Zeller G, Bork P. 2017. Subspecies in the global human gut
microbiome. *Mol Syst Biol* **13**:960. doi:10.15252/msb.20177589
da Silva Filho AC, Raittz RT, dos Santos-Weiss ICR. 2018. Comparative Analysis of
Genomic Island Prediction Tools. *Front Genet* **9**. doi:10.3389/fgene.2018.00619
Delmont TO, Kiefl E, Kilinc O, Esen OC, Uysal I, Rappé MS, Giovannoni S, Eren AM. 2019.
Single-amino acid variants reveal evolutionary processes that shape the
biogeography of a global SAR11 subclade. *eLife* **8**. doi:10.7554/elife.46497
Dittmar T, Lennartz ST, Buck-Wiese H, Hansell DA, Santinelli C, Vanni C, Blasius B,
Hehemann J-H. 2021. Enigmatic persistence of dissolved organic matter in the
ocean. *Nat Rev Earth Environ* **2**:570–583. doi:10.1038/s43017-021-00183-7
Drost H-G, Gabel A, Grosse I, Quint M. 2015. Evidence for Active Maintenance of
Phylotranscriptomic Hourglass Patterns in Animal and Plant Embryogenesis. *Mol Biol*
Evol **32**:1221–1231. doi:10.1093/molbev/msv012
Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Comput Biol* **7**:e1002195.
doi:10.1371/journal.pcbi.1002195
García-Martínez J, Rodríguez-Valera F. 2000. Microdiversity of uncultured marine

prokaryotes: the SAR11 cluster and the marine Archaea of Group I. *Mol Ecol* **9**:935–
48. doi:10.1046/j.1365-294x.2000.00953.x
Giovannoni SJ. 2017. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Annu*
Rev Mar Sci **9**:231–255. doi:10.1146/annurev-marine-010814-015934
Giovannoni SJ, Tripp HJ, Givan S, Podar M, Vergin KL, Baptista D, Bibbs L, Eads J,
Richardson TH, Noordewier M, Rappé MS, Short JM, Carrington JC, Mathur EJ.
2005. Genome streamlining in a cosmopolitan oceanic bacterium. *Science*
309:1242–5. doi:10.1126/science.1114057
Gokhale CS, Iwasa Y, Nowak MA, Traulsen A. 2009. The pace of evolution across fitness
valleys. *J Theor Biol* **259**:613–620. doi:10.1016/j.jtbi.2009.04.011
Grote J, Thrash JC, Huggett MJ, Cameron Thrash J, Huggett MJ, Landry ZC, Carini P,
Giovannoni SJ, Rappé MS. 2012. Streamlining and Core Genome Conservation
among Highly Divergent Members of the SAR11 Clade. *mBio* **3**:1–13.
doi:10.1128/mBio.00252-12.Editor
Haft DH, Loftus BJ, Richardson DL, Yang F, Eisen JA, Paulsen IT, White O. 2001.
TIGRFAMs: a protein family resource for the functional identification of proteins.
Nucleic Acids Res **29**:41–43. doi:10.1093/nar/29.1.41
Haro-Moreno JM, López-Pérez M, de la Torre JR, Picazo A, Camacho A, Rodriguez-Valera
F. 2018. Fine metagenomic profile of the Mediterranean stratified and mixed water
columns revealed by assembly and recruitment. *Microbiome* **6**:128.
doi:10.1186/s40168-018-0513-5
Haro-Moreno JM, López-Pérez M, Rodriguez-Valera F. 2021. Enhanced Recovery of
Microbial Genes and Genomes From a Marine Water Column Using Long-Read
Metagenomics. *Front Microbiol* **12**:708782. doi:10.3389/fmicb.2021.708782
Haro-Moreno JM, Rodriguez-Valera F, López-Pérez M. 2019. Prokaryotic Population
Dynamics and Viral Predation in a Marine Succession Experiment Using
Metagenomics. *Front Microbiol* **10**:2926. doi:10.3389/fmicb.2019.02926
Haro-Moreno JM, Rodriguez-Valera F, Rosselli R, Martinez-Hernandez F, Roda-Garcia JJ,
Gomez ML, Fornas O, Martinez-Garcia M, López-Pérez M. 2020. Ecogenomics of
the SAR11 clade. *Environ Microbiol* **22**:1748–1763. doi:10.1111/1462-2920.14896
Holt KE, Lassalle F, Wyres KL, Wick R, Mostowy RJ. 2020. Diversity and evolution of
surface polysaccharide synthesis loci in Enterobacteriales. *ISME J* **14**:1713–1730.
doi:10.1038/s41396-020-0628-0
Hsieh TC, Ma KH, Chao A. 2016. iNEXT: an R package for rarefaction and extrapolation of
species diversity (Hill numbers). *Methods Ecol Evol* **7**:1451–1456. doi:10.1111/2041-
210X.12613
Hu D, Fuller NR, Caterson ID, Holmes AJ, Reeves PR. 2022. Single-gene long-read
sequencing illuminates *Escherichia coli* strain dynamics in the human intestinal
microbiome. *Cell Rep* **38**:110239. doi:10.1016/j.celrep.2021.110239
Huang Y, Sheth RU, Zhao S, Cohen LA, Dabaghi K, Moody T, Sun Y, Ricaurte D,
Richardson M, Velez-Cortes F, Blazejewski T, Kaufman A, Ronda C, Wang HH.
2023. High-throughput microbial culturomics using automation and machine learning.
Nat Biotechnol **41**:1424–1433. doi:10.1038/s41587-023-01674-2
Huszczynski SM, Lam JS, Khursigara CM. 2019. The Role of Pseudomonas aeruginosa
Lipopolysaccharide in Bacterial Pathogenesis and Physiology. *Pathog 2020 Vol 9*
Page 6 **9**:6. doi:10.3390/PATHOGENS9010006
Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:
prokaryotic gene recognition and translation initiation site identification. *BMC*
Bioinformatics **11**:119. doi:10.1186/1471-2105-11-119
Kalynych S, Morona R, Cygler M. 2014. Progress in understanding the assembly process of
bacterial O-antigen. *FEMS Microbiol Rev* **38**:1048–1065. doi:10.1111/1574-
6976.12070
Kashtan N, Roggensack SE, Rodrigue S, Thompson JW, Biller SJ, Coe A, Ding H, Marttinen
P, Malmstrom RR, Stocker R, Follows MJ, Stepanauskas R, Chisholm SW. 2014.
Single-Cell Genomics Reveals Hundreds of Coexisting Subpopulations in Wild

Prochlorococcus. Science **344**:416–420. doi:10.1126/science.1248575

Kauffmann F. 1947. The serology of the coli group. *J Immunol Baltim Md 1950* **57**:71–100.

Kenyon JJ, Cunneen MM, Reeves PR. 2017. Genetics and evolution of Yersinia
pseudotuberculosis O-specific polysaccharides: A novel pattern of O-antigen
diversity. *FEMS Microbiol Rev* **41**:200–217. doi:10.1093/femsre/fux002

Konstantinidis KT, Tiedje JM. 2005. Genomic insights that advance the species definition for
prokaryotes. *Proc Natl Acad Sci U S A* **102**:2567–72. doi:10.1073/pnas.0409727102

Layoun P, López-Pérez M, Haro-Moreno JM, Haber M, Thrash JC, Henson MW, Kavagutti
VS, Ghai R, Salcher MM. 2024. Flexible genomic island conservation across
freshwater and marine Methylophilaceae. *ISME J* **18**:wrad036.
doi:10.1093/ismejo/wrad036

Lerouge I, Vanderleyden J. 2002. O-antigen structural variation: mechanisms and possible
roles in animal/plant–microbe interactions. *FEMS Microbiol Rev* **26**:17–47.
doi:10.1111/j.1574-6976.2002.tb00597.x

Letarov AV. 2023. Bacterial Virus Forcing of Bacterial O-Antigen Shields: Lessons from
Coliphages. *Int J Mol Sci*. doi:10.3390/ijms242417390

Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**:W242–W245.
doi:10.1093/nar/gkw290

Liu B, Furevi A, Perepelov AV, Guo X, Cao H, Wang Q, Reeves PR, Knirel YA, Wang L,
Widmalm G. 2019. Structure and genetics of Escherichia coli O antigens. *FEMS*
Microbiol Rev **44**:655–683. doi:10.1093/femsre/fuz028

Liu B, Knirel YA, Feng L, Perepelov AV, Senchenkova SN, Reeves PR, Wang L. 2014.
Structural diversity in Salmonella O antigens and its genetic basis. *FEMS Microbiol*
Rev **38**:56–89. doi:10.1111/1574-6976.12034

Lively CM. 2010. A Review of Red Queen Models for the Persistence of Obligate Sexual
Reproduction. *J Hered* **101**:S13–S20. doi:10.1093/jhered/esq010

López-Pérez M, Haro-Moreno JM, Coutinho FH, Martinez-Garcia M, Rodriguez-Valera F.
2020a. The Evolutionary Success of the Marine Bacterium SAR11 Analyzed through
a Metagenomic Perspective. *mSystems* **5**. doi:10.1128/mSystems.00605-20

López-Pérez M, Haro-Moreno JM, Gonzalez-Serrano R, Parras-Moltó M, Rodriguez-Valera
F. 2017. Genome diversity of marine phages recovered from Mediterranean
metagenomes: Size matters. *PLoS Genet* **13**:e1007018.
doi:10.1371/journal.pgen.1007018

López-Pérez M, Haro-Moreno JM, Iranzo J, Rodriguez-Valera F. 2020b. Genomes of the
"Candidatus Actinomarinales" Order: Highly Streamlined Marine Epipelagic
Actinobacteria. *mSystems* **5**. doi:10.1128/mSystems.01041-20

López-Pérez M, Martin-Cuadrado AB, Rodriguez-Valera F. 2014. Homologous
recombination is involved in the diversity of replacement flexible genomic Islands in
aquatic prokaryotes. *Front Genet* **5**:1–1. doi:10.3389/fgene.2014.00147

López-Pérez M, Rodriguez-Valera F. 2016. Pangenome evolution in themarine bacterium
alteromonas. *Genome Biol Evol* **8**:1556–1570. doi:10.1093/gbe/evw098

Lowe TM, Eddy SR. 1996. TRNAscan-SE: A program for improved detection of transfer RNA
genes in genomic sequence. *Nucleic Acids Res* **25**:955–964.
doi:10.1093/nar/25.5.0955

Martin-Cuadrado AB, Pašić L, Rodriguez-Valera F. 2015. Diversity of the cell-wall associated
genomic island of the archaeon Haloquadratum walsbyi. *BMC Genomics* **16**:603.
doi:10.1186/s12864-015-1794-8

Martínez de Tejada G, Pizarro-Cerdá J, Moreno E, Moriyón I. 1995. The outer membranes
of Brucella spp. are resistant to bactericidal cationic peptides. *Infect Immun* **63**:3054–
3061. doi:10.1128/iai.63.8.3054-3061.1995

Molina-Pardines C, Haro-Moreno JM, López-Pérez M. 2023. Phosphate-related genomic
islands as drivers of environmental adaptation in the streamlined marine
alphaproteobacterial HIMB59. *mSystems* **8**:e00898-23.
doi:10.1128/msystems.00898-23

Mostowy RJ, Holt KE. 2018. Diversity-Generating Machines: Genetics of Bacterial Sugar-
Coating. *Trends Microbiol* **26**:1008–1021. doi:10.1016/j.tim.2018.06.006
Neuenschwander SM, Ghai R, Pernthaler J, Salcher MM. 2018. Microdiversification in
genome-streamlined ubiquitous freshwater Actinobacteria. *ISME J* **12**:185–198.
doi:10.1038/ismej.2017.156
Ngugi DK, Stingl U. 2012. Combined Analyses of the ITS Loci and the Corresponding 16S
rRNA Genes Reveal High Micro- and Macrodiversity of SAR11 Populations in the
Red Sea. *PLoS ONE* **7**:e50274. doi:10.1371/journal.pone.0050274
Pachiadaki MG, Brown JM, Brown J, Bezuidt O, Berube PM, Biller SJ, Poulton NJ, Burkart
MD, La Clair JJ, Chisholm SW, Stepanauskas R. 2019. Charting the Complexity of
the Marine Microbiome through Single-Cell Genomics. *Cell* **179**:1623-1635.e11.
doi:10.1016/j.cell.2019.11.017
Pagnout C, Sohm B, Razafitianamaharavo A, Caillet C, Offroy M, Leduc M, Gendre H,
Jomini S, Beaussart A, Bauda P, Duval JFL. 2019. Pleiotropic effects of rfa-gene
mutations on Escherichia coli envelope properties. *Sci Rep* **9**:9696.
doi:10.1038/s41598-019-46100-3
Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz
P. 2018. A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat Biotechnol* **36**:996. doi:10.1038/nbt.4229
Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**:1043–55. doi:10.1101/gr.186072.114
Perlovsky LI, Frank-Kamenetskii MD. 2002. Statistical Limitations on Molecular Evolution. *J
Biomol Struct Dyn* **19**:1031–1043. doi:10.1080/07391102.2002.10506806
Pernitzsch SR, Alzheimer M, Bremer BU, Robbe-Saule M, De Reuse H, Sharma CM. 2021.
Small RNA mediated gradual control of lipopolysaccharide biosynthesis affects
antibiotic resistance in Helicobacter pylori. *Nat Commun* **12**:4433.
doi:10.1038/s41467-021-24689-2
Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.
2013. The SILVA ribosomal RNA gene database project: Improved data processing
and web-based tools. *Nucleic Acids Res* **41**:D590–D596. doi:10.1093/nar/gks1219
R Development Core Team R, Team RDC. 2011. R: A Language and Environment for
Statistical Computing. *R Found Stat Comput*, R Foundation for Statistical Computing.
doi:10.1007/978-3-540-74686-7
Raetz CRH, Whitfield C. 2002. Lipopolysaccharide Endotoxins. *Annu Rev Biochem* **71**:635–
700. doi:10.1146/annurev.biochem.71.110601.135414
Reeves P. 1995. Role of O-antigen variation in the immune response. *Trends Microbiol*
3:381–386. doi:10.1016/S0966-842X(00)88983-0
Richter M, Rossello-Mora R. 2009. Shifting the genomic gold standard for the prokaryotic
species definition. *Proc Natl Acad Sci* **106**:19126–19131.
doi:10.1073/pnas.0906412106
Rocap G, Distel DL, Waterbury JB, Chisholm SW. 2002. Resolution of Prochlorococcus and
Synechococcus Ecotypes by Using 16S-23S Ribosomal DNA Internal Transcribed
Spacer Sequences. *Appl Environ Microbiol* **68**:1180–1191.
doi:10.1128/AEM.68.3.1180-1191.2002
Roda-Garcia JJ, Haro-Moreno JM, Rodriguez-Valera F, Almagro-Moreno S, López-Pérez M.
2023. Single-amplified genomes reveal most streamlined free-living marine bacteria.
Environ Microbiol **25**:1136–1154. doi:10.1111/1462-2920.16348
Rodriguez-Valera F, Martin-Cuadrado A-B, López-Pérez M. 2016. Flexible genomic islands
as drivers of genome evolution. *Curr Opin Microbiol* **31**:154–160.
doi:10.1016/j.mib.2016.03.014
Rodriguez-Valera F, Martin-Cuadrado A-B, Rodriguez-Brito B, Pasić L, Thingstad TF,
Rohwer F, Mira A. 2009. Explaining microbial population genomics through phage
predation. *Nat Rev Microbiol* **7**:828–36. doi:10.1038/nrmicro2235
Rostøl JT, Marraffini L. 2019. (Ph)ighting Phages: How Bacteria Resist Their Parasites. *Cell*

Host Microbe **25**:184–194. doi:10.1016/j.chom.2019.01.009

Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, Yooseph S, Wu D, Eisen
JA, Hoffman JM, Remington K, Beeson K, Tran B, Smith H, Baden-Tillson H, Stewart
C, Thorpe J, Freeman J, Andrews-Pfannkoch C, Venter JE, Li K, Kravitz S,
Heidelberg JF, Utterback T, Rogers Y-H, Falcón LI, Souza V, Bonilla-Rosso G,
Eguiarte LE, Karl DM, Sathyendranath S, Platt T, Bermingham E, Gallardo V,
Tamayo-Castillo G, Ferrari MR, Strausberg RL, Nealson K, Friedman R, Frazier M,
Venter JC. 2007. The Sorcerer II Global Ocean Sampling expedition: northwest
Atlantic through eastern tropical Pacific. *PLoS Biol* **5**:e77.
doi:10.1371/journal.pbio.0050077

Samuel G, Reeves P. 2003. Biosynthesis of O-antigens: Genes and pathways involved in
nucleotide sugar precursor synthesis and O-antigen assembly. *Carbohydr Res*
338:2503–2519. doi:10.1016/j.carres.2003.07.009

Schwartz DA, Lindell D. 2017. Genetic hurdles limit the arms race between Prochlorococcus
and the T7-like podoviruses infecting them. *ISME J* **11**:1836–1851.
doi:10.1038/ismej.2017.47

Segata N, Börnigen D, Morgan XC, Huttenhower C. 2013. PhyloPhlAn is a new method for
improved phylogenetic and taxonomic placement of microbes. *Nat Commun* **4**:2304.
doi:10.1038/ncomms3304

Sharon I, Morowitz MJ, Thomas BC, Costello EK, Relman DA, Banfield JF. 2013. Time
series community genomics analysis reveals rapid shifts in bacterial species, strains,
and phage during infant gut colonization. *Genome Res* **23**:111–120.
doi:10.1101/gr.142315.112

Shibl AA, Thompson LR, Ngugi DK, Stingl U. 2014. Distribution and diversity of
Prochlorococcus ecotypes in the Red Sea. *FEMS Microbiol Lett* **356**:118–126.
doi:10.1111/1574-6968.12490

Sintes E, del Giorgio PA. 2014. Feedbacks between protistan single-cell activity and
bacterial physiological structure reinforce the predator/prey link in microbial
foodwebs. *Front Microbiol* **5**. doi:10.3389/fmicb.2014.00453

Tamura K, Stecher G, Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. *Mol Biol Evol* **38**:3022–3027. doi:10.1093/molbev/msab120

Tatusov RL, Natale DA, Garkavtsev IV, Tatusova TA, Shankavaram UT, Rao BS, Kiryutin B,
Galperin MY, Fedorova ND, Koonin EV. 2001. The COG database: new
developments in phylogenetic classification of proteins from complete genomes.
Nucleic Acids Res **29**:22–8. doi:10.1093/nar/29.1.22

Thompson LR, Haroon MF, Shibl AA, Cahill MJ, Ngugi DK, Williams GJ, Morton JT, Knight
R, Goodwin KD, Stingl U. 2019. Red Sea SAR11 and Prochlorococcus single-cell
genomes reflect globally distributed pangenomes. *Appl Environ Microbiol* **85**.
doi:10.1128/AEM.00369-19

Thrash J.C, Temperton B, Swan BK, Landry ZC, Woyke T, DeLong EF, Stepanauskas R,
Giovannoni SJ, Thrash J Cameron, Temperton B, Swan BK, Landry ZC, Woyke T,
DeLong EF, Stepanauskas R, Giovannoni SJ. 2014. Single-cell enabled comparative
genomics of a deep ocean SAR11 bathytype. *ISME J* **8**:1440–1451.
doi:10.1038/ismej.2013.243

Truong DT, Tett A, Pasolli E, Huttenhower C, Segata N. 2017. Microbial strain-level
population structure & genetic diversity from metagenomes. *Genome Res* **27**:626–
638. doi:10.1101/gr.216242.116

Tsementzi D, Wu J, Deutsch S, Nath S, Rodriguez-R LM, Burns AS, Ranjan P, Sarode N,
Malmstrom RR, Padilla CC, Stone BK, Bristow LA, Larsen M, Glass JB, Thamdrup B,
Woyke T, Konstantinidis KT, Stewart FJ. 2016. SAR11 bacteria linked to ocean
anoxia and nitrogen loss. *Nature* **536**:179–183. doi:10.1038/nature19068

Viklund J, Martijn J, Ettema TJG, Andersson SGE. 2013. Comparative and phylogenomic
evidence that the alphaproteobacterium HIMB59 is not a member of the oceanic
SAR11 clade. *PLoS ONE* **8**. doi:10.1371/journal.pone.0078858

Viver T, Conrad RE, Rodriguez-R LM, Ramírez AS, Venter SN, Rocha-Cárdenas J, Llabrés

M, Amann R, Konstantinidis KT, Rossello-Mora R. 2024. Towards estimating the
number of strains that make up a natural bacterial population. *Nat Commun* **15**:544.
doi:10.1038/s41467-023-44622-z
Wilhelm LJ, Tripp HJ, Givan SA, Smith DP, Giovannoni SJ. 2007. Natural variation in SAR11
marine bacterioplankton genomes inferred from metagenomic data. *Biol Direct* **2**:1–
19. doi:10.1186/1745-6150-2-27
Zaragoza-Solas A, Haro-Moreno JM, Rodriguez-Valera F, López-Pérez M. 2022. Long-Read
Metagenomics Improves the Recovery of Viral Diversity from Complex Natural
Marine Samples. *mSystems* **7**:e0019222. doi:10.1128/msystems.00192-22
Zark M, Christoffers J, Dittmar T. 2017. Molecular properties of deep-sea dissolved organic
matter are predictable by the central limit theorem: Evidence from tandem FT-ICR-
MS. *Mar Chem* **191**:9–15. doi:10.1016/j.marchem.2017.02.005
Zhao Z, Amano C, Reinthaler T, Orellana MV, Herndl GJ. 2024. Substrate uptake patterns
shape niche separation in marine prokaryotic microbiome. *Sci Adv* **10**:eadn5143.
doi:10.1126/sciadv.adn5143
866
867

885 **Table 1.** Taxonomic classification of PacBio CCS reads containing a 23S rRNA gene

886 (hence an OBC) and classified to the order Pelagibacterales.

	Total OBCs	*495*	*458*	*642*	*533*	*652*
Subclade	*Genomospecies*	**Med-OCT2021-15m**	**Med-OCT2021-75m**	**Med-SEP2022-60m**	**MedWinter-JAN2019**	**MedWinter-FEB2022**
Ia	*Ia.1/I*	5	4	14	5	4
	Ia.3/VII (gMED)	**127**	**32**	**65**	**134**	**153***
	Ia.3/VII (gWID)	22	9	8	11	17
	Ia.3/Unclassifed	70	38	168	132	115
	Ia.4/I	1	4	1	0	1
	Ia.4/II	0	0	0	1	0
	Ia.4/III	17	5	16	16	16
	Ia.4/Unclassified	79	45	93	79	95
	Ia.Unclassified	14	80	45	16	20
Ib	*Ib.1*	5	1	2	1	2
	Ib.2	17	6	16	17	27
	Ib.3	0	0	0	0	0
	Ib.4	7	3	10	12	16
	Ib.5	22	8	23	9	12
	Ib.Unclassified	10	44	19	15	17

Ic	*Ic.1*	0	0	0	0	0
	Ic.2	**5**	**112***	**29**	**16**	**10**
IIa	-	90	61	127	66	140
IIb	-	0	5	3	2	4
IIIa	*IIIa.1*	0	0	2	0	3

*CCS reads used for rarefaction and extrapolation curves.

FIGURE LEGENDS

Figure 1. A. Cladogram-based classification of the 806 genomes that shared OBC-

type, (the complete cladogram is shown in **Figure S6**). The outer rings represent the

scale of the cladogram, measured as AAI among genomes (hierarchical clustering,

average linkage). Red and blue circles mark the Pelagibacterales limits for species

suggested in reference (Konstantinidis and Tiedje, 2005) (85% ANI) and the more

common standard 95% respectively (at these levels of similarity AAI and ANI are

nearly equal). The most relevant species names are used from reference (Haro-

Moreno et al., 2020) and labeled at the corresponding branching line. Inner

connections (>90 % shared genes, dashed line, >99 % – continuous line) indicate

shared OBC-types between genomes. Connector colors represent the average

identity values of the OBC-shared genes. The major Pelagibacterales subclades are

shaded: Ia – blue, Ib – green, II – purple, and IIIa – yellow. A purple dot near the

genome name indicates that it comes from a BATS single sample. **B.** Variation of the

number of different genes for the same OBC-type between pairs of genomes,

expressed as the genomic distance (100 - AAI, %). **C.** Frequency of sampling the

same OBC-type as a function of the phylogenetic distance between a pair of

genomes.

Figure 2. A. Scatterplot of the relationship between genome distance and locus

distance, both expressed as 100 - AAI (%), for only those genomes sharing an OBC-

type. The dashed line in the scatterplot represents the linear regression line. The

red-shaded area indicates recent horizontal gene transfer events (genome distance

is at least twice the OBC distance). Dots colored by taxonomy. **B.** Scatterplot of the

relationship between the dN/dS values for OBCs and genomes. Red dashed line

represents the regression line, whereas the black dashed line indicates the y=x.

Figure 3. Rarefaction (solid line) and extrapolation (dashed line) curves based on

OBC diversity against the number of sequences from **A.** All SAGs (black), and SAGs

coming from the single BATS sample, considering all sequences together (red line)

or only the four most representative genomospecies (remaining colored lines); **B.**

PacBio Sequel II metagenomic reads from a set of Mediterranean samples for two

abundant genomospecies, gMED in MedWinter-FEB2022 (purple line), and Ic.2 in

Med-OCT2021-75m (yellow line). The bottom plots represent how well the number of

detected and predicted sequences covered the diversity of OBCs. The dispersion

area shows a 95 % confidence interval.

SUPPORTING INFORMATION

Figure S1. A. Upper pie chart indicates the total number of Pelagibacterales

genomes screened (pale color), and the number of genomes on which we could

identify the O-chain biosynthetic gene cluster (OBC) (black area). The bottom pie

chart distributes the 806 OBCs according to their completeness: A – complete OBC;

B – the boundaries of the OBC, i.e. the 23S rRNA gene on the left-hand side and the

5S rRNA gene on the right-hand side, were detected, but in two different contigs

from the same genome; C – only the left-hand side; D – only the right-hand side. **B.**

Number of OBCs recovered by oceanic region. **C.** Taxonomic classification of

Pelagibacterales-containing OBCs based on a maximum-likelihood phylogenetic tree

from shared proteins (see methods). The resulting phylogenetic groups follow the

nomenclature described in Haro-Moreno et. al., 2020 (Haro-Moreno et al., 2020).

Figure S2. A. Genomic comparison of four selected complete OBCs from members

of the Ib.4 phylogroup. Note that in this case, the glycosylation island is located

between the 16S-ITS-23S rRNA genes and the tRNA-Val, tRNA-Met genes, and the

core genes involved in the peptidoglycan biosynthesis. **B.** Reconstruction of a partial

genome, Ib4-rB, in a single contig after the co-assembly of 3 nearly identical (>99 %

ANI) SAGs. The locations of the *dna*A, 16S, 23S, and 5S rRNA genes are indicated.

The metagenomic fragment recruitment of this genome in the Mediterranean Sea

(Med-SEP2014-60m) confirmed the location of the glycosylation island.

Figure S3. Genomic properties of the 163 complete OBCs and their corresponding

genomes. Boxplot on the left summarises the median, first and third quartiles of the

length of the OBC, considering all sequences as a unit (leftmost boxplot) or divided

by subgroups. Numbers below the taxonomic name indicate the number of OBCs for

each group. Boxplots on the right show the difference in the GC content and the

intergenic spacer (upper and lower panels, respectively) between the OBC and its

genome. Stars indicate the p-value (*** p-value < 0.001, ** p-value < 0.01).

Figure S4. A. Histogram resulted from the all-vs-all comparison of the 806 OBCs at

50 % amino acid identity threshold. X-axis indicates the percentage of orthologous

genes (OF) of the shortest sequence (genes shared between two OBCs), in groups

of 5 % OF.

Figure S5. Examples of OBC-type sharing among Pelagibacterales genomes. OBCs

are aligned and AAI color-coded.

Figure S6. Cladogram-based classification of the 806 genomes containing an OBC,

represented as in **Figure 1.**

Figure S7. Number of OBCs found in one or several taxonomic groups (>85 % AAI

or >95% AAI).

Figure S8. Rarefaction (solid line) curves based on OBC diversity against the

number of sequences from a set of five PacBio Sequel II metagenomic reads

collected from the Mediterranean Sea.

Table S1. Summary of Pelagibacterales genomes (SAGs and isolates) with an O-

chain Biosynthetic gene Cluster (OBC).

Table S2. Summary of pairwise comparisons among O-chain Biosynthetic gene

clusters (OBCs).

A

Similarity (AAI, %)

IIIa IIa Ib

Ib.3
Ib.2
Ib.1

Ia.4/II

gWID
gMED

Ia

SAG Origin
● BATS (single sample)

Comparison among Islands (AAI)
─── ≥ 99 %
─── 95 - 99 %
─── 85 - 95 %
─── ≤ 85 %
─── ≥ 99 % conserved genes
╌╌╌ 90 - 99 % conserved genes

B

Number of different genes vs Genome Distance (%)

C

Frequency of sampling the same OBC (%)

Pairwise genome AAI (%)

Taxonomic group
● All SAGs
● Ia.3
● Ia.3/VII (gMED)
● Ia.3/V (gWID)
● Ia.4
● Ib.1
● Ib.2
● Ib.4