Relationship between weapon size and six key behavioural and physiological traits

in males of the European earwig

Samantha E.M. Blackwell[1], Laura Pasquier[1], Simon Dupont[1], Séverine Devers[1], Charlotte

Lécureuil[1], Joël Meunier[1*]

[1]Institut de Recherche sur la Biologie de l'Insecte (IRBI), UMR CNRS 7261, Université de Tours,

France.

ORCID numbers :

• L Pasquier : 0009-0000-1346-2069

• S Dupont : 0000-0003-4760-6811

• C Lécureuil : 0000-0002-7339-9185

• J Meunier : 0000-0001-6893-2064

*Corresponding author: J Meunier, joel.meunier@univ-tours.fr

SUMMARY

In many animals, male weapons are large and extravagant morphological structures that typically enhance fighting ability and reproductive success. It is generally assumed that growing and carrying large weapons is costly, thus only males in the best condition can afford it. In the European earwig, males carry weapons in the form of forceps-like cerci, which can vary widely in size within populations. While long forceps appear to increase male's access to females, it is unknown whether it also correlates with other important male life-history traits. This information is important, however, in determining the potential reliability of forceps length as an indicator of male quality and the stability of this signalling system. Here, we tested whether forceps length is associated with six important behavioural and physiological traits in males of the European earwig. We sampled hundreds of males from two populations, selected 60 males with the longest and shortest forceps from each population, and then measured locomotor performance, boldness, aggregation behaviour, survival under harsh conditions, sperm storage, and survival after pathogen exposure. Contrary to our predictions, we detected no main association between forceps length and the traits measured. This lack of association was consistent between the two populations, although there were population-specific levels of boldness, aggregation and survival in harsh conditions (for long-forceps males only). Overall, these results challenge our current understanding of the function and quality signal of forceps length in this species and raise questions about the evolutionary drivers that could explain the maintenance of weapon size diversity within and between populations.

Keywords: Behaviour; Insect; Metarhizium; Ornament; Sexual selection; Weapons

INTRODUCTION

Animal reproduction often requires males to engage in physical competition and courtship to attract females in search of mates (Davies et al., 2012). From vertebrates to arthropods, these two needs have often led to the evolution of male weapons and ornaments through sexual selection (Emlen, 2008; Goldberg et al., 2019; McCullough et al., 2016). These weapons and ornaments are typically large and extravagant morphological structures that can grow on different parts of the male's body, take a variety of forms (such as antlers, horns, spurs, fangs and tusks), and work to enhance the male's fighting ability and/or attractiveness to females (Emlen, 2008). Textbook examples of this enhancement can be found in the white-tailed deer *Odocoileus virginianus*, where males growing the largest antlers are more likely to win fights with other males and have higher annual breeding success (Newbolt et al., 2017), and in the stalk-eyed flies *Cyrtodiopsis whitei* and *C. dahnanni*, where females prefer to mate with males exhibiting the longest eye stalks (Wilkinson et al., 1998).

However, not all males display extravagant weapons or ornaments (Emlen, 2008; Goldberg et al., 2019; McCullough et al., 2016), as the development and maintenance of these sexually selected traits often comes at a cost to males. This cost can arise from the fact that carrying heavy, bulky weapons (or ornaments) makes males less mobile and more visible to both predators and prey (Oufiero & Garland, 2007). For example, males with experimentally-enlarged wing spots have lower survival rates due to increased conspicuity to both visually orienting predators and visually orienting prey in the rubyspot damselfly *Hetaerina americana* (Grether, 1997). The cost of carrying large morphological structures can also arise because it may impose investment trade-offs with physiological functions ranging from metabolism to muscle development and spermatogenesis,

which can be crucial for male fitness and survival (Emlen, 2001). For example, males with the

largest hind leg weapon pay the highest resting metabolic rate and energy costs in the Hemiptera

Leptoscelis tricolor (Somjee et al., 2018). Similarly, growing large mandibles come with low flight-

muscle mass in the stag beetle *Cyclommatus metallifer* (Mills et al., 2016), and males who invest

in extravagant sexual displays show a more rapid decline in spermatogenesis than males who

invest less in these displays in the houbara bustard *Chlamydotis undulata* (Preston et al., 2011). As

a result, only the males in the very best condition are expected to possibly afford the development

and maintenance of extravagant weapons and ornaments (Otte & Stayman, 1979; Emlen &

Nijhout, 2000). Determining the reliability of these morphological structures as indicators of the

male quality is therefore crucial to understanding the stability of these signalling systems

(Berglund et al., 1996).

Earwigs (Dermaptera) are textbook examples of insects with males displaying a sexually

selected weapon. In this taxonomical group, females have relatively short, straight and hardened

forceps-like cerci typically used to defend the clutch against predators (Meunier, 2024), whereas

males display elongated, curved and hardened forceps-like cerci (hereafter referred to as forceps)

involved in mating courtship and fights against other males (Briceño & Eberhard, 1995; Kamimura,

2014; Walker & Fell, 2001). There are several lines of evidence to suggest that long forceps provide

males with benefits in terms of mating success (Eberhard & Gutierrez, 1991; Kamimura, 2014;

Tomkins & Brown, 2004). For example, males with long forceps are more likely to win fights with

other males by squeezing them between their cerci, and thus gain better access to females in the

toothed earwig *Vostox apicedentatus* (Moore & Wilson, 1993). In the European earwig *Forficula*

auricularia, male forceps are also used in male-male contests as a weapon to deter competitors

prior to mating (Styrsky & Rhein, 1999) or to interrupt mating individuals by non-copulating males

(Forslund, 2000, 2003; Walker & Fell, 2001). Long-forceps males are also generally more

aggressive, more readily accepted by females for mating and copulate longer compared to short-

forceps males (reviewed in Kamimura, 2014). Although forceps are involved in male courtship

(Walker & Fell, 2001), females do not seem to select their mate exclusively on the basis of forceps

length (Forslund, 2000, 2003; Radesäter & Halldórsdóttir, 1993; Walker & Fell, 2001).

While the relationship between male forceps length and mating success has been well

documented in earwigs (reviewed in Kamimura, 2014), it remains unclear whether having long or

short forceps is associated with other important life-history traits in these males. However, this is

important information to determine the potential reliability of forceps length as an indicator of

male quality and the stability of this possible signalling system. The very few studies that have

addressed this question focused on male immunity in the European earwig *F. auricularia* (Körner

et al., 2017; Rantala et al., 2007). On the one hand, they show that males with long forceps have

lower basal levels of certain components of the immune system, such as lysozyme activity and

hemocyte concentrations. This suggests that having long forceps comes with an immune cost for

males. On the other hand, they show no association between forceps length and other

components of the immune response, such as encapsulation rate and phenoloxidase activity.

Taken together, these findings suggest that the immune costs of having long forceps may be

relatively limited and raise the question of whether other important life-history traits can be

associated with forceps length.

Here, we investigated whether long and short forceps are associated with six important

behavioural and physiological traits in males of the European earwig. We sampled hundreds of

males in two natural populations of earwigs separated by 400 km and in each population, we

selected the 30 males with the longest and the 30 males with the shortest forceps. We then

measured their level of expression of three important behaviours, their survival rate in two distinct

harsh conditions and their sperm quantity. The first behaviour was their locomotor performance

(Cheutin et al., 2024), which reflects the ability of males to walk long distances (to forage, hide or

find a mate) while carrying long and heavy (or short and light) appendages. The second behaviour

was their likelihood to flee after a physical disturbance (i.e., boldness), which shows how males

react when disturbed by a predator attack (Thesing et al., 2015). The third behaviour was their

propensity to aggregate with conspecifics. This is an important parameter in this gregarious

species, as adults typically live in groups of up to several hundred individuals and social isolation

can have detrimental effects on their physiology (Kohlmeier et al., 2016; Van Meyel & Meunier,

2022). We also measured their survival rate in a harsh condition where they were isolated with no

access to a food source for 31 days, and then their survival rate after exposure to the common

entomopathogenic fungus *Metarhizium brunneum* (Vogelweith et al., 2017). Finally, we measured

the level of sperm storage in seminal vesicles of each male, a parameter that is often important in

the context of male-male competition for mating (Shuker & Simmons, 2014). If forceps length is a

reliable signal of good quality, we predict that long forceps males would 1) have higher locomotor

performance, 2) be bolder and thus less likely to flee after a simulated attack, 3) be more social

and thus more likely to aggregate with conspecifics, 4) be more resilient to harsh environmental

conditions and thus have a higher survival rate after 31 days of starvation, and/or would 5) have

higher immune defences and thus survive better after a pathogen infection than short-forceps

males. We also predict that 6) long-forceps males would produce more sperm and contain more

sperm in their seminal vesicles than short forceps males. This last prediction would be consistent

with the longer duration of copulation reported for the long-forceps males (Kamimura, 2014),

even if this longer duration may also reflect other male mating strategies, such as mate guarding.

MATERIAL AND METHODS

Animal rearing and selection

In July 2022, we field sampled 3000 males and females in a cultivated peach orchard near Valence,

France (Lat 44.9772790, Long 4.9286990) and 400 males and females in a non-cultivated area at

the edge of a forest near Cinais, France (Lat 47.1606970, Long 0.1763663). All these individuals

belong to *Forficula auricularia* Linneaus, 1758, also called *Forficula auricularia* clade A (González-

Miguéns et al., 2020). Immediately after field sampling, we distributed them by population of

origin into 50 plastic terrariums (37 × 22 × 25 cm, balanced sex ratio) to homogenize nutrition,

habitation, and access to mates for the males (Körner et al., 2017). Three months later (i.e., at the

end of the reproductive season), we removed the females from all the terrariums to mimic their

natural dispersal. One month later, for each population, we visually selected the 30 males with the

longest forceps and the 30 males with the shortest forceps (Körner et al., 2017) and isolated them

in individual Petri dishes (diameter 5 cm) for later use in behavioural and physiological

measurements (A timeline of the experimental design and the detailed sample sizes can be found

in Figure 1). Two days after isolation, we confirmed the robustness of our two forceps categories

by measuring the mean of the left and right outer forceps of the 120 males. These measurements

were done to the nearest 0.001 mm using the Leica Application Suite 4.5 software (Leica

Microsystems, Wetzlar, Germany) on pictures taken under a binocular scope (Leica, MZ 12.5).

These measurements confirmed that there was no overlap between the two male categories

(Figure 2). All the remaining males and females that were not used in this experiment were

involved in other experiments that are not presented here.

From field sampling to isolation, we fed animals with artificial food containing mostly

pollen, carrots, and cat food (see details in Kramer et al. 2015). During these three months, we

offered this food *ad libitum* to homogenise the nutritional condition of the males when we started

our measurements. We then kept the isolated males without access to food from the day they

were isolated until 31 days after their isolation (Figure 1) to test whether resistance to both

starvation and social isolation (i.e., harsh environmental conditions) were population and/or

forceps-length specific, while ensuring that good rearing conditions did not mask any potential

investment trade-offs between forceps length and other life history traits. We then provided food

to males after day 31 to measure their survival following pathogen exposure. We kept all animals

on a 12h:12h light:dark schedule at 18°C, and both the terrarium and Petri dishes contained a layer

of moist sand.

Behavioural measurements

From the 5th to the 24th day after isolation, we measured three behaviours in the 120 isolated

males: the locomotor performance, the likelihood of fleeing after a physical disturbance, and the

level of aggregation. All these measurements followed standard protocols for earwigs (Honorio et

al., 2023; Merleau et al., 2022).

First, we measured male's locomotor performance 5 days after isolation. On that day, we

transferred each male to a 3D-printed circular arena (Open field; diameter = 8 cm, height 0.4 cm)

with a cover made of glass, placed on an infrared table and kept in complete darkness. We then

video recorded males' locomotion for 20 minutes (Camera: BASLER BCA 1300, Germany; Media

Recorder v4.0, Noldus Information Systems, Netherland) and subsequently analyzed the resulting

videos with the software Ethovision XT 16 (Noldus Information Systems, Netherland). We defined

the locomotor performance of each male as the total distance he walked (in cm) during the entire

recording (Honorio et al., 2023; Merleau et al., 2022).

Second, we measured the likelihood of fleeing after a physical disturbance 10 days after

isolation. On that day, we carefully opened each Petri dish, pricked the male on the pronotum with

a glass capillary and then recorded whether or not the male's first reaction was to move more

than one body length away from its initial position (i.e., flee).

Finally, we measured the level of aggregation of each male 24 days after isolation. We

placed each male in a 3D-printed arena used in Van Meyel & Meunier, 2022, consisting of four

linearly aligned circular chambers (diameter 4 cm). Three of the chambers were connected by 0.5

cm wide corridors allowing earwigs to move between chambers. The width of the corridor

connecting to the fourth (outer) chamber was reduced to 0.15 cm, which prevented earwig

movement while allowing the circulation of odours and antennal contacts between individuals on

both sides (Van Meyel & Meunier, 2022). We started the experiment by placing two naive males

and one female from the same population (but not involved in any other experiment or

measurement) in this isolation chamber, and the tested male in one of the connected chambers.

We recorded whether the tested male was in the chamber next to the group of conspecifics (yes

or no) and repeated this measurement by taking pictures every hour for 48 hours using infrared

cameras and the software Pylon Viewer v5.1.0 (Basler©, Ahrensburg, Germany). For each tested

male, we thus obtained an aggregation score ranging from 0 to 48, which was defined as the total

number of pictures in which a male was in the chamber next to the group of conspecifics. After

the 48h of aggregation test, we returned each male to its Petri dish until its use for subsequent

measurements (see below). For ease of handling, all manipulated earwigs were anaesthetised with

CO_2 during the set-up of this last measurement.

Sperm storage measurement

Of the 91 males still alive on day 31 after isolation (Figure 1), we used a random subset of 39 to

measure sperm storage as the number of sperm present in their seminal vesicles (Figure 1). This

counting occurred about two months after the males were separated from the females, which is

probably long enough for the males to rebuild their sperm reserves, regardless of their previous

mating rate. We counted sperm numbers following the protocol detailed in Damiens et al. (2002).

In brief, we dissected each male under a dissecting microscope, placed their seminal vesicles on a

slide with 15 µL of 1x Phosphate Buffered Saline and then pierced it to release all the sperm. We

subsequently dried the plate on a heating block, after which the smear of sperm was sealed with

a 70% ethanol solution and allowed to dry again at ambient temperature. The slides were then

stored at 3°C. Two days later, we deposited 20µL of DAPI dye (concentration = 10 µmol/L) on the

smear, covered it with a small piece of plastic wrap to allow the dye to infiltrate the cells while not

drying out, and 10 minutes later, we replaced the plastic wrap with a glass slide cover. The slide

was then viewed under the microscope at 20x magnification, and we took pictures of 5 different

fields of the smear. Using an Olympus micrometre calibration slide and the software Image J

(Schneider et al., 2012), we then counted the number of spermatozoa in all 5 fields for each male.

Finally, we used all these numbers to calculate the number of spermatozoa per millimetre square

and multiplied this number by the total area of the smear to obtain the total number of sperm per

male. It should be noted that sperm storage was measured in males that survived 31 days in

isolation without access to food, so that this value represents the sperm storage of individuals best

adapted to these two stressful conditions.

Survival in harsh environments and after exposure to pathogens

We measured male survival under two types of harsh conditions. The first type of harsh condition

was the absence of any food source (starvation) combined with social isolation, which is known to

have detrimental effects on this gregarious species (Kohlmeier et al., 2016; Van Meyel & Meunier,

2022). We assessed the male survival rate under these conditions by recording whether each of

the 120 males tested was still alive on day 31 after isolation (Figure 1).

The second type of harsh condition was exposure to pathogens. We measured survival rate

after pathogen exposure in the 52 males that were still alive on day 31 and were not used to

measure sperm storage (Figure 1). We exposed each male to spores of the entomopathogenic

fungus *Metharizium brunneum* (formerly *M. anisopliae*). This fungus is a natural and lethal

pathogen of *F. auricularia* (Arcila & Meunier, 2020; Coulm & Meunier, 2021; Günther & Herter,

1974). The infection followed a standard protocol detailed in Kohlmeier et al. (2016). In brief, we

immersed each male in an Eppendorf tube containing 1.5 mL of a conidiospore suspension of *M.*

brunneum diluted in 0.05% Tween 20 (Sigma P-1379) at a concentration of 10^6 spores/mL. We

then gently swirled the tube from side to side for 4 seconds, removed the male and placed it back

in its original Petri dish with standard food (see above) that was changed twice a week. We then

kept the infected males at 20°C on a 12:12 light:dark schedule. We checked them daily for

mortality over the course of 45 days. As with sperm storage, survival after pathogen exposure was

measured in males that survived 31 days in isolation without access to food, so that it represents

the survival rate of individuals best adapted to two these stressful conditions.

Statistical analyses

We conducted all statistical analyses using the software R v4.3.2 (https://www.r-project.org/)

loaded with the packages *DHARMa* (Hartig, 2020), *car* (Fox & Weisberg, 2019), *survival* (Therneau,

2020) and *emmeans* (Lenth, 2022). We analysed locomotor performance, sperm count and

aggregation score using three general linear models (*lm* in R), while we analysed the likelihood to

flee (yes or no) and whether males were dead 31 days after isolation (yes or no) using two

generalized linear model (*glm* in R) with binomial error distributions. Finally, we analysed the

survival rate of the pathogen-exposed males using a Cox proportional hazard regression model

allowing for censored data (*Coxph* in R), i.e., males still alive 45 days after exposure to the

pathogen. In these six models, we entered the type of forceps length (long or short), the

population of origin of the males (Cinais or Valence) and the interaction between these two

variables as explanatory factors. Overall, we checked that all model assumptions were met (e.g.,

homoscedasticity and normality of residuals) using the *DHARMa* package. To this end, we log-

transformed male locomotor performance and log+1-transformed the aggregation scores. In the

model where we found a significant interaction, we conducted pairwise comparisons using the

estimated marginal means of the models and we corrected P values for multiple testing using the

Tukey method, as implemented in the *emmeans* R package. Finally, we conducted power analyses

for each of our statistical models using the packages *pscl* (Jackman et al., 2024), *pwr* (Champely et

al., 2020) and *powerSurvEpi* (Qiu et al., 2021).

RESULTS

Overall, we detected no main association between forceps length and the six traits measured

(Table 1). This applied to male locomotor performance, likelihood of fleeing after a physical

disturbance, aggregation score, sperm storage and survival after pathogen infection (Figure 3;

Table 1). There was also no difference between males with short and long forceps in terms of

survival under harsh environmental conditions, although males with long forceps survived better

when they came from Cinais compared to Valence (this trend was not present in males with short

forceps; Figure 3; interaction in Table 1; pairwise comparisons: Long forceps Cinais vs Valence: Z =

-3.311; P = 0.005; Short forceps Cinais vs Valence: Z = -0.611; P = 0.929). Regardless of forceps

length, males from Cinais were less likely to flee after a physical disturbance and less gregarious

than males from Valence (Table 1). These differences between the two populations were absent

for all other traits measured (Table 1). There was no interaction between the population of origin

and forceps length for any of the traits measured, except for survival under harsh environmental

conditions (Table 1). The statistical power of each analysis is shown in Table 1.

DISCUSSION

The display of large, extravagant weapons often comes at a cost to the males (Emlen, 2008; Goldberg et al., 2019; McCullough et al., 2016). It is therefore expected that only males in the best condition can afford them and thus that these large weapons signal good male quality (Otte & Stayman, 1979; Emlen & Nijhout, 2000). Here, we tested this prediction in the European earwig by investigating whether male forceps length is associated with six important behavioural and physiological traits. Contrary to predictions, our experiment did not allow us to detect an association between forceps length and locomotor performance, boldness (i.e., likelihood to flee after a physical disturbance), aggregation behaviour, sperm production, and male survival after pathogen infection. These findings were consistent between the two populations, although some of the traits measured were population specific: males from Cinais were generally bolder, less gregarious and (only if they carried long-forceps) had a better chance of surviving in harsh conditions than males from Valence.

Our results contrast with much of the literature reporting associations between sexually selected male attributes and physiological, behavioural or immunological traits in arthropods (Emlen, 2008; Goldberg et al., 2019; McCullough et al., 2016). However, the European earwig is not the only case where such associations appear to be lacking (Emlen, 2008; Swallow & Husak, 2011). For example, carrying giant claw is not associated with the efficiency of escape behavior and the level of metabolic costs in two fiddler crabs (Pena & Levinton, 2021; Tullis & Straube, 2017). Similarly, bearing large horns does not reflect male growth, mobility, or immunity in the rhinoceros beetle *Trypoxylus dichotomus* (McCullough & Emlen, 2013; McCullough & Tobalske, 2013). It has been suggested that this apparent lack of general association may be due to the fact

that the performance of each weapon size depend on the environment in which the weapons are

used and/or because weapon sizes reflect alternative reproductive tactics (McCullough et al.,

2016; McCullough & Emlen, 2013). This could be the case with the European earwig (Tomkins et

al., 2005). In this species, males with large forceps have been suggested to be more active in

guarding of females, while males with small forceps are more active in sneaking into copulations

(Kamimura, 2014; Tomkins & Brown, 2004). Having long forceps should therefore be rewarded

more frequently when the encounter rate between competitors is high (Hunt & Simmons, 2001),

such as in high population densities. Consistent with this prediction, Tomkins & Brown (2004)

found that the proportion of long forceps males increased with population densities across 46

island and mainland sites in the UK. Another possible explanation for our results is that the

association between male forceps length and the six traits measured could have been masked by

differences in the amount of resources available to each male and/or used by each male for these

traits (van Noordwijk & de Jong, 1986). In our study, all males were kept under identical laboratory

conditions and fed *ad libitum* for three months prior to our measurement. This explanation would

therefore suggest that the variation in resources influencing investment decisions is determined

prior to field sampling, e.g. during development or early adulthood. Previous data suggest that it

could be the case in the European earwig. In this species, early life conditions have long-term

effects on the physiology and behaviour of adults (Raveh et al., 2016; Thesing et al., 2015; Wong

& Kölliker, 2014) and inter-individual variation in female condition can mask investment trade-off

between egg quantity and quality (Koch & Meunier, 2014). Overall, our findings call for future

studies to confirm whether males with short or long forceps have alternative reproductive tactics,

whether their reproductive success depends on population densities and whether early life

conditions can affect the likelihood to detect the association between forceps length and the six

traits measured. They also call for further research to quantify other potential costs of carrying

long forceps in this species, for example in terms of predation rates and ability to fly (Crumb &

Eide, 1941). These notwithstanding, our findings emphasize that forceps size is not a good

predictor of the six behavioural and physiological traits measured in males.

While we found no overall difference between short- and long-forceps males, our data

reveal population differences in terms of males' boldness, aggregation level and resistance to

starvation. Cinais had males that were generally bolder, less gregarious, as well as males with long

forceps that survived food deprivation better than Valence. These effects are unlikely to reflect a

plastic response to the direct environment of the males tested (e.g. population differences in

terms of nutritional status), as they were all reared under common laboratory conditions for

months prior to the start of our experiments. Instead, it could reflect population idiosyncrasies

that have affected their development, such as climatic conditions (Valence is warmer than Cinais),

environmental conditions and/or exposure to phytosanitary products (e.g. Valence is a cultivated

orchard, whereas Cinais is an uncultivated forest edge), or population-specific genetic background.

In line with this hypothesis, several life history traits of the European earwig are shaped by the

duration of cold during egg development, the level of warm temperatures during nymph

development and, more generally, by seasonal parameters encountered by offspring during

development (Coulm & Meunier, 2021; Körner et al., 2018; Tourneur & Meunier, 2020). Similarly,

recent studies have shown that the European earwig can be sensitive to exposure to even low

levels of pesticides, which can lead to populations specificities in terms of earwig physiology,

morphology and behaviour (Fricaux et al., 2023; Le Navenant et al., 2019; Malagnoux, Capowiez,

et al., 2015; Malagnoux, Marliac, et al., 2015; Mauduit et al., 2021; Meunier et al., 2020).

Regardless of the nature of these population idiosyncrasies, our study shows that they do not

affect the (lack of) association between forceps length and the six traits measured.

Overall, our findings questions the robustness of our understanding of forceps length

diversity in terms of function, maintenance and use as a possible quality signal in the European

earwig (McCullough et al., 2016). To date, our knowledge of these questions is exclusively based

on laboratory experiments in which females were presented to one or two males (reviewed in

Kamimura, 2014). While the results of these studies suggest that males can gain fitness benefit

from growing longer forceps (Forslund, 2000, 2003; Radesäter & Halldórsdóttir, 1993; Styrsky &

Rhein, 1999; Walker & Fell, 2001), they contain two important limitations. First, they considered

mating success (i.e., success in gaining copulation) and not reproductive success (i.e. success in

producing offspring). As these two parameters are not necessarily related (Thompson et al., 2011),

it cannot be excluded that short forceps males have a similar or even higher reproductive output

than their counterparts. This could be because mating is more efficient in short compared to long

forceps males (Brown, 2006), or because short forceps males have alternative reproductive tactics

to long forceps males (see above; Tomkins & Brown, 2004). One study used a genetic approach to

study reproductive success and found no association between male forceps length and the

number of offspring sired (Sandrin et al., 2015). However, this study only focused on length

variation within short forceps males. The second limitation is that all these studies have examined

mating success in groups of a maximum of two males. This number is far less than the hundreds

of individuals that typically constitute earwig aggregates (Meunier, 2024) and thus cannot rule out

the possibility that the apparent advantage of long-forceps males in pairs or trios may disappear

or even become a disadvantage under different demographic conditions (Hunt & Simmons, 2001;

Oliveira et al., 2008; Tomkins & Brown, 2004). Hence, our current understanding of the

relationship between forceps length and male fitness should be treated with caution in the

European earwig. Our results thus call for future studies to determine the function and reliability

of male forceps length under natural conditions, and the evolutionary drivers that explain the

maintenance of its diversity within and between populations.

367 Acknowledgments

368   We thank Lucas Marie-Orléach for his comments on a previous version of this manuscript. We also

369   thank Jean-Christophe Lenoir and Armand Guillermin for their help with field sampling, and the

370   INRAE unité expérimentale Recherche Intégrée Gotheron for giving us access to their orchards for

371   earwig field sampling. Finally, we thank the two anonymous reviewers and Luna Grey for their

372   comments on this manuscript.

374 Funding

375   This project has been funded by the French National Research Agency (ANR-20-CE02-0002 to J.M).

377 Data, script and code availability

378   All data and R script are available online: https://doi.org/10.5281/zenodo.11469628

380 Conflict of interest disclosure

381   The authors declare that they have no financial conflict of interest with the content of this article.

382   J.M is a recommender for PCI Zoology.

References

Arcila, F., & Meunier, J. (2020). Friend or foe? The apparent benefits of gregarine (Apicomplexa:
Sporozoa) infection in the European earwig. *International Journal for Parasitology*, *50*(6–
7), 461–469. https://doi.org/10.1016/j.ijpara.2020.01.007

Berglund, A., Bisazza, A., & Pilastro, A. (1996). Armaments and ornaments: An evolutionary
explanation of traits of dual utility. *Biological Journal of the Linnean Society*, *58*(4), 385–
399. https://doi.org/10.1111/j.1095-8312.1996.tb01442.x

Briceño, R. D., & Eberhard, W. G. (1995). The functional morphology of male cerci and associated
characters in 13 species of tropical earwigs (Dermaptera: Forficulidae, Labiidae,
Carcinophoridae, Pygidicranidae). *Smithsonian Contributions to Zoology*, *555*, 1–63.
https://doi.org/10.5479/si.00810282.555

Brown, G. S. (2006). *Sperm competition and male forceps dimorphism in the European earwig*
Forficula auricularia *(Dermaptera: Forficulina)* (Issue October). University of St. Andrews.

Champely, S., Ekstrom, C., Dalgaard, P., Gill, J., Weibelzahl, S., Anandkumar, A., Ford, C., Volcic, R.,
& Rosario, H. D. (2020). *pwr: Basic Functions for Power Analysis* (1.3-0) [Computer
software]. https://cran.r-project.org/web/packages/pwr/index.html

Cheutin, M.-C., Leclerc, B., & Meunier, J. (2024). With or without you: Gut microbiota does not
predict aggregation behavior in European earwig females. *Behavioral Ecology*, *35*(3),
arae022. https://doi.org/10.1093/beheco/arae022

Coulm, M., & Meunier, J. (2021). Effects of temperature, fungal infection and weight on intermoult
duration and survival of starving larvae in the European earwig. *Journal of Insect*
Physiology, *132*, 104262. https://doi.org/10.1016/j.jinsphys.2021.104262

Crumb, S. E., & Eide, P. M. (1941). The European earwig. *United State Department of Agriculture.*
Technical Bulletin, *766*, 1–73.

Damiens, D., Bressac, C., Brillard, J.-P., & Chevrier, C. (2002). Qualitative aspects of sperm stock in
males and females from Eupelmus orientalis and Dinarmus basalis (Hymenoptera:
Chalcidoidea) as revealed by dual fluorescence. *Physiological Entomology*, *27*(2), 97–102.
https://doi.org/10.1046/j.1365-3032.2002.00274.x

Davies, N. B., Krebs, J. R., & West, S. A. (2012). *An introduction to behavioural ecology*. Wiley-
Blackwell.

Eberhard, W. G., & Gutierrez, E. E. (1991). Male dimorphisms in beetles and earwigs and the
question of developmental constraints. *Evolution*, *45*(1), 18–28.
https://doi.org/10.2307/2409478

Emlen, D. J. (2001). Costs and the diversification of exaggerated animal structures. *Science*,
291(5508), 1534–1536. https://doi.org/10.1126/science.291.5508.1534

Emlen, D. J. (2008). The evolution of animal weapons. *Annual Review of Ecology, Evolution, and*
Systematics, *39*(1), 387–413. https://doi.org/10.1146/annurev.ecolsys.39.110707.173502

Forslund, P. (2000). Male-male competition and large size mating advantage in European earwigs,
Forficula auricularia. *Animal Behaviour*, *59*(4), 753–762.
https://doi.org/10.1006/anbe.1999.1359

Forslund, P. (2003). An experimental investigation into status-dependent male dimorphism in the
European earwig, *Forficula auricularia*. *Animal Behaviour*, *65*(2), 309–316.
https://doi.org/10.1006/anbe.2003.2046

Fox, J., & Weisberg, S. (2019). *An {R} Companion to Applied Regression.* Thousand Oaks CA: Sage.

Fricaux, T., Le Navenant, A., Siegwart, M., Rault, M., Coustau, C., & Le Goff, G. (2023). The
Molecular Resistance Mechanisms of European Earwigs from Apple Orchards Subjected to
Different Management Strategies. *Insects*, *14*(12), 944.
https://doi.org/10.3390/insects14120944

Goldberg, D. L., Landy, J. A., Travis, J., Springer, M. S., & Reznick, D. N. (2019). In love and war: The
morphometric and phylogenetic basis of ornamentation, and the evolution of male display
behavior, in the livebearer genus *Poecilia*. *Evolution*, *73*(2), 360–377.
https://doi.org/10.1111/evo.13671

González-Miguéns, R., Muñoz-Nozal, E., Jiménez-Ruiz, Y., Mas-Peinado, P., Ghanavi, H. R., &
García-París, M. (2020). Speciation patterns in the *Forficula auricularia* species complex:
Cryptic and not so cryptic taxa across the western Palaearctic region. *Zoological Journal of*
the Linnean Society, *190*(3), 788–823. https://doi.org/10.1093/zoolinnean/zlaa070

Grether, G. F. (1997). Survival cost of an intrasexually selected ornament in a damselfly.
Proceedings of the Royal Society of London. Series B: Biological Sciences, *264*(1379), 207–
210. https://doi.org/10.1098/rspb.1997.0029

Günther, K., & Herter, K. (1974). Dermaptera (Ohrwürmer). In J. G. Helmcke, D. Starck, & H.
Wermuth (Eds.), *Handbuch der Zoologie*.

Hartig, F. (2020). *DHARMa: Residual diagnostic for hierarchical (Multi-level / mixed) regression*
models. (p. v 0.3.3.0). http://florianhartig.github.io/DHARMa/

Honorio, R., Moreau, S., Lécureuil, C., & Meunier, J. (2023). Cadmium does not affect post-hatching
maternal care or early offspring development in earwigs. *Behavioral Ecology*, *34*(3), 399–
407. https://doi.org/10.1093/beheco/arad011

Hunt, J., & Simmons, L. W. (2001). Status-dependent selection in the dimorphic beetle
Onthophagus taurus. *Proceedings of the Royal Society of London. Series B: Biological*
Sciences, *268*(1484), 2409–2414. https://doi.org/10.1098/rspb.2001.1758

Jackman, S., Tahk, A., Zeileis, A., Maimone, C., Fearon, J., & Meers, Z. (2024). *pscl: Political Science*
Computational Laboratory (1.5.9) [Computer software]. https://cran.r-
project.org/web/packages/pscl/index.html

Kamimura, Y. (2014). Pre- and postcopulatory sexual selection and the evolution of sexually
dimorphic traits in earwigs (Dermaptera). *Entomological Science*, *17*(2), 139–166.
https://doi.org/10.1111/ens.12058

Koch, L. K., & Meunier, J. (2014). Mother and offspring fitness in an insect with maternal care:
Phenotypic trade-offs between egg number, egg mass and egg care. *BMC Evolutionary*
Biology, *14*(1), 125. https://doi.org/10.1186/1471-2148-14-125

Kohlmeier, P., Holländer, K., & Meunier, J. (2016). Survival after pathogen exposure in group-living
insects: Don't forget the stress of social isolation! *Journal of Evolutionary Biology*, *29*(9),
1867–1872. https://doi.org/10.1111/jeb.12916

Körner, M., Foitzik, S., & Meunier, J. (2018). Extended winters entail long-term costs for insect
offspring reared in an overwinter burrow. *Journal of Thermal Biology*, *74*, 116–122.
https://doi.org/10.1016/j.jtherbio.2018.03.021

Körner, M., Vogelweith, F., Foitzik, S., & Meunier, J. (2017). Condition-dependent trade-off
between weapon size and immunity in males of the European earwig. *Scientific Reports*,
7(1), 7988. https://doi.org/10.1038/s41598-017-08339-6

Kramer, J., Thesing, J., & Meunier, J. (2015). Negative association between parental care and
sibling cooperation in earwigs: A new perspective on the early evolution of family life?
Journal of Evolutionary Biology, *28*(7), 1299–1308. https://doi.org/10.1111/jeb.12655

Le Navenant, A., Siegwart, M., Maugin, S., Capowiez, Y., & Rault, M. (2019). Metabolic mechanisms
and acetylcholinesterase sensitivity involved in tolerance to chlorpyrifos-ethyl in the
earwig *Forficula auricularia*. *Chemosphere*, *227*, 416–424.
https://doi.org/10.1016/j.chemosphere.2019.04.065

Lenth, R. V. (2022). *Emmeans: Estimated Marginal Means, aka Least-Squares Means. R package*
version 1.7.2. Https://CRAN.R-project.org/package=emmeans [Computer software].
https://cran.r-project.org/package=emmeans

Malagnoux, L., Capowiez, Y., & Rault, M. (2015). Impact of insecticide exposure on the predation
activity of the European earwig *Forficula auricularia*. *Environmental Science and Pollution*
Research, *22*(18), 14116–14126. https://doi.org/10.1007/s11356-015-4520-9

Malagnoux, L., Marliac, G., Simon, S., Rault, M., & Capowiez, Y. (2015). Management strategies in
apple orchards influence earwig community. *Chemosphere*, *124*(1), 156–162.
https://doi.org/10.1016/j.chemosphere.2014.12.024

Mauduit, E., Lécureuil, C., & Meunier, J. (2021). Sublethal exposure to deltamethrin stimulates
reproduction and has limited effects on post-hatching maternal care in the European
earwig. *Environmental Science and Pollution Research*, *28*(29), 39501–39512.
https://doi.org/10.1007/s11356-021-13511-7

McCullough, E. L., & Emlen, D. J. (2013). Evaluating the costs of a sexually selected weapon: Big
horns at a small price. *Animal Behaviour*, *86*(5), 977–985.
https://doi.org/10.1016/j.anbehav.2013.08.017

McCullough, E. L., Miller, C. W., & Emlen, D. J. (2016). Why sexually selected weapons are not
ornaments. *Trends in Ecology & Evolution*, *31*(10), 742–751.
https://doi.org/10.1016/j.tree.2016.07.004

McCullough, E. L., & Tobalske, B. W. (2013). Elaborate horns in a giant rhinoceros beetle incur
negligible aerodynamic costs. *Proceedings of the Royal Society B: Biological Sciences*,
280(1758), 20130197. https://doi.org/10.1098/rspb.2013.0197

Merleau, L.-A., Larrigaldie, I., Bousquet, O., Devers, S., Keller, M., Lécureuil, C., & Meunier, J.
(2022). Exposure to pyriproxyfen (juvenile hormone agonist) does not alter maternal care
and reproduction in the European earwig. *Environmental Science and Pollution Research*,
29(48), 72729–72746. https://doi.org/10.1007/s11356-022-20970-z

Meunier, J. (2024). The biology and social life of earwigs (Dermaptera). *Annual Review of
Entomology*, *69*, 259–276. https://doi.org/10.1146/annurev-ento-013023015632

Meunier, J., Dufour, J., Van Meyel, S., Rault, M., & Lécureuil, C. (2020). Sublethal exposure to
deltamethrin impairs maternal egg care in the European earwig *Forficula auricularia*.
Chemosphere, *258*, 127383. https://doi.org/10.1016/j.chemosphere.2020.127383

Mills, M. R., Nemri, R. S., Carlson, E. A., Wilde, W., Gotoh, H., Lavine, L. C., & Swanson, B. O. (2016).
Functional mechanics of beetle mandibles: Honest signaling in a sexually selected system.
Journal of Experimental Zoology Part A: Ecological Genetics and Physiology, *325*(1), 3–12.
https://doi.org/10.1002/jez.1961

Moore, A. J., & Wilson, P. (1993). The evolution of sexually dimorphic earwig forceps: Social
interactions among adults of the toothed earwig, *Vostox apicedentatus*. *Behavioral
Ecology*, *4*(1), 40–48. https://doi.org/10.1093/beheco/4.1.40

Newbolt, C. H., Acker, P. K., Neuman, T. J., Hoffman, S. I., Ditchkoff, S. S., & Steury, T. D. (2017).
Factors influencing reproductive success in male white-tailed deer: Male Breeding Success
in Deer. *The Journal of Wildlife Management*, *81*(2), 206–217.
https://doi.org/10.1002/jwmg.21191

Oliveira, R. F., Taborsky, M., & Brockmann, H. J. (2008). *Alternative reproductive tactics: An
integrative approach*. Cambridge University Press.

Oufiero, C. E., & Garland, T. (2007). Evaluating performance costs of sexually selected traits.
Functional Ecology, *21*(4), 676–689. https://doi.org/10.1111/j.1365-2435.2007.01259.x

Pena, R. C., & Levinton, J. (2021). Two species of fiddler crab show no cost of bearing a sexually
selected giant claw in predator escape behaviors. *Journal of Experimental Marine Biology
and Ecology*, *541*, 151570. https://doi.org/10.1016/j.jembe.2021.151570

Preston, B. T., Jalme, M. S., Hingrat, Y., Lacroix, F., & Sorci, G. (2011). Sexually extravagant males
age more rapidly: Sexually extravagant males age rapidly. *Ecology Letters*, *14*(10), 1017–
1024. https://doi.org/10.1111/j.1461-0248.2011.01668.x

Qiu, W., Chavarro, J., Lazarus, R., Rosner, B., & Ma, J. (2021). *powerSurvEpi: Power and Sample Size*
Calculation for Survival Analysis of Epidemiological Studies (0.1.3) [Computer software].
https://cran.r-project.org/web/packages/powerSurvEpi/index.html

Radesäter, T., & Halldórsdóttir, H. (1993). Two male types of the Common earwig: Male-male
competition and mating success. *Ethology*, *95*(2), 89–96. https://doi.org/10.1111/j.1439-
0310.1993.tb00460.x

Rantala, M. J., Roff, D. a, & Rantala, L. M. (2007). Forceps size and immune function in the earwig
Forficula auricularia L. *Biological Journal of the Linnean Society*, *36*, 331–332.
https://doi.org/10.1111/j.1095-8312.2007.00741.x

Raveh, S., Vogt, D., & Kölliker, M. (2016). Maternal programming of offspring in relation to food
availability in an insect (*Forficula auricularia*). *Proceedings of the Royal Society of London*
B: Biological Sciences, *283*(1828), 20152936. https://doi.org/10.1098/rspb.2015.2936

Sandrin, L., Meunier, J., Raveh, S., Walser, J.-C., & Kölliker, M. (2015). Multiple paternity and mating
group size in the European earwig, *Forficula auricularia*. *Ecological Entomology*, *40*(2),
159–166. https://doi.org/10.1111/een.12171

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image
analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Shuker, D. M., & Simmons, L. W. (2014). *The evolution of insect mating systems*. Oxford University
Press.

Somjee, U., Woods, H. A., Duell, M., & Miller, C. W. (2018). The hidden cost of sexually selected
traits: The metabolic expense of maintaining a sexually selected weapon. *Proceedings of*
the Royal Society B: Biological Sciences, *285*(1891), 20181685.
https://doi.org/10.1098/rspb.2018.1685

Styrsky, J. D., & Rhein, S. V. (1999). Forceps size does not determine fighting success in European
earwigs. *Journal of Insect Behavior*, *12*(4), 475–482.

Swallow, J., & Husak, J. (2011). Compensatory traits and the evolution of male ornaments.
Behaviour, *148*(1), 1–29. https://doi.org/10.1163/000579510X541265

Therneau, T. M. (2020). *A Package for Survival Analysis in R* [Computer software]. https://cran.r-
project.org/package=survival

Thesing, J., Kramer, J., Koch, L. K., & Meunier, J. (2015). Short-term benefits, but transgenerational
costs of maternal loss in an insect with facultative maternal care. *Proceedings of the Royal*
Society of London B: Biological Sciences, *282*(1817), 20151617.
https://doi.org/10.1098/rspb.2015.1617

Thompson, D. J., Hassall, C., Lowe, C. D., & Watts, P. C. (2011). Field estimates of reproductive
success in a model insect: Behavioural surrogates are poor predictors of fitness. *Ecology*
Letters, *14*(9), 905–913. https://doi.org/10.1111/j.1461-0248.2011.01655.x

Tomkins, J. L., & Brown, G. S. (2004). Population density drives the local evolution of a threshold
dimorphism. *Nature*, *431*, 1099–1103. https://doi.org/10.1038/nature02936.1.

Tomkins, J. L., Kotiaho, J. S., & LeBas, N. R. (2005). Matters of scale: Positive allometry and the
evolution of male dimorphisms. *American Naturalist*, *165*(3), 389–402.
https://doi.org/10.1086/427732

Tourneur, J.-C., & Meunier, J. (2020). Variations in seasonal (not mean) temperatures drive rapid
adaptations to novel environments at a continent scale. *Ecology*, *101*(4), e02973.
https://doi.org/10.1002/ecy.2973

Tullis, A., & Straube, C. H. T. (2017). The metabolic cost of carrying a sexually selected trait in the
male fiddler crab *Uca pugilator*. *Journal of Experimental Biology*, *220*, 3641–3648.
https://doi.org/10.1242/jeb.163816

Van Meyel, S., & Meunier, J. (2022). Costs and benefits of isolation from siblings during family life
in adult earwigs. *Animal Behaviour*, *193*, 91–99.
https://doi.org/10.1016/j.anbehav.2022.09.003

van Noordwijk, A. J., & de Jong, G. (1986). Acquisition and allocation of resources: Their influence
on variation in life history tactics. *The American Naturalist*, *128*(1), 137–142.

Vogelweith, F., Foitzik, S., & Meunier, J. (2017). Age, sex, mating status, but not social isolation
interact to shape basal immunity in a group-living insect. *Journal of Insect Physiology*, *103*,
64–70. https://doi.org/10.1016/j.jinsphys.2017.10.007

Walker, K. A., & Fell, R. D. (2001). Courtship roles of male and female European earwigs, *Forficula*
auricularia L. (Dermaptera: Forficulidae), and sexual use of forceps. *Journal of Insect*
Behavior, *14*(1), 1–17.

Wilkinson, G. S., Kahler, H., & Baker, R. H. (1998). Evolution of female mating preferences in stalk-
eyed flies. *Behavioral Ecology*, *9*(5), 525–533. https://doi.org/10.1093/beheco/9.5.525

Wong, J. W. Y., & Kölliker, M. (2014). Effects of food restriction across stages of juvenile and early
adult development on body weight, survival and adult life history. *Journal of Evolutionary*
Biology, *27*(11), 2420–2430. https://doi.org/10.1111/jeb.12484

Table 1 – Effects of male forceps length and population of origin of the three physiological and

three behaviours measured in this study. Significant p-values are in bold.

	Population of origin	Forceps length	Interaction	Statistical power
Locomotor performance	$F_{1,115} = 0.21$ P = 0.648	$F_{1,115} = 1.14$ P = 0.287	$F_{1,115} = 1.25$ P = 0.265	0.241
Likelihood of fleeing away after a physical disturbance	LR $\chi^2_1 = 8.74$ P = **0.003**	LR $\chi^2_1 = 1.06$ P = 0.303	LR $\chi^2_1 = 0.57$ P = 0.450	0.239
Aggregation score	$F_{1,88} = 3.83$ P = **0.054**	$F_{1,88} = 0.004$ P = 0.948	$F_{1,88} = 0.02$ P = 0.878	0.347
Survival rate under harsh environmental conditions	LR $\chi^2_1 = 7.88$ P = **0.005**	LR $\chi^2_1 = 0.02$ P = 0.885	LR $\chi^2_1 = 4.22$ P = **0.039**	0.447
Sperm count	$F_{1,35} = 0.42$ P = 0.522	$F_{1,35} = 0.19$ P = 0.667	$F_{1,35} = 0.75$ P = 0.391	0.155
Survival rate after infection	LR $\chi^2_1 = 0.02$ P = 0.891	LR $\chi^2_1 = 0.15$ P = 0.695	LR $\chi^2_1 = 0.46$ P = 0.496	0.432

597 **Figure 1** – Timeline of the experimental design and evolution of sample size over the course of the

598 experiment. The observed decrease in sample size over time reflects their mortality during

599 isolation and in the absence of any food source. Males are distributed according to their forceps

600 length category and population of origin.

			Survival to isolation & starvation				
		D0 Isolation	D5 Locomotor performance	D10 Response to disturbance	D24 Aggregation	D31 Sperm count	D31 Survival to infection
Long forceps	Cinais	N=30	30	30	18	9	8
	Valence	N=30	29	29	28	10	18
Short forceps	Cinais	N=30	30	30	22	10	12
	Valence	N=30	30	30	24	10	14

Figure 2 – Forceps length distribution of the 120 selected males with (blue) short and (orange)

long forceps in (A) Cinais and (B) Valence populations. Dashed lines show the mean values per

forceps category for each population.

Figure 3 – Association between forceps length and male (A) locomotor performance, (B) likelihood

of fleeing after a physical disturbance, (C) propensity to aggregate with conspecifics, (D) likelihood

of being alive after 31 days in social isolation and without food access, (E) sperm storage in the

vesicle and (F) survival rate after pathogen infection. Males came from Cinais (square) or Valence

(triangle). Different letters *P* <0.05.