Ether lipids influence cancer cell fate by modulating iron uptake

Whitney S. Henry[1], Sebastian Müller[2], Jia-Shu Yang[3], Sarah Innes-Gold[4], Sunny Das[1], Ferenc Reinhardt[1], Kim Sigmund[1], Vaishnavi V. Phadnis[1,5], Zhengpeng Wan[6], Elinor Eaton[1], Julio L. Sampaio[7], George W. Bell[1], Amartya Viravalli[8], Paula T. Hammond[9,10,15], Roger D. Kamm[6,12,15], Adam E. Cohen[4,12,15], Natalie Boehnke[8,9,15], Victor W. Hsu[3,15], Kandice R. Leventhal[13,15], Raphaël Rodriguez[2,15,*] and Robert A. Weinberg[1,5,14,15,*,#]

1. Whitehead Institute for Biomedical Research, Cambridge, MA 02142, USA
2. Institut Curie, CNRS, INSERM, PSL Research University, Equipe Labellisée Ligue Contre le Cancer, Paris 75005, France
3. Division of Rheumatology, Inflammation and Immunity, Brigham and Women's Hospital, and Dept. of Medicine, Harvard Medical School, Boston, MA 02115, USA
4. Dept. of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA
5. Dept. of Biology, MIT, Cambridge, MA 02139, USA
6. Dept. of Biological Engineering, MIT, Cambridge, MA 02139, USA
7. Institut Curie, INSERM, Mines ParisTech, Paris 75005, France
8. Dept. of Chemical Engineering and Materials Science, University of Minnesota Minneapolis, MN 55455, USA
9. Koch Institute for Integrative Cancer Research, MIT, Cambridge, MA 02139, USA
10. Dept. of Chemical Engineering, MIT, Cambridge, MA 02139, USA
11. Dept. of Mechanical Engineering, MIT, Cambridge, MA 02139, USA
12. Dept. of Physics, Harvard University, Cambridge, MA 02138, USA
13. Dept. of Molecular Physiology and Biological Physics, Center for Membrane and Cell Physiology, University of Virginia, Charlottesville, VA 22903, USA
14. Ludwig Center for Molecular Oncology, Cambridge, MA 02139, USA
15. Senior author

#Lead contact

*Correspondence: Raphaël Rodriguez, PhD (raphael.rodriguez@curie.fr) and Robert A. Weinberg, PhD (weinberg@wi.mit.edu)

ABSTRACT

Cancer cell fate has been widely ascribed to mutational changes within protein-coding genes associated with tumor suppressors and oncogenes. In contrast, the mechanisms through which the biophysical properties of membrane lipids influence cancer cell survival, dedifferentiation and metastasis have received little scrutiny. Here, we report that cancer cells endowed with a high metastatic ability and cancer stem cell-like traits employ ether lipids to maintain low membrane tension and high membrane fluidity. Using genetic approaches and lipid reconstitution assays, we show that these ether lipid-regulated biophysical properties permit non-clathrin-mediated iron endocytosis via CD44, leading directly to significant increases in intracellular redox-active iron and enhanced ferroptosis susceptibility. Using a combination of in vitro three-dimensional microvascular network systems and in vivo animal models, we show that loss of ether lipids also strongly attenuates extravasation, metastatic burden and cancer stemness. These findings illuminate a mechanism whereby ether lipids in carcinoma cells serve as key regulators of malignant progression while conferring a unique vulnerability that can be exploited for therapeutic intervention.

KEYWORDS

Ether lipids, membrane tension, endocytosis, CD44, iron, metastasis, ferroptosis.

INTRODUCTION

Cancer cells have the capacity to undergo dynamic changes in identity, structure, and function, making them remarkably versatile and adaptable. Alterations in the lipid composition of cell membranes is one key element contributing to the phenotypic plasticity of cells. The distinctive physicochemical properties and subcellular localization of various lipids within cell membranes influence a range of biological processes, including cellular trafficking, signaling and metabolism[1]. Despite our growing knowledge of lipid biology, our understanding of how specific lipid subtypes impact cancer cell fate remains limited.

Emerging studies have demonstrated that therapy-resistant mesenchymal-like carcinoma cells exhibit an elevated vulnerability to ferroptosis[2-4], an iron-dependent form of cell death characterized by the unrestricted accumulation of oxidized membrane phospholipids[5-7]. Indeed, in previous work we showed that the natural product salinomycin can selectively eliminate otherwise therapy resistant, mesenchymal-enriched cancer stem cells (CSC), doing so by targeting lysosomal iron to promote an iron-dependent cell death[4, 8, 9]. In this context, we found that such CSC-enriched cells exhibit a high intracellular iron load compared to their non-CSC-like counterparts, rendering them especially vulnerable to elimination by induced ferroptosis[13,14].

Ferroptosis can also be instigated by pharmacologic inhibition of ferroptosis suppressors, such as glutathione peroxidase 4 (GPX4)[5, 10], ferroptosis-suppressor protein 1 (FSP1, previously known as AIFM2)[11, 12], as well as through downregulation of reduced glutathione (GSH)[13, 14]. Activated CD8$^+$ T cells may also induce ferroptosis in cancer cells[9,10]. Beyond cancer, ferroptosis has been implicated in the pathogenesis of several neurodegenerative diseases and acute injury of the kidney, liver and heart[15-18].

In previous work, we undertook an unbiased, genome-wide CRISPR/Cas9 screen with the goal of identifying genes that govern ferroptosis susceptibility in high-grade human serous ovarian cancer cells[19]. This screen revealed a previously unrecognized role for ether lipid-synthesizing enzymes, such as alkylglycerone phosphate synthase (AGPS), in modulating ferroptosis susceptibility. The ether phospholipids generated by these enzymes represent a unique subclass of glycerophospholipids characterized by an ether-linked hydrocarbon group formed at the *sn*-1 position of the glycerol backbone[20]. This phospholipid subtype constitutes ~20% of the total phospholipid pool in many types of mammalian cells.

The significance of ether lipid species in human health is underscored by the severe inherited peroxisomal disorders caused by their deficiency[20]. This often manifests as profound developmental abnormalities, such as neurological defects, visual and hearing loss, and reduced lifespan. In the context of cancer, elevated ether lipid levels have been correlated with increased metastatic potential of carcinoma cells[21-24]. Despite these pathological associations, the mechanism(s) by which ether lipids affect cancer progression remain(s) elusive. Furthermore, exactly why loss of ether lipids results in decreased ferroptosis susceptibility required further investigation.

Our previous work and that of others ascribed a role to polyunsaturated ether phospholipids as chemical substrates prone to the iron-mediated oxidation that triggers ferroptotic cell death[19, 25]. Here, we demonstrate that ether lipids also play an unrelated biophysical role, doing so by facilitating iron endocytosis in carcinoma cells. This represents an unappreciated mechanism of intracellular signaling in which a lipid

139   contributes to the intracellular level of a critical metal-based signaling species - iron. In
140   addition, our findings highlight the functional importance of this poorly studied lipid
141   subtype in enabling a variety of malignancy-associated cell phenotypes including
142   metastasis and tumor-initiating abilities. Together, these results establish a role for ether
143   lipids as critical effectors of cancer cell fate.
144
145   **RESULTS**
146   **Ether lipids play a key role in maintaining a ferroptosis susceptible cell state**
147   In order to investigate the mechanism(s) by which ether lipid deficiency reduces
ferroptosis susceptibility, we employed CRISPR/Cas9 to knockout (KO) the *AGPS* gene,
in ferroptosis-sensitive TGF-β-treated PyMT-1099 murine breast cancer cells[26] **(Fig. 1a,**
1b, Extended Data Fig. 1a). The *AGPS* gene encodes a rate-limiting enzyme critical for
ether lipid biosynthesis[20]. Consistent with our prior studies[19], loss of ether lipids via
AGPS KO significantly decreased the susceptibility of these cancer cells to ferroptosis
induced by treatment with the GPX4 inhibitors ML210 or RSL3 **(Fig. 1c, Extended Data**
Fig. 1b).
155   By performing lipidomic analysis, we validated that knockout of *AGPS* resulted in
a significant reduction in total ether abundance in these cells **(Fig. 1d)**. More than half of
the identified ether lipids contained polyunsaturated fatty acyl groups which are highly
prone to free radical attack **(Fig. 1e)**. Based on this observation, we speculated that loss
of ether lipids could attenuate ferroptosis susceptibility by depleting the pool of available
ether lipid substrates for lipid peroxidation. Therefore, we performed oxidized lipidomic
analysis on two ferroptosis-sensitive breast cancer cell lines that were treated with a
GPX4 inhibitor. These experiments indicated that ether lipids could indeed be oxidized
following ferroptosis induction **(Fig. 1f, Extended Data Fig. 1c)**.
164   Given that ether lipids only constitute about ~20% of total lipids, we also
investigated whether the relative abundance of non-ether-linked polyunsaturated
phospholipids, were impacted by loss of ether lipids. Surprisingly, our analyses revealed
that ether lipid deficiency actually increased the relative abundance of several
polyunsaturated diacyl phospholipids with putative pro-ferroptosis function[27, 28] **(Fig. 1g)**.
To ensure that these findings were not an idiosyncrasy of our TGF-β-treated PyMT-1099
AGPS KO cells, we confirmed this observation in PyMT-MMTV-derived pB3 murine
AGPS KO breast cancer cells[29] **(Fig. 1h, 1i, Extended Data Fig. 1d)**. Importantly, re-
expression of AGPS (i.e., "addback") could restore the relative levels of these non-
ether-linked polyunsaturated diacyl phospholipids to levels comparable to pB3 WT cells
(Fig. 1j). These findings strongly argue against the notion that ether deficiency
attenuates ferroptosis susceptibility simply by decreasing the global level of
polyunsaturated phospholipids and further underscores the importance of
polyunsaturated ether phospholipids in maintaining a ferroptosis susceptible cell state.
178
179   **Ether lipids regulate cellular redox-active iron levels in cancer cells**
180   The above observations together with our oxidized ether phospholipidomic
analysis supported the notion that ether lipids could modulate ferroptosis susceptibility,
at least in part, by serving as substrates for lipid peroxidation. These observations,
however, failed to address the formal possibility that alterations of ether lipid
composition could also affect intracellular levels of redox-active iron, the central

mediator of the lipid peroxidation that drives ferroptosis. For this reason, we investigated
whether alterations in ether lipid composition actually affected the intracellular levels of
redox-active iron. To address this possibility, we used two orthogonal analyses to
assess intracellular iron levels. Since the endolysosomal compartment is a key reservoir
of iron within cells[30-32], we used a lysosomal iron (II)-specific fluorescent probe,
RhoNox-M[33], to gauge the levels of iron within these cells. In addition, we used
inductively coupled plasma mass spectrometry (ICP-MS) to quantify total intracellular
iron levels[32, 34, 35].
Remarkably, loss of AGPS reduced intracellular iron levels in all murine cancer
cell lines tested, whereas re-expression of AGPS (i.e., "addback") restored intracellular
iron to levels comparable to those seen in parental ferroptosis-sensitive cancer cells
(Fig. 2a-2d, Extended Data Fig. 2a). This provided the first indication that changes in
ether phospholipids directly affect the levels of intracellular iron. Indeed, there was no
direct precedent for the ability of a membrane-associated phospholipid to serve as an
enhancer of the levels of an intracellular metal ion.
Supporting this conclusion, we found that a reduction in the levels of ether lipids,
achieved via knockout of the genes encoding either the AGPS or the fatty acid
reductase 1 (FAR1) enzymes[20] **(Extended Data Fig. 2b)**, also resulted in a significant
decrease in intracellular iron levels, in this case in the mesenchymal-enriched OVCAR8
human high-grade serous ovarian cancer cell line **(Fig. 2e, 2f).** Furthermore, we noted
that treatment of AGPS KO cells with ferric ammonium citrate (FAC)[36, 37], which provides
an exogenous source of ferric ions re-sensitized cultured AGPS KO mesenchymal
breast and ovarian carcinoma cells to ferroptosis induction, doing so even in the
absence of elevated ether phospholipids **(Fig. 2g, 2h)**.
We further extended this analysis by studying the behavior of mammary
carcinoma cells forming tumors in vivo. Consistent with our in vitro data, inductively
coupled plasma mass spectrometry indicated that total iron levels are reduced in breast
tumors derived from implanted pB3 AGPS KO carcinoma cells relative to those arising
from pB3 wildtype (WT) or control pB3 AGPS-addback cells **(Fig. 2i)**. Such observations
reinforced the notion that ether lipids are critical regulators of intracellular iron levels, a
rate-limiting component of ferroptosis.
216
Ether lipids facilitate CD44-mediated iron endocytosis
We proceeded to investigate the mechanism(s) by which membrane-associated
ether lipids regulate intracellular iron content. This led us to examine the behavior of two
proteins that act as major mediators of cellular iron import - transferrin receptor 1
(TfR1)[38] and CD44[32] – and whether their functioning was altered in response to loss of
ether phospholipids. While CD44 is best known as a cell-surface cancer stem-cell
marker[39, 40], recent research revealed its critical role in mediating endocytosis of iron-
bound hyaluronates in CSC-enriched cancer cells and in activated immune cells[32, 41]. To
monitor these two alternative iron import mechanisms, we performed endocytosis
kinetics experiments using fluorescently labeled transferrin as a proxy for TfR1
internalization and fluorescently labeled hyaluronic acid (HA), whose main plasma
membrane receptor is CD44, as a marker for CD44 internalization[42, 43].
Here we observed that the rate of endocytosis of TfR1 was marginally affected by
a deficiency of ether lipids in pB3 breast cancer cells **(Fig. 3a)**. In stark contrast, CD44-

mediated endocytosis was significantly impaired when the *AGPS* gene was knocked out
in this ether lipid-deficient, cancer cells **(Fig. 3b)**. Conversely, CD44-dependent iron
import could be restored to normal levels by introduction of a functional *AGPS* gene into
these *AGPS* KO cells **(Fig. 3b)**. Reduction in the rate of internalization of CD44 but not
TfR1 could also be observed in ether lipid-deficient PyMT-1099 TGF-β-treated breast
cancer cells **(Fig. 3c, 3d)**. Hence, in these cells, ether lipids played a critical role in
modulating intracellular iron concentration by regulating endocytosis of CD44 but not
transferrin receptor. These findings were consistent with our previous observations that
TfR1 and CD44 localize to distinct endocytic vesicles in CSC-enriched cancer cells[32, 44]
making plausible that their internalization was governed by independent endocytic
mechanisms.
To further support the role of CD44 in promoting iron uptake – acting via its
endocytosis of HA – we demonstrate that knocking out the gene encoding CD44 or,
alternatively, treating cancer cells with hyaluronidase, led to a significant reduction in
intracellular iron levels **(Fig. 3e)**. Conversely, supplementing these cells with hyaluronic
acid increased intracellular iron levels **(Fig. 3e)**. Similar observations were seen in
human OVCAR8 cells **(Extended Data Fig. 3a)**. Taken together, these observations
further supported the influential role of CD44 in mediating iron uptake in these cancer
cells[32].
We then studied whether the observed defect in CD44 endocytosis observed in
ether lipid deficient cells was limited to CD44 or, instead, reflected a general impairment
in the endocytosis of a variety of plasma membrane-associated glycoproteins. In fact,
CD44 is known to undergo a type of clathrin- and dynamin-independent form of
endocytosis[44-46]. This alternative mechanism of endocytosis differs from the one
regulating TfR1 recycling, which undergoes clathrin-mediated endocytosis, in which
small invaginations of clathrin-coated pits undergo scission facilitated by the GTPase
dynamin[45, 47].
To test whether loss of ether phospholipids had a wider effect on the clathrin- and
dynamin-independent mode of endocytosis, we examined the rate of uptake of dextran
(70 kDa), a branched polysaccharide known to undergo endocytosis by a clathrin-
independent mechanism[45]. Similar to CD44, we observed that loss of AGPS also
exhibited a significant reduction in the rate of dextran endocytosis; this behavior could
be reversed by restoration of ether phospholipids levels achieved by AGPS
complementation **(Fig. 3f, 3g)**. Moreover, the loss of ether lipids had a negligible effect
on the rate of EGFR endocytosis which, like TfR1, is known to rely on clathrin-mediated
endocytosis **(Fig. 3h, Extended Data Fig. 3b)**.
Hence, these observations reinforced the notion that the internalization of
extracellular and cell-surface molecules is mediated by at least two distinct mechanisms
that differ in their dependence on ether phospholipids. More specifically, these findings
provided strong support for the involvement of a widely acting clathrin- and dynamin-
independent form of endocytosis, on which CD44 internalization depended[32, 48-50] and is
significantly compromised in ether lipid-deficient cells.
273
Ether lipid deficiency impairs membrane biophysical properties
The above observations did not provide mechanistic insights into how changes in
the composition of membrane ether lipids could exert an effect on CD44 internalization.

As observed by others, non-clathrin-mediated endocytosis, which is employed by CD44,
is particularly sensitive to changes in the physicochemical properties of the lipid bilayer
forming plasma membranes[46, 51-56]. Such changes can influence membrane tension,
membrane fluidity and stability, and formation of lipid rafts, all of which, in turn, impact
the assembly and dynamics of clathrin-independent, cell-surface endocytic structures.
Hence, we hypothesized that ether lipids alter the biophysical properties of the lipid
bilayer of the plasma membrane to facilitate elevated iron endocytosis via CD44.
Membrane deformability can be gauged by the parameter of membrane tension,
which measures the forces exerted on a defined cross-section of the plasma
membrane. It is influenced by both the in-plane tension of the lipid bilayer and the
attachment of the plasma membrane to the underlying cell cortex[57, 58]. Indeed,
alterations in membrane tension have long been demonstrated to affect endocytosis[59-
64], prompting us to assess the effects of loss of ether lipids on plasma membrane
tension. To quantify membrane tension directly, we generated a membrane tether using
an optically trapped bead and measured the pulling force (f) and the tube radius (R) to
calculate membrane tension (σ) of living cells[48, 65] **(Fig. 4a, 4b).** We found that depletion
of ether phospholipids led to a significant increase in membrane tension in pB3 AGPS
KO cells relative to the corresponding pB3 WT cancer cells **(Fig. 4c)**. This shift was
largely attenuated upon restoration of AGPS expression in pB3 AGPS KO cells or upon
exposure of cultured cells to liposomes composed of polyunsaturated ether
phospholipids **(Fig. 4c)**. Treatment of pB3 AGPS KO cells with these ether lipid-
containing liposomes also increased the rate of CD44 endocytosis to levels comparable
to those of pB3 WT cells **(Fig. 4d)**. No changes were observed in the rate of clathrin-
dependent TfR1 endocytosis under these conditions **(Fig. 4e)**. Taken together, these
results provided the first indication that ether lipids facilitate CD44-mediated iron
endocytosis in cancer cells, in part, by decreasing membrane tension.
Membrane lipid packing can also impact endocytosis[66-68]. It is related to the
fluidity or viscosity of the lipid bilayer, with higher lipid packing correlating with higher
viscosity. This, in turn, affects the ease with which proteins and lipids undergo lateral
diffusion and conformational changes within a lipid bilayer, thereby affecting
endocytosis-related signaling[48]. This motivated us to investigate the contribution of ether
lipids to membrane lipid packing. To do so, we used the C-laurdan lipid-based, polarity-
sensitive dye, which yields a spectral emission shift dependent on the degree of lipid
packing[69]. These measurements are used to calculate a unitless index, termed
generalized polarization (GP), where a higher GP indicates increased lipid packing[69].
Our measurements using the C-laurdan dye indicated that a reduction in ether lipid
levels resulted in a measurable, significant increase in membrane packing **(Fig. 4f-4h)**,
which, like increase increases in membrane tension, negatively affects membrane
plasticity[66].
A third parameter governing the biophysical properties of lipid bilayers involves
the stability (size and lifetime) of lipid rafts. This parameter can be gauged by monitoring
the miscibility transition temperature (T_{misc}) of these membranes[70]. The association of
CD44 with lipid rafts, which are dynamically formed plasma membrane microdomains, is
known to be critical for CD44-mediated HA endocytosis[71]. We reasoned that a decrease
in the stability of lipid rafts, and thus a decrease in T_{misc}, would result in impairment of
CD44 endocytosis[70]. Thus, we measured the effect of loss of ether lipids on the

miscibility transition temperature. In fact, we observed a decrease in T_{misc} upon loss of
AGPS in pB3 cancer cells, which indicated a decrease in lipid raft stability. This finding
supports a role for ether lipids in maintaining the plasma membrane organization
through lipid raft microdomains **(Fig. 4i)**, revealing yet another biophysical property of
lipid bilayers that can influence CD44 endocytosis.
It is noteworthy that clathrin-independent endocytosis exhibits a greater
dependency on the membrane biophysical properties assessed above[46, 51-56]. This may
explain why loss of ether lipids can exert a significant effect on the rate of CD44-
mediated iron endocytosis but negligible effects on the clathrin-mediated TfR1
endocytosis. Furthermore, these findings illuminated a mechanism by which membrane-
associated ether lipids could govern a major mechanism of iron internalization, which in
turn could impact the vulnerability of cancer cells to ferroptosis inducers.
335
Loss of ether lipids decreases metastasis and cancer cell stemness
Prior studies have demonstrated that reduced membrane tension and elevated
intracellular iron can promote cancer metastasis[72-76]. These findings of others caused us
to investigate whether changes in the ether lipid composition of cancer cells impacted
key steps of the multi-step invasion-metastasis cascade, notably extravasation
efficiency, post-extravasation proliferation[77], as well as the functionally critical trait of
cancer cell stemness, i.e., tumor-initiating ability.
We measured extravasation efficiency by employing an in vitro three-dimensional
microvascular network system composed of human umbilical vein endothelial cells
(HUVECs) and normal human lung fibroblasts. This system has been shown to
accurately model some of the complex biological processes associated with cancer cell
extravasation[78-82]. Using this defined experimental system, we found that loss of ether
lipids significantly decreased extravasation efficiency **(Fig. 5a-5c)**. Furthermore, we
observed a strong reduction in overall metastatic burden following intracardiac injection
in syngeneic hosts of the pB3 AGPS KO breast cancer cells relative to corresponding
wildtype cells **(Fig. 5d-5f)**. As an important control in these experiments, we determined
that ether lipid deficiency in these cells had a modest effect on primary tumor growth
kinetics, making it unlikely that the loss of ether phospholipids had a significant effect on
post-extravasation proliferation of disseminated tumor cells **(Fig. 5g-5i)**. A decrease in
metastatic burden was also observed upon knockout of *AGPS* and *FAR1* in OVCAR8
cancer cells, and upon loss of CD44 in pB3 cancer cells **(Extended Data Fig. 4a-4c)**.
Given that high CD44 expression and elevated intracellular iron levels are
positively correlated with cancer cell stemness[32, 39, 83], we investigated whether ether
lipid deficiency also affects the tumor-initiating capacity of cancer cells as gauged by an
experimental limiting dilution tumor-implantation assay. These experiments indicated
that loss of ether lipids in pB3 breast cancer cells decreases cancer cell stemness **(Fig.
5j, Extended Data Fig. 4d)**, which, as we have found in other investigations, serve as a
reliable marker of metastasis-initiating capacity[84]. In addition, we found that loss of ether
lipids significantly attenuates the tumor-initiating potential and metastatic capacity of
PyMT-1099 AGPS KO TGF-β-treated (ether lipid deficient) cancer cells following
implantation into the orthotopic site — the mammary stromal fat pad **(Fig. 5k-5m)**.
Hence, ether lipids play critical roles in promoting cancer cell stemness and resulting
post-extravasation colonization.

369
DISCUSSION

The present findings indicate the need to consider the complex interplay between
genetics and the biophysical properties of cell membranes as determinants of cancer
cell fate and emphasize the potential role of lipids and metals in this process.
Membrane-associated phospholipids have previously been implicated as important
mediators of cell transformation, in large part through the actions of inositol
phospholipids and their derivatives[85]. In the present study, we shed light on an entirely
different and poorly studied role of lipids in influencing cell fate through their effects on
membrane biophysical properties and their impact on iron homeostasis. Specifically, we
uncover a mechanism whereby alterations in ether lipids affect the biophysical
properties of the plasma membrane to impact distinct cell-biological processes — iron
uptake and neoplasia-related phenotypes, notably metastasis and cancer cell
stemness/tumor-initiating ability. Importantly, this biochemical configuration creates a
unique vulnerability of cancer cells to ferroptosis, and suggests that targeting lipid
metabolism and iron homeostasis could be exploited to suppress subpopulations of
highly metastatic and drug-tolerant carcinoma cells[86].
Ether phospholipids have been widely portrayed as participants in ferroptosis
through their role as substrates prone to iron-catalyzed peroxidation. However, our
findings indicate an entirely different mechanism whereby ether lipids directly modulate
the levels of intracellular iron, a rate-limiting component governing ferroptosis
susceptibility[4, 32, 73]. By emphasizing the role of membrane biophysical properties in
governing iron uptake, we depart from the conventional focus limited to portraying
phospholipids as substrates for peroxidation. This shift in perspective has the potential
to open new avenues for research, as it challenges researchers to explore the
biophysical aspects of membranes as a new dimension in the regulation of this cell
death program.
Alterations in intracellular iron level can impact gene expression via various
mechanisms including modulation of chromatin-modifying enzyme activity[32, 87, 88]. For
example, increase of intracellular iron levels has been shown to promote the activity of
iron-dependent demethylases[32, 87, 88], impacting gene expression profiles underlying cell
plasticity[32] and immune cell activation[41]. Our finding that ether lipid deficiency reduces
intracellular iron levels explains, at least in part, how loss of ether lipids may impact
cancer-associated transcriptional programs, acting at the epigenetic level and enabling
a variety of malignancy-associated cell phenotypes including metastasis and cancer
stemness. Such mechanisms may act in concert with non-iron-dependent processes,
which are also regulated by ether lipids to affect cancer malignancy traits.
The implications of our research findings may extend far beyond the realm of
cancer pathogenesis. We speculate that the biophysical modulation of membranes and
its intersection with iron biology could be a widely-acting determinant of cell fate,
impacting processes such as differentiation, immune activation, wound healing, and
embryonic development.

ACKNOWLEDGMENTS

413   We thank all members of the Weinberg, Farese and Walther labs for insightful
414   discussions and reagents. We are grateful to Brent Stockwell, Laurie Boyer and Fabien

Sindikubwabo for helpful discussions. We acknowledge technical support from Caroline A. Lewis, Zon W. Lai and Marina Plays, the CurieCoreTech Metabolomics and Lipidomics Technology Platform at the Institut Curie, the following facilities at the Whitehead Institute: Metabolite Profiling Core, W.M. Keck Microscopy Core, Flow Cytometry Core, as well as the Koch Institute's Robert A. Swanson (1969) Biotechnology Center, specifically the MIT Koch Institute small animal imaging core and MIT Tang Histology facility.

This work was funded in part by the National Institutes of Health: R37GM058615 (VWH), the National Cancer Institute K99CA255844/R00CA255844 (NB), and the National Cancer Institute (RDK). Additional funding support was received from MIT Stem Cell (RAW), Brendan Bradley Gift (RAW), Nile Albright Research Foundation (RAW), Samuel Waxman Cancer Research Foundation (RAW), Virginia, D.K. Ludwig Fund for Cancer Research Center (RAW), European Research Council under the European Union's Horizon 2020 research and innovation programme grant agreement No 647973 (RR), Foundation Charles Defforey-Institut de France (RR), Ligue Contre le Cancer Equipe Labellisée (RR), Fondation Bettencourt Schueller (RR), Marble Center for Cancer Nanomedicine (PTH), Jane Coffin Childs Memorial Fund (WSH), Ludwig Center at MIT's Koch Institute for Integrative Cancer Research (WSH) and the New Horizon UROP Fund/MIT (VVP).

AUTHOR CONTRIBUTIONS

COMPETING INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Fig. 1. Ether lipids play a key role in maintaining a ferroptosis susceptible cell state. See also Extended Data Fig.1.

a. Schematic of experimental model for lipidomic analysis.
b. Immunoblot analysis for AGPS expression in PyMT-1099 WT or AGPS KO cells. Cells were treated with TGF-β (2 ng/ml) for 10 d where indicated.
c. Cell viability following treatment with the GPX4 inhibitor ML210 for 72 h. 1099 WT or AGPS KO cells were pretreated with TGF-β (2 ng/ml) for 10 d prior to assay. Graph is representative of two independent biological replicates.
d. Bar graph showing percent of total lipids constituted by ether lipids following AGPS KO in untreated wildtype (WT) or TGF-β-treated (2 ng/ml;10 d) PyMT-1099 cells.

e. Pie chart showing the relative proportion of ether lipids with various total numbers of
double bonds.
f. Amount in pmol of oxidized phosphatidylethanolamine (Oxi. PE) ether and ester
phospholipids in PyMT-1099 TGF-β cells treated with ML210 for 24 h. Five biological
replicates per condition.
g. Volcano plot showing the $\log_2$ fold change in the relative abundance of various lipid
species upon knockout of AGPS in PyMT-1099 TGF-β-treated cells. Blue indicates
non-ether linked polyunsaturated phospholipids with a total of at least 3 double
bonds; orange indicates all ether lipids identified in lipidomic analysis and black
denotes all other lipids identified.
h. Volcano plot showing the $\log_2$ fold change in the relative abundance of various lipid
species upon knockout of AGPS in pB3 cells. Blue indicates non-ether linked
polyunsaturated phospholipids with a total of at least 3 double bonds; orange
indicates all ether lipids identified in lipidomic analysis and black denotes all other
lipids identified.
i. Bar graph showing the percent of total lipids constituted by ether lipids in pB3 WT,
pB3 AGPS KO and pB3 AGPS addback cells.
j. Bar graph showing the effects of ether lipids on the relative abundance of selected
polyunsaturated diacyl phospholipids in pB3 cells.
Unless stated otherwise, all samples were analyzed in technical triplicates and shown
as the mean +/− SEM. Statistical significance was calculated using unpaired, two-tailed
t-test. For figures 1h-1j: pB3 WT and AGPS KO cells were transduced with the
respective vector control plasmids. pB3 AGPS addback cells are derivatives of AGPS
KO cells transduced with a murine AGPS expression vector.
484
Fig. 2. Ether lipids regulate cellular redox-active iron levels in cancer cells. See
also Extended Data Fig. 2.
a. Relative lysosomal iron levels based on Rhodox-M fluorescence intensity normalized
to the fluorescence intensity of lysotracker. Fold change is calculated relative to
untreated PyMT-1099 wild-type (WT) cells.
b. Relative lysosomal iron levels based on Rhodox-M fluorescence intensity normalized
to the fluorescence intensity of lysotracker. Fold change is calculated relative to pB3
WT cells.
c. Inductively coupled plasma-mass spectrometry (ICP-MS) of cellular iron in PyMT-
1099 WT or AGPS KO cells pretreated with 2 ng/ml TGF-β for 10 d.
d. Inductively coupled plasma-mass spectrometry (ICP-MS) of cellular iron in pB3 cell
line derivatives.
e. Relative lysosomal iron levels in OVCAR8 NT sg, FAR1 KO or AGPS KO cells
pretreated with FAC (50 µg/ml) for 24 h. Data shown are based on Rhodox-M
fluorescence intensity normalized to lysotracker fluorescence intensity. Fold change
is calculated relative to NT sg.
f. Inductively coupled plasma-mass spectrometry (ICP-MS) of cellular iron in OVCAR8
NT sg, FAR1 KO or AGPS KO cells pretreated with FAC (50 µg/ml) for 24 h.
g. Cell viability of OVCAR8 NT sg, FAR1 KO or AGPS KO cells pretreated with FAC (50
µg/ml) for 24 h followed by ML210 treatment for 72 h.

h. Cell viability in response to ML210 treatment. PyMT-1099 WT or AGPS KO cells
were pretreated with TGF-β (2 ng/ml) for 10 days followed by FAC treatment (100
µg/ml) for an additional 24 h. Cells were then treated with ML210 in the presence or
absence of liproxstatin-1 (0.2µM) and cell viability was assessed after 72 h.
i. ICP-MS of cellular iron from primary tumors derived from pB3 WT, pB3 AGPS KO,
and pB3 AGPS addback cells. Mean +/- SEM from 3 independent tumor samples per
condition. Each datapoint represents the average iron measurement from 5 technical
replicates per tumor sample.
All samples were analyzed with 3-6 technical replicates and shown as the mean +/−
SEM unless stated otherwise. Statistical significance was calculated using unpaired,
two-tailed t-test. Abbreviation: NT – nontargeting.
516
Fig. 3. Ether lipids facilitate CD44-mediated iron endocytosis. See also Extended
Data Fig. 3.
a. Endocytic transport of fluorescently labeled transferrin as assessed by quantitative
colocalization with an early endosomal marker (EEA1) in pB3 cells.
b. Endocytic transport of fluorescently labeled hyaluronate probe as assessed by
quantitative colocalization with an early endosomal marker (EEA1) in pB3 cells.
c. Endocytic transport of fluorescently labeled transferrin as assessed by quantitative
colocalization with an early endosomal marker (EEA1) in PyMT-1099 WT or AGPS
KO cells pretreated with 2 ng/ml TGF-β for 10 d.
d. Endocytic transport of fluorescently labeled hyaluronate probe as assessed by
quantitative colocalization with an early endosomal marker (EEA1) in PyMT-1099
WT or AGPS KO cells pretreated with 2 ng/ml TGF-β for 10 d.
e. ICP-MS of cellular iron following treatment with either hyaluronan or hyaluronidase in
PyMT-1099 WT or CD44 KO cells pretreated with 2 ng/ml TGF-β for 10 d.
f. Endocytic transport of dextran as assessed by quantitative colocalization with the
early endosomal marker EEA1 in pB3 cells.
g. Endocytic transport of dextran as assessed by quantitative colocalization with the
early endosomal marker EEA1 in PyMT-1099 WT or AGPS KO cells pretreated with
2 ng/ml TGF-β for 10 d.
h. Endocytosis of EGFR as assessed by quantitative colocalization of internalized EGF
with an early endosomal marker (EEA1) in PyMT-1099 WT or AGPS KO cells
pretreated with 2 ng/ml TGF-β for 10 days. Cells were treated with 2 ng/ml EGF.
All data shown as mean +/− SEM and statistical significance was calculated using
unpaired, two-tailed t-test; Examined n=10 fields of cells per experimental sample for all
endocytosis-related experiments and n=4 replicates for ICP-MS.
542
Fig. 4. Ether lipid deficiency impairs membrane biophysical properties.
a. Schematic of membrane tether pulling assay and fluorescence image showing a
tether pulled from the plasma membrane of a pB3 cell using an optically trapped 4
µm anti-Digoxigenin coated polystyrene bead.
b. Graph showing tether radius (R) and tether force measurements (f) in pB3 WT,
AGPS KO, and AGPS addback cells. All data shown as mean +/− SD.

c. Membrane tension measurements in pB3 WT, AGPS KO pretreated with 20µM of the
indicated ether phospholipid liposomes, and AGPS addback cells. All data shown as
mean +/− SEM.
d. Endocytic transport of fluorescently labeled hyaluronate probe as assessed by
quantitative colocalization with an early endosomal marker (EEA1) in pB3 WT or
AGPS KO cells pretreated with 20µM of the indicated ether phospholipid liposomes.
All data shown as mean +/− SEM.
e. Endocytic transport of fluorescently labeled transferrin as assessed by quantitative
colocalization with an early endosomal marker (EEA1) in pB3 WT or AGPS KO cells
pretreated with 20µM of the indicated ether phospholipid liposomes. All data shown
as mean +/− SEM.
f. GP values of C-Laurdan-labeled plasma membranes from pB3 WT, AGPS KO and
AGPS addback cells. Data is shown as mean GP +/− SD.
g. GP values of C-Laurdan-labeled intracellular membranes from pB3 WT, AGPS KO
and AGPS addback cells. Data is shown as mean GP +/− SD.
h. GP maps of C-Laurdan-labeled intracellular membranes from PyMT-1099 WT or
AGPS KO cells treated with 2 ng/ml TGF-β for 10 d. Data shown as mean GP +/−
SD.
i. Representative curves showing a leftward shift of the phase separation curve in
GPMVs from pB3 AGPS KO cells in comparison to wildtype pB3 control cells. This is
indicative of less stable phase separation upon loss of AGPS. Curves were
generated by counting >/= 20 vesicles/temperature/condition at >4 temperature. The
data was fit to a sigmoidal curve to determine the temperature at which 50% of the
vesicles were phase-separated (T_{misc}). Data shown as the average fits of 3
independent experiments. Inset showing a decrease in miscibility transition
temperatures (T_{misc}) upon loss of AGPS in pB3 cells. Graph shows the mean +/-
SEM of 3 independent experiments.
Statistical significance was calculated using unpaired, two-tailed t-test.
577
Fig. 5. Loss of ether lipids decreases metastasis and cancer cell stemness. See
also Extended Data Fig. S4.
a. Representative confocal images of extravasated tdTomato-labeled pB3 WT and
AGPS KO cells from an in vitro microvascular network established using HUVEC
(green) and normal human lung fibroblasts (unlabeled), over a time period of 24 h.
b. Quantification of extravasated tdTomato-labeled pB3 WT and AGPS KO cells from
an in vitro microvascular network established using HUVEC (green) and normal
human lung fibroblasts (unlabeled), over a time period of 24 h. Each datapoint
represents number of extravasated cells per device. Data is representative of two
independent biological replicates. Graph shows the mean +/- SEM and statistical
significance was calculated using unpaired, two-tailed t-test.
c. Quantification of extravasated tdTomato-labeled PyMT-1099 cell line derivatives
from an in vitro microvascular network established using HUVEC (green) and normal
human lung fibroblasts (unlabeled), over a time period of 24 h. Each datapoint
represents number of extravasated cells per device. Data is representative of two
independent biological replicates. Graph shows the mean +/- SEM and statistical
significance was calculated using unpaired, two-tailed t-test.

d. Representative IVIS images of overall metastatic burden in C57BL/6 female mice
following intracardiac injection of GFP-luciferized pB3 WT (n=5) and AGPS KO (n=5)
cells. Mean +/- SEM.
e. Quantification of overall metastatic burden in C57BL/6 female mice following
intracardiac injection of GFP-luciferized pB3 WT (n=5) and AGPS KO (n=5) cells.
Mean +/- SEM.
f. Representative images of H&E-stained sections of harvested kidneys from C57BL/6
female mice following intracardiac injection of pB3 WT or pB3 AGPS KO cells.
g. Gross images of primary tumors derived from pB3 WT control cells and pB3 AGPS
KO cells.
h. Tumor growth kinetics of primary tumors derived from pB3 WT control cells and pB3
AGPS KO cells. (n=5 mice per group).
i. Bar graph showing the average weight from primary tumors derived from pB3 WT
control cells and pB3 AGPS KO cells. Data shows the mean +/- SEM.
j. Estimated number of cancer stem cells (CSCs) per 10,000 cells as calculated by
extreme limiting dilution analysis (ELDA) software. Tumor-initiating capacity was
assessed following implantation of indicated amounts of pB3 WT or pB3 AGPS KO
cells into the mammary fat pad of C57BL/6 mice. *P* values, χ2 pairwise test.
k. Table showing the number of mice with palpable primary tumors at 121 d post
orthotopic implantation of PyMT-1099 WT or AGPS KO cells pretreated with 2 ng/ml
TGF-β for 10 d into NSG female mice.
l. Quantification of lung metastases for aforementioned experiment. Data shows the
mean number of lung metastases +/- SEM.
m. Representative images of H&E-stained lungs harvested from C57BL/6 female mice
following orthotopic injection of PyMT-1099 WT or AGPS KO cells pretreated with 2
ng/ml TGF-β for 10 d. Lungs were harvested after 121 d post-injection.
621
622   **EXTENDED DATA FIGURE LEGENDS**
623
<u>Extended Data Fig.1</u>
a. Schematic of peroxisomal-ether lipid biosynthetic pathway.
b. Cell viability following treatment with the GPX4 inhibitor RSL3 for 72 h. PyMT-1099
WT or AGPS KO cells were pretreated with TGF-β (2 ng/ml) for 10 d prior to assay.
Graph is representative of two independent biological replicates.
c. Amount in pmol of oxidized phosphatidylethanolamine (Oxi. PE) ether and ester
phospholipids in pB3 cells treated with RSL3 for 24 hours. Five biological replicates
per condition.
d. Immunoblot analysis for AGPS expression in mesenchymal-enriched pB3 WT, AGPS
KO, and AGPS addback cells. pB2 cells served as a control for expression of
epithelial-like markers.
635
<u>Extended Data Fig. 2</u>
a. Inductively coupled plasma-mass spectrometry (ICP-MS) of cellular iron in the
mesenchymal-enriched 687g WT and AGPS KO murine breast cancer cell line.
b. Immunoblot analysis of OVCAR8 AGPS KO, FAR1 KO or nontargeting sg (control)
cells.

Unless stated otherwise, all samples were analyzed in technical triplicates and shown
as the mean +/− SEM. Statistical significance was calculated using unpaired, two-tailed
t-test.
644
Extended Data Fig. 3
a. ICP-MS of cellular iron following treatment with either hyaluronan or hyaluronidase in
OVCAR8 WT or CD44 KO cells.
b. Endocytosis of EGFR as assessed by quantitative colocalization of internalized EGF
with an early endosomal marker (EEA1) in PyMT-1099 WT or AGPS KO cells
pretreated with 2 ng/ml TGF-β for 10 days. Cells were treated with 200 ng/ml EGF.
All data shown as mean +/− SEM and statistical significance was calculated using
unpaired, two-tailed t-test; Examined n=10 fields of cells per experimental sample for all
endocytosis-related experiments and n=4 replicates for ICP-MS.
654
Extended Data Fig. 4
a. Bright-field (top) and fluorescence images (bottom) showing reduced mesenteric
metastases from athymic nude mice injected with tdTomato-labeled OVCAR8 NT sg,
AGPS KO and FAR1 KO cells via the intraperitoneal route.
b. Representative IVIS images of overall metastatic burden in C57BL/6 female mice
following intracardiac injection of GFP-luciferized pB3 WT (n=5) and CD44 KO (n=6)
cells. Mean +/- SEM.
c. Quantification of overall metastatic burden in C57BL/6 female mice following
intracardiac injection of GFP-luciferized pB3 WT (n=5) and CD44 KO (n=6) cells.
Mean +/- SEM.
d. Table showing number of cells implanted per mice for limiting dilution assay.
666
METHODS
Cell lines
The pB3 MMTV-PyMT-derived murine breast cancer cell line was a kind gift from the
laboratory of Harold L. Moses[29]. 687g cells (also called EpCAM[Lo]Snail-YFP[Hi]) were
originally established from tumors that developed in the MMTV-PyMT-Snail-IRES-YFP
reporter mouse model, previously developed by the Weinberg lab[89]. pB3 and 687g cell
lines were cultured in 1:1 DMEM/F12 medium containing 5% adult bovine serum with
1% penicillin-streptomycin and 1% non-essential amino acids[29]. The PyMT-1099 murine
breast cancer cell line was a kind gift from the laboratory of Gerald Christofori and
cultured in DMEM supplemented with 10% fetal bovine serum, 1% penicillin-
streptomycin and 1% glutamine[26]. These cells were treated with 2 ng/ml of TGF-β for 10
days, prior to performing any subsequent analyses. OVCAR8 cells were obtained from
the laboratory of Joan Brugge and cultured in 1:1 MCDB 105 medium/Medium 199
Earle's Eagles medium supplemented with 10% fetal bovine serum and 1% penicillin-
streptomycin. All cells were cultured in a humidified incubator at 37°C with 5% CO_2. All
cells were negative for mycoplasma. Human cell line authentication (CLA) analysis of
OVCAR8 cells were performed by the Duke University DNA Analysis facility.
Established murine lines have not been STR profiled.
685
Animal studies

All animal studies were conducted according to the MIT Committee on Animal Care
protocol. For primary tumor growth studies: 1 million cells were resuspended in 20%
Matrigel/PBS and injected into the mammary fat pad of 6-8 weeks old female mice.
C57BL/6 mice (Jackson Laboratories) were used for in vivo experiments with pB3 cells.
NSG mice were used for in vivo experiments with PyMT-1099 cells. These cells were
pretreated with TGF-β (2 ng/ml) for 10 days prior to injection. Tumor size was measured
once a week using a vernier caliper and tumor volume was calculated using the
formula: Tumor volume = length x width^2/2, where length represents the largest tumor
diameter and width represents the perpendicular tumor diameter. For limiting dilution
tumor-initiating assays: pB3 WT or pB3 AGPS KO cells were resuspended in 20%
Matrigel/PBS and injected into the mammary fat pad of 6-8 weeks old female C57BL/6
mice (Jackson Laboratories) at the following dilutions: 100,000, 10,000, 1000, 100 cells.
Animals were assessed for palpable tumors after 39 days post injection. The estimated
number of CSCs was calculated using the extreme limiting dilution analysis (ELDA)
software. For experimental metastasis involving pB3 cell lines, 0.2 million cells were
resuspended in 200μl of PBS and injected into the left ventricle of 6-8 weeks old female
C57BL/6 mice. Metastatic burden was measured after 10 d post-injection via
bioluminescence in live animals using the IVIS Spectrum in vivo imaging system.
Images were analyzed using Living Image software (PerkinElmer). For OVCAR8 cells,
1.5 million cells were resuspended in PBS and implanted into 6-8 weeks old female
athymic nude mice (Jackson Laboratories) via intraperitoneal injections. Metastatic
burden was assessed after 6 weeks using a fluorescence dissecting microscope.
709
Generation of gene-edited cell lines using CRISPR/Cas9
With the exception of OVCAR8 cells, all AGPS KO single cell clones were generated via
transient transfection with mouse AGPS CRISPR/Cas9 KO Plasmids (Catalog no. sc-
432759, Santa Cruz) according to the manufacturer's instructions. GFP-positive cells
were sorted via fluorescence-activated cell sorting (FACS) into 96-well plates with one
cell per well and single cell clones were subsequently expanded. AGPS KO single cell
clones were assessed for loss of AGPS expression via western blot analysis. pB3
AGPS addback cells were generated by transducing AGPS KO single cell clone with
pLV[Exp]-Puro-EF1A-mAgps lentiviral vector (VectorBuilder). pB3 AGPS KO cells
expressing pLV[Exp]-Puro-EF1A-Stuffer_300bp (VectorBuilder) were established as
controls and noted as pB3 AGPS KO + EV in the manuscript. Lentivirus was produced
by transfecting HEK293T cells with viral envelope (VSVG, Addgene) and packaging
plasmids (psPAX2, Addgene). Viral supernatant was collected after 48 h and filtered
through a 0.45μm filter. Stably transduced cells were selected with 2 μg/ml puromycin.
OVCAR8 FAR1 KO and AGPS KO single cell clones as well as nontargeting control
cells, were established as previously described[19]. pB3 CD44 KO cells (bulk) were
generated using human CD44 CRISPR/Cas9 KO Plasmids (Catalog no. sc-419558,
Santa Cruz) according to the manufacturer's instructions. After 48 h post-transfection,
cells were sorted by flow cytometry for GFP positive cells, expanded in culture, and re-
sorted twice for CD44 negative cells using Alexa Fluor® 647 anti-mouse/human CD44
Antibody. Cells were maintained as bulk CD44 KO cells.
731
732

Plasmids	Source
pLV[Exp]-Puro-EF1A-Agps (mouse)	VectorBuilder
pLV[Exp]-Puro-EF1A-Stuffer_300bp	VectorBuilder
Mouse AGPS CRISPR/Cas9 KO Plasmids	Catalog no. sc-432759, Santa Cruz
Mouse CD44/HCAM CRISPR/Cas9 KO Plasmids	Catalog no. sc-419558, Santa Cruz
LentiCRISPRv2-puro-Nontargeting	Published[19]
LentiCRISPRv2-puro-human AGPS sg	Published[19]
LentiCRISPRv2-puro-human FAR1 sg	Published[19]
pLV-EF1A-eGFP-LUC	VectorBuilder
pCDH-EF1-Luc2-P2A-tdTomato	Plasmid #72486, Addgene

Lipidomics analysis

Lipid extraction for mass spectrometry lipidomics
Mass spectrometry-based lipid analysis was performed by Lipotype GmbH (Dresden,
Germany) as described[90]. Lipids were extracted using a chloroform/methanol
procedure[91]. Samples were spiked with internal lipid standard mixture containing:
cardiolipin 14:0/14:0/14:0/14:0 (CL), ceramide 18:1;2/17:0 (Cer), diacylglycerol
17:0/17:0 (DAG), hexosylceramide 18:1;2/12:0 (HexCer), lyso-phosphatidate 17:0
(LPA), lyso-phosphatidylcholine 12:0 (LPC), lyso-phosphatidylethanolamine 17:1 (LPE),
lyso-phosphatidylglycerol 17:1 (LPG), lyso-phosphatidylinositol 17:1 (LPI), lyso-
phosphatidylserine 17:1 (LPS), phosphatidate 17:0/17:0 (PA), phosphatidylcholine
17:0/17:0 (PC), phosphatidylethanolamine 17:0/17:0 (PE), phosphatidylglycerol
17:0/17:0 (PG), phosphatidylinositol 16:0/16:0 (PI), phosphatidylserine 17:0/17:0 (PS),
cholesterol ester 20:0 (CE), sphingomyelin 18:1;2/12:0;0 (SM), triacylglycerol
17:0/17:0/17:0 (TAG). After extraction, the organic phase was transferred to an infusion
plate and dried in a speed vacuum concentrator. The dry extract was re-suspended in
7.5 mM ammonium formiate in chloroform/methanol/propanol (1:2:4, V:V:V). All liquid
handling steps were performed using Hamilton Robotics STARlet robotic platform with
the Anti Droplet Control feature for organic solvents pipetting.

MS data acquisition
Samples were analyzed by direct infusion on a QExactive mass spectrometer (Thermo
Scientific) equipped with a TriVersa NanoMate ion source (Advion Biosciences).
Samples were analyzed in both positive and negative ion modes with a resolution of
$R_{m/z=200}=280000$ for MS and $R_{m/z=200}=17500$ for MSMS experiments, in a single
acquisition. MSMS was triggered by an inclusion list encompassing corresponding MS
mass ranges scanned in 1 Da increments[92]. Both MS and MSMS data were combined
to monitor CE, DAG and TAG ions as ammonium adducts; LPC, LPC O-, PC, PC O-, as
formiate adducts; and CL, LPS, PA, PE, PE O-, PG, PI and PS as deprotonated anions.

763 MS only was used to monitor LPA, LPE, LPE O-, LPG and LPI as deprotonated anions;
764 Cer, HexCer and SM as formiate adducts.
765
Data analysis and post-processing
Data were analyzed with in-house developed lipid identification software based on
LipidXplorer[93, 94]. Only lipid identifications with a signal-to-noise ratio >5, and a signal
intensity 5-fold higher than in corresponding blank samples were considered for further
data analysis. Simple imputation of missing values was performed by replacing missing
values with 0.5 * minimum non-zero value for each lipid assayed. Relative level of total
ether lipids was determined by summing the pmol value of all ether lipids identified
followed by normalization to total lipids.
774
Oxidized lipidomics
100,000 cells per condition were plated in 6-well plates 24 h prior to the experiment. For
1099, cells were treated with TGF-β for 10 d. pB3 cells were treated with 500 nM RSL3,
OVCAR8 cells with 2 µM ML210 and TGF-β-treated 1099 cells with 10 µM ML210 for 24
h. Cells were subsequently washed with 1x PBS and then with 150 mM ammonium
bicarbonate. Cells were then scraped and resuspended in 150 mM ammonium
bicarbonate and centrifuged at 300 g for 5 min. The supernatant was removed and cells
were resuspended in 1 mL of 150 mM ammonium bicarbonate. The solutions were
centrifuged at 12,000 RPM for 10 min and the supernatant was removed. 200 µL of 150
mM sodium bicarbonate was added to the pellet and samples were flash frozen in liquid
nitrogen. Cells were prepared in 5 independent biological replicates and lipidomics
analysis was performed on the same day for all the replicates. For lipidomics analysis,
the 200 µL cell lysates were spiked with 1.4 µL of internal standard lipid mixture
containing 300 pmol of phosphatidylcholine 17:0-17:0, 50 pmol of
phosphatidylethanolamine 17:0-17:0, 30 pmol of phosphatidylinositol 16:0-16:0, 50 pmol
of phosphatidylserine 17:0-17:0, 30 pmol of phosphatidylglycerol 17:0-17:0 and 30 pmol
of phosphatidic acid 17:0-17:0 and subjected to lipid extraction at 4 °C, as previously
described[95]. The sample was then extracted with 1 mL of chloroform-methanol (10:1) for
2 h. The lower organic phase was collected, and the aqueous phase was re-extracted
with 1 mL of chloroform-methanol (2:1) for 1 h. The lower organic phase was collected
and evaporated in a SpeedVac vacuum concentrator. Lipid extracts were dissolved in
100 µL of infusion mixture consisting of 7.5 mM ammonium acetate dissolved in
propanol:chloroform:methanol [4:1:2 (vol/vol)]. Samples were analyzed by direct
infusion in a QExactive mass spectrometer (Thermo Fisher Scientific) equipped with a
TriVersa NanoMate ion source (Advion Biosciences). 5 µL of sample were infused with
gas pressure and voltage set to 1.25 psi and 0.95 kV, respectively. PC, PE, PEO, PCOx
and PEOx were detected in the 10:1 extract, by positive ion mode FTMS as protonated
adducts by scanning m/z= 580–1000 Da, at $R_{m/z=200}$=280 000 with lock mass activated at
a common background (m/z=680.48022) for 30 s. Every scan is the average of 2 micro-
scans, automatic gain control (AGC) was set to 1E6 and maximum ion injection time
(IT) was set to 50 ms. PG and PGOx were detected as deprotonated adducts in the
10:1 extract, by negative ion mode FTMS by scanning m/z= 420–1050 Da, at
$R_{m/z=200}$=280 000 with lock mass activated at a common background (m/z=529.46262)
for 30 s. Every scan is the average of 2 micro-scans. Automatic gain control (AGC) was

set to 1E6 and maximum ion injection time (IT) was set to 50ms. PA, PAOx, PI, PIOx,
PS and PSOx were detected in the 2:1 extract, by negative ion mode FTMS as
deprotonated ions by scanning m/z= 400–1100 Da, at $R_{m/z=200}$=280 000 with lock mass
activated at a common background (m/z=529.46262) for 30 s. Every scan is the
average of 2 micro-scans, automatic gain control (AGC) was set to 1E6 and maximum
ion IT was set to 50 ms. All data was acquired in centroid mode. All lipidomics data
were analyzed with the lipid identification software, LipidXplorer[93]. Tolerance for MS and
identification was set to 2 ppm. Data were normalized to internal standards.
817

Immunoblotting
Cells were washed with ice-cold PBS and lysed in 1X Cell lysis buffer (Cell Signaling
Technology, Cat. #9803S) containing 1mM PMSF protease inhibitor (Cell Signaling
Technology, Cat. #8553S). Protein samples were prepared with NuPAGE LDS Sample
Buffer (Thermo Fischer Scientific, Cat. #NP0007), NuPage Sample Reducing Agent
(Thermo Fischer Scientific, Cat. #NP0004), and heated at 70 ºC for 10 minutes.
Samples were resolved by SDS-PAGE, transferred to nitrocellulose (Bio-Rad) and
blocked in 5% milk/TBST for 1 h at room temperature. Membranes were incubated
overnight with the respective primary antibodies at 4 ºC, washed with 1X TBST,
incubated with HRP-conjugated secondary antibodies and developed using
SuperSignal™ West Dura Extended Duration Substrate (Thermo Fischer Scientific, Cat.
#34076).
830

831   **Antibody information**
832   The table below indicates the antibodies used for western blotting and flow cytometry
833   analyses.

Antibody	Source	Catalog No.
E-Cadherin	Cell Signaling Technology	3195S
Zeb1	Cell Signaling Technology	3396S
N-cadherin	Cell Signaling Technology	13116S
AGPS	Invitrogen	PA5-56400
FAR1	Novus Bio.	NBP1-89847
Snail	Cell Signaling Technology	3879S
GAPDH	Cell Signaling Technology	2118S
CD44 (Western Blot)	Abcam	ab189524
Alexa Fluor® 647 anti-mouse/human CD44 Antibody	BioLegend	103017
Anti-rabbit IgG, HRP-linked antibody	Cell Signaling Technology	7074S
Anti-mouse IgG, HRP-linked antibody	Cell Signaling Technology	7076S

Cell viability assay

Cells were seeded in 96-well black clear bottom plates (Corning) at 2000 or 3000 cells (1099 +/- TGF-β) and 6000 cells (OVCAR8) per well. Approximately, 12-16 h post-seeding, cells were treated with various drug concentrations using an HP D300e Digital Dispenser unless stated otherwise. Cell viability was assessed at 72 h post-treatment by performing CellTiter-Glo Luminescent Cell Viability Assays (Promega) according to the manufacturer's instructions. Relative viability was calculated by normalizing to untreated controls unless stated otherwise. Non-linear regression models were applied to generate the regression fit curves using GraphPad Prism. Drug compounds were purchased as indicated: RSL3 (Selleck Chem), ML210 (Sigma Aldrich), and Liproxstatin-1 (Fisher Scientific). For experiments involving ferric ammonium citrate (FAC), FAC (Sigma) was prepared fresh in sterile 1× PBS and manually added directly to cell culture media at the indicated concentrations at the time of seeding into 96-well plates. Unless stated otherwise, cells were pretreated with FAC for 24 h prior to ML210 treatment.

Inductively coupled plasma mass spectrometry (ICP-MS)

Cells were treated for 24 h with Hyaluronic acid (Carbosynth, FH45321, 600-1000 kDa, 1 mg/mL) or Hyaluronidase (HD, Sigma-Aldrich, H3884, 0.1 mg/mL) as indicated. Glass vials equipped with Teflon septa were cleaned with nitric acid 65% (VWR, Suprapur, 1.00441.0250), washed with ultrapure water (Sigma-Aldrich, 1012620500) and dried. Cells were harvested and washed twice with 1× PBS. Cells were then counted using an automated cell counter (Entek) and transferred in 200μL 1× PBS to the cleaned glass vials. The same volume of 1× PBS was transferred into separate vials for the background subtraction, at least in duplicate per experiment. For tumor samples, small pieces of the tumors were added into pre-weighed cleaned glass vials. Samples were lyophilized using a freeze dryer (CHRIST, 22080). Glass vials with lyophilized tumor samples were weighed to determine the dry weight for normalization. Samples were subsequently mixed with nitric acid 65% and heated at 80°C overnight. Samples were diluted with ultrapure water to a final concentration of 0.475 N nitric acid and transferred to metal-free centrifuge vials (VWR, 89049-172) for subsequent ICP-MS analyses. Amounts of metals were measured using an Agilent 7900 ICP-QMS in low-resolution mode, taking natural isotope distribution into account. Sample introduction was achieved with a micro-nebulizer (MicroMist, 0.2 mL/min) through a Scott spray chamber. Isotopes were measured using a collision-reaction interface with helium gas (5 mL/min) to remove polyatomic interferences. Scandium and indium internal standards were injected after inline mixing with the samples to control the absence of signal drift and matrix effects. A mix of certified standards was measured at concentrations spanning those of the samples to convert count measurements to concentrations in the solution. Values were normalized against cell number or dry weight.

Iron measurements using Rhodox-M

The lysosome-specific fluorescent Fe(II) probe RhoNox-M was synthesized in 3 steps according to a previously published procedure [33]. ^{1}H NMR (300 MHz, CDCl$_3$) δ 7.93 (1H, d, J = 2.0 Hz), 7.45 (1H, dd, J = 8.5 Hz, 2.0 Hz), 7.40–7.30 (3H, m), 7.05 (1H, d, J

= 8.5 Hz), 6.90 (1H, d, *J* = 7.0 Hz), 6.80 (1H, d, *J* = 8.0 Hz), 6.50–6.44 (2H, m), 5.28–
5.35 (2H, m), 3.62 (6H, m), 2.97 (6H, s). MS (ESI) m/z: calcd. for $C_{24}H_{25}N_2O_3$ [M+H]$^+$
389.19, found: 389.35. Cells were incubated with 1 µM Rhonox-M for 1 h or lysotracker
deep red (Thermo Fisher Scientific L12492) according to the manufacturer's instructions
for 1 h. Cells were then washed twice with ice-cold 1× PBS and suspended in
incubation buffer prior to being analysed by flow cytometry. For each condition, at least
10000 cells were counted. Data were recorded on a BD Accuri C6 (BD Biosciences)
and processed using Cell Quest (BD Biosciences) and FlowJo (FLOWJO, LLC). The
signal for Rhonox-M was normalized against the signal of lysotracker of cells treated in
parallel.
890
Endocytosis experiments
Antibody against EEA1 was purchased from BD Biosciences (Catalog no. 610456).
Conjugated transferrin (mouse)-Alexa546, dextran-Alexa555, and EGF-Alexa555 were
purchased from Invitrogen. Cy3-conjugated hyaluronan was synthesized in-house. Cy2-
or Cy3-conjugated donkey antibodies against mouse IgG were purchased from Jackson
ImmunoResearch. For receptor-mediated endocytosis, cells were washed with serum-
free medium and then incubated in this medium with Cy3-conjugated hyaluronan (0.1
mg/ml), Alexa 555-conjugated EGF (either 2 ng/ml or 200 ng/ml), or Alexa 546-
conjugated transferrin (5 µg/ml) for 1 hr at 4° C. Cells were then washed to clear
unbound ligand, and shifted to 37 °C for times indicated in the figures. Cells were
stained for EEA1, followed by confocal microscopy to assess the arrival of ligand to the
early endosome. To assess fluid-phase uptake, Alexa 555-conjugated dextran (0.2
mg/ml) was added to complete medium and cells were incubated 37 °C for times
indicated in the figures. Cells were then stained for EEA1, followed by confocal
microscopy to assess the arrival of this probe to the early endosome.
906
Confocal microscopy
Colocalization studies were performed with the Zeiss Axio Observer Z1 Inverted
Microscope having a Plan-Apochromat 63× objective, the Zeiss LSM 800 with Airyscan
confocal package with Zeiss URGB (488- and 561-nm) laser lines, and Zen 2.3 blue
edition confocal acquisition software. For quantification of colocalization, ten fields of
cells were examined, with each field typically containing about 5 cells. Images were
imported into the NIH ImageJ v.1.50e software, and then analyzed through a plugin
software (https://imagej.net/Coloc_2). Under the 'image' tab, the 'split channels' option
was selected. Under the 'plugins' tab, 'colocalization analysis' option was selected, and
within this option, the 'colocalization threshold' option was selected. Menders Coefficient
was used for colocalization analysis. Colocalization values were calculated by the
software, and expressed as the fraction of protein of interest colocalized with EEA1.
919
Synthesis of HA-Cy3 probe
Hyaluronic acid (HA, 2 mg, Sigma 75044, Lot #BCBM2884) was dissolved in a 1:1
solution of dimethylsulfoxide (DMSO) and water (0.4 mL) for a stock concentration of 5
mg/mL. The polymer was sonicated under heating to ensure full solubilization. The HA
solution was then diluted into HEPES (50 mM final HEPES concentration for a total
reaction volume of 2 mL once all components are combined). Sulfo-Cyanine3 amine

(2.36 mg, Lumiprobe) was separately dissolved in DMSO (0.236 mL) for a stock
concentration of 10 mg/mL. N-(3-Dimethylaminopropyl)-N′-ethylcarbodiimide
hydrochloride (EDC, 0.253 mg, Sigma) was separately dissolved in 50 mM HEPES
(0.051 mL) for a stock concentration of 5 mg/mL. The HA and EDC solutions were then
combined under stirring, followed by addition of the dye solution. The reaction was
stirred, protected from light, at room temperature for 12 h. Following, unreacted dye was
removed via Amicon Ultra-0.5 Centrifugal Filters (Millipore Sigma). Manufacturer
guidelines were followed to select purification spin speeds and times: 14000 rcf, 15 min
per wash step (water) until washes were clear and colorless. The purified HA-Cy3 probe
was stored in water at 4 °C until used.
936
937   **Preparation of liposomes**
938   Ether lipid liposomes were prepared as previously described [19]. C18(Plasm)-20:4PE
939   (Catalog. no 852804) and C18(Plasm)-18:1PE (Catalog. no. 852758) were purchased
940   from Avanti Polar Lipids Inc.
941
942   **Characterization of liposomes**
943   Table shows the hydrodynamic diameter and polydispersity of liposomes used in this
944   study. The average and standard deviation of three technical repeats is provided. A
945   Malvern ZS90 Particle Analyzer was used for size measurements reported.

Liposome	Hydrodynamic diameter (nm)	Polydispersity index
C18(Plasm) - 20:4 PE	174.8 ± 2.25	0.273 ± 0.01
C18(Plasm) - 18:1 PE	128.7 ± 2.07	0.287 ± 0.03

946
Membrane tension
Tether pulling experiments were performed on a home-built optical trap, following
principles described elsewhere [65, 96]. Briefly, 4 µm anti-Digoxigenin coated polystyrene
beads (Spherotech) were trapped with a 1064 nm, Ytterbium laser (IPG Photonics)
focused through a 60x 1.2 NA objective (Olympus). Forces on the beads were
measured by the deflection of backscattered trapping laser light onto a lateral effect
position sensor (Thorlabs) and calibrated using the viscous drag method [97]. To measure
tether radii (R), cell lines were transiently transfected with a membrane-targeted
fluorescent protein (glycosylphosphatidylinositol-anchored eGFP, Addgene #32601)
using a TransIT-X2 transfection kit (Mirus). Tether radius was obtained by comparing
tether fluorescence to fluorescence counts from a known area of the parent cell
membrane, as described [48]. Tether force (f) and fluorescence measurements were
performed simultaneously. Membrane tension was calculated using the following
equation:
961
962    $$\text{Membrane tension, } \sigma = \frac{f}{4\pi R}$$
963
Ether lipid liposome reconstitution assays
Adherent cells were treated with ether lipid liposomes 16-18 h prior to performing
respective membrane tension or endocytosis assays. Lipid liposomes were added

directly to the culture medium for a final concentration of 20 µM. Cells were switched
from liposome-containing media to "extracellular imaging buffer" (HEPES buffer with
dextrose, NaCl, KCl, MgCl2, CaCl2) during membrane tension experiments.
970
Miscibility transition temperatures (T_{misc}) measurements
Miscibility transition temperatures (T_{misc}) measurements were performed as previously
reported [98, 99]. Briefly, cells were washed in PBS, and cell membranes were labeled with
5 µg/ml fluorescent disordered/nonraft phase marker FAST DiO (Thermo Fisher
Scientific) for 10 min on ice. Cells were then washed twice in GPMV buffer (10 mM
HEPES, 150 mM NaCl, 2 mM CaCl2, pH 7.4), and then incubated with GPMV buffer
supplemented with 25 mM paraformaldehyde (PFA) and 2 mM dithiothreitol (DTT) for 1
h at 37 °C. Vesicles were imaged at 40× on an inverted epifluorescence microscope
(Leica DMi8) under temperature-controlled conditions using a microscope stage
equipped with a Peltier element (Warner Instruments). GPMVs were imaged from 4° C-
28 °C, counting phase-separated and uniform vesicles at each temperature. For each
temperature, 25-50 vesicles were counted and the percent of phase-separated vesicles
were calculated, plotted versus temperature, and a fitted to a sigmoidal curve to
determine the temperature at which 50% of the vesicles were phase-separated (T_{misc}).
985
C-laurdan spectral imaging
C-Laurdan imaging was performed as previously described [98-102]. Briefly, cells were
washed with PBS and stained with 10 µg/mL C-Laurdan for 10 min on ice, then imaged
using confocal microscopy on a Leica SP8 with spectral imaging at 60× (water
immersion, NA= X) and excitation at 405 nm. The emission was collected as two
images: 420–460 nm and 470–510 nm. MATLAB (MathWorks, Natick, MA) was used to
calculate the two-dimensional (2D) GP map, where GP for each pixel was calculated as
previously described [102]. Briefly, each image was background subtracted and
thresholded to keep only pixels with intensities greater than 3 standard deviations of the
background value in both channels. The GP image was calculated for each pixel using
Eq.1. GP maps (pixels represented by GP value rather than intensity) were imported
into ImageJ. To calculate the average PM GP, line scans drawn across individual cells.
PM GP values were taken as peak GP values from the periphery of the cell, whereas
internal membranes were calculated as the average of all values outside the PM peak.
The average GP of the internal membranes was calculated by determining the average
GP of all pixels in a mask drawn on each cell just inside of the PM.

1002
$$GP = \frac{\sum_{420}^{460} I_x - \sum_{470}^{510} I_x}{\sum_{420}^{460} I_x + \sum_{470}^{510} I_x}$$

1003

1004 **Extravasation assay**

1005 *Cells and reagents*: Immortalized human umbilical vein endothelial cells (ECs)
1006 expressing BFP[82] were cultured in VascuLife VEGF Endothelial Medium (Lifeline Cell
1007 Technology). Normal human lung fibroblasts (FBs) (Lonza, P7) were cultured in
1008 FibroLife S2 Fibroblast Medium (Lifeline Cell Technology).
1009

Microfluidic device: 3D cell culture chips (AIM Biotech) were used to generate in vitro microvascular networks (MVNs). The AIM chip body was made of cyclic olefin polymer (COP) with a type of gas-permeable plastic serving as the bottom film. AIM Biotech chips contained three parallel channels: a central gel channel flanked by two media channels. Microposts separated fluidic channels and serve to confine the liquid gelling solution in the central channel by surface tension before polymerization. The gel channel was 1.3 mm wide and 0.25 mm tall, the gap between microposts was 0.1 mm, and the width of media channels was 0.5 mm.

Microvascular network formation: To generate perfusable MVNs, ECs and FBs were seeded into the microfluidic chip using a two-step method [103]. Briefly, ECs and FBs were concentrated in VascuLife containing thrombin (4 U/mL). For the first step seeding, the outer layer EC solution was made with a final concentration of 10×10^6/mL. After mixed with fibrinogen (3 mg/mL final concentration) at a 1:1 ratio, the outer layer EC solution was pipetted into the gel inlet, immediately followed by aspirating from the gel outlet, leaving only residual solution around the microposts. For the second step, another solution with final concentrations of 5×10^6/mL ECs and 1.5×10^6/mL fibroblasts was similarly mixed with fibrinogen and then pipetted into the same chip through the gel outlet. The device was placed upside down to polymerize in a humidified enclosure and allowed to polymerize at 37 °C for 15 min in a 5% CO_2 incubator. Next, VascuLife culture medium was added to the media channels and changed daily in the device. After 7 days, MVNs were ready for further experiments.

Tumor cell perfusion in MVNs: 1099 or pB3 cell line derivatives expressing pCDH-EF1-Luc2-P2A-tdTomato (Plasmid #72486, Addgene) were resuspended at a concentration of 1×10^6/mL in culture medium. To perfuse these tumor cells into in vitro MVNs, the culture medium in one media channel was aspirated, followed by injection of a 20 µL tumor cell suspension in the MVNs and repeated twice. Microfluidic devices were then placed at 37 °C for 15 min in a 5% CO_2 incubator for 15 min. After that, the tumor cell medium was aspirated from the media channels to remove the unattached cells, and Vasculife was replenished. Devices were then placed back to the incubator. 24 h later, devices were fixed, washed, and imaged using an Olympus FLUOVIEW FV1200 confocal laser scanning microscope with a 10× objective and an additional 2× zoom-in function. Z-stack images were acquired with a 5 µm step size. All images shown are collapsed Z-stacks, displayed using range-adjusted Imaris software, unless otherwise specified. Extravasation percentage was calculated by dividing the cell number of extravasated tumor cells with the total number of tumor cells in the same imaging region of interest.

Histology
Harvested tissues were fixed by incubating with 10% neural-buffered formalin (VWR Scientific) at 4°C for 16–18 h. Fixed samples were then transferred to 70% ethanol and submitted to Hope Babette Tang Histology Facility at the Koch Institute at MIT for paraffin-embedding and H&E staining. Metastatic burden was quantified using QuPath software [104] and Image J [105].

Statistical Analysis

For statistical analyses, Mann-Whitney U test or unpaired, two-tailed t-test were performed using GraphPad Prism.

Supplemental Data Fig. file S1: Lipidomic analysis of ether lipid deficient cells

Correspondence and requests for materials should be addressed to Robert A. Weinberg (weinberg@wi.mit.edu).

REFERENCES

1. Levental, I. & Lyman, E. Regulation of membrane protein structure and function by their lipid nano-environment. *Nat Rev Mol Cell Biol* **24**, 107-122 (2023).
2. Viswanathan, V.S. *et al.* Dependency of a therapy-resistant state of cancer cells on a lipid peroxidase pathway. *Nature* **547**, 453-457 (2017).
3. Hangauer, M.J. *et al.* Drug-tolerant persister cancer cells are vulnerable to GPX4 inhibition. *Nature* **551**, 247-250 (2017).
4. Mai, T.T. *et al.* Salinomycin kills cancer stem cells by sequestering iron in lysosomes. *Nat Chem* **9**, 1025-1033 (2017).
5. Stockwell, B.R. *et al.* Ferroptosis: A Regulated Cell Death Nexus Linking Metabolism, Redox Biology, and Disease. *Cell* **171**, 273-285 (2017).
6. Yang, W.S. & Stockwell, B.R. Ferroptosis: Death by Lipid Peroxidation. *Trends Cell Biol* **26**, 165-176 (2016).
7. Dixon, S.J. *et al.* Ferroptosis: an iron-dependent form of nonapoptotic cell death. *Cell* **149**, 1060-1072 (2012).
8. Gupta, P.B. *et al.* Identification of selective inhibitors of cancer stem cells by high-throughput screening. *Cell* **138**, 645-659 (2009).
9. Antoszczak, M. *et al.* Iron-Sensitive Prodrugs That Trigger Active Ferroptosis in Drug-Tolerant Pancreatic Cancer Cells. *J Am Chem Soc* **144**, 11536-11545 (2022).
10. Yang, W.S. *et al.* Regulation of ferroptotic cancer cell death by GPX4. *Cell* **156**, 317-331 (2014).
11. Doll, S. *et al.* FSP1 is a glutathione-independent ferroptosis suppressor. *Nature* **575**, 693-698 (2019).
12. Bersuker, K. *et al.* The CoQ oxidoreductase FSP1 acts parallel to GPX4 to inhibit ferroptosis. *Nature* **575**, 688-692 (2019).
13. Yang, W.S. & Stockwell, B.R. Synthetic lethal screening identifies compounds activating iron-dependent, nonapoptotic cell death in oncogenic-RAS-harboring cancer cells. *Chem Biol* **15**, 234-245 (2008).
14. Dixon, S.J. *et al.* Pharmacological inhibition of cystine-glutamate exchange induces endoplasmic reticulum stress and ferroptosis. *Elife* **3**, e02523 (2014).
15. Li, Q. *et al.* Inhibition of neuronal ferroptosis protects hemorrhagic brain. *JCI Insight* **2**, e90777 (2017).
16. Fang, X. *et al.* Ferroptosis as a target for protection against cardiomyopathy. *Proc Natl Acad Sci U S A* **116**, 2672-2680 (2019).

17.   Yu, Y. *et al.* Hepatic transferrin plays a role in systemic iron homeostasis and liver
ferroptosis. *Blood* **136**, 726-739 (2020).
18.   Friedmann Angeli, J.P. *et al.* Inactivation of the ferroptosis regulator Gpx4 triggers
acute renal failure in mice. *Nat Cell Biol* **16**, 1180-1191 (2014).
19.   Zou, Y. *et al.* Plasticity of ether lipids promotes ferroptosis susceptibility and
evasion. *Nature* (2020).
20.   Dean, J.M. & Lodhi, I.J. Structural and functional roles of ether lipids. *Protein Cell*
9, 196-206 (2018).
21.   Benjamin, D.I. *et al.* Ether lipid generating enzyme AGPS alters the balance of
structural and signaling lipids to fuel cancer pathogenicity. *Proc Natl Acad Sci U*
S A **110**, 14912-14917 (2013).
22.   Snyder, F. & Wood, R. Alkyl and alk-1-enyl ethers of glycerol in lipids from normal
and neoplastic human tissues. *Cancer Res* **29**, 251-257 (1969).
23.   Albert, D.H. & Anderson, C.E. Ether-linked glycerolipids in human brain tumors.
Lipids **12**, 188-192 (1977).
24.   Snyder, F., Blank, M.L. & Morris, H.P. Occurrence and nature of O-alkyl and O-
alk-I-enyl moieties of glycerol in lipids of Morris transplanted hepatomas and
normal rat liver. *Biochim Biophys Acta* **176**, 502-510 (1969).
25.   Cui, W., Liu, D., Gu, W. & Chu, B. Peroxisome-driven ether-linked phospholipids
biosynthesis is essential for ferroptosis. *Cell Death Differ* **28**, 2536-2551 (2021).
26.   Saxena, M., Kalathur, R.K.R., Neutzner, M. & Christofori, G. PyMT-1099, a
versatile murine cell model for EMT in breast cancer. *Sci Rep* **8**, 12123 (2018).
27.   Doll, S. *et al.* ACSL4 dictates ferroptosis sensitivity by shaping cellular lipid
composition. *Nat Chem Biol* **13**, 91-98 (2017).
28.   Kagan, V.E. *et al.* Oxidized arachidonic and adrenic PEs navigate cells to
ferroptosis. *Nat Chem Biol* **13**, 81-90 (2017).
29.   Ye, X. *et al.* Distinct EMT programs control normal mammary stem cells and
tumour-initiating cells. *Nature* **525**, 256-260 (2015).
30.   Pantopoulos, K., Porwal, S.K., Tartakoff, A. & Devireddy, L. Mechanisms of
mammalian iron homeostasis. *Biochemistry* **51**, 5705-5724 (2012).
31.   Muckenthaler, M.U., Rivella, S., Hentze, M.W. & Galy, B. A Red Carpet for Iron
Metabolism. *Cell* **168**, 344-361 (2017).
32.   Muller, S. *et al.* CD44 regulates epigenetic plasticity by mediating iron
endocytosis. *Nat Chem* **12**, 929-938 (2020).
33.   Niwa, M., Hirayama, T., Okuda, K. & Nagasawa, H. A new class of high-contrast
Fe(II) selective fluorescent probes based on spirocyclized scaffolds for
visualization of intracellular labile iron delivered by transferrin. *Org Biomol Chem*
12, 6590-6597 (2014).
34.   Houk, R.S. Mass spectrometry of inductively coupled plasmas. *Analytical*
Chemistry **58**, 97A-105A (1986).
35.   Figueroa, J.A., Stiner, C.A., Radzyukevich, T.L. & Heiny, J.A. Metal ion transport
quantified by ICP-MS in intact cells. *Sci Rep* **6**, 20551 (2016).
36.   Hoepken, H.H., Korten, T., Robinson, S.R. & Dringen, R. Iron accumulation, iron-
mediated toxicity and altered levels of ferritin and transferrin receptor in cultured
astrocytes during incubation with ferric ammonium citrate. *J Neurochem* **88**,
1194-1202 (2004).

37. Bauckman, K.A., Haller, E., Flores, I. & Nanjundan, M. Iron modulates cell
survival in a Ras- and MAPK-dependent manner in ovarian cells. *Cell Death Dis*
4, e592 (2013).
38. Gammella, E., Buratti, P., Cairo, G. & Recalcati, S. The transferrin receptor: the
cellular iron gate. *Metallomics* **9**, 1367-1375 (2017).
39. Al-Hajj, M., Wicha, M.S., Benito-Hernandez, A., Morrison, S.J. & Clarke, M.F.
Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci*
U S A **100**, 3983-3988 (2003).
40. Yan, Y., Zuo, X. & Wei, D. Concise Review: Emerging Role of CD44 in Cancer
Stem Cells: A Promising Biomarker and Therapeutic Target. *Stem Cells Transl*
Med **4**, 1033-1043 (2015).
41. Solier, S. *et al.* A druggable copper-signalling pathway that drives inflammation.
Nature **617**, 386-394 (2023).
42. Aruffo, A., Stamenkovic, I., Melnick, M., Underhill, C.B. & Seed, B. CD44 is the
principal cell surface receptor for hyaluronate. *Cell* **61**, 1303-1313 (1990).
43. Hua, Q., Knudson, C.B. & Knudson, W. Internalization of hyaluronan by
chondrocytes occurs via receptor-mediated endocytosis. *J Cell Sci* **106 (Pt 1)**,
365-375 (1993).
44. Bendris, N. *et al.* SNX9 promotes metastasis by enhancing cancer cell invasion
via differential regulation of RhoGTPases. *Mol Biol Cell* (2016).
45. Thottacherry, J.J., Sathe, M., Prabhakara, C. & Mayor, S. Spoiled for Choice:
Diverse Endocytic Pathways Function at the Cell Surface. *Annu Rev Cell Dev*
Biol **35**, 55-84 (2019).
46. Saric, A. & Freeman, S.A. Endomembrane Tension and Trafficking. *Front Cell*
Dev Biol **8**, 611326 (2020).
47. McMahon, H.T. & Boucrot, E. Molecular mechanism and physiological functions
of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **12**, 517-533 (2011).
48. Pinot, M. *et al.* Lipid cell biology. Polyunsaturated phospholipids facilitate
membrane deformation and fission by endocytic proteins. *Science* **345**, 693-697
(2014).
49. Howes, M.T. *et al.* Clathrin-independent carriers form a high capacity endocytic
sorting system at the leading edge of migrating cells. *J Cell Biol* **190**, 675-691
(2010).
50. Lakshminarayan, R. *et al.* Galectin-3 drives glycosphingolipid-dependent
biogenesis of clathrin-independent carriers. *Nat Cell Biol* **16**, 595-606 (2014).
51. Galbiati, F., Razani, B. & Lisanti, M.P. Emerging themes in lipid rafts and
caveolae. *Cell* **106**, 403-411 (2001).
52. Doherty, G.J. & McMahon, H.T. Mechanisms of endocytosis. *Annu Rev Biochem*
78, 857-902 (2009).
53. Boucrot, E. *et al.* Endophilin marks and controls a clathrin-independent endocytic
pathway. *Nature* **517**, 460-465 (2015).
54. Holst, M.R. *et al.* Clathrin-Independent Endocytosis Suppresses Cancer Cell
Blebbing and Invasion. *Cell Rep* **20**, 1893-1905 (2017).
55. Thottacherry, J.J. *et al.* Mechanochemical feedback control of dynamin
independent endocytosis modulates membrane tension in adherent cells. *Nat*
Commun **9**, 4217 (2018).

56. Baba, T. *et al.* Clathrin-dependent and clathrin-independent endocytosis are
differentially sensitive to insertion of poly (ethylene glycol)-derivatized cholesterol
in the plasma membrane. *Traffic* **2**, 501-512 (2001).
57. Kozlov, M.M. & Chernomordik, L.V. Membrane tension and membrane fusion.
Curr Opin Struct Biol **33**, 61-67 (2015).
58. Sitarska, E. & Diz-Munoz, A. Pay attention to membrane tension:
Mechanobiology of the cell surface. *Curr Opin Cell Biol* **66**, 11-18 (2020).
59. Wu, X.S. *et al.* Membrane Tension Inhibits Rapid and Slow Endocytosis in
Secretory Cells. *Biophys J* **113**, 2406-2414 (2017).
60. Gottlieb, T.A., Ivanov, I.E., Adesnik, M. & Sabatini, D.D. Actin microfilaments play
a critical role in endocytosis at the apical but not the basolateral surface of
polarized epithelial cells. *J Cell Biol* **120**, 695-710 (1993).
61. Dai, J. & Sheetz, M.P. Regulation of endocytosis, exocytosis, and shape by
membrane tension. *Cold Spring Harb Symp Quant Biol* **60**, 567-571 (1995).
62. Hirama, T. *et al.* Membrane curvature induced by proximity of anionic
phospholipids can initiate endocytosis. *Nat Commun* **8**, 1393 (2017).
63. Loh, J. *et al.* An acute decrease in plasma membrane tension induces
macropinocytosis via PLD2 activation. *J Cell Sci* **132** (2019).
64. Bucher, D. *et al.* Clathrin-adaptor ratio and membrane tension regulate the flat-to-
curved transition of the clathrin coat during endocytosis. *Nat Commun* **9**, 1109
(2018).
65. Dai, J. & Sheetz, M.P. Mechanical properties of neuronal growth cone
membranes studied by tether formation with laser optical tweezers. *Biophys J* **68**,
988-996 (1995).
66. Levental, K.R. *et al.* Lipidomic and biophysical homeostasis of mammalian
membranes counteracts dietary lipid perturbations to maintain cellular fitness.
Nat Commun **11**, 1339 (2020).
67. Halbleib, K. *et al.* Activation of the Unfolded Protein Response by Lipid Bilayer
Stress. *Mol Cell* **67**, 673-684 e678 (2017).
68. Ernst, R., Ejsing, C.S. & Antonny, B. Homeoviscous Adaptation and the
Regulation of Membrane Lipids. *J Mol Biol* **428**, 4776-4791 (2016).
69. Levitan, I. Evaluating membrane structure by Laurdan imaging: Disruption of lipid
packing by oxidized lipids. *Curr Top Membr* **88**, 235-256 (2021).
70. Veatch, S.L. *et al.* Critical fluctuations in plasma membrane vesicles. *ACS Chem
Biol* **3**, 287-293 (2008).
71. Thankamony, S.P. & Knudson, W. Acylation of CD44 and its association with lipid
rafts are required for receptor and hyaluronan endocytosis. *J Biol Chem* **281**,
34601-34609 (2006).
72. Li, P. *et al.* The iron chelator Dp44mT suppresses osteosarcoma's proliferation,
invasion and migration: in vitro and in vivo. *Am J Transl Res* **8**, 5370-5385
(2016).
73. Basuli, D. *et al.* Iron addiction: a novel therapeutic target in ovarian cancer.
Oncogene **36**, 4089-4099 (2017).
74. Guo, W. *et al.* An important role of the hepcidin-ferroportin signaling in affecting
tumor growth and metastasis. *Acta Biochim Biophys Sin (Shanghai)* **47**, 703-715
(2015).

75. Chi, Y. *et al.* Cancer cells deploy lipocalin-2 to collect limiting iron in
leptomeningeal metastasis. *Science* **369**, 276-282 (2020).
76. Tsujita, K. *et al.* Homeostatic membrane tension constrains cancer cell
dissemination by counteracting BAR protein assembly. *Nat Commun* **12**, 5930
(2021).
77. Lambert, A.W., Pattabiraman, D.R. & Weinberg, R.A. Emerging Biological
Principles of Metastasis. *Cell* **168**, 670-691 (2017).
78. Chen, M.B. *et al.* On-chip human microvasculature assay for visualization and
quantification of tumor cell extravasation dynamics. *Nat Protoc* **12**, 865-880
(2017).
79. Chen, M.B., Whisler, J.A., Jeon, J.S. & Kamm, R.D. Mechanisms of tumor cell
extravasation in an in vitro microvascular network platform. *Integr Biol (Camb)* **5**,
1262-1271 (2013).
80. Frose, J. *et al.* Epithelial-Mesenchymal Transition Induces Podocalyxin to
Promote Extravasation via Ezrin Signaling. *Cell Rep* **24**, 962-972 (2018).
81. Chen, M.B., Lamar, J.M., Li, R., Hynes, R.O. & Kamm, R.D. Elucidation of the
Roles of Tumor Integrin beta1 in the Extravasation Stage of the Metastasis
Cascade. *Cancer Res* **76**, 2513-2524 (2016).
82. Wan, Z. *et al.* A robust vasculogenic microfluidic model using human
immortalized endothelial cells and Thy1 positive fibroblasts. *Biomaterials* **276**,
121032 (2021).
83. Prince, M.E. *et al.* Identification of a subpopulation of cells with cancer stem cell
properties in head and neck squamous cell carcinoma. *Proc Natl Acad Sci U S A*
104, 973-978 (2007).
84. Lambert, A.W. *et al.* DeltaNp63/p73 drive metastatic colonization by controlling a
regenerative epithelial stem cell program in quasi-mesenchymal cancer stem
cells. *Dev Cell* **57**, 2714-2730 e2718 (2022).
85. Fruman, D.A., Meyers, R.E. & Cantley, L.C. Phosphoinositide kinases. *Annu Rev*
Biochem **67**, 481-507 (1998).
86. Rodriguez, R., Schreiber, S.L. & Conrad, M. Persister cancer cells: Iron addiction
and vulnerability to ferroptosis. *Mol Cell* (2021).
87. Suzuki, T. *et al.* Crucial role of iron in epigenetic rewriting during adipocyte
differentiation mediated by JMJD1A and TET2 activity. *Nucleic Acids Res* (2023).
88. Herr, C.Q. & Hausinger, R.P. Amazing Diversity in Biochemical Roles of Fe(II)/2-
Oxoglutarate Oxygenases. *Trends Biochem Sci* **43**, 517-532 (2018).
89. Dongre, A. *et al.* Epithelial-to-Mesenchymal Transition Contributes to
Immunosuppression in Breast Carcinomas. *Cancer Res* **77**, 3982-3989 (2017).
90. Surma, M.A. *et al.* Mouse lipidomics reveals inherent flexibility of a mammalian
lipidome. *Sci Rep* **11**, 19364 (2021).
91. Ejsing, C.S. *et al.* Global analysis of the yeast lipidome by quantitative shotgun
mass spectrometry. *Proc Natl Acad Sci U S A* **106**, 2136-2141 (2009).
92. Surma, M.A. *et al.* An automated shotgun lipidomics platform for high throughput,
comprehensive, and quantitative analysis of blood plasma intact lipids. *Eur J*
Lipid Sci Technol **117**, 1540-1549 (2015).

93.   Herzog, R. *et al.* A novel informatics concept for high-throughput shotgun
lipidomics based on the molecular fragmentation query language. *Genome Biol*
12, R8 (2011).
94.   Herzog, R. *et al.* LipidXplorer: a software for consensual cross-platform
lipidomics. *PLoS One* **7**, e29851 (2012).
95.   Sampaio, J.L. *et al.* Membrane lipidome of an epithelial cell line. *Proc Natl Acad*
Sci U S A **108**, 1903-1907 (2011).
96.   Ashkin, A. Acceleration and Trapping of Particles by Radiation Pressure. *Physical*
Review Letters **24**, 156-159 (1970).
97.   Kuo, S.C. & Sheetz, M.P. Optical tweezers in cell biology. *Trends Cell Biol* **2**, 116-
118 (1992).
98.   Levental, K.R. & Levental, I. Isolation of giant plasma membrane vesicles for
evaluation of plasma membrane structure and protein partitioning. *Methods Mol*
Biol **1232**, 65-77 (2015).
99.   Sezgin, E. *et al.* Elucidating membrane structure and protein behavior using giant
plasma membrane vesicles. *Nat Protoc* **7**, 1042-1051 (2012).
100.   Levental, K.R. *et al.* Polyunsaturated lipids regulate membrane domain stability
by tuning membrane order. *Biophys J* **110(8)**, 1800-1810 (2016).
101.   Levental, K.R. *et al.* omega-3 polyunsaturated fatty acids direct differentiation of
the membrane phenotype in mesenchymal stem cells to potentiate osteogenesis.
Sci Adv **3**, eaao1193 (2017).
102.   Sezgin, E., Waithe, D., Bernardino de la Serna, J. & Eggeling, C. Spectral
imaging to measure heterogeneity in membrane lipid packing. *Chemphyschem*
16, 1387-1394 (2015).
103.   Wan, Z. *et al.* A Robust Method for Perfusable Microvascular Network Formation
In Vitro. *Small Methods* **6**, e2200143 (2022).
104.   Bankhead, P. *et al.* QuPath: Open source software for digital pathology image
analysis. *Sci Rep* **7**, 16878 (2017).
105.   Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. NIH Image to ImageJ: 25 years
of image analysis. *Nat Methods* **9**, 671-675 (2012).
1313

Fig. 1

Fig. 2

Fig. 3

Fig. 4

a Membrane tension

$$\sigma = \frac{f}{4\pi R}$$

Fig. 5

Extended Data Fig.1

a

Peroxisome

ER

Fatty acid

Long chain
Fatty alcohol

FAR1

DHAP
Acyl-CoA

GNPAT

AGPS

Ether phospholipid
precursor

Ether phospholipid
precursor

Ether phospholipid

b

WT ⬤ AGPS KO
WT + TGF-β ⬤ AGPS KO + TGF-β

% Viability

0.0 0.1 0.2 0.3 0.4 0.5

[RSL3], μM

c

pB3

pmol

5×10^4
4×10^4
3×10^4
2×10^4
1×10^4
0

RSL3 − + − +

Oxi. (PE)   Oxi. (PE)
Ether-PLs   Ester-PLs

d

pB3

pB2 WT AGPS KO AGPS Addback

E-cadherin
Vimentin
Zeb1
Snail
Agps
Gapdh

Extended Data Fig. 2

a

p=0.0088

Fe per 1x10^6 cells (ng)

30
20
10
0

WT   AGPS KO

b

NT sg   AGPS KO   FAR1 KO

Agps
Far1
Gapdh

Extended Data Fig. 4

a

d

No. of cells implanted	100	1,000	10,000	100,000
WT	3/3	3/3	3/3	3/3
AGPS KO	0/3	3/3	3/3	3/3