1 Title

2 Sub-cone visual resolution by active, adaptive sampling in the human foveola

4 Authors

5 Jenny L. Witten, Veronika Lukyanova, Wolf M. Harmening*

7 Affiliations

8 Rheinische Friedrich-Wilhelms-Universität Bonn, Department of Ophthalmology, Bonn,
9 Germany

10 *Corresponding author: Jenny Lorén Witten, jennylwitten@gmail.com

12 Abstract

The foveated architecture of the human retina and the eye's mobility enable prime spatial
vision, yet the interplay between photoreceptor cell topography and the constant motion of
the eye during fixation remains unexplored. With *in vivo* foveal cone-resolved imaging and
simultaneous microscopic photo stimulation, we examined visual acuity in both eyes of 16
participants while precisely recording the stimulus path on the retina. We find that resolution
thresholds were correlated with the individual retina's sampling capacity, and exceeded what
static sampling limits would predict by 18 %, on average. The length and direction of
fixational drift motion, previously thought to be primarily random, played a key role in
achieving this sub-cone diameter resolution. The oculomotor system finely adjusts drift
behavior towards retinal areas with higher cone densities within only a few hundred
milliseconds to enhance retinal sampling.

25 Introduction

Assessing visual abilities was already important in historic times[1], and the precise
measurement of visual acuity, our ability to resolve fine spatial detail by eye, has great
importance for many real life scenarios and is up to this day the primary diagnostic tool to
determine visual function in a clinical and optometric setting. Quite surprisingly, the widely-
believed assumption that the packing density and arrangement of retinal photoreceptors at
the foveal center set the limit to this ability has never been experimentally confirmed.

Fovealization, the morphological and functional specialization of the cellular architecture of
the light sensitive retina optimizes the human eye for high-acuity daytime vision[2,3]. Within the
central one-degree diameter of the fovea, termed foveola, postreceptoral neurons are
displaced centrifugally and the area is free of potentially shadowing blood vessels and glia
cells[4,5]. The outer segments of foveolar cone photoreceptors are maximally thinned and
densely packed for peak spatial sampling[6–8], which at the same time makes these cells the
most difficult to study *ex vivo*[9] as well as *in vivo*[10]. Each foveolar cone synapses to one ON-
and one OFF-midget bipolar cell, which in turn synapse exclusively upon single ON- and
OFF-midget ganglion cells, a circuit that is adult like before birth[11]. This establishes an
undisturbed *private line* from individual foveal receptors to central processing stages.

Based on indirect comparisons between histological and psychophysical data, the
hypothesis that cone spacing imposes the fundamental limit for visual resolution has been
put forward[8,12]. It is well established that cone spacing, especially in the central fovea, is
highly variable between individuals[12–15], making general comparisons between acuity
measurements and foveolar density estimated from histological samples susceptible to error.
One of the main reasons why the hypothesis lacks direct experimental proof is that because,
under natural viewing conditions, both visual resolution and experimental access to foveal
photoreceptors is blurred by the imperfect optics of the human eye[16,17]. Here, we have
overcome the optical barrier of the human eye by employing adaptive optics cell-resolved *in*
vivo retinal imaging in conjunction with micro-psychophysics to study directly whether the
individual's mosaic of foveolar cones determines visual performance in a high-acuity
resolution task.

While acuity is assumed to be mainly limited by the resolving capacity of the eye's optics and
retinal mosaic, it is well established that, for different visual tasks, performance thresholds
can be substantially lower than the sampling grain of photoreceptors. This phenomenon has
been termed hyperacuity[18] and depends on the neural visual system's ability to extract
subtle differences within the spatial patterns of the optical image on the retina[19]. Thus, the
visual system already incorporates mechanisms to detect relative spatial offsets an order of
magnitude smaller than the spatial granularity of the retina. To make use of those fine
distinctions in a resolution task, the neuronal system needs to go beyond purely spatial
coding of incoming signals.

Unlike a camera, the visual system depends on temporal transients arising in the receptor's
cellular signals. Neurons in the retina, thalamus and later stages of the visual pathways
respond strongly to temporal changes[20,21]. Thusly, the fovealized retinal architecture in
humans is accompanied by a dynamic sampling behavior that, by quick and precise
movements of the eye, brings retinal images of objects of interest to land in the foveola[22].

Even during steady fixation, for example of a distant face or a single letter of this text,

incessant fixational eye movements slide tens to hundreds of foveolar photoreceptors across

the retinal image, thereby introducing temporal modulations that translate spatial activation

patterns into the temporal domain[23]. Small and rapid gaze shifts known as microsaccades

relocate gaze within the foveola during periods of fixation[22], and between microsaccades,

the eyes perform a more continuous, seemingly random motion termed fixational drift[24,25].

Computational work suggested that fixational eye motion would introduce noise and thus

impair visual acuity[26,27]. Contrarily, recent studies on human psychophysics demonstrated

fixational eye motion to be beneficial for fine spatial vision[25,28,29]. Especially drift motion has

been increasingly argued to not just be randomly refreshing neural activity, but rather

structuring it[25,30,31] and being under central control[32].

The incessant motion of the eye conveys fine spatiotemporal detail that requires deciphering

of continuously changing photoreceptor signals, which are linked by the geometry of the

photoreceptor array and by how the eye moves. For instance, luminance modulation in

individual cones will scale with drift length. Larger luminance variations on single receptors

also yields more neuronal activity within the range of temporal frequencies parvocellular

ganglion cells are sensitive to. Selective spatial frequencies can thus be amplified by varying

drift length[25]. While the neuronal mechanisms that generate fixational drift are still not fully

understood[33], its consequence to visual perception has been demonstrated. Drift was shown

to improve visual performance in resolution tasks[25,28,34]. Indeed, considerable differences in

ocular drift between individuals exist[31,35], and subjects exhibiting less drift were shown to

have better acuity[31]. If such differences are a consequence of an active, adaptive

mechanism, however, and how drift behavior is related to the photoreceptors that sample the

retinal image is unknown.

The direct experimental access to the foveolar center, when other limiting factors like image

blur or retinal motion are taken out of the equation or can be precisely measured, will allow

to confirm or reject the long-standing hypothesis about the individual limits of vision. This will

help to understand the fundamental physiological limitations of the visual system and will

have important implications for clinical studies of retinal health.

Results

Resolution is finer than single cone sampling limits

We investigated the limitations of the photoreceptor packing density on individual visual resolution acuity by overcoming the optical aberrations of the eye with adaptive optics scanning laser ophthalmoscopy (AOSLO), while simultaneously performing psychophysical measurements and recording the fixational retinal motion (Figure 1a, b and c). In a four-alternative forced-choice task, 16 healthy participants indicated the orientation of an E-optotype while inspecting the stimulus with their individually preferred fraction of foveolar photoreceptors. These cone photoreceptors were simultaneously imaged and it was later identified which cells contributed to resolving the stimulus (Figure 1c, f). A psychometric fit to the data expressed as percentage correct from 100 trials was used to compute visual acuity thresholds (see online Methods and Figure 1d,e). In this near diffraction-limited testing condition, participants reached visual acuity thresholds between 20.6 and 28.5 arcsec (mean ± SD: 24.1 ± 2.4 arcsec), which compares to 20/8 vision (logMAR = -0.4). All participants reached thresholds better than 20/10 vision (logMAR = -0.3), the last line of a typical clinical Snellen chart or projectors of acuity optotypes that are used in clinical as well as optometric daily routine.

Cone densities at the cone density centroid (CDC) ranged between 10,692 and 16,997 cones/deg^2, with an average density of 13,640 cones/deg^2 (Average peak cone density, PCD: 13,944 cones/deg^2, range: 10,823 to 17,309 cones/deg^2), comparable to previous reports[13,14,36–39]. The median sampling cone density ranged between 10,297 and 16,104 cones/deg^2 (mean: 13,149 cones/deg^2). Two experimental runs of the eyes with highest and lowest sampling density are exemplarily shown in Supplementary Movie 1 and 2. The two foveolar cone mosaic images were also visualized and overlayed with a Snellen E stimulus at average threshold size (Figure 2a). A static, theoretical prediction given by the Nyquist sampling limit would assume the high-density retina where each single cone diameter is smaller than the Snellen E's gap or bar is able to resolve the stimulus, whereas the low-density retina fails in identifying the correct orientation (schematic representation in Figure 2b). However, for our 788 nm testing condition, all participants reached individual resolution thresholds well below their Nyquist limit predicted by the spacing between rows of cones (Figure 2c, d). On average, visual acuity thresholds exceeded this theoretical prediction by 20 % and 16 % in dominant and non-dominant eyes, respectively. When participants performed the same resolution task with a longer infrared wavelength (840 nm) imaging background, the absolute thresholds were slightly higher and thus closer to the Nyquist limit. Visual acuity thresholds were on average 7 % below and 2 % above the Nyquist limit for dominant and non-dominant eyes, respectively. These absolute visual acuity thresholds

were the only case where noteworthy differences arose between the 788 nm and 840 nm

experimental condition. For all other analyses, we found qualitatively similar results for either

wavelength and therefore only report the 788 nm results throughout the manuscript.

For the first time, we could measure the direct relation between the individual foveolar cone

photoreceptor sampling density and participants visual resolution thresholds. We found the

diffraction limited visual acuity thresholds to be strongly correlated to the foveolar sampling

density in dominant as well as fellow eyes (Figure 2d). The higher the cone density, the

smaller the visual stimulus that could be resolved. The degree of correlation slightly differed

for dominant (r^2 = 0.45, p = 0.005) and non-dominant eyes (r^2 = 0.28, p = 0.036), suggesting

that up to 45 % of the variance in inter-subject visual acuity can be explained by the

individual cone sampling densities. Overall, participants reached significantly lower

thresholds with their dominant eyes (average: 1.5 arcsec, SD ± 1.1; paired t-test, p < 0.001).

Nevertheless, visual acuity thresholds were strongly correlated between dominant and non-

dominant eyes (r^2 = 0.80, p < 0.001, Figure 2e). To test whether the effect of different

absolute thresholds might be explained by underlying differences in the sampling cone

density, fellow eyes densities were compared to each other. Sampling densities had a very

strong correlation between fellow eyes (r^2 = 0.85, p < 0.001, Figure 2e), but did not differ

between right and left eyes (p = 0.38) nor when grouping them according to ocular

dominance (p = 0.88). This compares well to previous studies that also showed strong

correlations between fellow eyes regarding both anatomical[14] as well as functional[15]

characteristics. Dominant eyes had a median of 78 cones/deg^2 higher densities compared to

their fellow eyes. To account for the 1.5 arcsec difference in acuity thresholds, a much

higher density difference of about 1,500 cones/deg^2 would have been needed. Based on

these results, we conclude that the spatial arrangement of foveal cones can only partially

predict resolution acuity. In the following, we show that ocular motion and its associated

temporal modulations also influence visual resolution.

Ocular drift is an active sampling mechanism

As the eye drifts, a visual stimulus projected onto the retina is processed as a spatiotemporal

luminance flow. The stimulus itself as well as the extent of drift motion determine the

characteristics of modulation. By analyzing the exact retinal locations sampling the stimulus

we show the impact of the traveled path length first (Figure 3), followed by the direction of

drift motion and its relation to anatomical and functional landmarks (Figure 4). In our

experiments we revealed that participants kept coming back to the same few hundreds of

cone photoreceptors (Figure 3a and Figure 3 – figure supplement 1). To focus on the

characteristics and implications of drift eye motion, trials containing microsaccades during
stimulus presentation were excluded from the analyses. During the short stimulus duration
however, microsaccades rarely occurred, as participants tend to suppress their
microsaccades, likely because they can be detrimental to fine-scale discrimination[40,41]. Drift
motion patterns varied greatly across, but also within participants. Examples of drift motion
paths for the eyes that performed the smallest and largest drift motion, on average, show a
great variability in shapes as well as extent of motion (Figure 3b). In our analyses, we chose
the drift length (sum of piecewise vector lengths) as the prime metric to describe the ocular
drift motion, because the randomness underlying alternative metrics of drift eye movements
becomes increasingly questionable (see also Discussion). Across all participants and
experimental trials, drift lengths ranged between 2.5 and 17.2 arcmin, with a median length
of 6.5 arcmin (which corresponds to a velocity of 5 to 34.5 arcmin/sec, median: 13
arcmin/sec, Figure 3c). The drift lengths are slightly smaller than in previous non-AO studies,
which is attributable to the viewing situation. The participants were looking at a very small
imaging field within a completely dark periphery without distracting structures or stimuli. The
smallest drift movement performed was similar among eyes (range: 2.5 – 5.4 arcmin),
whereas the largest individual drifts differed more than three times as much (range: 7.7 –
17.2 arcmin). Therefore, the individual drift span was rather driven by the larger drift lengths
of an eye and there was a strong correlation between median drift length and drift range
(dominant eyes: $r^2 = 0.55$, $p = 0.002$, non-dominant eyes: $r^2 = 0.34$, $p = 0.02$, Figure 3d).

Between fellow eyes, which were measured consecutively, drift lengths had a very strong
correlation ($r^2 = 0.86$, $p < 0.001$, Figure 3e) with no significant difference between eyes
(paired t-test, $p = 0.2$). The median drift lengths of all eyes varied between 4.8 and 8.5
arcmin (mean ± SD: 6.6 ± 1.1 arcmin). Individual visual acuity thresholds were significantly
correlated with drift lengths (dominant: $r^2 = 0.25$, $p = 0.04$; non-dominant: $r^2 = 0.29$, $p = 0.03$,
Figure 3g), with a trend towards better visual acuity for small ocular drift motion. On a
photoreceptor resolved scale, this confirms recent findings which showed individual acuity
thresholds to be correlated with the drift motion during a non-AO acuity task, closely related
to the drift measured in a sustained fixation task [31].

Considering the previously shown correlation between visual acuity and sampling cone
density, one could assume those two aspects to go along with an increase of ocular drift for
lower cone densities, whereas higher densities potentially need less drift to translate the
stimulus over the same number of cones. However, we don't find the drift motion to be tuned
in a way to always let the stimulus slip across a similar number of cones. There was no
significant correlation between cone densities and drift length (Figure 3f, dominant: $r^2 = 0.07$,
$p = 0.3$; non-dominant: $r^2 = 0.06$, $p = 0.4$). Also, we do observe similar drift length across

stimulus sizes. We note however that in all our experimental trials, stimulus sizes were quite
similar. If drift length tuning existed, it may have been more pronounced with a larger
dispersion of stimulus sizes. In the following we show that drift direction is indeed tuned to
optimize sampling.

Drift is adaptive and directed

Ocular drift has long been assumed to be a persistent jittery motion that follows random
trajectories. Recent work showed that the amount of drift can vary and may be adapted to
the task that has to be performed[25,31]. We here investigated if beyond this, humans are able
to actively tune their ocular drift direction to exploit their prime spatial retinal processing
properties. We therefore registered the individual drift motion trajectories with the
photoreceptor mosaic, tracked them from the retinal location where the stimulus turned on
(onset) to where it turned off after 500 ms (offset), and related these trajectories to foveolar
landmarks (Figure 4a, b). Because of the individual retinal locations used for fixation before
stimulus onset, we registered that, across all eyes, drift motion occurred towards all
directions during stimulus inspection, and no general trend in drift eye movements towards a
particular cardinal direction across participants occurred (Figure 4c). Individual eyes,
however, showed different drift behavior mostly directed towards one or two of the four
quadrants. All four cardinal directions were represented. Participant $P8_{right}$, for example,
drifted towards the nasal or superior fovea in 90 % of all trials. $P14_{right}$, on the other hand,
drifted towards the temporal fovea in 75 % of all trials.

When the frame of reference was rotated in each trial to register the motion from the onset
location relative to the CDC, we found a clear directional bias in which the drift was likely to
move the stimulus closer to the CDC. The drift directionality was evaluated by measuring the
relative angle between drift onset to drift offset and drift onset to CDC. We observed a strong
trend of drift directionality; 49 % of all drift episodes moved the stimulus towards the CDC ±
45° (Figure 4d). The directionality was not pronounced directly after stimulus onset but
increased with presentation duration (Rayleigh test for circular non-uniformity, $p < 0.001$ for
all conditions, see Figure 4 – figure supplement 2). Among eyes, the individual fractions
ranged between 16 and 80 % of trials. Only two eyes drifted towards the CDC less
frequently than given by chance (Figure 4 – figure supplement 1). We computed the
directionality tuning as the ratio of relative drift towards the CDC ± 45° (purple quadrant in
Figure 4d) and the mean relative drift towards the 3 other quadrants. A ratio of 1 indicated
the same relative frequency of drift towards all cardinal directions, whereas for a tuning ratio
of 2 the retina moved the CDC towards the stimulus twice as often compared to each of the

other 3 cardinal directions. The directionality tuning ratios ranged between 0.6 and 11.8 with

a median value of 3. Directionality tuning ratios had a significant effect on how much the

resolution threshold exceeded the Nyquist limit. Participants with highly tuned drift reached

larger differences between the Nyquist limit and their visual acuity threshold (dominant eyes:

$r^2 = 0.45$, p = 0.01; non-dominant eyes: $r^2 = 0.27$, p = 0.04, Figure 4e). Drift directionality was

mostly similar between eyes, and if intra-ocular differences occurred, they were not related

to ocular dominance. Also, we did not observe an effect of training on drift directionality: one

of the two trained observers had a very strong drift directionality (7 and 11.8 in the dominant

and non-dominant eye, respectively) while the other one exhibited a tuning ratio below

average (2.1 and 2.3 in the dominant and non-dominant eye, respectively).

Next to the CDC, two other foveolar landmarks are often reported as anchor locations

describing the center of the fovea. We here show that the CDC has the strongest relevance

with respect to drift tuning. When relating the drift trajectories to the preferred retinal locus of

fixation (PRL) or the location of peak cone density (PCD), we found a weaker approximation

towards both. The retinae moved the stimulus towards the PRL or PCD location in 42 % or

35 % of all trials, respectively (Figure 4f). Therefore, the observed directionality was

strongest towards the CDC. In a considerable number of trials, the stimulus onset was

further displaced from all of the 3 retinal locations and therefore a directed drift motion

resulted in approximation towards CDC as well as PRL and PCD. Also, in some eyes, 2 or

all of these retinal locations lay very close together, which results in very similar effects.

Nevertheless, in some eyes with particularly stable fixation that had at least a few arcmin

distance between their PRL and CDC we repeatedly observed a stimulus onset close to PRL

followed by a directional drift towards CDC with a resulting stimulus offset closer to the CDC.

Across participants this also resulted in a significant reduction of the isoline contour area

(ISOA) size between stimulus onset and offset (p = 0.02, Figure 3 – figure supplement 1,

Figure 4b and Supplementary Movie 3). The median ISOA for stimulus onset locations was

92.5 arcmin^2 which was reduced to 68.2 arcmin^2 for stimulus offset locations. This decrease

in size of the area of all retinal landing points supports the view of a certain retinal cone or

very small area of a few arcmin^2 to be the target region of the drift eye motion in a resolution

task.

When we looked at how much the individual drift trajectory decreased the distance from

either location, the median distance convergence (onset/offset distance) towards CDC, PRL

and PCD was about 12 %, 7 % and 3 %, respectively. While no participant had an average

convergence of more than 30 % towards PRL or PCD, the maximum convergence ratio

towards CDC was about 50 %. An adaptive drift behavior was also found in the relative drift

lengths exhibited in each stimulus presentation. Although the individual drift lengths could

vary substantially from trial to trial, we found that, across all participants and experimental

trials, eyes exhibited significantly larger drift lengths when the stimulus onset location was

further away from the CDC ($\rho^2 = 0.06$, $p < 0.001$, Figure 4g). The onset distance was not

correlated with drift directionality (Figure 4g). Across all trials, the average sampling cone

density increased between stimulus onset and offset for most of the participants. This

sampling gain was computed as the ratio between the maximum sampling density during the

trial and the sampling density at the stimulus onset location. The sampling gain was

significantly correlated with the potential retinal sampling gain of individuals in dominant ($r^2 =$

0.35, $p = 0.02$) as well as non-dominant eyes ($r^2 = 0.50$, $p = 0.002$, Figure 4h). Observers

exploited on average 30 % of their potential sampling gain in both fellow eyes. Interestingly,

one observer combined all the previously described sampling features particularly strong in

his dominant eye (P08_R). It had a steep cone density gradient, exhibited strong directional

tuning towards the CDC and had large drift lengths for stimulus onsets far from the CDC.

This eye was excluded from the sampling gain analysis because fixation behavior differed by

more than 4 standard deviations from the group average.

Discussion

By using synchronous adaptive optics imaging and visual stimulation of the foveola, we find

that the human visual system is capable of resolving spatial orientation of E optotypes

smaller than a single photoreceptor diameter and uncover a fixational eye motor behavior

that optimizes retinal sampling in accordance with the individual photoreceptor mosaic.

Spatial vision, and in particular visual acuity, is the most tested and used performance metric

with a close relation to everyday vision. It provides the main behavioral outcome for clinical

studies of vision. Measured in daily routine or clinical studies, best corrected visual acuity of

young and healthy adults is usually between 20/20 and 20/12.5 (60 and 37.5 arcsec)[42,43].

Even if lower order aberrations are corrected by e.g. glasses or contact lenses, higher order

aberrations inherently blur the retinal image, depending on their magnitude[43]. Adaptive

optics induce a close-to-diffraction limited optical correction, where the optical improvement

is significantly correlated with an increase in visual acuity thresholds[17]. By correcting

aberrations with AOSLO, we measured Snellen-E thresholds that were up to half the size

(between 20/10 and 20/6.9; 30 to 20.6 arcsec) compared to the natural viewing condition.

This is slightly lower than previously presented data[42], very likely because of different

wavelengths used for experimentation (Figure 2c). It might be surprising to learn that the

neural machinery of human vision is able to resolve such tiny stimuli, because natural

viewing is blurred by the eye's optics. Even though observers are, to some degree, adapted

to their own aberrations[44], best subjective image quality is seen when on average 88 % of

the aberrations are corrected[45]. This may indicate that, under normal viewing conditions,

optical aberrations and not cone topography may play the dominant role in limiting the eye's

acuity.

By removing most aberrations in our experiments, we can study in how far resolution

thresholds are linked to or limited by the optimized but at the same time individual

morphology of the human foveola. While in the periphery, midget retinal ganglion cell

sampling dominates resolution, resolution of the foveal center was estimated to be governed

by the cone sampling limit[8,46]. By first-time direct experimental validation in the same

participants, we here confirm the hypothesis that the individual spacing of cones can predict

the resolution capacity of our foveola when optical influences are bypassed (Figure 2). We

found that the individual spatial arrangement of cones was highly correlated to the visual

acuity of participants and explains up to 45 % of its variance (Figure 2d). Eyes with higher

foveolar sampling capacity reached lower thresholds than eyes with less densely packed

cone photoreceptors. Moreover, all participants reached resolution thresholds that exceeded

the static Nyquist sampling limit when tested with near infrared, 788 nm light. Natural vision

is comprised of multiwavelength stimuli, thus, using 788 nm in isolation is at the top end of

our retinal sensitivity. In a first part of our study, participants also performed experiments

with 840 nm light. Thresholds were rather approximating the Nyquist limit with this longer

near-infrared wavelength (Figure 2c). The L- and M-cone photopigment absorbance for 840

nm is about 1.4 log unit lower than for 788 nm[47]. The decreased cone sensitivity combined

with a larger Airy-Disk size of about 7 % are likely to be detrimental for the longer, 840nm,

wavelength. We would expect a potential for even lower thresholds for shorter wavelengths.

Otherwise, a potential for lower thresholds is only expected in eyes with higher angular cone

densities. Perhaps contrary at first sight, this could potentially be the case for observers with

higher myopia. Myopic eyes, despite retinal stretching, generally have a higher angular

sampling density in and around the foveola, compared to emmetropes[13]. Therefore, we

would expect acuity thresholds to be lower for myopic participants, in the case that (a)

angular cone density is increased like previously suggested and (b) AO correction and

display resolution are still sufficient to completely resolve the foveolar cone mosaic.

Psychophysical data for more participants with higher myopia and longer axial lengths would

be needed to verify this assumption.

Theoretical predictions of the Nyquist resolution limit are implying stationary sampling. If the

retinal image is under-sampled, aliasing occurs at the frequency of the receptor mosaic,

which may obscure the original image, especially its orientation (compare example

snapshots in Fig. 1f). While prior knowledge of the stimulus has been shown to theoretically

help to de-alias under-sampled signals even in a static condition[48], we believe that retinal
image motion plays a significant role in deciphering orientation at the limits of spatial
sampling. Fixational eye movements continuously modulate the luminance flow on individual
cones and postreceptoral neuronal activity. Drift motion has long been presumed as a
random jitter, a result of limited precision of the oculomotor system[49,50]. More recent work
revealed that drift motion is neither random[51] nor detrimental due to the introduction of
noise[26,27], but rather a fine tuned motion, beneficial for psychophysical measures of visual
acuity in the parafovea[34] as well as foveola[25]. As also observed in other sensory organs[52],
neurons in the visual system are strongly selective not just for spatial patterns, but also for
temporally changing stimuli[53], a finding that is also supported by computational modeling,
suggesting that the visual system may utilize principles comparable to those used in
computational imaging for achieving super-resolution via camera motion[54]. Within the past
decades, the interdisciplinary term "geometrical super-resolution" which is devoted to the
filtering properties of sensor systems has become common[55]. These resolution advantages
may be achieved in the visual system by incorporating mechanisms that allow for the
recognition of positional differences smaller than a single cell. That such mechanism exist is
exemplified in a phenomenon known as hyperacuity. Fine localization discriminations of only
a few seconds of arc are performed by identification of the centroid of the retinal light
distributions[56] of the involved pattern components. In a diffraction limited resolution task, the
visual system seems to be able to translate the temporal luminance modulation in individual
photoreceptors by ocular drift to additional spatial information about the stimulus position
and shape. Contrary, the indirect suppression of natural fixational eye motion by retinal
stabilization techniques impairs visual acuity outside the foveolar center[25,28]. For prolonged
static stimulus presentations, retinal spiking decays over time, while drift motion keeps the
luminance change active, continuously refreshes the receptive field input and sustains
neuronal activity[23].

We found a significant correlation of drift motion and visual acuity thresholds between
individuals, indicating that drift motion may be one of the key elements in reaching sub-cone
resolution thresholds. Interestingly, acuity improved for smaller fixational drift and decreased
in participants who exhibited larger drift motion, on average. The fact that less drift is
beneficial to reach the lowest possible acuity thresholds reflects the characteristics of
spatiotemporal luminance changes introduced by smaller or larger drift motion. Smaller drifts
induce luminance changes with higher spatial frequencies and models of retinal ganglion cell
activity suggest a higher contrast sensitivity for high spatial frequency motion and less for
low spatial frequencies compared to a static retina[23,57]. This is supported by other recent

work which also showed that visual acuity thresholds can even be predicted from drift

magnitudes measured in a sustained fixation task[31].

There is evidence that fixational eye motion might have systematic components in primates.

A previous study in macaque monkeys revealed a systematic directional drift response only

a few dozens of milliseconds after various visual transients[58]. In our study, we reveal that a

certain drift directionality can not only be triggered by particular visual transients, but that

human observers are capable to adapt their drift direction to enact an oculomotor strategy

that takes advantage of the maximum resolution capacity provided within the retina. Our

participants precisely moved their eye to have the stimulus slip across the most densely

packed cone cells within their foveola. We hereby shed light on a mechanism that is

potentially particularly active during fine discrimination tasks. This confirms that drift can be

quickly adjusted in a continuous closed-loop control[59], while, as other recent work suggests,

being at the same time able to quickly switch to an open-loop process, as specific task

knowledge influences the dominant orientation of drift, even in the sudden absence of visual

information[60].

Yet, the underlying neuronal control of drift motion remains not fully understood. Recent

work suggested, based on brainstem recordings in rhesus monkeys, that the origin can be

found mostly upstream of the ocular motoneurons. It can likely be explained as diffusion in

the oculomotor integrator which is mainly driven by noise, but additionally affected by

mechanisms within the visual motor pathway (e.g. feedback mechanisms)[33]. An

incorporation of a visual feedback loop to that model was shown to modulate the statistics of

eye motion, given a time lag of about 100 ms (mainly due to synaptic processing delays, of

order 60-80 ms[58]). This fits our results well. Our presentation time of 500 ms sufficed for a

modulation of the fixational drift motion towards retinal areas of higher cone sampling (also

see Figure 4 – figure supplement 1-3). Our data supports the view that some aspects of the

statistics of drift motion can be influenced by the visual task[25,33,58,61]. The superior colliculus

seems to play a major role in modulating drift motion in a feedback loop to visual inputs[30]. It's

not only involved in controlling large eye motions[62] and microsaccades[63], but also reflects

neural responses to fixational drift that are likely a result of sensory input[64].

So, even though the CDC is displaced from the PRL in a way to be beneficial for natural

binocular vision[15], constant visual feedback allows to adapt the drift direction and therefore

also the task related PRL. Commonly, the term PRL is used for describing the retinal

location that is preferably used in fixational tasks. It is still a matter of debate what factors

drive the development of this very reproducible[15] retinal location and in how far it might

provide enhanced visual function. Sensitivity to small light spots in the foveola seems to be

rather plateau-like and not particularly pronounced at the PRL[65]. As recently shown, the PRL

slightly differs between different tasks but has a larger interindividual variability[41]. The here

shown results indicate that also when measuring visual resolution, the PRL is not

necessarily the center of the sampling drift motion. The directional drift motion leads to a

shift of the preferred retinal location for a resolution task towards the CDC (Figure 4 – figure

supplement 3 and Supplementary Movie 3). Previous work that compared active versus

passive fixation did not show a systematic offset in a similar experimental setup. However, 5

out of 8 participants also shifted their PRL in a Snellen E task closer to the CDC compared

to the PRL for fixating a static disk stimulus[41], the conditions that are best comparable to our

study. The main difference to our visual acuity experiments was that automatically paced

random time intervals between presentations (0.5 – 1.5 sec) were applied to not allow the

participants to anticipate the next trial whereas in our study, participants self-paced the

stimulus output to be able to prepare and focus for the next trial. It might be that this

extremely fine-tuned usage of the visual feedback loop can only be kept active for rather

short time intervals. By shifting the stimulus towards the CDC in 50 % of cases, the potential

sampling gain within individual eyes was exploited by 30%, on average, which goes along

with a cone density increase of 3 % or 285 cones/deg^2. Even though this increase in cone

density alone would not account for the difference between acuity thresholds and Nyquist

limit, this and the simultaneous spatiotemporal luminance modulation contribute to achieving

sub-cone visual acuity thresholds.

Between fellow eyes we found very strong correlations for all the measured parameters.

While drift lengths and directionality as well as cone densities are very symmetric between

dominant and non-dominant eyes (Figure 2e and 3e), significantly lower acuity thresholds of

1.5 arcsec, on average, were observed in the dominant eyes of participants (Figure 2e). The

dominant eyes' visual input has a tendency to be preferred during binocular viewing, but has

not been shown to exhibit relevant differences in visual function in healthy eyes with low

refractive errors[66,67]. Partially this may be due to limited accuracy in the mainly used clinical

methods (e.g. Snellen Chart or projection have ~ 10 arcsec steps between optotype rows).

This very fine binocular difference between eyes emphasizes that some remaining factors

which especially comprise the neural postprocessing steps, also play an important role and

may facilitate the slight functional advantage of dominant eyes.

For clinical studies of retinal health and in new therapeutical approaches, photoreceptor

health and visual acuity can be related to other more standard clinical measures as OCT-

derived measures of outer segment length or retinal thickness which have been shown to

serve for estimates of cone density[68]. Therefore, building a larger dataset on photoreceptor

resolved foveolar maps and associated visual function measures may help to, on the one

hand, better understand the interplay between structural and functional changes to draw

conclusions about disease progression, intervention efficiency or the interpretation of retinal

imaging data in studies aimed at vision restoration. On the other hand, detailed examination

of psychophysical measures with knowledge about the exact neural sampling characteristics

offers a great potential to answer further questions about e.g. resolution limits in myopia, the

effect of image stabilization in the very center of the foveola or implications for binocular

viewing that could previously only be hypothesized. The awareness of the oculomotor

system being able to finely adjust the drift motion behavior for a particular task may guide

future interpretation of fixational eye motion.

Material and Methods

Participants

A total of 38 participants with White ethnicity underwent a preliminary screening where ocular biometry, ophthalmologic status, fixational eye motion and adaptive optics correction as well as foveolar image quality were tested. From those, 20 participants with normal ophthalmologic status, resolvable foveolar cones and ocular anatomy that allowed for a 7 mm pupil aperture during experimentation were chosen for subsequent examination. All 6 male and 14 female observers (17 adults [age: 18 – 42], 3 children [age: 10, 12 and 14]) had no or only mild refractive errors (SE: ± 2.5 diopters). The children and 15 adults were naïve participants and two adults were experienced observers. More detailed cone topography and eye motion characteristics of the here studied population have been shown previously[15]. The experiments were conducted under two different light conditions (16 participants 788 nm, 12 participants 840 nm). Eight participants took part in both experimental conditions. We mainly report the data acquired for the 788 nm condition in this manuscript and show 840 nm data for comparison where noteworthy differences arise.

Written informed consent was obtained from each participant and all experimental procedures adhered to the tenets of the Declaration of Helsinki, in accordance with the guidelines of the independent ethics committee of the medical faculty at the Rheinische Friedrich-Wilhelms-Universität of Bonn.

Ocular dominance

Ocular dominance was determined by a Miles Test prior to pupil dilation and visual acuity testing. The experimenter stood in a distance of 6 m in front of the participant and asked them to form a small opening between thumbs and forefingers with both hands. The participant was then asked to extend their arms in front of them to look through the formed hole at the experimenter's face with both eyes open. This procedure was conducted 3-5 times to determine the dominant (= uncovered) eye in a 3/3 or at least 3/5 condition.

AOSLO retinal imaging

In vivo images of the complete foveolar cone mosaic were recorded using a custom-built adaptive optics scanning laser ophthalmoscope (AOSLO). The general setup of the AOSLO has been described previously[69] and pertinent differences as well as the method of determination of the preferred retinal locus of fixation (PRL) have been described in a recent publication[15].

In brief, the front-end of the Adaptive Optics Scanning Laser Ophthalmoscope (AOSLO) was
equipped with three f = 500mm focal telescopes. These telescopes were specifically
designed for point-scanning an adaptive optics-corrected focal light spot across the retina,
ensuring diffraction-limited resolution in both incident and reflected beams. The system
incorporated a magnetic actuator-driven deformable mirror (DM97-07, 7.2mm pupil diameter,
ALPAO, Montbonnot-Saint-Martin, France) positioned in a retinal conjugate plane. The
deformable mirror was controlled by the wavefront error signals from a 25x25 lenslet Shack
Hartmann sensor (SHSCam AR-S-150-GE, Optocraft GmbH, Erlangen, Germany) in closed-
loop. Imaging and wavefront correction utilized wavelengths of either 788 nm (±12 nm) or
840 nm (±12 nm) light, achieved through serial dichroic and bandpass filtering of a
supercontinuum source (SuperK Extreme EXR-15, NKT Photonics, Birkerød, Denmark). The
imaging field of view was 0.85 x 0.85 degrees of visual angle. The digital lateral resolution
was about 0.1 arcmin, the size of one pixel in the recorded videos and images. Light
reflected from the retina was detected by a photomultiplier tube (PMT, H7422-50,
Hamamatsu Photonics, Hamamatsu, Japan), positioned behind a confocal pinhole (Pinhole
diameter = 20 mm, equivalent to 0.47 (840nm) and 0.5 (788nm) Airy disk diameters).
Continuous sampling of the PMT signal was carried out using a field programmable gate
array (FPGA), resulting in a 512 x 512-pixel video at 30 Hz (600 pixels per degree of visual
angle). Through rapid acousto-optic intensity modulation of the imaging lights, the square
AOSLO imaging field was used as retinal display, where each pixel could be individually
controlled to produce the visual stimuli.

Cone map generation and computation of sampling characteristics

The best PRL videos acquired were selected to create spatially registered, high signal-to-
noise ratio images of the foveal center, which served as master retinal images for cone
labeling as well as referencing of stimulus motion trajectories. This study includes only
participants for whom the master retinal image was of sufficient quality to label all cones
across the image. Cone centers were identified and labeled semi manually, as previously
described[15,70]. Cone density was computed in two different ways. First, for deriving landmark
metrics of the foveolar cone map, we then computed Voronoi tessellation, estimating a patch
with certain area for each individual cone and summed the nearest 150 cone patches around
each image pixel. The number of cells was divided by the resulting area to derive a pixel-
resolved map of cone densities. Based on this map, the peak cone density (PCD) is defined
as the highest cone density value of the map with it's according retinal location. The cone
density centroid (CDC) is computed as the weighted centroid of the 20th percentile of highest
cone densities within the map. We refer to the CDC as the anatomical center and the anchor

for further spatial analyses in this study. The CDC has been shown to be a more robust and
reproducible metric to describe the anatomical center than the more routinely reported peak
cone density (PCD). While the PCD has value in reporting its quantity, namely the maximum
cone density of a retina, using it as a landmark is however not advised, for it being too
vulnerable against small changes in the analysis of cone density[15,71].

Second, for analyzing the relation between individual sampling limits and resolution acuity,
cone density was computed based on the cone cells contributing to the sampling process.
To identify the cones interacting in stimulus sampling, a simple model of cone light capture
was employed. Each cone was described by an associated light acceptance aperture with its
diameter estimated as 48 % of the average spacing between the cone and all of its
neighbors. The efficiency of the aperture along its diameter was approximated as Gaussian
profiles. Also, a model of the stimulus retinal image was computed by convolving the eye's
point spread function (diffraction limited at 788 nm for a 7 mm pupil) with the stimulus
bitmap. The complete two-dimensional model of cone apertures was then multiplied by
models of the presented stimuli to arrive at the cone-level light distribution based on the
different stimulus positions, sizes and orientations. The light distribution within each cone
was integrated across the entire cone aperture. This value was then normalized to the
degree to which the aperture was filled. Cone stimulation was considered to be maximal if
the entire aperture was filled. Using this method, a cone activation pattern could be
generated for each point in time (e.g. Figure 1f). To arrive at a task-related cone density
estimate for each frame (sampling cone density), the number of cones identified to interact
with the stimulus was divided by their summed cone area. In the presented analyses, the
median sampling density of all trials is analyzed and standard deviations are shown as grey
lines (Figure 2d, e). This stimulus related cone density was chosen to closely represent the
sampling process; however, the results do not qualitatively differ from using the cone density
map based on the 150 nearest cones.

We assumed a perfect hexagonal cell mosaic to estimate the average inter-cone-distance
(ICD) between neighboring cells and to compute the theoretical Nyquist sampling limit, which
is based on the spacing between rows of cones, and given by $N = \frac{\sqrt{3}}{2} \times ICD$.

Experimental procedures

For psychophysical acuity testing, participants reported the orientation of a Snellen-E
stimulus in a four-alternative forced-choice (4 AFC) task under unrestricted eye motion.
Psychophysical experiments were performed monocularly in both eyes. The non-dominant
eye was tested first and the dominant eye after a 15-30 minutes break. This protocol was

chosen because in pilot experiments in 7 participants (which were performed with a random

order) less time was needed and hence less fatigue was reported by the participants when

the second eye was the dominant one. In these pilot experiments, the same qualitative

difference of acuity thresholds between non-dominant and dominant were found.

Mydriasis and cycloplegia were established by two drops of 1% tropicamide, instilled into the

eyelid about 25 and 20 minutes prior to experiments. If experimentation took longer than 40

minutes, another drop of tropicamide was instilled. A customized dental impression mold

(bite bar) was used to immobilize and adjust the head position and thus to align the

participants eye in front of the imaging system to ensure optimal adaptive optics correction

and image quality. The participants were encouraged to take breaks at any time. We found

that proper resting is one of the most crucial factors during the rather complex AOSLO

experimentation. Frequent breaks ensure constant, high-level compliance and excellent

image quality as the basis for artefact-free and reproducible results.

Before recording experimental runs, each participant performed 3 test runs to get used to the

test procedure and the appearance of the stimuli. The stimuli were displayed as "off-stimuli"

on the infrared background by switching the displayed intensity via an acousto-optic

modulator [72] (AOM, TEM-250-50-10-840-2FP, Brimrose, Sparks Glencoe, MD, USA)(Figure

1a). Because of ocular diffraction, the stimulus contrast varied between 0.61 and 0.80 for an

18 arcsec versus 36 arcsec gap sized stimulus (3 and 6 pixel of the scanning raster,

respectively). The visual acuity testing followed the Bayesian adaptive procedure QUEST[73–

75]. Stimulus progression was self-paced by the participant. The stimuli were presented for

500 ms to avoid limitations by insufficient temporal summation[76]. Around each trial, a one

second AOSLO video was recorded, with the stimulation onset at around 300ms after video

onset. Visual acuity thresholds were estimated by pooling results from 5 consecutively run

staircases, with each containing 20 trials. A psychometric function was fitted using

psignifit4[77] to derive threshold estimates for further analysis. The expected threshold

variance is described and visualized by the 95 % confidence interval (Figure 1d, e and 2d).

Video processing and eye motion analysis

The AOSLO used a raster scanning technique where each frame was acquired over time.

The recorded videos were stabilized after psychophysical testing using custom settings

within the *MATLAB* based stabilization software from Stevenson et al.[78]. To acquire eye

traces at higher temporal resolution than the 30 Hz frame rate, each frame of the AOSLO

movie is broken into 32 horizontal strips of 16 pixels height and cross-correlated against a

reference frame. The reference frame was generally chosen automatically and exchanged

by a manually chosen frame in cases were stabilization failed despite good overall image
quality. This method allowed extraction of eye motion traces at temporal frequencies up to
960 Hz.

The frame-wise (30 Hz) stimulus position was encoded as a white cross marker in each
video. As single strip alignments can have small errors due to noise in the strip or retinal
torsion (particularly affecting the horizontal motion estimate)[79], we compute the average
offsets from the cross-containing strip and 2 previous/subsequent strips. These steps yielded
more accurate trajectories in retinal coordinates for every trial. All individual trial AOSLO
frames and the corresponding trajectories are then referenced to the single master retinal
image used for cone map generation.

To quantify the retinal motion across the stimulus, drift length was defined as the
concatenated vector sum of all frame-wise motion vectors within the 500 ms stimulus
duration (see also Figure 3c). Trials that contained microsaccades or blinks during stimulus
presentation were excluded from further analyses. Microsaccade occurrence varied highly
between participants (mean ± SD: 14 ± 10 % of trials, range: 2 % - 41 %). If not stated
differently, we here report the median drift length of all trials for individual eyes (e.g. of all
traces shown in Figure 3a). To quantify drift direction, the angle between each trajectory's
starting coordinate (coordinate center in Figure 4c) and end coordinate was computed. To
check for potential motion bias, the drift angles were first analyzed in retinal coordinates
(Figure 4c), and then as the relative angle, $\theta_{CDC,}$ formed between the drift vector and the line
connecting the retinal onset location and the CDC (Figure 4d). To compare directionality
towards other locations of interest, the same was done for PRL and PCD locations (Figure
4f).

Statistical information

All statistical analyses were conducted using custom written MATLAB code and significance
levels were set at 0.05. To assess the normal distribution of the dataset, a two-sided
Shapiro–Wilk test was employed. This test is recognized to be appropriate for small sample
sizes. The paired samples t-test was utilized to assess whether there were significant
differences between the means of normally distributed paired observations. For non-
parametric data, the Wilcoxon Signed-Rank test was employed. Linear correlations were
computed to examine the relationships between variables. For variables demonstrating
normal distribution, Pearson's correlation coefficient was employed, while for non-normally
distributed data, Spearman's rank correlation coefficient was utilized. Pearson's correlation
is sensitive to linear relationships, assuming bivariate normality, whereas Spearman's

correlation is a non-parametric measure suitable for monotonic relationships and is robust

against outliers and non-normal distributions.

Author contributions

J.L.W and W.M.H conceived the research idea and developed the data analysis pipeline.

J.L.W performed the experiments and data analyses. J.L.W, V. L. and W.M.H discussed the

results and wrote the manuscript.

Competing interests

The authors declare no competing interests.

Acknowledgments

This work is supported by the German Research Foundation, Emmy Noether-Program

HA5323/5-1; Carl Zeiss Foundation, HC-AOSLO; Novartis Pharma GmbH, EYENovative

research award and the Open Access Publication Fund of the University of Bonn.

References

1. Bohigian, G. M. An Ancient Eye Test-Using the Stars. *Surv. Ophthalmol.* **53**, 536–539 (2008).

2. Caves, E. M., Brandley, N. C. & Johnsen, S. Visual Acuity and the Evolution of Signals. *Trends Ecol. \& Evol.* **33**, 1–15 (2018).

3. Tuten, W. S. & Harmening, W. M. Foveal vision. *Curr Biol* **31**, R701–R703 (2021).

4. Hendrickson, A. E. & Yuodelis, C. The morphological development of the human fovea. *Ophthalmology* **91**, 603–612 (1984).

5. Syrbe, S. *et al.* Müller glial cells of the primate foveola: An electron microscopical study. *Exp. Eye Res.* **167**, 110–117 (2018).

6. Williams, D. R. & Coletta, N. J. Cone spacing and the visual resolution limit. *J. Opt. Soc. Am. A* **4**, 1514–1523 (1987).

7. Hirsch, J. & Curcio, C. A. The spatial resolution capacity of human foveal retina. *Vision Res.* **29**, 1095–1101 (1989).

8. Rossi, E. A. & Roorda, A. The relationship between visual resolution and cone spacing in the human fovea. *Nat Neurosci* **13**, 156–157 (2010).

9. Curcio, C. A., Packer, O. & Kalina, R. E. A whole mount method for sequential analysis of photoreceptor and ganglion cell topography in a single retina. *Vision Res.* **27**, 9–15 (1987).

10. Rossi, E. A. *et al.* Imaging retinal mosaics in the living eye. *Eye* **25**, 301–308 (2011).

11. Zhang, C. *et al.* Circuit reorganization shapes the developing human foveal midget connectome toward single-cone resolution. *Neuron* **108**, 905–918 (2020).

12. Curcio, C. A., Sloan, K. R., Kalina, R. E. & Hendrickson, A. E. Human photoreceptor topography. *J. Comp. Neurol.* **292**, 497–523 (1990).

13. Wang, Y. *et al.* Human foveal cone photoreceptor topography and its dependence on eye length. *Elife* **8**, 1–21 (2019).

14. Cava, J. A. *et al.* Assessing Interocular Symmetry of the Foveal Cone Mosaic. *Investig. Opthalmology Vis. Sci.* **61**, 1–11 (2020).

15. Reiniger, J. L., Domdei, N., Holz, F. G. & Harmening, W. M. Human gaze is systematically offset from the center of cone topography. *Curr. Biol.* **31**, 4188–4193 (2021).

16. Campbell, F. W. & Green, D. G. Optical and retinal factors affecting visual resolution. *J. Physiol.* **181**, 576–93 (1965).

17. Marcos, S., Sawides, L., Gambra, E. & Dorronsoro, C. Influence of adaptive-optics ocular aberration correction on visual acuity at different luminances and contrast polarities. *J. Vis.* **8**, 1–12 (2008).

18. Westheimer, G. Visual Acuity and Hyperacuity. *Invest. Ophthalmol.* **64**, 570–572 (1975).

19.   Westheimer, G. Optical superresolution and visual hyperacuity. *Prog. Retin. Eye Res.* **31**, 467–480 (2012).

20.   Nagano, T. Temporal sensitivity of the human visual system to sinusoidal gratings. *J. Opt. Soc. Am.* **70**, 711–716 (1980).

21.   Kaplan, E. & Benardete, E. The dynamics of primate retinal ganglion cells. *Prog. Brain Res.* **134**, 17–34 (2001).

22.   Ko, H. K., Poletti, M. & Rucci, M. Microsaccades precisely relocate gaze in a high visual acuity task. *Nat. Neurosci.* **13**, 1549–1554 (2010).

23.   Kuang, X., Poletti, M., Victor, J. D. & Rucci, M. Temporal encoding of spatial information during active visual fixation. *Curr. Biol.* **22**, 510–514 (2012).

24.   Krauskopf, J., Cornsweet, T. N. & Riggs, L. A. Analysis of eye movements during monocular and binocular fixation. *J. Opt. Soc. Am.* **50**, 572–578 (1960).

25.   Intoy, J. & Rucci, M. Finely tuned eye movements enhance visual acuity. *Nat. Commun.* **11**, 1–11 (2020).

26.   Burak, Y., Rokni, U., Meister, M. & Sompolinsky, H. Bayesian model of dynamic image stabilization in the visual system. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 19525–19530 (2010).

27.   Pitkow, X., Sompolinsky, H. & Meister, M. A neural computation for visual acuity in the presence of eye movements. *PLoS Biol.* **5**, 2898–2911 (2007).

28.   Rucci, M., Iovin, R., Poletti, M. & Santini, F. Miniature eye movements enhance fine spatial detail. *Nature* **447**, 851–854 (2007).

29.   Rolfs, M. Microsaccades: Small steps on a long way. *Vision Res.* **49**, 2415–2441 (2009).

30.   Hafed, Z. M., Chen, C. Y., Tian, X., Baumann, M. P. & Zhang, T. Active vision at the foveal scale in the primate superior colliculus. *J. Neurophysiol.* **125**, 1121–1138 (2021).

31.   Clark, A. M., Intoy, J., Rucci, M. & Poletti, M. Eye drift during fixation predicts visual acuity. *Proc. Natl. Acad. Sci. U. S. A.* **119**, 1–10 (2022).

32.   Herrmann, C. J. J., Metzler, R. & Engbert, R. A self-avoiding walk with neural delays as a model of fixational eye movements. *Sci. Rep.* **7**, 1–17 (2017).

33.   Ben-Shushan, N., Shaham, N., Joshua, M. & Burak, Y. Fixational drift is driven by diffusive dynamics in central neural circuitry. *Nat. Commun.* **13**, 1697 (2022).

34.   Ratnam, K., Domdei, N., Harmening, W. M. & Roorda, A. Benefits of retinal image motion at the limits of spatial vision. *J. Vis.* **17**, 1–11 (2017).

35.   Cherici, C., Kuang, X., Poletti, M. & Rucci, M. Precision of sustained fixation in trained and untrained observers. *J. Vis.* **12**, 1–16 (2012).

36.   Putnam, N. M. *et al.* The locus of fixation and the foveal cone mosaic. *J. Vis.* **5**, 632–639

(2005).

37. Wilk, M. A. *et al.* Assessing the spatial relationship between fixation and foveal specializations. *Vision Res.* **132**, 53–61 (2017).

38. Zhang, T. *et al.* Variability in human cone topography assessed by adaptive optics scanning laser ophthalmoscopy. *Am. J. Ophthalmol.* **160**, 290–300 (2015).

39. Wells-Gray, E. M., Choi, S. S., Bries, A. & Doble, N. Variation in rod and cone density from the fovea to the mid-periphery in healthy human retinas using adaptive optics scanning laser ophthalmoscopy. *Eye* **30**, 1135–1143 (2016).

40. Intoy, J., Mostofi, N. & Rucci, M. Fast and nonuniform dynamics of perisaccadic vision in the central fovea. *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

41. Bowers, N. R., Gautier, J., Lin, S. & Roorda, A. Fixational eye movements in passive versus active sustained fixation tasks. *J. Vis.* **21**, 1–16 (2021).

42. Rossi, E. a, Weiser, P., Tarrant, J. & Roorda, A. Visual performance in emmetropia and low myopia after correction of high-order aberrations. *J. Vis.* **7**, 14 (2007).

43. Reiniger, J. L. *et al.* Habitual higher order aberrations affect Landolt but not Vernier acuity. *J. Vis.* **19**, 1–15 (2019).

44. Artal, P. *et al.* Neural compensation for the eye's optical aberrations. *J. Vis.* **4**, 4 (2004).

45. Chen, L., Artal, P., Gutierrez, D. & Williams, D. R. Neural compensation for the best aberration correction. *J. Vis.* **7**, 9.1-9 (2007).

46. Williams, D. R. Aliasing in human foveal vision. *Vision Res.* **25**, 195–205 (1985).

47. Stockman, A. & Rider, A. T. Formulae for generating standard and individual human cone spectral sensitivities. *Color Res. Appl.* **48**, 818–840 (2023).

48. Ruderman, D. L. & Bialek, W. Seeing Beyond the Nyquist Limit. *Neural Comput.* **4**, 682–690 (1992).

49. Ditchburn, R. W. & Ginsborg, B. L. Involuntary eye movements during fixation. *J. Physiol.* **119**, 1–17 (1953).

50. Cornsweet, T. N. Determination of the stimuli for involuntary drifts and saccadic eye movements. *J. Opt. Soc. Am.* **46**, 987–993 (1956).

51. Rucci, M. & Poletti, M. Control and Functions of Fixational Eye Movements. *Annu. Rev. Vis. Sci.* **1**, 499–518 (2015).

52. Ahissar, E. & Arieli, A. Figuring space by time. *Neuron* **32**, 185–201 (2001).

53. Ahissar, E. & Arieli, A. Seeing via Miniature Eye Movements: A Dynamic Hypothesis for Vision. *Front. Comput. Neurosci.* **6**, 1–27 (2012).

54. Anderson, A. G., Ratnam, K., Roorda, A. & Olshausen, B. A. High-acuity vision from retinal

image motion. *J. Vis.* **20**, 1–19 (2020).

55. Zalevsky, Z. Exceeding the diffraction and the geometric limits of imaging systems: A review. in *Dolev, S., Oltean, M. (Eds.), Optical Supercomputing 2010. Lecture Notes in Computer Science* vol. 6748 119–130 (Springer-Verlag, 2011).

56. Westheimer, G. & McKee, S. P. Spatial configurations for visual hyperacuity. *Vision Res.* **17**, 941–7 (1977).

57. Rucci, M. & Victor, J. D. The unsteady eye: An information-processing stage, not a bug. *Trends in Neurosciences* vol. 38 195–206 (2015).

58. Malevich, T., Buonocore, A. & Hafed, Z. M. Rapid stimulus-driven modulation of slow ocular position drifts. *Elife* **9**, 1–21 (2020).

59. Gruber, L. Z. & Ahissar, E. Closed loop motor-sensory dynamics in human vision. *PLoS One* **15**, 1–18 (2020).

60. Lin, Y.-C., Intoy, J., Clark, A. M., Rucci, M. & Victor, J. D. Cognitive influences on fixational eye movements. *Curr. Biol.* **33**, 1606–1612 (2023).

61. Zhao, Z., Ahissar, E., Victor, J. D. & Rucci, M. Inferring visual space from ultra-fine extra-retinal knowledge of gaze position. *Nat. Commun.* **14**, 1–12 (2023).

62. Bergeron, A., Matsuo, S. & Guitton, D. Superior colliculus encodes distance to target, not saccade amplitude, in multi-step gaze shifts. *Nat. Neurosci.* **6**, 404–413 (2003).

63. Hafed, Z. M., Goffart, L. & Krauzlis, R. J. A neural mechanism for microsaccade generation in the primate superior colliculus. *Science (80-.).* **323**, 940–943 (2009).

64. Chen, C.-Y., Hoffmann, K.-P., Distler, C. & Hafed, Z. M. The Foveal Visual Representation of the Primate Superior Colliculus. *Curr. Biol.* **29**, 2109–2119 (2019).

65. Domdei, N., Reiniger, J. L., Holz, F. G. & Harmening, W. M. The Relationship Between Visual Sensitivity and Eccentricity, Cone Density and Outer Segment Length in the Human Foveola. *Invest. Ophthalmol. Vis. Sci.* **62**, 1–16 (2021).

66. Ehrenstein, W. H., Arnold-Schulz-Gahmen, B. E. & Jaschinski, W. Eye preference within the context of binocular functions. *Graefe's Arch. Clin. Exp. Ophthalmol.* **243**, 926–932 (2005).

67. Zhou, D. *et al.* Association of Visual Acuity with Ocular Dominance in 2045 Myopic Patients. *Curr. Eye Res.* **42**, 1155–1159 (2017).

68. Domdei, N. *et al.* Cone Density Is Correlated to Outer Segment Length and Retinal Thickness in the Human Foveola. *Invest. Ophthalmol. Vis. Sci.* **64**, 1–11 (2023).

69. Roorda, A. *et al.* Adaptive optics scanning laser ophthalmoscopy. *Opt. Express* **10**, 405–412 (2002).

70. Cunefare, D. *et al.* Open source software for automatic detection of cone photoreceptors in adaptive optics ophthalmoscopy using convolutional neural networks. *Sci. Rep.* **7**, 1–11

(2017).

71.   Wynne, N. *et al.* Intergrader agreement of foveal cone topography measured using adaptive
        optics scanning light ophthalmoscopy. *Biomed. Opt. Express* **13**, 4445–4454 (2022).

72.   Poonja, S., Patel, S., Henry, L. & Roorda, A. Dynamic visual stimulus presentation in an
        adaptive optics scanning laser ophthalmoscope. *J Refract Surg* **21**, 575–580 (2005).

73.   Watson, A. B. & Pelli, D. G. QUEST: a Bayesian adaptive psychometric method. *Percept.
        Psychophys.* **33**, 113–120 (1983).

74.   Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

75.   Pelli, D. G. The VideoToolbox software for visual psychophysics: transforming numbers into
        movies. *Spat. Vis.* **10**, 437–442 (1997).

76.   McAnany, J. J. The effect of exposure duration on visual acuity for letter optotypes and
        gratings. *Vision Res.* **105**, 86–91 (2014).

77.   Schütt, H. H., Harmeling, S., Macke, J. H. & Wichmann, F. A. Painfree and accurate Bayesian
        estimation of psychometric functions for (potentially) overdispersed data. *Vision Res.* **122**,
        105–123 (2016).

78.   Stevenson, S. B. & Roorda, A. Correcting for miniature eye movements in high-resolution
        scanning laser ophthalmoscopy. in *Ophthalmic Technologies XV* vol. 5688 145–151 (SPIE,
        2005).

79.   Hofmann, J., Domdei, L., Jainta, S. & Harmening, W. M. Assessment of binocular fixational
        eye movements including cyclotorsion with split-field binocular scanning laser
        ophthalmoscopy. *J. Vis.* **22**, 1–13 (2022).

Figures

Figure 1. Cone-resolved adaptive optics micro-psychophysics. a, Schematic of cell-resolved visual acuity testing in the human foveola with an adaptive optics scanning laser ophthalmoscope (AOSLO). Stimuli were dark Snellen-E optotypes presented at variable size and four orientations in the center of the 788 nm AOSLO imaging raster. Participants responded by indicating stimulus orientation during natural viewing, i.e. unrestricted eye motion. **b**, Exemplary single trial retinal motion trace and strip-wise image stabilization of a single AOSLO frame (shown here during a microsaccade for better visibility). Trials containing microsaccades or blinks during the 500 ms stimulus presentation (gray shaded area) were excluded. The x-axis grid represents individual video frames (33 ms). **c**, Foveolar retinal cone mosaic with exemplary single trial retinal motion across the stimulus. Time is represented by color from stimulus onset to offset (purple to yellow). The cone density centroid (CDC) is shown as a red circle with white fill. **d**, Typical psychophysical data of 5 consecutive runs in one eye. Each run followed a QUEST procedure with 20 trials. **e**, Psychometric function fit to the data (about 100 trials). Acuity thresholds were estimated at 62.5 % correct responses. **f**, Exemplary retinal images (upper rows) and corresponding cone activation patterns (lower rows) of one experimental run (20 trials from top left to bottom right). Cone activation patterns are shown for a representative single frame. See Supplementary Movie 1 and 2 for a real-time video representation.

Figure 2. Visual acuity depends on foveolar sampling capacity. a, Foveolar cone mosaics of the two eyes with highest and lowest cone densities, overlayed with the physical stimulus at an average threshold size (24 arcsec). **b**, Nyquist limit: critical details equaling or larger than the spacing of cones are resolvable. **c**, Visual acuity thresholds measured with 788 or 840 nm infrared light, normalized to the eyes' Nyquist limits. **d**, Correlation between participants individual visual acuity thresholds and cone density. Thresholds exceeded the Nyquist sampling limit and were significantly lower in eyes with higher cone densities. Dominant eyes are shown as filled, non-dominant eyes as open markers. The gray horizontal and vertical bars at each point represent standard deviations of sampling cone density and the 95 % confidence intervals for acuity thresholds. The theoretical Nyquist limit is represented by a dashed green line. **e**, Correlation between dominant and non-dominant eyes in visual acuity (top) and cone density (bottom). Dominant eyes reached, on average, 1.5 arcmin lower thresholds than non-dominant eyes, whereas cone density (at the retinal locations that sampled the stimulus) was very similar between fellow eyes.

Figure 3. Fixational drift and the contribution to visual acuity. a, Ocular drift during stimulus presentation (participant 16, left eye). Single AOSLO frame captured during Snellen E presentation (top left) and all single stimulus positions (colored dots) of 5 experimental runs shown on the corresponding cone mosaic (panel 2-6). White iso-lines delimit cone density percentile areas (90th to 50th percentile visible). Time is represented by color from stimulus onset to offset (purple to yellow). Individual drift trajectories for all eyes are shown in figure 3 supplement 1. **b**, Individual motion traces highlighting intra- and inter-subject drift variability. Traces are from one run in the participant with the lowest (upper rows) and highest (lower rows) average drift lengths. **c**, Computation of drift length as a sum of interframe motion vectors (top) and the relative frequency of occurrences among all participants and trials (bottom). **d**, Median drift length and drift length range showed a moderate correlation in dominant as well as non-dominant eyes (top). The minimum drift length was similar between participants (3.8 ± 0.8 arcmin) whereas the maximum length varied about three times as much (12.0 ± 2.7 arcmin). **e**, Drift lengths in fellow eyes had a very strong correlation. **f**, Cone density and drift length did not show a significant correlation in dominant or non-dominant eyes. **g**, The median drift length had a moderate correlation with visual acuity threshold in dominant as well as non-dominant eyes. Dominant eyes are indicated by filled, non-dominant eyes by open markers.

Figure 4. Drift moves stimuli to higher cone density areas. a, Five exemplary motion traces relative to CDC, PRL and PCD location on the Voronoi tessellated cone mosaic of one participant. **b**, All single trial motion traces of one eye shown on the corresponding cone mosaic (95 trials containing drift only). One-SD isoline areas (ISOA) are shown for all stimulus onset (blue) and offset (yellow) locations, indicating a trend of directional drift towards higher cone densities during 500 ms stimulus presentation. **c**, Polar histogram of all individual motion traces (n = 2739) shows the relative frequency of motion angles, θ_{Retina}, between start (coordinate center) and end of motion in retinal coordinates. The inset indicates θ sign. **d**, Same data as in c, where θ_{CDC} was computed relative to the line connecting drift start location and CDC, see inset. The pink quarter indicates the angular space used for the computation of the tuning ratio. For more details on the drift directionality of individual eyes, see figure supplement 1. **e**, The difference between acuity threshold and Nyquist limit showed a significant trend to be larger for stronger directionality tuning. The tuning ratio was computed as the ratio between the relative frequency of intra-participant drift motion towards the CDC (± 45 deg) and the average of drift motion towards the remaining 3 quadrants. **f**, Relative frequency of drift direction relative to CDC (top), PRL (middle) and PCD (bottom), respectively. For more details on the temporal progression of drift directionality, see figure supplement 2. **g**, Across all participants and trials, drift length correlated with stimulus onset distance from CDC. There was no clear effect of stimulus onset distance on motion directionality (data color corresponding to θ_{CDC}). **h**, The achieved sampling gain due to the performed drift motion is significantly correlated to the potential sampling gain in individuals. In both dominant and non-dominant eyes the potential sampling gain is on average exploited by 30 %, respectively. Due to shifting the CDC towards the stimulus, participants had different PRLs for a sustained fixation task and the visual resolution task (see figure supplement 3).

Supplementary figures

Figure 3 – figure supplement 1. Drift trajectories on foveolar mosaics. Participants (P01-16) are sorted according to acuity thresholds in their dominant eye (best to worst). All single trial motion traces are shown on the corresponding cone mosaic, color coded by elapsed time. One-SD isoline areas (ISOA) are shown for all stimulus onset (blue) and offset (yellow) locations. The medians of all onset and offset locations are shown as blue and yellow squares, respectively. The CDC is indicated by a red circle with white fill.

Figure 4 – figure supplement 1. Individual drift directionality. Sorted according to their dominant eyes CDC to PRL distance (largest to smallest) eyes drift directionality is shown relative to the CDC. The CDC is indicated by a red circle with white fill. Participant numbers (P01 to P16) are assigned based on the dominant eyes visual acuity threshold (best to worst). The upper left plot shows the angular labels exemplarily for all eyes. The tuning ratio was computed as the ratio between the relative frequency of intra-participant drift motion towards the CDC (± 45 deg), indicated by the pink quarter, and the average of drift motion towards the remaining 3 quadrants.

Figure 4 – figure supplement 2. Time course of drift directionality. The angles θCDC, θPRL and θPCD for all eyes were computed over time, relative to the stimulus onset and are shown after 33 ms (1 frame), 165 ms (5 frames), 330 ms (10 frames) and 462 ms (14 frames). The respective tuning ratio (TR) is shown in the bottom part of each subplot. The tuning ratio was computed as the ratio between the relative frequency of inter-participant drift motion towards the CDC (± 45 deg) and the average of drift motion towards the remaining 3 quadrants. For all cases, the circular data are not distributed uniformly around the circle (Rayleigh test, p < 0.001, with increasing significance for longer durations). The distributions were compared against each other by the Kolmogorov-Smirnov test. Only CDC vs. PRL (33 ms) and PRL vs. PCD (33 ms) show no statistically significant difference. The

distribution towards CDC compared to PCD is significantly different for 33 ms (p = 0.002). For all other timepoints, all distributions are significantly different from each other, with increasing significance for longer durations (for 462 ms, all p < 0.001).

Figure 4 – figure supplement 3. Different retinal locations used in a fixation or resolution task. Median PRL locations of all participants' right and left eyes plotted in retinal coordinates relative to the CDC. Retinal locations are plotted in teal for the fixation task and in black for the resolution task. A black line connects retinal locations of the same participant.

a AOSLO imaging and microstimulation | Fixational eye motion | Stimulus on retina

Retina
Foveola
5 arcmin

b

Stim.

Retinal position (arcmin)

— x — y

Time (ms)

Raw video frame | Stabilized video frame

c Stimulated cones during single trial

2 arcmin

Time (ms)
0   500

d

Gap size (arcsec)

○ Correct   ● Incorrect

Trial

e

Proportion correct (%)

Threshold estimate
(95 % CI)

N = 100

Gap size (arcsec)

f Experimental run (20 trials)

Physical stimulus

Single frame cone activation

Stimulus at threshold

5 arcmin

a

Max. sampling (16104 c/deg²)

2 arcmin

Min. sampling (10297 c/deg²)

2 arcmin

b

Signal   0   1

c

Normalized threshold

N   16   16   12   12
788 nm   840 nm

d

Nyquist limit

Visual acuity threshold (arcsec)

Acuity threshold (logMAR)

Cone density (cones/deg²)

■ Dominant (r² = 0.45, p = 0.005)
□ Non-dominant (r² = 0.28, p = 0.036)

e

Visual acuity (arcsec)

Non-dominant eye

r² = 0.80
p < 0.001

Dominant eye

Cone density (cones/deg²)

Non-dominant eye

*10³

r² = 0.85
p < 0.001

Dominant eye

*10³

a

Single frame — 5 arcmin

Run 1 — Time (ms) 0 – 500

Run 2

Run 3

Run 4

Run 5

b **Example motion paths**

Lowest average drift length (P02)

Highest average drift length (P14)

5 arcmin

c Drift length computation

$$\sum_{i=0}^{n} \sqrt{\Delta x_i^2 + \Delta y_i^2}$$

Δx
Δy

Time (frames)

N = 2739

Relative frequency / Drift length (arcmin)

d

Drift length range (arcmin)

Dom. ($r^2 = 0.55$, $p = 0.001$)
Non-dom. ($r^2 = 0.34$, $p = 0.02$)

Relative frequency / Drift length (arcmin)

Min / Max

e

Drift non-dominant eye (arcmin)

$r^2 = 0.86$
$p < 0.001$

Drift dominant eye (arcmin)

f

Drift length (arcmin)

Dom. ($r^2 = 0.07$, $p = 0.3$)
Non-dom. ($r^2 = 0.06$, $p = 0.4$)

Cone density (cones/deg²) *10³

g

Visual acuity threshold (arcsec)

Dominant ($r^2 = 0.25$, $p = 0.04$)
Non-dominant ($r^2 = 0.29$, $p = 0.03$)

Drift length (arcmin)

P01_R P01_L P09_R P09_L

Time (ms)
0   500

ISOA Onsets
ISOA Offsets

P02_R P02_L P10_R P10_L

P03_R P03_L P11_R P11_L

P04_R P04_L P12_R P12_L

P05_R P05_L P13_R P13_L

P06_R P06_L P14_R P14_L

P07_R P07_L P15_R P15_L

P08_R P08_L P16_R P16_L

10 arcmin

a

Time (ms) 0 — 500 | 2.5 arcmin | PCD | PRL | CDC

b

Onset | Offset | 10 arcmin

c Drift direction in retinal coordinates (θ_{Retina})

6 % | 4 % | 2 % | 45° | 0° | 315° | 225°

n = 2739 trials (32 eyes)

d Drift direction relative to CDC (θ_{CDC})

6 % | 4 % | 2 % | 45° | 0° ≐ | 315° | 225°

e Directionality tuning towards CDC

Nyquist - Acuity difference (arcsec)

Tuning ratio towards CDC ± 45°

Dom ($\rho^2 = 0.45$, p = 0.01) ■
Non-dom ($\rho^2 = 0.27$, p = 0.04) □

f Tuning comparison

Relative frequency (%) | Absolute frequency

CDC — Tuning ratio: 3.0
PRL — Tuning ratio: 2.3
PCD — Tuning ratio: 1.7

θ (°)
150 100 50 0 -50 -100 -150

g Drift length vs. distance to CDC

Drift length (arcmin)

Δ stimulus onset location to CDC (arcmin)

θ_{CDC} 0° — 180°

$\rho^2 = 0.06$
p < 0.001

h Exploited sampling gain

Sampling gain (%)

Potential sampling gain (%)

Dom ($r^2 = 0.35$, p = 0.02)
y = 0.30x + 0.76
Non-dom ($r^2 = 0.50$, p = 0.002)
y = 0.29x + 0.74

Right eyes

Left eyes

Offset from CDC (arcmin)

Offset from CDC (arcmin)

○ CDC   ● PRL (fixation)   ● PRL (resolution)