Title

Sub-cone visual resolution by active, adaptive sampling in the human foveola

Authors

Jenny L. Witten, Veronika Lukyanova, Wolf M. Harmening*

Affiliations

Rheinische Friedrich-Wilhelms-Universität Bonn, Department of Ophthalmology, Bonn,
Germany

*Corresponding author: Jenny Lorén Witten, jennylwitten@gmail.com

Abstract

The foveated architecture of the human retina and the eye's mobility enable prime spatial
vision, yet the interplay between photoreceptor cell topography and the constant motion of
the eye during fixation remains unexplored. With *in vivo* foveal cone-resolved imaging and
simultaneous microscopic photo stimulation, we examined visual acuity in both eyes of 16
participants while precisely recording the stimulus path on the retina. We find that resolution
thresholds were correlated with the individual retina's sampling capacity, and exceeded what
static sampling limits would predict by 18 %, on average. The amplitude and direction of
fixational drift motion, previously thought to be primarily random, played a key role in
achieving this sub-cone diameter resolution. The oculomotor system finely adjusts drift
behavior towards retinal areas with higher cone densities within only a few hundred
milliseconds to enhance retinal sampling.

Introduction

Assessing visual abilities was already important in historic times[1], and the precise
measurement of visual acuity, our ability to resolve fine spatial detail by eye, has great
importance for many real life scenarios and is up to this day the primary diagnostic tool to
determine visual function in a clinical and optometric setting. Quite surprisingly, the widely-
believed assumption that the packing density and arrangement of retinal photoreceptors at
the foveal center set the limit to this ability has never been experimentally confirmed.

Foveolization, the morphological and functional specialization of the cellular architecture of the light sensitive retina optimizes the human eye for high-acuity daytime vision[2,3]. Within the central one-degree diameter of the fovea, termed foveola, postreceptoral neurons are displaced centrifugally and the area is free of potentially shadowing blood vessels and glia cells[4,5]. The outer segments of foveolar cone photoreceptors are maximally thinned and densely packed for peak spatial sampling[6–8], which at the same time makes these cells the most difficult to study *ex vivo*[9] as well as *in vivo*[10]. Each foveolar cone synapses to one ON- and one OFF-midget bipolar cell, which in turn synapse exclusively upon single ON- and OFF-midget ganglion cells, a circuit that is adult like before birth[11]. This establishes an undisturbed *private line* from individual foveal receptors to central processing stages.

Based on indirect comparisons between histological and psychophysical data, the hypothesis that cone spacing imposes the fundamental limit for visual resolution has been put forward[8,12]. It is well established that cone spacing, especially in the central fovea, is highly variable between individuals[12–15], making general comparisons between acuity measurements and foveolar density estimated from histological samples susceptible to error. One of the main reasons why the hypothesis lacks direct experimental proof is that because, under natural viewing conditions, both visual resolution and experimental access to foveal photoreceptors is blurred by the imperfect optics of the human eye[16,17]. Here, we have overcome the optical barrier of the human eye by employing adaptive optics cell-resolved *in vivo* retinal imaging in conjunction with micro-psychophysics to study directly whether the individual's mosaic of foveolar cones determines visual performance in a high-acuity resolution task. Our findings may also resolve another, so far only indirectly tested, hypothesis if a potential visual resolution advantage arises in myopic eyes. Myopes, despite retinal stretching, generally have a higher angular sampling density in and around the fovea compared to emmetropes[13].

While acuity is assumed to be mainly limited by the resolving capacity of the eye's optics and retinal mosaic, it is well established that, for different visual tasks, performance thresholds can be substantially lower than the sampling grain of photoreceptors. This phenomenon has been termed hyperacuity[18] and depends on the neural visual system's ability to extract subtle differences within the spatial patterns of the optical image on the retina[19]. Thus, the visual system already incorporates mechanisms to detect relative spatial offsets an order of magnitude smaller than the spatial granularity of the retina. To make use of those fine distinctions in a resolution task, the neuronal system needs to go beyond purely spatial coding of incoming signals.

Unlike a camera, the visual system depends on temporal transients arising in the receptor's cellular signals. Neurons in the retina, thalamus and later stages of the visual pathways

respond strongly to temporal changes[20,21]. Thusly, the fovealized retinal architecture in
humans is accompanied by a dynamic sampling behavior that, by quick and precise
movements of the eye, brings retinal images of objects of interest to land in the foveola[22,23].
Even during steady fixation, for example of a distant face or a single letter of this text,
incessant fixational eye movements slide tens to hundreds of foveolar photoreceptors across
the retinal image, thereby introducing temporal modulations that translate spatial activation
patterns into the temporal domain[24]. Small and rapid gaze shifts known as microsaccades
relocate gaze within the foveola during periods of fixation[22], and between microsaccades,
the eyes perform a more continuous, seemingly random motion termed fixational drift[25,26].
Computational work suggested that fixational eye motion would introduce noise and thus
impair visual acuity[27,28]. Contrarily, recent studies on human psychophysics demonstrated
fixational eye motion to be beneficial for fine spatial vision[26,29,30]. Especially drift motion has
been increasingly argued to not just be randomly refreshing neural activity, but rather
structuring it[26,31,32] and being under central control[33].

The incessant motion of the eye conveys fine spatiotemporal detail that requires deciphering
of continuously changing photoreceptor signals, which are linked by the geometry of the
photoreceptor array and by how the eye moves. For instance, luminance modulation in
individual cones will scale with drift amplitude. Larger luminance variations on single
receptors also yields more neuronal activity within the range of temporal frequencies
parvocellular ganglion cells are sensitive to. Selective spatial frequencies can thus be
amplified by varying drift amplitude[26]. While the neuronal mechanisms that generate
fixational drift are still not fully understood[34], its consequence to visual perception has been
demonstrated. Drift was shown to improve visual performance in resolution tasks[26,29,35], and
a recent model of early retinal signals suggests that if drift amplitude is tuned to object size,
visual acuity would be enhanced[36]. Indeed, considerable differences in ocular drift between
individuals exist[32,37], and subjects exhibiting less drift were shown to have better acuity[32]. If
such differences are a consequence of an active, adaptive mechanism, however, and how
drift behavior is related to the photoreceptors that sample the retinal image is unknown.

The direct experimental access to the foveolar center, when other limiting factors like image
blur or retinal motion are taken out of the equation or can be precisely measured, will allow
to confirm or reject the long-standing hypothesis about the individual limits of vision. This will
help to understand the fundamental physiological limitations of the visual system and will
have important implications for clinical studies of retinal health.

Results

Resolution is finer than single cone sampling limits

We investigated the limitations of the photoreceptor packing density on individual visual resolution acuity by overcoming the optical aberrations of the eye with adaptive optics scanning laser ophthalmoscopy (AOSLO), while simultaneously performing psychophysical measurements and recording the fixational retinal motion (Fig. 1a, b and c). In a four-alternative forced-choice task, 16 healthy participants indicated the orientation of an E-optotype while inspecting the stimulus with their individually preferred fraction of foveolar photoreceptors. These cone photoreceptors were simultaneously imaged and it was later identified which cells contributed to resolving the stimulus (Fig. 1c, f). A psychometric fit to the data expressed as percentage correct from 100 trials was used to compute visual acuity thresholds (see online Methods and Fig. 1d,e). In this near diffraction-limited testing condition, participants reached visual acuity thresholds between 20.6 and 28.5 arcsec (mean ± SD: 24.1 ± 2.4 arcsec), which compares to 20/8 vision (logMAR = -0.4). All participants reached thresholds better than 20/10 vision (logMAR = -0.3), the last line of a typical clinical Snellen chart or projectors of acuity optotypes that are used in clinical as well as optometric daily routine.

Fig. 1 | Cone-resolved adaptive optics micro-psychophysics. a, Schematic of cell-resolved visual acuity testing in the human foveola with an adaptive optics scanning laser ophthalmoscope (AOSLO). Stimuli were dark Snellen-E optotypes presented at variable size and four orientations in the center of the 788 nm AOSLO imaging raster. Participants responded by indicating stimulus orientation during natural viewing, i.e. unrestricted eye motion. **b**, Exemplary single trial retinal motion trace and strip-wise image stabilization of a single AOSLO frame (shown here during a microsaccade for better visibility). Trials containing microsaccades or blinks during the 500 ms stimulus presentation (gray shaded area) were excluded. The x-axis grid represents individual video frames (33 ms). **c**, Foveolar retinal cone mosaic with exemplary single trial retinal motion across the stimulus. Time is represented by color from stimulus onset to offset (purple to yellow). The cone density centroid (CDC) is shown as a red circle with white fill. **d**, Typical psychophysical data of 5 consecutive runs in one eye. Each run followed a QUEST procedure with 20 trials. **e**, Psychometric function fit to the data (about 100 trials). Acuity thresholds were estimated at 62.5 % correct responses. **f**, Exemplary retinal images (upper rows) and corresponding cone activation patterns (lower rows) of one experimental run (20 trials from top left to bottom right). Cone activation patterns are shown for a representative single frame. See Supplementary Movie 1 and 2 for a real-time video representation.

Cone densities at the CDC ranged between 10,692 and 16,997 cones/deg^2, with an average density of 13,640 cones/deg^2 (PCD mean: 13,944 cones/deg^2, range: 10,823 to 17,309 cones/deg^2), comparable to previous reports[13,14,38–41]. The median sampling cone density ranged between 10,297 and 16,104 cones/deg^2 (mean: 13,149 cones/deg^2). Two experimental runs of the eyes with highest and lowest sampling density are exemplarily shown in Supplementary Movie 1 and 2. The two foveolar cone mosaic images were also visualized and overlayed with a Snellen E stimulus at average threshold size (Fig. 2a). The theoretical prediction, given by the Nyquist sampling limit, would assume the high-density retina where each single cone diameter is smaller than the Snellen E's gap or bar is able to resolve the stimulus, whereas the low-density retina fails in identifying the correct orientation (schematic representation in Fig. 2b). However, for our 788 nm testing condition, all participants reached individual resolution thresholds below their Nyquist limit predicted by the spacing between rows of cones (Fig. 2c, d). On average, visual acuity thresholds exceeded the theoretical prediction by 20 % and 16 % in dominant and non-dominant eyes, respectively. When participants performed the same resolution task with a longer infrared wavelength (840 nm) imaging background, the absolute thresholds were slightly higher and thus closer to the Nyquist limit. Visual acuity thresholds were on average 7 % below and 2 % above the Nyquist limit for dominant and non-dominant eyes, respectively. These absolute visual acuity thresholds were the only case where noteworthy differences arose between the 788 nm and 840 nm experimental condition. For all other analyses, we found qualitatively similar results for either wavelength and therefore only report the 788 nm results throughout the manuscript.

Fig. 2 | Visual acuity depends on foveolar sampling capacity. a, Foveolar cone mosaics of the two
eyes with highest and lowest cone densities, overlayed with the physical stimulus at an average
threshold size (24 arcsec). **b**, Nyquist limit: critical details equaling or larger than the spacing of cones
are resolvable. **c**, Visual acuity thresholds measured with 788 or 840 nm infrared light, normalized to
the eyes' Nyquist limits. **d**, Correlation between participants individual visual acuity thresholds and
cone density. Thresholds exceeded the Nyquist sampling limit and were significantly lower in eyes
with higher cone densities. Dominant eyes are shown as filled, non-dominant eyes as open markers.
The gray horizontal and vertical bars at each point represent standard deviations of sampling cone
density and the 95 % confidence intervals for acuity thresholds. The theoretical Nyquist limit is
represented by a dashed green line. **e**, Correlation between dominant and non-dominant eyes in
visual acuity (top) and cone density (bottom). Dominant eyes reached, on average, 1.5 arcmin lower
thresholds than non-dominant eyes, whereas cone density (at the retinal locations that sampled the
stimulus) was very similar between fellow eyes.

For the first time, we could measure the direct relation between the individual foveolar cone
photoreceptor sampling density and participants visual resolution thresholds. We found the
diffraction limited visual acuity thresholds to be strongly correlated to the foveolar sampling
density in dominant as well as fellow eyes (Fig. 2d). The higher the cone density, the smaller
the visual stimulus that could be resolved. The degree of correlation slightly differed for
dominant ($r^2 = 0.45$, $p = 0.005$) and non-dominant eyes ($r^2 = 0.28$, $p = 0.036$), suggesting
that up to 45 % of the variance in inter-subject visual acuity can be explained by the
individual cone sampling densities. Overall, participants reached significantly lower
thresholds with their dominant eyes (average: 1.5 arcsec, SD ± 1.1; paired t-test, $p < 0.001$).
Nevertheless, visual acuity thresholds were strongly correlated between dominant and non-
dominant eyes ($r^2 = 0.80$, $p < 0.001$, Fig. 2e, see supplementary discussion). To test whether
the effect of different absolute thresholds might be explained by underlying differences in the
sampling cone density, fellow eyes densities were compared to each other. Sampling

densities had a very strong correlation between fellow eyes ($r^2 = 0.85$, $p < 0.001$, Fig. 2e),

but did not differ between right and left eyes ($p = 0.38$) nor when grouping them according to

ocular dominance ($p = 0.88$). This compares well to previous studies that also showed

strong correlations between fellow eyes regarding both anatomical[14] as well as functional[15]

characteristics. Dominant eyes had a median of 78 cones/deg^2 higher densities compared to

their fellow eyes. To account for the 1.5 arcsec difference in acuity thresholds, a much

higher density difference of about 1,500 cones/deg^2 would have been needed. Next to the

spatial cell arrangement, that only partially predicted the achievable resolution acuity, ocular

motion and its associated temporal modulations also highly influence visual resolution.

Ocular drift is an active sampling mechanism

As the eye drifts, a visual stimulus projected onto the retina is processed as a spatiotemporal

luminance flow. The stimulus itself as well as the extent of drift motion determine the

characteristics of modulation. In our experiments, analyzing the exact retinal locations

sampling the stimulus, we revealed that participants kept coming back to the same few

hundreds of cone photoreceptors (Fig. 3a). To focus on the characteristics and implications

of drift eye motion, trials containing microsaccades during stimulus presentation were

excluded from the analyses. During the short stimulus duration however, microsaccades

rarely occurred, as participants tend to suppress their microsaccades, likely because they

can be detrimental to fine-scale discrimination[42,43]. Drift motion patterns varied greatly

across, but also within participants. Examples of drift motion paths for the eyes that

performed the smallest and largest drift motion, on average, show a great variability in

shapes as well as extent of motion (Fig. 3b). In our analyses, we chose the sum of piecewise

drift amplitude as the prime metric to describe the ocular drift motion, because the

randomness underlying alternative metrics of drift eye movements becomes increasingly

questionable (see also Discussion). Across all participants and experimental trials, drift

amplitudes ranged between 2.5 and 17.2 arcmin, with a median amplitude of 6.5 arcmin

(which corresponds to a velocity of 5 to 34.5 arcmin/sec, median: 13 arcmin/sec, Fig. 3c).

The drift amplitudes are slightly smaller than in previous non-AO studies, which is

attributable to the viewing situation. The participants were looking at a very small imaging

field within a completely dark periphery without distracting structures or stimuli. The smallest

drift movement performed was similar among eyes (range: 2.5 – 5.4 arcmin), whereas the

largest individual drifts differed more than three times as much (range: 7.7 – 17.2 arcmin).

Therefore, the individual drift span was rather driven by the larger drift amplitudes of an eye

and there was a strong correlation between median drift amplitude and drift range (dominant

eyes: $r^2 = 0.55$, $p = 0.002$, non-dominant eyes: $r^2 = 0.34$, $p = 0.02$, Fig. 3d).

Fig. 3 | Fixational drift and the contribution to visual acuity. a, Ocular drift during stimulus
presentation (participant 16, left eye). Single AOSLO frame captured during Snellen E presentation
(top left) and all single stimulus positions (colored dots) of 5 experimental runs shown on the
corresponding cone mosaic (panel 2-6). White iso-lines delimit cone density percentile areas (90th to
50th percentile visible). Time is represented by color from stimulus onset to offset (purple to yellow). **b**,
Individual motion traces highlighting intra- and inter-subject drift variability. Traces are from one run in
the participant with the lowest (upper rows) and highest (lower rows) average drift amplitudes. **c**,
Computation of drift amplitude as a sum of interframe motion vectors (top) and the relative frequency
of occurrences among all participants and trials (bottom). **d**, Median drift amplitude and drift amplitude
range showed a moderate correlation in dominant as well as non-dominant eyes (top). The minimum
drift amplitude was similar between participants (3.8 ± 0.8 arcmin) whereas the maximum amplitude
varied about three times as much (12.0 ± 2.7 arcmin). **e**, Drift amplitudes in fellow eyes had a very
strong correlation. **f**, Cone density and drift amplitude did not show a significant correlation in
dominant or non-dominant eyes. **g**, The median drift amplitude had a moderate correlation with visual
acuity threshold in dominant as well as non-dominant eyes. Dominant eyes are indicated by filled,
non-dominant eyes by open markers.

In fellow eyes, which were measured consecutively, drift amplitudes had a very strong
correlation (r^2 = 0.86, p < 0.001, Fig. 3e) with no significant difference between eyes (paired
t-test, p = 0.2). The median drift amplitudes of all eyes varied between 4.8 and 8.5 arcmin
(mean ± SD: 6.6 ± 1.1 arcmin). Individual visual acuity thresholds were significantly
correlated with drift amplitudes (dominant: r^2 = 0.25, p = 0.04; non-dominant: r^2 = 0.29, p =

0.03, Fig. 3g), with a trend towards better visual acuity for small ocular drift motion. On a
photoreceptor resolved scale, this confirms recent findings which showed individual acuity
thresholds to be correlated with the drift motion during a non-AO acuity task, closely related
to the drift measured in a sustained fixation task [32].

Considering the previously shown correlation between visual acuity and sampling cone
density, one could assume those two aspects to go along with an increase of ocular drift for
lower cone densities, whereas higher densities potentially need less drift to translate the
stimulus over the same number of cones. However, we don't find the eye motion to be tuned
in a way to always let the stimulus slip across a similar number of cones. There was no
significant correlation between cone densities and drift amplitude (dominant: $r^2 = 0.07$, p =
0.3; non-dominant: $r^2 = 0.06$, p = 0.4). Other aspects are likely to also influence the trial-wise
motion, e.g. drift motion might be particularly tuned for stimulus sizes close to the threshold
and less crucial for larger gap sizes as suggested in a recent model of early visual signal
processing[36].

Drift is adaptive and directed

Ocular drift has long been assumed to be a persistent jittery motion that follows random
trajectories. Recent work showed that the amount of drift can vary and may be adapted to
the task that has to be performed[26,32]. We here investigated if beyond this, humans are able
to actively tune their ocular drift direction to exploit their prime spatial retinal processing
properties. We therefore registered the individual drift motion trajectories with the
photoreceptor mosaic, tracked them from the retinal location where the stimulus turned on
(onset) to where it turned off after 500 ms (offset), and related these trajectories to foveolar
landmarks (Fig. 4a, b). Because of the individual retinal locations used for fixation before
stimulus onset, we registered that, across all eyes, drift motion occurred towards all
directions during stimulus inspection, and no general trend in drift eye movements towards a
particular cardinal direction across participants occurred (Fig. 4c). Individual eyes, however,
showed different drift behavior mostly directed towards one or two of the four quadrants. All
four cardinal directions were represented. Participant P8$_{right}$, for example, drifted towards the
nasal or superior fovea in 90 % of all trials. P14$_{right}$, on the other hand, drifted towards the
temporal fovea in 75 % of all trials. When the frame of reference was rotated in each trial to
register the motion from the onset location relative to the CDC, we found a clear directional
bias in which the drift was likely to move the stimulus closer to the CDC. The drift
directionality was evaluated by measuring the relative angle between drift onset to drift offset
and drift onset to CDC. We observed a strong trend of drift directionality; 49 % of all drift

episodes moved the stimulus towards the CDC ± 45° (Fig. 4d). Among eyes, the individual

fractions ranged between 16 and 80 % of trials. Only two eyes drifted towards the CDC less

frequently than given by chance (Fig. S2). We computed the directionality tuning as the ratio

of relative drift towards the CDC ± 45° (purple quadrant in Fig. 4d) and the mean relative drift

towards the 3 other quadrants. A ratio of 1 indicated the same relative frequency of drift

towards all cardinal directions, whereas for a tuning ratio of 2 the retina moved the CDC

towards the stimulus twice as often compared to each of the other 3 cardinal directions. The

directionality tuning ratios ranged between 0.6 and 11.8 with a median value of 3.

Directionality tuning ratios had a significant effect on how much the resolution threshold

exceeded the Nyquist limit. Participants with highly tuned drift reached larger differences

between the Nyquist limit and their visual acuity threshold (dominant eyes: $r^2 = 0.45$, p =

0.01; non-dominant eyes: $r^2 = 0.27$, p = 0.04, Fig. 4e). Drift directionality was mostly similar

between eyes, and if intra-ocular differences occurred, they were not related to ocular

dominance. Also, we did not observe an effect of training on drift directionality: one of the

two trained observers had a very strong drift directionality (7 and 11.8 in the dominant and

non-dominant eye, respectively) while the other one exhibited a tuning ratio below average

(2.1 and 2.3 in the dominant and non-dominant eye, respectively).

Fig. 4 | Drift moves stimuli to higher cone density areas. a, Five exemplary motion traces relative

to CDC, PRL and PCD location on the Voronoi tessellated cone mosaic of one participant. **b**, All

single trial motion traces of one eye shown on the corresponding cone mosaic (95 trials containing

drift only). One-SD isoline areas (ISOA) are shown for all stimulus onset (blue) and offset (yellow)

locations, indicating a trend of directional drift towards higher cone densities during 500 ms stimulus

presentation. **c**, Polar histogram of all individual motion traces (n = 2739) shows the relative frequency
of motion angles, θ_{Retina}, between start (coordinate center) and end of motion in retinal coordinates.
The inset indicates θ sign. **d**, Same data as in c, where θ_{CDC} was computed relative to the line
connecting drift start location and CDC, see inset. The pink quarter indicates the angular space used
for the computation of the tuning ratio. **e**, The difference between acuity threshold and Nyquist limit
showed a significant trend to be larger for stronger directionality tuning. The tuning ratio was
computed as the ratio between the relative frequency of intra-participant drift motion towards the CDC
(± 45 deg) and the average of drift motion towards the remaining 3 quadrants. **f**, Relative frequency of
drift direction relative to CDC (top), PRL (middle) and PCD (bottom), respectively. **g**, Across all
participants and trials, drift amplitude correlated with stimulus onset distance from CDC. There was no
clear effect of stimulus onset distance on motion directionality (data color corresponding to θ_{CDC}). **h**,
The achieved sampling gain due to the performed drift motion is significantly correlated to the
potential sampling gain in individuals. In both dominant and non-dominant eyes the potential sampling
gain is on average exploited by 30 %, respectively.

Next to the CDC, two other foveolar landmarks are often reported as anchor locations
describing the center of the fovea. When relating the drift trajectories to the preferred retinal
locus of fixation (PRL) or the location of peak cone density (PCD), we found a weaker
approximation towards both. The retinae moved the stimulus towards the PRL or PCD
location in 42 % or 35 % of all trials, respectively (Fig. 4f). Therefore, the observed
directionality was strongest towards the CDC. In a considerable number of trials, the
stimulus onset was further displaced from all of the 3 retinal locations and therefore a
directed drift motion resulted in approximation towards CDC as well as PRL and PCD. Also,
in some eyes 2 or all of these retinal locations lay very close together, which results in very
similar effects. Nevertheless, in some eyes with particularly stable fixation that had at least a
few arcmin distance between their PRL and CDC we repeatedly observed a stimulus onset
close to PRL followed by a directional drift towards CDC with a resulting stimulus offset
closer to the CDC (see also supplementary discussion). Across participants this also
resulted in a significant reduction of the isoline contour area (ISOA) size between stimulus
onset and offset (p = 0.02, Fig. 4b, Fig. S1 and Supplementary Movie 3). The median ISOA
for stimulus onset locations was 92.5 arcmin^2 which was reduced to 68.2 arcmin^2 for
stimulus offset locations. This decrease in size of the area of all retinal landing points
supports the view of a certain retinal cone or very small area of a few arcmin^2 to be the
target region of the drift eye motion in a resolution task.

When we looked at how much the individual drift trajectory decreased the distance from
either location, the median distance convergence (onset/offset distance) towards CDC, PRL
and PCD was about 12 %, 7 % and 3 %, respectively. While no participant had an average

convergence of more than 30 % towards PRL or PCD, the maximum convergence ratio

towards CDC was about 50 %. An adaptive drift behavior was also found in the relative drift

amplitudes exhibited in each stimulus presentation. Although the individual drift amplitudes

could vary substantially from trial to trial, we found that, across all participants and

experimental trials, eyes exhibited significantly larger drift amplitudes when the stimulus

onset location was further away from the CDC ($\rho^2 = 0.06$, p < 0.001, Fig. 4g). The onset

distance was not correlated with drift directionality (Fig. 4g). Across all trials, the average

sampling cone density increased between stimulus onset and offset for most of the

participants. This sampling gain was computed as the ratio between the maximum sampling

density during the trial and the sampling density at the stimulus onset location. The sampling

gain was significantly correlated with the potential retinal sampling gain of individuals in

dominant ($r^2 = 0.35$, p = 0.02) as well as non-dominant eyes ($r^2 = 0.50$, p = 0.002, Fig. 4h).

Observers exploited on average 30 % of their potential sampling gain in both fellow eyes.

Interestingly, one observer combined all the previously described sampling features

particularly strong in his dominant eye (P08_R). It had a steep cone density gradient,

exhibited strong directional tuning towards the CDC and had large drift amplitudes for

stimulus onsets far from the CDC. This eye was excluded from the sampling gain analysis

because fixation behavior differed by more than 4 standard deviations from the group

average.

Discussion

By using synchronous adaptive optics imaging and visual stimulation of the foveola, we find

that the human visual system is capable of resolving spatial features smaller than a single

photoreceptor diameter and uncover a fixational eye motor behavior that optimizes retinal

sampling in accordance with the individual photoreceptor mosaic.

Spatial vision and, in particular, visual acuity is the most tested and used performance metric

with a close relation to everyday vision. It provides the main behavioral outcome for clinical

studies of vision. Measured in daily routine or clinical studies, best corrected visual acuity of

young and healthy adults is usually between 20/20 and 20/12.5 (60 and 37.5 arcsec)[44,45].

Even if lower order aberrations are corrected by e.g. glasses or contact lenses, higher order

aberrations inherently blur the retinal image, depending on their magnitude[45]. Adaptive

optics induce a close-to-diffraction limited optical correction, where the optical improvement

is significantly correlated with an increase in visual acuity thresholds[17]. By correcting

aberrations with AOSLO, we measured Snellen-E thresholds that were up to half the size

(between 20/10 and 20/6.9; 30 to 20.6 arcsec) compared to the natural viewing condition.

This is slightly lower than previously presented data[44], very likely because of different

wavelengths used for experimentation (Fig. 2c). It might be surprising to learn that the neural

machinery of human vision is able to resolve such tiny stimuli, because natural viewing is

blurred by the eye's optics. Even though observers are, to some degree, adapted to their

own aberrations[46], best subjective image quality is seen when on average 88 % of the

aberrations are corrected[47].

In how far are those resolution thresholds linked to or limited by the optimized but at the

same time individual morphology of the human foveola? While in the periphery, midget

retinal ganglion cell sampling dominates resolution, resolution of the foveal center was

estimated to be governed by the cone sampling limit[8,48]. By first-time direct experimental

validation in the same participants, we here confirm the hypothesis that the individual

spacing of cones can predict the resolution capacity of our foveola when optical influences

are bypassed (Fig. 2). We found that the individual spatial arrangement of cones was highly

correlated to the visual acuity of participants and explains up to 45 % of its variance (Fig.

2d). Eyes with higher foveolar sampling capacity reached lower thresholds than eyes with

less densely packed cone photoreceptors. Moreover, all participants reached resolution

thresholds that exceeded the Nyquist sampling limit when tested with near infrared, 788 nm

light. Natural vision is comprised of multiwavelength stimuli, thus, using 788 nm in isolation is

at the top end of our retinal sensitivity. In a first part of our study, participants also performed

experiments with 840 nm light. Thresholds were rather approximating the Nyquist limit with

this longer near-infrared wavelength (Fig. 2c). The L- and M-cone photopigment absorbance

for 840 nm is about 1.4 log unit lower than for 788 nm[49]. The decreased cone sensitivity

combined with a larger Airy-Disk size of about 7 % are likely to be detrimental for the longer,

840nm, wavelength. We would expect a potential for even lower thresholds for shorter

wavelengths.

Otherwise, a potential for lower thresholds is only expected in eyes with higher angular cone

densities. Perhaps contrary at first sight, this could potentially be the case for observers with

higher myopia. Most likely, the myopic eye growth lies between models of global expansion

and an equatorial stretching, which results in increased cone densities on an angular scale

and decreased cone densities on a linear scale for increased axial length[13]. The only two

myopic participants in our population had mild myopia (spherical equivalent between -1 and -

2 D) and axial lengths between 24.5 and 25 mm. Those two myopes had rather average

acuity thresholds, which is, in conjunction with the parallel finding of an overall below Nyquist

resolution, explainable by their average angular cone densities. Therefore, we would expect

acuity thresholds to be lower for myopic participants, in the case that (a) also the angular

cone density is increased like previously suggested and (b) the AO correction and display

resolution are still sufficient to completely resolve the foveolar cone mosaic. Psychophysical
data for more participants with higher myopia and longer axial lengths would be needed to
verify this assumption.

Theoretical predictions of the Nyquist resolution limit are implying stationary sampling. In
reality, however, our eyes are never at rest, even when we attempt to maintain steady
fixation. Fixational eye movements continuously modulate the luminance flow on individual
cones and postreceptoral neuronal activity. Drift motion has long been presumed as a
random jitter, a result of limited precision of the oculomotor system[50,51]. More recent work
revealed that drift motion is neither random nor detrimental due to the introduction of
noise[27,28], but rather a fine tuned motion, beneficial for psychophysical measures of visual
acuity in the parafovea[35] as well as foveola[26,36]. Neurons in the visual system are strongly
selective not just for spatial patterns, but also for temporally changing stimuli, a finding that is
also supported by computational modeling, suggesting that the visual system may utilize
principles comparable to those used in computational imaging for achieving super-resolution
via camera motion[52]. Within the past decades, the interdisciplinary term "geometrical
super-resolution" which is devoted to the filtering properties of sensor systems has become
common[53]. These resolution advantages may be achieved in the visual system by
incorporating mechanisms that allow for the recognition of positional differences smaller than
a single cell. That such mechanism exist is exemplified in a phenomenon known as
hyperacuity. Fine localization discriminations of only a few seconds of arc are performed by
identification of the centroid of the retinal light distributions[54] of the involved pattern
components. In a diffraction limited resolution task, the visual system seems to be able to
translate the temporal luminance modulation in individual photoreceptors by ocular drift to
additional spatial information about the stimulus position and shape. Contrary, the indirect
suppression of natural fixational eye motion by retinal stabilization techniques impairs visual
acuity outside the foveolar center[26,29]. For prolonged static stimulus presentations, retinal
spiking decays over time, while drift motion keeps the luminance change active, continuously
refreshes the receptive field input and sustains neuronal activity[24].

We found a significant correlation of drift motion and visual acuity thresholds between
individuals, indicating that drift motion may be one of the key elements in reaching sub-cone
resolution thresholds. Interestingly, acuity improved for smaller fixational drift and decreased
in participants who exhibited larger drift motion, on average. The fact that less drift is
beneficial to reach the lowest possible acuity thresholds reflects the characteristics of
spatiotemporal luminance changes introduced by smaller or larger drift motion. Smaller drifts
induce luminance changes with higher spatial frequencies and models of retinal ganglion cell
activity suggest a higher contrast sensitivity for high spatial frequency motion and less for

low spatial frequencies compared to a static retina[24,55]. This is supported by other recent
work which also showed that visual acuity thresholds can even be predicted from drift
magnitudes measured in a sustained fixation task[32].

There is evidence that fixational eye motion might have systematic components in primates.
A previous study in macaque monkeys revealed a systematic directional drift response only
a few dozens of milliseconds after various visual transients[56]. In our study, we reveal that a
certain drift directionality can not only be triggered by particular visual transients, but that
human observers are capable to adapt their drift direction to enact an oculomotor strategy
that takes advantage of the maximum resolution capacity provided within the retina. Our
participants precisely moved their eye to have the stimulus slip across the most densely
packed cone cells within their foveola. We hereby shed light on a mechanism that is
potentially particularly active during fine discrimination tasks. Thus, drift is not the long-
assumed random walk process between corrective saccades or microsaccades. And yet, the
underlying mechanism to drift motion remains not fully understood. Recent work suggested,
based on brainstem recordings in rhesus monkeys, that the origin can be found mostly
upstream of the ocular motoneurons. It can likely be explained as diffusion in the oculomotor
integrator which is mainly driven by noise, but additionally affected by mechanisms within the
visual motor pathway (e.g. feedback mechanisms)[34]. An incorporation of a visual feedback
loop to that model was shown to modulate the statistics of eye motion, given a time lag of
about 100 ms (mainly due to synaptic processing delays, of order 60-80 ms[56]). This fits our
results well. Our presentation time of 500 ms sufficed for a modulation of the fixational drift
motion towards retinal areas of higher cone sampling (also see supplementary discussion of
PRL displacement). This supports the view that the statistics of motion, but not the
superdiffusive nature of fixational drift can be influenced by the visual task[26,34,56,57]. The
superior colliculus seems to play a major role in modulating drift motion in a feedback loop to
visual inputs[31]. It's not only involved in controlling large eye motions[58] and microsaccades[59],
but also reflects neural responses to fixational drift that are likely a result of sensory input[60].

So, even though the CDC is displaced from the PRL in a way to be beneficial for natural
binocular vision[15], constant visual feedback allows to adapt the drift direction and therefore
also the task related PRL. Commonly, the term PRL is used for describing the retinal
location that is preferably used in fixational tasks. It is still a matter of debate what factors
drive the development of this very reproducible[15] retinal location and in how far it might
provide enhanced visual function. Sensitivity to small light spots in the foveola seems to be
rather plateau like and not particularly pronounced at the PRL[61]. As recently shown, the PRL
slightly differs between different tasks but has a larger interindividual variability[43]. The here
shown results indicate that also when measuring visual resolution, the PRL is not

necessarily the center of the sampling drift motion. The directional drift motion leads to a

shift of the preferred retinal location for a resolution task towards the CDC (Fig. S3 and

Supplementary Movie 3). Previous work that compared active versus passive fixation did not

show a systematic offset in a similar experimental setup. However, 5 out of 8 participants

also shifted their PRL in a Snellen E task closer to the CDC compared to the PRL for fixating

a static disk stimulus[43], the conditions that are best comparable to our study. The main

difference to our visual acuity experiments was that automatically paced random time

intervals between presentations (0.5 – 1.5 sec) were applied to not allow the participants to

anticipate the next trial whereas in our study participants self-paced the stimulus output to be

able to prepare and focus for the next trial. It might be that this extremely fine-tuned usage of

the visual feedback loop can only be kept active for rather short time intervals. By shifting the

stimulus towards the CDC in 50 % of cases the potential sampling gain within individual eyes

was exploited by 30%, on average, which goes along with a cone density increase of 3 % or

285 cones/deg^2. Even though this increase in cone density alone would not account for the

difference between acuity thresholds and Nyquist limit, this and the simultaneous

spatiotemporal luminance modulation contribute to achieving sub cone visual acuity

thresholds.

Between fellow eyes we found very strong correlations for all the measured parameters.

While drift amplitudes and directionality as well as cone densities are very symmetric

between dominant and non-dominant eyes (Fig. 2e and 3e), significantly lower acuity

thresholds of 1.5 arcsec, on average, were observed in the dominant eyes of participants

(Fig. 2e). The dominant eyes visual input has a tendency to be preferred during binocular

viewing, but has not been shown to exhibit relevant differences in visual function in healthy

eyes with low refractive errors[62,63]. Partially this may be due to limited accuracy in the mainly

used clinical methods (e.g. Snellen Chart or projection have ~ 10 arcsec steps between

optotype rows). This very fine binocular difference between eyes emphasizes that some

remaining factors which especially comprise the neural postprocessing steps, also play an

important role and may facilitate the slight functional advantage of dominant eyes.

For clinical studies of retinal health and in new therapeutical approaches, photoreceptor

health and visual acuity can be related to other more standard clinical measures as OCT-

derived measures of outer segment length or retinal thickness which have been shown to

serve for estimates of cone density[64]. Therefore, building a larger dataset on photoreceptor

resolved foveolar maps and associated visual function measures may help to, on the one

hand, better understand the interplay between structural and functional changes to draw

conclusions about disease progression, intervention efficiency or the interpretation of retinal

imaging data in studies aimed at vision restoration. On the other hand, detailed examination

of psychophysical measures with knowledge about the exact neural sampling characteristics

offers a great potential to answer further questions about e.g. resolution limits in myopia, the

effect of image stabilization in the very center of the foveola or implications for binocular

viewing that could previously only be hypothesized. The awareness of the oculomotor

system being able to finely adjust the drift motion behavior for a particular task may guide

future interpretation of fixational eye motion.

Material and Methods

Participants

A total of 38 participants underwent a preliminary screening where ocular biometry, ophthalmologic status, fixational eye motion and adaptive optics correction as well as foveolar image quality were tested. From those, 20 participants with normal ophthalmologic status, resolvable foveolar cones and ocular anatomy that allowed for a 7 mm pupil aperture during experimentation were chosen for subsequent examination. All 6 male and 14 female observers (17 adults [age: 18 – 42], 3 children [age: 10, 12 and 14]) had no or only mild refractive errors (SE: ± 2.5 diopters). The children and 15 adults were naïve participants and two adults were experienced observers. More detailed cone topography and eye motion characteristics of the here studied population have been shown previously[15]. The experiments were conducted under two different light conditions (16 participants 788 nm, 12 participants 840 nm). Eight participants took part in both experimental conditions. We mainly report the data acquired for the 788 nm condition in this manuscript and show 840 nm data for comparison where noteworthy differences arise.

Written informed consent was obtained from each participant and all experimental procedures adhered to the tenets of the Declaration of Helsinki, in accordance with the guidelines of the independent ethics committee of the medical faculty at the Rheinische Friedrich-Wilhelms-Universität of Bonn.

Ocular dominance

Ocular dominance was determined by a Miles Test prior to pupil dilation and visual acuity testing. The experimenter stood in a distance of 6 m in front of the participant and asked them to form a small opening between thumbs and forefingers with both hands. The participant was then asked to extend their arms in front of them to look through the formed hole at the experimenter's face with both eyes open. This procedure was conducted 3-5 times to determine the dominant (= uncovered) eye in a 3/3 or at least 3/5 condition.

AOSLO retinal imaging

In vivo images of the complete foveolar cone mosaic were recorded using a custom-built adaptive optics scanning laser ophthalmoscope (AOSLO). The general setup of the AOSLO has been described previously[65] and pertinent differences as well as the method of determination of the preferred retinal locus of fixation (PRL) have been described in a recent publication[15].

In brief, the front-end of the Adaptive Optics Scanning Laser Ophthalmoscope (AOSLO) was

equipped with three f = 500mm focal telescopes. These telescopes were specifically

designed for point-scanning an adaptive optics-corrected focal light spot across the retina,

ensuring diffraction-limited resolution in both incident and reflected beams. The system

incorporated a magnetic actuator-driven deformable mirror (DM97-07, 7.2mm pupil diameter,

ALPAO, Montbonnot-Saint-Martin, France) positioned in a retinal conjugate plane. The

deformable mirror was controlled by the wavefront error signals from a 25x25 lenslet Shack

Hartmann sensor (SHSCam AR-S-150-GE, Optocraft GmbH, Erlangen, Germany) in closed-

loop. Imaging and wavefront correction utilized wavelengths of either 788 nm (±12 nm) or

840 nm (±12 nm) light, achieved through serial dichroic and bandpass filtering of a

supercontinuum source (SuperK Extreme EXR-15, NKT Photonics, Birkerød, Denmark). The

imaging field of view was 0.85 x 0.85 degrees of visual angle. The digital lateral resolution

was about 0.1 arcmin, the size of one pixel in the recorded videos and images. Light

reflected from the retina was detected by a photomultiplier tube (PMT, H7422-50,

Hamamatsu Photonics, Hamamatsu, Japan), positioned behind a confocal pinhole (Pinhole

diameter = 20 mm, equivalent to 0.47 (840nm) and 0.5 (788nm) Airy disk diameters).

Continuous sampling of the PMT signal was carried out using a field programmable gate

array (FPGA), resulting in a 512 x 512-pixel video at 30 Hz (600 pixels per degree of visual

angle). Through rapid acousto-optic intensity modulation of the imaging lights, the square

AOSLO imaging field was used as retinal display, where each pixel could be individually

controlled to produce the visual stimuli.

Cone map generation and computation of sampling characteristics

The best PRL videos acquired were selected to create spatially registered, high signal-to-

noise ratio images of the foveal center, which served as master retinal images for cone

labeling as well as referencing of stimulus motion trajectories. This study includes only

participants for whom the master retinal image was of sufficient quality to label all cones

across the image. Cone centers were identified and labeled semi manually, as previously

described[15,66]. Cone density was computed in two different ways. First, for deriving landmark

metrics of the foveolar cone map, we then computed Voronoi tessellation, estimating a patch

with certain area for each individual cone and summed the nearest 150 cone patches around

each image pixel. The number of cells was divided by the resulting area to derive a pixel-

resolved map of cone densities. Based on this map, the peak cone density (PCD) is defined

as the highest cone density value of the map with it's according retinal location. The cone

density centroid (CDC) is computed as the weighted centroid of the 20[th] percentile of highest

cone densities within the map. We refer to the CDC as the anatomical center and the anchor

for further spatial analyses in this study. The CDC has been shown to be a more robust and
reproducible metric to describe the anatomical center than the more routinely reported peak
cone density (PCD)[15,67].

Second, for analyzing the relation between individual sampling limits and resolution acuity,
cone density was computed based on the cone cells contributing to the sampling process.
To identify the cones interacting in stimulus sampling, a simple model of cone light capture
was employed. Each cone was described by an associated light acceptance aperture with its
diameter estimated as 48 % of the average spacing between the cone and all of its
neighbors. The efficiency of the aperture along its diameter was approximated as Gaussian
profiles. Also, a model of the stimulus retinal image was computed by convolving the eye's
point spread function (diffraction limited at 788 nm for a 7 mm pupil) with the stimulus
bitmap. The complete two-dimensional model of cone apertures was then multiplied by
models of the presented stimuli to arrive at the cone-level light distribution based on the
different stimulus positions, sizes and orientations. The light distribution within each cone
was integrated across the entire cone aperture. This value was then normalized to the
degree to which the aperture was filled. Cone stimulation was considered to be maximal if
the entire aperture was filled. Using this method, a cone activation pattern could be
generated for each point in time (e.g. Fig. 1f). To arrive at a task-related cone density
estimate for each frame (sampling cone density), the number of cones identified to interact
with the stimulus was divided by their summed cone area. In the presented analyses, the
median sampling density of all trials is analyzed and standard deviations are shown as grey
lines (Fig. 2d, e). This stimulus related cone density was chosen to closely represent the
sampling process; however, the results do not qualitatively differ from using the cone density
map based on the 150 nearest cones.

We assumed a perfect hexagonal cell mosaic to estimate the average inter-cone-distance
(ICD) between neighboring cells and to compute the theoretical Nyquist sampling limit, which
is based on the spacing between rows of cones, and given by $N = \frac{\sqrt{3}}{2} \times ICD$.

Experimental procedures

For psychophysical acuity testing, participants reported the orientation of a Snellen-E
stimulus in a four-alternative forced-choice (4 AFC) task under unrestricted eye motion.
Psychophysical experiments were performed monocularly in both eyes. The non-dominant
eye was tested first and the dominant eye after a 15-30 minutes break. This protocol was
chosen because in pilot experiments in 7 participants (which were performed with a random
order) less time was needed and hence less fatigue was reported by the participants when

the second eye was the dominant one. In these pilot experiments, the same qualitative
difference of acuity thresholds between non-dominant and dominant were found.

Mydriasis and cycloplegia were established by two drops of 1% tropicamide, instilled into the
eyelid about 25 and 20 minutes prior to experiments. If experimentation took longer than 40
minutes, another drop of tropicamide was instilled. A customized dental impression mold
(bite bar) was used to immobilize and adjust the head position and thus to align the
participants eye in front of the imaging system to ensure optimal adaptive optics correction
and image quality. The participants were encouraged to take breaks at any time. We found
that proper resting is one of the most crucial factors during the rather complex AOSLO
experimentation. Frequent breaks ensure constant, high-level compliance and excellent
image quality as the basis for artefact-free and reproducible results.

Before recording experimental runs, each participant performed 3 test runs to get used to the
test procedure and the appearance of the stimuli. The stimuli were displayed as "off-stimuli"
on the infrared background by switching the displayed intensity via an acousto-optic
modulator [68] (AOM, TEM-250-50-10-840-2FP, Brimrose, Sparks Glencoe, MD, USA)(Fig.
1a). Because of ocular diffraction, the stimulus contrast varied between 0.61 and 0.80 for an
18 arcsec versus 36 arcsec gap sized stimulus (3 and 6 pixel of the scanning raster,
respectively). The visual acuity testing followed the Bayesian adaptive procedure QUEST[69–
71]. Stimulus progression was self-paced by the participant. The stimuli were presented for
500 ms to avoid limitations by insufficient temporal summation[72]. Around each trial, a one
second AOSLO video was recorded, with the stimulation onset at around 300ms after video
onset. Visual acuity thresholds were estimated by pooling results from 5 consecutively run
staircases, with each containing 20 trials. A psychometric function was fitted using
psignifit4[73] to derive threshold estimates for further analysis. The expected threshold
variance is described and visualized by the 95 % confidence interval (Fig. 1d, e and 2d).

Video processing and eye motion analysis

The AOSLO used a raster scanning technique where each frame was acquired over time.
The recorded videos were stabilized after psychophysical testing using custom settings
within the *MATLAB* based stabilization software from Stevenson et al.[74]. To acquire eye
traces at higher temporal resolution than the 30 Hz frame rate, each frame of the AOSLO
movie is broken into 32 horizontal strips of 16 pixels height and cross-correlated against a
reference frame. The reference frame was generally chosen automatically and exchanged
by a manually chosen frame in cases were stabilization failed despite good overall image

quality. This method allowed extraction of eye motion traces at temporal frequencies up to
960 Hz.

The frame-wise (30 Hz) stimulus position was encoded as a white cross marker in each
video. As single strip alignments can have small errors due to noise in the strip or retinal
torsion (particularly affecting the horizontal motion estimate)[75], we compute the average
offsets from the cross-containing strip and 2 previous/subsequent strips. These steps yielded
more accurate trajectories in retinal coordinates for every trial. All individual trial AOSLO
frames and the corresponding trajectories are then referenced to the single master retinal
image used for cone map generation.

To quantify the retinal motion across the stimulus, drift amplitude was defined as the
concatenated vector sum of all frame-wise motion vectors within the 500 ms stimulus
duration (see also Fig. 3c). Trials that contained microsaccades or blinks during stimulus
presentation were excluded from further analyses. Microsaccade occurrence varied highly
between participants (mean ± SD: 14 ± 10 % of trials, range: 2 % - 41 %). If not stated
differently, we here report the median drift amplitude of all trials for individual eyes (e.g. of all
traces shown in Fig. 3a). To quantify drift direction, the angle between each trajectory's
starting coordinate (coordinate center in Fig. 4c) and end coordinate was computed. To
check for potential motion bias, the drift angles were first analyzed in retinal coordinates (Fig.
4c), and then as the relative angle, $\theta_{CDC,}$ formed between the drift vector and the line
connecting the retinal onset location and the CDC (Fig. 4d). To compare directionality
towards other locations of interest, the same was done for PRL and PCD locations (Fig. 4f).

Statistical information

All statistical analyses were conducted using custom written MATLAB code and significance
levels were set at 0.05. To assess the normal distribution of the dataset, a two-sided
Shapiro–Wilk test was employed. This test is recognized to be appropriate for small sample
sizes. The paired samples t-test was utilized to assess whether there were significant
differences between the means of normally distributed paired observations. For non-
parametric data, the Wilcoxon Signed-Rank test was employed. Linear correlations were
computed to examine the relationships between variables. For variables demonstrating
normal distribution, Pearson's correlation coefficient was employed, while for non-normally
distributed data, Spearman's rank correlation coefficient was utilized. Pearson's correlation
is sensitive to linear relationships, assuming bivariate normality, whereas Spearman's
correlation is a non-parametric measure suitable for monotonic relationships and is robust
against outliers and non-normal distributions.

Author contributions

700   J.L.W and W.M.H conceived the research idea and developed the data analysis pipeline.

701   J.L.W performed the experiments and data analyses. J.L.W, V. L. and W.M.H discussed the

702   results and wrote the manuscript.

Competing interests

705   The authors declare no competing interests.

References

1. Bohigian, G. M. An Ancient Eye Test-Using the Stars. *Surv. Ophthalmol.* **53**, 536–539 (2008).

2. Caves, E. M., Brandley, N. C. & Johnsen, S. Visual Acuity and the Evolution of Signals. *Trends Ecol. Evol.* **33**, 1–15 (2018).

3. Tuten, W. S. & Harmening, W. M. Foveal vision. *Curr. Biol.* **31**, R701–R703 (2021).

4. Hendrickson, A. E. & Yuodelis, C. The morphological development of the human fovea. *Ophthalmology* **91**, 603–612 (1984).

5. Syrbe, S. *et al.* Müller glial cells of the primate foveola: An electron microscopical study. *Exp. Eye Res.* **167**, 110–117 (2018).

6. Williams, D. R. & Coletta, N. J. Cone spacing and the visual resolution limit. *J. Opt. Soc. Am. A* **4**, 1514–1523 (1987).

7. Hirsch, J. & Curcio, C. A. The spatial resolution capacity of human foveal retina. *Vision Res* **29**, 1095–1101 (1989).

8. Rossi, E. A. & Roorda, A. The relationship between visual resolution and cone spacing in the human fovea. *Nat Neurosci* **13**, 156–157 (2010).

9. Curcio, C. A., Packer, O. & Kalina, R. E. A whole mount method for sequential analysis of photoreceptor and ganglion cell topography in a single retina. *Vision Res* **27**, 9–15 (1987).

10. Rossi, E. A. *et al.* Imaging retinal mosaics in the living eye. *Eye* **25**, 301–308 (2011).

11. Zhang, C. *et al.* Circuit Reorganization Shapes the Developing Human Foveal Midget Connectome toward Single-Cone Resolution. *Neuron* **108**, 905–918 (2020).

12. Curcio, C. A., Sloan, K. R., Kalina, R. E. & Hendrickson, A. E. Human photoreceptor topography. *J. Comp. Neurol.* **292**, 497–523 (1990).

13. Wang, Y. *et al.* Human foveal cone photoreceptor topography and its dependence on eye length. *Elife* **8**, 1–21 (2019).

14. Cava, J. A. *et al.* Assessing Interocular Symmetry of the Foveal Cone Mosaic. *Invest Ophthalmol Vis. Sci.* **61**, 1–11 (2020).

15. Reiniger, J. L., Domdei, N., Holz, F. G. & Harmening, W. M. Human gaze is systematically offset from the center of cone topography. *Curr. Biol.* **31**, 4188–4193 (2021).

16. Campbell, F. W. & Green, D. G. Optical and retinal factors affecting visual resolution. *J. Physiol.* **181**, 576–93 (1965).

17. Marcos, S., Sawides, L., Gambra, E. & Dorronsoro, C. Influence of adaptive-optics ocular aberration correction on visual acuity at different luminances and contrast polarities. *J Vis* **8**, 1–12 (2008).

18. Westheimer, G. Visual Acuity and Hyperacuity. *Invest. Ophthalmol.* **64**, 570–572 (1975).

19. Westheimer, G. Optical superresolution and visual hyperacuity. *Prog. Retin. Eye Res.* **31**, 467–480 (2012).

20. Nagano, T. Temporal sensitivity of the human visual system to sinusoidal gratings. *J. Opt. Soc. Am.* **70**, 711–716 (1980).

21. Kaplan, E. & Benardete, E. The dynamics of primate retinal ganglion cells. *Prog. Brain Res.* **134**, 17–34 (2001).

22. Ko, H. K., Poletti, M. & Rucci, M. Microsaccades precisely relocate gaze in a high visual acuity task. *Nat. Neurosci.* **13**, 1549–1554 (2010).

23. Poletti, M., Rucci, M. & Carrasco, M. Selective attention within the foveola. *Nat. Neurosci.* **20**, 1413–1417 (2017).

24. Kuang, X., Poletti, M., Victor, J. D. & Rucci, M. Temporal encoding of spatial information during active visual fixation. *Curr. Biol.* **22**, 510–514 (2012).

25. Krauskopf, J., Cornsweet, T. N. & Riggs, L. A. Analysis of eye movements during monocular and binocular fixation. *J. Opt. Soc. Am.* **50**, 572–578 (1960).

26. Intoy, J. & Rucci, M. Finely tuned eye movements enhance visual acuity. *Nat. Commun.* **11**, 1–11 (2020).

27. Burak, Y., Rokni, U., Meister, M. & Sompolinsky, H. Bayesian model of dynamic image stabilization in the visual system. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 19525–19530 (2010).

28. Pitkow, X., Sompolinsky, H. & Meister, M. A neural computation for visual acuity in the presence of eye movements. *PLoS Biol.* **5**, 2898–2911 (2007).

29. Rucci, M., Iovin, R., Poletti, M. & Santini, F. Miniature eye movements enhance fine spatial detail. *Nature* **447**, 851–854 (2007).

30. Rolfs, M. Microsaccades: Small steps on a long way. *Vision Res* **49**, 2415–2441 (2009).

31. Hafed, Z. M., Chen, C. Y., Tian, X., Baumann, M. P. & Zhang, T. Active vision at the foveal scale in the primate superior colliculus. *J. Neurophysiol.* **125**, 1121–1138 (2021).

32. Clark, A. M., Intoy, J., Rucci, M. & Poletti, M. Eye drift during fixation predicts visual acuity. *Proc. Natl. Acad. Sci. U. S. A.* **119**, 1–10 (2022).

33. Herrmann, C. J. J., Metzler, R. & Engbert, R. A self-avoiding walk with neural delays as a model of fixational eye movements. *Sci. Rep.* **7**, 1–17 (2017).

34. Ben-Shushan, N., Shaham, N., Joshua, M. & Burak, Y. Fixational drift is driven by diffusive dynamics in central neural circuitry. *Nat. Commun.* **13**, 1697 (2022).

35. Ratnam, K., Domdei, N., Harmening, W. M. & Roorda, A. Benefits of retinal image motion at the limits of spatial vision. *J Vis* **17**, 1–11 (2017).

36. Nghiem, T.-A. E., Dufoura, O., Reiniger, J. L., Harmening, W. M. & da Silveira, R. A. Fixational

eye movements as active sensation for high visual acuity. *PNAS - to be Updat. when Publ.* (2022).

37. Cherici, C., Kuang, X., Poletti, M. & Rucci, M. Precision of sustained fixation in trained and untrained observers. *J Vis* **12**, 1–16 (2012).

38. Putnam, N. M. *et al.* The locus of fixation and the foveal cone mosaic. *J Vis* **5**, 632–639 (2005).

39. Wilk, M. A. *et al.* Assessing the spatial relationship between fixation and foveal specializations. *Vision Res* **132**, 53–61 (2017).

40. Zhang, T. *et al.* Variability in Human Cone Topography Assessed by Adaptive Optics Scanning Laser Ophthalmoscopy. *Am. J. Ophthalmol.* **160**, 290–300 (2015).

41. Wells-Gray, E. M., Choi, S. S., Bries, A. & Doble, N. Variation in rod and cone density from the fovea to the mid-periphery in healthy human retinas using adaptive optics scanning laser ophthalmoscopy. *Eye* **30**, 1135–1143 (2016).

42. Intoy, J., Mostofi, N. & Rucci, M. Fast and nonuniform dynamics of perisaccadic vision in the central fovea. *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

43. Bowers, N. R., Gautier, J., Lin, S. & Roorda, A. Fixational eye movements depend on task and target. *bioRxiv* (2021) doi:10.1101/2021.04.14.439841.

44. Rossi, E. a, Weiser, P., Tarrant, J. & Roorda, A. Visual performance in emmetropia and low myopia after correction of high-order aberrations. *J Vis* **7**, 14 (2007).

45. Reiniger, J. L. *et al.* Habitual higher order aberrations affect Landolt but not Vernier acuity. *J Vis* **19**, 1–15 (2019).

46. Artal, P. *et al.* Neural compensation for the eye's optical aberrations. *J Vis* **4**, 4 (2004).

47. Chen, L., Artal, P., Gutierrez, D. & Williams, D. R. Neural compensation for the best aberration correction. *J Vis* **7**, 9.1-9 (2007).

48. Williams, D. R. Aliasing in human foveal vision. *Vision Res* **25**, 195–205 (1985).

49. Stockman, A. & Rider, A. T. Formulae for generating standard and individual human cone spectral sensitivities. *Color Res. Appl.* **48**, 818–840 (2023).

50. Ditchburn, R. W. & Ginsborg, B. L. Involuntary eye movements during fixation. *J. Physiol.* **119**, 1–17 (1953).

51. Cornsweet, T. N. Determination of the stimuli for involuntary drifts and saccadic eye movements. *J. Opt. Soc. Am.* **46**, 987–993 (1956).

52. Anderson, A. G., Ratnam, K., Roorda, A. & Olshausen, B. A. High-acuity vision from retinal image motion. *J Vis* **20**, 1–19 (2020).

53. Zalevsky, Z. Exceeding the diffraction and the geometric limits of imaging systems: A review. in *Dolev, S., Oltean, M. (Eds.), Optical Supercomputing 2010. Lecture Notes in Computer*

Science vol. 6748 119–130 (Springer-Verlag, 2011).

54. Westheimer, G. & McKee, S. P. Spatial configurations for visual hyperacuity. *Vision Res* **17**, 941–7 (1977).

55. Rucci, M. & Victor, J. D. The unsteady eye: An information-processing stage, not a bug. *Trends Neurosci.* **38**, 195–206 (2015).

56. Malevich, T., Buonocore, A. & Hafed, Z. M. Rapid stimulus-driven modulation of slow ocular position drifts. *Elife* **9**, 1–21 (2020).

57. Zhao, Z., Ahissar, E., Victor, J. D. & Rucci, M. Fine oculomotor knowledge enhances high-acuity vision. *bioRxiv* doi:10.1101/2022.03.17.483854.

58. Bergeron, A., Matsuo, S. & Guitton, D. Superior colliculus encodes distance to target, not saccade amplitude, in multi-step gaze shifts. *Nat. Neurosci.* **6**, 404–413 (2003).

59. Hafed, Z. M., Goffart, L. & Krauzlis, R. J. A neural mechanism for microsaccade generation in the primate superior colliculus. *Science (80-.).* **323**, 940–943 (2009).

60. Chen, C.-Y., Hoffmann, K.-P., Distler, C. & Hafed, Z. M. The Foveal Visual Representation of the Primate Superior Colliculus. *Curr. Biol.* **29**, 2109–2119 (2019).

61. Domdei, N., Reiniger, J. L., Holz, F. G. & Harmening, W. M. The Relationship Between Visual Sensitivity and Eccentricity, Cone Density and Outer Segment Length in the Human Foveola. *Invest Ophthalmol Vis Sci* **62**, 1–16 (2021).

62. Ehrenstein, W. H., Arnold-Schulz-Gahmen, B. E. & Jaschinski, W. Eye preference within the context of binocular functions. *Graefe's Arch. Clin. Exp. Ophthalmol.* **243**, 926–932 (2005).

63. Zhou, D. *et al.* Association of Visual Acuity with Ocular Dominance in 2045 Myopic Patients. *Curr. Eye Res.* **42**, 1155–1159 (2017).

64. Domdei, N. *et al.* Cone Density Is Correlated to Outer Segment Length and Retinal Thickness in the Human Foveola. *Invest Ophthalmol Vis Sci* **64**, 1–11 (2023).

65. Roorda, A. *et al.* Adaptive optics scanning laser ophthalmoscopy. *Opt Express* **10**, 405–412 (2002).

66. Cunefare, D. *et al.* Open source software for automatic detection of cone photoreceptors in adaptive optics ophthalmoscopy using convolutional neural networks. *Sci. Rep.* **7**, 1–11 (2017).

67. Wynne, N. *et al.* Intergrader agreement of foveal cone topography measured using adaptive optics scanning light ophthalmoscopy. *Biomed. Opt. Express* **13**, 4445–4454 (2022).

68. Poonja, S., Patel, S., Henry, L. & Roorda, A. Dynamic visual stimulus presentation in an adaptive optics scanning laser ophthalmoscope. *J Refract Surg* **21**, 575–580 (2005).

69. Watson, A. B. & Pelli, D. G. QUEST: a Bayesian adaptive psychometric method. *Percept. Psychophys.* **33**, 113–120 (1983).

70.   Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

71.   Pelli, D. G. The VideoToolbox software for visual psychophysics: transforming numbers into movies. *Spat. Vis.* **10**, 437–442 (1997).

72.   McAnany, J. J. The effect of exposure duration on visual acuity for letter optotypes and gratings. *Vision Res* **105**, 86–91 (2014).

73.   Schütt, H. H., Harmeling, S., Macke, J. H. & Wichmann, F. A. Painfree and accurate Bayesian estimation of psychometric functions for (potentially) overdispersed data. *Vision Res* **122**, 105–123 (2016).

74.   Stevenson, S. B. & Roorda, A. Correcting for miniature eye movements in high-resolution scanning laser ophthalmoscopy. in *Ophthalmic Technologies XV* vol. 5688 145–151 (SPIE, 2005).

75.   Hofmann, J., Domdei, L., Jainta, S. & Harmening, W. M. Assessment of binocular fixational eye movements including cyclotorsion with split-field binocular scanning laser ophthalmoscopy. *J Vis* **22**, 1–13 (2022).