Museomics of *Carabus* giant ground beetles evidences an Oligocene origin and *in situ* Alpine diversification

Authors

Marie T. Pauli[1,2,†], Jérémy Gauthier[1,2,†], Marjorie Labédan[1,3], Mickael Blanc[1], Julia Bilat[1], Emmanuel F.A. Toussaint[1,*]

Affiliations

[1] *Natural History Museum of Geneva, Geneva, Switzerland*
[2] *Cantonal Museum of Natural Sciences, Lausanne, Switzerland*
[3] *Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland*

† Joint first authors

* Correspondence: Emmanuel F.A. Toussaint, Natural History Museum of Geneva, 1208 Geneva, Switzerland. Email: emmanuel.toussaint@ville-ge.ch.

ORCID numbers
Marie T. Pauli (0000-0003-1488-9851)
Jérémy Gauthier (0000-0001-6666-1002)
Marjorie Labédan (0000-0002-2643-1277)
Mickael Blanc (0009-0005-1833-0854)
Julia Bilat (0009-0000-9249-7103)
Emmanuel F.A. Toussaint (0000-0002-8439-1285)

Abstract: The development of museomics represents a major paradigm shift in the use of natural
history collection specimens for systematics and evolutionary biology. New approaches in this
field allow the sequencing of hundreds to thousands of loci from across the genome using historical
DNA. HyRAD-X, a recently introduced capture method using bench-top designed probes, has
proved very efficient to recover genomic-scale datasets using natural history collection specimens.
Using this technique, we infer at both the intra- and interspecific levels, the most robust phylogeny
of Arcifera to date, an ecologically and morphologically diverse clade of *Carabus* giant ground
beetles. We successfully generated a genomic dataset of up to 1965 HyRAD-X loci for all
described species, permitting to infer a robust dated phylogenomic tree of this clade. Our species
delimitation and population genomic analyses suggest that the current classification in Arcifera is
in line with its evolutionary history. Our results suggest an origin of Arcifera in the late Oligocene
followed by speciation events during the warm mid-Miocene unlinked to Pleistocene glaciations.
The dynamic paleogeographic history of the Palearctic region likely contributed to the
diversification of this lineage with a relatively ancient colonization of the proto-Alps followed by
in situ speciation where most species of Arcifera are currently found sometimes syntopically likely
as a result of post-glaciations secondary contacts.

Keywords: Arcifera; Beetle evolution; Carabinae; Historical DNA; HyRAD-X; Phylogenomics;
Palearctic biogeography; Pleistocene glaciations

47
48
49

Introduction

Recent developments in museomics are opening new prospects allowing samples from natural history collections (NHC) to enter the era of genomics (reviewed in (Raxworthy & Smith 2021; Card *et al.* 2021). Specimens held in the collections are crucial for the study of systematics and taxonomy, but also for the study of ecology and evolution (Duchenne *et al.* 2020). Sampling from NHC specimens is also a major asset to study groups that are currently rare in the wild, for which authorizations to collect new specimens are difficult to obtain, or for which a comprehensive taxonomic and geographic sampling would require extensive fieldwork campaigns. Such a strategy is therefore very powerful when working on taxonomic groups presenting a wide geographical range. In extreme cases, and when species are believed to be extinct, NHC represent the only potential source of genetic data (Toussaint *et al.* 2021; de-Dios *et al.* 2023).

Innovative approaches are now making it possible to obtain genetic information from NHC specimens for which it has long been impossible to recover DNA. The DNA in these specimens, referred to as historical DNA (hDNA), is in low quantity, fragmented, has undergone chemical modifications over time and contains contaminants linked to the history of the collection (Raxworthy & Smith 2021). Improvements in extraction methods, sequencing technologies but above all the development of new capture methods has enabled an increasing amount of genetic information to be recovered. They allow to overcome the difficulties associated with highly degraded and fragmented hDNA from NHC samples, which prevents conventional amplification using standard molecular primers (Landry *et al.* 2023). Among these methods, Ultra Conserved Elements (Blaimer *et al.* 2016; Faircloth 2017) or anchored hybrid enrichment of conserved regions (AHE, Lemmon *et al.* 2012; Mayer *et al.* 2021), are based on the capture of informative loci previously designed from existing genomic data and generally target fairly conserved regions in order to perform large phylogenies. Applying these approaches to NHC specimens allows to integrate samples that are complicated to obtain in the field. In order to work on non-model species for which no prior genomic data is available, the HyRAD (Suchan *et al.* 2016) and HyRAD-X (Schmid *et al.* 2017) approaches enable probes to be designed directly from a few phylogenetically close fresh samples. These approaches, based on bench-top production of probes, also make it possible to dispense with the high cost of probe synthesis. Probes are designed using ddRADseq protocol (Peterson *et al.* 2012) allowing to target thousands of loci randomly distributed along the genome. This approach is suitable for integrating NHC samples into population-scale studies (Gauthier *et al.* 2020) or for phylogenetic studies of taxa that have recently diverged, such as within a genus (Gauthier *et al.* 2023). The HyRAD-X approach designs probes on fresh RNA extractions. By targeting only expressed gene loci, the HyRAD-X approach makes it possible to investigate phylogenetic questions at older evolutionary scales than the HyRAD approach (Toussaint *et al.* 2021). Using these probe sets, hDNA is then captured by hybridization and sequenced using NGS technologies. This allows only the targeted loci to be recovered while eliminating all unwanted fragments such as contaminants. After sequencing, the loci are reconstructed and aligned using appropriate bioinformatic pipelines in order to make phylogenetic inferences (Toussaint *et al.*

2021). Unlike random Whole Genome Sequencing (WGS) of all the extracted DNA, these targeted
approaches enable better recovery of loci and integration of a larger number of NHC samples into
the phylogenetic inferences made in fine (Toussaint *et al.* 2021). Although the efficiency of
HyRAD-X has been tested at higher taxonomic levels, an empirical investigation of its
performance at the interface between population and species levels is needed.
The genus *Carabus*, Linnaeus 1758 (Coleoptera: Carabidae), is a monophyletic highly
diversified lineage comprising *ca.* 970 species classified into 91 subgenera (Deuve 2019, 2021).
This genus, together with its sister genus *Calosoma* (cosmopolitan, 130 species) form the tribe
Carabini (Osawa *et al.* 2004; Toussaint Fls & Gillett 2018; Toussaint *et al.* 2021; Sota *et al.* 2022).
Within *Carabus*, the clade named Arcifera Imura, 1996 is sister to the very diversified clade
Eucarabi Deuve, 2013 (Deuve *et al.* 2012; Deuve 2021). This clade is mainly Palearctic, ranging
in the west from southwest England to Ukraine and Turkey in the east. The range of this group
notably encompasses the Carpathian mountains as well as the Swiss, Italian, Austrian and Dinaric
Alps. It currently includes four subgenera: *Carabus* (*Hygrocarabus*) Thomson, 1875, *Carabus*
(*Platycarabus*) Morawitz, 1886, *Carabus* (*Chaetocarabus*) Thomson, 1875 and *Carabus*
(*Heterocarabus*) Morawitz, 1886 (Deuve 2019, 2021). Within Arcifera, the subgenus *Carabus*
(*Hygrocarabus*) contains two species found from France to Ukraine, the status of which has been
extensively debated over the past decades due to reduced morphological differences and
inconsistent genetic admixture patterns (Müller-Kroehling *et al.* 2006; Müller-Kroehling *et al.*
2014); Matern *et al.* 2009, 2010; Mossakowski *et al.* 2020). These hygrophilous nocturnal species
live in river banks and hunt close or in the water of cold forest streams. The two species are in
relative allopatry with *Carabus nodulosus* Creutzer, 1799 being found from eastern France to
Austria and western Balkans, and *Carabus variolosus* Fabricius, 1787 from Slovakia to Ukraine
and Bulgaria (Kulijer 2019; Deuve 2021; Bekchiev *et al.* 2022; Hristovski *et al.* 2023). Despite
the protection status of their habitat (Annexes II and IV of the European Union's Habitats
Directive), these two species appear to be declining due to anthropogenic activities and their
consequences (Tyszecka *et al.* 2023). The subgenus *Carabus* (*Chaetocearabus*) only contains two
allopatric species following Deuve (Deuve 2019, 2021), the widespread *Carabus intricatus* Linné,
1761 found from western France to Ukraine and Greece, and the Greek endemic *Carabus*
arcadicus Gistl, 1850. These two species are found in sympatry in Greece where hybrids are
known from example at the Katara Pass in the Epirus region. Additionally the status of several
subspecies in both taxa has been debated, and some authors recognize *Carabus arcadicus merlini*
Schaum, 1861 (Greece), *Carabus intricatus lefebvrei* Dejean, 1826 (southern Italy including
Sicily) and *Carabus intricatus krueperi* Reitter, 1896 (Greece) as separate species within which
additional subspecific taxa have been described (e.g., Cavazzuti & Ghiretti 2020). Perhaps the
most debated taxon of the three being *Carabus intricatus lefebvrei* found south of Umbria to
northern Sicily which is largely allopatric from the rest of the Italian populations found only in the
extreme north of Italy from Piemonte to Friuli (Cavazzuti & Ghiretti 2020). The subgenus *Carabus*
(*Heterocarabus*) contains a unique species, *Carabus marietti* Cristofori & Jan, 1837, that is found
in southern Bulgaria near the Black Sea and in Anatolia (Turkey), however its ecology and

relationships between the numerous described subspecies remain poorly known (Gueorguiev &
Gueorguiev 1995; Hieke & Wrase 2008). Finally, the subgenus *Carabus* (*Platycarabus*) is
composed of five currently accepted species: *Carabus creutzeri* Fabricius, 1801, *Carabus*
cychroides Baudi, 1860, *Carabus depressus* Bonelli, 1811, *Carabus fabricii* Panzer, 1812 and
Carabus irregularis Fabricius, 1792. These beetles are characterized by a flattened morphology,
long legs, and elytra generally covered with small foveoli (except in *Carabus depressus lucens*
Schaum, 1857). They are most widely distributed in Central and Eastern Europe, generally at high
altitudes, in mountain forests and alpine pastures. The subgenus contains helicophagous species
that exhibit different hunting techniques related to the morphology of their mandibles and
prothorax (Casale *et al.* 1998). For instance, *Carabus cychroides* with a thin, elongated head and
prothorax, is adapted to enter gastropod shells and has undergone a process known as
"cychrization". This species has a very restricted range in the Piedmont region of Italy, is
endangered and the focus of reinforced conservation programmes (Anselmo & Rizzioli 2022a; b).
In contrast, the species *C. irregularis* presents a "licinization" resulting in a large head likely
adapted to cracking snail shells (Casale *et al.* 1998). The relationships among species of the
subgenus *Carabus* (*Platycarabus*) are still debated, and the various taxonomic divisions, both
species and subspecies, have yet to be clarified (Casale *et al.* 1998; Deuve 2021). Natural hybrids
have been suggested between *Carabus fabricii* and *Carabus depressus*, *Carabus creutzeri* and
Carabus irregularis, *Carabus creutzeri* and *Carabus depressus*, and *Carabus depressus* and
Carabus cychroides (Casale *et al.* 1998; Camard & Leplat 2004; Casale & Rapuzzi 2015),
indicating the need for an in-depth study of possible hybridization in this group.
One of the earliest attempts to elucidate the phylogeny of Arcifera was conducted by
Ishikawa (Ishikawa 1984), using 21 morphological characters. This study supported the
monophyly of Arcifera and placed *Carabus* (*Hygrocarabus*) as sister to the rest of the group, in
which *Carabus* (*Chaetocarabus*) was sister to *Carabus* (*Heterocarabus*) and *Carabus*
(*Platycarabus*). The first placement of Arcifera members in a molecular phylogeny was based on
a single mitochondrial fragment (i.e. ND5), and recovered *Carabus* (*Chaetocarabus*) and *Carabus*
(*Platycarabus*) as sister lineages, close to *Carabus* (*Limnocarabus*) Géhin, 1876 and *Carabus*
(*Euleptocarabus*) Nakane, 1956 (Imura *et al.* 1998). A subsequent study with the same gene
fragment but increased taxon sampling recovered a paraphyletic Arcifera due to the placement of
Carabus (*Hygrocarabus*) as sister to *Carabus* (*Limnocarabus*) and *Carabus* (*Euleptocarabus*)
(Imura *et al.* 2000). In the same study, *Carabus* (*Heterocarabus*) was sister to *Carabus*
(*Chaetocarabus*) and *Carabus* (*Platycarabus*). Using the same gene fragment, another study
inferred *Carabus* (*Platycarabus*) as sister to *Carabus* (*Chaetocarabus*) and *Carabus*
(*Heterocarabus*), within a largely unresolved *Carabus* clade (Su *et al.* 2003). A subsequent study
using two nuclear gene fragments recovered Arcifera, represented by *Carabus* (*Chaetocarabus*)
and *Carabus* (*Platycarabus*) as sister to the rest of the genus (=Eucarabi) (Sota & Ishikawa 2004).
More recently, Deuve *et al.* (2012) used ten loci to recover Arcifera as sister to the Eucarabi and
within Arcifera, they recovered *Carabus* (*Hygrocarabus*) as sister to *Carabus* (*Chaetocarabus*)
and *Carabus* (*Platycarabus*). Phylogenetic relationships among *Carabus* (*Platycarabus*) were also

investigated using Sanger sequencing data (Casale *et al.* 1998), suggesting that *C. irregularis* is
sister to the rest of the subgenus with *C. fabricii* and *C. depressus* being the most derived lineages
in the tree. In parallel to a moderate refinement in the phylogenetic inferences of Arcifera, the
estimation of divergence times in the clade has made some progress. Estimates for the origin of
Arcifera based on few loci range from the mid-Miocene (ca. 14 Ma, Deuve *et al.* 2012) to the early
Oligocene (ca. 30 Ma, Schmidt *et al.* 2023). No major improvement in our understanding of
Arcifera systematics and evolution has been achieved in the past decade and there is a need to infer
a robust evolutionary tree for this section of *Carabus* to better understand the morphological,
ecological and geographical evolution of constituent lineages.
In this study, we take advantage of the HyRAD-X approach to integrate a large number of
samples throughout the large geographical range of Arcifera. We rely on phylogenomic inferences,
species delimitations and population genomics approaches to clarify the taxonomy and elucidate
the evolutionary history of this complex group of species. In particular, we use this new genomic
framework to test which abiotic factors may have fostered the diversification of Arcifera through
space and time in the Cenozoic.
185
Material and methods
Taxon sampling and DNA extraction
The initial sampling was designed in order to sample major lineages within the Arcifera group
comprising four subgenera *Carabus* (*Chaetocarabus*), *Carabus* (*Heterocarabus*), *Carabus*
(*Hygrocarabus*) and *Carabus* (*Platycarabus*) (Deuve 2019). A total of 96 samples were initially
collected, mainly from NHC samples (87 samples, i.e. 90% of the dataset) but also from a few
fresh samples (9 samples, i.e. 10% of the dataset) when these were available (Supplementary Table
1). Multiple specimens of the same taxa and geographic populations were initially selected to
mitigate the risk of hDNA degradation that can result in specimens not being processed. NHC
specimens used in this study are kept at the Natural History Museum of Geneva (MHNG, 76
specimens) and Zoologische Staatssammlung München (ZSM-SNSB, 10 specimens). Eight
specimens collected in 96% ethanol were also used and have been deposited in the MHNG
collections. DNA was extracted destructively from a single leg using a QIAamp DNA Micro kit
(Qiagen, Hilden, Germany). Quantity and quality of the purified DNA were assessed with a
Fragment Analyzer. Based on DNA quality and concentrations, 38 specimens were not included
in the final samples selected for capture, enrichment and sequencing (*ca.* 40% of DNA extractions
not processed). Overall, a total of 56 Arcifera specimens were sequenced de novo for this study,
representing all Arcifera subgenera and species, several subspecies for the most widespread
species as well as a good geographical representation of each species range (taking into account
12 specimens that were eventually not included in the decisive datasets, see Results). Commonly,
early sampling erosion and discarded samples are not discussed in the framework of museomics
studies but we believe that this is critical to understand the limitations and cost of such approaches
in modern phylogenomic studies. The initial sampling in this study was specifically designed to
accommodate a *ca.* 40–50% specimen loss during DNA quality/quantity assessment (e.g.,

Toussaint *et al.* 2021), and therefore the resulting sampling is well-suited to tackle the focal
taxonomic and evolutionary questions in Arcifera. The final taxon sampling was complemented
by eight samples of *Carabus* (including one of *C. irregularis* and one of *C. variolosus*) and one of
Calosoma sycophanta (Linné, 1758) retrieved from (Toussaint *et al.* 2021) (see Supplementary
Table 1 for more details).
215
HyRAD-X protocol
The HyRAD protocol was applied as in (Toussaint *et al.* 2021). For fresh specimens a shearing
step with NEBNext dsDNA Fragmentase (New England Biolabs) was performed before library
preparation. Shotgun libraries were prepared based on the protocol developed in (Tin *et al.* 2014).
Purified DNA was phosphorylated with T4 Polynucleotide Kinase. After heat-denaturation into
single-stranded DNA, G-tailing was performed with Terminal Transferase and second strand DNA
was synthesized with Klenow Fragment (3′->5′exo-) using a poly-C oligonucleotide. Blunt-end
reaction was performed with T4 DNA Polymerase and barcoded adapters were ligated to the
phosphorylated end with T4 DNA ligase. After adapter fill-in with Bst DNA Polymerase (Large
Fragment), PCR were run using Phusion U Hot Start DNA Polymerase (Thermo Scientific) and
indexed PCR primers. Libraries were pooled in equimolar quantities based upon their respective
concentrations. Hybridization capture for enrichment of shotgun libraries was based on the
MYbaits protocol (Arbor Biosciences) modified as in (Toussaint *et al.* 2021) to include a two-step
capture at different temperatures (Li *et al.* 2013). Final library sequencing was performed on
Illumina NovaSeq 6000 SP using a paired-end protocol (Lausanne Genomic Technologies Facility,
Switzerland).
232
Illumina sequencing data cleanup and processing
Raw reads were demultiplexed according to indexes and barcodes using CutAdapt2 (Martin 2011).
Reads were cleaned using CutAdapt2 (Martin 2011) and quality was assessed all along the process
using fastqc (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Cleaned reads were
individually mapped onto the loci catalog using BWA-MEM (Li 2013) (Supplementary Figure 1).
The GATK (GenomeAnalysisTK) IndelRealigner tool (McKenna *et al.* 2010) realigned the indels
and deamination were corrected using *mapDamage2.0* (Jónsson *et al.* 2013). For each sample and
each locus, a consensus sequence was generated from the mapping file using samtools mpileup,
bcftools and *vcfutils.pl* (Li *et al.* 2009). Consensuses were generated keeping the majority allele at
each position. Twelve samples with too much missing data (more than 80% of N), were identified
using seqtk and removed (Supplementary Table 1). Two thresholds of minimum coverage
(min_cov) were applied to keep positions: min_cov=3 and min_cov=6. To test different levels of
missing data, decisive datasets were generated applying three thresholds for the minimum number
of samples per locus (min_sample): min_sample=10, min_sample=17 and min_sample=32. As a
result, six datasets were generated: Dataset A (min_cov=6, min_sample=10, 50 taxa, 1'481 loci),
Dataset B (min_cov=6, min_sample=10, 52 taxa, 1'965 loci), Dataset C (min_cov=6,
min_sample=17, 50 taxa, 1'014 loci), Dataset D (min_cov=3, min_sample=17, 52 taxa, 1'291

loci), Dataset E (min_cov=6, min_sample=32, 50 taxa, 366 loci) and Dataset F (min_cov=3,
min_sample=32, 52 taxa, 478 loci). The consensus sequences were combined and aligned with
MAFFT using the --auto option. Eight samples from (Toussaint *et al.* 2021) were integrated at the
alignment step. The final datasets only differ at the taxon sampling level with respect to *Carabus*
arcadicus merlini CBX0176 and *Carabus cychroides* CBX0082 that were included only in
Datasets B, D and F (these two taxa were systematically discarded because of low genomic
coverage when generating loci with a min_cov=6, i.e., in Datasets A, C and E).
MitoFinder (Allio *et al.* 2020) was used to identify mitochondrial genes among all
sequenced loci. The kept genes were shared by at least half of the samples. The genes were aligned,
and the sequences were cleaned in Geneious. Individual locus haplotype networks were built in
SplitsTree v.4.19.1 (Huson & Bryant 2006). The networks were reconstructed using calculated
uncorrected p-distances and the NeighborNet algorithm. All non-Arcifera outgroups were removed
before analyses. In parallel, a SNP calling was performed on the mapping files from the Arcifera
species using GATK (McKenna *et al.* 2010) in order to perform complementary population
genomic analyses and compare the results with those obtained from the locus oriented approach,
avoiding any bias linked to locus reconstruction (Dataset H).
266
Phylogenetic inferences
For each dataset, phylogenetic inferences were performed using IQ-TREE v2.0.5 (Minh *et al.*
2020) using the edge-linked partition model (Chernomor *et al.* 2016). First, the best partitioning
schemes were estimated using PartitionFinder v2.1.1 (Lanfear *et al.* 2017) with the rcluster
algorithm under the Akaike Information Criterion corrected (AICc), with a rcluster-max of 2,000
and a rcluster-percent of 20. The resulting partitioning schemes were then used in IQ-TREE to
select corresponding models of nucleotide substitution using ModelFinder (Kalyaanamoorthy *et*
al. 2017) and the AICc across all available models in IQ-TREE. To avoid local optima, we
performed 100 independent tree searches for each dataset in IQ-TREE. To estimate branch support,
we calculated 1,000 ultrafast bootstraps along with 1,000 SH-aLRT tests in IQ-TREE (Guindon *et*
al. 2010; Hoang *et al.* 2018). We used the hill-climbing nearest-neighbour interchange topology
search strategy to avoid severe model violations leading to biased ultrafast bootstrap estimations
(Hoang et al., 2018). The best tree for each analysis was selected based on the comparison of
maximum likelihood scores. Coalescent species trees were inferred using ASTRAL-hybrid (Zhang
& Mirarab 2022). We first performed individual locus trees using IQ-TREE v2.0.5 (Minh *et al.*
2020) and branch supports were assessed using 1,000 ultrafast bootstraps. Best substitution model
for each locus was estimated using ModelFinder (Kalyaanamoorthy *et al.* 2017). Species tree
reconstruction was performed combining gene trees using the weighted-ASTRAL optimization
algorithm (Zhang & Mirarab 2022) taking into account phylogenetic uncertainty by relying on
branch length and branch support across locus trees. As a complement to the locus reconstruction
approach, we performed phylogenetic inferences based on the SNP set used for the population
genomic approaches. Bi-allelic SNP shared by at least four samples were extracted and all
invariant sites removed. Species trees were inferred with RAxML-NG (Kozlov *et al.* 2019) using

GTR+G+ASC_LEWIS model for ascertainment bias correction and branch supports were assessed
using 1,000 bootstraps.

Divergence time estimation
Divergence time estimation was performed in BEAST 1.10.4 (Suchard *et al.* 2018) based on a
subset of loci selected using a gene-shopping approach to make these analyses tractable on a
bioinformatic cluster. SortaDate (Smith *et al.* 2018) was used with default settings to select 100
loci sorted using the following criteria: clock-likeness, tree length, and least topological conflict
with the IQ-TREE species tree on dataset E. The selected loci were then concatenated into a
Dataset G for relaxed-clock Bayesian divergence time estimation. The best partitioning scheme
and substitution models were determined with PartitionFinder2 (Lanfear *et al.* 2017) using the
greedy algorithm with the parameter minsubset-size = 2000 and the Bayesian information criterion
algorithm to choose between competing models. Clock partitioning was implemented by 1) a
single clock for all partitions and 2) a clock for each partition (eight in total; see Results). A
Bayesian lognormal relaxed clock model was assigned to the different clock partitions. Different
tree models were tested using a Yule pure birth model (Yule 1925; Gernhard 2008), a birth-death
model (Drummond *et al.* 2006; Gernhard 2008) as well as a Constant population size coalescent
model (Kingman 1982). Since the fossil record of *Carabus* is scarce, we relied on secondary
calibrations from a study focusing on Adephaga evolution based on 23 beetle fossil calibrations
(Baca *et al.* 2021). According to this study, the separation between the genera *Calosoma* and
Carabus occurred about 41.4 [37.1–46.1] million years ago (Ma). This age was used as a secondary
calibration for the corresponding node in our topology (split *Calosoma*/*Carabus*, in this case the
root). A second calibration was used to constrain the crown of *Carabus*. Following (Baca *et al.*
2021), this node was constrained to match the recovered age in their study at about 25.4 [22.8–
28.2] Ma. The analyses were conducted for 50 million generations, sampling parameters and trees
every 5000 generations. The maximum clade credibility tree for each analysis was generated in
TreeAnnotator 1.10.4.

Species delimitation and hybridisation
We used a combination of species delimitation methods and population genomic approaches to
test species and subspecies limits. For these analyses we excluded the six non-Arcifera outgroup
specimens resulting in a dataset composed of 44 samples. We extracted the 44 Arcifera samples
from Dataset E composed of 366 shared loci which present the lowest level of missingness (Table
1). First, BPP (Flouri *et al.* 2018) was used with the A11 option, using inverse–gamma
distributed diffuse priors (α = 3) for the population sizes (θ) and root ages ($\tau0$). Analysis was
run for 100,000 generations, sampling every 100 generations after a burnin of 8,000
generations. Second, the multi-locus species delimitation using Bayesian model comparison
implemented in the TR2 package (Fujisawa *et al.* 2016) has been applied on the same dataset.
Locus trees generated with IQ-TREE v2.0.5 (Minh *et al.* 2020) and previously used for the

weighted-ASTRAL approach were used as well as the maximum likelihood IQ-TREE consensus
tree on dataset A as guide tree. Outgroups were removed from gene trees and the guide tree.
From the SNP (Dataset H), population clustering was assessed using STRUCTURE 2.3.3
(Pritchard *et al.* 2000). Bi-allelic SNPs shared by at least 40% of the samples were extracted using
VCFtools v0.1.12a (Danecek *et al.* 2011). Because markers are supposed to be unlinked, we
extracted randomly only one SNP by locus. K-values from 1 to 15 were tested with no prior
population information and performed three times for each of them to verify a convergence of
estimations. A burn-in of 100,000 runs was used followed by 500,000 iterations. The most likely
number of clusters was determined using the Evanno method (Evanno *et al.* 2005) implemented
in Structure Harvester (Earl & vonHoldt 2012). The replicates were then combined and the figures
generated using CLUMPAK server (Kopelman *et al.* 2015). To investigate putative admixture
between species or subspecies we estimated Patterson's D statistic (ABBA-BABA test) (Patterson
et al. 2012) for all subspecies/species quartets using the Dsuite (Malinsky *et al.* 2021). The
analyses were performed on bi-allelic SNPs shared by at least 40% of the samples composed of
6,743 SNPs. Z-scores and associated p-values were calculated to assess the significance of the
results.
345
Results
Museomic approach efficiency
The combination of historical and fresh samples enabled to compare the effectiveness of
museomics methods. The DNA concentrations obtained from a single leg are very variable
between fresh samples (mean = 8.37 ng/µL; sd = 6.82 ng/µL) and NHC samples (mean = 1.18
ng/µL; sd = 2.45 ng/µL). There was a significant correlation between the quantity of DNA
extracted and the age of the specimens (Figure 1A). For the NHC samples, this concentration was
not homogeneous, with some samples nevertheless showing a high concentration. Forty samples
with a concentration below the detection thresholds, were excluded from the rest of the capture
process. It should be noted that some samples with very low DNA concentrations, such as *Carabus*
fabricii CBX0094 captured in 1977 with a concentration of only 0.08 ng/µL, were reliably placed
into the final phylogenetic inferences. For specimens with measurable DNA, the capture process
worked efficiently, allowing the sequencing of an average of 8.4 million reads per sample (sd =
9.2 million). There was a large difference between the average number of reads obtained from
fresh samples (mean = 23.8 millions; sd = 11.0 millions) and NHC samples (mean = 5.4 millions;
sd = 5.0 millions). The age of the specimens also had an influence on the number of reads obtained,
as there was a significant correlation between the age of the specimens and the number of reads
obtained (Figure 1B).
After locus reconstruction, the difference between fresh and NHC samples persists, with an
average of 1765 loci recovered in fresh samples (sd = 553) and 629 in NHC samples (sd = 447)
(Figure 1C). This difference is of the same order when looking at shared loci (Figure 1D). There
is a large heterogeneity in the number of loci recovered between NHC samples, largely linked to
the age of the specimen. Samples with too few loci (< 150 loci), i.e. 12 samples, had to be excluded

from the final datasets. For 35 NHC samples, the number of loci recovered, on average 793 (sd =
400), was sufficient to include them in subsequent analyses. Although strict filtering steps reduced
the number of NHC samples, they also ensured the reliability of the dataset for downstream
inferences.
373
374   *Phylogenomic inferences*
375
We inferred the phylogenetic relationships among Arcifera using six different datasets under
various taxon sampling and/or gene sampling strategies (Figure 2). The results of analyses based
on a concatenation approach performed in IQ-TREE and on a coalescent species-tree approach
conducted in wASTRAL are consistent except for the placement of *Carabus marietti*, the
branching pattern within *C.* (*Chaetocarabus*) and relationships between *C. creutzeri* and *C.*
fabricii. The subgenus *C.* (*Hygrocarabus*) is recovered as monophyletic in all analyses (including
wASTRAL) and as sister to the rest of Arcifera, however all analyses failed to recover *C.*
nodulosus and *C. variolosus* as reciprocally monophyletic. In all IQ-TREE inferences except the
one based on Dataset E and all wASTRAL analyses except the one based on Datasets A and B, *C.*
marietti is recovered as sister to *C.* (*Chaetocarabus*) with heterogeneous levels of branch support.
In the IQ-TREE analysis of Dataset E, this taxon is recovered as sister to the genus *Carabus* as a
whole, whereas in wASTRAL analyses of Datasets A and B it is recovered as sister to Arcifera
except *C.* (*Hygrocarabus*) with low branch support. The subgenus *C.* (*Chaetocarabus*) is always
recovered as monophyletic but internal relationships differ between analyses. A minority of
analyses recovered *C. arcadicus* and *C. intricatus* as reciprocally monophyletic (for instance no
wASTRAL analysis recovered this relationship). The subspecies *C. intricatus lefebvrei* is
recovered as sister to the nominal subspecies in all analyses. When *Carabus arcadicus merlini*
is included (Datasets B, D and F only), it never groups with other specimens of the nominal
subspecies resulting in *Carabus arcadicus* being consistently inferred as paraphyletic when this
taxon is included (Supplementary Figure 2). The subgenus *C.* (*Platycarabus*) is recovered as
monophyletic and with identical interspecific relationships across all IQ-TREE analyses but some
contention in wASTRAL ones. The alpine endemic *C. cychroides* is recovered as sister to the rest
of the subgenus in all analyses with strong branch support (IQ-TREE and wASTRAL). The species
C. depressus is inferred as the next lineage branching off in *C.* (*Platycarabus*) across all IQ-TREE
analyses and most wASTRAL analyses (except in Dataset A and E where it is recovered as sister
to *C. irregularis* with low branch support). The subspecies *C. depressus lucens* is recovered as
sister to the nominal subspecies in all analyses. The placement of the three remaining *C.*
(*Platycarabus*) species is identical across all IQ-TREE analyses with strong branch support, with
C. creutzeri being sister to *C. fabricii* and *C. irregularis*. The wASTRAL analyzes recover
different relationships but with low branch support, with a weakly supported sister relationship
between *C. creutzeri* and *C. fabricii* in analyses of Datasets D, E and F. The subspecies *C. fabricii*
malachiticus is recovered as nested within the nominal subspecies in all analyses. The subspecies
C. irregularis montandoni is recovered as sister to *C. irregularis bucephalus* and *C. irregularis*

irregularis in all IQ-TREE analyses whereas it is *C. irregularis bucephalus* that is inferred as sister to *C. irregularis irregularis* and *C. irregularis montandoni* in all wASTRAL analyses. Overall the IQ-TREE and wASTRAL inferences are highly compatible when collapsing the weakly supported relationships in wASTRAL species trees (gray and red asterisks in Figure 2). In particular, inconsistent relationships in wASTRAL compared to IQ-TREE always received poor branch support. We observe that branch support and overall phylogenetic resolution appears positively correlated to gene and taxon sampling (i.e., including less taxa and less loci to improve matrix completeness likely results in a loss of resolution).

Divergence time estimation

The BEAST dating analysis revealed consistent results for the four main nodes, i.e. the root, *Carabus*, Arcifera and *Carabus* (*Platycarabus*) nodes, according to the three models tested Yule, Birth-Death model, and Constant population size coalescent (Figures 3 and 4). The coalescent model including eight Bayesian log-normal relaxed clocks received the best marginal likelihood as calculated using stepping-stone sampling in BEAST and was therefore selected hereafter. This inference suggests an origin of Arcifera at 26.07 Ma (95% HPD: 22.77 - 29.67 Ma) and 14.56 Ma (95% HPD: 12.52 - 16.76 Ma) for the *Carabus (Platycarabus)* subgenus.

Species delimitation and putative hybridization

The different approaches to species delimitation produced contrasting results. The analysis performed with BPP is the most consistent with morphology and the current classification. The two species *Carabus variolosus* and *C. nodulosus* are well separated even though *C. nodulosus* is not monophyletic in our phylogeny. TR2 approach proposes an oversplit of the three *C. nodulosus* samples. Conversely, the STRUCTURE approach groups the two species in a single cluster (Supplementary Figure 6). *Carabus marietti*, the only representative of *Carabus (Heterocarabus)* is delineated as a species in all three approaches. Within *Carabus (Chaetocarabus)*, the two species *C. arcadicus* and *C. intricatus* are delineated by BPP but are merged by TR2 and STRUCTURE, potentially for the same reasons as in *Carabus (Hygrocarabus)*. It should be noted that the two subspecies of *C. intricatus*, i.e. *C. intricatus lefebvrei* and *C. intricatus intricatus*, are never delineated as distinct species. The species *C. cychroides* was well discriminated in two of the three approaches, with only TR2 proposing an additional split of the most basal sample. The two subspecies of *C. depressus*, *C. depressus depressus* and *C. depressus lucens* are not grouped together in the BPP approach and are identified as two distinct species. The results of the three methods are fully consistent with the morphology for *C. creutzeri* and *C. fabricii*. For *C. irregularis*, the situation is similar for two of the three methods, i.e. BPP and Structure. Among the 85 trios analysed, high D-statistics values, > 0.25, with significant p-values were observed for three trios. For two of these, *C. cychroides* was observed in P1 and *C. arcadicus* in P3. Despite this, no f-branch signal significantly different from zero could be identified (Supplementary Figure

7). These results suggest an absence of past introgression between the different species and
subspecies.

Discussion

Using museomics to obtain an extensive dataset

The HyRAD and HyRAD-X methods are unique in that they allow in-house production of probes
using a ddRAD protocol, either directly on the DNA of a few fresh samples (Suchan *et al.* 2016)
or on their RNA (Schmid *et al.* 2017). These approaches allow to target several thousand loci and
generate in turn high-resolution phylogenomic inferences (Young & Gillung 2020). In this study,
we obtained 1,965 loci for the most extensive dataset. These loci were informative enough to
resolve both the deep relationships between outgroups and the more recent relationships at the
intrageneric and intraspecific scales. In addition, the identification of SNPs on these loci also
enabled population genomic approaches such as the study of genetic structure and admixture.
In addition, the HyRAD-X approach made it possible to integrate samples with extremely
low initial DNA quantities. However, out of 96 samples from which DNA was extracted, 40 had
an undetectable quantity of DNA. In the context of museomics projects, it is therefore instrumental
to plan for redundancy in the sampling, with several samples per targeted taxon, in order to
compensate for any failures. Furthermore, the ability to generate genetic information from hDNA
is not entirely predictable. The recovery of meaningful genomic data does not seem to be linked
to the age of samples (Figure 1), in line with existing observations (Toussaint *et al.* 2021; Nunes
et al. 2022). In that vein, large amounts of genomic data could be obtained from older NHC
samples when almost none could be obtained from more recent samples. The quality and quantity
of DNA that can be extracted from NHC specimens is linked to factors that we cannot control,
such as the conditions of collection and preservation process (Post *et al.* 1993; Dillon *et al.* 1996;
Ruppert *et al.* 2023).

Systematics and species delimitation in Arcifera

Our results provide a robust phylogenomic tree of Arcifera for the first time (Figure 2). Overall
we support the view that Arcifera represents a monophylum within which all four subgenera form
clades. The monophyly of Arcifera is also supported by the presence of a hook-shaped ligulum
(i.e., arculus) at the base of the endophallus, a strong morphological character that unites all
constituents of this lineage (Imura *et al.* 2000; Deuve *et al.* 2012). Our study is the first to provide
strong evidence for these relationships while including all species of the group. Other studies based
on reduced genomic sampling, often a single gene fragment, either failed to recover Arcifera as
monophyletic (Imura *et al.* 2000; Osawa *et al.* 2004), or had too limited a taxon sampling to
properly test the placement and otherwise monophyly of each subgenus (Su *et al.* 2003; Sota &
Ishikawa 2004; Deuve *et al.* 2012). Except for a minority of analyses, our results strongly suggest

that *Carabus* (*Hygrocarabus*) is sister to the rest of Arcifera, with *Carabus* (*Platycarabus*) as sister
to a clade formed by *Carabus* (*Chaetocarabus*) and *Carabus* (*Heterocarabus*).

Within *Carabus* (*Hygrocarabus*), we recover *C. variolosus* nested within *C. nodulosus*.
This result contrasts with the ones of Mossakowski *et al.* (2020) where the two species were
suggested to be well differentiated genetically. In their study, these authors argued based on the
analysis of two gene fragments that both taxa form distinct clades although several specimens
caused each species to be paraphyletic. Some tests of mating between the two candidate species
were also performed in this study and suggested that the two lineages do not mate. However, the
scale and conditions of these trials do not allow to conclusively rule out potential mating. We argue
that in the current state of our knowledge it is not yet possible to definitively test species
boundaries, past introgression and signature of hybridization between *Carabus nodulosus* and *C.*
variolosus. A desired approach would be to combine a large geographical sampling as in
Mossakowski *et al.* (2020) with a genomic scale dataset as developed in the present study to revisit
the systematic conundrum within this subgenus at the population level.

Within *Carabus* (*Chaetocarabus*), we recover *C. arcadicus* as sister to *C. intricatus* in most
analyses (Figure 2). These two species are allopatric, morphologically well-differentiated and little
doubt exists with respect to their status as distinct species. Surprisingly our species delimitation
analyses only partly support the two species hypothesis, with TR2 and STRUCTURE considering
that *Carabus* (*Chaetocarabus*) is a unique species. Considering the low genomic coverage of some
taxa included in the analyses (see below), the clear morphological and geographical split between
these lineages and the support from BPP analyses, we argue that the validity of these two species
is uncontroversial. Natural hybrids with an intermediate morphology and usually green dorsal
pattern are known to exist along the limits of their respective ranges in northern Greece (i.e., at the
Katara pass) where *Carabus intricatus macedonicus* (not sampled here) and *C. arcadicus*
arcadicus co-occur. Both *Carabus arcadicus* and *C. intricatus* also comprise geographically
restricted subspecies in Greece that have been considered valid species by some authors. In the
south of Greece, the melanistic subspecies *Carabus arcadicus merlini* is endemic to the
Peloponnese peninsula and allopatric from the nominal subspecies present in the north. One
specimen of this taxon was sequenced but genomic coverage was low and therefore it was only
included in the less stringent Datasets B, D and F. In the phylogenetic analyses of these datasets,
the inclusion of *Carabus arcadicus merlini* systematically results in all three *C. arcadicus*
specimens forming a phylogenetic grade within which *C. intricatus* is nested. We argue that this
is an artifact possibly caused by missing genomic sampling and that both species are reciprocally
monophyletic as recovered in all other analyses and as suggested by morphology. However, it is
possible that *Carabus arcadicus merlini* represents a distinct evolutionary lineage since it is always
recovered as sister to the rest of *C. (Chaetocarabus)*. Additional taxon sampling is needed to test
the placement of this morphologically distinct taxon within the subgenus. Across its range,
Carabus intricatus is represented by the nominal subspecies from western France and UK to
northern Greece. In the south of Italy and Sicily, this species is represented by the allopatric
Carabus intricatus lefebvrei. The status of this taxon is debated and some authors consider it a

valid species. In our results, we recover this subspecies as sister to the nominal subspecies
represented by specimens from France and Piemonte. Our phylogenetic inferences support the
view of *Carabus intricatus lefebvrei* as a possible distinct species but our species delimitation
analyses reject this hypothesis. To properly test species boundaries within *Carabus intricatus*,
additional taxon sampling is needed including a much denser geographical sampling of the
nominal subspecies along with all described valid subspecies (Deuve 2019). In the Balkans, several
subspecies of *Carabus intricatus* have been described and represented more or less isolated
populations restricted to northern Greece. Despite our efforts we could not obtain DNA of good
quality for *Carabus intricatus krueperi* endemic to eastern Thessaly and considered by some
authors to be a valid species. Here as well, a denser taxon sampling is needed to properly test
species boundaries in this group. The placement of *Carabus* (*Heterocarabus*) *marietti* as sister to
C. (*Chaetocarabus*) receives support from most analyses in this study. Despite a relatively
circumscribed geographic range in northern Turkey and southern Bulgaria, numerous taxa have
been described in this subgenus even though currently a single species is considered valid (Turin
et al. 2003; Deuve 2019). Increasing the taxon sampling for this group by covering all its
geographical range would allow testing the match between morphological and genetic diversity
and better understand the evolution of this unique lineage at the inter- and intraspecific interface.
Within *Carabus* (*Platycarabus*), we recover *C. cychroides* as sister to the rest of the
subgenus. This result is unexpected because this species is a very narrowly restricted endemic to
Piemonte mountain ranges where it lives in alpine meadows and scree >2000m. The species was
only included once in a phylogenetic framework by Casale *et al.* (1998) who recovered it as a
derived lineage close to *Carabus depressus* and *C. fabricii*. Interestingly, a sister relationship of
this species to the rest of *Carabus* (*Platycarabus*) was suggested by the analysis of morphological
characters in Casale *et al.* (1998). Indeed, this species is morphologically quite different from the
rest of the subgenus in that it is one of the most extreme examples of cychrization in *Carabus*, a
process by which the pronotum is narrowed to allow predation inside snail shells (= stenocephalic
morphology). All species of the subgenus present a stenocephalic morphology, although less
marked than in *Carabus cychroides*, except for *C. irregularis* which is macrocephalic. Our
phylogenetic inferences are therefore important to understand the evolution of predation strategies
and associated morphology across the genus *Carabus* in which both types of morphologies exist
(Sota & Ishikawa 2004). Most malacophagous and helicophagous species in *Carabus* are
macrocephalic and use their enlarged pronotum, head and robust mandibles to break snail shells.
Cases of stenocephaly are most notably observed in *Carabus* (*Platycarabus*) but also in *C.*
(*Damaster*) Kollar, 1836 and *C.* (*Macrothorax*) Desmarest, 1850. The fact that *Carabus*
irregularis, the only *C.* (*Platycarabus*) macrocephalic species, is recovered as the most derived
species in the subgenus, indicates that macrocephaly possibly evolved from a stenocephalic
morphology unlike what was suggested in Casale *et al.* (1998). In the case of *Carabus cychroides*,
it is not closely related to any other species of the subgenus as suggested by previous authors, and
despite rare known natural hybrids with *C. depressus* in the Cottian Alps (i.e., Colle delle Finestre,
Monte Morefreddo, Monte Albergian), these species do not share an immediate recent common

ancestry (Sturani 1962; Casale *et al.* 1998; Anselmo & Rizzioli 2022a; b). The rest of *Carabus*
(*Platycarabus*) species and most sampled subspecies are found monophyletic (Figure 2). We
recover the subspecies *Carabus depressus lucens* as sister to the nominal subspecies in all analyses
and with robust branch support. This subspecies is morphologically quite divergent from the
nominal subspecies and *C. depressus bonellii* as it completely lacks elytral foveoli. It is also
allopatric from the rest of the *C. depressus* populations, being found in a small translapine region
between France and Italy (i.e., French Queyras to Italian Alpi Marittime), and its status as a valid
species even though rejected by three out of four species delimitation analyses should be revisited
with enhanced population sampling. Our taxon sampling within *Carabus creutzeri* does not allow
testing subspecies monophyly and relationships in detail but species delimitation analyses
unambiguously support a single species (Figure 3). Within *Carabus fabricii*, we recover the
Carpathian populations of *C. fabricii* (ssp *fassati* = nominal ssp, and spp *malachiticus*) nested
within Alpine populations of the nominal subspecies. This is unexpected to some extent as
Carabus fabricii presents a disjunct distribution between the Alps and the Carpathians (i.e., it is
not currently found in the Danube valley). Our results suggest that despite an allopatric range, gene
flow has been maintained between all populations of this species, however branch supports for
internal relationships in *Carabus fabricii* are moderate and an enhanced taxon sampling is needed
to understand the past and present connectivity between populations. All species delimitation
analyses support a unique species. One of the most interesting subspecific cases is recovered in
Carabus irregularis. This species is the most widespread of the subgenus ranging from eastern
France to Romania and Ukraine. It comprises three valid subspecies, one of which *Carabus*
irregularis montandoni from the Carpathians, was suggested to be a valid species based on
molecular evidence (Homburg *et al.* 2013). Our results support to some extent this view with *C.*
irregularis montandoni being found sister to the rest of populations in all IQ-TREE analyses but
not in wASTRAL analyses where the other subspecies *C. irregularis bucephalus* is found as sister
to the rest of the clade. There seems to be a genetic differentiation between the three recognized
subspecies of *C. irregularis* but our species delimitation analyses support the view of a single
species.
597
598 *Evolution of the Arcifera group*
599
The divergence time estimation analyses all recover an origin of Arcifera ca. 26 Ma in the
Oligocene. We did not perform a biogeographic estimation of ancestral ranges in the group because
several species are very widespread and initial attempts resulted in unresolved patterns. The fact
that species boundaries within *Carabus* (*Hygrocarabus*) are unstable also prevented a proper
reconstruction. However, it is possible to discuss several phylogenetic splits in the framework of
our results. The stem branch connecting *Carabus* (*Hygrocarabus*) to the rest of Arcifera is long,
potentially representing periods of extinction in this lineage. Currently the two recognized species
in the subgenus occur in temperate forests where adults live and hunt near and in good quality
streams. The reconfiguration of such habitats in the past 25 million years due to climatic

oscillations (Westerhold *et al.* 2020) may have extirpated populations and pushed others in their
current ranges. Considering the specificity of these two lineages to their habitat, and predictions
of global warming and their impact on such ecosystems (Capon *et al.* 2021; Bonacina *et al.* 2023),
it is likely that they may be increasingly threatened in the future.

With respect to biogeography, one of the most interesting lineages in Arcifera is the clade
composed of *Carabus* (*Chaetocarabus*) and *Carabus* (*Heterocarabus*). Because *Carabus*
(*Heterocarabus*) *marietti* is restricted to eastern Bulgaria and western Turkey, and *Carabus*
(*Chaetocarabus*) distributed in Greece (*C. arcadicus* is endemic to Greece), it is likely that the
ancestors of this clade originated in the geologically highly complex Aegean area. The split
between the two subgenera ca. 17 Ma predates the timing of the opening of the Aegean sea in the
Tortonian ca. 8 Ma (i.e., opening of the Mid-Aegean Trench or Aegean barrier; van Hinsbergen &
Schmid 2012), rejecting the hypothesis of geographic vicariance in the south as suggested in other
lineages (Poulakakis *et al.* 2015). Interestingly, both subgenera have very marginally overlapping
distributions in the Thrace basin with *Carabus* (*Heterocarabus*) currently distributed on the
southern Black Sea coast where *Carabus intricatus* is also represented by the subspecies *C.*
intricatus starensis (Gueorguiev & Gueorguiev 1995). At the time of divergence in the early
Miocene (i.e., Burdigalian), the Thrace basin formed a connection between the eastern Balkan
peninsula and Anatolia (Rögl 1997, 1999; Sachsenhofer *et al.* 2017; Erbil *et al.* 2021). It is possible
that ancestral populations dispersed in the Balkan Peninsula and/or in Anatolia where they evolved
independently. Under this scenario, the close geographic ranges of these two species would likely
represent secondary contact associated with more recent colonization of the Thrace basin. A more
robust population-level taxon sampling, especially of *Carabus* (*Heterocarabus*), might elucidate
the fine-scale biogeographic history of this clade in the future. Within *Carabus* (*Chaetocarabus*),
the two currently recognized species are mostly allopatric with only a short overlap in western
Greece (e.g. Katara pass). There is no clear geological barrier that may have fostered vicariant
diversification at the time of speciation ca. 7 Ma. Further diversification appears to be occurring
at the population level with *Carabus intricatus lefebvrei* endemic of southern Italy and allopatric
from the nominal subspecies. Similarly, *Carabus arcadicus merlini* endemic to Peloponnese is
morphologically quite divergent from the nominal subspecies and might represent a case of
ongoing speciation. The wide dispersal of *Carabus intricatus* across the western Palearctic region
is likely recent and may be explained by the generalist habitat preference of this species. Additional
geographical and taxon sampling will likely yield more robust inferences of evolutionary patterns
and processes within this clade in the future.

The evolutionary history of the subgenus *Carabus* (*Platycarabus*) is also revealed by our
analyses. We recover the narrowly endemic *Carabus cychroides* as sister to the rest of the
subgenus. This is surprising as it was not suggested by the molecular inference of Casale *et al.*
(1998). This placement has strong implications for our understanding of alpine biogeography in
this group. Only *Carabus irregularis* has lowland populations and its derived placement in the
phylogeny indicates that alpine specialization was likely ancestral in the subgenus with recent shift
in that species to lower habitats. This phylogenomic pattern and the origin of the subgenus *ca.* 15

Ma during the warmest period of the Neogene seems to indicate that ancestors of *Carabus*
(*Platycarabus*) may have been less specialized than nowadays and were distributed in mountain
regions. In the mid Miocene, mountain ranges across the Alps had the same elevation as nowadays
(Campani *et al.* 2012; Krsnik *et al.* 2021), however ecosystems were different due to significantly
warmer climatic conditions. When the climate progressively turned colder these beetles adapted
to ensuing conditions and became alpine specialists. It is possible that species of the subgenus
diverged due to competition, niche filling and/or host specialization as observed in *Carabus*
cychroides for instance. We hypothesize that in the latest sequence of their evolutionary history,
Pleistocene glaciations played a limited role in speciation since all current species had already
diverged (Figure 3).

Although natural hybrids are known between different species of the subgenus, our results
recover no hybridization signal between them. The most significant case concerns the species *C.*
fabricii and *C. irregularis*, whose ranges largely overlap in Switzerland, Austria and Slovakia. It
is in these sympatric areas that several cases of natural hybridisation have been identified (e.g. at
the Radstädter Tauern Pass in Austria, Mandl 1960). However, our genetic results do not show
any hybridisation signals between the species, either on genetic structure, where the two clusters
are well separated, or in the approach using Dsuite, which seeks to trace admixture signals in the
lineages. These results suggest that these sporadic hybridization events are not conserved in
populations and could imply a potential infertility of F1s (Casale *et al.* 1998). Furthermore, the
networks obtained with the three mitochondrial genes (Supplementary Figure 4) group the samples
of the *C. irregularis* and *C. fabricii* species in the same cluster. These mitonuclear discordance
patterns are frequent in the literature and can be explained by the specific biological properties of
mitochondrial DNA (uniparental inheritance and reduced recombination; Birky 2001) or
differences in the evolutionary histories of nuclear and mitochondrial markers including
incomplete lineage sorting and gene flow among species (Sota & Vogler 2001; Suchan *et al.* 2017).
The results obtained with the nuclear loci are sufficiently robust to be able to consider that the
hybridisations observed are either localised or do not induce lasting admixture between the species.
A more detailed analysis of hybrids, local populations and the implications of hybridisation on the
fitness of individuals could provide a better understanding of the mechanisms involved.

Integrating current species distribution, genetic isolation of these alpine species was
already in place when glaciation cycles struck the Alps. As a result, dispersal of populations in
peripheral glacial refugia as observed in *C. irregularis* (Homburg *et al.* 2013) did not result in
genetic homogenization despite species being placed in secondary contact. It is also possible in
the case of the more alpine-adapted species (all but *C. irregularis*) that dispersal occurred in
nunataks rather than peripheral glacial refugia (Holderegger & Thiel-Egenter 2009; Schönswetter
& Schneeweiss 2019; Kosiński *et al.* 2019), which would have resulted in an increased genetic
differentiation among populations as suggested by our analyses. Coupling a more extensive
geographic sampling of these five alpine species with niche modeling analyses may help testing
more specifically the different scenarios that governed range and genetic evolution of these
populations during Pleistocene glaciations.

Acknowledgements

We warmly thank Michael Balke for the loan of material from the ZSM-SNSB. We thank Elsa Ricossa for the digitization of specimens housed at the Natural History Museum of Geneva. We thank Céline Rochet for assistance in fieldwork. We thank the city of Geneva for an internal student grant awarded to MTP. We thank Conrad Gillett for allowing the use of his photographs and drawings in this article. We thank Ivan Rapuzzi for fruitful discussions and feedback on taxonomic aspects of this work.

Funding

This study was partly funded by a Master student grant awarded by the City of Geneva. EFAT is funded by a FNS grant 310030_200491.

Conflict of interest disclosure

The authors declare no conflict of interest.

Data, script, code, and supplementary information availability
Raw reads are available on the NCBI BioProject PRJNA1086379. The data underlying this article (final alignments and trees) and bioinformatic scripts are available on Github repository (https://github.com/JeremyLGauthier/Arcifera_phylogeny).

Figure captions
Table 1. Alignment statistics for each dataset, including the number of taxa, the number of loci, the minimum coverage, the minimum number of taxa, the alignment length, the percentage of missing data, the numbers and percentages of variable sites and of parsimony informative sites, and the GC content.

Figure 1. Statistical summary of locus recovery. Plots representing the relationship between the collection year and DNA concentration (A), number of sequenced reads (B), number of loci recovered for each sample (C), and number of shared loci in final dataset B. In each plot, ethanol-preserved samples are shown in green, samples from museums with an age < 30 years in blue and samples from museums with an age > 30 years in classic yellow. Correlations were tested with Spearman's correlation tests and adjusted coefficients of determination R-squared were estimated using a linear model.

Figure 2. Summary of phylogenetic inferences across Arcifera based on HyRAD-X data. The presented topology is derived from a maximum likelihood analysis performed in IQ-TREE using Dataset A. Branch support from this analysis is shown for all branches. Branch support retrieved in different analyses is shown for major branches according to the inserted caption. Sample type is indicated according to the inserted caption. Abbreviations at the end of each taxon label correspond to the following countries: AT, Austria, CH, Switzerland, FR, France, GR, Greece, HU, Hungary, IT, Italy, KP, Carpathians (Slovakia to Romania), RO, Romania, SI, Slovenia, SK, Slovakia, TK, Turkey. An illustration of a male *Carabus (Platycarabus) cychroides* is presented (Drawing: Conrad Gillett).

Figure 3. Bayesian divergence time estimates for the subgenus *Carabus (Platycarabus)* and Arcifera group. Maximum clade credibility tree obtained from a BEAST analysis using eight Bayesian log-normal relaxed clocks and a Coalescent Constant Size tree model. Node estimates are postburn in median ages, with 95% credibility intervals. Histogram represents the number of loci recovered for each sample and sample type are indicated according to the inserted caption. The section on the right shows the results of species delimitations identified using the different methods indicated above. The shades of gray represent the concordance between the different approaches with black being a total consensus. Habitus of three representative species (1) *Carabus nodulosus nodulosus* (credit : Conrad Gillett), (2) *Carabus intricatus intricatus* (credit : Conrad Gillett) and (3) *Carabus irregularis irregularis* (credit : Conrad Gillett) are shown.

Figure 4. Comparison of divergence time estimation between competing tree models and relaxed-clock partitioning strategies. Box-plots indicate for each analysis (color-coding inserted as a caption on the right side of the figure) the median age of the focal node (see X axis) and associated 95% age credibility interval. BD, birth-death model; CS, constant population size coalescent model.

Supplementary Table 1. Descriptive statistics for each included and non-included sample, including historical sample data, molecular biology information (DNA concentrations), sequencing and loci reconstruction statistics.

Supplementary Figure 1. Schematic representation of the bioinformatic pipeline.

Supplementary Figure 2. Maximum likelihood trees for each dataset: Dataset A (min_cov=6, min_sample=10, 50 taxa, 1'481 loci), Dataset B (min_cov=6, min_sample=10, 52 taxa, 1'965 loci), Dataset C (min_cov=6, min_sample=17, 50 taxa, 1'014 loci), Dataset D (min_cov=3, min_sample=17, 52 taxa, 1'291 loci), Dataset E (min_cov=6, min_sample=32, 50 taxa, 366 loci) and Dataset F (min_cov=3, min_sample=32, 52 taxa, 478 loci). Node supports indicate SH-aLRT and UFBoot values.

Supplementary Figure 3. Species trees obtained with wASTRAL on each dataset: Dataset A (min_cov=6, min_sample=10, 50 taxa, 1'481 loci), Dataset B (min_cov=6, min_sample=10, 52 taxa, 1'965 loci), Dataset C (min_cov=6, min_sample=17, 50 taxa, 1'014 loci), Dataset D (min_cov=3, min_sample=17, 52 taxa, 1'291 loci), Dataset E (min_cov=6, min_sample=32, 50 taxa, 366 loci) and Dataset F (min_cov=3, min_sample=32, 52 taxa, 478 loci). Node supports indicate SH-aLRT and UFBoot values. Node supports indicate LPP values.

Supplementary Figure 4. Individual locus haplotype networks (A. CO1, B. CO3 and C. CYTB). Networks were generated in SplitsTree using calculated uncorrected p-distances and the NeighborNet algorithm. The colour coding for the different morphological groups is identical to the one used (photo credit : Marie Pauli).

Supplementary Figure 5. Structure plots estimated on unlinked shared SNPs for K=1 to K = 15. For each K, the Mean(LnProb) is indicated.

Supplementary Figure 6. F4-branch statistic plotted as a heatmap. The tree topology is plotted above, and on the left, every branch of the tree is displayed (including internal branches).

References

Allio R, Schomaker-Bastos A, Romiguier J *et al.* (2020) MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics. *Molecular ecology resources*.

Anselmo L, Rizzioli B (2022a) Side Threats: Further Possible Effects Of Warming On The High Alpine Narrow Endemic Carabus *Cychroides* (Coleoptera: Carabidae). *Nature Conservation Research. Заповедная наука*, 7, 88–94.

Anselmo L, Rizzioli B (2022b) The small range and the great threat: extinction risk assessment of the narrow endemism *Carabus cychroides* under climate change. *Journal of insect conservation*, 26, 17–27.

Baca SM, Gustafson GT, Alexander AM, Gough HM, Toussaint EFA (2021) Integrative phylogenomics reveals a Permian origin of Adephaga beetles. *Systematic entomology*, 46, 968–990.

Bekchiev R, Antov M, Boyadzhiev P *et al.* (2022) Carabus variolosus (Fabricius, 1787) (Coleoptera: Carabidae) in Bulgaria: rediscovered after 111 years. *Historia naturalis Bulgarica*.

Birky CW Jr (2001) The inheritance of genes in mitochondria and chloroplasts: laws, mechanisms,

and models. *Annual review of genetics*, 35, 125–148.

Blaimer BB, Lloyd MW, Guillory WX, Brady SG (2016) Sequence Capture and Phylogenetic
Utility of Genomic Ultraconserved Elements Obtained from Pinned Insect Specimens. *PloS*
one, 11, e0161531.

Bonacina L, Fasano F, Mezzanotte V, Fornaroli R (2023) Effects of water temperature on
freshwater macroinvertebrates: a systematic review. *Biological reviews of the Cambridge*
Philosophical Society, 98, 191–221.

Camard A, Leplat J (2004) *Hybrides du genre Carabus*. Magellanes, France.

Campani M, Mulch A, Kempf O, Schlunegger F, Mancktelow N (2012) Miocene paleotopography
of the Central Alps. *Earth and planetary science letters*, 337-338, 174–185.

Capon SJ, Stewart-Koster B, Bunn SE (2021) Future of Freshwater Ecosystems in a 1.5°C Warmer
World. *Frontiers of Environmental Science & Engineering in China*, 9.

Card DC, Shapiro B, Giribet G, Moritz C, Edwards SV (2021) Museum Genomics. *Annual review*
of genetics.

Casale A, Pruser F, Arndt E, Mossakowski D (1998) Phylogenetic relationships in the subgenus
Platycarabus MORAWITZ,1886 (Coleoptera: Carabidae: Carabini). *Mus. reg. Sci. nat.*
Torina, Atti, 1998, 429–448.

Casale A, Rapuzzi I (2015) Due nuovi ibridi fra specie del genere Carabus Linné, 1758, e nuova
località italiana dell'ibrido *C. (Platycarabus) depressus depressus* Bonelli, 1811 x *C.*
(Platycarabus) fabricii fabricii Panzer, 1812 (Coleoptera Carabidae). *Rivista piemontese di*
Storia naturale, 157–169.

Cavazzuti P, Ghiretti D (2020) *Carabus d'Italia* (NE Scientifiche, Ed,). Natura Edizione
Scientifiche, Bologna.

Chernomor O, von Haeseler A, Minh BQ (2016) Terrace Aware Data Structure for Phylogenomic
Inference from Supermatrices. *Systematic biology*, 65, 997–1008.

Danecek P, Auton A, Abecasis G *et al.* (2011) The variant call format and VCFtools.
Bioinformatics , 27, 2156–2158.

Deuve T (2019) Classification du genre Carabus L., 1758. Liste Blumenthal 2018-2019
(Coleoptera, Carabidae). *Coléoptères*, 25, 33–102.

Deuve T (2021) *Carabus of the World [Holarctic Region]* (Magellanes, Ed,).

Deuve T, Cruaud A, Genson G, Rasplus J-Y (2012) Molecular systematics and evolutionary
history of the genus Carabus (Col. Carabidae). *Molecular phylogenetics and evolution*, 65,
259–275.

Dillon N, Austin AD, Bartowsky E (1996) Comparison of preservation techniques for DNA
extraction from hymenopterous insects. *Insect molecular biology*, 5, 21–24.

de-Dios T, Fontsere C, Renom P *et al.* (2023) Whole-genomes from the extinct Xerces Blue
butterfly can help identify declining insect species. *eLife*, 12.

Drummond AJ, Ho SYW, Phillips MJ, Rambaut A (2006) Relaxed phylogenetics and dating with
confidence. *PLoS biology*, 4, e88.

Duchenne F, Thébault E, Michez D *et al.* (2020) Phenological shifts alter the seasonal structure of

pollinator assemblages in Europe. *Nature ecology & evolution*, 4, 115–121.

Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation genetics resources*, 4, 359–361.

Erbil Ü, Okay AI, Hakyemez A (2021) Late Oligocene—Early Miocene shortening in the Thrace Basin, northern Aegean. *International Journal of Earth Sciences*, 110, 1921–1936.

Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular ecology*, 14, 2611–2620.

Faircloth BC (2017) Identifying conserved genomic elements and designing universal bait sets to enrich them. *Methods in ecology and evolution / British Ecological Society*, 8, 1103–1112.

Flouri T, Jiao X, Rannala B, Yang Z (2018) Species Tree Inference with BPP Using Genomic Sequences and the Multispecies Coalescent. *Molecular biology and evolution*, 35, 2585–2593.

Fujisawa T, Aswad A, Barraclough TG (2016) A Rapid and Scalable Method for Multilocus Species Delimitation Using Bayesian Model Comparison and Rooted Triplets. *Systematic biology*, 65, 759–771.

Gauthier J, Borer M, Toussaint EFA *et al.* (2023) Museomics reveals evolutionary history of Oreina alpine leaf beetles (Coleoptera: Chrysomelidae). *Systematic entomology*, 48, 658–671.

Gauthier J, Pajkovic M, Neuenschwander S *et al.* (2020) Museomics identifies genetic erosion in two butterfly species across the 20th century in Finland. *Molecular ecology resources*, 20, 1191–1205.

Gernhard T (2008) The conditioned reconstructed process. *Journal of theoretical biology*, 253, 769–778.

Gueorguiev VB, Gueorguiev BV (1995) *Catalogue of the Ground-beetles of Bulgaria (Coleoptera; Carabidae)* (Pensoft Publishers, Ed,). Pensoft Publishers.

Guindon S, Dufayard J-F, Lefort V *et al.* (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic biology*, 59, 307–321.

Hieke F, Wrase DW (2008) Faunistik der laufkäfer bulgariens. (Coleoptera, Carabidae). *Mitteilungen aus dem museum fur naturkunde in Berlin. Deutsche entomologische zeitschrift*, 35, 1–171.

van Hinsbergen DJJ, Schmid SM (2012) Map view restoration of Aegean–West Anatolian accretion and extension since the Eocene. *Tectonics*, 31.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular biology and evolution*, 35, 518–522.

Holderegger R, Thiel-Egenter C (2009) A discussion of different types of glacial refugia used in mountain biogeography and phylogeography. *Journal of biogeography*, 36, 476–480.

Homburg K, Drees C, Gossner MM *et al.* (2013) Multiple glacial refugia of the low-dispersal ground beetle Carabus irregularis: molecular data support predictions of species distribution models. *PloS one*, 8, e61185.

Hristovski S, Mésàros G, Komnenov M, Cvetkovska-Gjorgjievska A (2023) Rediscovery of Carabus (Hygrocarabus) variolosus nodulosus Creutzer, 1799 on Shar Planina Mt. *Macedonian Journal of Ecology and Environment*.

Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary studies. *Molecular biology and evolution*, 23, 254–267.

Imura Y, Kim C-G, Su Z-H, Osawa S (1998) An Attempt at the Higher Classification of the Carabina (Coleoptera, Carabidae) Based on Morphology and Molecular Phylogeny, with Special Reference to Apotomopterus, Limnocarabus and Euleptocarabus. *Elytra, Tokyo*, 26, 17–35.

Imura Y, Su Z-H, Osawa S (2000) Phylogenetic Relationships in the Division Arciferi (Coleoptera, Carabidae). *Elytra, Tokyo*, 28, 235–239.

Ishikawa R (1984) Phylogeny and subgeneric classification of the genus Chaetocarabus THOMSON (Coleoptera, Carabidae). *Kontyu, Tokyo*, 94–109.

Jónsson H, Ginolhac A, Schubert M, Johnson PLF, Orlando L (2013) mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* , 29, 1682–1684.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature methods*, 14, 587–589.

Kingman JFC (1982) The coalescent. *Stochastic Processes and their Applications*, 13, 235–248.

Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I (2015) Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular ecology resources*, 15, 1179–1191.

Kosiński P, Sękiewicz K, Walas Ł, Boratyński A, Dering M (2019) Spatial genetic structure of the endemic alpine plant *Salix serpillifolia*: genetic swamping on nunataks due to secondary colonization? *Alpine Botany*, 129, 107–121.

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A (2019) RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* , 35, 4453–4455.

Krsnik E, Methner K, Campani M *et al.* (2021) Miocene high elevation in the Central Alps. *Solid Earth*, 12, 2615–2631.

Kulijer D (2019) New records and distribution of threatened *Carabus (variolosus) nodulosus* Creutzer, 1799 in Bosnia and Herzegovina (Coleoptera: Carabidae). *Acta entomologica slovenica (Ljubljana)*, 27.

Landry B, Bilat J, Hayden J *et al.* (2023) The identity of Argyrialacteella (Fabricius, 1794) (Lepidoptera, Pyraloidea, Crambinae), synonyms, and related species revealed by morphology and DNA capture in type specimens. *ZooKeys*, 1146, 1–42.

Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) PartitionFinder 2: New Methods for Selecting Partitioned Models of Evolution for Molecular and Morphological Phylogenetic Analyses. *Molecular biology and evolution*, 34, 772–773.

Lemmon AR, Emme SA, Lemmon EM (2012) Anchored hybrid enrichment for massively high-

throughput phylogenomics. *Systematic biology*, 61, 727–744.

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-bio.GN]*.

Li H, Handsaker B, Wysoker A *et al.* (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* , 25, 2078–2079.

Li C, Hofreiter M, Straube N, Corrigan S, Naylor GJP (2013) Capturing protein-coding genes across highly divergent species. *BioTechniques*, 54, 321–326.

Malinsky M, Matschiner M, Svardal H (2021) Dsuite - Fast D-statistics and related admixture evidence from VCF files. *Molecular ecology resources*, 21, 584–595.

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, 17, 10–12.

Matern A, Desender K, Drees C *et al.* (2009) Genetic diversity and population structure of the endangered insect species *Carabus variolosus* in its western distribution range: Implications for conservation. *Conservation genetics* , 10, 391–405.

Matern A, Drees C, Vogler AP, Assmann T (2010) Linking Genetics and Ecology: Reconstructing the History of Relict Populations of an Endangered Semi-Aquatic Beetle. In: *Relict Species*, pp. 253–265. Springer Berlin Heidelberg.

Mayer C, Dietz L, Call E *et al.* (2021) Adding leaves to the Lepidoptera tree: capturing hundreds of nuclear genes from old museum specimens. *Systematic entomology*, 46, 649–671.

McKenna A, Hanna M, Banks E *et al.* (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research*, 20, 1297–1303.

Minh BQ, Schmidt HA, Chernomor O *et al.* (2020) IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular biology and evolution*, 37, 1530–1534.

Mossakowski D, Bérces S, Hejda R *et al.* (2020) High molecular diversity in *Carabus (Hygrocarabus) variolosus* and *C. nodulosus*. *Acta zoologica Academiae Scientiarum Hungaricae* , 66, 147–168.

Müller-Kroehling, S. (2006): Ist der Gruben-Großlaufkäfer *Carabus (variolosus) nodulosus* ein Taxon des Anhanges II der FFH-Richtlinie in Deutschland? – Waldökologie online 3: 52-57 *ResearchGate*.

Müller-Kroehling, S. (2014): Remarks on the current situation of *Carabus variolosus nodulosus* relating to the interpretation of its Habitats Directive status, the 2013 report under that directive, and its threat level in Germany and Central Europe – Angewandte Carabidologie 10: 97-100 *ResearchGate*.

Nunes R, Storer C, Doleck T *et al.* (2022) Predictors of sequence capture in a large-scale anchored phylogenomics project. *Frontiers in Ecology and Evolution*, 10.

Osawa S, Su ·-R, Imura Y (2004) *Molecular Phylogeny and Evolution of Carabid Ground Beetles* (SJ Kk, Ed,).

Patterson N, Moorjani P, Luo Y *et al.* (2012) Ancient admixture in human history. *Genetics*, 192,

1065–1093.

Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double digest RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PloS one*, 7, e37135.

Post RJ, Flook PK, Millest AL (1993) Methods for the preservation of insects for DNA studies. *Biochemical systematics and ecology*, 21, 85–92.

Poulakakis N, Kapli P, Lymberakis P *et al.* (2015) A review of phylogeographic analyses of animal taxa from the Aegean and surrounding regions. *Journal of zoological systematics and evolutionary research = Zeitschrift fur zoologische Systematik und Evolutionsforschung*, 53, 18–32.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics*, 155, 945–959.

Raxworthy CJ, Smith BT (2021) Mining museums for historical DNA: advances and challenges in museomics. *Trends in ecology & evolution*.

Rögl F (1997) Palaeogeographic Considerations for Mediterranean and Paratethys Seaways (Oligocene to Miocene). *Annalen des Naturhistorischen Museums in Wien. Serie A. Mineralogie und Petrographie, Geologie und Palaeontologie, Anthropologie und Praehistorie*, 99, 279–310.

Rögl F (1999) Mediterranean and Paratethys; facts and hypotheses of an Oligocene to Miocene paleogeography; short overview. *Geologica Carpathica (Bratislava)*, 339–349.

Ruppert L-S, Segelbacher G, Staab M, Winiger N (2023) Gauging DNA degradation among common insect trap preservatives. *Entomologia experimentalis et applicata*.

Sachsenhofer RF, Popov SV, Bechtel A *et al.* (2017) Oligocene and Lower Miocene source rocks in the Paratethys: palaeogeographical and stratigraphic controls. *Geological Society*, 464, 267–306.

Schmid S, Genevest R, Gobet E *et al.* (2017) HyRAD-X, a versatile method combining exome capture and RAD sequencing to extract genomic information from ancient DNA. *Methods in ecology and evolution / British Ecological Society*, 8, 1374–1388.

Schmidt J, Opgenoorth L, Mao K, Baniya CB, Hofmann S (2023) Molecular phylogeny of mega-diverse *Carabus* attests late Miocene evolution of alpine environments in the Himalayan–Tibetan Orogen. *Scientific reports*, 13, 1–16.

Schönswetter P, Schneeweiss GM (2019) Is the incidence of survival in interior Pleistocene refugia (nunataks) underestimated? Phylogeography of the high mountain plant *Androsace alpina* (Primulaceae) in the European Alps revisited. *Ecology and evolution*, 9, 4078–4086.

Smith SA, Brown JW, Walker JF (2018) So many genes, so little time: A practical approach to divergence-time estimation in the genomic era. *PloS one*, 13, e0197433.

Sota T, Ishikawa R (2004) Phylogeny and life-history evolution in *Carabus* (subtribe Carabina: Coleoptera, Carabidae) based on sequences of two nuclear genes. *Biological journal of the Linnean Society. Linnean Society of London*, 81, 135–149.

Sota T, Takami Y, Ikeda H *et al.* (2022) Global dispersal and diversification in ground beetles of

the subfamily Carabinae. *Molecular phylogenetics and evolution*, 167, 107355.

Sota T, Vogler AP (2001) Incongruence of mitochondrial and nuclear gene trees in the Carabid
beetles Ohomopterus. *Systematic biology*, 50, 39–59.

Sturani M (1962) *Osservazioni e ricerche biologiche sul genere Carabus Linnaeus (sensu lato)*
(Coleoptera Carabidae) (F Pagano, Ed,).

Suchan T, Espíndola A, Rutschmann S *et al.* (2017) Assessing the potential of RAD-sequencing
to resolve phylogenetic relationships within species radiations: The fly genus *Chiastocheta*
(Diptera: Anthomyiidae) as a case study. *Molecular phylogenetics and evolution*, 114, 189–
198.

Suchan T, Pitteloud C, Gerasimova NS *et al.* (2016) Hybridization Capture Using RAD Probes
(hyRAD), a New Tool for Performing Genomic Analyses on Collection Specimens. *PloS one*,
11, e0151651.

Suchard MA, Lemey P, Baele G *et al.* (2018) Bayesian phylogenetic and phylodynamic data
integration using BEAST 1.10. *Virus evolution*, 4, vey016.

Su Z-H, Imura Y, Zhou H-Z, Okamoto M, Osawa S (2003) Mode of morphological differentiation
in the Latitarsi-ground beetles (Coleoptera, Carabidae) of the world inferred from a
phylogenetic tree of mitochondrial ND5 gene sequences. *Genes & genetic systems*, 78, 53–
70.

Tin MM-Y, Economo EP, Mikheyev AS (2014) Sequencing degraded DNA from non-
destructively sampled museum specimens for RAD-tagging and low-coverage shotgun
phylogenetics. *PloS one*, 9, e96793.

Toussaint Fls EFA, Gillett CPDT (2018) Rekindling Jeannel's Gondwanan vision? Phylogenetics
and evolution of Carabinae with a focus on Calosoma caterpillar hunter beetles. *Biological*
journal of the Linnean Society. Linnean Society of London, 123, 191–207.

Toussaint EFA, Gauthier J, Bilat J *et al.* (2021) HyRAD-X Exome Capture Museomics Unravels
Giant Ground Beetle Evolution. *Genome biology and evolution*, 13.

Turin H, Penev L, Casale A (2003) *The Genus Carabus in Europe: A Synthesis*. Sofia/Leiden.

Tyszecka K, Zając K, Kadej M (2023) Habitat preferences of the mountain population of the
endangered beetle *Carabus variolosus ssp. variolosus* indicate its vulnerability to climate
change. *Global Ecology and Conservation*, 46, e02601.

Westerhold T, Marwan N, Drury AJ *et al.* (2020) An astronomically dated record of Earth's
climate and its predictability over the last 66 million years. *Science*, 369, 1383–1387.

Young AD, Gillung JP (2020) Phylogenomics — principles, opportunities and pitfalls of big-data
phylogenetics. *Systematic entomology*, 45, 225–247.

Yule GU (1925) II.—A mathematical theory of evolution, based on the conclusions of Dr. J. C.
Willis, F. R. S. *Philosophical Transactions of the Royal Society of London. Series B,*
Containing Papers of a Biological Character, 213, 21–87.

Zhang C, Mirarab S (2022) Weighting by Gene Tree Uncertainty Improves Accuracy of Quartet-
based Species Trees. *Molecular biology and evolution*, 39, msac215.

1047

Table 1

Dataset	No. of taxa	No. of loci	Min. cov	Min. taxa	Align. length	Missing data	Variable sites	Parsimony informative sites	GC_content
A	50	1,481	6	10	346,687	66.7%	40,743 (11.8%)	19,061 (5.5%)	0.469
B	52	1,965	3	10	458,090	67.9%	59,905 (13.1%)	28,256 (6.2%)	0.468
C	50	1,014	6	17	251,069	59.9%	32,461 (12.9%)	15,871 (6.3%)	0.469
D	52	1,291	3	17	317,337	60.4%	45,834 (14.4%)	22,711 (7.2%)	0.468
E	50	366	6	32	93,904	43.2%	12,566 (13.4%)	6,350 (6.8%)	0.474
F	52	478	3	32	126,457	45.3%	18,843 (14.9%)	9,655 (7.6%)	0.471
G	50	100	6	10	29,855	44.5%	29,855 (17.7%)	2,801 (9.4%)	0.463
H	44	na	3	4	26,201	73.0%	26,201 (100.0%)	13,539 (52.0%)	0.507

Figure 1.

Cal. (Calosoma) sycophanta CBX0022 / FR
C. (Orinocarabus) concolor CBX0073 / IT
'Eucarabi'
C. (Carabus) granulatus CBX0043 / FR
C. (Archicarabus) nemoralis CBX0072 / FR
C. (Procrustes) coriaceus CBX0042 / FR
C. (Chrysocarabus) auronitens CBX0027 / FR
C. (Hygrocarabus) nodulosus nodulosus CBX0190 / AT
C. (Hygrocarabus) nodulosus nodulosus CBX0192 / IT
C. (Hygrocarabus) nodulosus nodulosus CBX0019 / FR
C. (Hygrocarabus) variolosus variolosus CBX0193 / SK
C. (Hygrocarabus) variolosus variolosus CBX0182 / RO
Carabus
C. (Heterocarabus) marietti muchei CBX0101 / TK
C. (Chaetocarabus) arcadicus arcadicus CBX0211 / GR
C. (Chaetocarabus) arcadicus arcadicus CBX0079 / GR
C. (Chaetocarabus) intricatus lefebvrei CBX0178 / IT
C. (Chaetocarabus) intricatus intricatus CBX0078 / IT
C. (Chaetocarabus) intricatus intricatus CBX0195 / FR
'Arcifera'
C. (Platycarabus) cychroides CBX0203 / IT
C. (Platycarabus) cychroides CBX0170 / IT
C. (Platycarabus) cychroides CBX0080 / IT
C. (Platycarabus) cychroides CBX0081 / IT
C. (Platycarabus) depressus lucens CBX0207 / IT
0.005
subst./site
C. (Platycarabus) depressus lucens CBX0110 / IT
C. (Platycarabus) depressus lucens CBX0183 / IT
C. (Platycarabus) depressus depressus CBX0111 / CH
C. (Platycarabus) depressus depressus CBX0205 / CH
C. (Platycarabus) depressus depressus CBX0206 / CH
C. (Platycarabus) depressus depressus CBX0085 / IT
C. (Platycarabus) depressus depressus CBX0108 FR
C. (Platycarabus) depressus depressus CBX0088 /FR
C. (Platycarabus) creutzeri baldensis CBX0092 / IT
C. (Platycarabus) creutzeri rinaldoi CBX0119 / IT
C. (Platycarabus) creutzeri viridimicans CBX0120 / AT
C. (Platycarabus) creutzeri kircheri CBX0106 / AT
C. (Platycarabus) fabricii fabricii CBX0094 / AT
C. (Platycarabus) fabricii fabricii CBX0112 / AT
C. (Platycarabus) fabricii fabricii (**PT** ssp *fassati*) CBX0095 / SK
C. (Platycarabus) fabricii fabricii CBX0093 / CH
C. (Platycarabus) fabricii fabricii CBX0113 / CH
C. (Platycarabus) fabricii fabricii (ssp *fassati*) CBX0171 / SK
C. (Platycarabus) fabricii malachiticus CBX0172 / RO
C. (Platycarabus) irregularis montandoni CBX0165 / KP
C. (Platycarabus) irregularis montandoni CBX0166 / SK
C. (Platycarabus) irregularis bucephalus CBX0098 / SI
C. (Platycarabus) irregularis bucephalus CBX0168 / SI
C. (Platycarabus) irregularis irregularis CBX0115 / AT
C. (Platycarabus) irregularis irregularis CBX0020 / FR
C. (Platycarabus) irregularis irregularis CBX0208 / CH
C. (Platycarabus) irregularis irregularis CBX0194 / HU
C. (Platycarabus) irregularis irregularis CBX0099 / FR

6X	3X	Minimum genomic coverage
A	B	A/ min. 10 samples / 1'481 loci / 50 taxa
		B/ min. 10 samples / 1'965 loci / 52 taxa
C	D	C/ min. 17 samples / 1'014 loci / 50 taxa
		D/ min. 17 samples / 1'291 loci / 52 taxa
E	F	E/ min. 32 samples / 366 loci / 50 taxa
		F/ min. 32 samples / 478 loci / 52 taxa

■ IQ-TREE SH-aLRT ≥ 80 and UFBoot ≥ 95
■ IQ-TREE SH-aLRT ≥ 80 or UFBoot ≥ 95
□ IQ-TREE SH-aLRT < 80 and UFBoot < 95
■ IQ-TREE Not recovered
* wASTRAL LPP ≥ 0.95
* wASTRAL LPP < 0.95
* wASTRAL Not recovered

○ Ethanol-preserved
▲ Dry-pinned (< 30-year-old)
★ Dry-pinned (≥ 30-year-old)
⬠ Dry-pinned (Unknown age)
PT Paratype