Museomics of *Carabus* giant ground beetles evidences an Oligocene origin and *in situ* Alpine diversification

Authors

Marie T. Pauli[1,2,†], Jérémy Gauthier[1,2,†], Marjorie Labédan[1,3], Mickael Blanc[1], Julia Bilat[1], Emmanuel F.A. Toussaint[1,*]

Affiliations

[1] *Natural History Museum of Geneva, Geneva, Switzerland*

[2] *Naturéum, State Museum of Natural Sciences, Lausanne, Switzerland*

[3] *Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland*

† Joint first authors

* Correspondence: Emmanuel F.A. Toussaint, Natural History Museum of Geneva, 1208 Geneva, Switzerland. Email: emmanuel.toussaint@ville-ge.ch.

ORCID numbers
Marie T. Pauli (0000-0003-1488-9851)
Jérémy Gauthier (0000-0001-6666-1002)
Marjorie Labédan (0000-0002-2643-1277)
Mickael Blanc (0009-0005-1833-0854)
Julia Bilat (0009-0000-9249-7103)
Emmanuel F.A. Toussaint (0000-0002-8439-1285)

Abstract: The development of museomics represents a major paradigm shift in the use of natural history collection specimens for systematics and evolutionary biology. New approaches in this field allow the sequencing of hundreds to thousands of loci from across the genome using historical DNA. HyRAD-X, a recently introduced capture method using bench-top designed probes, has proved very efficient to recover genomic-scale datasets using natural history collection specimens. Using this technique, we infer at both the intra- and interspecific levels, the most robust phylogeny of Arcifera to date, an ecologically and morphologically diverse clade of *Carabus* giant ground beetles. We successfully generated a genomic dataset of up to 1965 HyRAD-X loci for all described species, permitting to infer a robust dated phylogenomic tree of this clade. Our species delimitation and population genomic analyses suggest that the current classification in Arcifera is in line with its evolutionary history. Our results suggest an origin of Arcifera in the late Oligocene followed by speciation events during the warm mid-Miocene unlinked to Pleistocene glaciations. The dynamic paleogeographic history of the Palearctic region likely contributed to the diversification of this lineage with a relatively ancient colonization of the proto-Alps followed by *in situ* speciation where most species of Arcifera are currently found sometimes syntopically likely as a result of post-glaciations secondary contacts.

Keywords: Arcifera; Beetle evolution; Carabinae; Historical DNA; HyRAD-X; Phylogenomics; Palearctic biogeography; Pleistocene glaciations

Introduction

Recent developments in museomics are opening new prospects allowing samples from natural history collections (NHC) to enter the era of genomics (reviewed in (Raxworthy & Smith 2021; Card *et al.* 2021). Specimens held in the collections are crucial for the study of systematics and taxonomy, but also for the study of ecology and evolution (Duchenne *et al.* 2020). Sampling from NHC specimens is also a major asset to study groups that are currently rare in the wild, for which authorizations to collect new specimens are difficult to obtain, or for which a comprehensive taxonomic and geographic sampling would require extensive fieldwork campaigns. Such a strategy is therefore very powerful when working on taxonomic groups presenting a wide geographical range. In extreme cases, and when species are believed to be extinct, NHC represent the only potential source of genetic data (Toussaint *et al.* 2021; de-Dios *et al.* 2023).

Innovative approaches are now making it possible to obtain genetic information from NHC specimens for which it has long been impossible to recover DNA. The DNA in these specimens, referred to as historical DNA (hDNA), is in low quantity, fragmented, has undergone chemical modifications over time and contains contaminants linked to the history of the collection (Raxworthy & Smith 2021). Improvements in extraction methods, sequencing technologies but above all the development of new capture methods has enabled an increasing amount of genetic information to be recovered. They allow to overcome the difficulties associated with highly degraded and fragmented hDNA from NHC samples, which prevents conventional amplification using standard molecular primers (Landry *et al.* 2023). Among these methods, Ultra Conserved Elements (Blaimer *et al.* 2016; Faircloth 2017) or anchored hybrid enrichment of conserved regions (AHE, Lemmon *et al.* 2012; Mayer *et al.* 2021), are based on the capture of informative loci previously designed from existing genomic data and generally target fairly conserved regions in order to perform large phylogenies. Applying these approaches to NHC specimens allows to integrate samples that are complicated to obtain in the field. In order to work on non-model species for which no prior genomic data is available, the HyRAD (Suchan *et al.* 2016) and HyRAD-X (Schmid *et al.* 2017) approaches enable probes to be designed directly from a few phylogenetically close fresh samples. These approaches, based on the bench-top production are considerably less expensive than commercially synthesized probes. The probes are designed using a ddRADseq protocol (Peterson *et al.* 2012) allowing to target thousands of loci randomly distributed along the genome. This approach is suitable for integrating NHC samples into population-scale studies (Gauthier *et al.* 2020) or for phylogenetic studies of taxa that have recently diverged, such as within a genus (Gauthier *et al.* 2023). The HyRAD-X approach designs probes on fresh RNA extractions. By targeting only expressed gene loci, the HyRAD-X approach makes it possible to investigate phylogenetic questions at older evolutionary scales than the HyRAD approach (Toussaint *et al.* 2021). Using these probe sets, hDNA is then captured by hybridization and sequenced using NGS technologies. This allows only the targeted loci to be recovered while eliminating all unwanted fragments such as

contaminants. After sequencing, the loci are reconstructed and aligned using appropriate
bioinformatic pipelines in order to make phylogenetic inferences (Toussaint *et al.* 2021). Unlike
random Whole Genome Sequencing (WGS) of all the extracted DNA, these targeted approaches
enable better recovery of loci and integration of a larger number of NHC samples into the
phylogenetic inferences (Toussaint *et al.* 2021). Although the efficiency of HyRAD-X has been
tested at higher taxonomic levels, an empirical investigation of its performance at the interface
between population and species levels is needed.

The genus *Carabus*, Linnaeus 1758 (Coleoptera: Carabidae), is a monophyletic highly
diversified lineage comprising *ca.* 970 species classified into 91 subgenera (Deuve 2019, 2021).
This genus, together with its sister genus *Calosoma* (cosmopolitan, 130 species) form the tribe
Carabini (Osawa *et al.* 2004; Toussaint Fls & Gillett 2018; Toussaint *et al.* 2021; Sota *et al.*
2022). Within *Carabus*, the clade named Arcifera Imura, 1996 is sister to the very diversified
clade Eucarabi Deuve, 2013 (Deuve *et al.* 2012; Deuve 2021). This clade is mainly Palearctic,
ranging in the west from southwest England to Ukraine and Turkey in the east. The range of this
group notably encompasses the Carpathian Mountains as well as the Swiss, Italian, Austrian and
Dinaric Alps. It currently includes four subgenera: *Carabus (Hygrocarabus)* Thomson, 1875,
Carabus (Platycarabus) Morawitz, 1886, *Carabus (Chaetocarabus)* Thomson, 1875 and
Carabus (Heterocarabus) Morawitz, 1886 (Deuve 2019, 2021). Within Arcifera, the subgenus *C.*
(Hygrocarabus) contains two species, *Carabus nodulosus* Creutzer, 1799 and *Carabus*
variolosus Fabricius, 1787, found from France to Ukraine, the status of which has been
extensively debated over the past decades due to reduced morphological differences and
inconsistent genetic admixture patterns (Müller-Kroehling *et al.* 2006, 2014; Matern *et al.* 2009,
2010; Mossakowski *et al.* 2020). These hygrophilous nocturnal species live in river banks and
hunt close or in the water of cold forest streams. The two species are in relative allopatry with *C.*
nodulosus being found from eastern France to Austria and western Balkans, and *C. variolosus*
from Slovakia to Ukraine and Bulgaria (Kulijer 2019; Deuve 2021; Bekchiev *et al.* 2022;
Hristovski *et al.* 2023). Despite the protection status of their habitat (Annexes II and IV of the
European Union's Habitats Directive), these two species appear to be declining due to
anthropogenic activities and their consequences (Tyszecka *et al.* 2023). The subgenus *C.*
(Chaetocearabus) only contains two allopatric species following Deuve (Deuve 2019, 2021), the
widespread *Carabus intricatus* Linné, 1761 found from western France to Ukraine and Greece,
and the Greek endemic *Carabus arcadicus* Gistl, 1850. These two species are found in sympatry
in Greece where hybrids are known (at the Katara Pass in the Epirus region for instance).
Additionally, the status of several subspecies in both taxa has been debated, and some authors
recognize *Carabus arcadicus merlini* Schaum, 1861 (Greece), *Carabus intricatus lefebvrei*
Dejean, 1826 (southern Italy including Sicily) and *Carabus intricatus krueperi* Reitter, 1896
(Greece) as separate species within which additional subspecific taxa have been described (e.g.,
Cavazzuti & Ghiretti 2020). Perhaps the most debated taxon of the three being *Carabus*
intricatus lefebvrei found south of Umbria to northern Sicily which is largely allopatric from the
rest of the Italian populations found only in the extreme north of Italy from Piemonte to Friuli

(Cavazzuti & Ghiretti 2020). The subgenus *C. (Heterocarabus)* contains a unique species, *Carabus marietti* Cristofori & Jan, 1837, that is found in southern Bulgaria near the Black Sea and in Anatolia (Turkey), however its ecology and relationships between the numerous described subspecies remain poorly known (Gueorguiev & Gueorguiev 1995; Hieke & Wrase 2008). Finally, the subgenus *C. (Platycarabus)* is composed of five currently accepted species: *Carabus creutzeri* Fabricius, 1801, *Carabus cychroides* Baudi, 1860, *Carabus depressus* Bonelli, 1811, *Carabus fabricii* Panzer, 1812 and *Carabus irregularis* Fabricius, 1792. These beetles are characterized by a flattened morphology, long legs, and elytra generally covered with small foveae (except in *Carabus depressus lucens* Schaum, 1857). They are most widely distributed in Central and Eastern Europe, generally at high altitudes, in mountain forests and alpine pastures. The subgenus contains helicophagous species that exhibit different hunting techniques related to the morphology of their mandibles and prothorax (Casale *et al.* 1998). For instance, *Carabus cychroides* with a thin, elongated head and prothorax, is adapted to enter gastropod shells and has undergone a process known as "cychrization". This species has a very restricted range in the Piedmont region of Italy, is endangered and the focus of reinforced conservation programs (Anselmo & Rizzioli 2022a; b). In contrast, the species *C. irregularis* presents a "licinization" resulting in a large head likely adapted to cracking snail shells (Casale *et al.* 1998). The relationships among species of the subgenus *C. (Platycarabus*) are still debated, and the various taxonomic divisions, both species and subspecies, have yet to be clarified (Casale *et al.* 1998; Deuve 2021). Natural hybrids have been suggested between *C. fabricii* and *C. depressus*, *C. creutzeri* and *C. irregularis*, *C. creutzeri* and *C. depressus*, and *C. depressus* and *C. cychroides* (Casale *et al.* 1998; Camard & Leplat 2004; Casale & Rapuzzi 2015), indicating the need for an in-depth study of possible hybridization in this group.

One of the earliest attempts to elucidate the phylogeny of Arcifera was conducted by Ishikawa (Ishikawa 1984), using 21 morphological characters. This study supported the monophyly of Arcifera and placed *C. (Hygrocarabus)* as sister to the rest of the group, in which *C.* (*Chaetocarabus*) was sister to *C. (Heterocarabus)* and *C. (Platycarabus)*. The first placement of Arcifera members in a molecular phylogeny was based on a single mitochondrial fragment (i.e. ND5), and recovered *C.* (*Chaetocarabus*) and *C. (Platycarabus)* as sister lineages, close to *Carabus* (*Limnocarabus*) Géhin, 1876 and *Carabus* (*Euleptocarabus*) Nakane, 1956 (Imura *et al.* 1998). A subsequent study with the same gene fragment but increased taxon sampling recovered a paraphyletic Arcifera due to the placement of *C. (Hygrocarabus)* as sister to *Carabus* (*Limnocarabus*) and *C.* (*Euleptocarabus*) (Imura *et al.* 2000). In the same study, *C. (Heterocarabus)* was sister to *C.* (*Chaetocarabus*) and *C. (Platycarabus)*. Using the same gene fragment, another study inferred *C. (Platycarabus)* as sister to *C.* (*Chaetocarabus*) and *C. (Heterocarabus)*, within a largely unresolved *Carabus* clade (Su *et al.* 2003). A subsequent study using two nuclear gene fragments recovered Arcifera, represented by *C.* (*Chaetocarabus*) and *C. (Platycarabus)* as sister to the rest of the genus (=Eucarabi) (Sota & Ishikawa 2004). More recently, Deuve *et al.* (2012) used ten loci to recover Arcifera as sister to the Eucarabi and within Arcifera, they recovered *C. (Hygrocarabus)* as sister to *C.* (*Chaetocarabus*) and *C.*

(Platycarabus). Phylogenetic relationships among *C. (Platycarabus)* were also investigated using Sanger sequencing data (Casale *et al.* 1998), suggesting that *C. irregularis* is sister to the rest of the subgenus with *C. fabricii* and *C. depressus* being the most derived lineages in the tree. In parallel to a moderate refinement in the phylogenetic inferences of Arcifera, the estimation of divergence times in the clade has made some progress. Estimates for the origin of Arcifera based on few loci range from the mid-Miocene (ca. 14 Ma, Deuve *et al.* 2012) to the early Oligocene (ca. 30 Ma, Schmidt *et al.* 2023). No major improvement in our understanding of Arcifera systematics and evolution has been achieved in the past decade and there is a need to infer a robust evolutionary tree for this section of *Carabus* to better understand the morphological, ecological, and geographical evolution of constituent lineages.

In this study, we take advantage of the HyRAD-X approach to integrate a large number of samples throughout the large geographical range of Arcifera. We rely on phylogenomic inferences, species delimitations and population genomics approaches to clarify the taxonomy and elucidate the evolutionary history of this complex group of species. In particular, we use this new genomic framework to test which abiotic factors may have fostered the diversification of Arcifera through space and time in the Cenozoic.

Material and methods

Taxon sampling and DNA extraction

The initial sampling was designed in order to sample major lineages within the Arcifera group comprising four subgenera *C. (Chaetocarabus)*, *C. (Heterocarabus)*, *C. (Hygrocarabus)* and *C. (Platycarabus)* (Deuve 2019). A total of 96 samples were initially collected, mainly from NHC samples (87 samples, i.e. 90% of the dataset) but also from a few fresh samples (9 samples, i.e. 10% of the dataset) when these were available (Supplementary Table 1). Multiple specimens of the same taxa and geographic populations were initially selected to anticipate the risk of failure linked to hDNA degradation that can result in specimens being excluded. NHC specimens used in this study are kept at the Natural History Museum of Geneva (MHNG, 76 specimens) and Zoologische Staatssammlung München (ZSM-SNSB, 10 specimens). Eight specimens collected in 96% ethanol were also used and have been deposited in the MHNG collections. DNA was extracted destructively from a single leg using a QIAamp DNA Micro kit (Qiagen, Hilden, Germany) and eluted in 20µL of ultrapure water. Quantity and quality of the purified DNA were assessed with a Fragment Analyzer. Based on DNA quality and concentrations, 38 specimens were not included in the final samples selected for capture, enrichment and sequencing (*ca.* 40% of DNA extractions not processed). Overall, a total of 56 Arcifera specimens were sequenced de novo for this study, representing all Arcifera subgenera and species, several subspecies for the most widespread species as well as a good geographical representation of each species range (taking into account 12 specimens that were eventually not included in the decisive datasets, see Results). Early sampling erosion and discarded samples are commonly not discussed in the framework of museomics studies, but we believe that this is critical to understand the limitations and cost of such approaches in modern phylogenomic studies. The initial sampling in this study

was specifically designed to accommodate a *ca.* 40–50% specimen loss during DNA
quality/quantity assessment (e.g., Toussaint *et al.* 2021), and therefore the resulting sampling is
well-suited to tackle the focal taxonomic and evolutionary questions in Arcifera. The final taxon
sampling was complemented by eight samples of *Carabus* (including one of *C. irregularis* and
one of *C. variolosus*) and one of *Calosoma sycophanta* (Linné, 1758) retrieved from (Toussaint
et al. 2021) (see Supplementary Table 1 for more details).
216
HyRAD-X protocol
The HyRAD protocol was applied as in (Toussaint *et al.* 2021) allowing to generate the same
probe set and therefore a backward compatibility with the data of this previous study. For fresh
specimens a shearing step with NEBNext dsDNA Fragmentase (New England Biolabs) was
performed before library preparation. Shotgun libraries were prepared based on the protocol
developed in (Tin *et al.* 2014). Purified DNA was phosphorylated with T4 Polynucleotide
Kinase. After heat-denaturation into single-stranded DNA, G-tailing was performed with
Terminal Transferase and second strand DNA was synthesized with Klenow Fragment (3□-
>5□exo-) using a poly-C oligonucleotide. Blunt-end reaction was performed with T4 DNA
Polymerase and barcoded adapters were ligated to the phosphorylated end with T4 DNA ligase.
After adapter fill-in with Bst DNA Polymerase (Large Fragment), PCR were run using Phusion
U Hot Start DNA Polymerase (Thermo Scientific) and indexed PCR primers. Libraries were
pooled in equimolar quantities based upon their respective concentrations. Hybridization capture
for enrichment of shotgun libraries was based on the MYbaits protocol (Arbor Biosciences)
modified as in (Toussaint *et al.* 2021) to include a two-step capture at different temperatures (Li
et al. 2013). Final library sequencing was performed on Illumina NovaSeq 6000 SP using a
paired-end protocol (Lausanne Genomic Technologies Facility, Switzerland).
234
Illumina sequencing data cleanup and processing
Raw reads were demultiplexed according to indexes and barcodes using CutAdapt2 (Martin
2011). Reads were cleaned using CutAdapt2 (Martin 2011) and quality was assessed all along
the process using fastqc (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Cleaned
reads were individually mapped onto the loci catalog using BWA-MEM (Li 2013)
(Supplementary Figure 1). The GATK (GenomeAnalysisTK) IndelRealigner tool (McKenna *et*
al. 2010) realigned the indels and deamination were corrected using *mapDamage2.0* (Jónsson *et*
al. 2013). For each sample and each locus, a consensus sequence was generated from the
mapping file using samtools mpileup, bcftools and *vcfutils.pl* (Li *et al.* 2009). Consensuses were
generated keeping the majority allele at each position. Twelve samples with too much missing
data (more than 80% of N), were identified using seqtk and removed (Supplementary Table 1).
Two thresholds of minimum coverage (min_cov) were applied to keep positions: min_cov=3 and
min_cov=6. To test different levels of missing data, decisive datasets were generated applying
three thresholds for the minimum number of samples per locus (min_sample): min_sample=10,
min_sample=17 and min_sample=32. As a result, six datasets were generated: Dataset A

(min_cov=6, min_sample=10, 50 taxa, 1,481 loci), Dataset B (min_cov=6, min_sample=10, 52
taxa, 1,965 loci), Dataset C (min_cov=6, min_sample=17, 50 taxa, 1,014 loci), Dataset D
(min_cov=3, min_sample=17, 52 taxa, 1,291 loci), Dataset E (min_cov=6, min_sample=32, 50
taxa, 366 loci) and Dataset F (min_cov=3, min_sample=32, 52 taxa, 478 loci) (Table 1). The
consensus sequences were combined and aligned with MAFFT using the --auto option. Eight
samples from (Toussaint *et al.* 2021) were integrated at the alignment step. The final datasets
only differ at the taxon sampling level with respect to *C. arcadicus merlini* CBX0176 and *C.*
cychroides CBX0082 that were included only in Datasets B, D and F (these two taxa were
systematically discarded because of low genomic coverage when generating loci with a
min_cov=6, i.e., in Datasets A, C and E).
MitoFinder (Allio *et al.* 2020) was used to identify mitochondrial genes among all
sequenced loci. The kept genes were shared by at least half of the samples. The genes were
aligned, and the sequences were cleaned in Geneious. Individual locus haplotype networks were
built in SplitsTree v.4.19.1 (Huson & Bryant 2006). The networks were reconstructed using
calculated uncorrected p-distances and the NeighborNet algorithm. All non-Arcifera outgroups
were removed before analyses. In parallel, a SNP calling was performed on the mapping files
from the Arcifera species using GATK (McKenna *et al.* 2010) in order to perform
complementary population genomic analyses and compare the results with those obtained from
the locus-oriented approach, avoiding any bias linked to locus reconstruction (Dataset H).
269
Phylogenetic inferences
For each dataset, phylogenetic inferences were performed using IQ-TREE v2.0.5 (Minh *et al.*
2020) using the edge-linked partition model (Chernomor *et al.* 2016). First, the best partitioning
schemes were estimated using PartitionFinder v2.1.1 (Lanfear *et al.* 2017) with the rcluster
algorithm under the Akaike Information Criterion corrected (AICc), with a rcluster-max of 2,000
and a rcluster-percent of 20. The resulting partitioning schemes were then used in IQ-TREE to
select corresponding models of nucleotide substitution using ModelFinder (Kalyaanamoorthy *et*
al. 2017) and the AICc across all available models in IQ-TREE. To avoid local optima, we
performed 100 independent tree searches for each dataset in IQ-TREE. To estimate branch
support, we calculated 1,000 ultrafast bootstraps along with 1,000 SH-aLRT tests in IQ-TREE
(Guindon *et al.* 2010; Hoang *et al.* 2018). We used the hill-climbing nearest-neighbour
interchange topology search strategy to avoid severe model violations leading to biased ultrafast
bootstrap estimations (Hoang et al., 2018). The best tree for each analysis was selected based on
the comparison of maximum likelihood scores. Coalescent species trees were inferred using
ASTRAL-hybrid (Zhang & Mirarab 2022). We first performed individual locus trees using IQ-
TREE v2.0.5 (Minh *et al.* 2020) and branch supports were assessed using 1,000 ultrafast
bootstraps. Best substitution model for each locus was estimated using ModelFinder
(Kalyaanamoorthy *et al.* 2017). Species tree reconstruction was performed combining gene trees
using the weighted-ASTRAL optimization algorithm (Zhang & Mirarab 2022) taking into
account phylogenetic uncertainty by relying on branch length and branch support across locus

trees. As a complement to the locus reconstruction approach, we performed phylogenetic inferences based on the SNP set used for the population genomic approaches. Bi-allelic SNPs shared by at least four samples were extracted and all invariant sites removed. Species trees were inferred with RAxML-NG (Kozlov *et al.* 2019) using GTR+G+ASC_LEWIS model for ascertainment bias correction and branch supports were assessed using 1,000 bootstraps.

Divergence time estimation

Divergence time estimation was performed in BEAST 1.10.4 (Suchard *et al.* 2018) based on a subset of loci selected using a gene-shopping approach to make these analyses tractable on a bioinformatic cluster. SortaDate (Smith *et al.* 2018) was used with default settings to select 100 loci sorted using the following criteria: clock-likeness, tree length, and least topological conflict with the IQ-TREE species tree on dataset E. The selected loci were then concatenated into a Dataset G for relaxed-clock Bayesian divergence time estimation. The best partitioning scheme and substitution models were determined with PartitionFinder2 (Lanfear *et al.* 2017) using the greedy algorithm with the parameter minsubset-size = 2000 and the Bayesian information criterion algorithm to choose between competing models. Clock partitioning was implemented by 1) a single clock for all partitions and 2) a clock for each partition (eight in total; see Results). A Bayesian lognormal relaxed clock model was assigned to the different clock partitions. Different tree models were tested using a Yule pure birth model (Yule 1925; Gernhard 2008), a birth-death model (Drummond *et al.* 2006; Gernhard 2008) as well as a Constant population size coalescent model (Kingman 1982). Since the fossil record of *Carabus* is scarce, we relied on secondary calibrations from a study focusing on Adephaga evolution based on 23 beetle fossil calibrations (Baca *et al.* 2021). According to this study, the separation between the genera *Calosoma* and *Carabus* occurred about 41.4 [37.1–46.1] million years ago (Ma). This age was used as a secondary calibration for the corresponding node in our topology (split *Calosoma*/*Carabus*, in this case the root). A second calibration was used to constrain the crown of *Carabus*. Following (Baca *et al.* 2021), this node was constrained to match the recovered age in their study at about 25.4 [22.8–28.2] Ma. The analyses were conducted for 50 million generations, sampling parameters and trees every 5000 generations. The maximum clade credibility tree for each analysis was generated in TreeAnnotator 1.10.4.

Species delimitation and hybridization

We used a combination of species delimitation methods and population genomic approaches to test species and subspecies limits. For these analyses we excluded the six non-Arcifera outgroup specimens resulting in a dataset composed of 44 samples. We extracted the 44 Arcifera samples from Dataset E composed of 366 shared loci which present the lowest level of missingness (Table 1). First, BPP (Flouri *et al.* 2018) was used with the A11 option, using inverse-gamma distributed diffuse priors (α = 3; ß = 1000) for the population sizes (θ) and root ages ($\tau 0$). Analysis was run for 100,000 generations, sampling every 100 generations after a burnin of 8,000 generations. Second, the multi-locus species delimitation using Bayesian model

comparison implemented in the TR2 package (Fujisawa *et al.* 2016) has been applied on the same dataset. Locus trees generated with IQ-TREE v2.0.5 (Minh *et al.* 2020) and previously used for the weighted-ASTRAL approach were used as well as the maximum likelihood IQ-TREE consensus tree on dataset A as guide tree. Outgroups were removed from gene trees and the guide tree.

From the SNPs (Dataset H), population clustering was assessed using STRUCTURE 2.3.3 (Pritchard *et al.* 2000). Bi-allelic SNPs shared by at least 40% of the samples were extracted using VCFtools v0.1.12a (Danecek *et al.* 2011). Because markers are supposed to be unlinked, we extracted randomly only one SNP by locus. K-values from 1 to 15 were tested with no prior population information and performed three times for each of them to verify a convergence of estimations. A burn-in of 100,000 runs was used followed by 500,000 iterations. The most likely number of clusters was determined using the Evanno method (Evanno *et al.* 2005) implemented in Structure Harvester (Earl & vonHoldt 2012). The replicates were then combined and the figures generated using CLUMPAK server (Kopelman *et al.* 2015). To investigate putative admixture between species or subspecies we estimated Patterson's D statistic (ABBA-BABA test) (Patterson *et al.* 2012) for all subspecies/species quartets using the Dsuite (Malinsky *et al.* 2021). The analyses were performed on bi-allelic SNPs shared by at least 40% of the samples composed of 6,743 SNPs. Z-scores and associated p-values were calculated to assess the significance of the results.

Results

Museomics approach efficiency

The combination of historical and fresh samples enabled us to compare the effectiveness of museomics methods. The DNA concentrations obtained from a single leg are very variable between fresh samples (mean = 8.37 ng/µL; sd = 6.82 ng/µL) and NHC samples (mean = 1.18 ng/µL; sd = 2.45 ng/µL). There was a significant correlation between the quantity of DNA extracted and the age of the specimens (Figure 1A). For the NHC samples, this concentration was not homogeneous, with some samples nevertheless showing a high concentration. Forty samples with a concentration below the detection thresholds, were excluded from the rest of the capture process. It should be noted that some samples with very low DNA concentrations, such as *C. fabricii* CBX0094 captured in 1977 with a concentration of only 0.08 ng/µL, were reliably placed into the final phylogenetic inferences. For specimens with measurable DNA, the capture process worked efficiently, allowing the sequencing of an average of 8.4 million reads per sample (sd = 9.2 million). There was a large difference between the average number of reads obtained from fresh samples (mean = 23.8 millions; sd = 11.0 millions) and NHC samples (mean = 5.4 millions; sd = 5.0 millions). The age of the specimens also had an influence on the number of reads obtained, as there was a significant correlation between the age of the specimens and the number of reads obtained (Figure 1B).

After locus reconstruction, the difference between fresh and NHC samples persists, with an average of 1765 loci recovered in fresh samples (sd = 553) and 629 in NHC samples (sd =

447) (Figure 1C). This difference is of the same order when looking at shared loci (Figure 1D).
There is a large heterogeneity in the number of loci recovered between NHC samples, largely
linked to the age of the specimen. Samples with too few loci (< 150 loci), i.e. 12 samples, had to
be excluded from the final datasets. For 35 NHC samples, the number of loci recovered, on
average 793 (sd = 400), was sufficient to include them in subsequent analyses. Although strict
filtering steps reduced the number of NHC samples, they also ensured the reliability of the
dataset for downstream inferences.
377
378 *Phylogenomic inferences*
379
We inferred the phylogenetic relationships among Arcifera using six different datasets under
various taxon sampling and/or gene sampling strategies (Figure 2). The results of analyses based
on a concatenation approach performed in IQ-TREE and on a coalescent species-tree approach
conducted in wASTRAL are consistent except for the placement of *C. marietti*, the branching
pattern within *C.* (*Chaetocarabus*) and relationships between *C. creutzeri* and *C. fabricii*. The
subgenus *C.* (*Hygrocarabus*) is recovered as monophyletic in all analyses (including wASTRAL)
and as sister to the rest of Arcifera, however all analyses failed to recover *C. nodulosus* and *C.*
variolosus as reciprocally monophyletic. In all IQ-TREE inferences except the one based on
Dataset E and all wASTRAL analyses except the one based on Datasets A and B, *C. marietti* is
recovered as sister to *C.* (*Chaetocarabus*) with heterogeneous levels of branch support. In the IQ-
TREE analysis of Dataset E, this taxon is recovered as sister to the genus *Carabus* as a whole,
whereas in wASTRAL analyses of Datasets A and B it is recovered as sister to Arcifera except
C. (*Hygrocarabus*) with low branch support. The subgenus *C.* (*Chaetocarabus*) is always
recovered as monophyletic but internal relationships differ between analyses. A minority of
analyses recovered *C. arcadicus* and *C. intricatus* as reciprocally monophyletic (for instance no
wASTRAL analysis recovered this relationship). The subspecies *C. intricatus lefebvrei* is
recovered as sister to the nominal subspecies in all analyses. When *C. arcadicus merlini* is
included (Datasets B, D and F only), it never groups with other specimens of the nominal
subspecies resulting in *C. arcadicus* being consistently inferred as paraphyletic when this taxon
is included (Supplementary Figure 2). The subgenus *C.* (*Platycarabus*) is recovered as
monophyletic and with identical interspecific relationships across all IQ-TREE analyses but
some contention in wASTRAL ones. The alpine endemic *C. cychroides* is recovered as sister to
the rest of the subgenus in all analyses with strong branch support (IQ-TREE and wASTRAL).
The species *C. depressus* is inferred as the next lineage branching off in *C.* (*Platycarabus*) across
all IQ-TREE analyses and most wASTRAL analyses (except in Dataset A and E where it is
recovered as sister to *C. irregularis* with low branch support). The subspecies *C. depressus*
lucens is recovered as sister to the nominal subspecies in all analyses. The placement of the three
remaining *C.* (*Platycarabus*) species is identical across all IQ-TREE analyses with strong branch
support, with *C. creutzeri* being sister to *C. fabricii* and *C. irregularis*. The wASTRAL analyzes
recover different relationships but with low branch support, with a weakly supported sister

relationship between *C. creutzeri* and *C. fabricii* in analyses of Datasets D, E and F. The
subspecies *Carabus fabricii malachiticus* is recovered as nested within the nominal subspecies in
all analyses. The subspecies *Carabus irregularis montandoni* is recovered as sister to *Carabus*
irregularis bucephalus and *Carabus irregularis irregularis* in all IQ-TREE analyses whereas it is
C. irregularis bucephalus that is inferred as sister to *C. irregularis irregularis* and *C. irregularis*
montandoni in all wASTRAL analyses. Overall, the IQ-TREE and wASTRAL inferences are
highly compatible when collapsing the weakly supported relationships in wASTRAL species
trees (gray and red asterisks in Figure 2). In particular, inconsistent relationships in wASTRAL
compared to IQ-TREE always received poor branch support. We observe that branch support and
overall phylogenetic resolution appears positively correlated to gene and taxon sampling (i.e.,
including fewer taxa and fewer loci to improve matrix completeness likely results in a loss of
resolution).
422
423 *Divergence time estimation*
424
The BEAST dating analysis revealed consistent results for the four main nodes, i.e. the root,
Carabus, Arcifera and *C. (Platycarabus)* nodes, according to the three models tested, Yule,
Birth-Death model, and Constant population size coalescent (Figures 3 and 4). The coalescent
model including eight Bayesian log-normal relaxed clocks received the best marginal likelihood
as calculated using stepping-stone sampling in BEAST and was therefore selected hereafter. This
inference suggests an origin of Arcifera at 26.07 Ma (95% HPD: 22.77 - 29.67 Ma) and 14.56
Ma (95% HPD: 12.52 - 16.76 Ma) for the *C. (Platycarabus)* subgenus.
432
433 *Species delimitation and putative hybridization*
434
The different approaches to species delimitation produced contrasting results. The analysis
performed with BPP is the most consistent with morphology and the current classification. The
two species *C. variolosus* and *C. nodulosus* are well separated even though *C. nodulosus* is not
monophyletic in our phylogeny. TR2 approach proposes an oversplit of the three *C. nodulosus*
samples. Conversely, the STRUCTURE approach groups the two species in a single cluster
(Supplementary Figure 6). STRUCTURE analyses reveal that the most probable number of
clusters is found for K=5 (deltaK = 1539.43) (Supplementary Figure 6). However, some clear
splits emerge at higher K values. This is the case for *C. (Heterocarabus) marietti*, represented
by a single sample which is identified as a separate cluster at K=7 or the split between *C.*
(Chaetocarabus) arcadicus - *C. (Chaetocarabus) intricatus* and *C. (Platycarabus) cychroides*
which emerges at K values greater than 5. The divisions we propose (Figure 3) is therefore a
combination of the logical splits identified by STRUCTURE for the different values of K. *C.*
marietti, the only representative of *C. (Heterocarabus)* is delineated as a species in all three
approaches. Within *C. (Chaetocarabus)*, the two species *C. arcadicus* and *C. intricatus* are
delineated by BPP but are merged by TR2 and STRUCTURE, potentially for the same reasons as

in *C. (Hygrocarabus)*. It should be noted that the two subspecies of *C. intricatus*, i.e. *C.*
intricatus lefebvrei and *C. intricatus intricatus*, are never delineated as distinct species. The
species *C. cychroides* was well discriminated in two of the three approaches, with only TR2
proposing an additional split of the most basal sample. The two subspecies of *C. depressus*, *C.*
depressus depressus and *C. depressus lucens* are not grouped together in the BPP approach and
are identified as two distinct species. The results of the three methods are fully consistent with
the morphology for *C. creutzeri* and *C. fabricii*. For *C. irregularis*, the situation is similar for two
of the three methods, BPP and Structure. Among the 85 trios analyzed, high D-statistics values,
> 0.25, with significant p-values were observed for three trios. For two of these, *C. cychroides*
was observed in P1 and *C. arcadicus* in P3. Despite this, no f-branch signal significantly
different from zero could be identified (Supplementary Figure 7). These results suggest an
absence of past introgression between the different species and subspecies.
462
463 **Discussion**
464
465 *Using museomics to obtain an extensive dataset*
466
The HyRAD and HyRAD-X methods are unique in that they allow in-house production of
probes using a ddRAD protocol, either directly on the DNA of a few fresh samples (Suchan *et al.*
2016) or on their RNA (Schmid *et al.* 2017). These approaches allow to target several thousand
loci and generate in turn high-resolution phylogenomic inferences (Young & Gillung 2020). In
this study, we obtained 1,965 loci for the most extensive dataset. These loci were informative
enough to resolve both the deep relationships between outgroups and the more recent
relationships at the intrageneric and intraspecific scales. In addition, the identification of SNPs
on these loci also enabled population genomic approaches such as the study of genetic structure
and admixture.
In addition, the HyRAD-X approach made it possible to integrate samples with extremely
low initial DNA quantities. However, out of 96 samples from which DNA was extracted, 40 had
an undetectable quantity of DNA. In the context of museomics projects, it is therefore
instrumental to plan for redundancy in the sampling, with several samples per targeted taxon, in
order to compensate for any failures. Furthermore, the ability to generate genetic information
from hDNA is not entirely predictable. The recovery of meaningful genomic data does not seem
to be linked to the age of samples (Figure 1), in line with existing observations (Toussaint *et al.*
2021; Nunes *et al.* 2022). In that vein, large amounts of genomic data could be obtained from
older NHC samples when almost none could be obtained from more recent samples. The quality
and quantity of DNA that can be extracted from NHC specimens is linked to factors that we
cannot control, such as the conditions of collection and preservation process (Post *et al.* 1993;
Dillon *et al.* 1996; Ruppert *et al.* 2023).
488
Systematics and species delimitation in Arcifera

Our results provide a robust phylogenomic tree of Arcifera for the first time (Figure 2). Overall,
we support the view that Arcifera represents a monophylum within which all four subgenera
form clades. The monophyly of Arcifera is also supported by the presence of a hook-shaped
ligulum (i.e., arculus) at the base of the endophallus, a strong morphological character that unites
all constituents of this lineage (Imura *et al.* 2000; Deuve *et al.* 2012). Our study is the first to
provide strong evidence for these relationships while including all species of the group. Other
studies based on reduced genomic sampling, often a single gene fragment, either failed to
recover Arcifera as monophyletic (Imura *et al.* 2000; Osawa *et al.* 2004), or had too limited a
taxon sampling to properly test the placement and otherwise monophyly of each subgenus (Su *et*
al. 2003; Sota & Ishikawa 2004; Deuve *et al.* 2012). Except for a minority of analyses, our
results strongly suggest that *C. (Hygrocarabus)* is sister to the rest of Arcifera, with *C.*
(Platycarabus) as sister to a clade formed by *C. (Chaetocarabus)* and *C. (Heterocarabus)*.

Within *C. (Hygrocarabus)*, we recover *C. variolosus* nested within *C. nodulosus*. This
result contrasts with the ones of Mossakowski *et al.* (2020) where the two species were
suggested to be well differentiated genetically. In their study, these authors argued based on the
analysis of two gene fragments that both taxa form distinct clades although several specimens
caused each species to be paraphyletic. Some tests of mating between the two candidate species
were also performed in this study and suggested that the two lineages do not mate. However, the
scale and conditions of these trials do not allow to conclusively rule out potential mating. We
argue that in the current state of our knowledge it is not yet possible to definitively test species
boundaries, past introgression and signature of hybridization between *C. nodulosus* and *C.*
variolosus. A desired approach would be to combine a large geographical sampling as in
Mossakowski *et al.* (2020) with a genomic scale dataset as developed in the present study to
revisit the systematic conundrum within this subgenus at the population level.

Within *C. (Chaetocarabus)*, we recover *C. arcadicus* as sister to *C. intricatus* in most
analyses (Figure 2). These two species are allopatric, morphologically well-differentiated and
little doubt exists with respect to their status as distinct species. Surprisingly our species
delimitation analyses only partly support the two species hypothesis, with TR2 and
STRUCTURE considering that *C. (Chaetocarabus)* is a unique species. Considering the low
genomic coverage of some taxa included in the analyses (see below), the clear morphological
and geographical split between these lineages and the support from BPP analyses, we argue that
the validity of these two species is uncontroversial. Natural hybrids with an intermediate
morphology and usually green dorsal pattern are known to exist along the limits of their
respective ranges in northern Greece (i.e., at the Katara pass) where *Carabus intricatus*
macedonicus (not sampled here) and *C. arcadicus arcadicus* co-occur. Both *C. arcadicus* and *C.*
intricatus also comprise geographically restricted subspecies in Greece that have been
considered valid species by some authors. In the south of Greece, the melanistic subspecies *C.*
arcadicus merlini is endemic to the Peloponnese peninsula and allopatric from the nominal
subspecies present in the north. One specimen of this taxon was sequenced but genomic

coverage was low and therefore it was only included in the less stringent Datasets B, D and F. In the phylogenetic analyses of these datasets, the inclusion of *C. arcadicus merlini* systematically results in all three *C. arcadicus* specimens forming a phylogenetic grade within which *C. intricatus* is nested. We argue that this is an artifact possibly caused by missing genomic sampling and that both species are reciprocally monophyletic as recovered in all other analyses and as suggested by morphology. However, it is possible that *C. arcadicus merlini* represents a distinct evolutionary lineage since it is always recovered as sister to the rest of *C. (Chaetocarabus)*. Additional taxon sampling is needed to test the placement of this morphologically distinct taxon within the subgenus. Across its range, *C. intricatus* is represented by the nominal subspecies from western France and UK to northern Greece. In the south of Italy and Sicily, this species is represented by the allopatric *C. intricatus lefebvrei*. The status of this taxon is debated and some authors consider it a valid species. In our results, we recover this subspecies as sister to the nominal subspecies represented by specimens from France and Piemonte. Our phylogenetic inferences support the view of *C. intricatus lefebvrei* as a possible distinct species but our species delimitation analyses reject this hypothesis. To properly test species boundaries within *C. intricatus*, additional taxon sampling is needed including a much denser geographical sampling of the nominal subspecies along with all described valid subspecies (Deuve 2019). In the Balkans, several subspecies of *C. intricatus* have been described and represented more or less isolated populations restricted to northern Greece. Despite our efforts we could not obtain DNA of good quality for *C. intricatus krueperi* endemic to eastern Thessaly and considered by some authors to be a valid species. Here as well, a denser taxon sampling is needed to properly test species boundaries in this group. The placement of *C. (Heterocarabus) marietti* as sister to *C.* (*Chaetocarabus*) receives support from most analyses in this study. Despite a relatively circumscribed geographic range in northern Turkey and southern Bulgaria, numerous taxa have been described in this subgenus even though currently a single species is considered valid (Turin *et al.* 2003; Deuve 2019). Increasing the taxon sampling for this group by covering all its geographical range would allow testing the match between morphological and genetic diversity and better understand the evolution of this unique lineage at the inter- and intraspecific interface.

Within *C. (Platycarabus)*, we recover *C. cychroides* as sister to the rest of the subgenus. This result is unexpected because this species is a very narrowly restricted endemic to Piemonte mountain ranges where it lives in alpine meadows and scree >2000m. The species was only included once in a phylogenetic framework by Casale *et al.* (1998) who recovered it as a derived lineage close to *C. depressus* and *C. fabricii*. Interestingly, a sister relationship of this species to the rest of *C. (Platycarabus)* was suggested by the analysis of morphological characters in Casale *et al.* (1998). Indeed, this species is morphologically quite different from the rest of the subgenus in that it is one of the most extreme examples of cychrization in *Carabus*, a process by which the pronotum is narrowed to allow predation inside snail shells (= stenocephalic morphology). All species of the subgenus present a stenocephalic morphology, although less marked than in *C. cychroides*, except for *C. irregularis* which is macrocephalic. Our

phylogenetic inferences are therefore important to understand the evolution of predation
strategies and associated morphology across the genus *Carabus* in which both types of
morphologies exist (Sota & Ishikawa 2004). Most malacophagous and helicophagous species in
Carabus are macrocephalic and use their enlarged pronotum, head and robust mandibles to break
snail shells. Cases of stenocephaly are most notably observed in *C. (Platycarabus)* but also in
Carabus (*Damaster*) Kollar, 1836 and *Carabus* (*Macrothorax*) Desmarest, 1850. The fact that *C.*
irregularis, the only *C. (Platycarabus)* macrocephalic species, is recovered as the most derived
species in the subgenus, indicates that macrocephaly possibly evolved from a stenocephalic
morphology unlike what was suggested in Casale *et al.* (1998). In the case of *C. cychroides*, it is
not closely related to any other species of the subgenus as suggested by previous authors, and
despite rare known natural hybrids with *C. depressus* in the Cottian Alps (i.e., Colle delle
Finestre, Monte Morefreddo, Monte Albergian), these species do not share an immediate recent
common ancestry (Sturani 1962; Casale *et al.* 1998; Anselmo & Rizzioli 2022a; b). The rest of
C. (Platycarabus) species and most sampled subspecies are found monophyletic (Figure 2). We
recover the subspecies *C. depressus lucens* as sister to the nominal subspecies in all analyses and
with robust branch support. This subspecies is morphologically quite divergent from the nominal
subspecies and *C. depressus bonellii* as it completely lacks elytral foveae. It is also allopatric
from the rest of the *C. depressus* populations, being found in a small transalpine region between
France and Italy (i.e., French Queyras to Italian Alpi Marittime), and its status as a valid species
even though rejected by three out of four species delimitation analyses should be revisited with
enhanced population sampling. Our taxon sampling within *C. creutzeri* does not allow testing
subspecies monophyly and relationships in detail but species delimitation analyses
unambiguously support a single species (Figure 3). Within *C. fabricii*, we recover the Carpathian
populations of *C. fabricii* (ssp *fassati* = nominal ssp, and spp *malachiticus*) nested within Alpine
populations of the nominal subspecies. This is unexpected to some extent as *C. fabricii* presents
a disjunct distribution between the Alps and the Carpathians (i.e., it is not currently found in the
Danube valley). Our results suggest that despite an allopatric range, gene flow has been
maintained between all populations of this species, however branch supports for internal
relationships in *C. fabricii* are moderate and an enhanced taxon sampling is needed to understand
the past and present connectivity between populations. All species delimitation analyses support
a unique species. One of the most interesting subspecific cases is recovered in *C. irregularis*.
This species is the most widespread of the subgenus ranging from eastern France to Romania and
Ukraine. It comprises three valid subspecies, one of which *C. irregularis montandoni* from the
Carpathians, was suggested to be a valid species based on molecular evidence (Homburg *et al.*
2013). Our results support to some extent this view with *C. irregularis montandoni* being found
sister to the rest of populations in all IQ-TREE analyses but not in wASTRAL analyses where
the other subspecies *C. irregularis bucephalus* is found as sister to the rest of the clade. There
seems to be a genetic differentiation between the three recognized subspecies of *C. irregularis*
but our species delimitation analyses support the view of a single species.
609

The divergence time estimation analyses all recover an origin of Arcifera ca. 26 Ma in the
Oligocene. We did not perform a biogeographic estimation of ancestral ranges in the group
because several species are very widespread and initial attempts resulted in unresolved patterns.
The fact that species boundaries within *C. (Hygrocarabus)* are unstable also prevented a proper
reconstruction. However, it is possible to discuss several phylogenetic splits in the framework of
our results. The stem branch connecting *C. (Hygrocarabus)* to the rest of Arcifera is long,
potentially representing periods of extinction in this lineage. Currently the two recognized
species in the subgenus occur in temperate forests where adults live and hunt near and in good
quality streams. The reconfiguration of such habitats in the past 25 million years due to climatic
oscillations (Westerhold *et al.* 2020) may have extirpated populations and pushed others into
their current ranges. Considering the specificity of these two lineages to their habitat, and
predictions of global warming and their impact on such ecosystems (Capon *et al.* 2021; Bonacina
et al. 2023), it is likely that they may be increasingly threatened in the future.
625   With respect to biogeography, one of the most interesting lineages in Arcifera is the clade
composed of *C. (Chaetocarabus)* and *C. (Heterocarabus)*. Because *C. (Heterocarabus) marietti*
is restricted to eastern Bulgaria and western Turkey, and *C. (Chaetocarabus)* distributed in
Greece (*C. arcadicus* is endemic to Greece), it is likely that the ancestors of this clade originated
in the geologically highly complex Aegean area. The split between the two subgenera ca. 17 Ma
predates the timing of the opening of the Aegean Sea in the Tortonian ca. 8 Ma (i.e., opening of
the Mid-Aegean Trench or Aegean barrier; van Hinsbergen & Schmid 2012), rejecting the
hypothesis of geographic vicariance in the south as suggested in other lineages (Poulakakis *et al.*
2015). Interestingly, both subgenera have very marginally overlapping distributions in the
Thrace basin with *C. (Heterocarabus)* currently distributed on the southern Black Sea coast
where *Carabus intricatus* is also represented by the subspecies *C. intricatus starensis*
(Gueorguiev & Gueorguiev 1995). At the time of divergence in the early Miocene (i.e.,
Burdigalian), the Thrace basin formed a connection between the eastern Balkan peninsula and
Anatolia (Rögl 1997, 1999; Sachsenhofer *et al.* 2017; Erbil *et al.* 2021). It is possible that
ancestral populations dispersed in the Balkan Peninsula and/or in Anatolia where they evolved
independently. Under this scenario, the close geographic ranges of these two species would
likely represent secondary contact associated with more recent colonization of the Thrace basin.
A more robust population-level taxon sampling, especially of *C. (Heterocarabus)*, might
elucidate the fine-scale biogeographic history of this clade in the future. Within *C.*
(Chaetocarabus), the two currently recognized species are mostly allopatric with only a short
overlap in western Greece (e.g. Katara pass). There is no clear geological barrier that may have
fostered vicariant diversification at the time of speciation ca. 7 Ma. Further diversification
appears to be occurring at the population level with *C. intricatus lefebvrei* endemic to southern
Italy and allopatric from the nominal subspecies. Similarly, *C. arcadicus merlini* endemic to
Peloponnese is morphologically quite divergent from the nominal subspecies and might

represent a case of ongoing speciation. The wide dispersal of *C. intricatus* across the western
Palearctic region is likely recent and may be explained by the generalist habitat preference of this
species. Additional geographical and taxon sampling will likely yield more robust inferences of
evolutionary patterns and processes within this clade in the future.

The evolutionary history of the subgenus *C. (Platycarabus)* is also revealed by our
analyses. We recover the narrowly endemic *C. cychroides* as sister to the rest of the subgenus.
This is surprising as it was not suggested by the molecular inference of Casale *et al.* (1998). This
placement has strong implications for our understanding of alpine biogeography in this group.
Only *C. irregularis* has lowland populations and its derived placement in the phylogeny indicates
that alpine specialization was likely ancestral in the subgenus with recent shift in that species to
lower habitats. This phylogenomic pattern and the origin of the subgenus *ca.* 15 Ma during the
warmest period of the Neogene seems to indicate that ancestors of *C. (Platycarabus)* may have
been less specialized than nowadays and were distributed in mountain regions. In the mid
Miocene, mountain ranges across the Alps had the same elevation as nowadays (Campani *et al.*
2012; Krsnik *et al.* 2021), however ecosystems were different due to significantly warmer
climatic conditions. When the climate progressively turned colder these beetles adapted to
ensuing conditions and became alpine specialists. It is possible that species of the subgenus
diverged due to competition, niche filling and/or host specialization as observed in *C. cychroides*
for instance. We hypothesize that in the latest sequence of their evolutionary history, Pleistocene
glaciations played a limited role in speciation since all current species had already diverged
(Figure 3).

Although natural hybrids are known between different species of the subgenus, our
results recover no hybridization signal between them. The most significant case concerns the
species *C. fabricii* and *C. irregularis*, whose ranges largely overlap in Switzerland, Austria and
Slovakia. It is in these sympatric areas that several cases of natural hybridization have been
identified (e.g. at the Radstädter Tauern Pass in Austria, Mandl 1960). However, our genetic
results do not show any hybridization signals between the species, either on genetic structure,
where the two clusters are well separated, or in the approach using Dsuite, which seeks to trace
admixture signals in the lineages. These results suggest that these sporadic hybridization events
are not conserved in populations and could imply a potential infertility of F1s (Casale *et al.*
1998). Furthermore, the networks obtained with the three mitochondrial genes (Supplementary
Figure 4) group the samples of the *C. irregularis* and *C. fabricii* species in the same cluster.
These mitonuclear discordance patterns are frequent in the literature and can be explained by the
specific biological properties of mitochondrial DNA (uniparental inheritance and reduced
recombination; Birky 2001) or differences in the evolutionary histories of nuclear and
mitochondrial markers including incomplete lineage sorting and gene flow among species (Sota
& Vogler 2001; Suchan *et al.* 2017). The results obtained with the nuclear loci are sufficiently
robust to be able to consider that the cases of hybridization observed are either localized or do
not induce lasting admixture between the species. A more detailed analysis of hybrids, local

populations and the implications of hybridization on the fitness of individuals could provide a
better understanding of the mechanisms involved.
Integrating current species distribution, genetic isolation of these alpine species was
already in place when glaciation cycles struck the Alps. As a result, dispersal of populations in
peripheral glacial refugia as observed in *C. irregularis* (Homburg *et al.* 2013) did not result in
genetic homogenization despite species being placed in secondary contact. It is also possible in
the case of the more alpine-adapted species (all but *C. irregularis*) that dispersal occurred in
nunataks rather than peripheral glacial refugia (Holderegger & Thiel-Egenter 2009; Schönswetter
& Schneeweiss 2019; Kosiński *et al.* 2019), which would have resulted in an increased genetic
differentiation among populations as suggested by our analyses. Coupling a more extensive
geographic sampling of these five alpine species with niche modeling analyses may help testing
more specifically the different scenarios that governed range and genetic evolution of these
populations during Pleistocene glaciations.
702

703 **Acknowledgements**
704

We warmly thank Michael Balke for the loan of material from the ZSM-SNSB. We thank Elsa
Ricossa for the digitization of specimens housed at the Natural History Museum of Geneva. We
thank Céline Rochet for assistance in fieldwork. We thank the city of Geneva for an internal
student grant awarded to MTP. We thank Conrad Gillett for allowing the use of his photographs
and drawings in this article. We thank Ivan Rapuzzi for fruitful discussions and feedback on
taxonomic aspects of this work. We also thank Michael Caterino, Julian Dupuis and an
anonymous reviewer for insightful comments on an earlier version of this article, as well as Felix
Sperling for his editorial work.
713

714 **Funding**
715

716 This study was partly funded by a Master student grant awarded by the City of Geneva. EFAT is
717 funded by a SNSF grant 310030_200491.
718

719 **Conflict of interest disclosure**
720

721 The authors declare no conflict of interest.
722

723 **Data, script, code, and supplementary information availability**
724 Raw reads are available on the NCBI BioProject PRJNA1086379. The data underlying this
725 article (final alignments and trees) and bioinformatic scripts are available on Github repository
726 (https://github.com/JeremyLGauthier/Arcifera_phylogeny).
727
728

Figure captions

Table 1. Alignment statistics for each dataset, including the number of taxa, the number of loci, the minimum coverage, the minimum number of taxa, the alignment length, the percentage of missing data, the numbers and percentages of variable sites and of parsimony informative sites, and the GC content.

Figure 1. Statistical summary of locus recovery. Plots representing the relationship between the collection year and DNA concentration (A), number of sequenced reads (B), number of loci recovered for each sample (C), and number of shared loci in final dataset B. In each plot, ethanol-preserved samples are shown in green, samples from museums with an age < 30 years in blue and samples from museums with an age > 30 years in classic yellow. Correlations were tested with Spearman's correlation tests and adjusted coefficients of determination R-squared were estimated using a linear model.

Figure 2. Summary of phylogenetic inferences across Arcifera based on HyRAD-X data. The presented topology is derived from a maximum likelihood analysis performed in IQ-TREE using Dataset A. Branch support from this analysis is shown for all branches. Branch support retrieved in different analyses is shown for major branches according to the inserted caption. Sample type is indicated according to the inserted caption. Abbreviations at the end of each taxon label correspond to the following countries: AT, Austria, CH, Switzerland, FR, France, GR, Greece, HU, Hungary, IT, Italy, KP, Carpathians (Slovakia to Romania), RO, Romania, SI, Slovenia, SK, Slovakia, TK, Turkey. An illustration of a male *Carabus (Platycarabus) cychroides* is presented (Drawing: Conrad Gillett).

Figure 3. Bayesian divergence time estimates for the subgenus *Carabus (Platycarabus)* and Arcifera group. Maximum clade credibility tree obtained from a BEAST analysis using eight Bayesian log-normal relaxed clocks and a Coalescent Constant Size tree model. Node estimates are postburn in median ages, with 95% credibility intervals. Histogram represents the number of loci recovered for each sample and sample type are indicated according to the inserted caption. The section on the right shows the results of species delimitations identified using the different methods indicated above. The shades of gray represent the concordance between the different approaches with black being a total consensus. Habitus of three representative species (1) *Carabus nodulosus nodulosus* (credit : Conrad Gillett), (2) *Carabus intricatus intricatus* (credit : Conrad Gillett) and (3) *Carabus irregularis irregularis* (credit : Conrad Gillett) are shown.

Figure 4. Comparison of divergence time estimation between competing tree models and relaxed-clock partitioning strategies. Box-plots indicate for each analysis (color-coding inserted as a caption on the right side of the figure) the median age of the focal node (see X axis) and associated 95% age credibility interval. BD, birth-death model; CS, constant population size coalescent model.

Supplementary Table 1. Descriptive statistics for each included and non-included sample, including historical sample data, molecular biology information (DNA concentrations), sequencing and loci reconstruction statistics.

Supplementary Figure 1. Schematic representation of the bioinformatic pipeline.

Supplementary Figure 2. Maximum likelihood trees for each dataset: Dataset A (min_cov=6, min_sample=10, 50 taxa, 1'481 loci), Dataset B (min_cov=6, min_sample=10, 52 taxa, 1'965 loci), Dataset C (min_cov=6, min_sample=17, 50 taxa, 1'014 loci), Dataset D (min_cov=3, min_sample=17, 52 taxa, 1'291 loci), Dataset E (min_cov=6, min_sample=32, 50 taxa, 366 loci) and Dataset F (min_cov=3, min_sample=32, 52 taxa, 478 loci). Node supports indicate SH-aLRT and UFBoot values.

Supplementary Figure 3. Species trees obtained with wASTRAL on each dataset: Dataset A (min_cov=6, min_sample=10, 50 taxa, 1'481 loci), Dataset B (min_cov=6, min_sample=10, 52 taxa, 1'965 loci), Dataset C (min_cov=6, min_sample=17, 50 taxa, 1'014 loci), Dataset D (min_cov=3, min_sample=17, 52 taxa, 1'291 loci), Dataset E (min_cov=6, min_sample=32, 50 taxa, 366 loci) and Dataset F (min_cov=3, min_sample=32, 52 taxa, 478 loci). Node supports indicate SH-aLRT and UFBoot values. Node supports indicate LPP values.

Supplementary Figure 4. Individual locus haplotype networks (A. CO1, B. CO3 and C. CYTB). Networks were generated in SplitsTree using calculated uncorrected p-distances and the NeighborNet algorithm. The color coding for the different morphological groups is identical to the one used (photo credit : Marie Pauli).

Supplementary Figure 5. Structure plots estimated on unlinked shared SNPs for K=1 to K = 15. For each K, the Mean(LnProb) is indicated.

Supplementary Figure 6. F4-branch statistic plotted as a heatmap. The tree topology is plotted above, and on the left, every branch of the tree is displayed (including internal branches).

References

Allio R, Schomaker-Bastos A, Romiguier J *et al.* (2020) MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics. *Molecular*

ecology resources.

Anselmo L, Rizzioli B (2022a) Side Threats: Further Possible Effects Of Warming On The High Alpine Narrow Endemic Carabus *Cychroides* (Coleoptera: Carabidae). *Nature Conservation Research. Заповедная наука*, 7, 88–94.

Anselmo L, Rizzioli B (2022b) The small range and the great threat: extinction risk assessment of the narrow endemism *Carabus cychroides* under climate change. *Journal of insect conservation*, 26, 17–27.

Baca SM, Gustafson GT, Alexander AM, Gough HM, Toussaint EFA (2021) Integrative phylogenomics reveals a Permian origin of Adephaga beetles. *Systematic entomology*, 46, 968–990.

Bekchiev R, Antov M, Boyadzhiev P *et al.* (2022) Carabus variolosus (Fabricius, 1787) (Coleoptera: Carabidae) in Bulgaria: rediscovered after 111 years. *Historia naturalis Bulgarica.*

Birky CW Jr (2001) The inheritance of genes in mitochondria and chloroplasts: laws, mechanisms, and models. *Annual review of genetics*, 35, 125–148.

Blaimer BB, Lloyd MW, Guillory WX, Brady SG (2016) Sequence Capture and Phylogenetic Utility of Genomic Ultraconserved Elements Obtained from Pinned Insect Specimens. *PloS one*, 11, e0161531.

Bonacina L, Fasano F, Mezzanotte V, Fornaroli R (2023) Effects of water temperature on freshwater macroinvertebrates: a systematic review. *Biological reviews of the Cambridge Philosophical Society*, 98, 191–221.

Camard A, Leplat J (2004) *Hybrides du genre Carabus*. Magellanes, France.

Campani M, Mulch A, Kempf O, Schlunegger F, Mancktelow N (2012) Miocene paleotopography of the Central Alps. *Earth and planetary science letters*, 337-338, 174–185.

Capon SJ, Stewart-Koster B, Bunn SE (2021) Future of Freshwater Ecosystems in a 1.5°C Warmer World. *Frontiers of Environmental Science & Engineering in China*, 9.

Card DC, Shapiro B, Giribet G, Moritz C, Edwards SV (2021) Museum Genomics. *Annual review of genetics.*

Casale A, Pruser F, Arndt E, Mossakowski D (1998) Phylogenetic relationships in the subgenus Platycarabus MORAWITZ,1886 (Coleoptera: Carabidae: Carabini). *Mus. reg. Sci. nat. Torina, Atti*, 1998, 429–448.

Casale A, Rapuzzi I (2015) Due nuovi ibridi fra specie del genere Carabus Linné, 1758, e nuova località italiana dell'ibrido *C. (Platycarabus) depressus depressus* Bonelli, 1811 x *C. (Platycarabus) fabricii fabricii* Panzer, 1812 (Coleoptera Carabidae). *Rivista piemontese di Storia naturale*, 157–169.

Cavazzuti P, Ghiretti D (2020) *Carabus d'Italia* (NE Scientifiche, Ed,). Natura Edizione Scientifiche, Bologna.

Chernomor O, von Haeseler A, Minh BQ (2016) Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. *Systematic biology*, 65, 997–1008.

Danecek P, Auton A, Abecasis G *et al.* (2011) The variant call format and VCFtools.
Bioinformatics , 27, 2156–2158.

Deuve T (2019) Classification du genre Carabus L., 1758. Liste Blumenthal 2018-2019
(Coleoptera, Carabidae). *Coléoptères*, 25, 33–102.

Deuve T (2021) *Carabus of the World [Holarctic Region]* (Magellanes, Ed,).

Deuve T, Cruaud A, Genson G, Rasplus J-Y (2012) Molecular systematics and evolutionary
history of the genus Carabus (Col. Carabidae). *Molecular phylogenetics and evolution*, 65,
259–275.

Dillon N, Austin AD, Bartowsky E (1996) Comparison of preservation techniques for DNA
extraction from hymenopterous insects. *Insect molecular biology*, 5, 21–24.

de-Dios T, Fontsere C, Renom P *et al.* (2023) Whole-genomes from the extinct Xerces Blue
butterfly can help identify declining insect species. *eLife*, 12.

Drummond AJ, Ho SYW, Phillips MJ, Rambaut A (2006) Relaxed phylogenetics and dating
with confidence. *PLoS biology*, 4, e88.

Duchenne F, Thébault E, Michez D *et al.* (2020) Phenological shifts alter the seasonal structure
of pollinator assemblages in Europe. *Nature ecology & evolution*, 4, 115–121.

Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. *Conservation*
genetics resources, 4, 359–361.

Erbil Ü, Okay AI, Hakyemez A (2021) Late Oligocene—Early Miocene shortening in the Thrace
Basin, northern Aegean. *International Journal of Earth Sciences*, 110, 1921–1936.

Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the
software STRUCTURE: a simulation study. *Molecular ecology*, 14, 2611–2620.

Faircloth BC (2017) Identifying conserved genomic elements and designing universal bait sets to
enrich them. *Methods in ecology and evolution / British Ecological Society*, 8, 1103–1112.

Flouri T, Jiao X, Rannala B, Yang Z (2018) Species Tree Inference with BPP Using Genomic
Sequences and the Multispecies Coalescent. *Molecular biology and evolution*, 35, 2585–
2593.

Fujisawa T, Aswad A, Barraclough TG (2016) A Rapid and Scalable Method for Multilocus
Species Delimitation Using Bayesian Model Comparison and Rooted Triplets. *Systematic*
biology, 65, 759–771.

Gauthier J, Borer M, Toussaint EFA *et al.* (2023) Museomics reveals evolutionary history of
Oreina alpine leaf beetles (Coleoptera: Chrysomelidae). *Systematic entomology*, 48, 658–
671.

Gauthier J, Pajkovic M, Neuenschwander S *et al.* (2020) Museomics identifies genetic erosion in
two butterfly species across the 20th century in Finland. *Molecular ecology resources*, 20,
1191–1205.

Gernhard T (2008) The conditioned reconstructed process. *Journal of theoretical biology*, 253,
769–778.

Gueorguiev VB, Gueorguiev BV (1995) *Catalogue of the Ground-beetles of Bulgaria*

(Coleoptera; Carabidae) (Pensoft Publishers, Ed,). Pensoft Publishers.

Guindon S, Dufayard J-F, Lefort V *et al.* (2010) New algorithms and methods to estimate
maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic*
biology, 59, 307–321.

Hieke F, Wrase DW (2008) Faunistik der laufkäfer bulgariens. (Coleoptera, Carabidae).
Mitteilungen aus dem museum fur naturkunde in Berlin. Deutsche entomologische
zeitschrift, 35, 1–171.

van Hinsbergen DJJ, Schmid SM (2012) Map view restoration of Aegean–West Anatolian
accretion and extension since the Eocene. *Tectonics*, 31.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the
Ultrafast Bootstrap Approximation. *Molecular biology and evolution*, 35, 518–522.

Holderegger R, Thiel-Egenter C (2009) A discussion of different types of glacial refugia used in
mountain biogeography and phylogeography. *Journal of biogeography*, 36, 476–480.

Homburg K, Drees C, Gossner MM *et al.* (2013) Multiple glacial refugia of the low-dispersal
ground beetle Carabus irregularis: molecular data support predictions of species distribution
models. *PloS one*, 8, e61185.

Hristovski S, Mésàros G, Komnenov M, Cvetkovska-Gjorgjievska A (2023) Rediscovery of
Carabus (Hygrocarabus) variolosus nodulosus Creutzer, 1799 on Shar Planina Mt.
Macedonian Journal of Ecology and Environment.

Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary studies.
Molecular biology and evolution, 23, 254–267.

Imura Y, Kim C-G, Su Z-H, Osawa S (1998) An Attempt at the Higher Classification of the
Carabina (Coleoptera, Carabidae) Based on Morphology and Molecular Phylogeny, with
Special Reference to Apotomopterus, Limnocarabus and Euleptocarabus. *Elytra, Tokyo*, 26,
17–35.

Imura Y, Su Z-H, Osawa S (2000) Phylogenetic Relationships in the Division Arciferi
(Coleoptera, Carabidae). *Elytra, Tokyo*, 28, 235–239.

Ishikawa R (1984) Phylogeny and subgeneric classification of the genus Chaetocarabus
THOMSON (Coleoptera, Carabidae). *Kontyu, Tokyo*, 94–109.

Jónsson H, Ginolhac A, Schubert M, Johnson PLF, Orlando L (2013) mapDamage2.0: fast
approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* , 29,
1682–1684.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nature methods*, 14, 587–589.

Kingman JFC (1982) The coalescent. *Stochastic Processes and their Applications*, 13, 235–248.

Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I (2015) Clumpak: a program
for identifying clustering modes and packaging population structure inferences across K.
Molecular ecology resources, 15, 1179–1191.

Kosiński P, Sękiewicz K, Walas Ł, Boratyński A, Dering M (2019) Spatial genetic structure of
the endemic alpine plant *Salix serpillifolia*: genetic swamping on nunataks due to secondary

colonization? *Alpine Botany*, 129, 107–121.

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A (2019) RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* , 35, 4453–4455.

Krsnik E, Methner K, Campani M *et al.* (2021) Miocene high elevation in the Central Alps. *Solid Earth*, 12, 2615–2631.

Kulijer D (2019) New records and distribution of threatened *Carabus (variolosus) nodulosus* Creutzer, 1799 in Bosnia and Herzegovina (Coleoptera: Carabidae). *Acta entomologica slovenica (Ljubljana)*, 27.

Landry B, Bilat J, Hayden J *et al.* (2023) The identity of Argyrialacteella (Fabricius, 1794) (Lepidoptera, Pyraloidea, Crambinae), synonyms, and related species revealed by morphology and DNA capture in type specimens. *ZooKeys*, 1146, 1–42.

Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) PartitionFinder 2: New Methods for Selecting Partitioned Models of Evolution for Molecular and Morphological Phylogenetic Analyses. *Molecular biology and evolution*, 34, 772–773.

Lemmon AR, Emme SA, Lemmon EM (2012) Anchored hybrid enrichment for massively high-throughput phylogenomics. *Systematic biology*, 61, 727–744.

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-bio.GN]*.

Li H, Handsaker B, Wysoker A *et al.* (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* , 25, 2078–2079.

Li C, Hofreiter M, Straube N, Corrigan S, Naylor GJP (2013) Capturing protein-coding genes across highly divergent species. *BioTechniques*, 54, 321–326.

Malinsky M, Matschiner M, Svardal H (2021) Dsuite - Fast D-statistics and related admixture evidence from VCF files. *Molecular ecology resources*, 21, 584–595.

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, 17, 10–12.

Matern A, Desender K, Drees C *et al.* (2009) Genetic diversity and population structure of the endangered insect species *Carabus variolosus* in its western distribution range: Implications for conservation. *Conservation genetics* , 10, 391–405.

Matern A, Drees C, Vogler AP, Assmann T (2010) Linking Genetics and Ecology: Reconstructing the History of Relict Populations of an Endangered Semi-Aquatic Beetle. In: *Relict Species*, pp. 253–265. Springer Berlin Heidelberg.

Mayer C, Dietz L, Call E *et al.* (2021) Adding leaves to the Lepidoptera tree: capturing hundreds of nuclear genes from old museum specimens. *Systematic entomology*, 46, 649–671.

McKenna A, Hanna M, Banks E *et al.* (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research*, 20, 1297–1303.

Minh BQ, Schmidt HA, Chernomor O *et al.* (2020) IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular biology and evolution*,

37, 1530–1534.

Mossakowski D, Bérces S, Hejda R *et al.* (2020) High molecular diversity in *Carabus*
(Hygrocarabus) variolosus and *C. nodulosus*. *Acta zoologica Academiae Scientiarum*
Hungaricae , 66, 147–168.

Müller-Kroehling, S. (2006): Ist der Gruben-Großlaufkäfer *Carabus (variolosus) nodulosus* ein
Taxon des Anhanges II der FFH-Richtlinie in Deutschland? – Waldökologie online 3: 52-
57 *ResearchGate*.

Müller-Kroehling, S. (2014): Remarks on the current situation of *Carabus variolosus nodulosus*
relating to the interpretation of its Habitats Directive status, the 2013 report under that
directive, and its threat level in Germany and Central Europe – Angewandte Carabidologie
10: 97-100 *ResearchGate*.

Nunes R, Storer C, Doleck T *et al.* (2022) Predictors of sequence capture in a large-scale
anchored phylogenomics project. *Frontiers in Ecology and Evolution*, 10.

Osawa S, Su ·-R, Imura Y (2004) *Molecular Phylogeny and Evolution of Carabid Ground*
Beetles (SJ Kk, Ed,).

Patterson N, Moorjani P, Luo Y *et al.* (2012) Ancient admixture in human history. *Genetics*, 192,
1065–1093.

Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double digest RADseq: an
inexpensive method for de novo SNP discovery and genotyping in model and non-model
species. *PloS one*, 7, e37135.

Post RJ, Flook PK, Millest AL (1993) Methods for the preservation of insects for DNA studies.
Biochemical systematics and ecology, 21, 85–92.

Poulakakis N, Kapli P, Lymberakis P *et al.* (2015) A review of phylogeographic analyses of
animal taxa from the Aegean and surrounding regions. *Journal of zoological systematics*
and evolutionary research = Zeitschrift fur zoologische Systematik und
Evolutionsforschung, 53, 18–32.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus
genotype data. *Genetics*, 155, 945–959.

Raxworthy CJ, Smith BT (2021) Mining museums for historical DNA: advances and challenges
in museomics. *Trends in ecology & evolution*.

Rögl F (1997) Palaeogeographic Considerations for Mediterranean and Paratethys Seaways
(Oligocene to Miocene). *Annalen des Naturhistorischen Museums in Wien. Serie A.*
Mineralogie und Petrographie, Geologie und Palaeontologie, Anthropologie und
Praehistorie, 99, 279–310.

Rögl F (1999) Mediterranean and Paratethys; facts and hypotheses of an Oligocene to Miocene
paleogeography; short overview. *Geologica Carpathica (Bratislava)*, 339–349.

Ruppert L-S, Segelbacher G, Staab M, Winiger N (2023) Gauging DNA degradation among
common insect trap preservatives. *Entomologia experimentalis et applicata*.

Sachsenhofer RF, Popov SV, Bechtel A *et al.* (2017) Oligocene and Lower Miocene source
rocks in the Paratethys: palaeogeographical and stratigraphic controls. *Geological Society*,

464, 267–306.

Schmid S, Genevest R, Gobet E *et al.* (2017) HyRAD□X, a versatile method combining exome
capture and RAD sequencing to extract genomic information from ancient DNA. *Methods*
in ecology and evolution / British Ecological Society, 8, 1374–1388.

Schmidt J, Opgenoorth L, Mao K, Baniya CB, Hofmann S (2023) Molecular phylogeny of mega-
diverse *Carabus* attests late Miocene evolution of alpine environments in the Himalayan–
Tibetan Orogen. *Scientific reports*, 13, 1–16.

Schönswetter P, Schneeweiss GM (2019) Is the incidence of survival in interior Pleistocene
refugia (nunataks) underestimated? Phylogeography of the high mountain plant *Androsace*
alpina (Primulaceae) in the European Alps revisited. *Ecology and evolution*, 9, 4078–4086.

Smith SA, Brown JW, Walker JF (2018) So many genes, so little time: A practical approach to
divergence-time estimation in the genomic era. *PloS one*, 13, e0197433.

Sota T, Ishikawa R (2004) Phylogeny and life-history evolution in *Carabus* (subtribe Carabina:
Coleoptera, Carabidae) based on sequences of two nuclear genes. *Biological journal of the*
Linnean Society. Linnean Society of London, 81, 135–149.

Sota T, Takami Y, Ikeda H *et al.* (2022) Global dispersal and diversification in ground beetles of
the subfamily Carabinae. *Molecular phylogenetics and evolution*, 167, 107355.

Sota T, Vogler AP (2001) Incongruence of mitochondrial and nuclear gene trees in the Carabid
beetles Ohomopterus. *Systematic biology*, 50, 39–59.

Sturani M (1962) *Osservazioni e ricerche biologiche sul genere Carabus Linnaeus (sensu lato)*
(Coleoptera Carabidae) (F Pagano, Ed,).

Suchan T, Espíndola A, Rutschmann S *et al.* (2017) Assessing the potential of RAD-sequencing
to resolve phylogenetic relationships within species radiations: The fly genus *Chiastocheta*
(Diptera: Anthomyiidae) as a case study. *Molecular phylogenetics and evolution*, 114, 189–
198.

Suchan T, Pitteloud C, Gerasimova NS *et al.* (2016) Hybridization Capture Using RAD Probes
(hyRAD), a New Tool for Performing Genomic Analyses on Collection Specimens. *PloS*
one, 11, e0151651.

Suchard MA, Lemey P, Baele G *et al.* (2018) Bayesian phylogenetic and phylodynamic data
integration using BEAST 1.10. *Virus evolution*, 4, vey016.

Su Z-H, Imura Y, Zhou H-Z, Okamoto M, Osawa S (2003) Mode of morphological
differentiation in the Latitarsi-ground beetles (Coleoptera, Carabidae) of the world inferred
from a phylogenetic tree of mitochondrial ND5 gene sequences. *Genes & genetic systems*,
78, 53–70.

Tin MM-Y, Economo EP, Mikheyev AS (2014) Sequencing degraded DNA from non-
destructively sampled museum specimens for RAD-tagging and low-coverage shotgun
phylogenetics. *PloS one*, 9, e96793.

Toussaint Fls EFA, Gillett CPDT (2018) Rekindling Jeannel's Gondwanan vision?
Phylogenetics and evolution of Carabinae with a focus on Calosoma caterpillar hunter
beetles. *Biological journal of the Linnean Society. Linnean Society of London*, 123, 191–

207.

Toussaint EFA, Gauthier J, Bilat J *et al.* (2021) HyRAD-X Exome Capture Museomics Unravels
Giant Ground Beetle Evolution. *Genome biology and evolution*, 13.

Turin H, Penev L, Casale A (2003) *The Genus Carabus in Europe: A Synthesis*. Sofia/Leiden.

Tyszecka K, Zając K, Kadej M (2023) Habitat preferences of the mountain population of the
endangered beetle *Carabus variolosus ssp. variolosus* indicate its vulnerability to climate
change. *Global Ecology and Conservation*, 46, e02601.

Westerhold T, Marwan N, Drury AJ *et al.* (2020) An astronomically dated record of Earth's
climate and its predictability over the last 66 million years. *Science*, 369, 1383–1387.

Young AD, Gillung JP (2020) Phylogenomics — principles, opportunities and pitfalls of
big⬜data phylogenetics. *Systematic entomology*, 45, 225–247.

Yule GU (1925) II.—A mathematical theory of evolution, based on the conclusions of Dr. J. C.
Willis, F. R. S. *Philosophical Transactions of the Royal Society of London. Series B,*
Containing Papers of a Biological Character, 213, 21–87.

Zhang C, Mirarab S (2022) Weighting by Gene Tree Uncertainty Improves Accuracy of Quartet-
based Species Trees. *Molecular biology and evolution*, 39, msac215.

1065

Cal. (*Calosoma*) *sycophanta* CBX0022 / FR

'Eucarabi'

C. (*Orinocarabus*) *concolor* CBX0073 / IT

C. (*Carabus*) *granulatus* CBX0043 / FR

C. (*Archicarabus*) *nemoralis* CBX0072 / FR

C. (*Procrustes*) *coriaceus* CBX0042 / FR

Carabus

C. (*Chrysocarabus*) *auronitens* CBX0027 / FR

C. (*Hygrocarabus*) *nodulosus nodulosus* CBX0190 / AT

C. (*Hygrocarabus*) *nodulosus nodulosus* CBX0192 / IT

C. (*Hygrocarabus*) *nodulosus nodulosus* CBX0019 / FR

C. (*Hygrocarabus*) *variolosus variolosus* CBX0193 / SK

C. (*Hygrocarabus*) *variolosus variolosus* CBX0182 / RO

C. (*Heterocarabus*) *marietti muchei* CBX0101 / TK

'Arcifera'

C. (*Chaetocarabus*) *arcadicus arcadicus* CBX0211 / GR

C. (*Chaetocarabus*) *arcadicus arcadicus* CBX0079 / GR

C. (*Chaetocarabus*) *intricatus lefebvrei* CBX0178 / IT

C. (*Chaetocarabus*) *intricatus intricatus* CBX0078 / IT

C. (*Chaetocarabus*) *intricatus intricatus* CBX0195 / FR

C. (*Platycarabus*) *cychroides* CBX0203 / IT

C. (*Platycarabus*) *cychroides* CBX0170 / IT

C. (*Platycarabus*) *cychroides* CBX0080 / IT

C. (*Platycarabus*) *cychroides* CBX0081 / IT

0.005
subst./site

C. (*Platycarabus*) *depressus lucens* CBX0207 / IT

C. (*Platycarabus*) *depressus lucens* CBX0110 / IT

C. (*Platycarabus*) *depressus lucens* CBX0183 / IT

C. (*Platycarabus*) *depressus depressus* CBX0111 / CH

C. (*Platycarabus*) *depressus depressus* CBX0205 / CH

C. (*Platycarabus*) *depressus depressus* CBX0206 / CH

C. (*Platycarabus*) *depressus depressus* CBX0085 / IT

C. (*Platycarabus*) *depressus depressus* CBX0108 FR

C. (*Platycarabus*) *depressus depressus* CBX0088 /FR

C. (*Platycarabus*) *creutzeri baldensis* CBX0092 / IT

C. (*Platycarabus*) *creutzeri rinaldoi* CBX0119 / IT

C. (*Platycarabus*) *creutzeri viridimicans* CBX0120 / AT

C. (*Platycarabus*) *creutzeri kircheri* CBX0106 / AT

C. (*Platycarabus*) *fabricii fabricii* CBX0094 / AT

C. (*Platycarabus*) *fabricii fabricii* CBX0112 / AT

C. (*Platycarabus*) *fabricii fabricii* (**PT** ssp. *fassati*) CBX0095 / SK

C. (*Platycarabus*) *fabricii fabricii* CBX0093 / CH

C. (*Platycarabus*) *fabricii fabricii* CBX0113 / CH

C. (*Platycarabus*) *fabricii fabricii* CBX0171 / SK

C. (*Platycarabus*) *fabricii malachiticus* CBX0172 / RO

C. (*Platycarabus*) *irregularis montandoni* CBX0165 / RO

C. (*Platycarabus*) *irregularis montandoni* CBX0166 / SK

C. (*Platycarabus*) *irregularis bucephalus* CBX0098 / SI

C. (*Platycarabus*) *irregularis bucephalus* CBX0168 / SI

C. (*Platycarabus*) *irregularis irregularis* CBX0115 / AT

C. (*Platycarabus*) *irregularis irregularis* CBX0020 / FR

C. (*Platycarabus*) *irregularis irregularis* CBX0208 / CH

C. (*Platycarabus*) *irregularis irregularis* CBX0194 / HU

C. (*Platycarabus*) *irregularis irregularis* CBX0099 / FR

6X	3X	Minimum genomic coverage
A	B	A/ min. 10 samples / 1'481 loci / 50 taxa
		B/ min. 10 samples / 1'965 loci / 52 taxa
C	D	C/ min. 17 samples / 1'014 loci / 50 taxa
		D/ min. 17 samples / 1'291 loci / 52 taxa
E	F	E/ min. 32 samples / 366 loci / 50 taxa
		F/ min. 32 samples / 478 loci / 52 taxa

■ IQ-TREE SH-aLRT ≥ 80 and UFBoot ≥ 95
■ IQ-TREE SH-aLRT ≥ 80 or UFBoot ≥ 95
■ IQ-TREE SH-aLRT < 80 and UFBoot < 95
■ IQ-TREE Not recovered
* wASTRAL LPP ≥ 0.95
* wASTRAL LPP < 0.95
* wASTRAL Not recovered

○ Ethanol-preserved
▲ Dry-pinned (< 30-year-old)
★ Dry-pinned (≥ 30-year-old)
⬟ Dry-pinned (Unknown age)
PT Paratype

Pleistocene Glaciations

Carabus (Hygrocarabus)
Carabus (Heterocarabus)
Carabus (Chaetocarabus)
Arcifera
Carabus (Platycarabus)

Interspecific Cladogenesis

Mid-Miocene Climatic Optimum

HyRAD-X loci / Dataset A
0   1000   2000

Morphology BPP TR2 Structure

C. nodulosus nodulosus CBX0190 / AT
C. nodulosus nodulosus CBX0192 / IT
C. nodulosus nodulosus CBX0019 / FR
C. variolosus variolosus CBX0193 / SK
C. variolosus variolosus CBX0182 / RO
C. marietti muchei CBX0101 / TK
C. arcadicus arcadicus CBX0211 / GR
C. arcadicus arcadicus CBX0079 / GR
C. intricatus lefebvrei CBX0178 / IT
C. intricatus intricatus CBX0078 / IT
C. intricatus intricatus CBX0195 / FR
C. cychroides CBX0203 / IT
C. cychroides CBX0170 / IT
C. cychroides CBX0080 / IT
C. cychroides CBX0081 / IT
C. depressus lucens CBX0207 / IT
C. depressus lucens CBX0110 / IT
C. depressus lucens CBX0183 / IT
C. depressus depressus CBX0111 / CH
C. depressus depressus CBX0205 / CH
C. depressus depressus CBX0206 / CH
C. depressus depressus CBX0085 / IT
C. depressus depressus CBX0108 FR
C. depressus depressus CBX0088 /FR
C. creutzeri baldensis CBX0092 / IT
C. creutzeri rinaldoi CBX0119 / IT
C. creutzeri viridimicans CBX0120 / AT
C. creutzeri kircheri CBX0106 / AT
C. fabricii CBX0094 / AT
C. fabricii CBX0112 / AT
C. fabricii fabricii CBX0095 / SK
C. fabricii fabricii CBX0093 / CH
C. fabricii fabricii CBX0113 / CH
C. fabricii fabricii CBX0171 / SK
C. fabricii malachiticus CBX0172 / RO
C. irregularis montandoni CBX0165 / ?
C. irregularis montandoni CBX0166 / SK
C. irregularis bucephalus CBX0098 / SI
C. irregularis bucephalus CBX0168 / SI
C. irregularis irregularis CBX0115 / AT
C. irregularis irregularis CBX0020 / FR
C. irregularis irregularis CBX0208 / CH
C. irregularis irregularis CBX0194 / HU
C. irregularis irregularis CBX0099 / FR

| OLIGO. | MIOCENE | P | P |

Geological Time (Ma)

30   25   20   15   10   5   0

■ Ethanol-preserved
■ Dry-pinned (≥ 30-year-old)
■ Dry-pinned (< 30-year-old)
■ Dry-pinned (Unknown age)

Table 1

Dataset	No. of taxa	No. of loci	Min. cov	Min. taxa	Align. length	Missing data	Variable sites	Parsimony informative sites	GC_content
A	50	1,481	6	10	346,687	66.7%	40,743 (11.8%)	19,061 (5.5%)	0.469
B	52	1,965	3	10	458,090	67.9%	59,905 (13.1%)	28,256 (6.2%)	0.468
C	50	1,014	6	17	251,069	59.9%	32,461 (12.9%)	15,871 (6.3%)	0.469
D	52	1,291	3	17	317,337	60.4%	45,834 (14.4%)	22,711 (7.2%)	0.468
E	50	366	6	32	93,904	43.2%	12,566 (13.4%)	6,350 (6.8%)	0.474
F	52	478	3	32	126,457	45.3%	18,843 (14.9%)	9,655 (7.6%)	0.471
G	50	100	6	10	29,855	44.5%	29,855 (17.7%)	2,801 (9.4%)	0.463
H	44	na	3	4	26,201	73.0%	26,201 (100.0%)	13,539 (52.0%)	0.507