Title: Reproductive modes in populations of late-acting self-incompatible
and self-compatible polyploid *Ludwigia grandiflora* subsp. *hexapetala* in
western Europe

Short running title: Mixed reproductive modes in *Ludwigia*

Authors: Solenn Stoeckel[1,2], Ronan Becheler[1,2], Luis Portillo-Lemus[1], Marilyne Harang[1], Anne-Laure
Besnard[1], Gilles Lassalle[1], Romain Causse-Védrines[3], Sophie Michon-Coudouel[3], Daniel J. Park[4],
Bernard J. Pope[4], Eric J. Petit[1], Dominique Barloy[1]

Affiliations

1 DECOD (Ecosystem Dynamics and Sustainability), Institut Agro, IFREMER, INRAE, Rennes, France.

2 IGEPP, INRAE, Institut Agro, Université de Rennes, Le Rheu, France.

3 Plateforme EcogenO, UAR3343 OSUR, Université de Rennes, CNRS, Rennes, France.

4 Melbourne Bioinformatics, The University of Melbourne, Parkville, Australia.

Corresponding author: solenn.stoeckel@inrae.fr

Abstract

Reproductive mode, *i.e.,* the proportion of individuals produced by clonality, selfing and outcrossing in populations, determines how hereditary material is transmitted through generations. It shapes genetic diversity and its structure over time and space, which can be used to infer reproductive modes.

Ludwigia grandiflora subsp. *hexapetala* (*Lgh*) is a partially clonal, polyploid, hermaphroditic, and heteromorphic plant that recently colonized multiple countries worldwide. In western Europe, individuals are either self-incompatible caused by a late-acting self-incompatibility (LSI) system developing long-styled flowers, or self-compatible (SC), with short-styled flowers.

In this study, we genotyped 53 long- and short-styled populations newly colonizing France and northern Spain using SNPs to estimate rates of clonality, selfing and outcrossing. We found that populations reproduced mainly clonally but with a high diversity of genotypes along with rates of sexuality ranging from 10% up to 40%. We also found evidence for local admixture between long- and short-styled populations in a background of genetic structure between floral morphs that was twice the level found within morphs. Long- and short-styled populations showed similar rates of clonality but short-styled populations presented significantly higher rates of selfing, as expected considering their breeding system, and despite the small rates of failure of the LSI system. Within the 53 studied populations, the 13 short-styled populations had fewer effective alleles, lower observed heterozygosity, and higher inbreeding coefficients, linkage disequilibrium and estimates of selfing than what was found in long-styled populations. These results emphasize the necessity to consider the variation of reproductive modes when managing invasive plant species. The overall maintenance of higher genetic diversity with the possibility of maintaining populations clonally in the absence of compatible partners may explain why long-styled individuals seem to be more prevalent in all newly expanding populations worldwide. Beyond *Lgh*, our methodological approach may inspire future studies to assess the reproductive modes in other autopolyploid populations.

Keywords: Partial clonality, Mating system, Late-acting self-incompatibility system, Selfing, Outcrossing, Autopolyploidy, Water primrose

Introduction

Plants reproduce using different mechanisms (i.e., breeding system), including clonality and different types of sexuality (i.e., mating system, including allogamy and selfing; Holsinger 2000). Reproductive mode corresponds to the actual balance between sexual and clonal reproduction, measured by the rate of clonality (De Meeus et al. 2007; Stoeckel et al. 2021a), and, within the proportion of offspring produced by sexuality, the balance between autogamy and geitonogamy (hereafter *selfing*), and allogamy (*outcrossing*), measured by the rate of selfing (Bürkli et al. 2017). Reproductive mode is one of the main drivers of genetic diversity in populations and species (Duminil et al. 2007, Ellegren & Galtier 2016, Glémin et al. 2019). It determines the way hereditary material is transmitted over generations, and thus constrains the range of genetic diversity that can evolve within populations and species (De Meeus et al. 2007, Fehrer 2010, Stoeckel & Masson 2014, Rouger et al. 2016). Its genetic consequences are deep enough to be discernible even in the presence of other affecting factors (Charlesworth 2003). Considering its influence on population genetic diversity and structure, reproductive mode is one of the major biological traits to decipher before interpreting other biological and ecological forces using molecular data (Duminil et al. 2007, Reichel et al. 2016, Bürkli et al. 2017, Orive & Krueger-Hadfield 2021). Given an adequate theoretical framework, population geneticists can use genotypic data to infer reproductive modes in populations (Halkett et al. 2005, Arnaud-Haond et al. 2007, Hardy 2016, Bürkli et al. 2017, Stoeckel et al. 2021a).

Many plant species reproduce using different breeding systems they adapt to different ecological contexts resulting into different reproductive modes (Richards 1997, Charlesworth 2006).

Uniparental reproduction, including clonality and selfing, may help plants to spread into new areas

(Baker's conjecture: Barrett et al. 2008, Pannel et al. 2015). Clonality consists of a parent producing

a new descendant that is a genetic copy of itself with the exception of rare somatic mutations and

mitotic recombinations (De Meeus et al. 2007). It results into an average excess of heterozygotes

compared to Hardy-Weinberg expectations (*i.e.,* negative F_{IS}), large variance of F_{IS} along the genome

and a small excess of linkage disequilibrium, increasing with decreasing population size (Navascues

et al. 2010, Stoeckel & Masson 2014, Stoeckel et al. 2021a). When parents can yield multiple

descendants, it may generate repeated genotypes in populations (Arnaud-Haond et al. 2007). These

effects vary with the rates of clonality and, in turn, can be used to infer the rates of clonality

themselves within populations (De Meeus et al. 2006, Becheler et al. 2020, Arnaud-Haond et al.

2020). Hermaphroditic plants have the possibility to sexually reproduce with themselves (selfing).

Selfing decreases effective population sizes and gene diversity resulting into high probability of allele

fixing (Wright et al. 2008, Roze 2016, Glémin et al. 2019). Selfing also prohibits genetic mixing

between different ancestral lineages, which decreases heterozygosity within individuals (David et

al. 2007, Hardy 2016) and strongly increases linkage disequilibrium between genes along genomes

(Golding & Strobeck 1980, Norborg 2000, Lucek & Willi 2021). Around half of hermaphrodite plants

restrict self-fertilization using a variety of molecular mechanisms grouped under the term "self-

incompatibility (SI) systems" which favour outcrossing within populations (De Nettancourt 2001,

Castric & Vekemans 2004, Gibbs 2014, Steinecke et al. 2022). Outcrossing is overall expected to

increase genetic diversity, to limit allele fixation and to reduce linkage disequilibrium when

compared to selfing, but only if linked to the genes involved in self-recognition (Glémin et al. 2001,

Stoeckel et al. 2006, Navascues et al. 2010). Within the different identified SI systems (Charlesworth

et al. 2005, Franklin-Tong 2008), the late-acting SI system (hereafter LSI) is still poorly studied

despite being identified in multiple taxonomic groups in angiosperms (De Nettancourt 1997,

Gibbs 2014). In LSI systems, self-pollen tubes grow and are only blocked shortly before penetrating

the ovule. Such species may present reduced female fertility due to self-pollen disabling ovules,

favouring the clonal regeneration of populations (Vaughton et al. 2010). LSI systems are also

characterized by low but recurrent failures of the self-recognition system due to its late mechanism,

leading to the production of a low amount of selfed seeds in populations (Seavey & Bawa 1986;

Chen et al. 2012, Gibbs 2014). In contrast, gametophytic and sporophytic self-recognition occurring

very early in the pistil drastically limit self-pollination and thus selfed seeds (Lawrence 1985, Gibbs

2014). However, we do not yet know if these rare selfed seeds contribute to the dynamics of genetic

diversity of LSI populations and species, especially in peripatric conditions or in any situation when

compatible partners may lack. Due to these selfed seeds, LSI system may be relatively ineffective at

driving genetic diversity within populations, contrary to gametophytic and sporophytic SI systems

(Brennan et al. 2002, Stoeckel et al. 2006, Koelling et al. 2011, Busch & Urban 2011). In such

situations, the emergence and maintenance of LSI systems appear as new evolutionary puzzles

among the reproductive systems.

Water primrose, *Ludwigia grandiflora* subsp. *hexapetala* (Hook. & Arn.) Nesom and Kartesz (2000),

hereafter *Lgh*, is an insect pollinated, partially clonal, polyploid, hermaphroditic and heteromorphic

plant. This species is one of the most invasive aquatic plants in the world (Thouvenot et al. 2013).

Lgh is a decaploid species (2n=10x=80 chromosomes), resulting from hybridization of different

ancestral diploid species, some of which are represented more than once in the total genome of

Lgh, which belongs to the genus *Ludwigia* L. section Jussiaea (Hoch et al. 2015, Barloy et al, 2024).

Interestingly, *Lgh* includes an autotetraploid set of chromosomes, shared with *L. peploides* subsp.

montevidensis (2n=2x=16), hereafter *Lpm*, that is clearly distinct from the other ancestral part of

the *Lgh* genome (Barloy et al. 2024). *Lpm* is a self-compatible-only diploid species with only one

common floral morphology (Estes & Thorp 1974, Grewel et al. 2016). *Lgh* presents heteromorphic

flowers corresponding to two floral morphs: L-morph individuals develop long-styled flowers and S-

morph individuals develop short-styled flowers, that cross and result into 100% viable and fertile F1

and F2 descendants (Portillo-Lemus 2021, Portillo-Lemus et al. 2021) while inter-species crosses

only result in a low number of chlorotic and unfertile descendants (Barloy et al. 2024).

All tested L-morph flowers expressed an active LSI in western European populations (Portillo-Lemus

et al. 2022). During the core flowering season (summer), in experimental greenhouse conditions, L-

morph individuals show a stable seemingly insignificant rate of autogamy (around 0.2‰ of the

available ovules) that increases at the end of the flowering season, during autumn, to 1‰, which is

common in LSI systems (Gibbs 2014). Due to the massive blossoming of this species, growing in very

dense populations, these selfed seeds would add up yearly in field populations to thousands of

seeds per square meter (Portillo-Lemus et al. 2021). This pattern of low rates of autogamy, that

increases at the end of the flowering season, may provide the advantage of reproductive assurance

(Goodwillie & Weber 2018). In contrast, all tested S-morph individuals were self-compatible (SC) in

western European populations but in their pistils, pollen tubes of the L-morph growed significantly

faster and were thus advantaged to fertilize ovules when in competition with self-pollen (Portillo-

Lemus et al. 2022). In addition, peripatric *Lgh* populations, including European populations, were

previously reported as exclusively clonal with few clonal lines (Dandelot 2004, Okada et al. 2009).

Recently established populations in France and northern Spain mostly present only one of the two

compatibility modes locally, sometimes with a population of the other type a few to tens of

kilometers away, which may result in effective allogamy (Portillo-Lemus et al. 2021, 2022). All these

different breeding systems make possible very different reproductive modes in populations,

comprising all possible quantitative combinations of mainly clonal, autogamous and allogamous

modes.

Here, we assessed the reproductive modes of 53 invasive Lgh populations across western Europe.

Considering the complex case of *Lgh* in western European populations, we hypothesized that L-

morph populations supposed to express a LSI should present typical genetic footprints of dominant

outcrossing, and perhaps higher rates of clonality due to the local lack of compatible partners and

self-pollen interferences, while S-morph populations supposed to be SC should present higher rates of sexuality prevailed by selfing. To understand the genetic impacts of reproductive modes, we also quantified the covariations of genetic indices and their importance to define the genetic diversity within the 53 genotyped populations, and compared these observations to the theoretical expectations obtained from a Wright-Fisher-like model extended to autotetraploids (Stoeckel et al. 2024). Finally, we took advantage from the fact that recent local populations in France and northern Spain still present only one of the two compatibility modes to compare and interpret the genetic differences found within and between L-morph (LSI) and S-morph (SC) populations, with the aim to assess the influence of LSI on genetic diversity, as similarly tackled in sporophytic self-incompatible and SC Brassicaceae species previously (Koelling et al. 2011).

Materials and methods

Sampling strategy and floral morphology of populations

We collected 795 stems of *Lgh* from 53 locations (52 locations in France and one in Catalonia, Spain), corresponding to an area that spans 580 kilometres east-to-west and 1,100 kilometres north-to-south (Figure SI1). At each location, we collected 15 stems (hereafter, 'individuals') along a linear transect of 40 meters. Along each transect, we randomly collected three stems at coordinates $X_1 = 0m$; $X_2 = 10m$; $X_3 = 20m$; $X_4 = 30m$ and $X_5 = 40m$ within a one meter-square quadrat. The young leaves of each sampled individual were stored after lyophilization until DNA extraction.

We visually identified floral morphologies of flowers found along each transect within the sampled *Lgh* populations. In a previous study on seven populations among the 53 studied here (underlined population names in Figure SI1), one hundred and five sampled individuals resulted to identify a binary distribution of floral morphology with formally-identified self-incompatibility types: all L-morph individuals were LSI typed and all S-morph individuals were SC typed (Portillo-Lemus et al

2021) while all these individuals succeeded to cross and give viable and fertile plants. Interestingly,

these two types of populations spatially distribute in monomorphic populations along different

rivers. We supposed for this study the LSI versus SC status of individuals and populations using their

floral morphologies: L-morph individuals were supposed to develop a LSI system and S-morph

individuals being SC. To support this conjecture, as done in Portillo-Lemus et al. (2021), we checked

the fruitset in each of the 53 sampled and genotyped populations: Low and even no fruitsets were

found in L-morph individuals and populations, while full fruitsets were found in S-morph individuals

and populations, in agreement with our conjecture.

In addition to the crossbreeding results in which all L- and S-morph individuals succeeded to cross,

giving full fruit set and 100% viable first- and second-generation descendants (Portillo-Lemus et al.

2021), we here counted the chromosome numbers on karyotypes of S-morph individuals sampled

in two fruitful populations and of L-morph individuals sampled in five fruitless populations to

validate that L- and S-morph individuals belong to *Lgh*. Between 50 to 150 kilometers separated two

consecutive samples (populations underlined Figure SI1). To prepare the karyotypes, we used the

method detailed in Barloy et al (2024) that already karyotyped a S-morph individual sampled near

the French Atlantic coast. The same method was used in Bou Manobens et al. (2019) to karyotype

a L-morph individual sampled in Catalunya.

Definition of the autotetraploid SNP marker set

As no molecular markers suitable for clonal discrimination were yet available for *Lgh* and *Lpm*, we

generated an original set of SNP markers via RAD-Seq (Baird et al., 2008) using two pools of 15

individuals each, respectively sampled across five *Lgh* and three *Lpm* western European populations.

RAD DNA library generation, sequencing and the analysis pipeline to identify *Lgh* SNP markers were

carried out as described in Delord et al. (2018). In brief, DNA of *Lgh* and *Lpm* were digested by *Sbfl*

restriction enzyme and used to prepare DNA libraries that were then sequenced using an Illumina

HiSeq3000 (150bp paired-end reads). A total of 14,233 and 34,287 RAD-Seq-determined SNPs were

filtered to yield 340 and 326 SNPs, one per aligned sequence, for *Lpm* and *Lgh*, respectively. Design

of primers compatible with Hi-Plex multiplexing was carried out by Melbourne Bioinformatics.

Finally, sixty and fifty SNP markers matched the quality criteria to design a Hi-Plex set of SNPs for

Lpm and *Lgh*, respectively (Hammet et al. 2019). In our study, we only considered polymorphic SNPs

that were shared between *Lgh* and *Lpm* to be sure they belong to the tetraploid part of *Lgh* genome

derived from *Lpm* (Barloy et al. 2024). We finally kept this set of 36 polymorphic and stable SNP

markers to genotype *Lgh* samples and analyze genetic diversity in each sampled population (primer

sequences are openly listed in Stoeckel et al. 2023).

To verify that this set of SNPs was really tetraploid, we computed the Akaike's information criterion

from the maximum likelihood of the best genotype considering the distribution of sequenced allele

countings among individuals and markers as a function of the ploidy level, using a similar approach

to that proposed by Burnham & Anderson (2002).

$$AIC(ploidy) = \sum_{l=1}^{l=36} \sum_{i=1}^{i=795} -2\log \mathcal{L}\left(genotype_{i,l} \middle| ploidy, data_{i,l}\right) + 2K$$

where $data_{i,l}$ is the distribution of sequenced alleles among A, C, G and T within individual i at locus

l, and K is the number of possible genotypes given the ploidy and the four possible alleles (A, C, G

and T). The likelihood of the possible genotypes $\mathcal{L}\left(genotype_{i,l} \middle| ploidy, data_{i,l}\right)$ follows a

multinomial distribution of the sequenced allele countings distributed between the four different

possible nucleobases (A, C, G and T).

DNA extraction, SNP marker production and genotyping

To genotype the 795 individually-tagged samples with these 36 SNPs, we extracted DNA from 25 to

30 mg of dried young leaf tissues. The DNA extractions were performed using the NucleoSpin Plant II

from MACHEREY-NAGEL kit, following the manufacturer's recommendations. We used L1 buffer as

lysis buffer. All individuals were genotyped from a solution of 20ng/µl of *Lgh* DNA, using a modified

version of the Hi-Plex protocol (Hammet et al. 2019; Besnard et al. 2023). Hi-Plex is an amplicon

sequencing technique (*sensu* Meek & Larson 2019) in which all SNPs are co-amplified in a multiplex

reaction before Illumina or Ion-Torrent sequencing. Here, we used Illumina. Intermediate steps

include dual indexing of individual samples used for demultiplexing. Reads were then assigned to

loci by aligning them to reference sequences with BWA-MEM 0.7.15-r1140 (Li & Durbin 2010) and

alleles were counted with Samtools 1.9 (Danecek et al. 2021).

Allele dosage

The posterior probabilities of each single SNP genotype within each individual (hereafter *single SNP*

genotype) were computed using the likelihood of all possible genotypes considering a multinomial

distribution of the number of times each nucleobase was genotyped at one SNP marker within one

individual. In order to obtain the most confident dataset possible, we only assigned a genotype

when its posterior probability of allele dosage exceeded 70%. When one SNP within one individual

presented a posterior probability of allele dosage equal or lower than 70%, we assigned and

analysed it afterward as a missing genotype.

Genotypic and genetic descriptors

We expected that populations reproducing clonally may yield repeated multi-locus genotypes

(MLGs, i.e., the same genotype at all the 36 SNPs found in multiple individuals). By possibly

producing these repeated genotypes and by varying the relative distribution of the number of

samples of each of these distinct genotypes, rates of clonality impact genotypic richness and

evenness in populations (Halkett et al. 2005; Arnaud-Haond et al. 2007).

We measured genotypic richness using the R index (Dorken & Eckert 2001, Arnaud-Haond et al.

2005), which is defined as:

$R = \frac{G-1}{N_g-1}$

where G is the number of distinct genotypes (genets) and N_g is the number of genotyped individuals.

We also measured genotypic evenness as the parameter β of the Pareto distribution, which

describes the slope of the power-law inverse cumulative distribution of the size of MLGs (Arnaud-

Haond et al. 2007):

$N \geq X = aX - \beta$

where N≥X is the number of sampled ramets belonging to genets containing X or more ramets in

the sample of the population studied, and the parameters *a* and *β* are fitted by regression analysis.

Low β-values indicate the dominance of one or few clonal lineages. Pareto β is less biased by the

sampling effort than clonal richness R (Stoeckel et al. 2021a). In our sample of 15 individuals, we

expected a population with only one repeated MLG to present a Pareto β value of 0.03 and a value

of 3 in samples with no repeated genotypes. β < 2 indicates a population reproducing with high rates

of clonality (greater than 0.8 to 0.9). For a sample size of 15 individuals, 2 < β < 3 indicates

intermediate to low rates of clonality (0.6 to 0.8). A β at 3, its maximum value, is indicative of a

mainly sexual population, with rates of clonality ranging from zero to 0.6, depending on the other

genetic indices.

Rates of clonality and selfing confer different effects on the range of within-population genetic

polymorphism as well as on probabilities of genetic identities within individuals. We thus also

estimated expected and observed heterozygosity H_E and H_O, allelic richness A_E, inbreeding

coefficient F_{IS}, and linkage disequilibrium within each sampled population.

The Wright (1931, 1949) inbreeding coefficient F_{IS} accounts for intraindividual genetic variation at

one locus as a departure from Hardy-Weinberg assumptions of the genotyped populations. We

computed one F_{IS} value per locus per population. We then reported the mean values of F_{IS} (MF_{IS})

and the inter-locus variance of F_{IS} (VarF$_{IS}$) for each population. Both MF$_{IS}$ and VarF$_{IS}$ are very

informative about the underlying reproductive systems (Stoeckel et al. 2006, David et al. 2007).

Clonality makes the MF$_{IS}$ values tend toward -1 along with high interlocus variance (Stoeckel &

Masson 2014, Stoeckel et al. 2021a). A moderate amount of sexual reproduction results in MF$_{IS}$

values around 0 (Balloux et al. 2003). VarF$_{IS}$ varies with rates of clonality, from very limited variance

expected in sexual populations to high variance in very clonal populations (Stoeckel & Masson 2014,

Stoeckel et al. 2021a). Positive MF$_{IS}$ values are expected in populations reproducing using

consanguinity and selfing (Castric et al. 2002, David et al. 2007). All sexual reproductive modes,

allogamous and autogamous, result in low VarF$_{IS}$ within a population as massive recombination

tends to homogenize intra-individual genetic identities along the genomes (Stoeckel et al. 2021a).

We measured linkage disequilibrium over all markers using the unbiased multilocus linkage

disequilibrium index, $\bar{r}_d$ (Agapow & Burt 2001). This mean correlation coefficient (*r*) of genetic

distances (*d*) between unordered alleles at *n* loci ranges from 0 to 1. This metric has the advantage

of limiting the dependency of the correlation coefficient on the number of alleles and loci, and it is

well suited to measure linkage disequilibrium in partially clonal populations (De Meeus & Balloux

2004, De Meeus et al. 2006). In general, LD is only slightly affected by clonality, except when

clonality is high (c > 0.9) and/or when genetic drift dominates over mutation rate (*e.g.,* when

population sizes are small N<50 compared to u=0.001; Navascues et al. 2010, Stoeckel et al. 2021a).

In contrast, inbreeding and selfing are efficient processes for quickly generating strong LD, after only

few generations. Finally, we measured genetic differentiation between populations of LSI, between

populations of SC and between LSI and SC populations using ρ$_{ST}$, an index adapted to study

autopolyploid populations (Ronfort et al. 1998). All these indices were computed using GENAPOPOP

(Stoeckel et al. 2024), a software dedicated to analysing genetic diversity and differentiation in

autopolyploid populations genotyped with confident allele dosage and reproducing through all

possible rates of clonality and selfing.

We also used SPAGEDI (v1.5, Hardy & Vekemans 2002) to estimate rates of selfing within

autopolyploid populations genotyped with confident allele dosage. This approach infers rate of

selfing (Sg) from identity disequilibrium coefficients (g_{2z} estimator), assuming that populations

reproduce by self-fertilization and random mating and are at inbreeding equilibrium (David et al.

2007 for diploids, Hardy 2016 for autopolyploids). Identity disequilibrium coefficients are measured

as the correlation in heterozygosity of distinct loci within the genome, and present the advantage

of being more robust to null alleles and genotyping errors than raw F_{IS} (David et al., 2007). The effect

of partial clonality on g_{2z} estimator is not yet defined. We thus considered for this study that clonality

would only marginally impact identity disequilibrium coefficients, the possibility to be at the

inbreeding equilibrium and the corresponding estimates of selfing. A synthesis of all these

hypotheses based on previous knowledge is provided in Table SI1.

Ranking populations by clonality and selfing

We proposed and computed a synthesis index (Σclon) calculated from Pareto β and $VarF_{IS}$ to rank

the studied populations from the less to the more clonal. These two population genetic indices are

known to vary with rates of clonality being unbiased in samples (Stoeckel et al. 2021a). To avoid

issues of scaling as the range of values of these descriptors are different by several orders of

magnitude, Σclon is computed in each population i as the sum of Pareto $\beta(i)$ and $VarFIS(i)$ values

that were previously normalized over the whole dataset, respectively $z_\beta(i)$ and $z_{VarFIS}(i)$.

309
$$z_\beta(i) = 1 - \frac{\beta(i) - min(\mathrm{B})}{max(\mathrm{B}) - min(\mathrm{B})}$$

310
$$z_{VarFIS}(i) = \frac{VarFIS(i) - min(\mathrm{VF})}{max(\mathrm{VF}) - min(\mathrm{VF})}$$

311
$$\Sigma clon(i) = {}^1\!/_2 \left[z_\beta(i) + z_{VarFIS}(i) \right]$$

where $\beta(i)$ and $VarFIS(i)$ are the measured Pareto β and variance of F_{IS} over loci in the population

i, and where $\mathrm{B} = \{\beta(1), \beta(2), \dots, \beta(55)\}$ and $VF = \{VarFIS(1), VarFIS(2), \dots, VarFIS(55)\}$ are the

respective sets of Pareto β and VarF$_{IS}$ over loci in the 53 populations used to obtain their minimum

(min) and maximum (max) values.

Using the same approach as for Σclon, we computed a synthesis index (Σself) calculated from $\bar{r}_d$,

MF$_{IS}$ and Sg to rank the studied populations from the less to the more selfed.

$$\Sigma\text{self}(i) = \frac{1}{3}\left[z_{\bar{r}_d}(i) + z_{\text{MFIS}}(i) + z_{\text{Sg}}(i)\right]$$

Statistical data analysis

First, to better understand the correlation between population genetic indices and estimates of

reproductive modes from a large field dataset and to compare with the theoretical correlations

obtained from simulations (Stoeckel et al. 2024), we computed a Principal Component Analysis

(PCA) to comprehend the covariations of genetic indices, the correlations and the redundancies

between the 17 genetic diversity indices (described above: G, R, D, Pareto β, $\bar{r}_d$, Ae, varAe, He,

varHe, Ho, varHo, MF$_{IS}$, varF$_{IS}$, PIDsib, PIDu, Sg, SE.Sg) measured among the 53 sampled populations,

and their link with reproductive modes including self-compatibility. We reported the amount of

variation retained by the first two principal components and the correlation circle on which we

plotted predictions of Σclon and Σself as supplementary variables. We also reported the score plot

to visualize how L-morph (LSI) and S-morph (SC) populations distribute along the principal

components of population genetic diversity. To avoid scaling issues, all descriptors were normalized

before analysis.

We then analyzed the relationship between genetic diversity indices and their relationships with

Σclon and Σself measured in the 53 sampled populations using Kendal partial rank-order correlation

tests. We reported the corresponding matrices of correlation.

To detect differences in distribution of population genetic indices, Σclon and Σself, among floral

morphs, we computed non-parametric Kruskal-Wallis tests that do not make any assumptions about

the type of distribution and about homogeneity of variances between the tested distributions.

When needed, we used post-hoc pairwise tests for multiple comparisons of mean rank sums

(Nemenyi's test).

All statistical tests were computed using PYTHON 3.11, SCIPY.STATS 1.9.3 (Virtanen et al. 2020) and

SCIKIT-POSTHOCS (Terpilowski 2019), except the PCA that was performed using R v4.2.2 and the library

FACTOMINER (Lê et al. 2008).

Results

Among the 53 populations, we found 40 populations with only L-morph individuals and 13

populations with only S-morph individuals (Table SI2, Figure SI1). Karyotypes of L-morph plants from

two populations and of S-morph plants from five populations all presented the same number of

chromosomes (2n=80, Figure SI2) confirming that L and S-morph individuals in France and L-morph

individuals in northern Spain (Bou Manobens et al. 2019) belong to *L. grandiflora* subsp. *hexapetala*

(2n=10X=80, Barloy et al. 2024). Akaike's information criterion on the distribution of allele counting

over all our data supported tetraploidy as the best ploidy level for this set of 36 SNPs (Figure SI3),

as expected from Barloy et al (2024).

Allele dosage and missing genotypes

Within the 795 individuals genotyped at 36 SNPs (resulting in a total of 28,620 single-SNP

genotypes), 99.97% (28,612) of SNPs were genotyped with posterior probability of allele dosage

superior to 70% (Table SI3). 785 individuals were genotyped with a full set of 36 SNPs with posterior

probabilities of allele dosage higher than 70%. Ten individuals distributed in nine populations

showed one of their SNP markers with posterior probabilities of allele dosage equal or lower than

70%, that we assigned therefore as missing genotypes.

Statistical power of the developed SNP marker set

Over the 36 polymorphic SNPs, we found an effective number of alleles (A_E) of 1.36 per SNP over all populations (Table SI2). Among populations, mean A_E values over the 36 SNPs were homogenous, ranging from 1.22 to 1.55 (median=1.34). We, however, found large standard deviation of A_E between SNPs within populations, ranging from 1.6 to 2.3 (median=2.1). Some SNPs were apparently fixed in some populations while polymorphic at the scale of the whole dataset. When not fixed, gene diversity H_E in polymorphic SNPs ranged from 0.14 to 0.33 (median=0.2).

These 36 SNPs in the autotetraploid part of *Lgh* would theoretically allow $4^{36}=4.7\times10^{21}$ different possible MLGs considering the four possible nucleobases and $2^{36}=6.9\times10^{10}$ different possible MLGs assuming two possible nucleobases per locus. Considering allele frequencies in the sampled populations, the probabilities of identities under panmixia ranged from 8.5×10^{-12} to 5.2×10^{-5} (median=2.1×10^{-7}) and the unbiased probabilities of identity between sibs P_{ID-SIB} ranged from 4.2×10^{-6} to 8.3×10^{-3} (median=5.6×10^{-4}; Table SI2). We then considered that the SNP set we used to genotype the 795 sampled individuals via Hi-Plex method showed sufficient statistical power to distinguish between true MLGs, and that individuals with identical MLGs were true repeated genotypes (ramets) of a clonal lineage (a genet).

Genetic and genotypic diversity

Across populations, we identified a total of 462 distinct MLGs (genets) within the 795 sampled individuals genotyped. Among them, we found 404 genets (88%) with a single ramet and only 58 genets (12%) with more than two ramets (Table SI4). Forty-eight genets had two to seven ramets distributed over one to six populations (median=2), seven genets with 10 to 33 ramets distributed over two to 17 populations (median=9), and one large genet of 99 ramets distributed over 24 populations (Figure 1).

Within populations, we found from three to 15 different genets (median=12) per population among

the 15 sampled individuals (Table SI4), implying the clear occurrence of repeated genotypes but also

a wide diversity of genets within most populations (Figure 1). Accordingly, genotypic richness (R)

ranged from 0.14 to 1 (median=0.79). The genotypic evenness, Pareto β, ranged from 0.056 to 3

(median=1.478, Table SI2).

Observed heterozygosity H_O was also high in most populations, slightly above expected

heterozygosity, ranging from 0.13 to 0.38 (median=0.26). The mean standard inbreeding

coefficients (MF_{IS}) averaged over all genotyped loci within populations were negative in all 53

populations and ranged from -0.33 to -0.126 with a very negative median of -0.274 (Table SI2).

Variances of F_{IS} between loci within population were very high, ranging from 1.75 to 36.60

(median=27.57). All these measures argued for reproductive modes implying high rates of clonality.

Estimates of selfing (S_g) were overall low with a handful of high values, ranging from 0 to 0.61

(median=0.06). Fifteen populations (28%) were estimated with no selfing. Nineteen populations

(36%) showed non-zero estimates under 0.1, 13 populations (25%) between 0.1 and 0.29, and six

populations (11%) with estimates between 0.42 and 0.61.

Finally, linkage disequilibria within populations between genotyped SNPs were overall low, with r_d

values ranging from 0 to 0.51 with a median value of 0.12. Forty-two populations (79%) were under

0.25, and only eight populations (21%) showed linkage disequilibrium between 0.25 and 0.51.

Analyses of covariations between population genetic indices

The first two components of the principal component analysis on the values of genetic diversity

found in the 53 genotyped populations accounted for 78.4% of the total variance between

populations (Figure SI4). Non-parametric Kendall partial rank-order correlations between genetic

indices (Table SI5) and correlations on the first two principal components from the 17 population

genetic indices measured showed three non-collinear groups of associated genetic indices (Figure

SI4.A) that are very similar to the theoretical groups of population genetic indices expected to covary

with different rates of clonality, selfing and outcrossing in autopolyploids (Stoeckel et al. 2024). A first cluster regrouped G, D*, R, Pareto β and $VarF_{IS}$, indices that are known to be sensitive to clonality (Stoeckel et al. 2021a) but also VarHe, VarHo, and VarAe. This cluster largely explains the first dimension (50.2% of the total variance) of the PCA and was collinear to Σ_{CLON} (also see Figure SI4.B). As expected under partial clonality, $VarF_{IS}$ was negatively correlated to genotypic diversity indices (R and β; Figure SI5). The second cluster regrouped P_{ID-SIB}, P_{ID-u}, H_E, A_E and H_O, indices that are linked to the general genetic diversity of populations (Figure SI4.A). The third cluster regrouped S_g, $SE.S_g$, MF_{IS} and r_D, indices that are usually used to identify, rank and estimate rates of selfing versus outcrossing in sexual populations (Castric et al. 2002, Bürkly et al. 2017). The second dimension of the PCA (23.4% of the total variance) was mostly correlated to Ho, Sg, He and r_D which was collinear to Σ_{SELF} (Figure SI4.C). The clusters of genetic indices were corroborated by the correlation between Σ_{CLON} and genetic indices (Figure SI5) and Σ_{SELF} and genetic indices (Figure SI6). The correlation across populations between Σ_{CLON} and Σ_{SELF} was negative and highly significant (r_s = -0.66; $p < 0.001$; Figure 2). This correlation within LSI populations was negative and highly significant (r_s = -0.65; $p < 0.001$) while it was nonsignificant in SC populations (r_s = -0.32; p =0.289).

Differences in genetic diversity and structure between L- and S-morph populations

The number of ramets per genet was similar in L-morph (LSI) and S-morph (SC) populations (H=1.85, p=0.173, Figure 1 & SI7), as were their genotypic richness (R, H=1.23, p=0.267) and evenness (Pareto's β, H=1.60, p=0.206; Table SI2). The distributions of other population genetic indices related to clonality were also not significantly different between in L-morph (LSI) and S-morph (SC) populations ($VarF_{IS}$: H=2.94, p=0.086; P_{ID-SIB}: H=1.98, p=0.160 and Σ_{CLON} H=2.28, p=0.131; Figure 3). Mean observed heterozygosity (Ho, H=5.27, p=0.022) and their variances (VarHo, H=13.85, p<0.001), mean effective number of alleles per locus (Ae, H=5.13, p=0.024) and its variance over

locus (VarAe, H=9.49, p=0.002), MF$_{IS}$ (H=11.85, p<0.001), linkage disequilibrium (r$_D$, H=5.55, p=0.018), estimate of selfing rates (Sg, H=12.39, p<0.001) and its standard error over loci (SESg, H=9.58, p=0.002) significantly differed between in L-morph (LSI) and S-morph (SC) populations. S-morph (SC) populations, compared to L-morph (LSI), showed lower mean observed heterozygosity (medians, L-morph: 0.262, S-morph: 0.210) and less variance between loci (medians, L-morph: 2.400, S-morph: 1.859), lower effective number of alleles (medians, L-morph: 1.341, S-morph: 1.283) and less variance between loci (medians, L-morph: 4.419, S-morph: 3.580), higher linkage disequilibrium (medians, L-morph: 0.113, S-morph: 0.152), higher estimates of selfing rate (medians, L-morph: 0.045, S-morph: 0.175) even if with higher standard error between loci (medians, L-morph: 0.058, S-morph: 0.156) and higher Σ$_{SELF}$ (medians, L-morph: 0.222, S-morph: 0.495).

Genetic differentiation (ρ$_{ST}$) among pairs of S-morph (SC) populations (median value of ρ$_{ST}$=0.27) was comparable to the genetic differentiation found among pairs of L-morph (LSI) populations (median value of ρ$_{ST}$=0.26; H=319.37, post-hoc p=0.322; Figure 4). The median of ρ$_{ST}$ between pairs of L-morph (LSI) and S-morph (SC) populations reached 0.65. The distribution of inter-morph ρ$_{ST}$ values differed both from the distribution of pairs of intra-L-morph (LSI) ρ$_{ST}$ values (post-hoc p<0.001) and from the distribution of pairs of intra-S-morph (SC) ρ$_{ST}$ values (post-hoc p<0.001). The minimum spanning tree of genetic distances between individuals computed with *GenAPoPop* showed quite clustered distributions of L-morph (LSI) and S-morph (SC) individuals but also with clear evidence of admixtures between their lineages (Figure 5).

Discussion

We used population genetics on autotetraploid SNPs with allele dosage to assess the rate of

clonality, of selfing and of outcrossing (i.e., reproductive mode) of 53 populations of *Ludwigia*

hexapetala subsp. *grandiflora* (*Lgh*) recently colonizing western European watersheds.

To achieve this goal, we developed reproducible codominant molecular markers (SNPs) that enabled

allele dosage-based genotyping of sampled individuals. We then used tailored computational

analyses to perform confident population genetic analyses in autopolyploids. These two

methodological steps allowed solving the remaining challenges to perform population genetic

analyses in autopolyploids (Dufresne et al. 2014), including *Lgh*. The resulting framework can be

applied to any autopolyploid species.

Lgh develops two floral morphs respectively associated with a Late-acting self-incompatible system

(L-morph) and a self-compatible system (S-morph, Portillo-Lemus et al. 2021, 2022). Interestingly,

the sampled populations in western Europe showed either L- or S-morph resulting in two groups of

sampled populations: one group of 40 L-morph (LSI) populations and one group of 13 S-morph (SC)

populations. They live in similar ecological conditions (Portillo-Lemus et al. 2021) which thereby

provides a rare opportunity to characterize the genetic consequences of an LSI system compared to

a group of SC populations in the same species and ecological context.

The five L-morph and the two S-morph water primrose populations we karyotyped, separated by 50

to 150 km from one to the other in France, all had 80 chromosomes corresponding to the species

Ludwigia grandiflora subsp. *hexapetala*. We didn't find yet individual with 48 chromosomes

corresponding to *Ludwigia grandiflora* subsp. *grandiflora*. These results agree with previous

observations of Dandelot et al (2005) and Barloy et al. (2024) in France, Bou Manobens et al. (2019)

in northern Spain and Armitage et al. (2013) in Great Britain.

Most of the genetic variance explained by clonality and selfing among *Lgh* populations

The correlations of population genetics indices on the 53 genotyped *Lgh* populations (Figure SI4)

were congruent with previous theoretical predictions on the variations of genetic indices with

different rates of clonality and selfing for autopolyploid populations (Stoeckel et al. 2024). Identical

covariations were already predicted by Wright-Fisher-like models adapted for diploid (Stoeckel et

al. 2021a) and haplodiplontic life-cycles (Stoeckel et al. 2021b), and validated respectively using

multiple field populations of marine phanerogams for diploids (Arnaud-Haond et al. 2020) and of

brown, green, red algae and mosses for haplodiplontics (Krueger-Hadfield et al. 2021). All these

predictions and observations argue for the primary importance of reproductive mode, especially of

high clonality, to drive variation in genetic diversity within populations (Duminil et al. 2007). They

support the methodological importance of accurate assessment of rates of clonality and of selfing

before starting to interpret genetic diversity.

Dominant clonality within the western European *Lgh* populations

Previous field and lab observations of peripatric *Lgh* populations reported massive production of

dispersing vegetative propagules, rapid expansions of patches, and an important capacity for

spontaneous cutting-planting (Dandelot et al. 2005, Thouvenot et al. 2013, Grewell et al. 2016,

Skaer-Thomason et al. 2018a,b). Okada et al. (2009) genotyped around 800 individuals sampled in

27 *Lgh* populations in Californian wetlands using a set of eight AFLP markers and reported an

extremely reduced clonal diversity: 95% of the samples had the same genotype and 18 populations

over 27 (67%) supported a single AFLP-genotype. All over the UK, only two haplotypes on 14

sampled stems were found using chloroplast sequences (Armitage et al. 2013). These haplotypes

were even shared with some *Lgh* samples invading California. With no measure of the probability

of identity to assess the marker set, these results could be due to the lack of resolution of the

markers used resulting in an artificially elevated measure of clonality (Waits et al. 2001, Villate et al.

2010). In any case, all these studies concluded that invasive populations of *Lgh* in Europe and in USA

reproduce by exclusive clonality, with a very narrow base of ancestral clones or being monoclonal

(Dandelot et al. 2005, Okada et al. 2009, Thouvenot et al. 2013, Grewell et al. 2016).

In our study, we chose to develop and use SNPs rather than AFLP markers. The latter can be

challenging in terms of precisely sizing fragments, leading to suboptimal reproducibility, particularly

across different platforms (Fry et al. 2009). The 36 SNPs we developed are easily reproducible,

accurate enough to distinguish between offspring of sibling mating as evidenced by the probabilities

of identity we obtained, and less expensive.

With these SNPs, we found typical genetic signatures of high clonality, including clear occurrence of

replicated genotypes and mean negative F_{IS} values with high interlocus variances, in all the 53

western European populations we genotyped. Globally, gene diversities measured in these invasive

Lgh populations were in the higher range of values commonly found for SNPs (Fischer et al. 2017,

Schmidt et al. 2021). Such levels of gene diversity are in line with strong rates of clonality that buffer

the loss of alleles due to genetic drift (Reichel et al. 2016, Stoeckel et al. 2021b). Among the 58 MLGs

with replicates, seven MLGs were found in a single population and 51 MLGs were found in multiple

sites. Clonal propagules of *Lgh* are known to disperse carried by river current and by birds (zoochory:

seedlings have the ability to stick to feathers; Grewel et al. 2016). Clonal reproduction by rhizomes

at the local scale and by the dispersal of clonal propagules would thus be the main source of

population growth and invasive spread of *Lgh* in western Europe.

Our results also report a previously-underestimated genotype diversity in the invasive populations

in western Europe. We found 462 distinct MLGs (genets) out of 795 individuals within these

populations and the large majority of these MLGs (404, thus 88%) were only sampled once in the 53

populations. Our results are still congruent with Dandelot (2004) unpublished measures obtained in

three Mediterranean populations using inter-simple sequence repeats (ISSRs). This data reports

three and seven genotypes over 11 sampling units within two *Lgh* populations and two genotypes

over nine sampling units in a third population, all sampled in the south-east of France. These results

could reveal contrasted situations between *Lgh* populations introduced in the USA and continental

Europe.

Between 10 and 40% of sexual reproduction in *Lgh* populations in western Europe

Beyond the qualitative indication of dominant clonality, we then aimed at estimating the rates of

sexuality in each of these 53 populations. All showed small to medium values of linkage

disequilibrium between SNPs. Such values are expected in highly but non-exclusive clonal

populations with large population sizes (Navascues et al. 2010, Stoeckel et al. 2021a). Thirty-two

populations showed Pareto β values of their distributions of clonal sizes under 2, which are only

found in theoretical populations with rates of clonality higher than 0.8 (Stoeckel et al. 2024). All 53

populations showed mean negative F_{IS} with high interlocus variance, with values observed in

theoretical populations with rates of clonality higher than 0.6 but under 0.9 (Balloux et al. 2003,

Stoeckel & Masson 2014, Reichel et al. 2016, Stoeckel et al. 2024). All these values of population

genetic indices were consistent with the interpretation that *Lgh* populations in western Europe must

reproduce with effective rates of clonality between 60% and 90%, thus with 10% to 40% sexuality

(De Meeus et al. 2006, Arnaud-Haond et al. 2020, Stoeckel et al. 2024). These estimates were also

supported by the local evidence of recombination between clonal lineages, and even between L-

morph (LSI) and S-morph (SC) lineages at the scale of a watershed (Figure 5). The diversity of

genotypes detected in western Europe is thus indicative of rare but significant local sexual events

rather than of a large clonal diversity that would have maintain and propagate by exclusive clonality

since its introduction. Our results newly advocate that sexual reproduction should not, therefore,

be overlooked in these invasive populations, especially in management plans.

Genetic consequences of LSI compared to self-compatible populations

The late-acting SI system remains one of the less studied breeding systems among the mechanisms

favouring outcrossing in plants (Gibbs 2014). Its efficiency to favour outcrossing and its

consequences on genetic diversity within populations, especially considering its low but common

failures, were not yet deciphered and not yet compared to SC populations in the same ecological

conditions, as already previously explored for gametophytic and sporophyte SI systems (Busch 2005,

Koelling et al. 2011).

The maintenance of SI systems is one of the most intriguing evolutionary puzzles (Porcher &

Lande 2005). Indeed, SI systems are fated to breakdown because SC individuals present the

advantage of reproductive assurance when compatible partners are limited, especially in peripatric

conditions (Eckert et al. 2006). This advantage is even absolute when no compatible partners are

available within pollination range, as we commonly found in *Lgh* populations in western Europe

(Figure SI1). Conversely, outcrossing imposed by SI systems decreases the probability of expressing

deleterious mutations in descendants as compared to selfing with the same genetic background

(Rice 2004, Navascues et al. 2010).

We found that the 40 populations with L-morph individuals (LSI) had a higher number of effective

alleles, higher gene diversity, higher observed heterozygosity and less linkage disequilibrium than

the 13 populations with S-morph individuals (SC; Figure 3).

These genetic differences are very unlikely to directly result from the consequences of the LSI

system hitchhiked to the whole genome in L-morph individuals and populations. Indeed, it would

imply either that all the 36 SNPs would be physically linked to the genes under negative frequency-

dependent selection coding for the LSI or that *Lgh* outcrossed for many generations in small

population sizes (Glémin et al. 2001, Navascues et al. 2010). *Lgh* in western Europe develops

including far more than thousands of stems per local population (Portillo-Lemus et al. 2021) and the

linkage disequilibrium values also argued for large effective population sizes. We found similar

estimated rates of sexuality and of clonality between L-morph and S-morph populations, using Σ_{CLON}

and all indices sensible to clonality (R, Pareto Beta and VarFis; Figure 3). We however found

significant difference in estimates of selfing between L-morph (LSI) and S-morph (SC) populations,

and in Σ_{SELF} and in all indices sensible to selfing (Ae, linkage disequilibrium, Ho, mean Fis) revealing

typical signatures of higher selfing rates in S-morph populations. Hardy (2016) method estimated a

median selfing rate (Sg) of 0.18 in S-morph populations versus 0.05 in L-morph populations.

Consequently, the genetic differences we found between S-morph and L-morph populations may

thus rather be due to the effects of selfing impacting S-morph (SC) populations than due to

outcrossing protecting the loss of genetic diversity or rates of clonality in L-morph (LSI) populations.

Advantages of dominant clonality with preferential allogamy in invasive *Lgh* populations

Uniparental clonal reproduction (including clonality and selfing) may help the demographical

maintenance of plants spreading out of their native range with limited or even without compatible

or less related sexual partner at pollination distance (Baker's conjecture: Barrett et al. 2008, Pannel

et al. 2015). If they rather reproduce using selfing, their descendants increase the probability to

express inbreeding depression and to lose heterozygosity. Some invasive populations develop with

low genetic diversity (He et al. 2024), questioning on the biological and environmental factors that

may explain their success (i.e., the genetic paradox of invasions, Allendorf & Lundquist 2003). But

many other plants spread out of their native ranges with substantial genetic diversity and using

reproductive modes that favour outcrossing (Roman & Darling 2007, Forsman 2014), like *Lgh*

populations in western Europe, mostly when developing in harsh and stressful conditions (Fox &

Reed, 2011) or when the costs of inbreeding depression expressed by selfing are superior to the

benefits of reproductive assurance (Layman et al. 2017).

Peripatric populations of *Lgh* in western Europe seem to solve all these problems and paradox by

mixing clonality with preferential allogamy: clonality allows the local maintenance and spreading of

population without losing heterozygosity and genetic diversity, and subtle but significant sexuality

with preferential allogamy, favoured by LSI and faster growth of crossed-pollen tubes, enables

recombination between lineages favouring the emergence of locally adapted genomes with

potential higher vigour and fertility (heterosis: Darwin 1876, Lippman & Zamir 2007, Birchleret al.

2010).

This reproductive mode, i.e. dominant clonality with preferential allogamy, is common in plants

(Vallejo-Marin et al. 2010). The micro-physiological mechanism(s) slowing down the growth of self-

pollen tubes, rather than blocking them, in simultaneous monoecious and hermaphrodite SC species

and resulting to favour allogamy when compatible pollen is available may also be common in plant,

although potentially overlooked (Glover 2007, Nasrallah 2017). In such peripatric conditions, selfing

would thus only present the limited or transient advantage to produce seeds that can maintain in

local seed banks and with different dispersal properties compared to clonal propagules. The limited

interest of selfing in this species may explain why, based on photos collected on the web, L-morph

(LSI) individuals seem more frequent in peripatric than in native populations, and why in western

Europe, around 76% of the populations are L-morph (LSI) against 24% of S-morph (SC; Portillo et al.

2021). Our results thus call for measuring the true proportions of LSI and SC in invasive versus native

worldwide *Lgh* populations with estimation of their rates of clonality and of selfing.

Unusual selfing syndrome in *Lgh* populations

Highly selfed populations of different plant species tend to share similar morphology and functions

resulting in a set of traits called *selfing syndrome*, including reduced flower size (Darwin 1876,

Tsuchimatsu & Fujii 2022). Selfing syndrome, including reduced flower size, seems to evolve rapidly,

as observed for example in five generations in *Mimulus guttatus* (Bodbyl-Roels & Kelly 2011), in four

generations in *Silene latifolia* (Delph et al. 2004), in three generations in *Phlox drummondii* (Lendvai

& Levin 2003) and after only two generations in *Eichhornia paniculata* (Worley & Barrett 2000).

On the contrary, S-morph (SC) individuals in *Lgh* develop larger flowers than allogamous L-morphs

(Portillo-Lemus et al. 2021). However, S-morph (SC) individuals and populations dominantly

reproduced using clonality and produced selfed offspring only when lacking crossed pollen. Both

clonality and faster growth of crossed-pollen tubes may delay or even compromise the emergence

of the first steps of a selfing syndrome. These two traits may call for further studies on the

emergence of selfing syndrome in partially clonal and selfed populations.

Conclusion

We found that peripatric populations of *Lgh* in western Europe reproduced using dominant clonality

with limited but stable significant sexuality with preferential outcrossing, in nearly all populations,

within a large clonal diversity. *Lgh* is one of the most invasive aquatic plants in the world, and

considerable efforts are made to limit its deleterious effects in the newly colonized ecosystems

(Thouvenot et al. 2013, Grewell et al. 2016, Portillo-Lemus et al. 2021). The rare sexual events,

allogamous when possible, occurring in invasive peripatric populations of *Lgh* may favour the

emergence of new genotypes, more adapted to local conditions. Managers should thus chiefly

concentrate their actions on the most sexual populations, and on the contact zones between S- and

L-morph populations in France. We also found variations of the rates of clonality and of selfing

among *Lgh* populations. Considering the importance of reproductive modes on the dynamics and

evolution of populations (Duminil et al. 2007, Ellegren& Galtier 2016, Glémin et al. 2019), our results

advocate that management actions should consider the local effective reproductive modes to

control *Lgh* population by population. Finally, knowing the reproductive modes of populations, the

distribution of clones and the self-compatibility across western European populations now allow

deciphering and interpreting their population structure, identifying their origin and routes, and

predicting their possible short-term dynamics.

Data Accessibility Statement

The data that support the findings of this study are openly available on Zenodo (https://doi.org/10.5281/zenodo.10246264).

Benefits Generated

Benefits from this research accrue from the sharing of our data and results on public databases as described above.

Acknowledgements

We thank Michel Bozec (UMR DECOD), Diane Corbin (FRAPNA Loire-Ecopôle du Forez), Guillaume Le Roux (Reserve Naturel Val d'Alier Châtel-de-Neuvre), Fabrice Dejoux (Service Agriculture Environnement Roannais Agglomération and Jordi Bou Manobens (University of Girona, Spain) for helping to collect *Lgh* samples. We thank Olivier Coriton and Virginie Huteau for their help acquiring karyotypes.

We warmly thank Ingrid Leveque, Marie-Therese Delaroche, Gervaise Fevrier and Patricia Nadan for their administrative support and facilitation.

Funding information

This work was supported by FEDER Région Centre-Val de Loire, by Agence de l'eau Loire-Bretagne (grant Nature 2045, programme 9025, AP 2015 9025) and by the French National Research Agency (project Clonix2D ANR-18-CE32-0001). FEDER funded the doctoral grant of L. Portillo-Lemus and salary of Marilyne Harang. Postdoctoral grant of Ronan Becheler was funded by Clonix2D ANR-18-CE32-0001.

Conflict of interest

The authors declare that they have no financial conflicts of interest based on the content of this

article.

Author contributions

DB and SS laid the foundation of this work, conceived the study and were responsible for funding

applications. LPL, MH and DB collected samples and performed field observations. LPL and DB read

karyotypes and counted chromosomes. EP and DB conceived and supervised RAD4SNP and Hi-Plex

methodology, BJP and DJP contributed to define the SNP set. LPL, MH and ALB performed RAD4SNP

approach to develop the SNPs. LPL and MH extracted DNA and genotyped the samples, SMC and

RCV sequenced the data. GL performed the bioinformatic analyses and SS computed the Bayesian

allele dosages. DB, LPL and SS performed early data explorations at the end of LPL PhD thesis that

was supervised by DB and SS. RB and SS performed computations and analyses, tracked and

managed the bibliography, produced tables, figures and interpretation. DB, RB and SS wrote the

manuscript. All authors read and approved the final manuscript.

References

Abdallah, D., Baraket, G., Perez, V., Ben Mustapha, S., Salhi-Hannachi, A., & Hormaza, J. I. (2019).

Analysis of self-incompatibility and genetic diversity in diploid and hexaploid plum

genotypes. *Frontiers in Plant Science*, *10*.

https://www.frontiersin.org/articles/10.3389/fpls.2019.00896

Agapow, P.-M., & Burt, A. (2001). Indices of multilocus linkage disequilibrium. *Molecular Ecology*

Notes, *1*(1–2), 101–102. https://doi.org/10.1046/j.1471-8278.2000.00014.x

Allendorf, F. W., & Lundquist, L. L. (2003). Introduction: Population Biology, Evolution, and

Control of Invasive Species. *Conservation Biology*, 17(1), 24–30.

https://doi.org/10.1046/j.1523-1739.2003.02365.x

Armitage, J. D., Könyves, K., Bailey, J. P., David, J. C., & Culham, A. (2013). A molecular,

morphological and cytological investigation of the identity of non-native *Ludwigia*

(Onagraceae) populations in Britain. *New Journal of Botany*, *3*(2), 88–95.

https://doi.org/10.1179/2042349713Y.0000000023

Arnaud-Haond, S., Alberto, F., Teixeira, S., Procaccini, G., Serrão, E. A., & Duarte, C. M. (2005).

Assessing genetic diversity in clonal organisms: low diversity or low resolution? combining

power and cost efficiency in selecting markers. *Journal of Heredity*, *96*(4), 434–440.

https://doi.org/10.1093/jhered/esi043

Arnaud-Haond, S., Duarte, C. M., Alberto, F., & Serrão, E. A. (2007). Standardizing methods to

address clonality in population studies. *Molecular Ecology*, *16*(24), 5115–5139.

https://doi.org/10.1111/j.1365-294X.2007.03535.x

Arnaud-Haond, S., Stoeckel, S., & Bailleul, D. (2020). New insights into the population genetics of

partially clonal organisms: When seagrass data meet theoretical expectations. *Molecular*

Ecology, *29*(17), 3248–3260. https://doi.org/10.1111/mec.15532

Baduel, P., Bray, S., Vallejo-Marin, M., Kolář, F., & Yant, L. (2018). The "polyploid hop": shifting

challenges and opportunities over the evolutionary lifespan of genome duplications. *Frontiers*

in Ecology and Evolution, *6*. https://www.frontiersin.org/articles/10.3389/fevo.2018.00117

Baird, N. A., Etter, P. D., Atwood, T. S., Currey, M. C., Shiver, A. L., Lewis, Z. A., Selker, E. U.,

Cresko, W. A., & Johnson, E. A. (2008). Rapid SNP Discovery and Genetic Mapping Using

Sequenced RAD Markers. *PLOS ONE*, 3(10), e3376.

https://doi.org/10.1371/journal.pone.0003376

Balloux, F., Lehmann, L., & de Meeûs, T. (2003). The population genetics of clonal and partially

clonal diploids. *Genetics*, *164*(4), 1635–1644. https://doi.org/10.1093/genetics/164.4.1635

Barloy, D., Lemus, L. P.-, Krueger-Hadfield, S. A., Huteau, V., & Coriton, O. (2024). Genomic

relationships among diploid and polyploid species of the genus Ludwigia L. section Jussiaea

using a combination of cytogenetic, morphological, and crossing investigations. *Peer*

Community in Evolutionary Biology, 4, e8. https://doi.org/10.24072/pci.evolbiol.100645.

Barrett, S. C. H., Colautti, R. I., & Eckert, C. G. (2008). Plant reproductive systems and evolution

during biological invasion. *Molecular Ecology*, 17(1), 373–383.

https://doi.org/10.1111/j.1365-294X.2007.03503.x

Barrett, S. C. H. (2010). Understanding plant reproductive diversity. *Philosophical Transactions of*

the Royal Society B: Biological Sciences, *365*(1537), 99–109.

https://doi.org/10.1098/rstb.2009.0199

Becheler, R., Guillemin, M.-L., Stoeckel, S., Mauger, S., Saunier, A., Brante, A., Destombe, C., &

Valero, M. (2020). After a catastrophe, a little bit of sex is better than nothing: Genetic

consequences of a major earthquake on asexual and sexual populations. *Evolutionary*

Applications, *13*(8), 2086–2100. https://doi.org/10.1111/eva.12967

Becheler, R., Masson, J.-P., Arnaud-Haond, S., Halkett, F., Mariette, S., Guillemin, M.-L., Valero,

M., Destombe, C., & Stoeckel, S. (2017). ClonEstiMate, a Bayesian method for quantifying

rates of clonality of populations genotyped at two-time steps. *Molecular Ecology Resources*,

17(6), e251–e267. https://doi.org/10.1111/1755-0998.12698

Beck, J. B., Al-Shehbaz, I. A., & Schaal, B. A. (2006). *Leavenworthia* (Brassicaceae) revisited:

testing classic systematic and mating system hypotheses. *Systematic Botany*, *31*(1), 151–159. https://doi.org/10.1600/036364406775971732

Besnard, A.-L., Park, D. J., Pope, B. J., Hammet, F., Michon-Coudouel, S., Biget, M., Krueger-Hadfield, S. A., Mauger, S., & Petit, E. J. (2023). Workflow for SNP genotyping using the Hi-Plex method. *Protocols.io*, https://doi.org/10.17504/protocols.io.8epv5jnnnl1b/v2

Birchler, J. A., Yao, H., Chudalayandi, S., Vaiman, D., & Veitia, R. A. (2010). Heterosis. *The Plant Cell*, *22*(7), 2105–2112. https://doi.org/10.1105/tpc.110.076133

Bodbyl Roels, S. A., & Kelly, J. K. (2011). Rapid evolution caused by pollinator loss in *Mimulus guttatus*. *Evolution*, *65*(9), 2541–2552. https://doi.org/10.1111/j.1558-5646.2011.01326.x

Booy, G., Hendriks, R.J.J., Smulders, M.J.M., Van Groenendael, J.M. and Vosman, B. (2000). Genetic Diversity and the Survival of Populations. *Plant Biology*, *2,* 379-395. https://doi.org/10.1055/s-2000-5958

Bou Manobens, J., Portillo-Lemus, L., & Curcó i Masip, A. (2019). New contributions to allochthonous Ludwigia species (Onagraceae) on Catalonia. Butlletí de La Institució Catalana d'Història Natural, 83, 45–48. https://doi.org/10.2436/20.1502.01.4

Brennan, A. C., Harris, S. A., Tabah, D. A., & Hiscock, S. J. (2002). The population genetics of sporophytic self-incompatibility in *Senecio squalidus* L. (Asteraceae) I: S allele diversity in a natural population. *Heredity*, *89*(6), Article 6. https://doi.org/10.1038/sj.hdy.6800159

Bürkli, A., Sieber, N., Seppälä, K., & Jokela, J. (2017). Comparing direct and indirect selfing rate estimates: When are population-structure estimates reliable? *Heredity*, *118*(6), Article 6. https://doi.org/10.1038/hdy.2017.1

Busch, J. W. (2005). Inbreeding depression in self-incompatible and self-compatible populations of *Leavenworthia alabamica*. *Heredity*, *94*(2), Article 2. https://doi.org/10.1038/sj.hdy.6800584

Busch, J. W., & Urban, L. (2011). Insights gained from 50 years of studying the evolution of self-compatibility in *Leavenworthia* (Brassicaceae). *Evolutionary Biology*, *38*(1), 15–27. https://doi.org/10.1007/s11692-010-9104-5

Cabin, R. J., Evans, A. S., Jennings, D. L., Marshall, D. L., Mitchell, R. J., & Sher, A. A. (1996). Using bud pollinations to avoid self-incompatibility: Implications from studies of three mustards. *Canadian Journal of Botany*, 74(2), 285–289. https://doi.org/10.1139/b96-034

Castric, V., Bernatchez, L., Belkhir, K., & Bonhomme, F. (2002). Heterozygote deficiencies in small lacustrine populations of brook charr *Salvelinus fontinalis* Mitchill (Pisces, Salmonidae): A test of alternative hypotheses. *Heredity*, *89*(1), Article 1. https://doi.org/10.1038/sj.hdy.6800089

Castric, V., & Vekemans, X. (2004). Plant self-incompatibility in natural populations: a critical assessment of recent theoretical and empirical advances. *Molecular Ecology*, *13*(10), 2873–2889. https://doi.org/10.1111/j.1365-294X.2004.02267.x

Charlesworth, D. (2003). Effects of inbreeding on the genetic diversity of populations. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *358*(1434), 1071–1084. https://doi.org/10.1098/rstb.2003.1296

Charlesworth, D. (2006). Evolution of Plant Breeding Systems. Current Biology, *16*(17), 726–735. https://doi.org/10.1016/j.cub.2006.07.068

Charlesworth, D., Vekemans, X., Castric, V., & Glémin, S. (2005). Plant self-incompatibility systems: A molecular evolutionary perspective. *New Phytologist*, *168*(1), 61–69. https://doi.org/10.1111/j.1469-8137.2005.01443.x

Chen, X., Hao, S., Wang, L., Fang, W., Wang, Y., & Li, X. (2012). Late-acting self-incompatibility in tea plant (*Camellia sinensis*). *Biologia*, *67*(2), 347–351. https://doi.org/10.2478/s11756-012-0018-9

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools. GigaScience, 10, giab008. https://doi.org/10.1093/gigascience/giab008

Dandelot, S. (2004). *Les Ludwigia spp. Invasives du Sud de la France: Historique, Biosystématique, Biologie et Ecologie* [PhD thesis, Aix-Marseille 3]. https://www.theses.fr/2004AIX30052

Dandelot, S., Verlaque, R., Dutartre, A., & Cazaubon, A. (2005). Ecological, Dynamic and

Taxonomic Problems Due to Ludwigia (Onagraceae) in France. Hydrobiologia, 551(1), 131–

136. https://doi.org/10.1007/s10750-005-4455-0

Darwin, C. R. (1876). *The effects of cross and self-fertilization in the vegetable kingdom*. John Murray

(London).

David, P., Pujol, B., Viard, F., Castella, V., & Goudet, J. (2007). Reliable selfing rate estimates from

imperfect population genetic data. *Molecular Ecology*, *16*(12), 2474–2487.

https://doi.org/10.1111/j.1365-294X.2007.03330.x

De Meeûs, T., & Balloux, F. (2004). Clonal reproduction and linkage disequilibrium in diploids: A

simulation study. *Infection, Genetics and Evolution*, *4*(4), 345–351.

https://doi.org/10.1016/j.meegid.2004.05.002

De Meeûs, T., Lehmann, L., & Balloux, F. (2006). Molecular epidemiology of clonal diploids: A

quick overview and a short DIY (do it yourself) notice. *Infection, Genetics and Evolution*,

6(2), 163–170. https://doi.org/10.1016/j.meegid.2005.02.004

De Meeûs, T., Prugnolle, F., & Agnew, P. (2007). Asexual reproduction: Genetics and evolutionary

aspects. *Cellular and Molecular Life Sciences*, *64*(11), 1355–1372.

https://doi.org/10.1007/s00018-007-6515-2

De Nettancourt, D. (1997). Incompatibility in angiosperms. *Sexual Plant Reproduction*, *10*(4), 185–

199. https://doi.org/10.1007/s004970050087

De Nettancourt, D. (2001). *Incompatibility and Incongruity in Wild and Cultivated Plants*. Springer.

https://doi.org/10.1007/978-3-662-04502-2

Dellaporta, S. L., & Calderon-Urrea, A. (1993). Sex Determination in Flowering Plants. *The Plant

Cell*, *5*(10), 1241–1251. https://doi.org/10.2307/3869777

Delord, C., Lassalle, G., Oger, A., Barloy, D., Coutellec, M.-A., Delcamp, A., Evanno, G., Genthon,

C., Guichoux, E., Le Bail, P.-Y., Le Quilliec, P., Longin, G., Lorvelec, O., Massot, M.,

Reveillac, E., Rinaldo, R., Roussel, J.-M., Vigouroux, R., Launey, S., & Petit, E. J. (2018). A

cost-and-time effective procedure to develop SNP markers for multiple species: A support for

community genetics. *Methods in Ecology and Evolution*, *9*(9), 1959–1974.

https://doi.org/10.1111/2041-210X.13034

Delph, L. F., Gehring, J. L., Frey, F. M., Arntz, A. M., & Levri, M. (2004). Genetic constraints on

floral evolution in a sexually dimorphic plant revealed by artificial selection. *Evolution*, *58*(9),

1936–1946. https://doi.org/10.1111/j.0014-3820.2004.tb00481.x

Dorken, M. E., & Eckert, C. G. (2001). Severely reduced sexual reproduction in northern populations

of a clonal plant, *Decodonverticillatus* (Lythraceae). *Journal of Ecology*, *89*(3), 339–350.

https://doi.org/10.1046/j.1365-2745.2001.00558.x

Dufresne, F., Stift, M., Vergilino, R., & Mable, B. K. (2014). Recent progress and challenges in

population genetics of polyploid organisms: An overview of current state-of-the-art molecular

and statistical tools. *Molecular Ecology*, *23*(1), 40–69. https://doi.org/10.1111/mec.12581

Duminil, J., Fineschi, S., Hampe, A., Jordano, P., Salvini, D., Vendramin, G. G., & Petit, R. J. (2007).

Can population genetic structure be predicted from life-history traits? *The American*

Naturalist, *169*(5), 662–672. https://doi.org/10.1086/513490

Eckert, C., Samis, K. & Dart, S. (2006). Reproductive Assurance and the Evolution of Uniparental

Reproduction in Flowering Plants. In Harder & Barrett (Eds), *Ecology and Evolution of*

Flowers, pp.183–200. Oxford, UK: Oxford University Press.

Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*, *17*(7),

Article 7. https://doi.org/10.1038/nrg.2016.58

Estes, J. R., & Thorp, R. W. (1974). Pollination in Ludwigia peploides ssp. Glabrescens

(Onagraceae). *Bulletin of the Torrey Botanical Club,* *101*(5), 272–276.

https://doi.org/10.2307/2484872

Fehrer, J. (2010). Unraveling the mysteries of reproduction. *Heredity*, *104*(5), Article 5.

https://doi.org/10.1038/hdy.2010.12

Fischer, M. C., Rellstab, C., Leuzinger, M., Roumet, M., Gugerli, F., Shimizu, K. K., Holderegger,

R., & Widmer, A. (2017). Estimating genomic diversity and population differentiation – an

empirical comparison of microsatellite and SNP variation in *Arabidopsis halleri*. *BMC*

Genomics, *18*(1), 69. https://doi.org/10.1186/s12864-016-3459-7

Flanagan, B. A., Krueger-Hadfield, S. A., Murren, C. J., Nice, C. C., Strand, A. E., & Sotka, E. E.

(2021). Founder effects shape linkage disequilibrium and genomic diversity of a partially

clonal invader. *Molecular Ecology*, *30*(9), 1962–1978. https://doi.org/10.1111/mec.15854

Forsman, A. (2014). Effects of genotypic and phenotypic variation on establishment are important

for conservation, invasion, and infection biology. *Proceedings of the National Academy of*

Sciences, 111(1), 302–307. https://doi.org/10.1073/pnas.1317745111

Fox, C. W., & Reed, D. H. (2011). Inbreeding Depression Increases with Environmental Stress: An

Experimental   Study   and   Meta-Analysis.   *Evolution*,   65(1),   246–258.

https://doi.org/10.1111/j.1558-5646.2010.01108.x

Franklin-Tong,   V.   E.   (2008).   *Self-Incompatibility   in   Flowering   Plants*.   Springer.

https://doi.org/10.1007/978-3-540-68486-2

Fry, N. K., Savelkoul, P. H., & Visca, P. (2009). Amplified Fragment Length Polymorphism

Analysis. In: Caugant, D. (eds) Molecular Epidemiology of Microorganisms. *Methods in*

Molecular Biology, vol 551, 89-104. Humana Press, Totowa, NJ. https://doi.org/10.1007/978-

1-60327-999-4_8

Gibbs, P. E. (2014). Late-acting self-incompatibility – the pariah breeding system in flowering plants.

New Phytologist, *203*(3), 717–734. https://doi.org/10.1111/nph.12874

Glémin, S., Bataillon, T., Ronfort, J., Mignot, A., & Olivieri, I. (2001). inbreeding depression in small

populations   of   self-incompatible   plants.   *Genetics*,   *159*(3),   1217–1229.

https://doi.org/10.1093/genetics/159.3.1217

Glémin, S., François, C. M., & Galtier, N. (2019). Genome Evolution in Outcrossing vs. Selfing vs.

Asexual   Species.   In   M.   Anisimova   (Ed.),   *Evolutionary   Genomics: Statistical   and*

Computational Methods (pp. 331–369). Springer. https://doi.org/10.1007/978-1-4939-9074-

0_11

Glover, B. J. (2007). Preventing Self-fertilization. In B. Glover (Ed.), Understanding Flowers and

Flowering: An integrated approach (p. 0). Oxford University Press.

https://doi.org/10.1093/acprof:oso/9780198565970.003.0012

Golding, G. B., & Strobeck, C. (1980). Linkage disequilibrium in a finite population that is partially

selfing. *Genetics*, *94*(3), 777–789. https://doi.org/10.1093/genetics/94.3.777

Goodwillie, C., & Weber, J. J. (2018). The best of both worlds? A review of delayed selfing in

flowering plants. *American Journal of Botany*, *105*(4), 641–655.

https://doi.org/10.1002/ajb2.1045

Grewell, B., Netherland, M., & Skaer Thomason, M. (2016). Establishing research and management

priorities for invasive water primroses *(Ludwigia spp.)*.

https://doi.org/10.13140/RG.2.1.5020.6482

Haldane, J. B. S. (1930). Theoretical genetics of autopolyploids. *Journal of Genetics*, 22(3), 359–

372. https://doi.org/10.1007/BF02984197

Halkett, F., Simon, J.-C., & Balloux, F. (2005). Tackling the population genetics of clonal and

partially clonal organisms. *Trends in Ecology & Evolution*, *20*(4), 194–201.

https://doi.org/10.1016/j.tree.2005.01.001

Hammet, F., Mahmood, K., Green, T. R., Nguyen-Dumont, T., Southey, M. C., Buchanan, D. D.,

Lonie, A., Nathanson, K. L., Couch, F. J., Pope, B. J., & Park, D. J. (2019). Hi-Plex2: A

simple and robust approach to targeted sequencing-based genetic screening. *BioTechniques*,

67(3), 118–122. https://doi.org/10.2144/btn-2019-0026

Hao, Y.-Q., Zhao, X.-F., She, D.-Y., Xu, B., Zhang, D.-Y., & Liao, W.-J. (2012). The Role of Late-

acting self-incompatibility and early-acting inbreeding depression in governing female

fertility in monkshood, *Aconitum kusnezoffii*. *PLOS ONE*, *7*(10), e47034.

https://doi.org/10.1371/journal.pone.0047034

Harder, L. D., & Barrett, S. C. H. (2006). *Ecology and Evolution of Flowers*. OUP Oxford.

Hardy, O. J. (2016). Population genetics of autopolyploids under a mixed mating model and the estimation of selfing rate. *Molecular Ecology Resources*, *16*(1), 103–117. https://doi.org/10.1111/1755-0998.12431

Hardy, O. J., & Vekemans, X. (2002). SPAGEDI: A versatile computer program to analyze spatial genetic structure at the individual or population levels. *Molecular Ecology Notes*, *2*(4), 618–620. https://doi.org/10.1046/j.1471-8286.2002.00305.x

He, Z.-Z., Stotz, G. C., Liu, X., Liu, J.-J., Wang, Y.-G., Yang, J., Li, L.-F., Zhang, W.-J., Nan, P., & Song, Z.-P. (2024). A global synthesis of the patterns of genetic diversity in endangered and invasive plants. *Biological Conservation*, 291, 110473. https://doi.org/10.1016/j.biocon.2024.110473

Herben, T., Suda, J., & Klimešová, J. (2017). Polyploid species rely on vegetative reproduction more than diploids: A re-examination of the old hypothesis. *Annals of Botany*, *120*(2), 341–349. https://doi.org/10.1093/aob/mcx009

Hoban, S., Campbell, C. D., da Silva, J. M., Ekblom, R., Funk, W. C., Garner, B. A., Godoy, J. A., Kershaw, F., MacDonald, A. J., Mergeay, J., Minter, M., O'Brien, D., Vinas, I. P., Pearson, S. K., Pérez-Espona, S., Potter, K. M., Russo, I.-R. M., Segelbacher, G., Vernesi, C., & Hunter, M. E. (2021). Genetic diversity is considered important but interpreted narrowly in country reports to the Convention on Biological Diversity: Current actions and indicators are insufficient. *Biological Conservation*, *261*, 109233. https://doi.org/10.1016/j.biocon.2021.109233

Hoch, P. C., Wagner, W. L., & Raven, P. H. (2015). The correct name for a section of Ludwigia L. (Onagraceae). *PhytoKeys*, *50*, 31–34. https://doi.org/10.3897/phytokeys.50.4887

Holsinger, K. E. (2000). Reproductive systems and evolution in vascular plants. *Proceedings of the National Academy of Sciences*, *97*(13), 7037–7042. https://doi.org/10.1073/pnas.97.13.7037

Husband, B. C., Ozimec, B., Martin, S. L., & Pollock, L. (2008). Mating consequences of polyploid evolution in flowering plants: current trends and insights from synthetic polyploids.

International Journal of Plant Sciences, *169*(1), 195–206. https://doi.org/10.1086/523367

Igic, B., Lande, R., & Kohn, J. R. (2008). Loss of self-incompatibility and its evolutionary consequences. *International Journal of Plant Sciences*, *169*(1), 93–104. https://doi.org/10.1086/523362

Koelling, V. A., Hamrick, J. L., & Mauricio, R. (2011). Genetic diversity and structure in two species of *Leavenworthia* with self-incompatible and self-compatible populations. *Heredity*, *106*(2), 310-318. https://doi.org/10.1038/hdy.2010.59

Krueger-Hadfield, S. A., Guillemin, M.-L., Destombe, C., Valero, M., & Stoeckel, S. (2021). Exploring the Genetic Consequences of Clonality in Haplodiplontic Taxa. *Journal of Heredity*, 112(1), 92–107. https://doi.org/10.1093/jhered/esaa063

Lawrence, M. J., Marshall, D. F., Curtis, V. E., & Fearon, C. H. (1985). Gametophytic self-incompatibility re-examined: A reply. *Heredity*, 54(1), 131–138. https://doi.org/10.1038/hdy.1985.17

Layman, N. C., Fernando, M. T. R., Herlihy, C. R., & Busch, J. W. (2017). Costs of selfing prevent the spread of a self-compatibility mutation that causes reproductive assurance. *Evolution*, 71(4), 884–897. https://doi.org/10.1111/evo.13167

Lê, S., Josse, J. & Husson, F. (2008). FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software,* *25*(1), 1-18. https://doi.org/10.18637/jss.v025.i01

Lendvai, G., & Levin, D. A. (2003). Rapid response to artificial selection on flower size in *Phlox*. *Heredity*, *90*(4), Article 4. https://doi.org/10.1038/sj.hdy.6800249

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics*, *26*, 589-595. https://doi.org/10.1093/bioinformatics/btp698.

Lippman, Z. B., & Zamir, D. (2007). Heterosis: Revisiting the magic. *Trends in Genetics*, *23*(2), 60–66. https://doi.org/10.1016/j.tig.2006.12.006

Lucek, K., & Willi, Y. (2021). Drivers of linkage disequilibrium across a species' geographic range. *PLOS Genetics*, *17*(3), e1009477. https://doi.org/10.1371/journal.pgen.1009477

Mable, B. K. (2008). Genetic causes and consequences of the breakdown of self-incompatibility:

Case studies in the Brassicaceae. *Genetics Research*, *90*(1), 47–60.

https://doi.org/10.1017/S0016672307008907

Meek, M. H., & Larson, W. A. (2019). The future is now: Amplicon sequencing and sequence capture

usher in the conservation genomics era. Molecular Ecology Resources, 19(4), 795–803.

https://doi.org/10.1111/1755-0998.12998

Meirmans, P. G., Liu, S., & van Tienderen, P. H. (2018). The analysis of polyploid genetic data.

Journal of Heredity, *109*(3), 283–296. https://doi.org/10.1093/jhered/esy006

Nasrallah, J. B. (2017). Plant mating systems: Self-incompatibility and evolutionary transitions to

self-fertility in the mustard family. Current Opinion in Genetics & Development, 47, 54–60.

https://doi.org/10.1016/j.gde.2017.08.005

Navascués, M., Stoeckel, S., & Mariette, S. (2010). Genetic diversity and fitness in small populations

of partially asexual, self-incompatible plants. *Heredity*, *104*(5), Article 5.

https://doi.org/10.1038/hdy.2009.159

Nesom, G.L. & Kartesz, J.T. (2000). Observations on the *Ludwigia uruguayensis* complex

(Onagraceae) in the United States. Castanea 65(2): 123-125.

http://www.jstor.org/stable/4034110

Nordborg, M. (2000). Linkage disequilibrium, gene trees and selfing: an ancestral recombination

graph with partial self-fertilization. *Genetics*, *154*(2), 923–929.

https://doi.org/10.1093/genetics/154.2.923

Okada, M., Grewell, B. J., & Jasieniuk, M. (2009). Clonal spread of invasive *Ludwigia hexapetala*

and *L. grandiflora* in freshwater wetlands of California. *Aquatic Botany*, *91*(3), 123–129.

https://doi.org/10.1016/j.aquabot.2009.03.006

Orive, M. E., & Krueger-Hadfield, S. A. (2021). Sex and asex: a clonal lexicon. *Journal of Heredity*,

112(1), 1–8. https://doi.org/10.1093/jhered/esaa058

Pannell, J. R., Auld, J. R., Brandvain, Y., Burd, M., Busch, J. W., Cheptou, P.-O., Conner, J. K.,

Goldberg, E. E., Grant, A.-G., Grossenbacher, D. L., Hovick, S. M., Igic, B., Kalisz, S., Petanidou, T., Randle, A. M., de Casas, R. R., Pauw, A., Vamosi, J. C., & Winn, A. A. (2015). The scope of Baker's law. *New Phytologist*, *208*(3), 656–667. https://doi.org/10.1111/nph.13539

Porcher, E., & Lande, R. (2005). The evolution of self-fertilization and inbreeding depression under pollen discounting and pollen limitation. *Journal of Evolutionary Biology*, *18*(3), 497–508. https://doi.org/10.1111/j.1420-9101.2005.00905.x

Portillo Lemus, L. O. (2021). Système de reproduction, polyploïdie et diversité génétique des populations invasives de Ludwigia grandiflora subsp hexapetala en France. *PhD Thesis*. http://www.theses.fr/2021NSARA088/document

Portillo Lemus, L. O., Bozec, M., Harang, M., Coudreuse, J., Haury, J., Stoeckel, S., & Barloy, D. (2021). Self-incompatibility limits sexual reproduction rather than environmental conditions in an invasive water primrose. *Plant-Environment Interactions*, *2*(2), 74–86. https://doi.org/10.1002/pei3.10042

Portillo Lemus, L. O., Harang, M., Bozec, M., Haury, J., Stoeckel, S., & Barloy, D. (2022). Late-acting self-incompatible system, preferential allogamy and delayed selfing in the heteromorphic invasive populations of *Ludwigia grandiflora subsp. Hexapetala*. *Peer Community Journal*, *2*. https://doi.org/10.24072/pcjournal.108

Pyšek, P. (1998). Is there a taxonomic pattern to plant invasions? *Oikos*, *82*(2), 282–294. https://doi.org/10.2307/3546968

Reichel, K., Masson, J.-P., Malrieu, F., Arnaud-Haond, S., & Stoeckel, S. (2016). Rare sex or out of reach equilibrium? The dynamics of FISin partially clonal organisms. *BMC Genetics*, *17*(1), 76. https://doi.org/10.1186/s12863-016-0388-z

Rice, S. H. (2004). *Evolutionary theory: Mathematical and conceptual foundations*. Sinauer Associates.

Richards, A. J. (1997). *Plant breeding systems*. Garland Science.

Roman, J., & Darling, J. A. (2007). Paradox lost: Genetic diversity and the success of aquatic invasions. *Trends in Ecology & Evolution*, 22(9), 454–464. https://doi.org/10.1016/j.tree.2007.07.002

Ronfort, J., Jenczewski, E., Bataillon, T., & Rousset, F. (1998). Analysis of population structure in autotetraploid species. *Genetics*, *150*(2), 921–930. https://doi.org/10.1093/genetics/150.2.921

Rouger, R., Reichel, K., Malrieu, F., Masson, J. P., & Stoeckel, S. (2016). Effects of complex life cycles on genetic diversity: Cyclical parthenogenesis. *Heredity*, *117*(5), Article 5. https://doi.org/10.1038/hdy.2016.52

Roze, D. (2016). Background selection in partially selfing populations. *Genetics*, *203*(2), 937–957. https://doi.org/10.1534/genetics.116.187955

Schmidt, T. L., Jasper, M.-E., Weeks, A. R., & Hoffmann, A. A. (2021). Unbiased population heterozygosity estimates from genome-wide sequence data. *Methods in Ecology and Evolution*, *12*(10), 1888–1898. https://doi.org/10.1111/2041-210X.13659

Seavey, S. R., & Bawa, K. S. (1986). Late-acting self-incompatibility in angiosperms. *The Botanical Review*, *52*(2), 195–219. https://doi.org/10.1007/BF02861001

Skaer Thomason, M. J., Grewell, B. J., & Netherland, M. D. (2018a). Dynamics of *Ludwigia hexapetala* invasion at three spatial scales in a regulated river. *Wetlands*, *38*(6), 1285–1298. https://doi.org/10.1007/s13157-018-1053-2

Skaer Thomason, M. J., McCort, C. D., Netherland, M. D., & Grewell, B. J. (2018b). Temporal and nonlinear dispersal patterns of *Ludwigia hexapetala* in a regulated river. *Wetlands Ecology and Management*, *26*(5), 751–762. https://doi.org/10.1007/s11273-018-9605-z

Steinecke, C., Gorman, C. E., Stift, M., & Dorken, M. E. (2022). Outcrossing rates in an experimentally admixed population of self-compatible and self-incompatible Arabidopsis lyrata. *Heredity*, *128*(1), Article 1. https://doi.org/10.1038/s41437-021-00489-8

Stoeckel, S., Arnaud-Haond, S., & Krueger-Hadfield, S. A. (2021b). The combined effect of haplodiplonty and partial clonality on genotypic and genetic diversity in a finite mutating

population. *Journal of Heredity*, *112*(1), 78–91. https://doi.org/10.1093/jhered/esaa062

Stoeckel, S., Barloy, D., Portillo Lemus, L., & Becheler, R. (2023). Genotyping measures and population genetic indices for assesing reproductive modes of polyploid Ludwigia grandiflora subsp. hexapetala in western Europe [Data set]. Zenodo. https://doi.org/10.5281/zenodo.10246264

Stoeckel, S., Becheler, R., Bocharova, E., & Barloy, D. (2023). GenAPoPop 1.0: A user-friendly software to analyse genetic diversity and structure from partially clonal and selfed autopolyploid organisms. Molecular Ecology Resources, 00, 1–11. https://doi.org/10.1111/1755-0998.13886

Stoeckel, S., Grange, J., Fernández-Manjarres, J. F., Bilger, I., Frascaria-Lacoste, N., & Mariette, S. (2006). Heterozygote excess in a self-incompatible and partially clonal forest tree species— Prunus avium L. *Molecular Ecology*, *15*(8), 2109–2118. https://doi.org/10.1111/j.1365-294X.2006.02926.x

Stoeckel, S., Castric, V., Mariette, S., & Vekemans, X. (2008). Unequal allelic frequencies at the self-incompatibility locus within local populations of Prunus avium L.: An effect of population structure? Journal of Evolutionary Biology, 21(3), 889–899. https://doi.org/10.1111/j.1420-9101.2008.01504.x

Stoeckel, S., & Masson, J.-P. (2014). The exact distributions of fis under partial asexuality in small finite populations with mutation. *PLOS ONE*, *9*(1), e85228. https://doi.org/10.1371/journal.pone.0085228

Stoeckel, S., Porro, B., & Arnaud-Haond, S. (2021a). The discernible and hidden effects of clonality on the genotypic and genetic states of populations: Improving our estimation of clonal rates. *Molecular Ecology Resources*, *21*(4), 1068–1084. https://doi.org/10.1111/1755-0998.13316

Terpilowski, (2019). scikit-posthocs: Pairwise multiple comparison tests in Python. Journal of Open Source Software, 4(36), 1169, https://doi.org/10.21105/joss.01169

Thouvenot, L., Haury, J., & Thiebaut, G. (2013). A success story: Water primroses, aquatic plant

pests. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *23*(5), 790–803.

https://doi.org/10.1002/aqc.2387

Tsuchimatsu, T., & Sota, F. (2022). The selfing syndrome and beyond: diverse evolutionary

consequences of mating system transitions in plants. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, 377, 20200510. https://doi.org/10.1098/rstb.2020.0510.

Vallejo-Marín, M., Dorken, M. E., & Barrett, S. C. H. (2010). The ecological and evolutionary

consequences of clonality for plant mating. *Annual Review of Ecology, Evolution, and*

Systematics, *41*(1), 193–213. https://doi.org/10.1146/annurev.ecolsys.110308.120258

Van Drunen, W. E., & Husband, B. C. (2019). Evolutionary associations between polyploidy, clonal

reproduction, and perenniality in the angiosperms. *New Phytologist*, *224*(3), 1266–1277.

https://doi.org/10.1111/nph.15999

Vaughton, G., Ramsey, M., & Johnson, S. D. (2010). Pollination and late-acting self-incompatibility

in Cyrtanthus breviflorus (Amaryllidaceae): Implications for seed production. *Annals of*

Botany, 106(4), 547–555. https://doi.org/10.1093/aob/mcq149

Villate, L., Esmenjaud, D., Van Helden, M., Stoeckel, S., & Plantard, O. (2010). Genetic signature

of amphimixis allows for the detection and fine scale localization of sexual reproduction

events in a mainly parthenogenetic nematode. *Molecular Ecology*, 19(5), 856–873.

https://doi.org/10.1111/j.1365-294X.2009.04511.x

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E.,

Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K.

J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., & van Mulbregt, P. (2020).

SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*,

17(3), Article 3. https://doi.org/10.1038/s41592-019-0686-2

Waits, L. P., Luikart, G., & Taberlet, P. (2001). Estimating the probability of identity among

genotypes in natural populations: Cautions and guidelines. *Molecular Ecology*, 10(1), 249–

256. https://doi.org/10.1046/j.1365-294X.2001.01185.x

Wang, Y.-J., Müller-Schärer, H., van Kleunen, M., Cai, A.-M., Zhang, P., Yan, R., Dong, B.-C., & Yu, F.-H. (2017). Invasive alien plants benefit more from clonal integration in heterogeneous environments than natives. *New Phytologist*, *216*(4), 1072–1078. https://doi.org/10.1111/nph.14820

Worley, A. C., & Barrett, S. C. H. (2000). Evolution of floral display in *Eichhornia paniculata* (Pontederiaceae): direct and correlated responses to selection on flower size and number. *Evolution*, *54*(5), 1533–1545. https://doi.org/10.1111/j.0014-3820.2000.tb00699.x

Wright, S. (1931). Evolution in Mendelian Populations. *Genetics*, *16*(2), 97–159.

Wright, S. (1949). The Genetical Structure of Populations. *Annals of Eugenics*, *15*(1), 323–354. https://doi.org/10.1111/j.1469-1809.1949.tb02451.x

Wright, S. I., Ness, R. W., Foxe, J. P., & Barrett, S. C. H. (2008). Genomic consequences of outcrossing and selfing in plants. *International Journal of Plant Sciences*, *169*(1), 105–118. https://doi.org/10.1086/523366

1098

1099

Figure 1: Distributions of Multi-Locus Genotypes (MLGs) among LSI and SC *Lgh* populations in western
Europe. In black, the number of unique MLGs; In grey, the number of MLGs with less than 7 ramets; In
color, MLGs with more than 10 ramets found in multiple populations. Proportions of each MLGs over all the
populations are plotted in the last right bar (All). More than half the samples are unique genotypes.

1100

1101

1102

1103

1104

1105

Figure 2: Correlations between Σ_{CLON} and Σ_{SELF} found in 13 SC populations (light grey points and regression line) and 40 LSI populations (black points and regression line). 95% confidence intervals are given for LSI and SC populations in dark grey and light grey, respectively. Spearman rank-order correlation coefficients (r_s) and probabilities that Σ_{CLON} and Σ_{SELF} would not be correlated (p) are reported for LSI and SC populations.

1111

1112
1113 **Figure 3:** Violin plots comparing distributions of genetic indices in LSI (L, grey) versus SC (S, white)
1114 populations. Violin plots are cut for their minimum and maximum values. Kruskall-Wallis tests are reported
1115 as H-statistics as well as the probability that LSI and SC population would present the same distribution of
1116 genetic indices.

1117
1118 **Figure 4:** Distributions of pairwise rho$_{st}$ within pairs of SC populations (white violin plot), pairs of LSI
1119 populations (light grey violin plot) and between pairs of LSI and SC populations (dark grey violin plot).
1120 Probabilities that pair of non-parametric distributions are equal are reported.

1121
1122   **Figure 5:** Minimum spanning tree of the genetic distances (number of different alleles) between LSI (grey
1123   points) and SC (white points) individuals. Arrows indicate some of the evidences of admixture between LSI
1124   and SC lineages.

Supplementary Information (SI)

Situation of *Lgh* in USA and Mediterranean populations in previous studies (Dandelot 2004, Okada et al. 2009, Armitage et al. 2013):
- nearly monoclonal populations arguing for the ecological advantage of clonality (fragmentation and rhizome propagation) in these invasive populations.
But
- using weakly polymorphic markers,
- before the demonstration of the heterostyly and of the Late-acting self-incompatibility system in *Lgh*

Situation of *Lgh* in western Europe, studied here:
- Spatial context: The other compatible type populations are away from hundreds of meters to several kilometers (Portillo-Lemus et al. 2021)
- Clonality through fragmentation and rhizomes, as in other populations (Portillo-Lemus 2021).
- Massive blossoming with a lot of pollinators harvesting flowers (Portillo-Lemus 2021).
- Faster pollen tube elongation of pollen from the other compatible type (Portillo-Lemus et al. 2022)

Our working hypotheses about the main, realized reproductive modes in population:
- **LSI populations:** A very limited production of seeds (Portillo-Lemus et al. 2021)
Hypothesis 1: **preferential clonality** due to lack of compatible partners to produce seeds → Baker's conjecture on reproductive modes (Pannell et al. 2015).
Hypothesis 2: The rare seeds really contribute to the populations → biological feature of seeds or recombination would have advantage over clones. If so, are they mainly:
    Selfed seeds → reproductive assurance and advantage of recombination over clonality in a uniparental context?
    Outcrossed seeds → advantage of outcrossed seeds due to heterosis and/or recombination despite the lack of locally-available compatible partners → selection of long pollination ranges

- **SC populations:** Production of a lot of seeds in natural populations
Hypothesis 1: despite a huge production of seeds, **preferential clonality** due to the physiological and/or ecological advantage of clones over seed germination process?
Hypothesis 2: the mass of seeds does contribute to the populations or some evolutionary advantage of recombination in offspring compared to clonal fragmentation and rhizome propagation to adapt local conditions? If so, are these seeds mainly:
    Selfed seeds → within uniparental reproduction, biological feature of seeds or genetic recombination would provide advantage over clones?
Outcrossed seeds → Faster pollen tube elongation of pollen from the other type gives real advantage or advantage to allogamy to adapt new local conditions?

Supplementary information1: Synthesis of the questionings of our study to estimate reproductive mode of

53 populations of invasive Lgh using genetic diversity on the autotetraploid part of its genome.

Genotypic or genetic index	Range of values	Main driver	Behavior	references
β*	$\in [0; +\infty]$	Rate of clonality	↘ when c ↗ $\beta \to 0$ when $c \to 1$	Halkett et al. 2005, Arnaud-Haond et al. 2007, 2020, Stoeckel et al. 2021
$varF_{IS}$	$\in [0; +\infty]$	Rate of clonality	↗ when c ↗ $VarF_{IS} \to 0$ when $c \to 0$	Stoeckel & Masson 2014, Stoeckel et al. 2021
r_d	$\in [0; 1]$	Rate of selfing	slightly ↗ when c ↗ ↗ when s ↗ $r_d \to 1$ when $s \to 1$	Weir and Cockerham 1973, Navascues et al. 2010, Halkett et al. 2005
H_O	$\in [0; 1]$	Rate of selfing	↘ when s ↗ $H_O \to 0$ when $s \to 1$	David et al. 2007, Halkett et al. 2005
F_{IS}	$\in [-1; 1]$	Both rates	↘ when c ↗ $FIS \to -1$ when $c \to 1$ $FIS \to +1$ when $s \to 1$	Stoeckel & Masson 2014, Balloux et al. 2003, Stoeckel et al. 2021
H_E	$\in [0; 1]$	Rate of selfing	↘ when s ↗ $H_E \to 0$ when $s \to 1$	Glémin et al. 2001, Ritland & Ganders 1987
P_{ID-SIB}	$\in [0; +\infty]$	Both rates	↘ when c ↗ ↘ when s ↗	Waits et al. 2001
S_g**	$\in [0; 1]$	Rate of selfing	↗ when s ↗	David et al. 2007, Hardy et al. 2016

* similar behavior of the clonal richness R
** inferred values of s, according to the method of Hardy (2016)

Table SI1: Hypotheses about the influence of clonality and selfing on the expected variations of genetic indices.

see the attached xls file

Table SI2: All genetic and genotypic indices computed on the 53 sampled populations. Each index was obtained from 15 individuals sampled in each population. (See spreadsheet supplementary material).

Population with missing genotype	nb_missing_genotypes	nb_individuals_concerned
Bléré_Ls	1	1
Chambéon_écopôle_du_Forez	2	2
Le_Port_Ls	1	1
Plougonver_Léguer	1	1
Quevillon	1	1
Rennes_apigné_prairie	1	1
Roost-Warendin	1	1
Saint_Aignan_Couflons_Ls	1	1
Vichy	1	1

Over single SNP genotypes	N	%age
single SNP genotype P(cad)>70%	28610	0.99965
single SNP genotype P(cad)≤70%	10	0.00035
total	28620	1

Over individuals	N	%age
nb_individuals_with_full_genotype	785	0.98742
nb_individuals_with_one_missing_genotype	10	0.01258
total	795	1

Table SI3: Summary of the allele dosage. First subtable lists populations with individual genotyped with at least one individual with missing genotype at one locus. Second subtable reports the frequency of single SNP genotypes with posterior probability of allele dosage strictly superior to 70%. Third subtable reports the frequency of individuals with their 36 SNP genotypes assigned with posterior probability superior to 70%.

Table SI4: Distribution of the different repeated Multi-Locus Genotypes (MLGs) found with 36 SNPs among populations considering their mating system (LSI or SC). Figure 1 presents a summarized plot of this table. (See spreadsheet supplementary material).

	G	H_O	H_E	A_E	R	β	F_{IS}	$VarF_{IS}$	r_d	PID_{SIB}	S_g	Σ_{clon}	Σ_{self}
G	-	0.438	**0.013**	**0.009**	***	***	***	***	0.057	**0.008**	**0.008**	***	***
H_O	0.076	-	***	***	0.397	0.462	0.122	**0.025**	0.653	***	**0.032**	0.065	0.142
H_E	**0.244**	**0.742**	-	***	**0.01**	**0.014**	0.212	***	**0.026**	***	0.844	***	0.372
A_E	**0.255**	**0.722**	**0.82**	-	**0.008**	**0.013**	0.353	***	0.094	***	0.347	***	0.845
R	**0.991**	0.083	**0.251**	0.258	-	***	***	***	0.052	**0.007**	**0.005**	***	***
β	**0.91**	0.071	**0.238**	**0.24**	0.919	-	***	***	0.126	**0.009**	**0.004**	***	***
F_{IS}	**0.602**	-0.144	0.116	0.086	**0.603**	0.557	-	***	***	0.163	***	***	***
$VarF_{IS}$	**-0.579**	**-0.209**	**-0.435**	**-0.346**	**-0.589**	**-0.551**	**-0.626**	-	***	***	***	***	***
r_d	0.187	0.042	**0.207**	0.156	0.19	0.148	**0.432**	-0.365	-	**0.021**	***	**0.006**	***
PID_{SIB}	**0.259**	**0.729**	**0.981**	**0.833**	**0.265**	**0.251**	0.13	**-0.443**	0.214	-	0.936	***	0.286
S_g	**0.264**	**-0.202**	-0.019	-0.089	**0.277**	**0.28**	0.542	-0.379	0.353	-0.008	-	***	***
Σ_{clon}	**-0.79**	**-0.171**	**-0.376**	**-0.347**	**-0.799**	**-0.809**	**-0.604**	**0.756**	-0.254	**-0.387**	**-0.34**	-	***
Σ_{self}	**0.388**	**-0.136**	0.083	0.018	**0.399**	**0.361**	**0.723**	**-0.524**	**0.631**	0.099	**0.69**	**-0.451**	-

Table SI5: Correlations between genetic indices using non-parametric Kendall partial rank-order correlation. Below the diagonal, the values of the coefficient of correlation τ are consigned. Above, the related p-values are provided. Values inferior to the threshold 0.05, and the related coefficient τ were bold. The parameters Σ_{CLON} and Σ_{SELF} correspond to the sum of normalized indices sensitive to clonality and selfing, respectively.

Figure SI1: Map of the locations of the 53 sampled populations across France and Spain. In orange,
populations with only L-morph individuals, supposed to mate under the control of a late-acting self-
incompatible (LSI); In purple, populations with only S-morph individuals, supposed to be self-
compatible (SC). In light blue, the Loire river system. In grey, the North-to-South (N-S) and West-to-
East (W-E) scale distances of the sampled populations. Underlined, the seven populations in which 15
individuals were tested for the self-incompatibility (Portillo-Lemus et al. 2021) and one individual was
karyotyped to count the chromosome numbers.

Figure SI2: Photos of somatic metaphase chromosomes indicating that L- and S-morph individuals
that succeed to mate and to give full fruit set and 100% viable first- and second-generation
descendants (Portillo-Lemus et al. 2021) also present the same number of chromosomes
(2n=10X=80), belonging to the same taxon *Ludwigia grandiflora* subsp. *hexapetala*. We still didn't
find *Ludwigia grandiflora* subsp. *grandiflora* karyotype in France (2n=6X=48, see Dandelot 2005,
Barloy et al. 2024). **A**: somatic metaphase chromosomes from L-morph (2n=10X=80); **B**: somatic
metaphase chromosomes from S-morph (2n=10X=80).

Figure SI3: Akaike's information criterion (AIC) as function of the ploidy level on the likelihoods of the
sequenced allele countings over all individuals. Tetraploidy, a ploidy 4x, presents the lowest AIC, and
thus the best supported model to explain distribution of sequenced allele countings over all
individuals, which confirm our initial SNP development to be sure of their ploidy level.

Figure SI4.A: Correlation circles of the principal component analysis on the 17 centered and scaled
genetic indices measured in 53 *Lgh* populations in western Europe. The first horizontal dimension
accounting for 50.24% of the variance regroups indices related to clonality (number of genotypes *G*,
genotypic diversity *R*, clonal heterogeneity *D*, clonal evenness *Beta_Pareto*, variance of Fis *varFis*,
variance of expected heterozygosity *varHe*, variance of observed heterozygosity *varHo* and variance
of effective number of alleles varAe between loci). The second vertical dimension accounting for
23.39% of the variance with a main contribution of estimate of selfing Sg. Other indices, *i.e.*, mean
observed heterozygosity *Ho*, mean expected heterozygosity *He*, effective number of alleles *Ae*,
inbreeding coefficient *Mfis*, linkage disequilibrium *rd*, standard error of estimate of selfing *SE.Sg*,
unbiased probability of identity under panmixia *PIDu* and between sibs *PIDsib*. In blue and dashed
lines, the predictions of Σclon (*Sclon*) and Σself (*Sself*) provided as supplementary variables.

37
38 **Figure SI4.B:** Bar plot of the genetic index contributions to the first principal component. The
39 expected average contribution *1/(number of genetic indices)* is plotted as the red dashed line. VarFis
40 and indices of genotypic diversity (Pareto , R, G & D) are the main contributors to the first
41 component.

42
Figure SI4.C: Bar plot of the genetic index contributions to the second principal component. The
expected average contribution *1/(number of genetic indices)* is plotted as the red dashed line.
Estimates of selfing (Sg), mean observed heterozygosity (H_O), mean effective number of alleles (A_E)
and probabilities of identities (pid_u and pid_{sib}) are the main contributors to the second component.

Figure SI4.D: Plot of the projection of each population onto the two first principal components. SC
populations are plotted as red triangles and LSI populations as black diamonds. Concentration
ellipses around LSI and SC populations are plotted in black and red, respectively. Barycenters of the
two groups are represented as bigger black diamond and red triangle, respectively.

Figure SI5: Relationship and correlation between Σ_{CLON} and genetic indices (R, Pareto β, MFis,
VarFis, Ho, Ae, Sg, pidsib and $\bar{r}_d$). Per graph, all the 53 populations were plotted as black
dots. *: p<0.05, **: p<0.01, ***: p<0.001.

Figure SI6: Relationship and correlation between Σ_{SELF} and genetic indices (R, Pareto β, MFis, VarFis, Ho, Ae, Sg, pidsib and $\bar{r}_d$). Per graph, all the 53 populations were plotted as black dots. *: p<0.05, **: p<0.01, ***: p<0.001.

62 **Figure SI7:** Distributions of the logarithms of the number of ramets per genet in LSI (grey) and
63 SC (white) populations. We report the Kruskal-Wallis probability that the two distributions
64 are identical.

Origin of the higher genetic diversity in LSI populations as compared to SC

A first hypothesis to explain the genetic differences found between LSI and SC populations may be due to the outcrossing consequences on LSI populations. Indeed, similar genetic differences between SC and SI were recurrently reported in populations mating with sporophytic and gametophytic SI systems (Igic et al. 2008, Mable 2008). For example, *Leavenworthia alabamica* and *L. crassa* are two sister-species mating with a sporophytic SI within a small genus of eight annual species in which self-fertilization has evolved at least four times (Beck et al. 2006). In these species, comparisons between SI and SC populations also showed higher mean genetic diversity in SI populations as compared to the SC, in an overall background of high genetic diversity (Busch 2005, Koelling et al. 2011). High Ho, negative Fis and high effective number of alleles per marker are also found in species mating with gametophytic SI systems, such as in the *Prunus* genius (Stoeckel et al. 2006, Stoeckel et al. 2008, Abdallah et al. 2019). However, theoretical predictions argue that this kind of higher genetic diversity favored and maintained in SI populations as compared to panmictic and SC populations must only be found on polymorphic markers highly physically linked to the genomic regions involved in the SI mechanisms that evolve under balancing selection, and also, at a lesser magnitude, on physically unlinked markers but only in small population, with less than 100 individuals including clones (Glémin et al. 2001, Navascues et al. 2010). The 36 SNPs we developed must randomly distribute over the 16 chromosomes inherited twice from a common ancestor with *Lpm*, i.e., 32 chromosomes out of 80 total chromosomes of *Lgh* (Barloy et al. 2024), with no reason that all aggregate around the genes involved in the LSI self-recognition. In addition, individuals and populations of *Lpm* are reported to massively use self-fertilization, thus with no SI system (Estes& Thorp 1974, Grewel et al. 2016). By parsimony, there is less chance that genes involved in the LSI reaction would be stored on

these chromosomes than on some of the 48 other chromosomes; and therefore, very unlikely

that our SNPs would be physically linked to genes under balanced selection due to LSI. *Lgh*

populations in western Europe include far more than one thousand stems per local population

(Portillo-Lemus et al. 2021). It is thus unlikely that differences in genetic diversity observed

between LSI and SC populations would directly result from the consequences of the LSI system

hitchhiked to the whole genome (Navascues et al. 2010).

The second hypothesis is that high genetic diversity in LSI populations may arise and be

maintained due to higher rates of clonality (Stoeckel & Masson 2014, Reichel et al. 2016,

Stoeckel et al. 2021a) as compared to SC populations. In LSI, self-pollen tubes physiologically

blocked at the micropyle may prevent outcrossed-pollen tubes to reach the available ovules,

decreasing female fertility and increasing the importance of clonal regeneration in

populations (Vaughton et al. 2010). In addition, clonality is another uniparental reproductive

mode that allows the local maintenance of populations in the absence of sexually compatible

partners, which is more probable in peripatric conditions (Pannel et al. 2015). Dominant clonal

reproduction and propagation are commonly found in peripatric populations extending their

living range into new areas (Pyšek 1998, Wang et al. 2017). Furthermore, there is an

established intriguing co-occurrence of clonality in SI populations of flowering plants (Vallejo-

Marin et al. 2010). We know that dominant clonal reproduction theoretically favours the

accumulation of deleterious mutations and that these deleterious mutations contribute to the

advantage of outcrossing, which decreases the probability of inbreeding depression in

offsprings, as compared to SC reproducers (Navascues et al. 2010). The observed genetic

differences between LSI and SC populations may thus result from higher clonality in LSI

populations lacking of compatible partners. But we found here no difference in estimates of

clonality between LSI and SC populations, in Σ_{CLON} and in all indices sensible to clonality (R,

Pareto Beta and VarFis; Figure 3).

We rather found significant difference in estimates of selfing between LSI and SC populations,

in Sg (Hardy 2016), in Σ_{SELF} and in all indices sensible to selfing (Ae, linkage disequilibrium, Ho,

mean Fis) revealing typical signatures of higher selfing rates in SC populations. Using Hardy's

method (2016) extended from David et al. (2007) for autopolyploids, SC populations were

estimated to reproduce using a mean selfing rate of 0.18 while LSI populations with only a

mean selfing rate of 0.05, both estimated with similar rates of sexuality. Congruent with higher

estimates of selfing rates and Σ_{SELF} in SC populations, we found higher linkage disequilibrium,

lower observed heterozygosity and less effective number of alleles in SC populations (Figure

3). The observed differences in genetic diversity between LSI and SC populations must thus be

rather caused by the genetic consequences of higher selfing in SC populations, decreasing

their genetic diversity, than due to a direct outcrossing effect of the LSI hitchhiked over all the

Lgh genome within these large populations of *Lgh* (see Navascues et al. 2010). Interestingly,

the outcrossing breeding system allowed by LSI, with a small rate of selfing allowed especially

at the end of the flowering season, was measurable at the population level in genotyping (5%

of selfing in LSI versus 18% in SC) and thus may be efficient at an eco-evolutionary scale.

The recent arrival of *Lgh* in western Europe may also have contributed to shape these genetic

differences. Recent founding events in SC populations are indeed expected to decrease the

overall genetic diversity all along the genomes, including fewer effective alleles, lower gene

diversity, and potentially more linkage disequilibrium as compared to outcrossing populations

(Flanagan et al. 2021). Despite the ρ_{ST} values we observed between pairs of SI and SC

populations being approximately double those measured within pairs of SC populations and

within pairs of LSI populations, we also found clues of admixture in the genetic distances

between LSI and SC individuals in western Europe (Figure 5). This result would thus tend to

minimize the contribution of the foundation in the differences in genetic diversity we

observed between LSI and SC populations, and stress again the importance of selfing to shape

the genetic differences we found between SC and LSI populations.

141