Title: **Reproductive modes of polyploid *Ludwigia grandiflora subsp. hexapetala* in western Europe: the effects of a late-acting self-incompatibility system and its absence on genetic diversity within populations**

Short running title: Mixed reproductive modes in *Ludwigia*

Authors: Solenn Stoeckel[1,2], Ronan Becheler[1,2], Luis Portillo-Lemus[1], Marilyne Harang[1], Anne-Laure Besnard[1], Gilles Lassalle[1], Romain Causse-Védrines[3], Sophie Michon-Coudouel[3], Daniel J. Park[4], Bernard J. Pope[4], Eric J. Petit[1], Dominique Barloy[1]

Affiliations

1 DECOD (Ecosystem Dynamics and Sustainability), Institut Agro, IFREMER, INRAE, Rennes, France.

2 IGEPP, INRAE, Institut Agro, Université de Rennes, Le Rheu, France.

3 Plateforme EcogenO, UAR3343 OSUR, Université de Rennes, CNRS, Rennes, France.

4 Melbourne Bioinformatics, The University of Melbourne, Parkville, Australia.

Corresponding author: solenn.stoeckel@inrae.fr

Abstract

Reproductive mode, *i.e.,* the proportion of individuals in populations produced by clonality, selfing and outcrossing in populations, determines how hereditary material is transmitted through generations. It shapes genetic diversity and its structure over time and space.

Ludwigia grandiflora subsp. *hexapetala* (*Lgh*) is a partially clonal, polyploid, hermaphroditic, and heteromorphic plant that recently colonized multiple countries worldwide. In western Europe, individuals in this species are either self-incompatible caused by a late-acting self-incompatibility (LSI) system developing long-styled flowers, or self-compatible (SC) developing short-styled flowers. We do not yet know if the LSI system is effective in driving genetic diversity within populations, as previously showed for gametophytic and sporophytic SI systems. In addition, determining the reproductive mode of newly established *Lgh* populations in Europe will contribute to understand their ecological and evolutionary roles at the margins of species distribution. However, we still lack of a full approach including the essential step to unfold population genetic indices to estimates rates of clonality, selfing and outcrossing in autopolyploids.

In this study, we proposed such an approach to measure genetic diversity to assess reproductive modes on 53 LSI and SC populations on newly colonizing France and northern Spain, using SNPs amplifying on a distinct autotetraploid part of the *Lgh* genome. SNPs are easily reproducible, adapted for intraspecific genetic studies and allowed confident allele dosage to genotype autopolyploid individuals. Such genotyping data in turn made it possible to use recent methodological developments to compute and interpret genetic diversity to assess reproductive mode in each population.

We found that populations reproduced mainly clonally but with a high diversity of genotypes and with substantial estimated rates of sexual events, up to 40%. We also found evidence for

local admixture between LSI and SC populations in a background of genetic structure between pairs of LSI and SC populations that was twice the ones found among pairs of LSI populations or pairs of SC populations, arguing that SC and LSI populations may follow distinct expansion dynamics. These evidences of sexual events in nearly all populations and of high diversity in clonal lineages changes the way we think about managing *Lgh* worldwide invasive populations.

We also took advantage of the spatial segregation of LSI and SC populations invading France to characterize the population genetic consequences of an LSI system. LSI and SC populations showed similar rates of clonality but significantly different rates of selfing, as expected considering their breeding system, and despite the small rates of failure of LSI system. Within the 53 studied populations, the 13 SC-only populations distinguished with fewer effective alleles, lower observed heterozygosity, and higher inbreeding coefficients, linkage disequilibrium and estimates of selfing than found in LSI populations. These results suggested that genetic differences significantly found between LSI and SC populations mainly came from the increased selfing rates in SC populations rather than due to the genome-wide consequences of outcrossing in LSI populations or due to different rates of clonality. The overall maintenance of higher genetic diversity, with the possibility of resorting to clonality, selfing and outcrossing altogether, may explain why LSI populations seem to be more prevalent in all newly expanding populations worldwide.

Beyond *Lgh*, our methodological approach may inspire future studies to assess reproductive mode of other autopolyploid eukaryote populations, and our results emphasize the necessity to consider the variations of reproductive modes when managing invasive plant species.

Keywords: Partial clonality, Mating system, Late-acting self-incompatibility system, Selfing, Outcrossing, Autopolyploidy, Water primrose

Introduction

Plants can reproduce using different mechanisms (i.e., breeding system), including clonality and different types of sexuality (i.e., mating system, including allogamy and selfing; Holsinger 2000). Reproductive mode corresponds to the actual balance between sexual and clonal reproduction, measured by the rate of clonality (De Meeus et al. 2007; Stoeckel et al. 2021a), and, within the proportion of offspring produced by sexuality, the balance between autogamy and geitonogamy (hereafter *selfing*) and allogamy (*outcrossing*) measured by the rate of selfing (Bürkli et al. 2017). Reproductive mode is one of the main drivers of genetic diversity in populations and species (Duminil et al. 2007, Ellegren & Galtier 2016, Glémin et al. 2019). It determines the way hereditary material is transmitted over generations, and thus constrains the range of genetic diversity that can evolve within populations and species (De Meeus et al. 2007, Fehrer 2010, Stoeckel & Masson 2014, Rouger et al. 2016). Its genetic consequences are deep enough to be discernible even in the presence of other affecting factors (Charlesworth 2003). Considering its influence on genetic diversity and its structure in populations, reproductive mode is one of the major biological traits to decipher before interpreting other biological and ecological forces using molecular data (Duminil et al. 2007, Reichel et al. 2016, Bürkli et al. 2017, Orive & Krueger-Hadfield 2021). Given an adequate theoretical framework, population geneticists can use genotypic data to infer reproductive modes in populations (Halkett et al. 2005, Arnaud-Haond et al. 2007, Hardy 2016, Bürkli et al. 2017, Stoeckel et al. 2021a).

Approximately 80% of plant species are partially clonal (Barrett 2010, Vallejo-Marin et al. 2010). Clonality consists of a parent producing a new descendant that is a genetic copy of itself with the exception of rare somatic mutations and mitotic recombinations (De Meeus et al.

2007). Clonality generates repeated genotypes in populations when parents can breed

multiple descendants (Arnaud-Haond et al. 2007). This yields an average excess of

heterozygotes compared to Hardy-Weinberg expectations (*i.e.,* negative F_{IS}), large variance of

F_{IS} among polymorphic markers along the genome and a small excess of linkage

disequilibrium, increasing with decreasing population size (Navascues et al. 2010, Stoeckel &

Masson 2014, Stoeckel et al. 2021a). These effects vary with the rates of clonality and, in turn,

can be used to infer the rates of clonality themselves within populations (De Meeus et al.

2006, Becheler et al. 2020, Arnaud-Haond et al. 2020).

Approximately 70% of angiosperm species are hermaphroditic (Dellaporta & Calderon-

Urrea 1993) and have thus the possibility to sexually reproduce with themselves (selfing).

Selfing decreases effective population sizes and thus fastens the effect of genetic drift, leading

gene diversity to decrease in populations and even to allele fixing (Wright et al. 2008, Roze

2016, Glémin et al. 2019). Selfing also prohibits genetic mixing between different ancestral

lineages, resulting in decreased heterozygosity within individuals (David et al. 2007,

Hardy 2016) and increased linkage disequilibrium between genes along genomes (Golding &

Strobeck 1980, Norborg 2000, Lucek & Willi 2021).

Around half of hermaphrodite plants restrict self-fertilization using a variety of molecular

mechanisms grouped under the term "self-incompatibility (SI) systems" which favour

outcrossing within populations (De Nettancourt 2001, Castric & Vekemans 2004, Gibbs 2014,

Steinecke et al. 2022). Outcrossing is overall expected to increase genetic diversity, to limit

allele fixation and to reduce linkage disequilibrium when compared to selfing. However, such

expectations depend on the genetic linkage between the considered loci and the genomic

region carrying the genes involved in self-recognition (Glémin et al. 2001, Stoeckel et al. 2006,

Navascues et al. 2010). Within the different identified SI systems (Charlesworth et al. 2005,

Franklin-Tong 2008), the late-acting SI system (hereafter LSI) is still poorly studied despite

being identified in multiple taxonomic groups in angiosperms (De Nettancourt 1997,

Gibbs 2014). In LSI systems, self-pollen tubes grow and are only blocked shortly before

penetrating the ovule, or at the micropyle, or, in some species, the ovules are penetrated but

their syngamy fails. Such species may present reduced female fertility due to self-pollen

disabling ovules, favouring the clonal regeneration of populations (Vaughton et al. 2010). LSI

systems are also characterized by low but recurrent failures of the self-recognition system due

to its late mechanism, leading to the production of a low amount of selfed seeds in populations

(Seavey & Bawa 1986; Chen et al. 2012, Gibbs 2014). Gametophytic and sporophytic self-

recognition mechanism occurring very early in the pistil drastically limit self-pollination thus

selfed seeds (Lawrence 1985, Gibbs 2014) and need specific chemical or physical treatments

to be overcome (Cabin et al. 1996, Ahmad et al. 2022). We however don't know yet if these

rare selfed seeds may contribute to the dynamics of genetic diversity of LSI populations and

species, especially in peripatric conditions or in any situation when compatible partners may

lack. Due to these selfed seeds, LSI system may even be less effective in driving genetic

diversity within populations, as already described previously for gametophytic and

sporophytic SI systems (Brennan et al. 2002, Stoeckel et al. 2006, Koelling et al. 2011, Busch &

Urban 2011). In such situation, the LSI systems could become a new evolutionary puzzle

among the reproductive systems.

Autopolyploidy is pervasive in plants whether it derives from whole genome duplication or

from multiple hybridization events (Baduel et al. 2018). Clonality and selfing tend to occur

more frequently in polyploids than in their diploid counterparts (Husband et al. 2008, Herben

et al. 2017, Van Drunen & Husband 2018). Autopolyploid species bring two more main

challenges when tackling population genetics (Dufresne et al. 2014): The first issue is achieving

confident allele dosage to obtain a robust genotype for each individual at each locus. For

instance, disentangling between AAAC, ACCC and AACC genotypes in an autotetraploid

requires a reproducible and robust quantitative approach to genotyping. Second, computing

values of genetic diversity, identities and heterozygosity within autopolyploid populations

requires adapted methods (Hardy 2016, Meirmans et al. 2018). For instance, in

autopolyploids, heterozygosity, i.e., the probability that two random alleles are not identical-

in-state, can take more different values than in the case of diploids (Haldane 1930, Hardy

2016). Autopolyploids require dedicated computations of population genetic indices to not

result in artefactual population genetic values. These dedicated methods can now be

commonly computed using dedicated software such as Spagedi (Hardy & Vekemans 2002) and

GenAPoPop (Stoeckel et al. 2024).

Water primrose, *Ludwigia grandiflora* subsp. *hexapetala* (Hook. & Arn.) Nesom and Kartesz

(2000), hereafter *Lgh*, is one of the most invasive aquatic plants in the world (Thouvenot et al.

2013). *Lgh* is an alloautodecaploid species (2n=10x=80 chromosomes), resulting from

hybridization of different ancestral diploid species, some of which are represented more than

once in the total genome of *Lgh*, which belongs to the genus *Ludwigia* L. section Jussiaea (Hoch

et al. 2015, Barloy et al, 2024). Interestingly, *Lgh* includes an autotetraploid set of

chromosomes, shared with *L. peploides* subsp. *montevidensis* (2n=2x=16), hereafter *Lpm*, that

is clearly distinct from the other ancestral part of *Lgh* genome (Barloy et al. 2024). *Lpm* is a

self-compatible-only diploid species with only one common floral morphology (Estes & Thorp

1974, Grewel et al. 2016). On its side, *Lgh* presents heteromorphic flowers corresponding to

two floral morphs: L-morph individuals develop long-styled flowers and S-morph individuals

develop short-styled flowers (Portillo-Lemus et al. 2021). L-morph flowers express an active

LSI in western European populations (Portillo-Lemus et al. 2022). During the core flowering

season (summer), in experimental greenhouse conditions, L-morph individuals show a stable

seemingly insignificant rate of autogamy (around 0.2‰ of the available ovules) that increases

at the end of the flowering season, during autumn, to 1‰, which is common in LSI systems

(Gibbs 2014. Due to the massive blossoming of this species, growing in very dense populations,

these selfed seeds would add up yearly in field populations to thousands of seeds per metre

square (Portillo-Lemus et al. 2021). This pattern of low rates of autogamy, that increase at the

end of the flowering season, may provide the advantage of reproductive assurance

(Goodwillie & Weber 2018). In contrast, S-morph individuals are self-compatible (SC) in

western European populations but pollen tubes of the L-morph grow significantly faster and

are thus advantaged to fertilize ovules when in competition with self-pollen (Portillo-Lemus

et al. 2022). In addition, peripatric *Lgh* populations, including European populations, were

previously reported as exclusively clonal with few clonal lines (Dandelot 2004, Okada et al.

2009). Recently established populations in France and northern Spain mostly present only one

of the two compatibility modes locally, sometimes with a population of the other type a few

to tens kilometers away, which may result in effective allogamy (Portillo-Lemus et al. 2021,

2022). All these different breeding systems make possible very different reproductive modes

in populations, comprising all possible quantitative combinations of mainly clonal,

autogamous and allogamous modes. Genotyping with confident allele dosage an

autotetraploid inherited genome, analyzing the genotyping to obtain population genetic

measures to estimate the respective and quantitative contributions of these different

reproductive modes still constitute challenging tasks in autopolyploid species as pointed out

by Dufresne et al. (2014)'s review and synthesis. If using genetic diversity within population to

estimate rates of clonality versus sexuality, and of selfing versus allogamy is now commonly

achieved for diploids and haplodiplontics (David et al. 2007, Arnaud-Haond et al. 2020,

Krueger-Hadfield et al. 2021, Stoeckel et al. 2021a,b), we still lack of such a framework to

estimate the reproductive modes in autopolyploid populations with their specificities, with an

applied case study which would demonstrate its effectiveness.

In this study, we proposed a complete framework, from genotyping autotetraploid part of *Lgh*

genome with allele dosage to the estimation of reproductive modes per population using

genetic data, compiling recent molecular and software developments (Besnard et al. 2023,

Hardy 2016, Stoeckel et al. 2024) to fill the gaps identified by Dufresne et al. (2014) for

confidently genotyping autopolyploid individuals. First, we developed a codominant and

reproducible set of single-nucleotide polymorphism (SNP) markers for this species that are

more adapted for intraspecific genetic studies, being able to distinguish between siblings and

different clonal lineages, and that allowed to perform reproducible and confident allele

dosage. Indeed, allele dosage at each marker within each individual is mandatory to compute

confident population genetic measures within populations that have no reason to follow

genotype frequencies expected under Hardy-Weinberg proportions, being partially clonal,

outcrossed and selfed. To be sure of the ploidy level of the studied markers and that the

genotyped markers did not interact with genomes of the other ancestral species of *Lgh*

genome, we genotyped SNPs only the autotetraploid part of *Lgh* genome shared with the

diploid species *Lpm*. The *Lpm* genome is sufficiently genetically distinct from the other

ancestral genomes of *Lgh* to avoid misgenotyping (Barloy et al. 2024).

We then used this approach to identify and study the variability of the reproductive modes of

Lgh populations within the invaded region of Western Europe considering their LSI and SC

status. To assess the reproductive modes of each sampled population and estimate their rates

of clonality and selfing, we measured gene diversity, heterozygosity, linkage disequilibrium

and genotypic diversities within 53 populations sampled in western Europe. Being able to

assess reproductive modes in each population, we will discuss the potential advantage of

uniparental reproduction including clonality and selfing, of recombination (selfing and

allogamy) and of heterozygosity in such peripatric invasive populations.

Second, due to the fact that recent local populations in France and northern Spain present

only one of the two compatibility modes, we took advantage of this contrasted situation

within the same species and in comparable ecological situations to compare genetic diversity

among LSI and SC populations, to assess the influence of LSI on genetic diversity as similarly

tackled in sporophytic self-incompatible and SC Brassicaceae species (Koelling et al. 2011).

Indeed, genetic diversity underlies the evolvability of species and their survival and adaptation

to environmental change and ecosystem resilience (Booy et al. 2008, Hoban et al. 2021). The

small proportions of selfed seeds allowed by the LSI mechanism may mitigate the difference

in genetic diversity between LSI and SC populations compared to sporophytic and

gamethophytic self-incompatible systems. Considering the complex case of *Lgh* in western

European populations, we hypothesized that LSI populations should present typical genetic

footprints of dominant outcrossing, and perhaps higher rates of clonality due to the local lack

of compatible partners and self-pollen interferences, while SC populations should present

higher rates of sexuality prevailed by selfing. To understand the genetic impacts of

reproductive modes, we also quantified the covariations of genetic indices and their

importance to define the genetic diversity within the 53 genotyped populations, and

compared these observations to the theoretical expectations obtained from a Wright-Fisher-

like model extended to autotetraploids (Stoeckel et al. 2024). Finally, we interpreted the

genetic differences found within and between LSI and SC populations to understand how

reproductive mode(s) contributed to the genetic diversity in these peripatric *Lgh* populations.

Materials and methods

Sampling strategy and floral morphology of populations

We collected 795 stems of *Lgh* from 53 locations (52 locations in France and one in Catalonia, Spain), corresponding to an area that spans 880 kilometres east-to-west and 1,100 kilometres north-to-south (Figure SI1). At each location, we collected 15 stems (hereafter individual) along a linear transect measuring 40 meters length. Along each transect, we randomly collected 3 stems at coordinates $X_1 = 0m$; $X_2 = 10m$; $X_3 = 20m$; $X_4 = 30m$ and $X_5 = 40m$ within a 1 meter-square quadrat. The young leaves of each sampled individual were stored after lyophilization until DNA extraction. We visually identified floral morphologies of flowers found along each transect within the sampled *Lgh* populations.

Definition of the SNP marker set

As no molecular markers suitable for clonal discrimination were yet available for *Lgh* and *Lpm*, we generated an original set of SNP markers using RAD-Seq strategy (Baird et al., 2008) from five and three *Lgh* and *Lpm* western European populations, respectively using a pool of 15 individuals sampled for each population. RAD DNA library, RAD-sequencing and pipeline to identify *Lgh* SNP markers were carried out as described in Delord et al. (2018). To summary, DNA of *Lgh* and *Lpm* were digested by *Sbfl* restriction enzyme and used to prepare DNA libraries then sequenced using a illumine HiSeq3000 (150bp paired-end reads). After different steps of filtering, a total of 14,233 and 34,287 RAD-Seq was found in *Lgh* and *Lpm* respectively. At the end, 340 and 326 SNPs were obtained for *Lpm* and *Lgh*, respectively. Design of primers compatible with Hi-Plex multiplexing was carried out at the Melbourne Bioinformatics lab. Finally, sixty and fifty SNP markers were obtained for *Lpm* and *Lgh*, respectively. In our study,

we only considered polymorphic SNPs that were shared between *Lgh* and *Lpm* to be sure they

belong to the tetraploid part of *Lgh* genome derived from *Lpm* (Barloy et al. 2024). We finally

kept a set of 36 polymorphic and stable SNP markers to genotype *Lgh* samples and analyze

genetic diversity in each sampled population (primer sequences are openly listed in Stoeckel

et al. 2023).

DNA extraction, SNP marker production and genotyping

To genotype the 795 individually-tagged samples with these 36 SNPs, we extracted DNA from

25 to 30 mg of dried young leaf tissues. The DNA extractions were performed using the

NucleoSpin Plant II from MACHEREY-NAGEL kit, following the manufacturer's

recommendations. We used L1 buffer as lysis buffer. All individuals were genotyped from a

solution of 20ng/µl of *Lgh* DNA, using a modified version of the Hi-Plex protocol (Hammet et

al. 2019; Besnard et al. 2023). Hi-Plex is an amplicon sequencing technique (*sensu* Meek &

Larson 2019) in which all SNPs are co-amplified in a multiplex reaction before Illumina or Ion-

Torrent sequencing. Here, we used Illumina. Intermediate steps include dual indexing of

individual samples used for demultiplexing. Reads were then assigned to loci by aligning them

to reference sequences with BWA-MEM 0.7.15-r1140 (Li & Durbin 2010) and alleles were

counted with Samtools 1.9 (Danecek et al. 2021).

Allele dosage

The posterior probabilities of each single SNP genotype within each individual (hereafter

single SNP genotype) were computed using the likelihood of all possible genotypes considering

a multinomial distribution of the number of times each nucleobase was genotyped at one SNP

locus within one individual. In order to obtain the most confident dataset possible, we only

assigned a genotype when its posterior probability of allele dosage exceeded 70%. When one

SNP within one individual presented a posterior probability of allele dosage equal or lower

than 70%, we labelled and analysed it afterward as a missing genotype.

Genotypic and genetic descriptors

We expected that populations reproducing clonally may yield repeated multi-locus genotypes

(MLGs, i.e., the same genotype at all the 36 SNPs found in multiple individuals). By possibly

producing these repeated genotypes and by varying the relative distribution of the number of

samples of each of these distinct genotypes, rates of clonality impact genotypic richness and

evenness in populations (Halkett et al. 2005; Arnaud-Haond et al. 2007).

We measured genotypic richness using the R index (Dorken & Eckert 2001, Arnaud-Haond et

al. 2005), which is defined as:

$R = \frac{G-1}{N_g - 1}$

where G is the number of distinct genotypes (genets) and N_g is the number of genotyped

individuals. We also measured genotypic evenness as the parameter β of the Pareto

distribution, which describes the slope of the power-law inverse cumulative distribution of the

size of MLGs (Arnaud-Haond et al. 2007):

$N \geq X = aX - \beta$

where N≥X is the number of sampled ramets belonging to genets containing X or more ramets

in the sample of the population studied, and the parameters *a* and *β* are fitted by regression

analysis. Low β-values indicate the dominance of one or few clonal lineages. Pareto β is less

biased by the sampling effort than clonal richness R (Stoeckel et al. 2021a). In our sample of

15 individuals, we expected a population with only one repeated MLG to present a Pareto β

value of 0.03 and a value of 3 in samples with no repeated genotypes. β < 2 indicates a

population reproducing with high rates of clonality (greater than 0.8 to 0.9). For a sample size

of 15 individuals, 2 < β < 3 indicates intermediate to low rates of clonality (0.6 to 0.8). A β at

3, its maximum value, is indicative of a mainly sexual population, with rates of clonality ranging

from zero to 0.6, depending on the other genetic indices.

Rates of clonality and selfing confer different effects on the range of within-population genetic

polymorphism as well as on probabilities of genetic identities within individuals. We thus also

estimated expected and observed heterozygosity H_E and H_O, allelic richness A_E, inbreeding

coefficient F_{IS}, and linkage disequilibrium within each sampled population.

The Wright (1931, 1949) inbreeding coefficient F_{IS} accounts for intraindividual genetic

variation at one locus as a departure from Hardy-Weinberg assumptions of the genotyped

populations. We computed one F_{IS} value per locus per population. We then reported the mean

values of F_{IS} (MF_{IS}) and the inter-locus variance of F_{IS} ($VarF_{IS}$) for each population. Both MF_{IS}

and $VarF_{IS}$ are very informative about the underlying reproductive systems (Stoeckel et al.

2006, David et al. 2007). Clonality makes the MF_{IS} values tend toward -1 along with high

interlocus variance (Stoeckel & Masson 2014, Stoeckel et al. 2021a). A moderate amount of

sexual reproduction results in MF_{IS} values around 0 (Balloux et al. 2003). $VarF_{IS}$ varies with

rates of clonality, from very limited variance expected in sexual populations to high variance

in very clonal populations (Stoeckel & Masson 2014, Stoeckel et al. 2021a). Positive MF_{IS}

values are expected in populations reproducing using consanguinity and selfing (Castric et al.

2002, David et al. 2007). All sexual reproductive modes, allogamous and autogamous, result

in low $VarF_{IS}$ within a population as massive recombination tends to homogenize intra-

individual genetic identities along the genomes (Stoeckel et al. 2021a).

We measured linkage disequilibrium over all markers using the unbiased multilocus linkage

disequilibrium index, $\bar{r}_d$ (Agapow & Burt 2001). This mean correlation coefficient (r) of genetic

distances (d) between unordered alleles at n loci ranges from 0 to 1. This metric has the

advantage of limiting the dependency of the correlation coefficient on the number of alleles

and loci, and it is well suited to measure linkage disequilibrium in partially clonal populations

(De Meeus & Balloux 2004, De Meeus et al. 2006). In general, LD is only slightly affected by

clonality, except when clonality is high (c > 0.9) and/or when genetic drift dominates over

mutation rate (*e.g.,* when population sizes are small N<50 compared to u=0.001; Navascues

et al. 2010, Stoeckel et al. 2021a). In contrast, inbreeding and selfing are efficient processes

for quickly generating strong LD, after only few generations. Finally, we measured genetic

differentiation between populations of LSI, between populations of SC and between LSI and

SC populations using ρ_{ST}, an index adapted to study autopolyploid populations (Ronfort et al.

1998). All these indices were computed using GENAPOPOP (Stoeckel et al. 2024), a software

dedicated to analysing genetic diversity and differentiation in autopolyploid populations

genotyped with confident allele dosage and reproducing through all possible rates of clonality

and selfing.

We also used SPAGEDI (v1.5, Hardy & Vekemans 2002) to estimate rates of selfing within

autopolyploid populations genotyped with confident allele dosage. This approach infers rate

of selfing (Sg) from identity disequilibrium coefficients (g_{2z} estimator), assuming that

populations reproduce by self-fertilization and random mating and are at inbreeding

equilibrium (David et al. 2007 for diploids, Hardy 2016 for autopolyploids). Identity

disequilibrium coefficients are measured as the correlation in heterozygosity of distinct loci

within the genome, and present the advantage of being more robust to null alleles and

genotyping errors than raw F_{IS} (David et al., 2007). The effect of partial clonality on g_{2z}

estimator is not yet defined. We thus considered for this study that clonality would only

marginally impact identity disequilibrium coefficients, the possibility to be at the inbreeding

equilibrium and the corresponding estimates of selfing. A synthesis of all these hypotheses

based on previous knowledge is provided in Table SI1.

Ranking populations by clonality and selfing

We proposed and computed a synthesis index (Σclon) calculated from Pareto β and VarF$_{IS}$ to

rank the studied populations from the less to the more clonal. These two population genetic

indices are known to vary with rates of clonality being unbiased in samples (Stoeckel et al.

2021a). To avoid issues of scaling as the range of values of these descriptors are different by

several orders of magnitude, Σclon is computed in each population i as the sum of Pareto $\beta(i)$

and VarFIS(i) values that were previously normalized over the whole dataset, respectively

$z_\beta(i)$ and $z_{\text{VarFIS}}(i)$.

367
$$z_\beta(i) = 1 - \frac{\beta(i) - min(\text{B})}{max(\text{B}) - min(\text{B})}$$

368
$$z_{\text{VarFIS}}(i) = \frac{\text{VarFIS}(i) - min(\text{VF})}{max(\text{VF}) - min(\text{VF})}$$

369
$$\Sigma\text{clon}(i) = \frac{1}{2}\left[z_\beta(i) + z_{\text{VarFIS}}(i)\right]$$

where $\beta(i)$ and VarFIS(i) are the measured Pareto β and variance of F$_{IS}$ over loci in the

population i, and where $\text{B} = \{\beta(1), \beta(2), \dots, \beta(55)\}$ and $VF =$

$\{\text{VarFIS}(1), \text{VarFIS}(2), \dots, \text{VarFIS}(55)\}$ are the respective sets of Pareto β and VarF$_{IS}$ over loci

in the 53 populations used to obtain their minimum (min) and maximum (max) values.

Using the same approach as for Σclon, we computed a synthesis index (Σself) calculated from

$\bar{r}_d$, MF$_{IS}$ and Sg to rank the studied populations from the less to the more selfed.

376
$$\Sigma\text{self}(i) = \frac{1}{3}\left[z_{\bar{r}_d}(i) + z_{\text{MFIS}}(i) + z_{\text{Sg}}(i)\right]$$

Statistical data analysis

First, to better understand the correlation between population genetic indices and estimates of reproductive modes from a large field dataset and to compare with the theoretical correlations obtained from simulations (Stoeckel et al. 2024), we computed a Principal Component Analysis (PCA) to comprehend the covariations of genetic indices, the correlations and the redundancies between the 17 genetic diversity indices (described above: G, R, D, Pareto β, $\bar{r}_d$, Ae, varAe, He, varHe, Ho, varHo, MF$_{IS}$, varF$_{IS}$, PIDsib, PIDu, Sg, SE.Sg) measured among the 53 sampled populations, and their link with reproductive modes including self-compatibility. We reported the amount of variation retained by the first two principal components and the correlation circle on which we plotted predictions of Σclon and Σself as supplementary variables. We also reported the score plot to visualize how LSI and SC populations distribute along the principal components of population genetic diversity. To avoid scaling issues, all descriptors were normalized before analysis.

We analyzed the relationship between genetic diversity indices and their relationships with Σclon and Σself measured in the 53 sampled populations using Kendal partial rank-order correlation tests. We reported the corresponding matrices of correlation.

To detect differences in distribution of population genetic indices, Σclon and Σself, among floral morphs, we computed non-parametric Kruskal-Wallis tests that do not make any assumptions about the type of distribution and about homogeneity of variances between the tested distributions. When needed, we used post-hoc pairwise tests for multiple comparisons of mean rank sums (Nemenyi's test).

399   All statistical tests were computed using PYTHON 3.11, SCIPY.STATS 1.9.3 (Virtanen et al. 2020)

400   and SCIKIT-POSTHOCS (Terpilowski 2019), except the PCA that was performed using R v4.2.2 and

401   the library FACTOMINER (Lê et al. 2008).

Results

Among the 53 populations, we found 40 populations with only L-morph individuals and 13 populations with only S-morph individuals (Table SI2, Figure SI1). The 36 SNP markers we developed for *Lgh* using RAD-Seq and used to genotype the 795 sampled individuals via Hi-Plex allowed confident allele dosage and sufficient statistical power to identify repeated genotypes due to clonal reproduction and to assess the respective proportions of clonality, selfing and allogamy in each population.

Allele dosage and missing genotypes

Within the 795 individuals genotyped at 36 SNPs (resulting in a total of 28,620 single-SNP genotypes), 99.97% (28,612) were genotyped with posterior probability of allele dosage superior to 70% (Table SI3). 785 individuals were thus genotyped with a full set of 36 SNPs with posterior probabilities of allele dosage higher than 70%. Ten individuals distributed in nine populations showed one of their SNP markers with posterior probabilities of allele dosage equal or lower than 70%, that we assigned therefore as missing genotypes.

Statistical power of the developed SNP marker set

Over the 36 polymorphic SNPs, we found an effective number of alleles (A_E) of 1.36 per SNP over all populations (Table SI2). Among populations, mean A_E values over the 36 SNPs were homogenous, ranging from 1.22 to 1.55 (median=1.34). We, however, found large standard deviation of A_E between SNPs within populations, ranging from 1.6 to 2.3 (median=2.1). Some SNPs were apparently fixed in some populations while polymorphic at the scale of the whole dataset. When not fixed, gene diversity H_E in polymorphic SNPs ranged from 0.14 to 0.33

(median=0.2).

These 36 SNPs in the autotetraploid part of *Lgh* would theoretically allow $4^{36}=4.7\times10^{21}$

different possible MLGs considering the four possible nucleobases and $2^{36}=6.9\times10^{10}$ different

possible MLGs assuming two possible nucleobases per locus. Considering allele frequencies in

the sampled populations, the probabilities of identities under panmixia ranged from 8.5×10^{-12}

to 5.2×10^{-5} (median=2.1×10^{-7}) and the unbiased probabilities of identity between sibs $P_{ID\text{-}SIB}$

ranged from 4.2×10^{-6} to 8.3×10^{-3} (median=5.6×10^{-4}; Table SI2). We then considered that the

SNP set we used showed sufficient statistical power to distinguish between true MLGs, and

that individuals with identical MLGs were true repeated genotypes (ramets) of a clonal lineage

(a genet).

Genetic and genotypic diversity

Across populations, we identified a total of 462 distinct MLGs (genets) within the 795 sampled

individuals genotyped. Among them, we found 404 genets (88%) with a single ramet and only

58 genets (12%) with more than two ramets (Table SI4). Forty-eight genets had two to seven

ramets distributed over one to six populations (median=2), seven genets with 10 to 33 ramets

distributed over two to 17 populations (median=9), and one large genet of 99 individuals

distributed over 24 populations (Figure 1).

Within populations, we found from three to 15 different genets (median=12) per population

among the 15 sampled individuals (Table SI4), implying the clear occurrence of repeated

genotypes but also a wide diversity of genets within most populations (Figure 1). Accordingly,

genotypic richness (R) ranged from 0.14 to 1 (median=0.79). The genotypic evenness, Pareto

β, ranged from 0.056 to 3 (median=1.478, Table SI2).

Observed heterozygosity H_O was also high in most populations, slightly above expected

heterozygosity, ranging from 0.13 to 0.38 (median=0.26). The mean standard inbreeding

coefficients (MF_{IS}) averaged over all genotyped loci within populations were negative in all 53

populations and ranged from -0.33 to -0.126 with a very negative median of -0.274 (Table SI2).

Variances of F_{IS} between loci within population were very high, ranging from 1.75 to 36.60

(median=27.57). All these measures argued for reproductive modes implying high rates of

clonality.

Estimates of selfing (S_g) were overall low with a handful of high values, ranging from 0 to 0.61

(median=0.06). Fifteen populations (28%) were estimated with no selfing. Nineteen

populations (36%) showed non-zero estimates under 0.1, 13 populations (25%) between 0.1

and 0.29, and six populations (11%) with estimates between 0.42 and 0.61.

Finally, linkage disequilibria within populations between genotyped SNPs were overall low,

with r_d values ranging from 0 to 0.51 with a median value of 0.12. Forty-two populations (79%)

were under 0.25, and only eight populations (21%) showed linkage disequilibrium between

0.25 and 0.51.

Analyses of covariations between population genetic indices

The first two components of the principal component analysis on the values of genetic

diversity found in the 53 genotyped populations accounted for 78.4% of the total variance

between populations (Figure SI2). Non-parametric Kendall partial rank-order correlations

between genetic indices (Table SI5) and correlations on the first two principal components

from the 17 population genetic indices measured showed three non-collinear groups of

associated genetic indices (Figure SI1.A) that are very similar to the theoretical groups of

population genetic indices expected to covary with different rates of clonality, selfing and

outcrossing in autopolyploids (Stoeckel et al. 2024). A first cluster regrouped G, D*, R, Pareto

β and $VarF_{IS}$, indices that are known to be sensitive to clonality (Stoeckel et al. 2021a) but also

VarHe, VarHo, and VarAe. This cluster largely explains the first dimension (50.2% of the total

variance) of the PCA and was collinear to Σ_{CLON} (also see Figure SI1.B). As expected under

partial clonality, $VarF_{IS}$ was negatively correlated to genotypic diversity indices (R and β; Figure

SI3). The second cluster regrouped P_{ID-SIB}, P_{ID-u}, H_E, A_E and H_O, indices that are linked to the

general genetic diversity of populations (Figure SI1). The third cluster regrouped S_g, SE.Sg, MF_{IS}

and r_D, indices that are usually used to identify, rank and estimate rates of selfing versus

outcrossing in sexual populations (Castric et al. 2002, Bürkly et al. 2017). The second

dimension of the PCA (23.4% of the total variance) was mostly correlated to Ho, Sg, He and r_D

which was collinear to Σ_{SELF} (Figure SI1.C). The clusters of genetic indices were corroborated

by the correlation between Σ_{CLON} and genetic indices (Figure SI2) and Σ_{SELF} and genetic indices

(Figure SI4).

The correlation across populations between Σ_{CLON} and Σ_{SELF} was negative and highly significant

(r_s = -0.66; p < 0.001; Figure 2). This correlation within LSI populations was negative and highly

significant (r_s = -0.65; p < 0.001) while it was nonsignificant in SC populations (r_s = -0.32; p

=0.289).

Differences in genetic diversity and structure between L- and S-morph populations

The number of ramets per genet was similar in LSI and SC populations (H=1.85, p=0.173,

Figure 1 & SI5), as well as were their genotypic richness (R, H=1.23, p=0.267) and evenness

(Pareto's β, H=1.60, p=0.206; Table SI2). The distributions of other population genetic indices

related to clonality were also not significantly different between LSI and SC populations

(VarF$_{IS}$: H=2.94, p=0.086; P$_{ID-SIB}$: H=1.98, p=0.160 and Σ$_{CLON}$ H=2.28, p=0.131; Figure 3).

Mean observed heterozygosity (Ho, H=5.27, p=0.022) and their variances (VarHo, H=13.85,

p<0.001), mean effective number of alleles per locus (Ae, H=5.13, p=0.024) and its variance

over locus (VarAe, H=9.49, p=0.002), MF$_{IS}$ (H=11.85, p<0.001), linkage disequilibrium (r$_D$,

H=5.55, p=0.018), estimate of selfing rates (Sg, H=12.39, p<0.001) and its standard error over

loci (SESg, H=9.58, p=0.002) significantly differed between LSI and SC populations. SC

populations, compared to LSI, showed lower mean observed heterozygosity (medians, LSI:

0.262, SC: 0.210) and less variance between loci (medians, LSI: 2.400, SC: 1.859), lower

effective number of alleles (medians, LSI: 1.341, SC: 1.283) and less variance between loci

(medians, LSI: 4.419, SC: 3.580), higher linkage disequilibrium (medians, LSI: 0.113, SC: 0.152),

higher estimates of selfing rate (medians, LSI: 0.045, SC: 0.175) even if with higher standard

error between loci (medians, LSI: 0.058, SC: 0.156) and higher Σ$_{SELF}$ (medians, LSI: 0.222, SC:

0.495).

Genetic differentiation (ρ$_{ST}$) among pairs of SC populations (median value of ρ$_{ST}$=0.27) was

comparable to the genetic differentiation found among pairs of LSI populations (median value

of ρ$_{ST}$=0.26; H=319.37, post-hoc p=0.322; Figure 4). The median of ρ$_{ST}$ between pairs of LSI and

SC populations reached 0.65. The distribution of inter-morph ρ$_{ST}$ values differed both from the

distribution of pairs of intra-LSI ρ$_{ST}$ values (post-hoc p<0.001) and from the distribution of pairs

of intra-SC ρ$_{ST}$ values (post-hoc p<0.001). The minimum spanning tree of LSI and SC individuals

computed with *GenAPoPop* showed quite clustered distributions of LSI and SC individuals but

also with clear evidence of admixtures between LSI and SC lineages (Figure 5).

Discussion

We used population genetics on autotetraploid SNPs with allele dosage to assess the rate of clonality, of selfing and of outcrossing (i.e., reproductive mode) of 53 populations of *Ludwigia hexapetala* subsp. *grandiflora* (*Lgh*) recently colonizing western European watersheds. To achieve such goal, we developed reproducible codominant molecular markers (SNPs) enabling allele dosage to genotype sampled individuals and used tailored computational analyses that answer the remaining challenges to perform confident population genetic analyses in autopolyploids (Dufresne et al. 2014), This framework can be applied to any autopolyploid species.

Lgh develops two floral morphs respectively associated with an active Late-acting self-incompatible system (L-morph) and a self-compatible system (S-morph, Portillo-Lemus et al. 2021, 2022). Interestingly, the sampled populations in western Europe showed either L or S-morph resulting in two groups of sampled populations: one group of 40 LSI (L-morph) populations and one group of 13 SC (S-morph) populations. They live in close ecological conditions (Portillo-Lemus et al. 2021) which thereby provides a rare opportunity to characterize the genetic consequences of an LSI system compared to a group of SC populations in the same species and ecological context.

Nearly three quarters of the genetic variance explained by different rates of clonality and selfing among *Lgh* populations

The correlation of variations of population genetics indices on the 53 genotyped *Lgh* populations were congruent with theoretical predictions on the variations of genetic indices with different rates of clonality and selfing for autopolyploid populations (Stoeckel et al.

2024). Similarly, such correlations were already predicted from Wright-Fisher-like models

adapted for diploid (Stoeckel et al. 2021a) and haplodiplontic life-cycles (Stoeckel et al.

2021b), and validated respectively using multiple field populations of marine phanerogams

for diploids (Arnaud-Haond et al. 2020) and of brown, green and red algae and mosses for

haplodiplontics (Krueger-Hadfield et al. 2021). These predictions and observations for

diploids, haplodiplontics and now autopolyploids argue again for the primary importance of

reproductive mode, especially of high clonality, to drive variation of genetic diversity among

populations (Duminil et al. 2007). They support the methodological importance to preliminary

estimate confident rates of clonality and of selfing before starting to interpret genetic diversity

with any other factor.

Underestimated clonal diversity in the western European populations of *Lgh*

In our study, we chose to develop and use SNPs rather than AFLP markers which variation in

the precision of sizing of fragments leads to suboptimal reproducibility, particularly across

different platforms (Fry et al. 2009). These 36 SNPs are easily reproducible, accurate enough

to distinguish between offspring of sibling mating as evidence by the probabilities of identity

we obtained, and less expensive in time and money. With these SNPs, we found typical genetic

signatures of high clonality, including clear occurrence of replicated genotypes and mean

negative F_{IS} values with high interlocus variances in all the 53 western European populations.

Previous field and lab observations of peripatric *Lgh* populations reported massive production

of dispersal propagules, rapid expansions of patches and an important capacity for

spontaneous cutting-planting (Dandelot et al. 2005, Thouvenot et al. 2013, Grewell et al. 2016,

Skaer-Thomason et al. 2018a, b). Okada et al. (2009) genotyped around 800 individuals

sampled in 27 *Lgh* populations in Californian wetlands using a set of eight AFLP markers and

reported an extremely reduced clonal diversity: 95% of the samples had the same genotype

and 18 populations over 27 (67%) supported a single AFLP-genotype. All over UK, only two

haplotypes on 14 sampled stems were found using chloroplasts sequences (Armitage et al.

2013). These haplotypes were even shared with some *Lgh* samples invading California. All

these studies concluded that invasive populations of *Lgh* in Europe and in USA reproduce by

exclusive clonality, with a very narrow base of ancestral clones or being monoclonal (Dandelot

et al. 2005, Okada et al. 2009, Thouvenot et al. 2013, Grewell et al. 2016).

However, we found 462 distinct MLGs (genets) out of 795 individuals within these populations

and the large majority of these MLGs (404, thus 88%) were only sampled once in the 53

populations. Our results report a previously-underestimated genotype diversity in the invasive

populations in western Europe, yet congruent with Dandelot (2004) unpublished measures

obtained in three Mediterranean populations using inter-simple sequence repeats (ISSRs).

Dandelot (2004) reported three and seven genotypes over 11 sampling units within two *Lgh*

populations and two genotypes over nine sampling units in a third population, all sampled in

the south-east of France. These results could reveal contrasted situations between *Lgh*

populations introduced in the USA and continental Europe.

Among the 58 MLGs with replicates, seven MLGs were found in a single population and 51

MLGs were found in multiple sites. Clonal propagules of *Lgh* are known to disperse carried by

river current and by birds (zoochory: seedlings have the ability to stick to feathers; Grewel et

al. 2016). Even rare, MLGs found in multiple sites are likely to blur the genetic structure among

Lgh populations if ignored as clonal dispersal. Preliminary detection of these MLGs distributed

among geographical locations will help to understand and interpret the invasive dynamics in

a subsequent study.

Between 10 to 40% of sexual reproduction in *Lgh* populations in western Europe

The large genotype diversity observed in western European populations may either result from an initial or continuous introduction of a large clonal diversity that could have propagated clonally in European watershed or be due to local sexual events mixing introduced lineages into new local genotypes. Beyond the qualitative indication of clonality, we thus aimed at estimating the quantitative rates of clonality versus sexuality using genetic indices in this autotetraploid dataset.

Thirty-two populations showed Pareto β values of their distributions of clonal sizes under 2, which are only found in theoretical populations with rates of clonality higher than 0.8 in diploids, haplodiplontics and autopolyploids (Stoeckel et al. 2021a,b, 2024). All 53 populations showed mean negative F_{IS} with high interlocus variance as expected in highly clonal populations (Balloux et al. 2003, Stoeckel & Masson 2014, Reichel et al. 2016). These values can be observed in theoretical populations with rates of clonality higher than 0.6 (Stoeckel et al. 2021a, b, 2024). All 53 populations also showed small to medium values of linkage disequilibrium between SNPs as expected in highly clonal populations with large population sizes (Navascues et al. 2010, Stoeckel et al. 2021a). Globally, gene diversities measured in these invasive *Lgh* populations were in the higher range of values commonly found for SNPs (Fischer et al. 2017, Schmidt et al. 2021). Such levels of gene diversity are in line with strong rates of clonality that buffer the loss of alleles due to genetic drift (Reichel et al. 2016, Becheler et al. 2020, Stoeckel et al. 2021b). All these values of population genetic indices were consistent with the interpretation that *Lgh* populations in western Europe must reproduce with effective rates of clonality between 60% and 90% (De Meeus et al. 2006, Arnaud-Haond et al. 2020, Stoeckel et al. 2021a, b), even if rates of clonality were variable within this range among populations. Clonal reproduction by rhizomes at the local scale and, in a lesser

proportion, by the dispersal of clonal propagules at larger (*i.e.,* inter-locations) scales would

thus be the main source of population growth and invasive spread of *Lgh* out of its native

range.

Despite being predominant, clonality is, however, not the exclusive reproductive mode. Our

population genetic measures didn't match with exclusive clonality, even being recent.

Despites estimation of rate of sexuality using temporal genotyping would be more resolutive

(Becheler et al. 2017), the values of population genetic indices rather fit between 10% to 40%

of sexuality (DeMeeus et al 2006, Stoeckel et al. 2021a,b, Stoeckel et al. 2024). We also found

local evidence of recombination between clonal lineages, and even rare crosses between LSI

and SC lineages as advocated by the genetic distances between individuals (Figure 5). The

diversity of genotypes detected in western Europe would thus rather match rare but

significant sexual events than the introduction of a large clonal diversity that would have

maintain and propagate by exclusive clonality. Our results newly advocate that sexual

reproduction should not, therefore, be overlooked in these invasive populations, especially in

management plans.

Genetic consequences of LSI compared to self-compatible populations

The late-acting SI system is still one of the less studied breeding systems among the

mechanisms favouring outcrossing in plants (Gibbs 2014). Its efficiency to favour outcrossing

and its consequences on genetic diversity within populations, especially considering its low

but common failures, were not yet deciphered and not yet compared to SC populations in the

same ecological conditions, as already previously explored for gametophytic and sporophyte

SI systems (Busch 2005, Koelling et al. 2011).

The maintenance of SI systems is one of the most intriguing evolutionary puzzles (Porcher &

Lande 2005). Indeed, SI systems are fated to breakdown because SC individuals present the

advantage of reproductive assurance when compatible partners are limited, especially in

peripatric conditions (Eckert et al. 2006). This advantage is even absolute when no compatible

partners are available within pollination range, as we commonly found in *Lgh* populations in

western Europe (see Figure SI1). Conversely, outcrossing imposed by SI systems decreases the

probability of expressing deleterious mutations in descendants as compared to selfing with

the same genetic background (Rice 2004, Navascues et al. 2010).

We found that the 40 *Lgh* populations with L-morph individuals (LSI) showed relatively more

effective alleles, higher gene diversity, higher observed heterozygosity and less linkage

disequilibrium than found in the 13 populations with S-morph individuals (SC). However, these

genetic differences may either be due to the consequences of outcrossing in LSI populations,

or due to different rates of clonality, with potentially higher rates of clonality in LSI populations

due to the local lack of compatible partners and reduced female fertility due to self-pollen

disabling the available ovules, or due to the consequences of autogamy in SC populations. We

develop and disentangle between these three hypotheses thereafter.

Origin of the higher genetic diversity in LSI populations as compared to SC

A first hypothesis to explain the genetic differences found between LSI and SC populations

may be due to the outcrossing consequences on LSI populations. Indeed, similar genetic

differences between SC and SI were recurrently reported in populations mating with

sporophytic and gametophytic SI systems (Igic et al. 2008, Mable 2008). For example,

Leavenworthia alabamica and *L. crassa* are two sister-species mating with a sporophytic SI

within a small genus of eight annual species in which self-fertilization has evolved at least four

times (Beck et al. 2006). In these species, comparisons between SI and SC populations also

showed higher mean genetic diversity in SI populations as compared to the SC, in an overall

background of high genetic diversity (Busch 2005, Koelling et al. 2011). High Ho, negative Fis

and high effective number of alleles per marker are also found in species mating with

gametophytic SI systems, such as in the *Prunus* genius (Stoeckel et al. 2006, Stoeckel et al.

2008, Abdallah et al. 2019). However, theoretical predictions argue that this kind of higher

genetic diversity favored and maintained in SI populations as compared to panmictic and SC

populations must only be found on polymorphic markers highly physically linked to the

genomic regions involved in the SI mechanisms that evolve under balancing selection, and

also, at a lesser magnitude, on physically unlinked markers but only in small population, with

less than 100 individuals including clones (Glémin et al. 2001, Navascues et al. 2010). The 36

SNPs we developed must randomly distribute over the 16 chromosomes inherited twice from

a common ancestor with *Lpm*, i.e., 32 chromosomes out of 80 total chromosomes of *Lgh*

(Barloy et al. 2024), with no reason that all aggregate around the genes involved in the LSI self-

recognition. In addition, *Lpm* is an SC-only species (Estes& Thorp 1974, Grewel et al. 2016). By

parsimony, there is less chance that genes involved in LSI reaction would be stored on these

chromosomes than on some of the 48 other chromosomes; And therefore, very unlikely that

our SNPs would be physically linked to genes under balanced selection due to LSI. *Lgh*

populations in western Europe include far more than one thousand stems per local population

(Portillo-Lemus et al. 2021). It is thus unlikely that differences in genetic diversity observed

between LSI and SC populations would directly result from the consequences of the LSI system

hitchhiked to the whole genome (Navascues et al. 2010).

The second hypothesis is that high genetic diversity in LSI populations may arise and be

maintained due to higher rates of clonality (Stoeckel & Masson 2014, Reichel et al. 2016,

Stoeckel et al. 2021a) as compared to SC populations. In LSI, self-pollen tubes physiologically

blocked at the micropyle may prevent outcrossed-pollen tubes to reach the available ovules,

decreasing female fertility and increasing the importance of clonal regeneration in

populations (Vaughton et al. 2010). In addition, clonality is another uniparental reproductive

mode that allows the local maintenance of populations in the absence of sexually compatible

partners, which is more probable in peripatric conditions (Pannel et al. 2015). Dominant clonal

reproduction and propagation are commonly found in peripatric populations extending their

living range into new areas (Pyšek 1998, Wang et al. 2017). Furthermore, there is an

established intriguing co-occurrence of clonality in SI populations of flowering plants (Vallejo-

Marin et al. 2010). We know that dominant clonal reproduction theoretically favours the

accumulation of deleterious mutations and that these deleterious mutations contribute to the

advantage of outcrossing, which decreases the probability of inbreeding depression in

offsprings, as compared to SC reproducers (Navascues et al. 2010). The observed genetic

differences between LSI and SC populations may thus result from higher clonality in LSI

populations lacking of compatible partners. But we found here no difference in estimates of

clonality between LSI and SC populations, in Σ_{CLON} and in all indices sensible to clonality (R,

Pareto Beta and VarFis; Figure 3).

We rather found significant difference in estimates of selfing between LSI and SC populations,

in Sg (Hardy 2016), in Σ_{SELF} and in all indices sensible to selfing (Ae, linkage disequilibrium, Ho,

mean Fis) revealing typical signatures of higher selfing rates in SC populations. Using Hardy's

method (2016) extended from David et al. (2007) for autopolyploids, SC populations were

estimated to reproduce using a mean selfing rate of 0.18 while LSI populations with only a

mean selfing rate of 0.05, both estimated with similar rates of sexuality. Congruent with higher

estimates of selfing rates and Σ_{SELF} in SC populations, we found higher linkage disequilibrium,

lower observed heterozygosity and less effective number of alleles in SC populations (Figure

3). The observed differences in genetic diversity between LSI and SC populations must thus be

rather caused by the genetic consequences of higher selfing in SC populations, decreasing

their genetic diversity, than due to a direct outcrossing effect of the LSI hitchhiked over all the

Lgh genome within these large populations of *Lgh* (see Navascues et al. 2010). Interestingly,

the outcrossing breeding system allowed by LSI, with a small rate of selfing allowed especially

at the end of the flowering season, was measurable at the population level in genotyping (5%

of selfing in LSI versus 18% in SC) and thus may be efficient at an eco-evolutionary scale.

The recent arriving of *Lgh* in western Europe may also have contributed to shape these genetic

differences. Recent founding events in SC populations are indeed expected to decrease the

overall genetic diversity all along the genomes, including fewer effective alleles, lower gene

diversity, and potentially more linkage disequilibrium as compared to outcrossing populations

(Flanagan et al. 2021). Despite the ρ_{ST} values we observed between pairs of SI and SC

populations being approximately double those measured within pairs of SC populations and

within pairs of LSI populations, we also found clues of admixture in the genetic distances

between LSI and SC individuals in western Europe (Figure 5). This result would thus tend to

minimize the contribution of the foundation in the differences in genetic diversity we

observed between LSI and SC populations, and stress again the importance of selfing to shape

the genetic differences we found between SC and LSI populations.

Advantages of partially clonal LSI populations over SC populations in *Lgh*

Interestingly, we found no correlation between estimates of rates of clonality and of selfing in

SC populations. Selfing rates rather decreased when outcrossing increased, while rates of

clonality remained similar in all populations.

In SC populations of *Lgh* (S-morph), pollen tubes of the L-morph (LSI type) grow significantly faster in SC individuals (Portillo-Lemus et al. 2022). It seems a sufficient and efficient mechanism to favour outcrossing when compatible pollens are available in western European *Lgh* SC populations. The micro-physiological mechanism(s) impacting self-pollen tube growing rather than blocking them in simultaneous (as opposed to dichogamy) monoecious or hermaphrodite species seems to be currently overlooked in literature (e.g. Glover 2007, Nasrallah 2017). Our result obtained on SC *Lgh* calls for a better appraisal of such mechanism in species in which outcrossed offspring seems to be favoured without the occurrence of a clear macro SI system.

Considering we found no differences in the estimated rates of clonality between LSI and SC populations, our results thus argue for an overall stable background rate of clonal propagation, common to the whole species in western Europe, and for an adjustment of the mating system to the local availability of the compatible partners, shifting from selfing allowing reproductive insurance, to outcrossing when possible. In such system, selfing advantage, if any, would result from a transient reproductive insurance, decreasing the cost of developing sexual organs when no outcrossed partners would be available at pollination range, rather than as an efficient uniparental reproductive mode to expand into new areas with no compatible partners, as compared to clonality. Heterozygosity rather than recombination in *Lgh* would thus present some ecological and/or evolutive advantage. This reproductive mode, dominant clonality with some allogamy, is commonly observed in multiple plants (Vallejo-Marin et al. 2010). In such system, we should find higher vigour and fertility of the outcrossed and most heterozygous individuals (heterosis) as compared to the selfed and more homozygous individuals (Darwin 1876, Lippman & Zamir 2007, Birchleret al. 2010).

Reduced allele diversity, heterozygosity and increased linkage disequilibrium between

markers in SC populations due to the higher selfing rates as compared to LSI populations may

explain why only 24% of the western European populations were SC against 76% of LSI.

(Portillo et al. 2021). Based on an analysis of photo collected on the web, Portillo et al (2021)

reported that LSI populations of *Lgh* were more frequent in peripatric populations than in

natives in which SC populations seem in majority. Some successful invasive populations

develop with low genetic diversity (He et al. 2024), questioning on the biological and

environmental factors that may explain their successes in such peripatric conditions (called

the genetic paradox of invasions, Allendorf & Lundquist 2003). This seems to not be the case

of *Lgh* in western Europe: reproductive modes avoiding the effects of inbreeding depression

in *Lgh* populations may on the opposite facilitate their peripatric spreading, as commonly

observed in multiple other invasive plants (Roman & Darling 2007, Forsman 2014), mostly

when developing in harsh and stressful conditions (Fox & Reed, 2011) or when the costs of

inbreeding depression expressed by selfing are superior to the benefits of reproductive

insurance (Layman et al. 2017). Peripatric populations of *Lgh* in western Europe by mixing

clonality with preferential allogamy would thereby solve the problem of maintaining

populations locally without the presence of compatible or less related sexual partner while

reducing the probability to express inbreeding depression (uniparental clonal reproduction,

Baker's conjecture: Barrett et al. 2008, Pannel et al. 2015), and would maximize the probability

of developing offspring with hybrid vigour thanks to LSI and faster growth of crossed-pollen

tubes. In such peripatric conditions, selfing in SC populations would only have limited or

transient advantage to provide behedging reproductive insurance. To disentangle between

the genetic advantages of outcrossing versus reproductive insurance and better state if *Lgh*

populations belong to the genetic paradox of invasions genetic, our results call for measuring

the true proportion of LSI and SC in invasive versus native worldwide *Lgh* populations with

estimation of the rates of clonality and of selfing.

Unusual selfing syndrome in *Lgh* populations

Surprisingly, in *Lgh*, SC populations reproducing with higher selfing rates develop S-morph

flowers that are significantly larger than L-morph flowers that are allogamous controlled by

an LSI system (Portillo-Lemus et al. 2021). Yet, highly selfed populations of plant species tend

to share similar morphology and function resulting in a set of traits called the 'selfing

syndrome', including reduced flower size (Darwin 1876, Tsuchimatsu & Fujii 2022). In *Lgh*,

such high selfing rates may only have occurred recently and endemically in these isolated and

founding European invasive populations, explaining such anomaly trait. Indeed, pioneering

Lgh populations in western Europe only appeared during the last century (Dandelot 2004). But

selfing syndrome, including reduced flower size, seems to evolve rapidly, as observed for

example in five generations in *Mimulus guttatus* (Bodbyl-Roels & Kelly 2011), in four

generations in *Silene latifolia* (Delph et al. 2004), in three generations in *Phlox drummondii*

(Lendvai & Levin 2003) and after only two generations in *Eichhornia paniculata* (Worley &

Barrett 2000). However, in *Lgh*, SC populations with higher selfing rates also reproduce using

intermediate to high rates of clonality, which may delay or even impact the emergence of the

first steps of a selfing syndrome. Our results in *Lgh* thus call for further studies on the

emergence of selfing syndrome in partially clonal and selfed populations.

Conclusion

Using reproducible molecular markers enabling allele dosage to genotype sampled individuals

and tailored computational analyses that answer the remaining challenges to perform

population genetic analyses in autopolyploids (Dufresne et al. 2014), we found that peripatric

populations of *Lgh* in western Europe reproduced using dominant clonality with subtle but

significant sexuality, in nearly all populations, within a large clonal diversity.

Lgh is one of the most invasive aquatic plants in the world, and considerable effort are made

to limit its deleterious effects on the newly colonized ecosystems and on their native species

(Thouvenot et al. 2013, Grewell et al. 2016, Portillo-Lemus et al. 2021). The rare sexual events,

allogamous when possible, occurring in invasive peripatric populations of *Lgh* may favour the

emergence of new genotypes, more adapted to local conditions. Managers should thus chiefly

concentrate their actions on the most sexual populations. We also found variations of the

rates of clonality and of selfing between *Lgh* populations. Considering the importance of

reproductive modes on the dynamics and evolution of populations (Duminil et al. 2007,

Ellegren & Galtier 2016, Glémin et al. 2019), our results advocate that management actions to

control *Lgh* should be built considering their effective reproductive modes, population by

population.

Finally, knowing the reproductive modes of populations, the distribution of clones and the

self-compatibility across western European populations will help to decipher and interpret

their current population structure in a next future, with the aim to identify the origin and the

routes of this recent invasion and to predict its possible short-term dynamics.

Data Accessibility Statement

The data that support the findings of this study are openly available on Zenodo (https://doi.org/10.5281/zenodo.10246264).

Benefits Generated

Benefits from this research accrue from the sharing of our data and results on public databases as described above.

Acknowledgements

We thank Michel Bozec (UMR DECOD), Diane Corbin (FRAPNA Loire-Ecopôle du Forez), Guillaume Le Roux (Reserve Naturel Val d'Alier Châtel-de-Neuvre), Fabrice Dejoux (Service Agriculture Environnement Roannais Agglomération and Jordi Bou Manobens (University of Girona, Spain) for helping to collect *Lgh* samples.

We warmly thank Ingrid Leveque, Marie-Therese Delaroche, Gervaise Fevrier and Patricia Nadan for their administrative support and facilitation.

Funding information

This work was supported by FEDER Région Centre-Val de Loire, by Agence de l'eau Loire-Bretagne (grant Nature 2045, programme 9025, AP 2015 9025) and by the French National Research Agency (project Clonix2D ANR-18-CE32-0001). FEDER funded the doctoral grant of L. Portillo-Lemus and salary of Marilyne Harang. Postdoctoral grant of Ronan Becheler was funded by Clonix2D ANR-18-CE32-0001.

Conflict of interest

The authors declare that they have no financial conflicts of interest based on the content of

this article.

Author contributions

DB and SS laid the foundation of this work, conceived the study and were responsible for

funding applications. LPL, MH and DB collected samples and performed field observations. EP

and DB conceived and supervised RAD4SNP and Hi-Plex methodology, BJP and DJP contributed

to define the SNP set. LPL, MH and ALB performed RAD4SNP approach to develop the SNPs.

LPL and MH extracted DNA and genotyped the samples, SMC and RCV sequenced the data. GL

performed the bioinformatic analyses and SS computed the Bayesian allele dosage. DB, LPL

and SS performed early data explorations at the end of LPL PhD thesis that was supervised by

DB and SS. RB and SS performed computations and analyses, tracked and managed the

bibliography, produced tables, figures and interpretation. DB, RB and SS wrote the

manuscript. All authors read and approved the final manuscript.

References

Abdallah, D., Baraket, G., Perez, V., Ben Mustapha, S., Salhi-Hannachi, A., & Hormaza, J. I.

(2019). Analysis of self-incompatibility and genetic diversity in diploid and hexaploid

plum genotypes. *Frontiers in Plant Science*, *10*.

https://www.frontiersin.org/articles/10.3389/fpls.2019.00896

Agapow, P.-M., & Burt, A. (2001). Indices of multilocus linkage disequilibrium. *Molecular*

Ecology Notes, *1*(1–2), 101–102. https://doi.org/10.1046/j.1471-8278.2000.00014.x

Allendorf, F. W., & Lundquist, L. L. (2003). Introduction: Population Biology, Evolution, and

Control of Invasive Species. *Conservation Biology*, 17(1), 24–30.

https://doi.org/10.1046/j.1523-1739.2003.02365.x

Armitage, J. D., Könyves, K., Bailey, J. P., David, J. C., & Culham, A. (2013). A molecular,

morphological and cytological investigation of the identity of non-native *Ludwigia*

(Onagraceae) populations in Britain. *New Journal of Botany*, *3*(2), 88–95.

https://doi.org/10.1179/2042349713Y.0000000023

Arnaud-Haond, S., Alberto, F., Teixeira, S., Procaccini, G., Serrão, E. A., & Duarte, C. M.

(2005). Assessing genetic diversity in clonal organisms: low diversity or low resolution?

combining power and cost efficiency in selecting markers. *Journal of Heredity*, *96*(4),

434–440. https://doi.org/10.1093/jhered/esi043

Arnaud-Haond, S., Duarte, C. M., Alberto, F., & Serrão, E. A. (2007). Standardizing methods

to address clonality in population studies. *Molecular Ecology*, *16*(24), 5115–5139.

https://doi.org/10.1111/j.1365-294X.2007.03535.x

Arnaud-Haond, S., Stoeckel, S., & Bailleul, D. (2020). New insights into the population

genetics of partially clonal organisms: When seagrass data meet theoretical

expectations. *Molecular Ecology*, *29*(17), 3248–3260.

https://doi.org/10.1111/mec.15532

Baduel, P., Bray, S., Vallejo-Marin, M., Kolář, F., & Yant, L. (2018). The "polyploid hop":
shifting challenges and opportunities over the evolutionary lifespan of genome
duplications. *Frontiers in Ecology and Evolution*, *6*.
https://www.frontiersin.org/articles/10.3389/fevo.2018.00117

Baird, N. A., Etter, P. D., Atwood, T. S., Currey, M. C., Shiver, A. L., Lewis, Z. A., Selker, E.
U., Cresko, W. A., & Johnson, E. A. (2008). Rapid SNP Discovery and Genetic
Mapping Using Sequenced RAD Markers. *PLOS ONE*, 3(10), e3376.
https://doi.org/10.1371/journal.pone.0003376

Balloux, F., Lehmann, L., & de Meeûs, T. (2003). The population genetics of clonal and
partially clonal diploids. *Genetics*, *164*(4), 1635–1644.
https://doi.org/10.1093/genetics/164.4.1635

Barloy, D., Lemus, L. P.-, Krueger-Hadfield, S. A., Huteau, V., & Coriton, O. (2024). Genomic
relationships among diploid and polyploid species of the genus Ludwigia L. section
Jussiaea using a combination of cytogenetic, morphological, and crossing
investigations. *Peer Community in Evolutionary Biology*, 4, e8.
https://doi.org/10.24072/pci.evolbiol.100645.

Barrett, S. C. H., Colautti, R. I., & Eckert, C. G. (2008). Plant reproductive systems and
evolution during biological invasion. *Molecular Ecology*, 17(1), 373–383.
https://doi.org/10.1111/j.1365-294X.2007.03503.x

Barrett, S. C. H. (2010). Understanding plant reproductive diversity. *Philosophical
Transactions of the Royal Society B: Biological Sciences*, *365*(1537), 99–109.
https://doi.org/10.1098/rstb.2009.0199

Becheler, R., Guillemin, M.-L., Stoeckel, S., Mauger, S., Saunier, A., Brante, A., Destombe,
C., & Valero, M. (2020). After a catastrophe, a little bit of sex is better than nothing:
Genetic consequences of a major earthquake on asexual and sexual populations.

Evolutionary Applications, *13*(8), 2086–2100. https://doi.org/10.1111/eva.12967

Becheler, R., Masson, J.-P., Arnaud-Haond, S., Halkett, F., Mariette, S., Guillemin, M.-L.,

Valero, M., Destombe, C., & Stoeckel, S. (2017). ClonEstiMate, a Bayesian method for

quantifying rates of clonality of populations genotyped at two-time steps. *Molecular*

Ecology Resources, 17(6), e251–e267. https://doi.org/10.1111/1755-0998.12698

Beck, J. B., Al-Shehbaz, I. A., & Schaal, B. A. (2006). *Leavenworthia* (Brassicaceae) revisited:

testing classic systematic and mating system hypotheses. *Systematic Botany*, *31*(1),

151–159. https://doi.org/10.1600/036364406775971732

Besnard, A.-L., Park, D. J., Pope, B. J., Hammet, F., Michon-Coudouel, S., Biget, M., Krueger-

Hadfield, S. A., Mauger, S., & Petit, E. J. (2023). Workflow for SNP genotyping using

the   Hi-Plex   method.   *Protocols.io*,

https://doi.org/10.17504/protocols.io.8epv5jnnnl1b/v2

Birchler, J. A., Yao, H., Chudalayandi, S., Vaiman, D., & Veitia, R. A. (2010). Heterosis. *The*

Plant Cell, *22*(7), 2105–2112. https://doi.org/10.1105/tpc.110.076133

Bodbyl Roels, S. A., & Kelly, J. K. (2011). Rapid evolution caused by pollinator loss in

Mimulus guttatus. *Evolution*, *65*(9), 2541–2552. https://doi.org/10.1111/j.1558-

5646.2011.01326.x

Booy, G., Hendriks, R.J.J., Smulders, M.J.M., Van Groenendael, J.M. and Vosman, B. (2000).

Genetic Diversity and the Survival of Populations. *Plant Biology*, *2*, 379-395.

https://doi.org/10.1055/s-2000-5958

Brennan, A. C., Harris, S. A., Tabah, D. A., & Hiscock, S. J. (2002). The population genetics

of sporophytic self-incompatibility in *Senecio squalidus* L. (Asteraceae) I: S allele

diversity   in   a   natural   population.   *Heredity*,   *89*(6),   Article 6.

https://doi.org/10.1038/sj.hdy.6800159

Bürkli, A., Sieber, N., Seppälä, K., & Jokela, J. (2017). Comparing direct and indirect selfing

rate estimates: When are population-structure estimates reliable? *Heredity*, *118*(6), Article 6. https://doi.org/10.1038/hdy.2017.1

Busch, J. W. (2005). Inbreeding depression in self-incompatible and self-compatible populations of *Leavenworthia alabamica*. *Heredity*, *94*(2), Article 2. https://doi.org/10.1038/sj.hdy.6800584

Busch, J. W., & Urban, L. (2011). Insights gained from 50 years of studying the evolution of self-compatibility in *Leavenworthia* (Brassicaceae). *Evolutionary Biology*, *38*(1), 15–27. https://doi.org/10.1007/s11692-010-9104-5

Cabin, R. J., Evans, A. S., Jennings, D. L., Marshall, D. L., Mitchell, R. J., & Sher, A. A. (1996). Using bud pollinations to avoid self-incompatibility: Implications from studies of three mustards. *Canadian Journal of Botany*, 74(2), 285–289. https://doi.org/10.1139/b96-034

Castric, V., Bernatchez, L., Belkhir, K., & Bonhomme, F. (2002). Heterozygote deficiencies in small lacustrine populations of brook charr *Salvelinus fontinalis* Mitchill (Pisces, Salmonidae): A test of alternative hypotheses. *Heredity*, *89*(1), Article 1. https://doi.org/10.1038/sj.hdy.6800089

Castric, V., & Vekemans, X. (2004). Plant self-incompatibility in natural populations: a critical assessment of recent theoretical and empirical advances. *Molecular Ecology*, *13*(10), 2873–2889. https://doi.org/10.1111/j.1365-294X.2004.02267.x

Charlesworth, D. (2003). Effects of inbreeding on the genetic diversity of populations. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *358*(1434), 1071–1084. https://doi.org/10.1098/rstb.2003.1296

Charlesworth, D., Vekemans, X., Castric, V., & Glémin, S. (2005). Plant self-incompatibility systems: A molecular evolutionary perspective. *New Phytologist*, *168*(1), 61–69. https://doi.org/10.1111/j.1469-8137.2005.01443.x

Chen, X., Hao, S., Wang, L., Fang, W., Wang, Y., & Li, X. (2012). Late-acting self-

incompatibility in tea plant (*Camellia sinensis*). *Biologia*, *67*(2), 347–351.

https://doi.org/10.2478/s11756-012-0018-9

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A.,

Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools

and BCFtools. GigaScience, 10, giab008. https://doi.org/10.1093/gigascience/giab008

Dandelot, S. (2004). *Les Ludwigia spp. Invasives du Sud de la France: Historique,*

Biosystématique,   Biologie   et   Ecologie   [PhD   thesis,   Aix-Marseille 3].

https://www.theses.fr/2004AIX30052

Dandelot, S., Verlaque, R., Dutartre, A., & Cazaubon, A. (2005). Ecological, Dynamic and

Taxonomic Problems Due to Ludwigia (Onagraceae) in France. Hydrobiologia, 551(1),

131–136. https://doi.org/10.1007/s10750-005-4455-0

Darwin, C. R. (1876). *The effects of cross and self-fertilization in the vegetable kingdom*. John

Murray (London).

David, P., Pujol, B., Viard, F., Castella, V., & Goudet, J. (2007). Reliable selfing rate estimates

from imperfect population genetic data. *Molecular Ecology*, *16*(12), 2474–2487.

https://doi.org/10.1111/j.1365-294X.2007.03330.x

De Meeûs, T., & Balloux, F. (2004). Clonal reproduction and linkage disequilibrium in

diploids: A simulation study. *Infection, Genetics and Evolution*, *4*(4), 345–351.

https://doi.org/10.1016/j.meegid.2004.05.002

De Meeûs, T., Lehmann, L., & Balloux, F. (2006). Molecular epidemiology of clonal diploids:

A quick overview and a short DIY (do it yourself) notice. *Infection, Genetics and*

Evolution, *6*(2), 163–170. https://doi.org/10.1016/j.meegid.2005.02.004

De Meeûs, T., Prugnolle, F., & Agnew, P. (2007). Asexual reproduction: Genetics and

evolutionary aspects. *Cellular and Molecular Life Sciences*, *64*(11), 1355–1372.

https://doi.org/10.1007/s00018-007-6515-2

De Nettancourt, D. (1997). Incompatibility in angiosperms. *Sexual Plant Reproduction*, *10*(4),

185–199. https://doi.org/10.1007/s004970050087

De Nettancourt, D. (2001). *Incompatibility and Incongruity in Wild and Cultivated Plants*.

Springer. https://doi.org/10.1007/978-3-662-04502-2

Dellaporta, S. L., & Calderon-Urrea, A. (1993). Sex Determination in Flowering Plants. *The*

Plant Cell, *5*(10), 1241–1251. https://doi.org/10.2307/3869777

Delord, C., Lassalle, G., Oger, A., Barloy, D., Coutellec, M.-A., Delcamp, A., Evanno, G.,

Genthon, C., Guichoux, E., Le Bail, P.-Y., Le Quilliec, P., Longin, G., Lorvelec, O.,

Massot, M., Reveillac, E., Rinaldo, R., Roussel, J.-M., Vigouroux, R., Launey, S., &

Petit, E. J. (2018). A cost-and-time effective procedure to develop SNP markers for

multiple species: A support for community genetics. *Methods in Ecology and Evolution*,

9(9), 1959–1974. https://doi.org/10.1111/2041-210X.13034

Delph, L. F., Gehring, J. L., Frey, F. M., Arntz, A. M., & Levri, M. (2004). Genetic constraints

on floral evolution in a sexually dimorphic plant revealed by artificial selection.

Evolution, *58*(9), 1936–1946. https://doi.org/10.1111/j.0014-3820.2004.tb00481.x

Dorken, M. E., & Eckert, C. G. (2001). Severely reduced sexual reproduction in northern

populations of a clonal plant, *Decodonverticillatus* (Lythraceae). *Journal of Ecology*,

89(3), 339–350. https://doi.org/10.1046/j.1365-2745.2001.00558.x

Dufresne, F., Stift, M., Vergilino, R., & Mable, B. K. (2014). Recent progress and challenges

in population genetics of polyploid organisms: An overview of current state-of-the-art

molecular   and   statistical   tools.   *Molecular   Ecology*,   *23*(1),   40–69.

https://doi.org/10.1111/mec.12581

Duminil, J., Fineschi, S., Hampe, A., Jordano, P., Salvini, D., Vendramin, G. G., & Petit, R. J.

(2007). Can population genetic structure be predicted from life-history traits? *The*

American Naturalist, *169*(5), 662–672. https://doi.org/10.1086/513490

Eckert, C., Samis, K. & Dart, S. (2006). Reproductive Assurance and the Evolution of

Uniparental Reproduction in Flowering Plants. In Harder & Barrett (Eds), *Ecology and*

Evolution of Flowers, pp.183–200. Oxford, UK: Oxford University Press.

Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*,

17(7), Article 7. https://doi.org/10.1038/nrg.2016.58

Estes, J. R., & Thorp, R. W. (1974). Pollination in Ludwigia peploides ssp. Glabrescens

(Onagraceae). *Bulletin of the Torrey Botanical Club, 101*(5), 272–276.

https://doi.org/10.2307/2484872

Fehrer, J. (2010). Unraveling the mysteries of reproduction. *Heredity*, *104*(5), Article 5.

https://doi.org/10.1038/hdy.2010.12

Fischer, M. C., Rellstab, C., Leuzinger, M., Roumet, M., Gugerli, F., Shimizu, K. K.,

Holderegger, R., & Widmer, A. (2017). Estimating genomic diversity and population

differentiation – an empirical comparison of microsatellite and SNP variation in

Arabidopsis halleri. *BMC Genomics*, *18*(1), 69. https://doi.org/10.1186/s12864-016-

3459-7

Flanagan, B. A., Krueger-Hadfield, S. A., Murren, C. J., Nice, C. C., Strand, A. E., & Sotka, E.

E. (2021). Founder effects shape linkage disequilibrium and genomic diversity of a

partially clonal invader. *Molecular Ecology*, *30*(9), 1962–1978.

https://doi.org/10.1111/mec.15854

Forsman, A. (2014). Effects of genotypic and phenotypic variation on establishment are

important for conservation, invasion, and infection biology. *Proceedings of the National*

Academy of Sciences, 111(1), 302–307. https://doi.org/10.1073/pnas.1317745111

Fox, C. W., & Reed, D. H. (2011). Inbreeding Depression Increases with Environmental Stress:

An Experimental Study and Meta-Analysis. *Evolution*, 65(1), 246–258.

https://doi.org/10.1111/j.1558-5646.2010.01108.x

Franklin-Tong, V. E. (2008). *Self-Incompatibility in Flowering Plants*. Springer.
https://doi.org/10.1007/978-3-540-68486-2

Fry, N. K., Savelkoul, P. H., & Visca, P. (2009). Amplified Fragment Length Polymorphism
Analysis. In: Caugant, D. (eds) Molecular Epidemiology of Microorganisms. *Methods*
in Molecular Biology, vol 551, 89-104. Humana Press, Totowa, NJ.
https://doi.org/10.1007/978-1-60327-999-4_8

Gibbs, P. E. (2014). Late-acting self-incompatibility – the pariah breeding system in flowering
plants. *New Phytologist*, *203*(3), 717–734. https://doi.org/10.1111/nph.12874

Glémin, S., Bataillon, T., Ronfort, J., Mignot, A., & Olivieri, I. (2001). inbreeding depression
in small populations of self-incompatible plants. *Genetics*, *159*(3), 1217–1229.
https://doi.org/10.1093/genetics/159.3.1217

Glémin, S., François, C. M., & Galtier, N. (2019). Genome Evolution in Outcrossing vs. Selfing
vs. Asexual Species. In M. Anisimova (Ed.), *Evolutionary Genomics: Statistical and*
Computational Methods (pp. 331–369). Springer. https://doi.org/10.1007/978-1-4939-
9074-0_11

Glover, B. J. (2007). Preventing Self-fertilization. In B. Glover (Ed.), Understanding Flowers
and Flowering: An integrated approach (p. 0). Oxford University Press.
https://doi.org/10.1093/acprof:oso/9780198565970.003.0012

Golding, G. B., & Strobeck, C. (1980). Linkage disequilibrium in a finite population that is
partially selfing. *Genetics*, *94*(3), 777–789. https://doi.org/10.1093/genetics/94.3.777

Goodwillie, C., & Weber, J. J. (2018). The best of both worlds? A review of delayed selfing in
flowering plants. *American Journal of Botany*, *105*(4), 641–655.
https://doi.org/10.1002/ajb2.1045

Grewell, B., Netherland, M., & Skaer Thomason, M. (2016). Establishing research and

management priorities for invasive water primroses *(Ludwigia spp.)*. https://doi.org/10.13140/RG.2.1.5020.6482

Haldane, J. B. S. (1930). Theoretical genetics of autopolyploids. *Journal of Genetics*, 22(3), 359–372. https://doi.org/10.1007/BF02984197

Halkett, F., Simon, J.-C., & Balloux, F. (2005). Tackling the population genetics of clonal and partially clonal organisms. *Trends in Ecology & Evolution*, *20*(4), 194–201. https://doi.org/10.1016/j.tree.2005.01.001

Hammet, F., Mahmood, K., Green, T. R., Nguyen-Dumont, T., Southey, M. C., Buchanan, D. D., Lonie, A., Nathanson, K. L., Couch, F. J., Pope, B. J., & Park, D. J. (2019). Hi-Plex2: A simple and robust approach to targeted sequencing-based genetic screening. *BioTechniques*, 67(3), 118–122. https://doi.org/10.2144/btn-2019-0026

Hao, Y.-Q., Zhao, X.-F., She, D.-Y., Xu, B., Zhang, D.-Y., & Liao, W.-J. (2012). The Role of Late-acting self-incompatibility and early-acting inbreeding depression in governing female fertility in monkshood, *Aconitum kusnezoffii*. *PLOS ONE*, *7*(10), e47034. https://doi.org/10.1371/journal.pone.0047034

Harder, L. D., & Barrett, S. C. H. (2006). *Ecology and Evolution of Flowers*. OUP Oxford.

Hardy, O. J. (2016). Population genetics of autopolyploids under a mixed mating model and the estimation of selfing rate. *Molecular Ecology Resources*, *16*(1), 103–117. https://doi.org/10.1111/1755-0998.12431

Hardy, O. J., & Vekemans, X. (2002). SPAGEDI: A versatile computer program to analyze spatial genetic structure at the individual or population levels. *Molecular Ecology Notes*, *2*(4), 618–620. https://doi.org/10.1046/j.1471-8286.2002.00305.x

He, Z.-Z., Stotz, G. C., Liu, X., Liu, J.-J., Wang, Y.-G., Yang, J., Li, L.-F., Zhang, W.-J., Nan, P., & Song, Z.-P. (2024). A global synthesis of the patterns of genetic diversity in endangered and invasive plants. *Biological Conservation*, 291, 110473.

https://doi.org/10.1016/j.biocon.2024.110473

Herben, T., Suda, J., & Klimešová, J. (2017). Polyploid species rely on vegetative reproduction more than diploids: A re-examination of the old hypothesis. *Annals of Botany*, *120*(2), 341–349. https://doi.org/10.1093/aob/mcx009

Hoban, S., Campbell, C. D., da Silva, J. M., Ekblom, R., Funk, W. C., Garner, B. A., Godoy, J. A., Kershaw, F., MacDonald, A. J., Mergeay, J., Minter, M., O'Brien, D., Vinas, I. P., Pearson, S. K., Pérez-Espona, S., Potter, K. M., Russo, I.-R. M., Segelbacher, G., Vernesi, C., & Hunter, M. E. (2021). Genetic diversity is considered important but interpreted narrowly in country reports to the Convention on Biological Diversity: Current actions and indicators are insufficient. *Biological Conservation*, *261*, 109233. https://doi.org/10.1016/j.biocon.2021.109233

Hoch, P. C., Wagner, W. L., & Raven, P. H. (2015). The correct name for a section of Ludwigia L. (Onagraceae). *PhytoKeys*, *50*, 31–34. https://doi.org/10.3897/phytokeys.50.4887

Holsinger, K. E. (2000). Reproductive systems and evolution in vascular plants. *Proceedings of the National Academy of Sciences*, *97*(13), 7037–7042. https://doi.org/10.1073/pnas.97.13.7037

Husband, B. C., Ozimec, B., Martin, S. L., & Pollock, L. (2008). Mating consequences of polyploid evolution in flowering plants: current trends and insights from synthetic polyploids. *International Journal of Plant Sciences*, *169*(1), 195–206. https://doi.org/10.1086/523367

Igic, B., Lande, R., & Kohn, J. R. (2008). Loss of self-incompatibility and its evolutionary consequences. *International Journal of Plant Sciences*, *169*(1), 93–104. https://doi.org/10.1086/523362

Koelling, V. A., Hamrick, J. L., & Mauricio, R. (2011). Genetic diversity and structure in two species of *Leavenworthia* with self-incompatible and self-compatible populations.

Heredity, *106*(2), 310-318. https://doi.org/10.1038/hdy.2010.59

Krueger-Hadfield, S. A., Guillemin, M.-L., Destombe, C., Valero, M., & Stoeckel, S. (2021). Exploring the Genetic Consequences of Clonality in Haplodiplontic Taxa. *Journal of Heredity*, 112(1), 92–107. https://doi.org/10.1093/jhered/esaa063

Lawrence, M. J., Marshall, D. F., Curtis, V. E., & Fearon, C. H. (1985). Gametophytic self-incompatibility re-examined: A reply. *Heredity*, 54(1), 131–138. https://doi.org/10.1038/hdy.1985.17

Layman, N. C., Fernando, M. T. R., Herlihy, C. R., & Busch, J. W. (2017). Costs of selfing prevent the spread of a self-compatibility mutation that causes reproductive assurance. *Evolution*, 71(4), 884–897. https://doi.org/10.1111/evo.13167

Lê, S., Josse, J. & Husson, F. (2008). FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software, 25*(1), 1-18. https://doi.org/10.18637/jss.v025.i01

Lendvai, G., & Levin, D. A. (2003). Rapid response to artificial selection on flower size in *Phlox*. *Heredity*, *90*(4), Article 4. https://doi.org/10.1038/sj.hdy.6800249

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics*, *26*, 589-595. https://doi.org/10.1093/bioinformatics/btp698.

Lippman, Z. B., & Zamir, D. (2007). Heterosis: Revisiting the magic. *Trends in Genetics*, *23*(2), 60–66. https://doi.org/10.1016/j.tig.2006.12.006

Lucek, K., & Willi, Y. (2021). Drivers of linkage disequilibrium across a species' geographic range. *PLOS Genetics*, *17*(3), e1009477. https://doi.org/10.1371/journal.pgen.1009477

Mable, B. K. (2008). Genetic causes and consequences of the breakdown of self-incompatibility: Case studies in the Brassicaceae. *Genetics Research*, *90*(1), 47–60. https://doi.org/10.1017/S0016672307008907

Meek, M. H., & Larson, W. A. (2019). The future is now: Amplicon sequencing and sequence capture usher in the conservation genomics era. Molecular Ecology Resources, 19(4),

795–803. https://doi.org/10.1111/1755-0998.12998

Meirmans, P. G., Liu, S., & van Tienderen, P. H. (2018). The analysis of polyploid genetic data. *Journal of Heredity*, *109*(3), 283–296. https://doi.org/10.1093/jhered/esy006

Nasrallah, J. B. (2017). Plant mating systems: Self-incompatibility and evolutionary transitions to self-fertility in the mustard family. Current Opinion in Genetics & Development, 47, 54–60. https://doi.org/10.1016/j.gde.2017.08.005

Navascués, M., Stoeckel, S., & Mariette, S. (2010). Genetic diversity and fitness in small populations of partially asexual, self-incompatible plants. *Heredity*, *104*(5), Article 5. https://doi.org/10.1038/hdy.2009.159

Nesom, G.L. & Kartesz, J.T. (2000). Observations on the *Ludwigia uruguayensis* complex (Onagraceae) in the United States. Castanea 65(2): 123-125. http://www.jstor.org/stable/4034110

Nordborg, M. (2000). Linkage disequilibrium, gene trees and selfing: an ancestral recombination graph with partial self-fertilization. *Genetics*, *154*(2), 923–929. https://doi.org/10.1093/genetics/154.2.923

Okada, M., Grewell, B. J., & Jasieniuk, M. (2009). Clonal spread of invasive *Ludwigia hexapetala* and *L. grandiflora* in freshwater wetlands of California. *Aquatic Botany*, *91*(3), 123–129. https://doi.org/10.1016/j.aquabot.2009.03.006

Orive, M. E., & Krueger-Hadfield, S. A. (2021). Sex and asex: a clonal lexicon. *Journal of Heredity*, *112*(1), 1–8. https://doi.org/10.1093/jhered/esaa058

Pannell, J. R., Auld, J. R., Brandvain, Y., Burd, M., Busch, J. W., Cheptou, P.-O., Conner, J. K., Goldberg, E. E., Grant, A.-G., Grossenbacher, D. L., Hovick, S. M., Igic, B., Kalisz, S., Petanidou, T., Randle, A. M., de Casas, R. R., Pauw, A., Vamosi, J. C., & Winn, A. A. (2015). The scope of Baker's law. *New Phytologist*, *208*(3), 656–667. https://doi.org/10.1111/nph.13539

Porcher, E., & Lande, R. (2005). The evolution of self-fertilization and inbreeding depression

under pollen discounting and pollen limitation. *Journal of Evolutionary Biology*, *18*(3),

497–508. https://doi.org/10.1111/j.1420-9101.2005.00905.x

Portillo Lemus, L. O., Bozec, M., Harang, M., Coudreuse, J., Haury, J., Stoeckel, S., & Barloy,

D. (2021). Self-incompatibility limits sexual reproduction rather than environmental

conditions in an invasive water primrose. *Plant-Environment Interactions*, *2*(2), 74–86.

https://doi.org/10.1002/pei3.10042

Portillo Lemus, L. O., Harang, M., Bozec, M., Haury, J., Stoeckel, S., & Barloy, D. (2022).

Late-acting self-incompatible system, preferential allogamy and delayed selfing in the

heteromorphic invasive populations of *Ludwigia grandiflora subsp. Hexapetala*. *Peer*

Community Journal, *2*. https://doi.org/10.24072/pcjournal.108

Pyšek, P. (1998). Is there a taxonomic pattern to plant invasions? *Oikos*, *82*(2), 282–294.

https://doi.org/10.2307/3546968

Reichel, K., Masson, J.-P., Malrieu, F., Arnaud-Haond, S., & Stoeckel, S. (2016). Rare sex or

out of reach equilibrium? The dynamics of FISin partially clonal organisms. *BMC*

Genetics, *17*(1), 76. https://doi.org/10.1186/s12863-016-0388-z

Rice, S. H. (2004). *Evolutionary theory: Mathematical and conceptual foundations*. Sinauer

Associates.

Roman, J., & Darling, J. A. (2007). Paradox lost: Genetic diversity and the success of aquatic

invasions. *Trends in Ecology & Evolution*, 22(9), 454–464.

https://doi.org/10.1016/j.tree.2007.07.002

Ronfort, J., Jenczewski, E., Bataillon, T., & Rousset, F. (1998). Analysis of population structure

in autotetraploid species. *Genetics*, *150*(2), 921–930.

https://doi.org/10.1093/genetics/150.2.921

Rouger, R., Reichel, K., Malrieu, F., Masson, J. P., & Stoeckel, S. (2016). Effects of complex

life cycles on genetic diversity: Cyclical parthenogenesis. *Heredity*, *117*(5), Article 5. https://doi.org/10.1038/hdy.2016.52

Roze, D. (2016). Background selection in partially selfing populations. *Genetics*, *203*(2), 937–957. https://doi.org/10.1534/genetics.116.187955

Schmidt, T. L., Jasper, M.-E., Weeks, A. R., & Hoffmann, A. A. (2021). Unbiased population heterozygosity estimates from genome-wide sequence data. *Methods in Ecology and Evolution*, *12*(10), 1888–1898. https://doi.org/10.1111/2041-210X.13659

Seavey, S. R., & Bawa, K. S. (1986). Late-acting self-incompatibility in angiosperms. *The Botanical Review*, *52*(2), 195–219. https://doi.org/10.1007/BF02861001

Skaer Thomason, M. J., Grewell, B. J., & Netherland, M. D. (2018a). Dynamics of *Ludwigia hexapetala* invasion at three spatial scales in a regulated river. *Wetlands*, *38*(6), 1285–1298. https://doi.org/10.1007/s13157-018-1053-2

Skaer Thomason, M. J., McCort, C. D., Netherland, M. D., & Grewell, B. J. (2018b). Temporal and nonlinear dispersal patterns of *Ludwigia hexapetala* in a regulated river. *Wetlands Ecology and Management*, *26*(5), 751–762. https://doi.org/10.1007/s11273-018-9605-z

Steinecke, C., Gorman, C. E., Stift, M., & Dorken, M. E. (2022). Outcrossing rates in an experimentally admixed population of self-compatible and self-incompatible Arabidopsis lyrata. *Heredity*, *128*(1), Article 1. https://doi.org/10.1038/s41437-021-00489-8

Stoeckel, S., Arnaud-Haond, S., & Krueger-Hadfield, S. A. (2021b). The combined effect of haplodiplonty and partial clonality on genotypic and genetic diversity in a finite mutating population. *Journal of Heredity*, *112*(1), 78–91. https://doi.org/10.1093/jhered/esaa062

Stoeckel, S., Barloy, D., Portillo Lemus, L., & Becheler, R. (2023). Genotyping measures and population genetic indices for assesing reproductive modes of polyploid Ludwigia

grandiflora subsp. hexapetala in western Europe [Data set]. Zenodo. https://doi.org/10.5281/zenodo.10246264

Stoeckel, S., Becheler, R., Bocharova, E., & Barloy, D. (2023). GenAPoPop 1.0: A user-friendly software to analyse genetic diversity and structure from partially clonal and selfed autopolyploid organisms. Molecular Ecology Resources, 00, 1–11. https://doi.org/10.1111/1755-0998.13886

Stoeckel, S., Grange, J., Fernández-Manjarres, J. F., Bilger, I., Frascaria-Lacoste, N., & Mariette, S. (2006). Heterozygote excess in a self-incompatible and partially clonal forest tree species—Prunus avium L. *Molecular Ecology*, *15*(8), 2109–2118. https://doi.org/10.1111/j.1365-294X.2006.02926.x

Stoeckel, S., Castric, V., Mariette, S., & Vekemans, X. (2008). Unequal allelic frequencies at the self-incompatibility locus within local populations of Prunus avium L.: An effect of population structure? Journal of Evolutionary Biology, 21(3), 889–899. https://doi.org/10.1111/j.1420-9101.2008.01504.x

Stoeckel, S., & Masson, J.-P. (2014). The exact distributions of fis under partial asexuality in small finite populations with mutation. *PLOS ONE*, *9*(1), e85228. https://doi.org/10.1371/journal.pone.0085228

Stoeckel, S., Porro, B., & Arnaud-Haond, S. (2021a). The discernible and hidden effects of clonality on the genotypic and genetic states of populations: Improving our estimation of clonal rates. *Molecular Ecology Resources*, *21*(4), 1068–1084. https://doi.org/10.1111/1755-0998.13316

Terpilowski, (2019). scikit-posthocs: Pairwise multiple comparison tests in Python. Journal of Open Source Software, 4(36), 1169, https://doi.org/10.21105/joss.01169

Thouvenot, L., Haury, J., & Thiebaut, G. (2013). A success story: Water primroses, aquatic plant pests. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *23*(5), 790–

803. https://doi.org/10.1002/aqc.2387

Tsuchimatsu, T., & Sota, F. (2022). The selfing syndrome and beyond: diverse evolutionary consequences of mating system transitions in plants. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 377, 20200510. https://doi.org/10.1098/rstb.2020.0510.

Vallejo-Marín, M., Dorken, M. E., & Barrett, S. C. H. (2010). The ecological and evolutionary consequences of clonality for plant mating. *Annual Review of Ecology, Evolution, and Systematics*, *41*(1), 193–213. https://doi.org/10.1146/annurev.ecolsys.110308.120258

Van Drunen, W. E., & Husband, B. C. (2019). Evolutionary associations between polyploidy, clonal reproduction, and perenniality in the angiosperms. *New Phytologist*, *224*(3), 1266–1277. https://doi.org/10.1111/nph.15999

Vaughton, G., Ramsey, M., & Johnson, S. D. (2010). Pollination and late-acting self-incompatibility in Cyrtanthus breviflorus (Amaryllidaceae): Implications for seed production. *Annals of Botany*, 106(4), 547–555. https://doi.org/10.1093/aob/mcq149

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., & van Mulbregt, P. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, *17*(3), Article 3. https://doi.org/10.1038/s41592-019-0686-2

Wang, Y.-J., Müller-Schärer, H., van Kleunen, M., Cai, A.-M., Zhang, P., Yan, R., Dong, B.-C., & Yu, F.-H. (2017). Invasive alien plants benefit more from clonal integration in heterogeneous environments than natives. *New Phytologist*, *216*(4), 1072–1078. https://doi.org/10.1111/nph.14820

Worley, A. C., & Barrett, S. C. H. (2000). Evolution of floral display in *Eichhornia paniculata*

(Pontederiaceae): direct and correlated responses to selection on flower size and

number. *Evolution*, *54*(5), 1533–1545. https://doi.org/10.1111/j.0014-

3820.2000.tb00699.x

Wright, S. (1931). Evolution in Mendelian Populations. *Genetics*, *16*(2), 97–159.

Wright, S. (1949). The Genetical Structure of Populations. *Annals of Eugenics*, *15*(1), 323–354.

https://doi.org/10.1111/j.1469-1809.1949.tb02451.x

Wright, S. I., Ness, R. W., Foxe, J. P., & Barrett, S. C. H. (2008). Genomic consequences of

outcrossing and selfing in plants. *International Journal of Plant Sciences*, *169*(1), 105–

118. https://doi.org/10.1086/523366

1270

1271

1272

Figure 1: Distributions of Multi-Locus Genotypes (MLGs) among LSI and SC *Lgh* populations in
western Europe. In black, the number of unique MLGs; In grey, the number of MLGs with less than 7
ramets; In color, MLGs with more than 10 ramets found in multiple populations. Proportions of each
MLGs over all the populations are plotted in the last right bar (All). More than half the samples are
unique genotypes.

1278

1279

Figure 2: Correlations between Σ_{CLON} and Σ_{SELF} found in 13 SC populations (light grey points and regression line) and 40 LSI populations (black points and regression line). 95% confidence intervals are given for LSI and SC populations in dark grey and light grey, respectively. Spearman rank-order correlation coefficients (r_s) and probabilities that Σ_{CLON} and Σ_{SELF} would not be correlated (p) are reported for LSI and SC populations.

1285

1286

Figure 3: Violin plots comparing distributions of genetic indices in LSI (L, grey) versus SC (S, white) populations. Violin plots are cut for their minimum and maximum values. Kruskall-Wallis tests are reported as H-statistics as well as the probability that LSI and SC population would present the same distribution of genetic indices.

1291
1292 **Figure 4:** Distributions of pairwise rho$_{st}$ within pairs of SC populations (white violin plot), pairs of LSI
1293 populations (light grey violin plot) and between pairs of LSI and SC populations (dark grey violin plot).
1294 Probabilities that pair of non-parametric distributions are equal are reported.

1295
Figure 5: Minimum spanning tree of the genetic distances (number of different alleles) between LSI
(grey points) and SC (white points) individuals. Arrows indicate some of the evidences of admixture
between LSI and SC lineages.

Supplementary Information (SI)

Situation of *Lgh* in USA and Mediterranean populations in previous studies (Dandelot 2004, Okada et al. 2009, Armitage et al. 2013):
- nearly monoclonal populations arguing for the ecological advantage of clonality (fragmentation and rhizome propagation) in these invasive populations.
But
- using weakly polymorphic markers,
- before the demonstration of the heterostyly and of the Late-acting self-incompatibility system in *Lgh*

Situation of *Lgh* in western Europe, studied here:
- Spatial context: The other compatible type populations are away from hundreds of meters to several kilometers (Portillo-Lemus et al. 2021)
- Clonality through fragmentation and rhizomes, as in other populations (Portillo-Lemus 2021).
- Massive blossoming with a lot of pollinators harvesting flowers (Portillo-Lemus 2021).
- Faster pollen tube elongation of pollen from the other compatible type (Portillo-Lemus et al. 2022)

Our working hypotheses about the main, realized reproductive modes in population:
- **LSI populations:** A very limited production of seeds (Portillo-Lemus et al. 2021)
Hypothesis 1: **preferential clonality** due to lack of compatible partners to produce seeds → Baker's conjecture on reproductive modes (Pannell et al. 2015).
Hypothesis 2: The rare seeds really contribute to the populations → biological feature of seeds or recombination would have advantage over clones. If so, are they mainly:
  Selfed seeds → reproductive insurance and advantage of recombination over clonality in a uniparental context?
  Outcrossed seeds → advantage of outcrossed seeds due to heterosis and/or recombination despite the lack of locally-available compatible partners → selection of long pollination ranges

- **SC populations:** Production of a lot of seeds in natural populations
Hypothesis 1: despite a huge production of seeds, **preferential clonality** due to the physiological and/or ecological advantage of clones over seed germination process?
Hypothesis 2: the mass of seeds does contribute to the populations or some evolutionary advantage of recombination in offspring compared to clonal fragmentation and rhizome propagation to adapt local conditions? If so, are these seeds mainly:
  Selfed seeds → within uniparental reproduction, biological feature of seeds or genetic recombination would provide advantage over clones?
Outcrossed seeds → Faster pollen tube elongation of pollen from the other type gives real advantage or advantage to allogamy to adapt new local conditions?

Supplementary information1: Synthesis of the questionings of our study to estimate reproductive

mode of 53 populations of invasive Lgh using genetic diversity on the autotetraploid part of its

genome.

Genotypic or genetic index	Range of values	Main driver	Behavior	references
$\beta*$	$\in [0; +\infty]$	Rate of clonality	↘ when c ↗ $\beta \to 0$ when $c \to 1$	Halkett et al. 2005, Arnaud-Haond et al. 2007, 2020, Stoeckel et al. 2021
$varF_{IS}$	$\in [0; +\infty]$	Rate of clonality	↗ when c ↗ $VarF_{IS} \to 0$ when $c \to 0$	Stoeckel & Masson 2014, Stoeckel et al. 2021
r_d	$\in [0; 1]$	Rate of selfing	slightly ↗ when c ↗ ↗ when s ↗ $r_d \to 1$ when $s \to 1$	Weir and Cockerham 1973, Navascues et al. 2010, Halkett et al. 2005
H_O	$\in [0; 1]$	Rate of selfing	↘ when s ↗ $H_O \to 0$ when $s \to 1$	David et al. 2007, Halkett et al. 2005
F_{IS}	$\in [-1; 1]$	Both rates	↘ when c ↗ $FIS \to -1$ when $c \to 1$ $FIS \to +1$ when $s \to 1$	Stoeckel & Masson 2014, Balloux et al. 2003, Stoeckel et al. 2021
H_E	$\in [0; 1]$	Rate of selfing	↘ when s ↗ $H_E \to 0$ when $s \to 1$	Glémin et al. 2001, Ritland & Ganders 1987
P_{ID-SIB}	$\in [0; +\infty]$	Both rates	↘ when c ↗ ↘ when s ↗	Waits et al. 2001
S_g**	$\in [0; 1]$	Rate of selfing	↗ when s ↗	David et al. 2007, Hardy et al. 2016

* similar behavior of the clonal richness R
** inferred values of s, according to the method of Hardy (2016)

1306 **Table SI1:** Hypotheses about the influence of clonality and selfing on the expected variations of
1307 genetic indices.

1308

see the attached xls file

Table SI2: All genetic and genotypic indices computed on the 53 sampled populations. Each index was obtained from 15 individuals sampled in each population. (See spreadsheet supplementary material).

Population with missing genotype	nb_missing_genotypes	nb_individuals_concerned
Bléré_Ls	1	1
Chambéon_écopôle_du_Forez	2	2
Le_Port_Ls	1	1
Plougonver_Léguer	1	1
Quevillon	1	1
Rennes_apigné_prairie	1	1
Roost-Warendin	1	1
Saint_Aignan_Couflons_Ls	1	1
Vichy	1	1

Over single SNP genotypes	N	%age
single SNP genotype P(cad)>70%	28610	0.99965
single SNP genotype P(cad)≤70%	10	0.00035
total	28620	1

Over individuals	N	%age
nb_individuals_with_full_genotype	785	0.98742
nb_individuals_with_one_missing_genotype	10	0.01258
total	795	1

Table SI3: Summary of the allele dosage. First subtable lists populations with individual genotyped with at least one individual with missing genotype at one locus. Second subtable reports the frequency of single SNP genotypes with posterior probability of allele dosage strictly superior to 70%. Third subtable reports the frequency of individuals with their 36 SNP genotypes assigned with posterior probability superior to 70%.

Table SI4: Distribution of the different repeated Multi-Locus Genotypes (MLGs) found with 36 SNPs among populations considering their mating system (LSI or SC). Figure 1 presents a summarized plot of this table. (See spreadsheet supplementary material).

	G	H$_O$	H$_E$	A$_E$	R	β	F$_{IS}$	VarF$_{IS}$	r$_d$	PID$_{SIB}$	S$_g$	Σ$_{clon}$	Σ$_{self}$
G	-	0.438	**0.013**	**0.009**	***	***	***	***	0.057	**0.008**	**0.008**	***	***
H$_O$	0.076	-	***	***	0.397	0.462	0.122	**0.025**	0.653	***	**0.032**	0.065	0.142
H$_E$	**0.244**	**0.742**	-	***	**0.01**	**0.014**	0.212	***	**0.026**	***	0.844	***	0.372
A$_E$	**0.255**	**0.722**	**0.82**	-	**0.008**	**0.013**	0.353	***	0.094	***	0.347	***	0.845
R	**0.991**	0.083	**0.251**	0.258	-	***	***	***	0.052	**0.007**	**0.005**	***	***
β	**0.91**	0.071	**0.238**	**0.24**	0.919	-	***	***	0.126	**0.009**	**0.004**	***	***
F$_{IS}$	**0.602**	-0.144	0.116	0.086	**0.603**	**0.557**	-	***	***	0.163	***	***	***
VarF$_{IS}$	**-0.579**	**-0.209**	**-0.435**	**-0.346**	**-0.589**	**-0.551**	**-0.626**	-	***	***	***	***	***
r$_d$	0.187	0.042	**0.207**	0.156	0.19	0.148	**0.432**	-0.365	-	**0.021**	***	**0.006**	***
PID$_{SIB}$	**0.259**	**0.729**	**0.981**	**0.833**	**0.265**	**0.251**	0.13	**-0.443**	0.214	-	0.936	***	0.286
S$_g$	**0.264**	**-0.202**	-0.019	-0.089	**0.277**	**0.28**	0.542	-0.379	0.353	-0.008	-	***	***
Σ$_{clon}$	**-0.79**	**-0.171**	**-0.376**	**-0.347**	**-0.799**	**-0.809**	**-0.604**	0.756	-0.254	**-0.387**	-0.34	-	***
Σ$_{self}$	**0.388**	**-0.136**	0.083	0.018	**0.399**	**0.361**	**0.723**	**-0.524**	**0.631**	0.099	**0.69**	**-0.451**	-

Table SI5: Correlations between genetic indices using non-parametric Kendall partial rank-order correlation. Below the diagonal, the values of the coefficient of correlation τ are consigned. Above, the related p-values are provided. Values inferior to the threshold 0.05, and the related coefficient τ were bold. The parameters Σ$_{CLON}$ and Σ$_{SELF}$ correspond to the sum of normalized indices sensitive to clonality and selfing, respectively.

2 **Figure SI1:** Map of the locations of the 53 sampled populations across France and Spain. In orange,
3 the late-acting self-incompatible (LSI) populations; In purple, the self-compatible (SC) populations. In
4 light blue, the Loire river system. In grey, the North-to-South (N-S) and West-to-East (W-E) scale
5 distances of the sampled populations.

Figure SI2.A: Correlation circles of the principal component analysis on the 17 centered and scaled
genetic indices measured in 53 *Lgh* populations in western Europe. The first horizontal dimension
accounting for 50.24% of the variance regroups indices related to clonality (number of genotypes *G*,
genotypic diversity *R*, clonal heterogeneity *D*, clonal evenness *Beta_Pareto*, variance of Fis *varFis*,
variance of expected heterozygosity *varHe*, variance of observed heterozygosity *varHo* and variance
of effective number of alleles varAe between loci). The second vertical dimension accounting for
23.39% of the variance with a main contribution of estimate of selfing Sg. Other indices, *i.e.*, mean
observed heterozygosity *Ho*, mean expected heterozygosity *He*, effective number of alleles *Ae*,
inbreeding coefficient *Mfis*, linkage disequilibrium *rd*, standard error of estimate of selfing *SE.Sg*,
unbiased probability of identity under panmixia *PIDu* and between sibs *PIDsib*. In blue and dashed
lines, the predictions of Σclon (*Sclon*) and Σself (*Sself*) provided as supplementary variables.

Figure SI2.B: Bar plot of the genetic index contributions to the first principal component. The
expected average contribution *1/(number of genetic indices)* is plotted as the red dashed line. VarFis
and indices of genotypic diversity (Pareto , R, G & D) are the main contributors to the first
component.

Figure SI2.C: Bar plot of the genetic index contributions to the second principal component. The
expected average contribution *1/(number of genetic indices)* is plotted as the red dashed line.
Estimates of selfing (Sg), mean observed heterozygosity (H_O), mean effective number of alleles (A_E)
and probabilities of identities (pid_u and pid_{sib}) are the main contributors to the second component.

Figure SI2.D: Plot of the projection of each population onto the two first principal components. SC
populations are plotted as red triangles and LSI populations as black diamonds. Concentration
ellipses around LSI and SC populations are plotted in black and red, respectively. Barycenters of the
two groups are represented as bigger black diamond and red triangle, respectively.

Figure SI3: Relationship and correlation between Σ_{CLON} and genetic indices (R, Pareto β, MFis,
VarFis, Ho, Ae, Sg, pidsib and $\bar{r}_d$). Per graph, all the 53 populations were plotted as black
dots. *: p<0.05, **: p<0.01, ***: p<0.001.

Figure SI4: Relationship and correlation between Σ$_{SELF}$ and genetic indices (R, Pareto β, MFis, VarFis, Ho, Ae, Sg, pidsib and $\bar{r}_d$). Per graph, all the 53 populations were plotted as black dots. *: p<0.05, **: p<0.01, ***: p<0.001.

Figure SI4: Distributions of the logarithms of the number of ramets per genet in LSI (grey) and
SC (white) populations. We report the Kruskal-Wallis probability that the two distributions
are identical.

47