2P-NucTag: on-demand phototagging for molecular analysis of functionally identified cortical neurons

Jingcheng Shi[1,2,3,*], Boaz Nutkovich[4,5,*], Dahlia Kushinsky[4,5,*], Bovey Y. Rao[1,2,3,*], Stephanie A. Herrlinger[1,2,*], Tiberiu S. Mihaila[1,2], Katayun Cohen-Kashi Malina[4,5], Cliodhna K. O'Toole[1,2], Margaret E. Conde Paredes[1,2,3,6], Hyun Choong Yong[1,2], Erdem Varol[6], Attila Losonczy[1,2,#, ‡], and Ivo Spiegel[4,#,‡]

[1] Department of Neuroscience, Columbia University, New York, NY, United States
[2] Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, United States
[3] Doctoral Program in Neurobiology and Behavior, Columbia University, New York, NY, United States
[4] Department of Brain Sciences, Weizmann Institute of Science, Rehovot, Israel
[5] Department of Molecular Neuroscience, Weizmann Institute of Science, Rehovot, Israel
[6] Tandon School of Engineering, New York University, New York, NY, United States

[*] Co-first authors
[#] Co-senior authors
[‡] Corresponding authors. Emails: al2856@columbia.edu, ivo.spiegel@weizmann.ac.il

Abstract:
Neural circuits are characterized by genetically and functionally diverse cell types. A mechanistic understanding of circuit function is predicated on linking the genetic and physiological properties of individual neurons. However, it remains highly challenging to map the functional properties of transcriptionally heterogeneous neuronal subtypes in mammalian cortical circuits *in vivo*. Here, we introduce a high-throughput two-photon nuclear phototagging (2P-NucTag) approach optimized for on-demand and indelible labeling of single neurons via a photoactivatable red fluorescent protein following *in vivo* functional characterization in behaving mice. We demonstrate the utility of this function-forward pipeline by selectively labeling and transcriptionally profiling previously inaccessible 'place' and 'silent' cells in the mouse hippocampus. Our results reveal unexpected differences in gene expression between these hippocampal pyramidal neurons with distinct spatial coding properties. Thus, 2P-NucTag opens a new way to uncover the molecular principles that govern the functional organization of neural circuits.

Main text:

Information processing in neural circuits requires precise interactions between molecularly and functionally diverse populations of neurons. Since gene expression ultimately dictates neuronal connectivity and function, a fundamental goal of neuroscience has been to characterize gene expression profiles of functionally defined neurons and to measure changes in gene expression associated with distinct functional states of neurons[1]. High-throughput transcriptomic approaches such as single-cell/-nucleus RNA-seq (sc/snRNA-seq) have greatly accelerated the identification of gene programs in molecularly distinct types of neurons at single-cell resolution[2-5]. However, subsequent *in vivo* and *ex vivo* functional and anatomical characterization of neuronal subtypes[6-10] remains staggeringly slow as it requires the generation and validation of subtype-specific molecular tools[11-13]. Recently, correlated *in vivo* Ca^{2+} imaging with *post hoc* spatial transcriptomics has been used to relate gene expression with *in vivo* function, but this approach is limited to spatially sparse GABAergic interneurons[14-17]. Therefore, a method to identify genes that are differently expressed in densely packed but functionally distinct glutamatergic pyramidal neurons (PNs), could significantly accelerate our understanding of how gene expression determines circuit function and behavior.

The inability to tag single functionally identified cortical PNs *in vivo* in behaving animals presents a significant challenge, as large-scale neural recordings have shown that PNs are highly heterogeneous in their physiological, anatomical, and response properties, and are spatially intermixed within neocortical and hippocampal circuits[13,18-24]. For example, PNs with distinct spatial coding properties are distributed throughout the dense cell body layer of the hippocampus[25-32]. However, the origin of this functional diversity in feature selectivity is largely unknown, and it remains unclear if gene expression differences are associated with discrete and transient functional cell states[32-37]. Thus, there is a critical unmet need for function-forward approaches that directly map *in vivo* physiological and transcriptional profiles in cortical circuits in behaving animals.

Previous attempts using Ca^{2+} and light-dependent labeling of transiently active neurons[38] were limited by their spatial resolution, deficiencies in targeting neurons with high baseline intracellular Ca^{2+} levels, and the inability to label neurons that decrease their activity in response to behavioral state or sensory stimuli. Similarly, immediate early gene-dependent labeling approaches[39-42] lack the temporal and spatial resolution to faithfully report the precise activity patterns and response properties of single neurons. Finally, previous attempts to tag cortical neurons with photoactivatable fluorescent proteins[43] at single-cell cellular precision have been deployed with limited success[44].

Here we introduce a robust *in vivo* pipeline (2P-NucTag), based on a photoactivatable red fluorescent protein (PAmCherry) and a genetically encoded green Ca^{2+} indicator (GCamP7f), that optimizes a previously described framework[45] mainly used *ex vivo*. Our approach combines large-scale *in vivo* two-photon (2P) functional imaging of cortical PNs with reliable and selective 2P phototagging of nuclei in a subset of neurons based on their functional properties. Using fluorescence-activated cell sorting (FACS) to isolate phototagged neuronal nuclei *post hoc*, combined with our recently developed Meso-seq approach for transcriptomics in ultra-sparse populations[46], we achieve previously unattainable molecular characterization of *functionally* identified PNs *in vivo* in behaving animals.

In vivo two-photon phototagging with 2P-NucTag

A major challenge in combining *in vivo* functional recording with stable tagging in the same neuron is the co-expression of an activity sensor and a photoactivatable tag with spectrally separable fluorescent imaging and photoactivation. We overcame this challenge by generating a bicistronic construct on a recombinant adeno-associated viral (rAAV) backbone that co-expresses cytosolic GCaMP7f [47] for 2P Ca²⁺ imaging and a nucleus-targeted photoactivatable red fluorescent protein (H2B-PAmCherry) for 2P phototagging[45] under a promoter that is selective for cortical glutamatergic neurons[48] (2P-NucTag, Fig. 1A). Upon injection of the 2P-NucTag rAAV into the CA1 region of the mouse dorsal hippocampus to label CA1 PNs, we found that GCaMP7f is properly expressed in the perinuclear space of the infected PNs (Fig. 1B). Targeting

Fig. 1. *In vivo* two-photon phototagging with 2P-NucTag. (A) Schematics of the 2P-NucTag pipeline. Top: bicistronic rAAV construct, injection to the hippocampus. Middle: *in vivo* two-photon (2P) GCaMP- Ca²⁺ population

imaging followed by 2P PAmCherry photoactivation, fluorescence-activated cell sorting (*FACS*), and mesoscale
sequencing (*Meso-seq*). (**B**) Top: representative *in vivo* time-averaged (6 frames average) 2P images of individual
cells before (*Pre*) and after (*Post*) *in vivo* two-photon PAmCherry photoactivation in the CA1 pyramidal layer of the
mouse dorsal hippocampus. Individual nuclei were photoactivated with 810-nm 2P laser chessboard scanning region-
of-interest (ROI, yellow boxes) over target nuclei (70 x 70 pixel for each ROI, 0.1 μm/px, 1.3 ms/px total pixel dwell
time, 6,370 ms total scan time per ROI. Laser power was 40 mW measured after the objective) with a 3-dimensional
acousto-optical deflector microscope (3D-AOD). Gray: GCaMP7f (940 nm excitation), magenta: PAmCherry, (1040
nm excitation). Scale bar: 50 μm. Middle and bottom: imprints of letters '*BI*' and '*ZI*' following patterned *in vivo* two-
photon photoactivation in the hippocampal CA1 pyramidal layer (scale bar, 50 μm). (**C**) Characterization of *in vivo*
2P photoactivation parameters for PAmCherry: duration, wavelength, laser power (measured after the objective) (n =
11-12 cells per condition). Relative change in PAmCherry red fluorescence (ΔF) is based on normalizing the tagged
nuclei fluorescence to the fluorescence of neighboring untagged nuclei measured with 1040 nm excitation. (**D**) *In vivo*
stability of the PAmCherry fluorescence signal over days after a single photoactivation scan (n = 8 cells). (**E**)
Representative time-averaged images from z-stacks of photoactivated nuclei *in vivo* (magenta: PAmCherry, scale bar:
100 μm). Middle: *ex vivo post hoc* confocal z-stack image of the same field of view (FOV, magenta: PAmCherry).
Right: registered *in vivo* and *ex vivo* images following non-rigid image transformation (magenta: *in vivo*, yellow: *ex*
vivo, see methods). (**F**) Left: 3D overlay of tagged nuclei registered between *in vivo* (magenta) and *ex vivo* (yellow)
z-stacks with increasing lateral resolution (as in *E*). Gray box represents the segmented area for subsequent images.
Right: normalized lateral (*x-y*, left) and axial (*z*, right) fluorescence profiles (mean ± s.e.m.) of tagged cells *in vivo*
(magenta, n = 1 mouse, 200 cells). Yellow: mean ± s.e.m. of *ex vivo* confocal images (as in *E* and *F,* same mouse and
nuclei). Inset: (x-y) top: average *in vivo* maximum z–projection, bottom: average *ex vivo* maximum z projection; (z)
top: average *in vivo* lateral projection, bottom: average *ex vivo* lateral projection. Scale bar: 10 μm. Boxplots show the
25^{th}, 50^{th} (median), and 75^{th} quartile ranges, with the whiskers extending to 1.5 interquartile ranges below or above
the 25^{th} or 75^{th} quartiles, respectively. Outliers are defined as values extending beyond the whisker ranges.
79
the nuclei of GCaMP-expressing neurons, we achieved rapid nuclear PAmCherry photoconversion
using 810-nm excitation light on a three-dimensional acousto-optical deflector microscope (3D-
AOD)[49,50] (Fig. 1B; supplementary movie 1, see methods) and orthogonal GCaMP-Ca^{2+} activity
imaging with 940 nm excitation light (fig. S1A,B). Photoconverted PAmCherry red fluorescence
was detected at >1000 nm excitation (1040 or 1070 nm, see methods) and was localized to the
targeted nuclei (Fig. 1B, top). 2P-NucTag enabled the imprinting of arbitrary tagging patterns into
the CA1 pyramidal cell layer with 3D-AOD scanning, showing spatially precise photoconversion
of H2B-PAmCherry with single-nuclear and even sub-nuclear resolution (Fig. 1B, middle and
bottom). We carried out a detailed characterization of wavelength, laser power, and duration-
dependence of PAmCherry photoactivation *in vivo* to identify the optimal parameters for spatially
precise photoactivation (Fig. 1C). Based on our results, we opted for 810 nm excitation light with
37-42 mW laser power (measured after the objective), 1.3 ms/pixel dwell time over 70 pixel x 70
pixel regions-of-interest (ROIs*,* with 0.1 μm/pixel resolution). The 810-nm wavelength is
spectrally separated from the GCaMP-based Ca^{2+} imaging wavelength at 940 nm, and these
photoactivation wavelengths, laser power, and duration parameters yielded robust increases in
PAmCherry fluorescence of targeted nuclei (192-379% increase in PAmCherry red fluorescence
visualized using1040 nm excitation: n = 36 cells, 295% ± 8% ΔF/F, mean ± s.e.m.) after single
scans while minimizing total scan time and power. We confirmed that phototagged nuclei
remained detectable over multiple days after photolabeling (Fig. 1D). We next photoactivated a
subset of CA1 PNs in a large (700 x 700 μm) field of view (FOV), similar to the FOV size used
for *in vivo* 2P population imaging experiments in CA1 (Fig. 1E, left). We found clear
photoactivation of individual target nuclei when visualized *in vivo*. We then confirmed that *in vivo*
photolabeling was preserved in *post hoc* histological slices (Fig. 1E, middle), and that phototagged
cells can be reliably registered across *in vivo* z-stacks and *post hoc* confocal images (Fig. 1E, right,
Fig. 1F, fig. S1C,D, supplementary movie 2). In addition, we segmented the nuclei from both the

in vivo and *ex vivo* z-stacks to observe the average axial and lateral fluorescence profiles of
phototagged nuclei, demonstrating single-nucleus resolution (Fig 1F, right). Thus, 2P-NucTag
enables high-throughput, indelible phototagging of neuronal nuclei *in vivo* that can also be
identified via our registration pipeline for *post hoc* analyses *ex vivo*.
109
Phototagging of functionally identified PNs in the hippocampus *in vivo*
To demonstrate the utility of 2P-NucTag for on-demand labeling of *functionally* defined PNs, we
deployed 2P-NucTag to label hippocampal PNs located in the dorsal CA1 region which have well-
established spatial coding heterogeneity[25-32]. Only a subset of hippocampal PNs ('place cells')
exhibit reliable spatial tuning for a location ('place field') during exploration[51]. Place cell identity
is thought to be randomly allocated within a seemingly homogenous population of PNs[52], and is
highly dynamic, with the majority of cells changing their spatial tuning properties over the
timescale of days[32-37]. The spatial intermingling of place cells with active-non-place and silent
PNs[53-56] without apparent topographical organization in the densely packed CA1 pyramidal
layer[32,56] allows us to test the utility of 2P-NucTag in selectively labeling PNs occupying these
distinct functional states.
To selectively label CA1 PNs with distinct spatial coding properties, we trained mice in a
spatial navigation task for water rewards in a linear virtual reality environment[57-59] and performed
in vivo 2P GCaMP- Ca^{2+} imaging of PNs (Fig. 2A,B). We reliably detected

Fig. 2. Selective phototagging of place cells in the hippocampus with 2P-NucTag. (**A**) Pipeline for two-photon (*2P*) phototagging of functionally identified hippocampal neurons during spatial navigation. (**B**) Left: schematics of 2P imaging setup in virtual reality (VR). Head-fixed mice are trained to run for a water reward in a 4-m long linear VR corridor projected onto LCD screens surrounding the animal. At the end of the corridor, mice are teleported back to the start position after a 2-second delay. Right: example 2P field of view (FOV) of GCaMP in the CA1 pyramidal layer. Scale bar: 100 μm.(**C**) Left: Traces of relative GCaMP-Ca^{2+} fluorescence changes (ΔF/F) from five example CA1 place cells during VR spatial navigation. Right: heatmaps of normalized ΔF/F activity from three example place cells over 20 laps during VR navigation. (**D**) Left: heatmap of all CA1PNs detected with Suite2p/Cellpose in the FOV shown in *B*. Identified place cells are marked with an orange box. Right: Zoomed-in heatmap of place cell tuning curves. (**E**) Left: spatial mask (orange) of identified place cells from *D* in the FOV. Right: PAmCherry fluorescence (magenta) of tagged nuclei after 2P phototagging. Scale bar: 100 μm. (**F**) Left: overlay of spatial masks of identified CA1PNs and tagged nuclei for the FOVs in *E*. Scale bar: 100 μm. Right: tagging efficacy, defined as the fraction of successfully tagged place cell nuclei (93.3% ± 4.2%, mean ± s.e.m., n = 5 mice). (**G**) Left: Proportion of single, double, and triple-tagged nuclei following phototagging of a single place cell. Right: relative change in PAmCherry red fluorescence (1070 nm excitation) for non-tagged cells in the FOV after 2P imaging (green), after 2P phototagging of targeted place cell nuclei (orange) and off-target nuclei (gray, n = 5 mice). Boxplots show the 25th, 50th (median), and 75th quartile ranges, with the whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles, respectively. Outliers are defined as values extending beyond the whisker ranges.

GCaMP-Ca^{2+} signals from individual PNs (Fig. 2C, fig. S1C,D). We analyzed the basic characteristics of GCaMP7f in the bicistronic 2P-NucTag construct and confirmed that the basic properties of the indicator are similar to those for a previously published single-construct version of GCaMP7f [47] (fig. S1C,D). We classified all PNs in the imaging FOV as place cells, active non-place cells, or silent cells (Fig. 2D, see methods). Following the functional identification of PNs, we generated spatial masks of place cell locations in the FOV and used these masks to guide 2P phototagging of all identified place cells in the imaging FOV with 3D-AOD (n = 5 mice, Fig. 2E, F). In a separate set of mice (n = 4 mice), we tagged a subset of CA1 PNs that exhibited no detectable activity during the imaging session ('silent cells' see methods, fig. S2, Table 1). We could reliably register photoactivated nuclei to the spatial masks of functional profiles that we generated for cells of interest (Fig. 2F, fig. S2D,E, fig. S3, see methods). We quantified the number of place cell nuclei that we successfully photoactivated and compared it to the number of spatial masks we generated for each animal. Across all animals, we were able to phototag 93.3% ± 4.2% (n = 5 mice) of all targeted place cell masks (Fig. 2F). Aiming to tag approximately equal numbers of place and silent cells, we did not attempt to tag all the silent cells, which are more abundant (Table 1).

To assess the accuracy of 2P photoactivation in the densely packed CA1 pyramidal layer, we quantified the number of tagged nuclei for each targeted cell and found that photoactivation is well restricted to the target cells' nuclei with limited off-target labeled nuclei, which had a smaller increase in mCherry fluorescence (Fig. 2G, fig. S2F). Furthermore, 2P GCaMP-Ca^{2+} imaging at 940 nm over the course of the imaging session (18-40 min, see methods) resulted in a minimal increase of mCherry red fluorescence (37.4% ± 16.2% ΔF/F, n = 5 mice, Fig. 2G, fig. S2F, 1070 nm excitation). Targeted place cells demonstrated higher mCherry fluorescence (487.7% ± 67% ΔF/F, n = 5 mice) compared to the background and off-target fluorescence (210.4% ± 32.1% ΔF/F, n = 5 mice). Mouse behavior and the quality of GCaMP recordings were consistent between the two groups of mice (silent and place, fig. S2G-I). In sum, 2P-NucTag offers on-demand *in vivo* labeling of functionally defined neurons with high efficacy and accuracy.

Transcriptional profiling of functionally identified CA1 PNs

The ability to tag single cells *in vivo* enables a powerful new form of hypothesis generation and testing by which, for the first time, functionally defined cells with known behavioral relevance can be isolated and characterized. To demonstrate this, we sought to interrogate transcriptomic signatures of the functional 'place' and 'silent' cell states of CA1 PN.

Brain tissue containing dorsal CA1 was collected, and nuclei were dissociated and stained with CoreLite 488-conjugated NeuN antibody and DAPI to identify neuronal nuclei (see methods). *In vivo* photoactivated nuclei were identified by bright mCherry fluorescence (Fig. 3A,B) and were separated out from the non-photolabeled neuronal nuclei by fluorescent-activated cell sorting (FACS), enabling an estimated 18-67% recovery of all photoactivated nuclei identified during *in vivo* imaging (Fig. 3C). To determine whether place cells and silent cells differed in their gene expression programs, RNA-seq was performed on both populations of sorted nuclei by Meso-seq, an approach that enables reliable identification

Fig. 3. *Post hoc* transcriptional profiling of phototagged place and silent cells. (A) Schematics of *in vivo* photoactivated nuclei. 'Place' cell sample and 'Silent' cell samples from different mice were collected for FACS and

Meso-seq. (**B**) Representative FACS graph. Gating for mCherry was set after the first 5000 events of DAPI+ nuclei
to the border of the 'dim' mCherry+ population to separate out the sparse and high-intensity mCherry+ population.
Bright mCherry+ NeuN+ populations were collected as the photoactivated nuclei. (**C**) Top: number of FACS sorted
nuclei from 'place' and 'silent' samples (n = 9, 17-79 sorted nuclei, 40.3 ± 6.43, mean ± s.e.m.). Bottom: Proportion
of FACS sorted nuclei compared to the number of *in vivo* photoactivated nuclei (n = 9, 18.45% to 66.39% FACS
recovery, 39.94 ± 6.30%, mean ± s.e.m.). (**D**) Volcano plot of Meso-seq differential expressed gene (DEG) analysis
for 'place' and 'silent' cells (significantly different genes are shown in orange and blue. Orange: enriched in place
cells; blue: enriched in silent cells). (**E**) Meso-seq MA plot depicting DeSeq2 normalized gene counts versus log2 fold
change of silent/place samples. Genes that are significantly different are labeled in orange and blue (same as above).
Genes shown in panels *F* & *G* are highlighted and labeled in *E*. (**F**) Bar graph showing the normalized counts for
genes that are not differentially expressed (FDR adjusted p-value. *<0.05, **<0.001, ***<0.001, PyDeSeq2.
Otherwise, comparisons are not significant). (**G**) Bar graph showing the normalized counts for differentially expressed
genes (FDR adjusted p-value. *<0.05, **<0.001,***<0.001, PyDeSeq2. Otherwise, comparisons are not significant).
(**H**) Gene ontology analysis performed on all differentially expressed genes. Vertical line: FDR-adjusted p value of
0.05. NES = normalized enrichment score. Boxplots show the 25th, 50th (median), and 75th quartile ranges, with the
whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles, respectively. Outliers are
defined as values extending beyond the whisker ranges.
202
of differentially expressed genes in ultra-low amounts for FACS-isolated neuronal nuclei (i.e., tens
of sorted nuclei, Fig. 3D)[46]. After FACS-isolating the photo-tagged nuclei, libraries were generated
with the Meso-seq protocol and were sequenced at a depth of 40-60 million reads per library (fig.
S4A). Reads were aligned to the mm39 mouse genome assembly with STAR, counted with HTSeq,
and gene expression patterns were compared between nuclei isolated from mice in which place
cells were tagged (n = 5) and mice in which silent cells were tagged (n = 4) via PyDESeq2.
In both populations, canonical CA1 PN marker genes (*Map2, Actb, Dlg4, Neurod6*) were
highly and non-differentially expressed, supporting the cellular precision of our tagging approach
(Fig. 3E,F). However, 219 genes were identified as differentially expressed in a significant manner
between silent and place cells (Fig. 3E, see Supplementary Table). In some cases, specific genes
were reliably identified in one group and absent in the other. For example, no counts were
measured for an inward rectifying potassium channel (*Kcnj12*) in place cell samples, but the
expression was present in all silent cell samples. Conversely, mRNA from two transcription factors
of the zinc finger protein family (*Zfp84, Zfp977*) and a protocadherin gene (*Pcdha8*) were
identified in every place cell sample and in none of the silent cell samples (Fig. 3G).
To ensure that the observed differences did not arise from other sources of variation
between our place and silent cell samples, we performed several additional analyses, investigating
potential contributions from anatomical positioning, transcriptional responses to neural activity,
cellular health, and sex. Genes for which expression has been shown to vary along the dorsoventral
and proximodistal axes of CA1 were not differentially expressed in our place and silent-cell nuclei
(fig. S4B)[26,60], suggesting that any observed differences are unlikely to have originated from
differences in anatomical positioning during cell tagging. Immediate early gene counts (*Fos, Arc,*
Egr1, Npas4) did not differ (Fig. 3F)[61-63], suggesting that the ≥25 hours between the last behavior
session and tissue collection was sufficient time to eliminate the impact on our data of immediate
transcriptional responses to neural activity. Apoptotic gene counts also did not differ between the
two groups (fig. S4), suggesting that there was no significant difference in cell health between the
silent and place cells[64]. Finally, to test for gross molecular differences between the cells in an
unbiased fashion, we performed gene set enrichment analysis on fifty hallmark gene sets from the
mouse molecular signatures database and found no significant differences between silent and place
cell samples (Fig. 3H). Since the sex of mice used for both groups was not well balanced in this
study (Table 1), it is possible that gene expression differences originated from the mouse's sex

rather than the functional identity of the tagged cells. To explore this possibility, we compared
gene expression in randomly tagged (function-blind) CA1 PN nuclei from both male (n = 2) and
female (n = 2) mice. Although we found that known sex-specific genes[65] were differentially
expressed, none of the genes identified as enriched in either silent or place cells had significant
sex associations (fig. S5C,D).
To assess if our differentially expressed genes-of-interest signified upregulation or
downregulation in place cells from the mean or, conversely, downregulation or upregulation in
silent cells, we compared gene expression in place cells and silent cells to randomly tagged cells
separately. We found that *Kcnj12* expression was enriched in silent cells compared to random
cells. *Zfp84* and *Zfp977* were significantly downregulated in silent cells compared to random cells,
while *Pcdha8* expression was significantly enriched in place cells compared to random cells (Fig.
S5A,B).
Finally, to confirm the compatibility of 2P-NucTag with other commonly used *post*
hoc approaches for molecular analyses, we performed immunohistochemistry and *in*
situ hybridization on dorsal CA1 tissues with randomly photoactivated nuclei (fig. S6). We
immunolabeled for Calbindin protein and targeted Calb1 transcripts using the RNAScope[(TM)]
Multiplex Fluorescent assay, and found that 2P-NucTag signal is indeed stable and compatible
with the PFA-fixed and fixed-frozen tissue processing commonly used for these imaging
approaches.
In summary, our data demonstrate that, for the first time, we have the ability to apply
differential gene expression analysis on populations of cells from the same brain region that differ
only by their functional identity. Moreover, our in-depth sequencing approach can indeed identify
differences in gene expression between functionally defined cells tagged *in vivo* with 2P-NucTag.
257
Discussion
Comprehensive molecular-functional characterization of cortical circuits remains elusive due to
the limited toolkit for correlated *in vivo* functional recording techniques and *post hoc* molecular
analyses. Here, we introduced 2P-NucTag, a robust function-forward approach combining *in vivo*
imaging, on-demand single-nucleus tagging, and *post hoc* transcriptomics. This pipeline
seamlessly integrates 2P Ca^{2+} imaging in behaving mice, 2P phototagging of functionally
identified neuronal nuclei, FACS-isolation of the tagged nuclei, and subsequent analyses of
isolated nuclei with a mesoscale protocol for in-depth transcriptomics in ultra-sparse neuronal
populations.
We demonstrated the utility of 2P-NucTag by selectively labeling and analyzing
transcriptional profiles of two functionally orthogonal subpopulations of hippocampal PNs, 'place
cells' and 'silent cells'. Unexpectedly, our data revealed a number of genes that are putatively
differentially expressed between these distinctly transient physiological cell states[32-37]. The low
counts for some of these genes indicate that further replicates may be necessary to confirm their
differential expression. Nevertheless, several of the identified genes generate new hypotheses
about the origin of these two functional cell identities, which had canonically been considered
otherwise indistinguishable. Specifically, the marked enrichment of an inward rectifying
potassium channel (*Kcnj12*) in silent cells calls for experiments to investigate whether silent cells
are intrinsically less excitable than other CA1 PNs[66]. Future experiments will need to test the
function of specific genes that we have identified with our tagging technique which could reveal a
non-random allocation of functional activity in CA1 PNs during learning and experience. In
particular, future application of 2P-NucTag to selectively tag PNs with highly distinct and shared
coding properties, including place field propensity[25,33,67,68], recruitment to 'replay' and 'preplay'

events[69-72] and participation in functionally interconnected subnetworks[73] may uncover long
hypothesized genetic backbones of functional heterogeneity among PNs.

Our results showed low levels of unintended PAmCherry photoactivation during functional
imaging as well as limited off-target labeling. Although the activation level of these background
and off-target neurons is lower and the actual targeted cells' photoactivation can be distinguished
through *in vivo* imaging, FACS, and *ex vivo* confocal imaging, this presents a complicating factor
when setting gating levels for FACS. Therefore, in this study, we sorted only the brightest
population of cells, which shows high levels of mCherry fluorescence. Future optimization of the
phototagging vector and the 2P tagging parameters can help refine photoactivation specificity.
Nevertheless, our proof-of-principle implementation of 2P-NucTag to functionally heterogeneous
hippocampal PNs that are spatially intermixed in the densely packed pyramidal cell layer
demonstrates that 2P-NucTag should be readily applicable to neocortical tissue with lower cell
density.

Beyond this first application to the mouse hippocampus, 2P-NucTag will have broad appeal
for correlated structural, molecular, and functional analysis of neural circuits. Our approach offers
the ability to bridge molecular and functional architecture in cortical circuits and facilitates the
identification of candidate genes that determine the distinct circuit functions of neurons. Our
pipeline should be potent for uncovering molecular signatures of phenomenologically described
neuronal subsets with distinct feature selectivity, task-related activity, and longitudinal stability
that are spatially intermixed in cortical circuits[18,20,22,74-78].

Our approach can also be extended in multiple directions. Firstly, as 2P-NucTag is
compatible with *post hoc* histological analysis of imaged tissue, it can be combined with spatially
resolved transcriptomics[14,15,79,80] following automated registration between *in vivo* and *post hoc*
images[81,82]. Secondly, *in vitro* electrophysiological recordings from tagged cells would allow for
detailed *ex vivo* physiological, molecular, and anatomical characterization of *in vivo* tagged cells[6-
9]. Thirdly, in principle, our pipeline is compatible with sc/snRNA-seq, given the single-cell
resolution achieved through targeted photoactivation. Fourthly, further optimization of
phototagging with multi-color fluorophores may enable simultaneous labeling of multiple
functionally defined cell types in the same animal, while cytosolic tags[43] can promote anatomical
labeling of subcellular axonal or dendritic compartments for downstream structural and
connectivity analyses. Lastly, the on-demand nature of photoactivation and its days-long stability
following single photoactivation makes our pipeline a valuable tool for precise analysis of
transcriptional trajectories following cellular plasticity events during, for example, behavioral
learning[49,63,83,84].

In summary, our novel phototagging approach aids in expanding our knowledge of the
fundamental relationship between the molecular and functional architecture of mammalian cortical
circuits and allows for the identification of candidate genes that determine the distinct circuit
functions of seemingly homogeneous populations of neurons. Beyond the proof-of-principle
implementation to hippocampal circuits, our approach provides a general framework and roadmap
for linking genes to cells across neural circuits and model organisms.

Materials and Methods
Animals
All animal care and experiment procedures were in accordance with the guidelines of the National Institute of Health. Animal protocols were approved by the Columbia University Institutional Animal Care and Use Committee and the Weizmann Institute of Science Institutional Animal Care and Use Committee. Mice were group-housed under normal lighting conditions in a 12-hour light/dark cycle. *Ad libitum* water was provided until the beginning of training for the spatial navigation task.

Plasmids and Viral Constructs
pAAV-CW3SL-GCaMP7f-4Ala-H2B-PAmCherry was generated by standard cloning techniques. GCaMP7f was PCR-amplified from Addgene plasmid #104492 with a 3' Primer that contained sequences encoding four Alanine residues and the 2PA sequence (both in frame with the coding sequence of GCaMP7f). H2B-PAmCherry was amplified from Addgene plasmid #133419. The PCR products were then subcloned by Gibson-assembly into Addgene plasmid #61463 after removing EGFP from this plasmid by restriction with ClaI and EcoRI. The sequence of the cloned plasmid was validated by Sanger sequencing and GCaMP7f + 4xAla, P2A and H2B-PAmCherry were all found to be in frame. The plasmid was packaged into AAVDJ at a viral titer of 7.26E+15 essentially as described[46].

Surgery
All procedures were performed with mice under anesthesia using isoflurane (4% induction, 1.5% maintenance in 95% oxygen). Mice's body temperature was maintained using a heating pad both during and after the procedure. Surgeries were performed on a stereotaxic instrument (Kopf Instruments). Before incision, mice were given subcutaneous meloxicam, as well as bupivacaine at the incision site. Doses were calculated based on the animal's weight. An incision above the skull was made to expose bregma and lambda for vertical alignment. Skull surfaces were cleaned and scored to improve dental cement adhesion. For viral injection, a glass capillary loaded with rAAV is attached to a Nanoject device (Drummond Scientific).

For all experimental mice, viruses were injected unilaterally in the left dorsal CA1 at 4 depths using the coordinates: -2.2 AP, -1.75 ML, and -1.2, -1.1, -1.0, -0.9 DV (relative to Bregma). At each depth, 75nl of AAVDJ-CW3SL-GCaMP7f-4Ala-H2B-PAmCherry was injected. After injection, surgical sites were closed with sutures. Three days after injection, the skull was exposed and a 3mm craniotomy was made centered at the same coordinate of the injection site. Dura was removed, and the cortex was slowly aspirated with continuous irrigation of cold 1X PBS until the fiber tract above the hippocampus was visible. A 3-mm imaging cannula fitted with a 3mm glass coverslip was implanted over the craniotomy site. Cannulas were secured by Vetbond. A custom titanium headpost for head-fixation was secured first with C&B Metabond (Parkell) and then dental acrylic. At the end of each procedure, the mice received a 1.0 ml saline injection subcutaneously and recovered in their home cage with heating applied. Mice were monitored for 3 days after the procedure.

Behavior paradigm
Mice were first water-deprived and habituated to handling and head fixation at least 7 days after implant surgery. They were then exposed to a 4-m long linear virtual reality (VR) corridor[57-59] that

stayed consistent in the training and recording. At the end of the environment, an inter-trial interval
of 2 seconds of blank screen was included before the start of the next lap. For the next 10-14 days,
mice were trained to run through the virtual environment and lick for a 5% sucrose reward. The
rewards were first randomly distributed across the environment, and the number of rewards was
slowly reduced from 30 at the beginning of the training to 2 when the mouse was deemed ready
for recording. The final reward location was fixed toward the end of the VR environment. Mice
were trained to run at least 30-60 laps in the environment. During behavioral imaging, mice were
imagined during a single VR session (range: 18-40 min, 26 ± 3 min, n = 9 mice).

In vivo two-photon imaging and data processing

2P functional imaging was conducted using an 8-kHz resonant scanner (Bruker) and a 16x near-
infrared (NIR) water immersion objective (Nikon, 0.8 NA, 3.0-mm working distance). For
population imaging, a field of view of 700 μm x 700 μm was acquired at 30 Hz, 512 x 512 pixels
using a 940-nm laser (Chameleon Ultra II, Coherent, 45-91 mW after the objective). Red
(PAmCherry) and green (GCaMP7f) channels were separated by an emission cube set (green,
HQ525/70 m-2p; red, HQ607/45 m-2p; 575dcxr, Chroma Technology), and fluorescence signals
were collected with GaAsP photomultiplier tube modules (7422P-40, Hamamatsu). Following the
acquisition of two-photon imaging data, Ca^{2+} imaging data was structured and aligned with
behavior data using the SIMA analysis package[85]. CA1 ROIs were detected using the Suite2p
(v0.14.2) package [86]. To allow detection of all potential ROIs regardless of their activities during
the recording. Suite2p was run with Cellpose ('anatomical_only')[87] for the ROI detection step. The
pre-trained cyto2 model included in the published Cellpose package was used for ROI detection.
When capturing two-photon z-stack images of photoactivated PAmCherry nuclei, a fixed
wavelength 1070-nm laser (Fidelity-2W, Coherent) was used for excitation.

In vivo two-photon phototagging

Photoactivation was conducted using a three-dimensional random-access acousto-optical (3D-
AOD) microscope (3D Atlas, Femtonics)[16,50]. Mice were head-fixed and anesthetized with
isoflurane to minimize motion and increase the spatial precision of phototagging. The same 16x
NIR water immersion objective was used to find the same field of view as in the functional
recordings. Photoactivation was performed at 810-nm (Chameleon Ultra II, Coherent). Two-
photon images of the mCherry red fluorescence were taken before and after photoactivation using
a 1040-nm excitation laser (Alcor 1040-5W, Spark Lasers). Red (mCherry) and green (GCaMP7f)
channels were separated by an emission cube set (green, HQ520/60 m-2p; red, HQ650/160 m-2p;
565dcxr, Chroma Technology), and fluorescence signals were collected with GaAsP
photomultiplier tube modules (7422P-40, Hamamatsu). Two-photon images of the mCherry red
fluorescence were taken before and after photoactivation using a 1040-nm excitation laser (Alcor
1040-5W, Spark Lasers). For photoactivation, a 7 x 7 μm, 0.1 μm /pixel scanning pattern was
placed on the cell to be photoactivated. Each pixel was activated for a total dwell time of 1.3 ms
with a laser power of 40 mW. This gave the total scanning time of each cell at 6,370 ms. Following
photoactivation, a z-stack was taken for each mouse to assess the photoactivation efficacy.

For phototagging of 'place cells' and 'silent cells', photoactivation experiments were conducted a
day after functional recording sessions. The imaging field of view of 700 μm x 700 μm was
matched between the 3D-AOD microscope and the time-averaged GCaMP image from functional
recording. After confirming the same field of view as functional imaging, the viewport was

zoomed in to a dimension of 250 x 250 μm for more effective identification of targeted cells
according to the generated spatial masks. Following photoactivation of all cells, a z-stack image
was taken for each mouse to confirm the tagging accuracy. Mice were given *ad libitum* water after
functional imaging for at least 12 hours before photoactivation. During the session, mice were
monitored every 10 min for breathing rate and reflexes. Heating and eye ointment were applied.
Following the photoactivation, mice were returned to the home cage to recover with a heating pad.
419
Tissue dissociation and preparation of nuclei for FACS
For all animals used in this manuscript, tissue collection was performed at the same time of the
day (5 pm) and at least one hour after the photoactivation. Mice were euthanized using CO_2. The
headpost and metal cannulas were removed. Dorsal CA1 of the hippocampus was collected by first
using a 3-mm biopsy punch to cut a circular section of tissue the same size as the craniotomy. A
microspatula was used to remove the shallow section of tissue that contained dorsal CA1. The
tissue was placed in a 1.5 ml RNase-free Eppendorf tube and snap-frozen in liquid nitrogen.
Tissues were stored at -80 °C until the start of the nuclei isolation.
428
Nuclei were prepared for FACS sorting essentially as previously described[46]. In short, to isolate
the nuclei, each collected tissue was transferred to a dounce tissue homogenizer (DWK Life
Sciences) with 1ml of homogenization buffer (10 mM Tris Buffer, 250mM Sucrose, 25 mM KCl,
5mM $MgCl_2$, 0.1mM DTT, 0.1% Triton X-100, 1X Protease Inhibitor Cocktail, and 40U/μl
RNAsin Plus RNase Inhibitor in nuclease-free water). Loose and tight pestles were used to break
apart the tissue 10 times each. Following the homogenization, 1 ml of homogenization buffer was
added to each dounce, and the homogenate was pipetted up and down to further break apart tissue
before being passed through a 30 μm cell strainer (Miltenyi Biotec) and collected in a 15 ml conical
tube.
438
The homogenate was centrifuged at 4 °C, 700g for 8 min. The supernatants were then removed
from the visible cell pellet. The cell pellet was resuspended with 800 μl of blocking buffer (1X
PBS, 1% BSA, and 40 U/μl RNasin Plus RNase Inhibitor in nuclease-free water). The
resuspension was incubated on ice for 15 min and transferred to a 1.5 ml tube. 2 μl of CoraLite
Plus 488-conjugated NeuN Monoclonal antibody (Proteintech) was added to the resuspension, and
incubated on an orbital rotator at 4 °C for 30 min. After incubation, nuclei were centrifuged at 4
°C, 700 g for 8 min. The supernatant was removed, and the cell pellet was resuspended with 1000
μl of blocking buffer. DAPI was added to the suspension at a final concentration of 0.001 mg/ml,
and the samples were passed through a 40- μm Flowmi Cell Strainer (Bel-Art). All samples were
kept on ice until the start of FACS.
449
Fluorescence-activated cell sorting (FACS)
The sorting was performed at the Zuckerman Institute Flow Cytometry Core using a MoFlo Astrios
Cell Sorter (Beckman Coulter). Event rates were kept between 5000-10000 events per second. The
cell sorter uses a linear array of lasers ordered as 640 nm, 488 nm, 561 nm, 532 nm, 405 nm and
355 nm from top to bottom. For experiments described in this manuscript, 488 nm, 561 nm and
405 nm lasers were used to detect the fluorescence of NeuN, mCherry and DAPI respectively. A
gating control sample was used to set the gates for DAPI, NeuN and mCherry. Dissociated nuclei
were passed through the cell sorter to collect those with high mCherry signals. Bright and Dim
mCherry gates were determined after the first 5000 events of DAPI-positive nuclei. Gating was

set to collect only the sparse and bright population that showed a high mCherry signal, and these
events were collected as the photoactivated nuclei ('Bright' mCherry). Nuclei were collected into
SMART-Seq CDS sorting buffer that contains 1X lysis buffer, SMART-Seq Oligo-dT and RNase
inhibitor. All sorted samples were kept on dry ice as recommended by the SMART-Seq protocol
until the start of first-strand synthesis. To increase the likelihood of collecting nuclei in this ultra-
sparse population, aborted events for 'Bright' mCherry and all events for other positive mCherry
were collected into the blocking buffer (200 ml). This suspension was passed through the sorter
again with the same fluorescence gating following the completion of the first sorting to capture
the bright nuclei.
468
Meso-seq
We followed the previously published Meso-seq protocol[46] with minor modifications. Sorted
nuclei were collected in a lysis buffer following the SMART-Seq mRNA LP (with UMIs) protocol.
Reverse transcription for cDNA was followed by cDNA amplification using 17-18 PCR cycles.
Purified cDNA was prepared for sequencing the library using the SMART-seq Library Preparation
Kit. Libraries were amplified using 14 PCR cycles. The concentration of the final library was
determined using a Qubit3.0 Fluorometer (Invitrogen), and the average DNA fragment size was
determined using a Bioanalyzer (Agilent). Sequencing was performed on a NextSeq 2000
sequencer with P2-100 reagents (Illumina). Libraries were diluted and pooled according to the
recommendation of the sequencing kit.
479
Tissue collection and processing for immunohistochemistry and *in situ* hybridization
Mice were anesthetized with isoflurane and transcardially perfused with 20 mL of ice-cold 0.01M
phosphate base saline (PBS, Sigma) followed by 20 mL ice-cold 4% paraformaldehyde (PFA,
Electron Microscopy Sciences) in PBS. Brains were post-fixed in 4% PFA for 24 hours and then
saturated with a 10%, 20%, and 30% sucrose solution sequentially over 48 hours until they sunk
to the bottom of each successive solution. 30% Sucrose-saturated brains were then embedded in
OCT (Optimal Cutting Temperature Compound, Sakura, cat#4583), frozen, stored overnight at -
80 °C, and sliced transversely at 20 µm thickness with a cryostat. Sections were stored at -80 °C
on slides and used for either immunohistochemical staining or RNAScope *in situ* hybridization.
489
Immunohistochemistry
Sections (20 µm) were washed with PBS for 3x5 min. They were then permeabilized with a PBS
and 0.3% Triton X-100 (Sigma-Aldrich) solution (PBST) for 2x20 min and blocked with a 10%
NDS (Normal Donkey Serum, Jackson ImmunoResearch Labs) in PBST solution for 45 min. The
sections were then incubated with a primary antibody for Calbindin (rabbit anti-Calbindin primary,
1:300, cat#: CB38a, Swant) diluted in PBST for 24 hours at 4 °C. The sections were then brought
to room temperature (RT) and washed with PBS for 3x5 min. Secondary antibodies (Donkey anti-
rabbit Alexa-647, 1:300, Jackson Labs, cat#: 711-605-152) diluted in PBST with a 1:1000 dilution
of DAPI (Sigma-Aldrich) to stain nuclei was then applied to the sections for 1 hour at RT. Sections
were then washed with PBS for 3x5 min, and the sections were coverslipped and sealed with
Fluoromount-G™ (Electron Microscopy Sciences, cat#: 17984-25). Sections were imaged in 3 µm
z-steps using an inverted confocal microscope (A1 HD25, Nikon Instruments Inc; excitation
filters: 405/488/568/647nm).
503
RNAScope *in situ* hybridization

20 µm fixed frozen sections of the frozen tissue block were taken and the RNAScope™ Multiplex
Fluorescent Reagent Kit v2 – User Manual was followed (cat#: 323100). The Calbindin targeting
probe was designed and generated by Advanced Cell Diagnostics Inc (cat#: 428431, Entrez Gene
ID: 12307, GenBank Accession #: NM_009788.4). Slides were sequentially dehydrated using
ethanol solutions of increasing concentrations (50%, 70% and 100%) for 5 min at RT. 5-8 drops
of H_2O_2 were added to each sample, followed by a 10 min incubation at RT. After rinsing with
distilled water, antigen retrieval was performed for 5 min at 99°C. To digest sections, RNAscope
Protease III was applied to the sections for 30 min at 40°C. The Calb1 probe was hybridized for 2
h at 40°C and amplified with AMP1 (30 min), AMP2 (30 min), AMP3 (15 min); each was
incubated at 40°C. The probe was fluorescently tagged with 1:2000 TSA Vivid Fluorophore 650
(PN 323273). Slides were counterstained with DAPI for a nuclear stain to identify viable
Calbindin(-) cells and mounted in ProLong Gold Antifade Mountant. 20 µm sections were imaged
in 3 µm z-steps using an inverted confocal microscope with 20x oil objective (A1 HD25, Nikon
Instruments Inc.).
519
Data analysis
Quantification and Statistical Analysis
All statistical details for comparisons are described in the text. No statistical methods were used
to determine sample sizes. Boxplots show the 25th, 50th (median), and 75th quartile ranges with the
whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles,
respectively. Outliers are defined as values extending beyond the whisker ranges. For comparisons
between two populations with non-normal distributions, the Mann-Whitney U test was used. For
comparisons between gene expression datasets, the Wald test followed by multiple corrections via
the Benjamini and Hochberg method was used as described in PyDESeq2[88].
529
Event detection
Fluorescence GCaMP traces were deconvolved using OASIS for fast nonnegative
deconvolution[89]. As in ref. 90, these putative spike events were filtered at 3 median absolute
deviations (MAD) above the raw trace, using a predetermined signal decay constant of 400 ms.
The binarized signal was used to qualify whether a neuron was active at the respective frame. In
our analysis, we do not claim to uncover true spiking events in these neurons but use deconvolution
for denoising and diminishing Ca^{2+} autocorrelation.
537
Spatial tuning curves
The virtual environment was divided into 100 evenly spaced bins (4 cm), which were then utilized
to bin a histogram of each cell's neuronal activity. Neuronal activity was filtered to include activity
from when the animal was running above 3 cm/s and to exclude activity during the 2-sec
teleportation at the end of the 4-m track. The spatial tuning curves were normalized for the animal's
occupancy and then smoothed with a Gaussian kernel ($\sigma = 12$ cm) to obtain a smoothed activity
estimate.
545
Place/silent cell detection
Place fields were detected by identifying locations in the virtual environment where a neuron was
more active than expected by chance. We circularly shifted each neuron's deconvolved spike trace
and recomputed the smoothed, trial-averaged spatial tuning curve of the shifted trace to generate
a shuffled null tuning curve per cell. We repeated this procedure 1000 times in order to calculate

the 95th percentile of null tuning values at every spatial bin to generate a threshold for a p<0.05
significance curve. Spatial tuning curves that surpassed the null threshold were marked as
candidate place fields, and the place field width was calculated as the total bins where the tuning
curve exceeded the shuffled null tuning curve. To restrict our analysis to neurons with specific
firing fields, we additionally required that place fields have a width greater than 8 cm and less than
one-third of the virtual environment (1.3 m). To ensure that the place field activity was stable, we
also required that all place cells had activity for at least 20 laps.
558
559 If the binarized signal trace for a cell did not have any detected events via OASIS deconvolution,
560 the cell was classified as silent.
561
PAmCherry fluorescence quantification
(Fig. 1C,D)
Red PAmCherry fluorescence was calculated as the tagged nuclei fluorescence versus the
background fluorescence of nearby untagged nuclei for the image taken post-tagging. This was
intentional to control for periodic two-photon imaging at 940 nm that may cause an increase in
fluorescence for every cell in the FOV.
568
Fluorescence intensity distribution analysis
(Fig. 1F, right)
Red PAmCherry tagged nuclei were segmented using Cellpose with manual curation performed
within the Cellpose GUI for the max axial projections of the *ex vivo* z-stack (excitation: 568 nm
nm) and *in vivo* z-stack (1070 nm) to generate masks. The masks were used to segment the 3-D
volumes of each cell and the respective fluorescence profiles were normalized and aligned based
on the peak value. The average and standard error were computed based on the aligned
fluorescence profiles of the cells from the respective *in vivo* and *ex vivo* volumes.
577
Two-photon background excitation fluorescence quantification
[Fig. 1G (2P imaging), fig. S2F (2P imaging)]
Changes in PAmCherry red fluorescence due to two-photon fluorescence excitation at 940 nm
from functional recordings were examined. By taking the average fluorescence of the pre-imaging
red channel image and post-imaging red channel image detected at 1070 nm, ΔF/F was computed
using the change in fluorescence between the average post-imaging red image and pre-imaging
red image divided by the pre-imaging red image average [(post – pre) / pre].
585
Tagged and Off-target Fluorescence Change Quantification
[Fig. 1G (target, off-target), fig. S2F (target, off-target)]
Tagged nuclei were segmented using Cellpose with manual curation performed within the
Cellpose GUI for the max axial projection of the *in vivo* -stacks from each mouse to generate
masks. These masks were used to segment a 3D volume for each nuclei that was targeted across
all mice. The number of off-target nuclei was manually quantified laterally and axially per targeted
nuclei.
593
594 Masks for off-target nuclei were hand-drawn on the max axial projection to exclude the targeted
595 nuclei and were at most 2 nuclei bodies away. Experimental background fluorescence masks were

drawn on the surrounding areas with successfully tagged cells and excluded all targeted and off-target nuclei.

The fluorescence values for all tagged and off-target nuclei were percentile-filtered to exclude the lower 10% of fluorescence values to account for vignetting effects and mask inhomogeneities over the max axial projection. The percentile filtered fluorescence values for tagged and off-target nuclei were averaged, and the $\Delta F/F$ was computed by taking the difference between the average tagged or off-target fluorescence value and the average background fluorescence and then normalizing by the average background fluorescence [tagged: (tagged – background)/background; off-target: (off-target – background)/ background]. Note this background value is distinct from the one described in the two-photon background excitation fluorescence quantification.

GCaMP quantifications
(fig. S1AB, fig. S2I)
Frequency: The total number of deconvolved events by OASIS was normalized by the total duration of the recording.

For each cell, the average transient was segmented within a 15-second time window and computed by averaging along aligned deconvolved spike times. If multiple detected events were within 50 frames, the events were treated as a single transient. A cell was only used if there were at least 3 detected transients within the total trace to exclude cells without obvious GCaMP-Ca^{2+} dynamics. For each average transient, we computed the median value based on 5 seconds pre peak transient and the range 5-10 seconds after the peak transient (given that the average transient took significantly less than 5 seconds to resolve) to act as the baseline.

Amplitude: The difference between the max value of the average transient and the baseline value was computed
Half-Rise Time: The time between the half-max value prior to the max and the max value of each transient was computed. We excluded cells where the average half-max value was not observed prior to the transient and performed 99th percentile filtering to remove extreme outliers. Due to sampling rate limitations, we cannot comment on the true half-rise time, so these are approximations.
Half-Decay Time: We computed the time between the max and the half-max value following the transient peak. To remove extreme outliers, we performed 99th percentile filtering.

Image denoising
To correct for vertical scanning artifacts, we utilized combined wavelet and Fourier filters described in ref. 91 [github: https://github.com/DHI-GRAS/rmstripes]. Symlet 20 wavelets were used with varying levels of decomposition (2-5) for discrete wavelet transform to perform vertical striping correction in static images.

***In vivo* and *ex vivo* image registration**
The *in vivo* and *ex vivo* images were transformed into 3D volumetric images for registration. The *in vivo* sequential images were concatenated across the z direction, i.e., depth, stacking the 2D images into a volumetric representation using MATLAB. The *ex vivo* images composed of 2D

slices in each section were concatenated into 3D volumetric data using a stitching algorithm
developed as a precursor for automatic *ex vivo* and *in vivo* registration[81]
[github: https://github.com/ShuonanChen/multimodal_image_registration]. The discontinuity
between *ex vivo* sections results in an unknown spatial correlation between them, requiring
registration between sections. Common cells between sections (i.e., sections one and two) were
used as reference markers for registration. The common cells were manually selected using a
Napari GUI in Python and were utilized as reference markers to inform the scaling and affine
transformations to be applied The scaling and affine transformations were run automatically,
transforming the second section to align with the first section. In the scaling transformation, the
relative distances between cells in a section were compared to the cells in the first section, inducing
an enlargement or shrinkage of the second section to match the first. In the affine transformation,
the second section was geometrically transformed to align correctly with the first section. The
transformations were obtained and applied to each slice in each section, and then each slice was
concatenated together across the z direction to form a volumetric image.
655
The *in vivo* and *ex vivo* 3D images were then adjusted to have a uniform pixel size in all dimensions
(1 μm in x,y and z), ensuring matching FOVs, and equivalent resolution across both images. Time-
averaged representations of *in vivo* and *ex vivo* volumetric stacks were attained by employing
maximum intensity projection (MIP) representations in FIJI, compressing the stacks into 2D
images[92]. The *in vivo* and *ex vivo* registration was carried out using a non-rigid registration
algorithm for the (i) 3D volumetric stacks and (ii) MIP (2D) images. All cells common to both *in*
vivo and *ex vivo* images were manually selected as the centroid of each cell using a Napari GUI in
Python. The common cells were utilized as features to inform the scaling, affine transformation,
and deformation transformation, which were applied to the *ex vivo* image. The scaling and affine
transformations were run automatically. In the scaling transformation, the distances between the
cells in the *ex vivo* image and matching cells in *in vivo* induce enlargement or shrinking of the *ex*
vivo image to match the *in vivo* image. In the affine transformation, the *ex vivo* image was
geometrically transformed to align with the *in vivo* image using the matching cells. The images in
the GUI are updated to reflect the changes induced by the scaling and affine transformation. The
deformation transformation uses a vector field, smoothed with Gaussian filtering, to move cells
and deform the image, ensuring features in the transformed *ex vivo* image align with the *in vivo*
image. The deformation transformation was iteratively employed, beginning with a Gaussian
kernel size of 100, reducing to a kernel size of less than 10, with the user inspecting the alignment
and making manual adjustments to the cell centroid position in the GUI. The completed
transformed *ex vivo* image was then overlaid with the *in vivo* image.
676
RNA-sequencing data analysis
Sequencing data from the Illumina Sequencer was first post-processed through the Illumina
DRAGEN secondary analysis pipeline to de-multiplex based on a unique index for each sample.
RNA-seq reads were aligned to the mouse genome (mm39) using STAR[93]. Unique reads were
counted using HTSeq[94]. HTSeq generated reads were then analyzed for differential expression
using PyDESeq2[88]. Following HTSeq counts, any genes with expression in less than 3 samples
were discarded. FDR adjusted p values were used to determine significantly different genes. Both
'place' and 'silent' cell samples were analyzed against randomly tagged, function-blind, and sex-
matched samples generated from mouse CA1 tissue in the same way as described in this methods
section above. The same comparisons were made to identify differentially expressed genes for

'place' versus 'random', and 'silent' versus 'random'. Within this 'random' dataset, male-female
samples were compared to identify differentially expressed genes influenced by sex. All DEGs
from these comparisons were cross-referenced to find common hits.
690
Gene set enrichment analysis
692 Gene set enrichment analysis was performed on HTSeq-generated counts using the gseapy
693 package in Python 3.11. Enrichment of the fifty hallmark pathways from the molecular signatures
694 database for *Mus musculus* (version 2023.2) was compared in place and silent cells. Comparisons
695 were done with t-test and 1000 permutations.
696

Acknowledgments
We thank Ira Schieren and Max Wallach (Columbia, Zuckerman Institute, ZI) and Dr. Efrat Hagai
(Flow Cytometry Unit, Weizmann Institute of Science) for help with FACS, Columbia University
Herbert Irvine Comprehensive Cancer Center Molecular Pathology Shared Resources for help with
the Bioanalyzer, the Bendesky lab in ZI and Dr. Hadas Keren-Shaul, Revital Ronen at the
Weizmann, Nancy and Stephen Grand Israel National Center (G-INCPM) for help with RNA-
sequencing, George Zakka (Losonczy Lab) for technical support, Erica Rodriguez (Salzman lab)
in ZI for help with tissue processing, the Lomvardas lab in ZI for sequencing advice and help, the
Gogos lab in ZI for molecular bench space, the Polleux lab in ZI for help with cell counter and
confocal microscopy. We thank Franck Polleux, Joseph Gogos, Steve Siegelbaum, and Darcy
Peterka (Columbia, ZI) for feedback on the manuscript.
708

Funding
SAH is supported by the Burroughs Wellcome Fund. DK is supported by a fellowship from the
Israel Ministry of Absorption (IMOAb) and by the Horowitz Foundation. AL is supported by
NIMHR01MH124047, NIMHR01MH124867, NINDSR01NS121106, NINDSU01NS115530,
NINDSR01NS133381, NINDSR01NS131728, NIARF1AG080818. IS is supported by an ISF
personal grant (2354/19) and a BSF US-Israel binational grant (2021281) and he is the incumbent
of the Friends and Linda and Richard Price Career Development Chair and a scholar in the
Zuckerman STEM leadership program.

Author contributions:

Conceptualization: JS, BN, DK, AL, IS
Methodology: JS, BN, DK, BR, SAH, KCK, MECP, HCY
Investigation: JS, BR
Software: BR, JS, TSM, CKO, EV
Formal analysis: JS, BR, TSM, AL
Validation: JS, BN, DK, KCK, BR
Visualization: JS, BR, TSM, SAH, CKO, MECP, KC-KM
Data curation: JS, BR, TSM
Funding acquisition: AL, IS
Resources: AL, IS
Project administration: AL, IS
Supervision: AL, IS
Writing – original draft: JS, BN, TSM, AL, IS
Writing – review & editing: all authors

Competing interests: Authors declare that they have no competing interests.
734
735   **Data and material availability** All data, code, and materials used to generate figures and perform
736   statistical tests are available at Zenodo or NWB upon publication. RNA-seq data will be made
737   freely available on NCBI GEO upon acceptance of our manuscript. Accession numbers for
738   publicly available reagents are included in supplementary materials and methods.

References

1   Zeng, H. What is a cell type and how to define it? *Cell* **185**, 2739-2755 (2022).
https://doi.org:10.1016/j.cell.2022.06.031
2   Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using
Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015). https://doi.org:10.1016/j.cell.2015.05.002
3   Tasic, B. *et al.* Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat*
Neurosci **19**, 335-346 (2016). https://doi.org:10.1038/nn.4216
4   Zeisel, A. *et al.* Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-
cell RNA-seq. *Science* **347**, 1138-1142 (2015). https://doi.org:10.1126/science.aaa1934
5   Harris, K. D. *et al.* Classes and continua of hippocampal CA1 inhibitory neurons revealed by single-
cell transcriptomics. *PLoS Biol* **16**, e2006387 (2018). https://doi.org:10.1371/journal.pbio.2006387
6   Cadwell, C. R. *et al.* Electrophysiological, transcriptomic and morphologic profiling of single
neurons using Patch-seq. *Nat. Biotechnol.* **34**, 199-203 (2016).
7   Cossell, L. *et al.* Functional organization of excitatory synaptic strength in primary visual cortex.
Nature **518**, 399-403 (2015). https://doi.org:10.1038/nature14182
8   Ko, H. *et al.* Functional specificity of local synaptic connections in neocortical networks. *Nature*
473, 87-91 (2011). https://doi.org:10.1038/nature09880
9   Scala, F. *et al.* Phenotypic variation of transcriptomic cell types in mouse motor cortex. *Nature*
(2020). https://doi.org:10.1038/s41586-020-2907-3
10   Bock, D. D. *et al.* Network anatomy and in vivo physiology of visual cortical neurons. *Nature* **471**,
177-182 (2011). https://doi.org:10.1038/nature09802
11   Taniguchi, H. *et al.* A resource of Cre driver lines for genetic targeting of GABAergic neurons in
cerebral cortex. *Neuron* **71**, 995-1013 (2011). https://doi.org:10.1016/j.neuron.2011.07.026
12   Vormstein-Schneider, D. *et al.* Viral manipulation of functionally distinct interneurons in mice,
non-human   primates   and   humans.   *Nat   Neurosci*   **23**,   1629-1636   (2020).
https://doi.org:10.1038/s41593-020-0692-9
13   O'Toole, S. M., Oyibo, H. K. & Keller, G. B. Molecularly targetable cell types in mouse visual
cortex have distinguishable prediction error responses. *Neuron* **111**, 2918-2928 e2918 (2023).
https://doi.org:10.1016/j.neuron.2023.08.015
14   Bugeon, S. *et al.* A transcriptomic axis predicts state modulation of cortical interneurons. *Nature*
607, 330-338 (2022). https://doi.org:10.1038/s41586-022-04915-7
15   Condylis, C. *et al.* Dense functional and molecular readout of a circuit hub in sensory cortex.
Science **375**, eabl5981 (2022). https://doi.org:10.1126/science.abl5981
16   Geiller, T. *et al.* Large-Scale 3D Two-Photon Imaging of Molecularly Identified CA1 Interneuron
Dynamics   in   Behaving   Mice.   *Neuron*   **108**,   968-983   e969   (2020).
https://doi.org:10.1016/j.neuron.2020.09.013
17   Vancura, B., Geiller, T., Grosmark, A., Zhao, V. & Losonczy, A. Inhibitory control of sharp-wave
ripple duration during learning in hippocampal recurrent networks. *Nat Neurosci* (2023).
https://doi.org:10.1038/s41593-023-01306-7
18   Cohen-Kashi Malina, K. *et al.* NDNF interneurons in layer 1 gain-modulate whole cortical columns
according   to   an   animal's   behavioral   state.   *Neuron*   **109**,   2150-2164.e2155   (2021).
https://doi.org:10.1016/j.neuron.2021.05.001
19   Hansel, D. & van Vreeswijk, C. The mechanism of orientation selectivity in primary visual cortex
without   a   functional   map.   *J   Neurosci*   **32**,   4049-4064   (2012).
https://doi.org:10.1523/JNEUROSCI.6284-11.2012
20   Hirokawa, J., Vaughan, A., Masset, P., Ott, T. & Kepecs, A. Frontal cortex neuron types
categorically   encode   single   decision   variables.   *Nature*   **576**,   446-451   (2019).
https://doi.org:10.1038/s41586-019-1816-9

21 Mancinelli, S. & Lodato, S. Decoding neuronal diversity in the developing cerebral cortex: from
single cells to functional networks. *Curr Opin Neurobiol* **53**, 146-155 (2018).
https://doi.org:10.1016/j.conb.2018.08.001
22 Ohki, K. & Reid, R. C. Specificity and randomness in the visual cortex. *Curr Opin Neurobiol* **17**,
401-407 (2007). https://doi.org:10.1016/j.conb.2007.07.007
23 Peng, H. *et al.* Morphological diversity of single neurons in molecularly defined cell types. *Nature*
598, 174-181 (2021). https://doi.org:10.1038/s41586-021-03941-1
24 Treves, A. & Rolls, E. T. Computational analysis of the role of the hippocampus in memory.
Hippocampus **4**, 374-391 (1994).
25 Buzsaki, G. & Mizuseki, K. The log-dynamic brain: how skewed distributions affect network
operations. *Nat Rev Neurosci* **15**, 264-278 (2014). https://doi.org:10.1038/nrn3687
26 Cembrowski, M. S. *et al.* Spatial Gene-Expression Gradients Underlie Prominent Heterogeneity of
CA1 Pyramidal Neurons. *Neuron* (2016). https://doi.org:10.1016/j.neuron.2015.12.013
27 Mallory, C. S. & Giocomo, L. M. Heterogeneity in hippocampal place coding. *Curr Opin Neurobiol*
49, 158-167 (2018). https://doi.org:10.1016/j.conb.2018.02.014
28 Pettit, N. L., Yap, E. L., Greenberg, M. E. & Harvey, C. D. Fos ensembles encode and shape stable
spatial maps in the hippocampus. *Nature* **609**, 327-334 (2022). https://doi.org:10.1038/s41586-022-
05113-1
29 Soltesz, I. & Losonczy, A. CA1 pyramidal cell diversity enabling parallel information processing
in the hippocampus. *Nat Neurosci* **21**, 484-493 (2018). https://doi.org:10.1038/s41593-018-0118-
0
30 Sosa, M., Gillespie, A. K. & Frank, L. M. Neural Activity Patterns Underlying Spatial Coding in
the Hippocampus. *Curr Top Behav Neurosci* (2016). https://doi.org:10.1007/7854_2016_462
31 Tanaka, K. Z. *et al.* The hippocampal engram maps experience but not place. *Science* **361**, 392-397
(2018). https://doi.org:10.1126/science.aat5397
32 Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nature neuroscience* **16**, 264-
266 (2013). https://doi.org:10.1038/nn.3329
33 Lee, J. S., Briguglio, J. J., Cohen, J. D., Romani, S. & Lee, A. K. The Statistical Structure of the
Hippocampal Code for Space as a Function of Time, Context, and Value. *Cell* **183**, 620-635 e622
(2020). https://doi.org:10.1016/j.cell.2020.09.024
34 Kentros, C. G., Agnihotri, N. T., Streater, S., Hawkins, R. D. & Kandel, E. R. Increased attention
to spatial context increases both place field stability and spatial memory. *Neuron* **42**, 283-295
(2004).
35 Grosmark, A. D., Sparks, F. T., Davis, M. J. & Losonczy, A. Reactivation predicts the consolidation
of unbiased long-term cognitive maps. *Nat Neurosci* **24**, 1574-1585 (2021).
https://doi.org:10.1038/s41593-021-00920-7
36 Keinath, A. T., Mosser, C. A. & Brandon, M. P. The representation of context in mouse
hippocampus is preserved despite neural drift. *Nat Commun* **13**, 2415 (2022).
https://doi.org:10.1038/s41467-022-30198-7
37 Colgin, L. L., Moser, E. I. & Moser, M.-B. Understanding memory through hippocampal
remapping. *Trends in neurosciences* **31**, 469-477 (2008). https://doi.org:10.1016/j.tins.2008.06.008
38 Moeyaert, B. *et al.* Improved methods for marking active neuron populations. *Nat. Commun.* **9**,
4440 (2018). https://doi.org:10.1038/s41467-018-06935-2
39 DeNardo, L. A. *et al.* Temporal evolution of cortical ensembles promoting remote memory
retrieval. *Nat Neurosci* **22**, 460-469 (2019). https://doi.org:10.1038/s41593-018-0318-7
40 Guenthner, C. J., Miyamichi, K., Yang, H. H., Heller, H. C. & Luo, L. Permanent genetic access
to transiently active neurons via TRAP: targeted recombination in active populations. *Neuron* **78**,
773-784 (2013). https://doi.org:10.1016/j.neuron.2013.03.025
41 Lee, D., Hyun, J. H., Jung, K., Hannan, P. & Kwon, H.-B. A calcium- and light-gated switch to
induce gene expression in activated neurons. *Nat. Biotechnol.* **35**, 858-863 (2017).

42 Sørensen, A. T. *et al.* A robust activity marking system for exploring active neuronal ensembles.
Elife **5**, e13918 (2016). https://doi.org:10.7554/eLife.13918

43 Datta, S. R. *et al.* The Drosophila pheromone cVA activates a sexually dimorphic neural circuit.
Nature **452**, 473-477 (2008). https://doi.org:10.1038/nature06808

44 Lien, A. D. & Scanziani, M. In vivo Labeling of Constellations of Functionally Identified Neurons
for Targeted in vitro Recordings. *Front Neural Circuits* **5**, 16 (2011).
https://doi.org:10.3389/fncir.2011.00016

45 Lee, D., Kume, M. & Holy, T. E. Sensory coding mechanisms revealed by optical tagging of
physiologically defined neuronal types. *Science* **366**, 1384-1389 (2019).
https://doi.org:10.1126/science.aax8055

46 Apelblat, D., Roethler, O., Bitan, L., Keren-Shaul, H. & Spiegel, I. Meso-seq for in-depth
transcriptomics in ultra-low amounts of FACS-purified neuronal nuclei. *Cell Rep Methods* **2**,
100259 (2022). https://doi.org:10.1016/j.crmeth.2022.100259

47 Dana, H. *et al.* High-performance calcium sensors for imaging activity in neuronal populations and
microcompartments. *Nat Methods* **16**, 649-657 (2019). https://doi.org:10.1038/s41592-019-0435-
6

48 Scheyltjens, I. *et al.* Evaluation of the expression pattern of rAAV2/1, 2/5, 2/7, 2/8, and 2/9
serotypes with different promoters in the mouse visual cortex. *J Comp Neurol* **523**, 2019-2042
(2015). https://doi.org:10.1002/cne.23819

49 Geiller, T., Priestley, J. B. & Losonczy, A. A local circuit-basis for spatial navigation and memory
processes in hippocampal area CA1. *Curr Opin Neurobiol* **79**, 102701 (2023).
https://doi.org:10.1016/j.conb.2023.102701

50 Katona, G. *et al.* Fast two-photon in vivo imaging with three-dimensional random-access scanning
in large tissue volumes. *Nat Methods* **9**, 201-208 (2012). https://doi.org:10.1038/nmeth.1851

51 O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit
activity in the freely-moving rat. *Brain Research* **34**, 171-175 (1971).

52 Witharana, W. K. *et al.* Nonuniform allocation of hippocampal neurons to place fields across all
hippocampal subfields. *Hippocampus* **26**, 1328-1344 (2016). https://doi.org:10.1002/hipo.22609

53 Rich, P. D., Liaw, H. P. & Lee, A. K. Place cells. Large environments reveal the statistical structure
governing hippocampal representations. *Science* **345**, 814-817 (2014).
https://doi.org:10.1126/science.1255635

54 Thompson, L. T. & Best, P. J. Place cells and silent cells in the hippocampus of freely-behaving
rats. *J Neurosci* **9**, 2382-2390 (1989). https://doi.org:10.1523/JNEUROSCI.09-07-02382.1989

55 Wilson, M. A. & McNaughton, B. L. Dynamics of the hippocampal ensemble code for space.
Science **261**, 1055-1058 (1993).

56 Zaremba, J. D. *et al.* Impaired hippocampal place cell dynamics in a mouse model of the 22q11.2
deletion. *Nat Neurosci* **20**, 1612-1623 (2017). https://doi.org:10.1038/nn.4634

57 Bowler, J. C. & Losonczy, A. Direct cortical inputs to hippocampal area CA1 transmit
complementary signals for goal-directed navigation. *Neuron* **111**, 4071-4085 e4076 (2023).
https://doi.org:10.1016/j.neuron.2023.09.013

58 Bowler, J. C. *et al.* behaviorMate: An Intranet of Things Approach for Adaptable Control of
Behavioral and Navigation-Based Experiments. *bioRxiv* (2023).
https://doi.org:10.1101/2023.12.04.569989

59 Priestley, J. B., Bowler, J. C., Rolotti, S. V., Fusi, S. & Losonczy, A. Signatures of rapid plasticity
in hippocampal CA1 representations during novel experiences. *Neuron* (2022).
https://doi.org:10.1016/j.neuron.2022.03.026

60 Cembrowski, M. S., Wang, L., Sugino, K., Shields, B. C. & Spruston, N. Hipposeq: a
comprehensive RNA-seq database of gene expression in hippocampal principal neurons. *Elife* **5**,
e14997 (2016). https://doi.org:10.7554/eLife.14997

61   Bloodgood, B. L., Sharma, N., Browne, H. A., Trepman, A. Z. & Greenberg, M. E. The activity-
dependent transcription factor NPAS4 regulates domain-specific inhibition. *Nature* **503**, 121-125
(2013). https://doi.org:10.1038/nature12743
62   Sheng, M. & Greenberg, M. E. The regulation and function of c-fos and other immediate early
genes in the nervous system. *Neuron* **4**, 477-485 (1990). https://doi.org:10.1016/0896-
6273(90)90106-p
63   Spiegel, I. *et al.* Npas4 regulates excitatory-inhibitory balance within neural circuits through cell-
type-specific gene programs. *Cell* **157**, 1216-1229 (2014).
https://doi.org:10.1016/j.cell.2014.03.058
64   Elmore, S. Apoptosis: a review of programmed cell death. *Toxicol Pathol* **35**, 495-516 (2007).
https://doi.org:10.1080/01926230701320337
65   Xu, J., Burgoyne, P. S. & Arnold, A. P. Sex differences in sex chromosome gene expression in
mouse brain. *Hum Mol Genet* **11**, 1409-1419 (2002). https://doi.org:10.1093/hmg/11.12.1409
66   Epsztein, J., Brecht, M. & Lee, A. K. Intracellular determinants of hippocampal CA1 place and
silent cell activity in a novel environment. *Neuron* **70**, 109-120 (2011).
https://doi.org:10.1016/j.neuron.2011.03.006
67   Mizuseki, K. & Buzsaki, G. Preconfigured, skewed distribution of firing rates in the hippocampus
and entorhinal cortex. *Cell Rep* **4**, 1010-1021 (2013). https://doi.org:10.1016/j.celrep.2013.07.039
68   Alme, C. B. *et al.* Place cells in the hippocampus: eleven maps for eleven rooms. *Proc Natl Acad
Sci U S A* **111**, 18428-18435 (2014). https://doi.org:10.1073/pnas.1421056111
69   Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by hippocampal cellular
assemblies. *Nature* **469**, 397-401 (2011). https://doi.org:10.1038/nature09633
70   Dragoi, G. & Tonegawa, S. Selection of preconfigured cell assemblies for representation of novel
spatial experiences. *Philos Trans R Soc Lond B Biol Sci* **369**, 20120522 (2014).
https://doi.org:10.1098/rstb.2012.0522
71   Grosmark, A. D. & Buzsaki, G. Diversity in neural firing dynamics supports both rigid and learned
hippocampal sequences. *Science* **351**, 1440-1443 (2016). https://doi.org:10.1126/science.aad1935
72   Huszar, R., Zhang, Y., Blockus, H. & Buzsaki, G. Preconfigured dynamics in the hippocampus are
guided by embryonic birthdate and rate of neurogenesis. *Nat Neurosci* **25**, 1201-1212 (2022).
https://doi.org:10.1038/s41593-022-01138-x
73   Geiller, T. *et al.* Local circuit amplification of spatial selectivity in the hippocampus. *Nature* **601**,
105-109 (2022). https://doi.org:10.1038/s41586-021-04169-9
74   Solstad, T., Boccara, C. N., Kropff, E., Moser, M. B. & Moser, E. I. Representation of geometric
borders in the entorhinal cortex. *Science* **322**, 1865-1868 (2008).
https://doi.org:10.1126/science.1166466
75   Taube, J. S., Muller, R. U. & Ranck, J. B., Jr. Head-direction cells recorded from the postsubiculum
in freely moving rats. I. Description and quantitative analysis. *J Neurosci* **10**, 420-435 (1990).
https://doi.org:10.1523/JNEUROSCI.10-02-00420.1990
76   Eichenbaum, H. Time cells in the hippocampus: a new dimension for mapping memories. *Nat Rev
Neurosci* **15**, 732-744 (2014). https://doi.org:10.1038/nrn3827
77   Driscoll, L. N., Duncker, L. & Harvey, C. D. Representational drift: Emerging theories for
continual learning and experimental future directions. *Curr Opin Neurobiol* **76**, 102609 (2022).
https://doi.org:10.1016/j.conb.2022.102609
78   Josselyn, S. A. & Tonegawa, S. Memory engrams: Recalling the past and imagining the future.
Science **367** (2020). https://doi.org:10.1126/science.aaw4325
79   Xu, S. *et al.* Behavioral state coding by molecularly defined paraventricular hypothalamic cell type
ensembles. *Science* **370** (2020). https://doi.org:10.1126/science.abb2494
80   Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. RNA imaging. Spatially
resolved, highly multiplexed RNA profiling in single cells. *Science* **348**, aaa6090 (2015).
https://doi.org:10.1126/science.aaa6090

Chen, S. *et al.* Multimodal Microscopy Image Alignment Using Spatial and Shape Information and a Branch-and-Bound Algorithm. *Proc IEEE Int Conf Acoust Speech Signal Process* **2023** (2023). https://doi.org/10.1109/icassp49357.2023.10096185

Rao, B. Y. *et al.* Non-parametric Vignetting Correction for Sparse Spatial Transcriptomics Images. *Med Image Comput Comput Assist Interv* **12908**, 466-475 (2021). https://doi.org/10.1007/978-3-030-87237-3_45

Bittner, K. C., Milstein, A. D., Grienberger, C., Romani, S. & Magee, J. C. Behavioral time scale synaptic plasticity underlies CA1 place fields. *Science* **357**, 1033-1036 (2017). https://doi.org/10.1126/science.aan3846

Magee, J. C. & Grienberger, C. Synaptic Plasticity Forms and Functions. *Annu Rev Neurosci* (2020). https://doi.org/10.1146/annurev-neuro-090919-022842

Kaifosh, P., Zaremba, J. D., Danielson, N. B. & Losonczy, A. SIMA: Python software for analysis of dynamic fluorescence imaging data. *Front Neuroinform* **8**, 80 (2014). https://doi.org/10.3389/fninf.2014.00080

Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *bioRxiv*, 061507 (2016). https://doi.org/10.1101/061507

Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for cellular segmentation. *Nat Methods* **18**, 100-106 (2021). https://doi.org/10.1038/s41592-020-01018-x

Muzellec, B., Teleńczuk, M., Cabeli, V. & Andreux, M. PyDESeq2: a python package for bulk RNA-seq differential expression analysis. *bioRxiv*, 2022.2012.2014.520412 (2022). https://doi.org/10.1101/2022.12.14.520412

Friedrich, J., Zhou, P. & Paninski, L. Fast online deconvolution of calcium imaging data. *PLoS Comput Biol* **13**, e1005423 (2017). https://doi.org/10.1371/journal.pcbi.1005423

O'Hare, J. K. *et al.* Compartment-specific tuning of dendritic feature selectivity by intracellular Ca(2+) release. *Science* **375**, eabm1670 (2022). https://doi.org/10.1126/science.abm1670

Münch, B., Trtik, P., Marone, F. & Stampanoni, M. Stripe and ring artifact removal with combined wavelet — Fourier filtering. *Opt. Express* **17**, 8567-8591 (2009). https://doi.org/10.1364/OE.17.008567

Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012). https://doi.org/10.1038/nmeth.2019

Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013). https://doi.org/10.1093/bioinformatics/bts635

Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015). https://doi.org/10.1093/bioinformatics/btu638

Supplementary Figures

Fig. S1. Additional data on 2P-NucTag. (**A**) Representative average GCaMP-Ca^{2+} transients from nine CA1 PNs,
vertical scale bar (50% ΔF/F). (**B**) Left: GCaMP amplitude (54.6% ± 0.3% ΔF/F, n = 8190 cells in 9 mice). Middle: GCaMP half
rise time (0.15s ± 0.001s, n = 7797 cells in 9 mice), Right: GCaMP half decay time (0.44s ± 0.003s, n = 8282 cells in 9 mice, 940
nm excitation, See methods for cell exclusion criterion). Boxplots show the 25th, 50th (median), and 75th quartile ranges,
with the whiskers extending to 1.5 interquartile ranges below or above the 25th or 75th quartiles, respectively. Outliers
are defined as values extending beyond the whisker ranges. (**C**) Left: *in vivo* 3D visualization of the entire field of
view (FOV). Middle: subset of *in vivo* 3D visualization. Right: representative cell from *in vivo* 3D visualization. (**D**)
Left: *ex vivo* confocal 3D visualization of entire FOV. Middle: subset of *ex vivo* 3D visualization. Right: representative
cell from *ex vivo* 3D visualization. *C* and *D* are corresponding to *Figure 1F*.

Fig. S2. *In vivo* 2P-NucTag of silent cells. (**A**) Pipeline for two-photon (*2P*) phototagging of functionally identified
'silent' neurons in CA1 during VR spatial navigation (as in *Fig. 1B*). (**B**) Example 2P imaging field of view (FOV) of
GCaMP in the CA1 pyramidal layer from a mouse with "silent" cells targeted. Scale bar: 100 μm. (**C**) CA1PNs
detected with Suite2p/Cellpose in the FOV shown in *B*. (**D**) Spatial mask (blue) of identified 'silent' cells from the
FOV in *B*. Scale bar: 100 μm. (**E**) Left: PAmCherry fluorescence (magenta) of tagged nuclei after 2P phototagging.
Right: overlay of spatial masks of identified CA1PNs and tagged nuclei for the FOVs in *D* and *E*. Note that we only
tagged a subset of silent cells present in the FOV. Scale bar: 100 μm. (**F**) Left: Proportion of single, double, and triple-
tagged nuclei following phototagging of a single silent cell. Right: relative PAmCherry fluorescence change for non-
tagged cells in the FOV after 2P imaging (green), after 2P phototagging of targeted silent cell nuclei (orange) and off-
target nuclei (gray, n = 4 mice). (**G**) Average velocity of the mice during virtual reality navigation task (Mann-Whitney
U test, p-value = 0.286). (**H**) Left: deconvolved events per minute from all cells across all mice from 2P GCaMP-

Ca^{2+}imaging (averaged across mice, n = 5 'Place' mice, n = 4 'Silent' mice, Mann-Whitney U Test, p-value = 0.28).
Right: deconvolved events per minute from place cells across all mice (averaged across mice, place n = 5, silent n =
4, Mann-Whitney U Test, p-value = 0.90). (**I**) Left: GCaMP transient amplitude of all cells between groups (averaged
across mice, n = 5 'Place' mice, n = 4 'Silent' mice, Mann-Whitney U Test, p-value = 0.14), Middle: GCaMP half
rise time of all cells between groups (averaged across mice, n = 5 'Place' mice, n = 4 'Silent' mice, Mann-Whitney U
Test, p-value = 0.14), Right: GCaMP half decay time of all cells between groups (averaged across mice, n = 5 'Place'
mice, n = 4 'Silent' mice, Mann-Whitney U Test, p-value = 0.81). Boxplots show the 25[th], 50[th] (median), and 75[th]
quartile ranges, with the whiskers extending to 1.5 interquartile ranges below or above the 25[th] or 75[th] quartiles,
respectively. Outliers are defined as values extending beyond the whisker ranges.

Fig. S3. Additional data on phototagging. From left to right by column: animal ID (animals in *Figure 2* and *figure*
S2 are not shown here), *in vivo* two-photon (*2P*) imaging fields of view (FOVs), functionally defined masks (orange
for place cells and blue for silent cells), PAmCherry fluorescence (magenta) of tagged nuclei after 2P phototagging
and overlay of spatial masks from identified CA1PNs and tagged nuclei for the respective FOV in the same row. Note
that we only tagged a subset of silent cells present in the FOV in 'Silent' mice, in order to approximate the number of
phototagged place cells in 'Place' mice. Scale bar: 100 μm.

Fig. S4. Additional data on transcriptomics analysis of place and silent cells. (**A**) Sequencing statistics for all
'place' and 'silent' cell samples. Total number of reads - 40 to 60 million reads, 54.64 ± 2.09, n = 9. Percent mapped
by STAR - 80.58 to 86.92%. 84.33 ± 0.73. Number of unique genes - 14176 to 20990, 17486 ± 714. (**B**) Normalized
counts for groups of genes plotted for 'place' versus 'silent'. Here we show that gene expression of apoptotic genes,
superficial CA1 genes, deep CA1 genes, housekeeping genes, proximal CA1 genes, and distal CA1 genes are not
different between the two groups (FDR adjusted p-value. *<0.05, **<0.001, ***<0.001, PyDeSeq2. All comparisons
in this figure are not significant).

Fig. S5. Additional data on transcriptomics analysis of place and silent cells. (**A**) Top: MA plot of 'place' versus 'random'. Differentially expressed genes (DEGs) are labeled in orange. Bottom: MA plot of 'silent' versus 'random'. DEGs are labeled in blue. Both: DEGs that are common for 'place' versus 'random' were highlighted and labeled. (**B**) Normalized counts for 4 example genes that are significantly differentially expressed across comparisons (FDR adjusted p-value. *<0.05, **<0.001, ***<0.001, PyDeSeq2. Showing here a comparison of 'place' versus 'random' or 'silent' versus 'random'. 'Place' versus 'silent' comparisons were shown in Fig. 3). (**C**) MA plot of male versus female for the 'random' dataset. Top: Y-linked genes that are differentially expressed between sex are highlighted and labeled. Bottom: same for genes in panel A and B are highlighted and labeled. They are not differentially expressed between sex (FDR adjusted p-value. *<0.05, **<0.001, ***<0.001, PyDeSeq2. Otherwise, comparisons are not significant). (**D**) Normalized counts for 3 example DEGs between male and female.

1022 **Fig. S6. *Post hoc* immunohistochemistry and *in situ* hybridization on tagged tissue.** Top and middle: confocal
1023 horizontal images of *post hoc* calbindin immunohistochemistry on tissue with tagged cells in the hippocampus. Scale
1024 bar: 50 μm (top), 20 μm (middle, higher magnification). Bottom: confocal horizontal images of tagged tissue
1025 hybridized with an RNAScope probe for Calb1, highlighting tagged cells with positive Calb1 signal. Scale bar: 20
1026 μm.

Table 1

Mouse	Sex	Group	Place #	Silent #	Active Non-Place #	Mask #	Tag #
12-10	Male	Place	119	208	1754	119	112
12-20	Female	Place	250	470	1232	218	168
12-21	Female	Place	89	1040	701	74	73
12-27	Female	Place	132	547	548	119	119
12-37	Female	Place	126	290	1440	119	115
12-9	Male	Silent	9	152	655	152	87
12-12	Male	Silent	28	230	287	230	89
12-31	Male	Silent	129	305	1238	200	97
12-34	Male	Silent	187	303	571	113	79

Supplementary Movie 1. Phototagging
Real-time movie of *in vivo* two-photon imaging and phototagging of neurons. Imaging was performed at 1040 nm to visualize change in PAmCherry fluorescence. During phototagging, the 810-nm laser was scanned over target nuclei and the PMT was blanked. A frame average of 64 frames was applied for resolution and clarity. Final video was edited to include scale bar and laser switches. Video time is not representative of actual recording time.

Supplementary Movie 2. Registered *in vivo* and *ex vivo* image stacks
Three-dimensional rendering of registered z-stacks of *in vivo* and *ex vivo* tissue volume of the CA1 pyramidal layer with phototagged nuclei. Corresponding to *Fig. 1F* and Fig *S1C,D*. Magenta: PAmCherry *in vivo* (captured on Bruker 2P microscope, wavelength = 1070 nm). Orange: PAmCherry *ex vivo* (captured on A1 HD25, Nikon Instruments Inc., wavelength = 568 nm)

Supplementary Table. List of DEGs between place and silent cells
List of DEGs between place and silent cells showing log 2-fold change, mean of normalized count (baseMean), and FDR adjusted p-value. All genes in the table have an FDR-adjusted p-value less than 0.05 when comparing silent versus place cells. log2FoldChange is computed for silent versus place (positive: enriched in silent cells, negative: enriched in place cells).
File: Supplementary_Table.