DNA damage drives antigen diversification through mosaic VSG formation in *Trypanosoma brucei*

Jaclyn E. Smith[1], Kevin J. Wang[1], Jill M.C. Hakim[1], Alexander K. Beaver[1,2], Aishwarya Magesh[1], Shane D. Gilligan-Steinberg[1,3], Jessica Zheng[1], Jaime So[1], Erin M. Kennedy[1], Bailin Zhang[1,4], Elgin Henry Akin[1], Lusajo Mwakibete[1], Monica R. Mugnier[1]*

[1]W. Harry Feinstone Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, United States of America

[2]Department of Pathology, Johns Hopkins School of Medicine, Baltimore, Maryland, United States of America

[3]Department of Bioengineering, University of Washington, Seattle, Washington, United States of America

[4]Scripps Research Department of Integrative Structural and Computational Biology, La Jolla, San Diego, California, United States of America

*Corresponding author
Email: mmugnie1@jhu.edu (MRM)

Summary

Antigenic variation, using large genomic repertoires of antigen-encoding genes, allows pathogens to evade host antibody. Many pathogens, including the African trypanosome *Trypanosoma brucei,* extend their antigenic repertoire through genomic diversification. While evidence suggests that *T. brucei* relies heavily on the generation of new variant surface glycoprotein (VSG) genes to maintain a chronic infection, a lack of experimentally tractable tools for studying this process has obscured its underlying mechanisms. Here, we present a highly sensitive targeted RNA sequencing approach for measuring VSG diversification. Using this method, we demonstrate that a Cas9-induced DNA double-strand break within the VSG coding sequence can induce VSG recombination with patterns identical to those observed during infection. These newly generated VSGs are antigenically distinct from parental clones and thus capable of facilitating immune evasion. Together, these results provide insight into the mechanisms of VSG diversification and an experimental framework for studying the evolution of antigen repertoires in pathogenic microbes.

Introduction

Pathogen survival in a host depends upon effective and continuous immune evasion. Several bacteria and eukaryotic pathogens have adopted strategies of antigenic variation to evade host immunity, a process in which they continuously alter antigenic surface proteins to escape the host's adaptive immune response. The African trypanosome *Trypanosoma brucei*, a unicellular eukaryotic parasite and causative agent of human and animal African trypanosomiasis, uses an especially sophisticated system of antigenic variation. The parasite, which remains extracellular throughout infection and thus faces a perpetual onslaught of host antibody, periodically "switches" expression of a surface coat consisting of 10^7 copies of a single, immunogenic protein known as the Variant Surface Glycoprotein (VSG). This process allows parasites to escape host antibody and maintain a chronic infection.

Although the *T. brucei* VSG repertoire contains thousands of VSGs, it may not be sufficient to maintain a chronic infection through VSG switching alone. During an infection, each *T. brucei* parasite expresses a single VSG at a time from one of ~15 telomeric Bloodstream Expression Sites (the "active" BES)[1]. The remaining thousands of VSG-encoding genes are stored in other expression sites, subtelomeric arrays, and minichromosomes, all of which remain transcriptionally silenced[2]. The parasite switches its VSG either by transcriptional activation of a silent BES (*in situ* switching) or through a gene conversion event in which a new VSG is copied into the active expression site. While gene conversion-based switching allows for the activation of VSGs outside of a BES, analysis of the *T. brucei* genome has shown that only ~20% of the VSGs in the parasite genome are full-length genes encoding a functional VSG protein. The remaining ~80% of VSGs in the parasite genome consist of pseudogenes or gene fragments[2] and cannot immediately be used for immune evasion through *in situ* or gene conversion switching. Moreover, the number of VSGs expressed at a single timepoint during experimental infection sometimes exceeds the total number of intact VSGs in the parasite genome[3,4], further indicating that the repertoire of intact VSGs is insufficient for the antigenic diversity required to maintain a chronic infection.

Evidence suggests that *T. brucei* deals with this shortage of antigens through diversification of the VSG repertoire. Many mouse studies have shown that novel VSGs, generated during infection, predominate at later stages of infection[3,5,6], while analysis of parasites from natural human infections revealed expressed VSGs that were nearly completely absent from the genomes of contemporary field isolates[7]. These observations suggest that the generation of new VSGs is likely to be a central feature of *T. brucei* antigenic variation.

There are two mechanisms thought to be responsible for extending the VSG repertoire: mosaic formation and *de novo* point mutation. Mosaic VSGs form when two or more VSG genes combine through segmental gene conversion to form a novel VSG. This mechanism allows parasites to access pseudogenes and VSG fragments within the repertoire. While mosaic VSGs have been observed in the literature, they are often found under extreme experimental conditions[8,9] or late during infection[3,5,10,11] making it difficult

to discern how exactly they arose. VSGs also appear to occasionally acquire *de novo*
point mutations though these can be difficult to distinguish from a small gene conversion
event using a similar donor VSG. The origins of these mutations also remain mysterious,
though newly acquired mutations seem rare[9,12–14]. Ultimately, *de novo* mutation of VSGs
would allow parasites to generate new VSG sequences regardless of the contents of their
repertoire, further amplifying diversity.
93
Despite its clear importance, the mechanisms driving VSG diversification, whether
by mutation or recombination, remain poorly understood. It is plausible that DNA damage
and repair may play a role in either mechanism, as expressed VSG genes sit between
two highly repetitive and damage-prone stretches of DNA, the conserved 70bp repeat and
the telomere[2,15]. DNA breaks are frequently observed within the 70bp repeats at the active
expression site[16,17] and near telomeres within silent BESs[17]. Experimental evidence
implicates DNA damage in VSG switching more generally, as a DNA double-strand break
induced upstream of the VSG induces a switch[16,17], but it is unknown whether DNA
damage could also generate new VSGs.
103
There is a dearth of experimental evidence for the mechanisms driving VSG
diversification because there has been no controlled and reproducible *in vitro* system for
studying the process. Instead, researchers have had to rely on observation alone,
characterizing the sequences of expressed VSGs that were isolated by chance. This
approach has been crucial for generating hypotheses but is insufficient for deciphering
mechanisms driving the process. Here, we have developed a comprehensive toolkit for
the controlled and reproducible study of the diversification of individual VSGs. Using a
highly sensitive barcode-based targeted RNA-seq approach, we show that DNA double-
strand breaks can trigger the formation of mosaic VSGs that are identical to those
observed *in vivo* during infection. Homology appears to drive donor VSG selection, with
microhomology between the parent and donor VSG flanking the site of recombination in
almost every event. Finally, we observe that break-induced diversification is most efficient
in the portion of the VSG gene encoding the exposed top lobe of the VSG N-terminal
domain. We suggest that this could represent a potential hypervariable region within the
VSG, facilitating diversification in the region of the VSG protein where host antibody is
most likely to bind.

Results

DNA double-strand breaks trigger the formation of mosaic VSGs

A number of studies have suggested that VSG diversification occurs, at least some of the time, within the active expression site[3,13], with the sequence of the actively expressed VSG being altered through recombination and/or mutation. For this reason, we focused our analyses on diversification of the actively expressed VSG. Because a double-strand DNA break upstream of the VSG within the active BES is known to induce switching[16,17], we hypothesized that a break within the VSG coding sequence might result in mosaic VSG formation. To investigate this, we engineered tetracycline-inducible Cas9 expressing EATRO1125 parasites, which express the VSG AnTat1.1, and induced breaks across the AnTat1.1 coding sequence using a set of guide RNAs.

To evaluate potential recombination outcomes, we developed VSG Anchored Multiplex PCR Sequencing (VSG-AMP-seq), a technique that overcomes previous obstacles to studying VSG diversification by providing both high-throughput and highly accurate sequences (1A). Our approach uses long unique molecular indexes (UMIs) to generate high-confidence, accurate consensus sequences[18]. Consolidating reads into consensus

Figure 1. VSG-AMP-seq is a tool to detect diversification within a VSG of interest
A) Schematic of the library prep for VSG-AMP-seq. Figure adapted from [51]. B) Locations of target specific primers for VSG AnTat1.1. SL = 5′ splice leader cap, 14-mer = 3′ sequence conserved in all VSG transcripts

sequences allows errors like PCR chimeras, which occur during later cycles of PCR and therefore represent a minority of sequences in a consensus group[19], to be eliminated while true events are retained. Libraries are prepared by fragmenting VSG-specific cDNA. Fragments are then end-repaired, A-tailed and ligated to universal adapters containing a

25bp unique molecular index (1A). A VSG target of interest is selected, and a series of
staggered primers are designed to cover the length of the target VSG's coding sequence
(1B). By pairing target-specific primers with a universal reverse adapter primer, target
VSG fragments are amplified within a sample regardless of their identity. Mosaic reads
are defined as those for whom a portion of the read matches the target and the remainder
matches another VSG (the "donor VSG") within the VSG repertoire (VSGnome) of the
strain being studied[2]. Due to the selective, target-specific amplification, this method can
sensitively detect thousands of rare diversification events, even from mixed samples
where diversified VSGs are a minority of the population.
175
To induce breaks across AnTat1.1, we induced Cas9 expression for 24 hours and
then transfected in DNA amplicons containing a T7 promoter and a guide RNA targeting
various regions throughout the AnTat1.1 coding sequence (cut positions, relative to the
5′ end of the VSG transcript: 243, 369, 694, 894, 978, and 1459) to induce breaks in the
VSG. Parasites were collected two days after transfection of the guide, and mosaic
derivatives of AnTat1.1 were analyzed by VSG-AMP-Seq (2A; Supplementary Figures
2A, 2B & 3C). We detected thousands of recombination events from two independently
generated Cas9 clones (C1 = 5956, C2 = 4488).
184
Our analysis revealed diverse mosaic recombination events centered around each
break site. Such events were virtually absent from the negative (no guide) control,
suggesting this process does not occur at a high rate in the absence of a trigger or in the
presence of Cas9 alone. Notably, as the DNA breaks progressed further away from the
center of the AnTat1.1 coding sequence, the frequency of mosaic formation decreased
dramatically (2B). This did not appear to be related to the guide sequences (Supplemental
Figure 2C-G) or to guide cutting efficiency (Supplemental Figure 2H). To ensure that the
observed mosaic events represented mosaic VSGs that were truly expressed by the
parasite, and not technical artifacts or VSGs incapable of being stably expressed by *T.*
brucei, we also obtained individual clones of parasites expressing AnTat1.1-derived
mosaic VSGs. Using parasite lines that stably express a VSG-targeted guide RNA after
7 days of Cas9 expression (2C; Supplemental Figure 2A), we isolated 26 mosaic-
expressing clones, 4 from guide 243 and 22 from guide 694, from 6 parental cell lines.
These parasites expressed VSGs containing recombination events identical to those
observed using VSG-AMP-Seq. These results suggest that DNA damage within the active
VSG can trigger the formation of mosaic VSGs, with recombination events centered
around the site of DNA damage.
202
203 **Homology drives mosaic VSG formation**
204
Analysis of the mosaic recombination events detected by VSG-AMP-seq revealed
an important role for sequence homology in the formation of mosaic VSGs. Almost all
(C1= 99.71%, C2 = 99.46%) recombination events occurred within a region of shared
sequence between AnTat1.1 and each donor VSG, with an average length of ~9 bps
(average C1 = 9.13 bp, average C2 = 9.44 bp, median = 6 bp) (2F; Supplemental Figure
1A). Within the isolated mosaic clones, we observe short insertions (< 200 bp, average =
47 bp, median = 23 bp) predominating among the events (2D). Although read lengths

limited our ability to detect larger insertions with VSG-AMP-seq, approximately 55-60%
of the mosaic recombination reads detected by VSG-AMP-seq contained the same short
insertions (average C1 = 46 bp, average C2 = 45.8 bp) (Supplemental Figure 3B).
215
Notably, only a small number of donor VSGs were used for the majority of
recombination events. Upon closer inspection, these donors are members of a 6-VSG
family that contains AnTat1.1 and represent the only sequences within the EATRO1125
VSGnome with significant homology to this VSG (2E). N-terminal recombination events
appear restricted to just these family members (C1 = 100%, C2 = 99.87%). Together,
these results suggest that VSG sequence homology influences the outcome of VSG
recombination after a DNA break.
223
To determine the outcome following DNA damage of a VSG when there are no
homologous donors available, we used the same Cas9 system in the commonly used
Lister427 *T. brucei* line to cut the actively expressed VSG, VSG-2, which lacks
homologous family members. After inserting stably expressed guides targeting VSG-2
into the genome, we induced Cas9 expression for 7 days. Parasite clones isolated after
a break had all switched VSG expression from VSG-2, with no evidence of mosaic
recombination (Supplemental Figure 3A; cut position 707, n = 4; cut position 1082, n =
3). This further supports the hypothesis that homology between donor VSGs and the
region surrounding the DNA break drives mosaic formation.
233
We sought to determine what proportion of the VSG repertoire contains VSGs that
are members of VSG families and thus capable of diversifying through break-induced
mosaic formation. We defined the phylogenetic distance between VSGs within each of
the EATRO and Lister427 strains using a generalized time reversible model and found
that the proportions of the repertoire within families in each strain were remarkably similar.
In both strains, ~75% of the known VSG sequences are within a family and likely capable
of diversifying through the mechanisms described here (2G).
241

Figure 2. DNA double-strand breaks trigger mosaic VSG formation if homology is available
A) A histogram of unique recombination events identified along the AnTat1.1 transcript. The Cas9 DNA break site is indicated by a vertical line. Sections of the histogram are colored to indicate donor VSG identified. R indicates a guide that binds to the reverse strand. The midpoint of the perfect homology between AnTat1.1 and the donor VSG at the recombination site is plotted. If a donor VSG could not be unambiguously identified, the average recombination position was plotted. B) Quantification of mosaic recombination events induced by DNA breaks. The number of recombination events detected within 250bp up or downstream of the cut site was normalized to the number of total unanchored reads aligning within that region compared to the unanchored read count from the region with the smallest coverage to control for sequencing depth. (n=2, two independent clones) Statistical significance was determined with a one-way ANOVA with post-hoc Tukey HSD (** p <0.01) C) Schematics of mosaic VSGs from clones isolated after DNA breaks within AnTat1.1. Representative sequences shown. D) A histogram of donor VSG insertion lengths identified in all mosaic VSG clones isolated. The insert length only includes newly inserted sequence and does not include recombination sites. E) A schematic of the AnTat1.1 family aligned to the AnTat1.1 transcript. Gray sequences are a perfect match to AnTat1.1. F) A histogram of the length of the shared identity between AnTat1.1 and the donor VSG at the recombination sites. G) Quantification of types of VSGs within the VSGnomes from EATRO1125 and Lister427 parasites. The Lister427 VSGnome has 12 VSGs which are duplicated but lack diversification. SL = 5′ splice leader cap, 14-mer = 3′ sequence conserved in all VSG transcripts

Mosaic formation relies on a template for break-induced VSG diversification

Although most AnTat1.1-derived mosaics were identical to the putative donor VSG, many of these insertions altered only a few bps in AnTat1.1. We thus reasoned that it was possible that these events were *de novo* mutations created during DNA repair that happened to match the putative genome-encoded donor VSG. To test this possibility, we engineered dox-inducible Cas9-expressing Lister427 parasites and replaced VSG-2 with AnTat1.1 at the active expression site, BES1. Lister 427 has a different repertoire of VSGs compared to EATRO1125[2,20]. While AnTat1.1 is not endogenously present, there are four VSGs nearly identical to the AnTat1.1 family member VSG-2986 (99.5%, 98%, 97.5%, and 95.5% nucleotide sequence identity) (3A) that could serve as donor VSGs after a break in AnTat1.1. We detected hundreds of recombination events in two independent clones (clone 1 = 398, clone 2 = 833; 3C). Upon cutting, all mosaic recombination events detected utilized donor VSGs exclusively found in the Lister427 genome (3B), indicating that the repair events that generate mosaic VSGs require a template.

Figure 3. Nucleotide changes observed in mosaic VSGs arise from templated insertions
A) Schematics of the EATRO1125 AnTat1.1 VSG family and the VSGs similar to AnTat1.1 found within Lister427. Each is aligned to the AnTat1.1 transcript. For regions that are identical between two donor VSGs, regions are shown in the same color. B) A histogram of unique recombination events identified within Lister427 expressing AnTat1.1 parasites after a cut at position 694 along the AnTat1.1 transcript. The midpoint of the perfect homology between AnTat1.1 and the donor VSG at the recombination site is plotted. All AnTat1.1 family members except VSG-2986 were included as potential identifiable donor VSGs (all Lister VSGs + VSG-228, VSG-3110, VSG-7358, and VSG-4156). C) Quantification of mosaic recombination events induced by Cas9 at position 694. EATRO data is from 2A and 2B. The number of recombination events detected within 250bp up or downstream of the cut site was normalized to the number of total unanchored reads aligning within that region compared to the unanchored read count from the region with the smallest coverage from the EATRO cut sites to control for sequencing depth. SL = 5′ splice leader cap, 14-mer = 3′ sequence conserved in all VSG transcripts

Mosaic VSGs generated *in vitro* are identical to a subset of mosaic VSGs formed *in vivo*

To investigate whether the mosaic VSGs we detected after break induction *in vitro* reflected the events that occur naturally *in vivo*, we performed VSG-AMP-seq on parasites isolated from wildtype mouse infections, on day 15 post-infection, when AnTat1.1 has been mostly eliminated from the blood (Supplemental Figure 4B & 4C). We observed a strong C-terminal bias to all recombination events (316 recombination events across 7 mice). Very few sequences aligned to AnTat1.1 within the N-terminus, suggesting that most of these mosaic VSGs were complete replacements of the VSG N-terminal domain (4A; Supplemental Figure 4A). These donor VSGs shared significantly less homology with AnTat1.1 than the AnTat1.1 family members, usually only sharing spans of 100bps or less of imperfect homology within the C-terminal region of the VSGs. Nevertheless, there was still a short (average = 12 bp) span of perfect identity between AnTat1.1 and the donor at the recombination site (4E). Since VSGs expressed by parasites at the second peak of parasitemia are antigenically distinct from AnTat1.1, we reasoned these N-terminal replacement VSGs ensured complete immune evasion since only a small portion of AnTat1.1 was retained.

We wondered whether this restriction within mouse infections reflected a mechanistic bias towards recombination within the VSG C-terminal domain or the effect of host antibody selection, with only the most evasive recombination events surviving. To investigate these possibilities, we repeated the experiment in µMT mice[21], which do not have B-cells and therefore do not generate antibodies. We again analyzed day 15 post-infection, though AnTat1.1 is never cleared from these infections and parasitemia remains high throughout (Supplemental Figure 4B & 4C). In µMT mice, mosaic recombination events span the full length of the VSG, and many are identical to mosaic VSGs observed *in vitro* (4A; Supplemental Figure 4A) (2144 recombination events across 7 mice). *In vivo,* we observe short insertions (average = 41 bp) using donors homologous to AnTat1.1 flanked by short (average = 13 bp, median = 9 bp) regions of perfect identity, matching the patterns we observed *in vitro* (4E & 4F). Interestingly, as in the wildtype infections, there are an additional subset of recombination events within the C-terminus which are largely absent *in vitro* and use a much more diverse set of donors (4B). These data demonstrate that our *in vitro* system recapitulates a subset of *in vivo* recombination events, but there may be other pathways facilitating recombination *in vivo* in addition to those that can be triggered by a single, blunt DNA double-strand break.

Mosaic VSGs form in extravascular spaces during infection

While AnTat1.1 is eliminated from the blood in wildtype mice, it persists within tissues until at least day 14[4]. Given that parasite clearance is delayed in extravascular spaces, we wondered if mosaic formation may occur more readily in this parasite niche. To investigate this, we analyzed all assembled VSGs from a previous study of extravascular parasite populations[4]. Again, we found AnTat1.1 mosaics present in tissues identical to those observed in both µMT mice and *in vitro* (4C & 4G). Moreover, we observe that mosaic derivates of AnTat1.1 increase over time within tissue spaces during

infection (4D). Together, these data suggest that mosaic VSGs form preferentially within extravascular spaces, possibly due to the slower VSG-specific parasite clearance in these spaces.

Figure 4. AnTat1.1 mosaic VSGs form *in vivo* and reside within tissues of WT mice
A) A histogram of unique recombination events identified in all mice in wildtype or µMT mice from Day 15 post-infection. Recombination events found in multiple mice are represented once. (WT n=7; µMT n=7, from 3 independent experiments) The midpoint of the perfect homology between AnTat1.1 and the donor VSG at the recombination site is plotted. If a donor VSG could not be unambiguously identified, the average recombination position was plotted. B) Quantification of the number of donor VSGs utilized in mosaic recombination events within the N and C terminal domain of AnTat1.1 in wildtype and µMT mice from A). Statistical significance determined by a pairwise Wilcoxon test (*p<0.05) C) A histogram of the mosaic recombination events identified from VSG ORFs assembled in Beaver et al. All recombination events identified are shown at all time points sampled during infection: D6, D10, and D14. If a mosaic recombination event was identified in more than one tissue within the same mouse, it was counted once (n=12, 4 mice per tissue time point). D) Quantification of the mosaic recombination events detected within wildtype mouse blood or tissue. Statistical significance determined by pairwise Wilcoxon within each timepoint. (**p<0.01) E) A histogram of the length of the shared identity between AnTat1.1 and the donor VSG at the recombination sites. F) A histogram of donor VSG insertion lengths identified within individual reads. The insert length only includes newly inserted sequence and does not include recombination sites. G) Representative schematics of mosaic VSGs from assembled ORFs in Beaver et al. SL = 5′ splice leader cap, 14-mer = 3′ sequence conserved in all VSG transcripts

Small changes in the VSG sequence provide substantial antibody evasion

Many AnTat1.1-derived mosaic VSGs differ from their original sequence by only a few amino acids. To determine how these changes impacted the antigenic character of the mosaic VSGs, we analyzed live parasites expressing AnTat1.1 mosaic derivatives by flow cytometry using a potent rabbit anti AnTat1.1 polyclonal antibody raised against purified AnTat1.1 protein (5A). Many of the mutations block about 50% of antibody binding. Interestingly, three VSG-2986 mosaic clones were isolated with increasing insertion length (5B). The smallest insertion (18bps, with 3 a.a. changes, Mosaic 2), appears to account for most of the antigenic change, with only small decreases in binding associated with each additional length of insertion. We modeled the structure of these mosaics using ColabFold[22] and their general structures matched AnTat1.1 (5C), except at the disordered top of the N-terminal lobe. A careful examination of the locations of the mutations on AnTat1.1 show that mutations from Mosaic 2 which have the largest effect on antibody binding are at the apex of the structure while other alterations can be found on the side of the monomer (5D). These results demonstrate that, although the insertions characteristic of mosaic recombination are quite short, even these small changes to the VSG can confer large consequences for host antibody binding depending on their position.

Figure 5. Mutations within VSG hypervariable region shift antibody binding

A) A histogram showing the Alexa Fluor 647 staining intensity for parental controls (gray) and mosaic clones colored by donor VSG. (n= 4, from 3 independent parental lineages derived from 2 Cas9 clones) Schematics showing the mosaic VSGs are to the right. Donor VSG and amino acid substitutions are specified. B) Quantification of staining intensity changes for individual clones, based on median staining intensity. The median Alexa Fluor 647 intensities were normalized to the average of the parental clone. Statistical significance was determined based on a one-way ANOVA with a post-hoc Tukey HSD. (***p<0.001) C) Overlapping ribbon structures of AnTat1.1 and Mosaic 1 in blue and AnTat1.1 and Mosaic 2, Mosaic 3, and Mosaic 4 in purple as predicted by ColabFold. D) A space filling model of AnTat1.1 highlighting the changed residues within the monomer. VSG-3110 in blue and VSG-2986 in purple.

Discussion

Chronic *T. brucei* infection relies on the generation of new VSGs, with novel VSGs dominating late stages of infection[3,5,6]. However, the new VSGs generated during infection are complex, possibly arising from several intermediate events, and selected within a host environment with immunity against numerous previously expressed VSGs. The inherent complexity of chronic infection thus obscures the underlying biological principles driving VSG diversification. Here, we demonstrate that VSG diversification can be induced *in vitro* using Cas9-mediated double-strand DNA breaks within the VSG coding sequence, reproducibly generating mosaic VSGs that faithfully recapitulate mosaics formed naturally *in vivo*. By selecting just one VSG and looking at thousands of recombination outcomes, we have defined patterns characteristic of mosaic recombination. Mosaic VSGs typically form through short, templated insertions, and homology drives this process, restricting donor VSGs within the N-terminus to those from a set of closely related family members. Finally, we demonstrate that mosaic VSGs provide substantial immune evasion, particularly when these changes occur at the top of the VSG N-terminus, which may reflect a hypervariable region within the VSG.

We have shown that DNA breaks, previously shown to drive VSG switching[16,17], can also result in VSG diversification through mosaic VSG formation. While the exact DNA repair mechanism that generates mosaics is unclear, the patterns we observe— short templated-insertions relying on short stretches of sequence homology—suggest a mechanism similar to microhomology-mediated end joining (MMEJ). MMEJ is known to function in *T. brucei* and is typically RAD51-independent[23]. In line with this, there is evidence for a RAD51-independent pathway capable of driving antigenic variation in the parasite[24]. If an MMEJ-mediated small insertion occurs following a blunt, double-strand break, other types of breaks, like 5′ or 3′ staggered double-strand breaks, may result in alternative repair outcomes[25]. In B-cells, which use DNA breaks to undergo class switching and somatic hypermutation, the type of DNA break appears to dictate repair pathway choice[26,27]. It is thus possible that the N-terminal replacements we observe *in vivo*, but not *in vitro*, originate from multiple breaks, or staggered breaks with single-strand overhangs, that are repaired through a pathway distinct from the one driving repair of Cas9-induced breaks *in vitro*.

While our data suggest most (75%) VSGs in the archive are capable of diversification through the mechanism we have described, this is not the case for all VSGs. Indeed, we find that mosaic recombination occurs only if a homologous donor VSG is available. Most studies of antigenic variation in *T. brucei* have focused on VSG-2, a unique VSG that does not readily diversify; this may explain, at least in part, why mosaic recombination has been observed so rarely *in vitro* until now. The proportion of VSGs within families is surprisingly similar between the EATRO1125 and Lister427 strains. Perhaps this ratio has evolved to provide a balance between diversification events, which may not always provide full immune evasion, and switching, which is more likely to completely escape pre-existing immunity.

While the members of the AnTat1.1 VSG provide numerous possibilities for

recombination and the generation of new VSGs, we found that the center portion of the
gene appears to be especially successful at generating recombinants. We considered
that this might be attributable to available donor sequences or guide efficiency, but there
was nothing apparently unique about this site compared to other break locations. Protein
modeling of AnTat1.1, however, revealed that this region encodes the top of the N-
terminal lobe of the VSG, a site directly exposed to host antibody. For AnTat1.1 and many
similar VSGs[2] (A2-type), this is an unstructured region[28], and indeed the apex of most
crystalized VSG appears unstructured despite a surprising variety of underlying
structures[28]. We propose that this disordered region may have evolved to be more
tolerant to amino acid changes, allowing diversification within the part of the protein most
likely to facilitate immune evasion. While many recombination events may occur after a
DNA break, we hypothesize that only a few, particularly those within disordered regions,
maintain VSG structural integrity. Since expression of the VSG is essential for cell cycle
progression[29], it is possible that many newly formed mosaic VSGs lead to cell death for
the parasites, allowing only a subset of the recombination events to survive and resulting
in an apparent increase in recombination within the regions of the VSG where mutation
is most tolerated.
393
Although mosaic insertion events are often small (average ~46 bp) and can result
in few amino acid changes, these small changes, especially within the top of the N-
terminal lobe, confer substantial immune evasion. Just three amino acid substitutions,
from an 18 bp templated replacement, block approximately 50% antibody binding from a
potent polyclonal antiserum. While we have shown that these recombination events can
result in dramatic antigenic changes, many are also likely to result in incomplete immune
evasion. Perhaps as a result, mosaic VSGs derived from AnTat1.1 in WT blood appear
to be mostly N-terminal replacements, where the AnTat1.1 N-terminal domain is
completely replaced. A wider diversity of AnTat1.1-derived mosaics can be observed
within the tissues, however, where AnTat1.1-expressing parasites linger due to
differences in immune pressure[4]. In μMT mice, where AnTat1.1 expressing parasites can
linger indefinitely, this observation is further amplified. We thus hypothesize that
generating new VSGs is an iterative process, where a series of recombination events
progressively shifts the character of a VSG until an immunologically distinct variant is
formed. The relatively "protected" tissue spaces may serve as a site for this iterative
process to occur. Another intriguing possibility is that partial immune evasion is
advantageous for the parasite: a recent study suggested that exposure to sublethal
antibody concentrations can trigger a VSG switch in *T. brucei*[30]. Perhaps parasites
expressing these partially evasive variants are more prone to switching to a completely
new VSG.
414
In addition to shedding light on the selective pressures imposed upon mosaic
variants, our mouse data also demonstrate that our *in vitro* model of antigenic
diversification recapitulates VSG diversification *in vivo*. The recombination events we
detect in μMT mice represent the full breadth of possible recombination events which
AnTat1.1 can form, and these are largely represented within our *in vitro* break-induced
mosaic VSG populations. We observe short insertions flanked by short regions of identity
in both contexts, but the dominance of the 694 break position we observe *in vitro* is not

reflected in our *in vivo* analysis. This could be for a variety of reasons, including non-
uniform break locations along the VSG *in vivo* or a variety of DNA break types driving a
different pattern of outcomes. Intriguingly, in the absence of an exogenous DNA break,
we do not observe any diversification *in vitro*, while diversification occurs continuously in
the μMT context. This suggests that some aspect of the host environment other than
antibody pressure induces diversification *in vivo.* It remains to be investigated whether an
in vivo cue for diversification exists.
429
The diversification mechanisms we have described may also be at play in other
gene families where diversity is critical. Many pathogens express variable antigens, with
the antigenic repertoire stored within subtelomeric regions of the genome where
continuous DNA damage might facilitate diversification. Mosaic antigen genes have been
described in plasmodium var genes[31–34] and giardia variant-specific surface proteins
(VSPs)[35,36], some of which resemble the small insertion mosaics we have observed here.
Alternative end joining pathways appear to be intact in these organisms[37], and may
facilitate mosaic formation. Many alternative forms of DNA repair are also conserved in
mammals[38]; it is thus plausible that large gene families like olfactory receptors[39,40] and
protocadherins[39], which depend upon diversity and are known to swap sequences
through gene conversion events[41–44], may also be diversifying through similar
mechanisms. A rigorous targeted approach, like we have used here, may be required to
determine the role this mechanism plays in diversification of other gene families.
443
Here we answer long-standing questions about how the *T. brucei* VSG repertoire
evolved to generate new antigens critical for maintaining a chronic infection. Our high-
throughput, sensitive technique has enabled the characterization of thousands of mosaic
diversification events, orders of magnitude more than was previously feasible. This
analysis revealed a pattern of short, homology-driven, and templated insertions around
DNA break sites that can shift the antigenic character of the VSG. More broadly, this study
provides an experimental framework for the hypothesis-driven exploration of antigen
diversification in *T. brucei* as well as in other pathogenic microbes.
452

Methods

Parasites:

Pleiomorphic EATRO1125 AnTat1.1 90-13 *T. brucei* parasites were maintained in HMI-9 media with 10% heat inactivated FBS and 10% Serum Plus. Parasites were passaged when they reached approximately $5*10^5$ cells/mL. Monomorphic Single Marker Lister427 VSG221 TetR T7RNAP bloodstream form (NR42011; LOT: 61775530)[45]. *T. brucei* were maintained in HMI-9 up to $1*10^6$ parasites/mL. VSG221 has since been renamed to VSG-2.

Plasmids:

Plasmids were synthesized with Gibson Assembly using the following templates, when specified. Gibson Assembly master mix was custom made[46]. Whole plasmid sequencing was performed by Plasmidsaurus using Oxford Nanopore Technology with custom analysis and annotation.

pLEW100v5-BSD-FLAG-La-Cas9 was synthesized from pLEW100v5-BSD and pRPaCas9[47]. A *T. brucei* codon optimized FLAG tag was added to the N-terminus of Cas9. pLEW100V5-BSD was a gift from George Cross (Addgene plasmid # 27658 ; http://n2t.net/addgene:27658 ; RRID:Addgene_27658). pRPaCa9 was a gift from David Horn (Addgene plasmid # 111819 ; http://n2t.net/addgene:111819 ; RRID:Addgene_111819).

pHH-HYG-AnTat1.1 and pHH-HYG-VSG-228 were synthesized from pHH-HYG-VSG-3-S317A.[48] This was a gift from Joey Verdi. AnTat1.1 sequence was obtained from AnTat1.1 specific cDNA from mouse infection D6 cloned into a pMiniT vector with the PCR Cloning Kit (NEB, E1202S). VSG-228 was partially amplified from VSG PCR (see below) derived from AnTat1.1 depleted D16 μMT parasites from blood. The partial fragment was amplified with an AnTat1.1 family specific forward primer (5' - ACTACACCCACAACAAGCTCTA-3') and a pan VSG reverse primer which binds to a conserved 14bp region of the 3' UTR (5'-GATTTAGGTGACACTATAGTGTTAAAATATATC-3') with AmpliTaq Gold (Applied Biosystems, 4398881) (anneal & extension 60C 1m, 35 cycles).The resulting PCR was cloned into the pMiniT backbone and the remainder of VSG-228 was *de novo* synthesized with Gibson Assembly.

The pT7sgRNA plasmids were obtained according to Rico et al[47]. Briefly, sticky-end annealed sgRNAs were ligated into a BbsI-HF (NEB, R0539S) digestion of pT7sgRNA. Supplemental Table 1 contains the sequences for the annealed guides. pT7sgRNA was a gift from David Horn. (Addgene plasmid # 111820 ; http://n2t.net/addgene:111820 ; RRID:Addgene_111820)

Transgenic parasites:

To obtain all transgenic parasites, 5 million parasites were electroporated with 5-10ug of digested plasmid with an AMAXA Nucleofector II using X-001 in Human T-cell Nucleofector Solution (Lonza VPA-1002).

For tetracycline-inducible Cas9 parasites, EATRO1125 or Lister427 parasites
were electroporated with pLEW100v5-BSD-FLAG-La-Cas9, digested with NotI-HF (NEB,
R3189S). Parasites were immediately selected in 5ug/mL Blasticidin (Thermo Scientific,
R21001) and maintained in selection unless otherwise specified.
To obtain AnTat1.1 and VSG-228 expressing Lister427 parasites, Lister427
parasites were electroporated with pHH-HYG-VSG plasmids digested with BamHI-HF
(NEB, R3136S). After 16-24 hours recovery, 25ug/mL of hygromycin (Fisher Scientific,
J67371-XF) was added to the culture. After obtaining colonies 5-7 days later, selection
was reduced to 5ug/mL and maintained unless otherwise specified.
Lister427 parasites with pLEW100v5-BSD-FLAG-La-Cas9 inserted were
additionally electroporated with pHH-HYG-VSG plasmids expressing Antat1.1. Clones
were obtained as above.
To obtain constitutive guide expressing parasite lines, parasites were
electroporated with pT7sgRNA guide containing plasmids digested with NotI-HF[47]. After
16-24 hours recovery, single colonies of parasites were selected with 2ug/mL phleomycin
(Sigma-Aldrich, SML3001). Parasites were maintained without phleomycin selection.
515
516 **T7-Guide Synthesis & Purification:**
517   DNA fragments containing a T7 promoter and guide RNA sequence were
518 synthesized as described previously[49]. Briefly, T7-guide primers and a G00 primer were
519 amplified with Phusion polymerase for 35 cycles (annealing temp 60C, extension 5s).
520 PCR products from 12 identical PCRs were pooled and purified via ethanol precipitation.
521
Cas9 Transient Electroporations:
Approximately 24 hours prior to electroporation, in a single flask, 1 million parasites
in 12mLs of media per sample was seeded and induced with 1ug/mL doxycycline
(Millipore Sigma, D9891-1G). Only blasticidin selection (Cas9) was maintained and for
Lister427 parasites expressing AnTat1.1, hygromycin (active VSG expression) was
removed at this stage. Either 8mLs (Lister427) or 10mLs (EATRO1125) of parasites was
spun down, media removed, and parasites were resuspended in 100uL Human T-cell
Nucleofector Solution. Each sample was electroporated using the X-001 program on the
AMAXA Nucleofector II with approximately 1-1.5ug of purified T7-guide in a volume less
than 10uL or a sample without any DNA as a negative control. Parasites were moved into
5mLs HMI-9 in 6-well plates to recover for 30 minutes then moved into 20mLs total in
flasks to recover overnight. About 24 hours after electroporation, parasites were counted
and split. For EATRO1125, 12 million cells (or all the cells if there were fewer than
12million) were seeded into 60mLs total with blasticidin. For Lister427, 2 million cells were
seeded into 20mLs total with blasticidin. At the 48 hour time point, parasites were counted,
collected, and stored in TRIzol (Invitrogen, 15596026) for subsequent RNA extraction.
538
Isolation of mosaic expressing parasites:
Multiple clones of parasites expressing T7sgRNA with guides were derived from
two distinct inducible Cas9-expressing parasite clones. Dilutions of parasites were plated
in 96-well plates in 5ug/mL blasticidin and 1ug/mL doxycycline. Individual clones were
visible after 7-14 days. Upon isolation of a parasite clone, all drugs were removed.
Parasites were analyzed by flow cytometry. VSG sequences for clones were determined

from extracted RNA. cDNA was synthesized using the Superscript III Reverse
Transcriptase (Invitrogen, 18080051) and a VSG-specific primer which binds to a
conserved 14-bp sequence within the 3' UTR. (5'-GTGTTAAAATATATC-3'). 2uL of
RNase-treated cDNA was amplified for 35 cycles with VSG specific primers: a spliced-
leader (5'-ACAGTTTCTGTACTATATTG-3') and SP6-VSG 14-mer (5'-
GATTTAGGTGACACTATAGTGTTAAAATATATC-3') using Phusion polymerase
(Thermo Fisher, F530L) (annealing temp 55C, extension 45s). PCR products were
cleaned using the Monarch PCR & DNA cleanup kit (NEB, T1030L). VSG sequences
were determined by amplicon nanopore sequencing performed by Plasmidsaurus using
Oxford Nanopore Technology with custom analysis and annotation or Sanger sequencing
with the sequencing primer (5'- AGAGAATACTAAGCTAGTTGGC-3') performed by
Azenta Life Sciences.

Mouse Infections:

C57BL/6 mice and μMT- (B6.129S2-Ighm[tm1Cgn]/J)[21] mice were delivered at 8
weeks old from Jackson Labs. Approximately 5 EATRO1125 parasites were injected into
mice intravenously through the tail vein. These parasites express either AnTat1.1 or VSG-
421. Starting at D4 post-infection, parasitemia was monitored within mice every two days
via tail bleed. Blood was harvested at D6 post-infection through a submandibular bleed.
An additional gel pack and in-cage food pellets were provided to mice during recovery
since it helped μMT mice survive the blood loss. At D15, post-infection, mice were
humanely euthanized with ketamine injection and blood was harvested through cardiac
puncture. Blood was stored in TRIzol LS (Invitrogen, 10296028) for RNA extraction.

RNA preparation:

Parasites were stored in TRIzol prior to RNA extraction. Blood with parasites was
stored in TRIzol LS. RNA was extracted via phenol/chloroform extraction according to the
manufacturers protocol. Purified RNA was DNase treated with Turbo Dnase (Thermo
Fisher, AM2239) and purified with 1.8X Mag-Bind TotalPure NGS Beads (Omega Bio-
tek, M1378-01). Verification of effective DNase treatment was performed via PCR of
Hygromycin (EATRO1125 only) (Fwd: 5'-ACAGCGGTCATTGACTGGAG-3'; Rev: 5'-
ATTTGTGTACGCCCGACAGT-3', annealing temp 52C, extension 30s) or HSP70
(Lister427 & EATRO1125) (Fwd: 5'-AGAACACTATCAATGACCCCAAC-3'; Rev: 5'-
CCATGCCCTGGTACATCT-3', annealing temp 50C, extension 15s) genes for 30 cycles
using OneTaq DNA Polymerase (NEB, M0480L).

VSG-seq:

VSG-seq was performed as described elsewhere[3,4]. Briefly, cDNA was
synthesized using the Superscript III Reverse Transcriptase and a VSG-specific primer
which binds to a conserved 14-bp sequence within the 3' UTR. (5'-GTGTTAAAATATATC-
3'). cDNA was treated with Rnase A (Quiagen, 19101) and RNase H (Invitrogen,
18080051) for 30 minutes then purified with 1.8X Mag-Bind Total NGS Beads (Omega
Bio-tek, M1378-01). Purified cDNA was amplified for 25 cycles with VSG specific primers:
a spliced-leader (5'-ACAGTTTCTGTACTATATTG-3') and SP6-VSG 14-mer (5'-
GATTTAGGTGACACTATAGTGTTAAAATATATC-3') using Phusion polymerase
(Thermo Scientific, F530L) (annealing temp 55C, extension 45s). This PCR product was

prepared for sequencing using the Nextera XT DNA Sample Prep Kit (Illumina, FC-131-
1096) according to the manufacturer's instructions. Libraries were sequenced with 100
bp single-end reads on a NovaSeq6000. Analysis was performed using the
VSGSeqPipeline found at github.com/mugnierlab.
595

VSG-AMP-seq library preparation:
VSG-AMP-seq was based upon AMP-seq[50] and GUIDE-seq[51]. cDNA was
synthesized from DNA-free RNA using Superscript III Reverse Transcriptase and a VSG-
specific primer which binds to a conserved 14-bp sequence within the 3' UTR. (5'-
GTGTTAAAATATATC-3'). Second Strand synthesis was performed with NEBNext
mRNA Second Strand Synthesis Module (E6111L). The resulting double stranded cDNA
was purified with purified with 1.8X Mag-Bind Total NGS Beads. cDNA was fragmented
briefly with NEBNext dsDNA Fragmentase (M0348L) for ten minutes at 37C to obtain
fragments of approximately 500bps. Fragments were purified with a double-sided Mag-
bind bead cleanup. First, fragments were incubated with 0.5X beads, DNA bound to the
beads were discarded and an equal volume of beads as before, then a 1X PEG
concentration, was added to the fragments and the cleanup proceeded as normal.
Fragments were end-repaired with Enzymatics low concentration end repair mix (Y9140-
LC-L) and A-tailed with Recombinant Taq (Life Technologies, 100021276). Y-Adapters
were pre-annealed by incubating a MiSeq Common Adapter with Adapters (A01-A10)
containing 25 bp Unique Molecular Indexes (UMIs) and a barcode at 10uM at 95C for 1
sec, 60C for 1s, and slowly cooled to 4C. Adapters were ligated to the A-tailed fragments
with T4 DNA Ligase (Enzymatics L6030-LC-L). The resulting fragments were cleaned up
with 0.8X Mag-Bind beads. Two target specific PCRs were performed on the fragments
with pools of target specific primers, one in the forward and one in the reverse direction.
All primers are listed in Supplemental Table 1. Target primers have spacers of varying
lengths to generate sequence diversity without the need for PhiX. Target specific primers
are paired with P5_2 and a sample specific P primer (P701-P710) which contains a
second barcode. Fragments are amplified with Platinum Taq (Life Technologies,
10966018) using the following program: 95C 5 mins, 35 cycles of 95C for 30s, 55C for
30s, and 72C for 30s, and 72 for 5 mins. The resulting products were cleaned up with
0.6X Mag-bind beads. Libraries were quantified using the Qubit dsDNA HS Kit (Life
Technologies, Q32854) and run on a 1% agarose gel to determine average length.
Libraries were sequenced on a MiSeq or for deep sequencing of μMT samples on a
NovaSeq6000 with custom index1 and read2 primers using the following cycle conditions:
"151|8|33|131" with the paired-end Nextera sequencing protocol.
627

VSG-AMP-seq analysis:
Index sequences were added to the names of sequencing reads. FASTQ files were
demultiplexed by target specific primer using cutadapt v=3.5[1] searching for multiple
target-specific primers at the 5' end of read2 in paired-end mode using flags --
action=retain --overlap 10. Then, reads were quality trimmed with trim_galore v0.6.4_dev
(github.com/FelixKrueger/TrimGalore) where adapter sequences, if present, were
removed with flags --dont_gzip --paired --trim1. Spacers were removed from read1 if
present using cutadapt by searching for the target-specific primer sequence at the 3' end
of the read with flags --action=retain --overlap 10. Dual-indexed barcodes were used to

demultiplex reads. Mismatch allowed was determined by the barcodes present, so that a
unique barcode could still be identified.
639
To consolidate reads, UMIs were extracted from large FASTQ files, split into
smaller files and grouped by 100% identity using cd-hit-est v4.8.1[53,54] with flags -c 1.0 -n
8 -M 16000 -d 0. UMIs were then grouped by 92% using flags -c 0.92 -n 8 -M 16000 -d 0.
Reads were sorted into consensus groups based on UMIs and a consensus sequence
was formed as the most popular base at each position of the read with a threshold quality
score of at least 2. If there was a tie, an N was used. Clusters with fewer than 3 reads
were removed.
647
The full sequence length of the AnTat1.1 transcript was determined by
Plasmidsaurus using Oxford Nanopore Technology with custom analysis and annotation
and the 5'-end was appended with alternative splicing information from control samples.
The positions were 0 indexed. Read2, the anchored read, was aligned to AnTat1.1. Read
pairs were removed if: the first 4/10 bases after the primer did not align to AnTat1.1, an
alternative VSG which contained the primer sequence was amplified, the wrong position
of AnTat1.1 was amplified by the anchor, if the reads were too short (<15 bps), contained
too many Ns (>5), if there was an inversion event or a duplication, or if the transcript was
alternatively spliced. Read1, the unanchored read, was also aligned to AnTat1.1 with a
mismatch of up to 1bp. If no alignment could be found, read1 was trimmed to remove
AnTat1.1 like sequences from the 5' and 3' ends, allowing up to 1 mismatch on each end
to be removed, leaving a fragment which can be used to search the VSGnome and
identify potential donor VSGs. A consensus sequence was generated from the reads, if
they overlapped, using their alignment positions within AnTa1.1. AnTat1.1, the consensus
read, and putative donor VSG were aligned, and recombination sites were identified. If
no consensus could be generated, read1 was used. Ambiguous recombination sites
where a single donor could not be identified were represented by the average position of
the potential recombination sites and identified as ambiguous.
666
VSG Clustering and family identification:
FASTA files of genes encoding VSGs from the Lister427 strain were derived from
TrypsRU   (vsgs_tb427_all_atleast150aas_cds.txt,   vsgs_tb427_
nodups_atleast250aas_cds.txt, vsgs_tb427_nodups_atleast250aas_cdsplusflanks.txt)
(George Cross) and TriTrypDB[55] (TriTrypDB-65 Lister Strain 427 2018)[20]. A combined
master FASTA file was standardized to include important attributes as part of the
sequence name (VSG name, GB Accession Code, Nucleotide Length, Status of
Sequencing, Chromosomal Location, Source, Parasite Line (Lister427), and any other
Extra Information) and the corresponding sequence. From the 2018 genome, the genes
encoding for VSGs were isolated by looking for the keywords, "variant" or "VSG" in the
description attribute. The genomic sequence for these VSG genes was extracted from
the Genome file and included 300 bp of flanking sequence up and downstream for further
analysis. Identical VSGs were clustered using cd-hit-est, with the following flags: -c 1.0 -
n 8 -M 16000 -d 0. A representative sequence, usually from the 2018 genome with the
flanking sequences, was selected for each VSG and was the longest sequence available.
Duplicate VSGs identified at two different chromosomal locations within the 2018 genome

were added to the file to represent all known Lister427 VSGs. AnTat1.1 was added to the
master Lister427 VSG file for further analysis.
EATRO1125 VSGs were obtained from TrypsRU (George Cross) and included 200
bp flanking sequences. (vsgs_tb1125_all_atleast150aas_cdsplusflanks.txt)
687
Multiple sequence alignment was performed using MAFFT[56], using the following
flags: --auto --reorder. Following multiple sequence alignment, a phylogenetic tree was
constructed using a generalized time reversible mode in FastTree[57,58] with the following
flags: -nt -gtr. Clustering of AnTat1.1 and its family members was used to identify a
threshold genetic distance to generate subsequent subtrees. Threshold distances of
0.24659 and 0.21677 were used for EATRO1125 and Lister427, respectively. Down
sampling and subtree generation of representative sequences was performed using
PARNAS (0.1.4)[59]. The command used for PARNAS included flag: --cover –radius
<threshold distance>. From the subtree generated, the number of unique VSGs and the
number of VSG families were identified. Duplicate VSGs from the Lister427 genome were
analyzed to determine if they were within a group that lacked any diversification.
699
Western Blotting:
5 million parasites were spun down and washed with 25C PBS. Then, pelleted
parasites were resuspended in 50uL RIPA buffer (50mM Tris, 150mM NaCl, 1% NP-40,
0.5% Sodium Deoxycholate, 0.1% SDS, pH = 7.4) + 2X Laemmli buffer. Lysed parasites
were boiled at 95C for 5 minutes. 5uL of lysates were separated on a Tris-glycine
polyacrylamide gel at 110V for 100 minutes in Tris/glycine running buffer (25mM Tris,
192mM glycine, 0.1% SDS, pH = 8.3). Proteins were then transferred onto a nitrocellulose
membrane using transfer buffer (25mM Tris, 192mM glycine, 20% methanol) overnight at
a 60mAmps per transfer box at 25C. For Cas9 (1:1000) and EF1α (1:1000), lysates were
separated on 10% polyacrylamide gels and transferred onto 0.45um membranes. For γ-
H2A (1:200), lysates were separated on a 15% polyacrylamide gel and transferred onto
a 0.2um membrane. After transfer, blots were blocked with 0.5% BSA in TBS with 0.05%
Triton X-100 (TBST, 20mM Tris, 150mM NaCl, pH = 7.6) for an hour at RT under constant
agitation. Blots were incubated with primary antibody (see above dilutions) for 2 hours at
25C. After 5 TBST washes, blots were stained for 1 hour with goat anti mouse (1:5000,
Cell Signaling, 7076S) or goat anti rabbit-HRP-conjugated secondary (1:5000, Cell
Signaling, 7074S). After another 5 washes, blots were incubated with ECL (Cytiva,
RPN2109) and film was exposed to blots in a dark room. Developed film was scanned
and images were processed with FIJI[60]. Primary antibodies used were m anti FLAG (M2
clone) (Millipore Sigma, F3165-1MG), m anti EF1α (CBP-KK1 clone) (Millipore Sigma,
05-235), and rb anti γ-H2A was a kind gift from Galadriel Hovel-miner based upon Glover
and Horn[61].
722
Flow Cytometry:
In 96 well plates, 200,000 parasites were stained with 1:20,000 rabbit anti
AnTat1.1 primary antibody (Jay Bangs) for ten minutes at 4C while shaking in PBS +
10mg/mL glucose. Parasites were washed once with 100uL PBS + glucose. Then,
parasites were stained with Alexa Fluor 647-conjugated goat anti rabbit IgG (H +L),
F(ab')2 Fragment (Cell Signaling, 4414S) at 1:1000 at 4C while shaking in PBS + glucose.

After washing again with 100uL PBS + glucose, parasites were resuspended in PBS + glucose + 1:20 Propidium Iodide (BD Biosciences, 556463) and analyzed on a Attune Nxt flow cytometer (Invitrogen). Data analysis was performed using FlowJo v10.

Analysis and Modeling of VSG N-terminal Domains:
Full length protein coding sequences from AnTat1.1 and its isolated mosaics were used for structural modeling. Only N-terminal domain sequences were used for protein structural prediction since this region of the VSG is the most well-defined experimentally. Signal peptides are cleaved from the mature VSG during processing, so we used SignalP 6.0[62] to predict and remove the Sec/SPI sequence (--organism eukarya, --mode fast) resulting in a FASTA file of mature proteins. To determine the coordinate of the N-terminal domain, we used a python analysis pipeline available at (https://github.com/mugnierlab/find_VSG_Ndomains). The script identifies the boundaries of the VSG N-terminal domain using the HMMscan function under HMMer version 3.1b2[63]. Query sequences are searched against an HMM profile containing 735 known N-terminal domain sequences from Cross et al[2]. and N-terminal domains defined by the largest envelope domain coordinate that meets E value threshold (1 x 10^{-5}, -domE 0.00001). The processed FASTA file containing only mature VSG N-terminal domain sequences was used as input for structural prediction of monomers using LocalColabFold[22] function colabfold_batch run using the following arguments: --amber, --templates, --num_recycle 3. The best ranked output model with the highest average predicted local distance test score (pLDDT), that is the highest confidence model, was used for downstream analyses. Model visualization and alignment were performed using with UCSF ChimeraX version 1.7.1[64], developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

Acknowledgements

We would like to thank Jay Bangs for generating and sending us primary rabbit anti-AnTat1.1 polyclonal antibody. Thanks to Galadriel Hovel-miner for the primary rabbit anti-γ-H2A antibody. We appreciate members of the Mugnier lab for their thoughtful review of this manuscript, particularly Carolina Duque and Sneider Gutierrez. We also would like to thank Richard McCulloch for thoughtful feedback about this manuscript. We would like to thank David Mohr and the JHU GCRF Genomics Core for sequencing services. Thanks to Joey Verdi for pHH-HYG-VSG-3-S317A and advice about the generation of transgenic parasites. Data analysis was carried out at the Advanced Research Computing at Hopkins (ARCH) core facility (rockfish.jhu.edu), which is supported by the National Science Foundation (NSF) grant number OAC 1920103. This work was supported by the JHU Catalyst Award. JES, JMCH, AKB, JS, BZ, and MRM were supported by Office of the Director, NIH (DP5OD023065). JES and JS were additionally supported NIH training grant T32GM007 and JES was supported by the National Science Foundation (NSF) Graduate Research Fellowship under grant numbers 1232825 and 1746891. JMCH, AKB, JS and EHA were additionally supported by NIH T32AI007417.

References

1. Hertz-Fowler, C. *et al.* Telomeric Expression Sites Are Highly Conserved in Trypanosoma brucei. *PLoS ONE* **3**, e3527 (2008).
2. Cross, G. A. M., Kim, H.-S. & Wickstead, B. Capturing the variant surface glycoprotein repertoire (the VSGnome) of Trypanosoma brucei Lister 427. *Mol. Biochem. Parasitol.* **195**, 59–73 (2014).
3. Mugnier, M. R., Cross, G. A. M. & Papavasiliou, F. N. The in vivo dynamics of antigenic variation in *Trypanosoma brucei. Science* **347**, 1470–1473 (2015).
4. Beaver, A. *et al.* Extravascular spaces are reservoirs of antigenic diversity in Trypanosoma bruci infection. *bioRxiv* (2022) doi:https://doi.org/10.1101/2022.06.27.497797.
5. Hall, J. P. J., Wang, H. & Barry, J. D. Mosaic VSGs and the Scale of Trypanosoma brucei Antigenic Variation. *PLoS Pathog.* **9**, e1003502 (2013).
6. Roth, C., Bringaud, F., Layden, R. E., Baltz, T. & Eisen, H. Active late-appearing variable surface antigen genes in Trypanosoma equiperdum are constructed entirely from pseudogenes. *Proc. Natl. Acad. Sci.* **86**, 9375–9379 (1989).
7. So, J. *et al.* VSGs Expressed during Natural T. b. gambiense Infection Exhibit Extensive Sequence Divergence and a Subspecies-Specific Bias towards Type B N-Terminal Domains. *mBio* **13**, e02553-22 (2022).
8. Pays, E. *et al.* Gene conversion as a mechanism for antigenic variation in Trypanosomes. *Cell* **34**, 371–381 (1983).
9. Baltz, T. *et al.* Exposed epitopes on a Trypanosoma equiperdum variant surface glycoprotein altered by point mutations. *EMBO J.* **10**, 1653–1659 (1991).
10. Kamper, S. M. & Barbet, A. F. Surface epitope variation via mosaic gene formation is potential key to long-term survival of Trypanosoma brucei. *Mol. Biochem. Parasitol.* **53**, 33–44 (1992).
11. Thon, G., Baltz, T. & Eisen, H. Antigenic diversity by the recombination of pseudogenes. *Genes Dev.* **3**, 1247–1254 (1989).
12. Lu, Y., Hall, T., Gay, L. S. & Donelson, J. E. Point mutations are associated with a gene duplication leading to the bloodstream reexpression of a trypanosome metacyclic VSG. *Cell* **72**, 397–406 (1993).
13. Rice-Ficht, A. C., Chen, K. K. & Donelson, J. E. Point mutations during generation of expression-linked extra copy of trypanosome surface glycoprotein gene. *Nature* **298**, 676–679 (1982).
14. Lu, Y., Alarcon, C. M., Hall, T., Reddy, L. V. & Donelson, J. E. A strand bias occurs in point mutations associated with variant surface glycoprotein gene conversion in Trypanosoma rhodesiense. **14**, 10 (1994).
15. Berriman, M. *et al.* The architecture of variant surface glycoprotein gene expression sites in Trypanosoma brucei. *Mol. Biochem. Parasitol.* **122**, 131–140 (2002).
16. Boothroyd, C. E. *et al.* A yeast-endonuclease-generated DNA break induces antigenic switching in Trypanosoma brucei. *Nature* **459**, 278–281 (2009).
17. Glover, L., Alsford, S. & Horn, D. DNA Break Site at Fragile Subtelomeres Determines Probability and Mechanism of Antigenic Variation in African Trypanosomes. *PLoS Pathog.* **9**, e1003260 (2013).

18. Kou, R. *et al.* Benefits and Challenges with Applying Unique Molecular Identifiers in
Next Generation Sequencing to Detect Low Frequency Mutations. *PLOS ONE* **11**,
e0146638 (2016).
19. Kanagawa, T. Bias and artifacts in multitemplate polymerase chain reactions (PCR).
J. Biosci. Bioeng. **96**, 317–323 (2003).
20. Müller, L. S. M. *et al.* Genome organization and DNA accessibility control antigenic
variation in trypanosomes. *Nature* **563**, 121–125 (2018).
21. Kitamura, D., Roes, J., Kühn, R. & Rajewsky, K. A B cell-deficient mouse by
targeted disruption of the membrane exon of the immunoglobulin µ chain gene.
Nature **350**, 423–426 (1991).
22. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat. Methods*
19, 679–682 (2022).
23. Glover, L., Jun, J. & Horn, D. Microhomology-mediated deletion and gene
conversion in African trypanosomes. *Nucleic Acids Res.* **39**, 1372–1380 (2011).
24. McCulloch, R. & Barry, J. D. A role for RAD51 and homologous recombination in
Trypanosoma brucei antigenic variation. *Genes Dev.* **13**, 2875–2888 (1999).
25. Scully, R., Panday, A., Elango, R. & Willis, N. A. DNA double-strand break repair-
pathway choice in somatic mammalian cells. *Nat. Rev. Mol. Cell Biol.* **20**, 698–714
(2019).
26. Ling, A. K. *et al.* Double-stranded DNA break polarity skews repair pathway choice
during intrachromosomal and interchromosomal recombination. *Proc. Natl. Acad.*
Sci. **115**, 2800–2805 (2018).
27. So, C. C. & Martin, A. DSB structure impacts DNA recombination leading to class
switching and chromosomal translocations in human B cells. *PLOS Genet.* **15**,
e1008101 (2019).
28. Đaković, S. *et al.* A structural classification of the variant surface glycoproteins of the
African trypanosome. *PLoS Negl. Trop. Dis.* **17**, e0011621 (2023).
29. Sheader, K. *et al.* Variant surface glycoprotein RNA interference triggers a
precytokinesis cell cycle arrest in African trypanosomes. *Proc. Natl. Acad. Sci.* **102**,
8716–8721 (2005).
30. Tenaglia, A. H. *et al. Antibodies to Protozoan Variable Surface Antigens Induce*
Antigenic Variation. http://biorxiv.org/lookup/doi/10.1101/2022.06.21.497077 (2022)
doi:10.1101/2022.06.21.497077.
31. Zhang, X. *et al.* Rapid antigen diversification through mitotic recombination in the
human malaria parasite Plasmodium falciparum. *PLOS Biol.* **17**, e3000271 (2019).
32. Bopp, S. E. R. *et al.* Mitotic Evolution of Plasmodium falciparum Shows a Stable
Core Genome but Recombination in Antigen Families. *PLoS Genet.* **9**, e1003293
(2013).
33. Claessens, A. *et al.* Generation of Antigenic Diversity in Plasmodium falciparum by
Structured Rearrangement of Var Genes During Mitosis. *PLoS Genet.* **10**, e1004812
(2014).
34. Bull, P. C. *et al. Plasmodium falciparum* antigenic variation. Mapping mosaic *var*
gene sequences onto a network of shared, highly polymorphic sequence blocks.
Mol. Microbiol. **68**, 1519–1534 (2008).
35. Yang, Y. & Adam, R. D. A group of Giardia lamblia variant-specific surface protein
(VP) genes with nearly identical 5 ' regions. *Mol. Biochem. Parasitol.* (1995).

36. Adam, R. D. *et al.* The Giardia lamblia vsp gene repertoire: characteristics, genomic
organization, and evolution. *BMC Genomics* **11**, 424 (2010).
37. Nenarokova, A. *et al.* Causes and Effects of Loss of Classical Nonhomologous End
Joining Pathway in Parasitic Eukaryotes. *mBio* **10**, e01541-19 (2019).
38. Schimmel, J., van Schendel, R., den Dunnen, J. T. & Tijsterman, M. Templated
Insertions: A Smoking Gun for Polymerase Theta-Mediated End Joining. *Trends*
Genet. **35**, 632–644 (2019).
39. Williams, D. L. *et al.* May the Odds Be Ever in Your Favor: Non-deterministic
Mechanisms Diversifying Cell Surface Molecule Expression. *Front. Cell Dev. Biol.* **9**,
720798 (2022).
40. Zozulya, S., Echeverri, F. & Nguyen, T. The human olfactory receptor repertoire.
Genome Biol. **2**, research0018.1 (2001).
41. Lai, P. C. *et al.* An olfactory receptor pseudogene whose function emerged in
humans: a case study in the evolution of structure–function in GPCRs. *J. Struct.*
Funct. Genomics **9**, 29–40 (2008).
42. Niimura, Y. & Nei, M. Evolution of olfactory receptor genes in the human genome.
Proc. Natl. Acad. Sci. **100**, 12235–12240 (2003).
43. Newman, T. & Trask, B. J. Complex Evolution of 7E Olfactory Receptor Genes in
Segmental Duplications. *Genome Res.* **13**, 781–793 (2003).
44. Noonan, J. P., Grimwood, J., Schmutz, J., Dickson, M. & Myers, R. M. Gene
Conversion and the Evolution of Protocadherin Gene Cluster Diversity. *Genome*
Res. **14**, 354–366 (2004).
45. Wirtz, E., Leal, S., Ochatt, C. & Cross, George A. M. A tightly regulated inducible
expression system for conditional gene knock-outs and dominant-negative genetics
in Trypanosoma brucei. *Mol. Biochem. Parasitol.* **99**, 89–101 (1999).
46. Isothermal Reaction (Gibson Assembly) Master Mix. *Cold Spring Harb. Protoc.*
2017, pdb.rec090019 (2017).
47. Rico, E., Jeacock, L., Kovářová, J. & Horn, D. Inducible high-efficiency CRISPR-
Cas9-targeted gene editing and precision base editing in African trypanosomes. *Sci.*
Rep. **8**, 7960 (2018).
48. Gkeka, A. *et al. Immunodominant Surface Epitopes Power Immune Evasion in the*
African Trypanosome. http://biorxiv.org/lookup/doi/10.1101/2021.07.20.453071
(2021) doi:10.1101/2021.07.20.453071.
49. Shaw, S., Knüsel, S., Hoenner, S. & Roditi, I. A transient CRISPR/Cas9 expression
system for genome editing in Trypanosoma brucei. *BMC Res. Notes* **13**, 268 (2020).
50. Zheng, Z. *et al.* Anchored multiplex PCR for targeted next-generation sequencing.
Nat. Med. **20**, 1479–1484 (2014).
51. Tsai, S. Q. *et al.* GUIDE-seq enables genome-wide profiling of off-target cleavage by
CRISPR-Cas nucleases. *Nat. Biotechnol.* **33**, 187–197 (2015).
52. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMB.journal* **17**,.
53. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).
54. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

55. Amos, B. *et al.* VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics
resource center. *Nucleic Acids Res.* **50**, D898–D911 (2022).
56. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version
7: Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).
57. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: Computing Large Minimum
Evolution Trees with Profiles instead of a Distance Matrix. *Mol. Biol. Evol.* **26**, 1641–
1650 (2009).
58. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately Maximum-
Likelihood Trees for Large Alignments. *PLoS ONE* **5**, e9490 (2010).
59. Markin, A. *et al.* PARNAS: Objectively Selecting the Most Representative Taxa on a
Phylogeny. *Syst. Biol.* **72**, 1052–1063 (2023).
60. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*
Methods **9**, 676–682 (2012).
61. Glover, L. & Horn, D. Trypanosomal histone γH2A and the DNA damage response.
Mol. Biochem. Parasitol. **183**, 78–83 (2012).
62. Teufel, F. *et al.* SignalP 6.0 predicts all five types of signal peptides using protein
language models. *Nat. Biotechnol.* **40**, 1023–1025 (2022).
63. Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput. Biol.* **7**, e1002195
(2011).
64. Meng, E. C. *et al.* UCSF CHIMERAX : Tools for structure building and analysis.
Protein Sci. **32**, e4792 (2023).
931
932

933 **Supplemental figures**

A Target:AGAAGCAGCCGCGACACTGTTAATTTACGCCACGCACAAAATACAAGAC
   Read: AGAAGCAGCCGCGACACTGTTAATTTACGTCAGGATGAAAGTGGAAGCA
   Donor: GCAAGAAGCCGGAACGCTGTTAATTTACGTCAGGATGAAAGTGGAAGCA

Supplemental Figure 1. Mosaic recombination site
A) A schematic of a recombination site. The region in yellow is the putative recombination site, shared by all 3 sequences.

935
936

Supplemental Figure 2. Cas9 system and sgRNA design and analysis
A) A schematic of the Cas9 induction experiment. Transient electroporation shown on the left while constitutive guide expression to isolate mosaic clones shown at right. B) An immunoblot showing FLAG-tagged Cas9 induction 24 hours following doxycycline treatment (DOX). DMSO was used as a vehicle control. EF1α was used as a loading control. C) A histogram showing the midpoint of all possible recombination sites 5bps or longer between AnTat1.1 and its family members. D) Quantification of the number of recombination sites within 250bps up or downstream of the cut site for each donor VSG. E) Quantification of the Levenshtein distance between AnTat1.1 and family members. This includes mismatches, insertions and deletions. F) Quantification of the number of mismatches at the guide binding site between AnTat1.1 and family members. G) Histogram of which donor VSGs can disrupt the PAM when used to repair AnTat1.1. H) An immunoblot of γ-H2A from doxycycline induced clones and uninduced controls. DMSO was used as a vehicle control. Samples were normalized to EF1α loading and γ-H2A induction following doxycycline treatment after 24 hours was quantified at right.

Supplemental Figure 3. DNA breaks result in switchers if a homologous donor is not present
A) A schematic showing identified switchers following a Cas9-induced double strand break within VSG-2 at locations 707 and 1082. B) Histograms of donor VSG insertion lengths identified in all mosaic VSG reads from Figure 2A). The insert length only includes newly inserted sequence and does not include recombination sites. Clone C1 and C2 sequenced via VSG-AMP-seq are shown separately. The limit of detection for an insertion is approximately 200 bp. C) A histogram with a summary of the unique recombination sites found within guide-induced break for each clone. Cut sites are indicated above the histograms as gaps within the black line. The midpoint of the perfect homology between AnTat1.1 and the donor VSG at the recombination site is plotted. If a donor VSG could not be unambiguously identified, the average recombination position was plotted. The legend for the donor VSG colors is to the right. SL = 5′ splice leader cap, 14-mer = 3′ sequence conserved in all VSG transcripts

938
939

Supplemental Figure 4. Mouse infection extended data
A) Quantification of the number of recombination events detected per mouse. Each mouse was normalized to the number of total consolidated, aligned, and unanchored reads compared to the consolidated, aligned, unanchored read count for one mouse to control for sequencing depth. This was performed separately for each genotype. B) A time course of the parasitemia for the mouse infections. Blood was harvested every two days. C) Quantification of the percent of parasites expressing the starting VSG at D6 and D15 post-infection as quantified by VSG-seq.

940
941

 Supplemental Table 1

Guide Primers	
Primer Name	Sequence
AnTat1.1_243	GAAATTAATACGACTCACTATAGGATTCAAAAACGGCCAAAC GCCGTTTTAGAGCTAGAAATAGC
AnTat1.1_369R	GAAATTAATACGACTCACTATAGGGGGCGTAAATTAACAGTGT CGGTTTTAGAGCTAGAAATAGC
AnTat1.1_694	GAAATTAATACGACTCACTATAGGAGTACAGACCCAGAAGCC AGGTTTTAGAGCTAGAAATAGC
AnTat1.1_894	GAAATTAATACGACTCACTATAGGACGCCGGTGTCGCAGCT AAACGTTTTAGAGCTAGAAATAGC
AnTat1.1_978R	GAAATTAATACGACTCACTATAGGGTCGTTGGCTGCTTGGAG TTGTTTTAGAGCTAGAAATAGC
AnTat1.1_1459	GAAATTAATACGACTCACTATAGGACCAATCCAGAAAAGTGC AAGTTTTAGAGCTAGAAATAGC
G00	AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGA CTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC

Annealed Guides for T7-sgRNA insertion		
Fragment Name	FWD	REV
AnTat1.1_243	AGGGATTCAAAAACGGCCAA ACGCC	AAACGGCGTTTGGCCGTTTT TGAAT
AnTat1.1_694	AGGGAGTACAGACCCAGAA GCCAG	AAACCTGGCTTCTGGGTCTG TACT
VSG-2_707	AGGGACCAACGGCCTCGGC AAAAG	AAACCTTTTGCCGAGGCCGT TGGT
VSG-2_1082	AGGGCCAGTGGCGCAAAAC CTGGT	AAACACCAGGTTTTGCGCCA CTGG

DNase Verification Primers		
Primer Name	FWD	REV
HSP-70 (Tb927.11.11330)	AGAACACTATCAATGACCCC AAC	CCATGCCCTGGTACATCT
Hyg	ACAGCGGTCATTGACTGGA G	ATTTGTGTACGCCCGACAGT

VSG-AMP-seq Primers	
Primer Name	Sequence
All-VSG-3'UTR	GTGTTAAAATATATC
Y-adapter[%]	[Phos]GATCGGAAGAGC*C*A
A01^	AATGATACGGCGACCACCGAGATCTACACTAGATCGC(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGA TC*T
A03^	AATGATACGGCGACCACCGAGATCTACACTATCCTCT(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGA

	TC*T
A04^	AATGATACGGCGACCACCGAGATCTACACAGAGTAGA(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGATC*T
A05^	AATGATACGGCGACCACCGAGATCTACACGTAAGGAG(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGATC*T
A06^	AATGATACGGCGACCACCGAGATCTACACACTGCATA(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGATC*T
A07^	AATGATACGGCGACCACCGAGATCTACACAAGGAGTA(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGATC*T
A08^	AATGATACGGCGACCACCGAGATCTACACCTAAGCCT(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGATC*T
A09^	AATGATACGGCGACCACCGAGATCTACACGACATTGT(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGATC*T
A10^	AATGATACGGCGACCACCGAGATCTACACACTGATGG(N:252 52525)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)(N)ACACTCTTTCCCTACACGACGCTCTTCCGATC*T
P5_2	AATGATACGGCGACCACCGAGATCTACAC
P701	CAAGCAGAAGACGGCATACGAGATTCGCCTTAGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
P703	CAAGCAGAAGACGGCATACGAGATTTCTGCCTGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
P704	CAAGCAGAAGACGGCATACGAGATGCTCAGGAGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
P705	CAAGCAGAAGACGGCATACGAGATAGGAGTCCGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
P706	CAAGCAGAAGACGGCATACGAGATCATGCCTAGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
P707	CAAGCAGAAGACGGCATACGAGATGTAGAGAGGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
P708	CAAGCAGAAGACGGCATACGAGATCCTCTCTGGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
P709	CAAGCAGAAGACGGCATACGAGATATCACGGTGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA

P710	CAAGCAGAAGACGGCATACGAGATCGATGTGCGTGACTGGAGTCCTCTCTATGGGCAGTCGGTGA
AnTat1.1_1F[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$CGCAAACACTACAACGAGCC
AnTat1.1_2F[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$CAGAATGCGACACGGAAAGC
AnTat1.1_3F[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$ACGCAGGCGGCTTCAAAACA
AnTat1.1_4F[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$AACAGCCGCAGCAACCAAAC
AnTat1.1_0R[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$GGCCACAAATGCGGCAGAAAC
AnTat1.1_1R[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$GCCATAAGCTGCGGTTTCGT
AnTat1.1_2R[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$GTTGTGTATGGTTAGCAGGC
AnTat1.1_3R[&]	CCTCTCTATGGGCAGTCGGTGAT$(N)_{0-7}$CTTGTATTTTGTGCGTGGCG
Index1[%]	ATCACCGACTGCCCATAGAGAGGACTCCAGTCAC
Read2[%]	GTGACTGGAGTCCTCTCTATGGGCAGTCGGTGAT
Miscellaneous Additional Primers	
Name	Sequence
SL-FWD	ACAGTTTCTGTACTATATTG
SP6-14mer-REV	GATTTAGGTGACACTATAGTGTTAAAATATATC
AnTat1.1 Sanger Sequencing Primer	AGAGAATACTAAGCTAGTTGGC
Pan AnTat1.1family FWD	ACTACACCCACAACAAGCTCTA

943 * = indicates a phosphorotiolate bond modification
944 [Phos] = 5' phosphorylation
945 ^ = hand mixing
946 & = machine mixing
947 % = HPLC purification

A

Mosaic VSG

Fragmentation

Adapter ligation

Universal adapter primer

Target specific PCR

Target specific primer

Final library

Index2
Read1   Read2

P5   UMI   Donor   Target   Index1   P7

B   AnTat1.1-specific
VSG-AMP-seq primers

0   400   800   1200   1600
14-mer

SL   N-term   C-term

A Target: AnTat1.1 (EATRO)

unique recombinations

WT

μMT

B WT   μMT

donor VSGs

Experiment
- Expt1
- Expt2
- Expt3

N   C   N   C
Domain

C # recombinations from assembled ORFs

WT blood

WT tissues

Position within AnTat1.1 transcript (bp)

SL— N-term   C-term   14-mer

D

recombinations

D6   D10   D14
Day

- Blood
- Tissues

Legend
- VSG-228
- VSG-2986
- VSG-3110
- VSG-7358
- VSG-4156
- AnTat1.1 family
- Other VSGs

E

recombination sites

Homology length (bp)

F

inserts

Insert length (bp)

G WT tissue mosaic VSGs

Perfect sequence match to AnTat1.1

A

Mosaic 1

Donor VSG: VSG-3110

S(204)A +(205)G

Mosaic 2

Donor VSG: VSG-2986

S(204)G   A(207)S   S(208)N

Mosaic 3

Donor VSG: VSG-2986

S(204)G   A(207)S   S(208)N
T(217)S   P(227)T   T(234)I

Mosaic 4

Donor VSG: VSG-2986

S(204)G   A(207)S   S(208)N
T(217)S   P(227)T   T(234)I
A(238)E   S(243)T   S(246)T
L(247)I   Q(248)K   P(252)T
                 T(253)A

Staining intensity

Alexa Fluor 647

B

Percent AnTat1.1 staining

ns
***
***
***

Parental   1   2   3   4
AnTat1.1   Mosaics

C

Overlapped models of AnTat1.1
and mosaic VSGs

D

Side-view

Model of AnTat1.1
with altered resides colored
by donor VSG

Top-view

VSG-3110

VSG-2986

3
2   # mosaics
1   altered