1 **Identifying *in vivo* genetic dependencies of melanocyte and melanoma**
2 **development**
3

4 Sarah Perlee[1,2], Yilun Ma[1,3,4], Miranda V. Hunter[1], Jacob B. Swanson[5,6], Nelly M. Cruz[1],
5 Zhitao Ming[7], Julia Xia[1], Timothée Lionnet[5,6,8], Maura McGrail[7], Richard M. White[1,9, *]
6
[1]Department of Cancer Biology and Genetics, Memorial Sloan Kettering Cancer Center,
New York, NY, USA
[2]Gerstner Sloan Kettering Graduate School of Biomedical Sciences, Memorial Sloan
Kettering Cancer Center, New York, NY, USA.
[3]Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program, Memorial
Sloan Kettering Cancer Center, New York, NY, USA.
[4]Cell and Developmental Biology Program, Weill Cornell Graduate School of Medical
Sciences, New York, NY, USA.
[5]Institute for Systems Genetics, NYU Grossman School of Medicine, New York, NY,
USA.
[6]Department of Cell Biology, NYU Grossman School of Medicine, New York, NY, USA.
[7]Department of Genetics, Development and Cell Biology, Iowa State University, IA,
USA.
[8]Department of Biomedical Engineering, NYU Tandon School of Engineering, Brooklyn,
NY.
[9]Nuffield Department of Medicine, Ludwig Institute for Cancer Research, University of
Oxford, Oxford, UK.
[*] Correspondence to Richard M. White: Richard.white@ludwig.ox.ac.uk

25
26
Abstract

The advent of large-scale sequencing in both development and disease has identified
large numbers of candidate genes that may be linked to important phenotypes. We
have developed a rapid, scalable system for assessing the role of candidate genes
using zebrafish. We generated transgenic zebrafish in which Cas9 was knocked-in to
the endogenous *mitfa* locus, a master transcription factor of the melanocyte lineage.
The main advantage of this system compared to existing techniques is maintenance of
endogenous regulatory elements. We used this system to identify both cell-autonomous
and non-cell autonomous regulators of normal melanocyte development. We then
applied this to the melanoma setting to demonstrate that loss of genes required for
melanocyte survival can paradoxically promote more aggressive phenotypes,
highlighting that *in vitro* screens can mask *in vivo* phenotypes. Our genetic approach
offers a versatile tool for exploring developmental processes and disease mechanisms
that can readily be applied to other cell lineages.

41
Introduction:
Cell type-specific knockout of genes is crucial for unraveling the intricate molecular
mechanisms underlying cell function, development, and transformation. Efficient
methods are critical as sequencing projects such as All of Us, the UK Biobank and the
Cancer DepMap continue to identify hundreds of new candidate genes on a regular

basis, many of which have yet to be characterized *in vivo*[1-3]. Zebrafish (*Danio rerio*)
have emerged as a powerful genetic model organism for studying these processes. The
large number of animals that can be easily generated make it advantageous (compared
to other systems like mice) for querying large numbers of genetic variants that may
have links to human diseases. However, most studies in this system use germline
knockout, which makes it difficult to determine whether the identified gene has a specific
function in a particular lineage, or whether the phenotype reflects a more general
developmental defect. Furthermore, many genes which might have lineage specific
functions cannot be studied using germline knockout, since embryonic lethality often
accompanies global knockout of essential genes[4]. To overcome these limitations,
several groups have explored tissue-specific and conditional knockout strategies in
zebrafish[5-13]. For example, the MAZERATI system uses tissue-specific promoter
transgenes to drive Cas9 expression, which are introduced via plasmid injection into
one-cell stage embryos[6,14]. This system has been shown to be versatile and powerful.
One potential limitation of promoter fragment transgenes is that these fragments could
lack certain regulatory regions. For this reason, it is common in systems such as mice to
knock in genetic cassettes (i.e. Cre recombinase) into endogenous loci.
64
In this study, we wished to develop such an endogenous system to understand *in vivo*
genetic dependencies, using melanocytes and melanoma as an exemplar. Skin color
pigmentation is amongst the most heterogeneous of genetic systems, with a large
number of loci linked to melanocyte development. Mutations in pleiomorphic genes such
as *sox10, ednrb, jam3b, tuba8l3, meox1* and *kit* yield clear defects in
melanocyte/melanophore development, but these mutants also have defects in other
tissues[15-21]. Many of these genes are also expressed in melanoma, but their specific *in*
vivo function in tumorigenesis remains poorly understood. By integrating the Cas9
nuclease into the endogenous melanocyte-specific *mitfa* locus, we achieved specific
gene disruption across the melanocyte developmental lineage. We used our *mitfa*[Cas9]
knockin animals to inactivate the pigmentation gene *albino* and the embryonic essential
genes *sox10, tuba1a,* and *ptena/b* in melanocytes without additional developmental or
off-target phenotypes. We also demonstrated that this system can be used to induce
melanomas in wild-type fish by inactivating the tumor suppressors *tp53* and *ptena/b*
within melanocytes also expressing the melanoma oncogene BRAF[V600E]. Inactivating
the neural crest transcription factor *sox10* in these tumors significantly decreased
melanoma initiation but resulted in rare invasive tumors that highly expressed the *sox9*
transcription factor, highlighting that unexpected *in vivo* phenotypes can emerge that
aren't predicted from *in vitro* studies[22,23]. Our system can thus be used to advance our
understanding of melanocyte biology and the genetic underpinnings of phenotypic
switching in melanoma, an approach that can readily be used for other cell types.
86
Results:
Targeted integration of Cas9 to the *mitfa* locus
To selectively disrupt genes in a melanocyte lineage-specific manner while retaining
natural regulatory elements, we established a transgenic zebrafish line harboring Cas9
at the endogenous *mitfa* locus. *Mitfa* expression is detectable by 18 hours
postfertilization (hpf) and is highly expressed in pigment cells including melanocytes,

xanthophores and their progenitors. To achieve knock-in, we utilized the GeneWeld
method, which enables efficient integration of DNA cassettes using homology mediated
end joining (HMEJ)[24,25]. The knock-in cassette, containing Cas9 and BFP driven by the
eye-specific γ-crystallin promoter, was targeted to exon 2 of the *mitfa* gene, which leads
to disruption of the endogenous *mitfa* gene (Figure 1A). We strategically selected this
genomic site for several reasons. First, this is the sole exon shared among the three
mitfa protein-coding transcripts. Furthermore, targeting Cas9 to exon 2 offers the
flexibility of replicating this knock-in strategy in *casper* zebrafish, a transparent fish
widely used for melanoma studies that harbor a premature stop codon in *mitfa* exon 3[26].
102
The knock-in vector, along with Cas9 protein and guide RNAs targeting either the *mitfa*
integration site or the sequences flanking the 5' and 3' homology arms, were injected
into 1-cell stage wild-type Tropical 5D (T5D) zebrafish embryos (Figure 1B)[27]. An
average of 19% of injected embryos had detectable ocular BFP expression at 3 days
postfertilization (dpf), indicating successful integration (Figure S1A). Embryos were
raised to adulthood then crossed with WT T5D fish. The resulting F1 embryos were
screened for BFP+ eyes and tail clippings were sequenced to confirm precise
integration of the knock-in cassette at the *mitfa* locus. We sequenced four F1 fish and
detected on-target integration at the *mitfa* locus in all four fish (Figure S1B). In
zebrafish, one copy of *mitfa* is sufficient for the normal development of neural crest-
derived melanocytes, but loss of both copies of *mitfa* results in loss of the melanocyte
lineage[26,28]. To determine the effect of a single copy of *mitfa*^{Cas9} on melanocytes,
mitfa^{Cas9} fish were in-crossed and the resulting siblings were imaged at various time
points (Figure S1C). WT and *mitfa*^{Cas9} fish exhibited indistinguishable melanocyte
patterning during both embryonic and adult stages. However, as expected, fish carrying
two copies of Cas9 (*mitfa*$^{Cas9/Cas9}$) resembled the *nacre* mutant (*mitfa*-/-), characterized
by complete loss of melanocytes, indicating the endogenous *mitfa* gene is disrupted on
knock-in alleles[28]. We used zebrafish harboring one knock-in allele (*mitfa*^{Cas9}) for all
subsequent experiments to allow us to test gene function in melanocytes.
122
To determine whether Cas9 expression was restricted to melanocytes, we performed
fluorescence in situ hybridization (FISH) on 3dpf WT and *mitfa*^{Cas9} embryos to assess
mitfa and Cas9 mRNA expression (Figure 1C and S1D). As expected, *mitfa* was
detected within the embryonic melanocyte stripe regions in both WT and *mitfa*^{Cas9} fish.
In *mitfa*^{Cas9} embryos, Cas9 expression was observed in an average of 86% of these
mitfa+ cells (Figure S1D). These results provide functional validation for on-target
integration and melanocyte lineage-restricted expression of Cas9.
130
Melanocyte-specific loss of the human skin color associated gene *slc45a2* results
in robust loss of pigmentation
To assess the effectiveness of our stable *mitfa*^{Cas9} line, we first targeted the *albino* gene
(*slc45a2*). Large scale GWAS studies have repeatedly identified SNPs in this gene as
tightly linked to human skin color variation, and germline disruption leads to a complete
loss of pigmentation in melanocytes[29-31]. This distinct phenotype enables the visual
observation of Cas9 activity in zebrafish injected with *albino* gRNA. To test our system,
we designed plasmids, denoted MG-gRNA, containing a zU6:gRNA cassette followed

by *mitfa*:GFP to enable visualization of cells expressing the *albino* gRNA. MG-*albino*
and MG-NT were injected into one-cell stage embryos from crosses between *mitfa*^Cas9
and WT fish, which express Cas9 in all melanocytes but exhibit mosaic gRNA
expression (Figure 2A).
143
Embryos were screened for BFP+ eyes to identify those harboring *mitfa*^Cas9. Upon
adulthood, 92% of F0 MG-*albino* fish had mosaic loss of pigmentation within the
melanocyte stripe regions (Figure 2B-C). Unpigmented regions maintained GFP
expression (Figure 2B), indicating that the breaks in melanocyte stripes arise from loss
of pigmentation in melanocytes rather than loss of the melanocyte lineage (Figure 2B).
As expected, control MG-NT clutch mates displayed normal pigmentation and GFP+
melanocytes were indistinguishable from GFP- regions on the same fish (Figure 2B-C).
151
We next aimed to determine the efficiency of *mitfa*^Cas9 in the F1 generation, in which
every melanocyte stably expresses the *albino* gRNA and *mitfa*^Cas9. To generate F1
lines, we outcrossed the F0 MG-gRNA fish to WT and sorted embryos for BFP+ eyes
and GFP+ melanocytes. F1 MG-*albino* fish exhibited a near complete loss of
pigmentation, demonstrating that our *mitfa*^Cas9 system is highly efficient in F1 fish
(Figure 2D-E). To study the effect of *albino* inactivation across various stages of
development, F1 MG-gRNA fish were once again outcrossed to WT, generating F2 MG-
gRNA fish expressing one copy of *mitfa*^Cas9. Imaging revealed a measurable
pigmentation loss in F2 MG-*albino* melanocytes as early as 3 days post-fertilization
(dpf) (Figure 2F-G). F2 MG-*albino* fish maintained loss of pigmentation throughout
development, indicating the sustained activity of *mitfa*^Cas9 (Figure S2A-D).
163
To confirm on-target cutting of the *albino* gene in the melanocyte lineage (and not other
tissues), the skin from F1 MG-*albino* fish was dissected and GFP+ and GFP- cells were
isolated using FACS (Figure 2H). GFP+ cells encompass *mitfa*-expressing
melanocytes, xanthophores, and their progenitors, while GFP- cells represent other
non-melanocytic, non-*mitfa* expressing skin cells, such as keratinocytes, fibroblasts, and
immune cells. We performed CRISPR-seq on DNA from both populations to directly
measure Cas9 efficiency and cell-type specificity, quantifying the proportion of indels at
the *albino* locus. Although we have previously shown that FACS sorting melanocytic
cells is possible, it is important to note that we often observe high levels of
contaminating keratinocytes in FACS sorted melanocyte populations, which may lead to
an underestimation of the true allelic frequency in our CRISPR-seq[32]. Despite this,
robust inactivation of the *albino* gene was observed in GFP+ (*mitfa*-expressing) cells,
confirming somatic inactivation (Figure 2I). The majority of indels were frame-shift
inducing mutations. A likely PCR or sequencing artifact resulting in a "T" insertion was
excluded from the frameshift indel percentage as it was observed in all conditions
including WT fish with no pigmentation phenotype (Figure S2E). These results confirm
that our *mitfa*^Cas9 system allows us to inactivate genes within the melanocytic lineage *in*
vivo in a cell-type specific manner.
182
The neural crest-related gene *sox10* has specific function in adult melanocyte
patterning and regeneration

Having confirmed that our *mitfa*[Cas9] system allows us to manipulate melanocyte gene
expression, we next wanted to target embryonic essential genes. Detecting "negative"
gene knockout phenotypes such as cell death can be challenging *in vivo* due to the
selective advantage that WT cells have over mutants.
189
To assess the effectiveness of our *mitfa*[Cas9] model in detecting negative phenotypes, we
utilized it to target *sox10*, a transcription factor vital to the neural crest lineage and
indispensable for overall survival of both zebrafish and mice[15,33]. Consequently, a global
knockout approach is not feasible for studying the function of *sox10* in adult
melanocytes *in vivo*. Germline knockout of *sox10* in zebrafish leads to severe defects in
peripheral nerve development and the complete absence of melanocytes, and the
animals die around day 14[15,34,35]. Due to the complete absence of melanocytes and
their precursor cells, melanoblasts, in *sox10*[-/-] fish, it has not been possible to study the
specific role of *sox10* on these more differentiated cell types *in vivo*. One of the key
unanswered questions is how *sox10* balances its role in maintaining melanoblast
stemness with promoting terminal differentiation, particularly in the context of adult
melanocyte regeneration.
202
We leveraged our *mitfa*[Cas9] system to specifically inactivate *sox10* in the melanocyte
lineage. *mitfa*[Cas9] fish injected with a MG-*sox10* plasmid exhibited noticeable gaps in
their melanocyte stripes (Figure 3A-B). Germline deletion of a *sox10* enhancer region
similarly affects adult stripe patterning in zebrafish[36]. In contrast to the *albino* F0 fish,
the gaps in the stripes of the *sox10* F0 fish were not occupied by unpigmented GFP+
melanocytes, indicating a potential defect in the survival or differentiation of
melanocytes upon loss of *sox10*. This phenotype became even more pronounced in the
stable F1 lines, where several gaps in the melanocyte stripes are visible (Figure 3C).
Zebrafish have four dark stripes occupied by melanocytes, which alternate with light
colored interstripes occupied by yellow pigmented xanthophores[37]. Typically,
melanocyte stripe 1D (dorsal) is wider than the interstripe X0 directly below it, as
observed in MG-NT F1 fish (Figure 3C-D). However, the opposite is true for MG-*sox10*
F1 fish, where we observed an apparent widening of the interstripe region and
narrowing of the melanocyte stripes (Figure 3C-D). Melanocytes were counted in fish
treated with epinephrine to aggregate melanosomes, revealing a significant reduction in
the number of melanocytes in the F1 *sox10* KO fish compared to F1 NT fish (Figure
3E). Reduction in GFP+ cells within the xanthophore interstripes (X0) suggested that
sox10 deficiency also negatively affected xanthophore populations, in line with previous
work indicating a role for *sox10* in specifying both xanthophores and melanocytes in
zebrafish embryos[38].
223
Melanocytes continually regenerate throughout life, typically from a progenitor
population of melanoblasts or melanocyte stem cells. We next assessed whether *sox10*
was required for this. We treated adult fish with neocuproine, a copper chelator
previously shown to kill mature, pigmented melanocytes[39]. We then compared
regeneration from these progenitors in MG-*sox10* and MG-NT F1 fish by counting the
percentage of melanocytes that emerged after 7, 15, and 70 days (Figure 3F). At days 7
and 15, when the melanocytes were still in the process of regenerating from

melanoblasts, there were no significant differences between the two groups. However,
by the time regeneration was completed we found a significant disparity between the
regenerative potential of MG-NT and MG-*sox10* fish. At 70 days post-neocuproine
treatment, MG-NT fish had regenerated an average of 82% of their melanocytes,
whereas only 53% of melanocytes had regenerated in MG-*sox10* fish (Figures 3G-H).
These findings demonstrate that *sox10* is required for adult melanocyte regeneration,
highlighting its requirement within melanoblast populations outside of neural crest
specification.
239
Non-autonomous functions of *tuba1a/tuba1c* on melanocytes
One of the significant challenges of conventional global knockout methods is the
inability to differentiate between cell-autonomous and non-autonomous phenotypes.
This distinction is particularly crucial in melanocyte studies due to their direct and
indirect interactions with diverse cell types. A notable example of genes with pleiotropic
effects is the tubulin gene family, whose α/β tubulin heterodimers form microtubules
essential for various cellular processes such as cell division, motility, and intracellular
transport across various cell types[40]. Microtubules are also critical for the transport of
melanosomes in melanocytic cells[41].
249
Mutation of tubulin genes such as *tuba8l3a* in zebrafish leads to highly pleiotropic
effects including defects in melanocyte patterning, CNS abnormalities, and altered
craniofacial morphology[42]. We chose to focus on the closely related α tubulin gene
tuba1a, which is highly expressed in many cell types of the developing zebrafish
including melanocytes, but whose specific function in these cells remains unexplored[43].
Non-cell type specific knockout of *tuba1a* with the Alt-R CRISPR Cas9 system results in
extensive embryo abnormalities including a curved tail phenotype, pericardial edema,
and melanocytes with more dispersed pigmentation (Figure 4A-B, S3A-B). We found
that compared to 71% survival of NT Alt-R embryos, only 10% of embryos injected with
tuba1a Alt-R survive to 14dpf (Figure 4C). During validation of *tuba1a* Alt-R, we
discovered that this guide also targets *tuba1c*, a gene with 99.78% homology to *tuba1a*
in zebrafish (Figure S3B). Despite robust cutting, neither *tuba1a*- nor *tuba1c*-specific
guides alone were sufficient to recreate the dispersed melanocyte phenotype (Figure
S3C-D). However, we were able to recapitulate our findings using a second guide RNA
that simultaneously targeted both *tuba1a* and *tuba1c*, suggesting possible redundancy
of these genes, with compensation masking the phenotype in single-gene knockouts
(Figure S4E-F).
267
Similar to *sox10,* the embryonic lethality of global *tuba1a/c* knockout prevents the study
of knockout phenotypes in adult fish. To overcome this limitation, we implemented our
mitfa[Cas9] system to specifically knock out *tuba1a/c* in melanocytes (Figure S3G-H).
CRISPR sequencing of GFP+ cells isolated from *mitfaCas9* MG-*tuba1a/c* fish confirmed
successful knockout of both *tuba1a* and *tuba1c*, with knockout efficiencies comparable
to those observed in MG-albino fish (Figure S3I). Unlike global knockout of *tuba1a/c*,
melanocyte specific loss of *tuba1a/c* resulted in viable adult fish with no obvious
abnormalities (Figure S3G-H). Surprisingly, we did not observe the dispersed
melanocyte phenotype in any *mitfa*[Cas9] MG-*tuba1a/c* embryos (Figure 4D-E). Based on

this observation, we hypothesized that *tuba1a/c* may be functioning in a cell non-
autonomous manner on melanocytes.
279
The dispersed pigmentation observed in *tuba1a/c* Alt-R embryos resembles that seen in
blind zebrafish, which fail to contract their melanosomes in response to light[44]. This
light-mediated camouflage involves retinal ganglion cells signaling hypothalamic
neurons to release melanin-concentrating hormone (MCH), which then binds to MCH
receptors on melanocytes, triggering melanosome aggregation via microtubules[44]. The
normal melanosome dispersion observed in *mitfa*[Cas9] MG-*tuba1a/c* fish suggests that
rather than a cell-intrinsic defect in melanosome aggregation, *tuba1a/c* may indirectly
influence melanocytes through potential defects in retinal and/or central nervous system
cells crucial for light-mediated camouflage. Interestingly, *TUBA1A* mutation in humans
can result in both ophthalmologic and brain abnormalities and morpholino-based
knockdown of *tuba1a* in zebrafish inhibits CNS development[45,46].
291
To further test this idea, we treated *tuba1a/c* Alt-R embryos with epinephrine to assess
their ability to contract melanosomes in response to a non-visual stimulus. *Tuba1a/c* Alt-
R embryos exposed to epinephrine had robust aggregation of melanosomes,
demonstrating that the loss of *tuba1a/c* did not impair melanosome transport (Figure 4F-
G). This finding further supports a non-autonomous role for *tuba1a/c* in melanocytes.
Utilizing the *mitfa*[Cas9] system distinguish cell-autonomous from non-autonomous gene
functions provides valuable insights into the complex dynamics of cell-cell interactions
and how genes function within cellular and organismal networks.
300
Targeting tumor suppressors with *mitfa*[Cas9] induces melanoma
We next turned our attention to melanoma. In humans, tumor suppressors such as
PTEN and *TP53* are somatically inactivated in melanoma[47,48]. In contrast, the most
commonly used zebrafish models of melanoma use a germline *tp53* mutation, which is
not typically seen in humans with the disease and precludes our ability to discern the
role of *tp53* inactivation specifically in melanocytes and melanoma[49]. While global *tp53*
loss is not lethal, it does lead to a wide range of non-melanoma tumors, which can
deteriorate fish health and confound melanoma studies[50]. Furthermore, current
zebrafish melanoma models typically involve the utilization of MiniCoopR *mitfa* rescue
cassettes into *casper* or *nacre* zebrafish, which are devoid of normal pigment cells.
Although this method is highly efficient, it does not accurately represent melanoma
initiation in the context of normal skin architecture and relies on artificial overexpression
of *mitfa*. To mimic human melanoma initiation more accurately, we used our *mitfa*[Cas9]
model to assess tumor initiation using melanoma-specific gene knockout in animals with
wild-type skin. Previous studies have shown that expression of BRAF[V600E] and loss of
tp53 in WT (non-*casper*) fish results in relatively low tumor burden, suggesting that the
casper strain might be especially tumor prone in the setting of MiniCoopR *mitfa* rescue
although the exact reasons for this are not clear. Based on this, to increase penetrance,
we decided to also target *ptena* and *ptenb*, the zebrafish orthologs of human *PTEN*[51].
Ptena/ptenb loss in combination with *tp53* has been shown to accelerate melanoma
formation in zebrafish[52,53].
322

To first address the effect of *ptena/b* loss on normal melanocytes, we generated melanocyte-specific knockouts of *ptena* and *ptenb* (Figure 5A). To visualize cells expressing both *ptena* and *ptenb* guides, we designed additional plasmids, denoted MTdT-gRNA, containing a zU6:gRNA cassette followed by *mitfa*:TdTomato and coinjected MG-*ptena* and MTdT-*ptenb* plasmids into *mitfa*[Cas9] embryos. Like *sox10* and *tuba1a/c*, germline knockout of *ptena/b* is embryonic lethal[54,55]. In our *ptena/ptenb* F0 melanocyte-specific KO fish, survival was normal. We observed an aberrant expansion of the melanocytes outside of the stripe regions they normally are restrained to, suggesting that inactivation of *ptena* and *ptenb* induces defects in melanocyte patterning (Figure 5A and B). However, loss of *ptena/b* alone was not sufficient to induce melanoma.

To generate tumors, we co-injected *mitfa*[Cas9] embryos with plasmids encoding *mitfa:*BRAF[V600E] and sgRNAs targeting *tp53* and *ptena/b* (Figure 5C). Larvae were sorted for BFP+ eyes, indicating mitfa-Cas9 integration, and monitored for tumors over 30 weeks. No tumors were detected in BFP- fish from any of the injection conditions (n=136). We also did not observe any tumors in BFP+ fish injected with either BRAF[V600E] alone (n=35) or ptena/b; *tp53* guides in the absence of BRAF[V600E] (n=17). In contrast, tumors developed in all BFP+ groups injected with both *mitfa:*BRAF[V600E] and gRNA plasmids. Approximately 5% of fish from the BRAF;*tp53* group developed tumors (Figure 5D). This aligns with a previous study where 6% of *tp53*-/- fish developed tumors when injected with a BRAF[V600E] construct[51]. Targeting of *ptena* and *ptenb* in *mitfa*[Cas9] fish co-injected with *mitfa*:BRAF[V600E] resulted in a higher tumor incidence of 29% (Figure 5D).

When all three tumor suppressors (*tp53, ptena, ptenb*) were targeted in combination with BRAF[V600E], 59% of fish developed tumors by 30 weeks post fertilization (wpf) (Figure 5D). Immunohistochemistry (IHC) revealed that all tumor samples expressed BRAF[V600E], while p-AKT, a hallmark of *PTEN* inactivation, was only detected in samples with *ptena/b* KO (Figure 5E). To confirm on-target cutting, DNA was extracted from tumors and CRISPR sequencing was performed on PCR amplicons for each tumor suppressor gene. A large proportion of reads contained indels at the target locus for each of the target genes (*ptena*, *ptenb*, and *tp53*) (Figure 5F and S4A-C). Some tumors had an array of mutations, while others appeared to be composed largely of a one dominant clone (Figure S4A-C). Thus, similar to the MAZERATI system, our endogenous knockin system allows us to rapidly dissect the melanocyte-specific function of both oncogenes (BRAF[V600E]), as well as putative tumor suppressors (*tp53, ptena/b*), while avoiding pleiotropic effects of global tumor suppressor loss.

Targeting of lineage-specific oncogenes decreases tumor initiation but promotes progression

Our next objective was to leverage the mitfa[Cas9] fish as a tool for identifying or validating potential genetic dependencies within melanoma. Genetic dependencies can be difficult to study *in vivo* since inactivation of a required gene inevitably leads to selection for non-mutant alleles, as has been previously shown[56]. Large scale in vitro efforts such as the DepMap can provide guidance as to what genes are likely to be important for tumor

proliferation, but cell line studies alone cannot recapitulate the complex in vivo behavior
that occurs from knocking individual genes out. In melanoma, the DepMap has
identified *SOX10* as the top *in vitro* genetic dependency necessary for tumor cell
proliferation, suggesting it acts as a lineage-specific oncogene[3,57]. In prior work in both
zebrafish and mice, loss of *sox10* is clearly associated with a loss of tumor initiating
potential and cell proliferation[56,57]. Yet its specific role *in vivo* remains unclear, since it is
required for neural crest and melanocyte specification, making it difficult to know
whether its effects reflect loss of the entire lineage[57]. Given our data above showing that
sox10 is required specifically in melanocyte patterning and regeneration, we wished to
investigate its function in melanoma.
379
To assess this, we generated plasmids containing multiple sgRNAs driven by three
distinct zU6 promoters (zU6A, zU6B, and zU6C) to simultaneously target both tumor
suppressors (*tp53, ptena/ptenb*) and tumor promoting genes (*sox10*) in the same cells,
which reduces the chance of selecting for non-mutant alleles[5]. We co-injected these
plasmids with *mitfa*:BRAF[V600E], sorted fish for BFP+ eyes, and tracked tumor free
survival over the course of 50 weeks (Figure 6A-B). Consistent with the DepMap
prediction, *sox10* inactivation markedly reduced tumor incidence, which likely reflects
loss of proliferation. Only 3/96 (3.1%) fish with *sox10* knockouts initiated melanomas
compared to the 24/94 fish (25.5%) from the NT condition that developed tumors
(Figure 6B). Using IHC, we validated that *sox10* KO tumors expressed lower levels of
Sox10 protein compared to NT tumors (Figure 6C and S5A). Quantification of Sox10
intensity with immunofluorescence confirmed this finding (Figure S5B).
392
Although *sox10* clearly reduced tumor initiation, we noted "escapers". These tumors
appeared morphologically somewhat distinct compared to the *sox10* intact tumors.
While not easily quantifiable, qualitatively based on prior experience, the tumor cells
appeared more mesenchymal and penetrated more deeply under the skin than we
typically see with the BRAF[V600E] melanomas in the fish (Figure 6C). This observation
raised the possibility that these tumors had undergone "phenotype switching", a form of
cell plasticity in which cells reversibly move between opposite extremes of proliferation
versus invasive states[22,23,57,58]. In melanoma, the proliferative state is thought to be
characterized by high expression of *SOX10*, whereas the mesenchymal, invasive state
is characterized by high expression of *SOX9*[23]. These two highly related transcription
factors have seemingly opposite and possibly antagonistic roles in melanoma. *In vitro*,
knockout of *SOX10* in a variety of human cell lines is associated with acquisition of a
SOX9[hi] state, which has been suggested to be linked to the invasive phenotype[23]. We
performed a re-analysis of publicly available RNA-seq data using 3 patient-derived
human melanoma cell lines treated with siRNA targeting *SOX10* confirmed this
observation, with the *SOX10* lines now becoming *SOX9*[hi] (Figure 6E to J, S5C to L)[23].
These data raised the possibility that the more mesenchymal, invasive appearing
tumors we see in our escaper fish might be due to gain of *sox9* expression. To test this,
we stained our *sox10* CRISPR tumors for *sox9* and discovered that 2 of the 3 *sox10* KO
tumors markedly upregulated *sox9* relative to control tumors expressing NT gRNA
(Figure 6C and D). While this data is certainly not definitive, when taken in context of
the above *in vitro* and *in vivo* data in other model systems, it appears consistent with the

idea that loss of *sox10* could result in more invasive tumors. This preliminary
observation we have made will need further mechanistic dissection, and our system
would be amenable to studying this phenomenon in further depth. This data also
highlights that *in vitro* genetic studies such as the DepMap, which rely on proliferation
as the readout, can mask more complex *in vivo* phenotypes.
420
Discussion:
The ability to model genetic dependencies *in vivo* is one of the most important uses of
model organisms. Forward genetic mutagenesis screens using compounds such as
ENU or EMS led to the identification of many key developmental genes in organisms
such as Drosophila, C. elegans, Saccharomyces, zebrafish, and many others[59,60].
These approaches have rapidly accelerated in the era of TALENs and CRISPR, readily
allowing for knockout of nearly any gene in a wide variety of non-model organisms.
Cre/Lox approaches have further augmented our ability to discern the cell-type specific
effects of these genes, although these are still laborious and time consuming, especially
in models such as mice[13].
431
Alongside these approaches have been remarkable advances in large scale sequencing
of human tissues. GWAS-like efforts such as the UK Biobank and the US All of Us
programs have identified thousands of germline variants (SNPs) that may be linked to
interesting phenotype variation[1,2]. A conceptually similar approach is also happening in
diseases such as cancer, in which efforts such as the TCGA, ICGC and DepMap have
continued to unveil new potential somatic variants (SNVs) linked to cancer[3,61,62].
438
Across these efforts, in both development and disease, the large number of candidate
variants makes it challenging to connect these genotypes to phenotypes. This is
especially acute in melanoma, where the large number of somatic variants generated by
UV radiation leads to a high number of background mutations that may or may not have
pathogenic function. Thus, there is a substantial need for *in vivo* models that allow for
rapid and efficient modeling of candidate genes.
445
Prior work in the zebrafish has shown that promoter fragments driving Cas9 can provide
efficient and scalable approaches to this problem[6,14]. For example, MAZERATI has
been nicely applied to melanoma models to studying genetic dependencies in the
melanocyte lineage[14]. The technology we describe in this work largely builds off the
logic of this and related systems. One difference between our *mitfa*-Cas9 knockin line
compared to MAZERATI is that it maintains the endogenous regulatory elements
needed for control of the *mitfa* gene. We do not yet fully know when or if this difference
would be important, but given the complex role of MITF transcription as a "rheostat" in
melanoma[63], it is reasonable to assume these endogenous elements could become
important in some circumstances. Because in our study we did not directly compare the
mitfa-Cas9 knockin approach versus MAZERATI, we cannot make direct statements
about which is more efficient. However, because both techniques reliably result in
melanoma, each researcher will be able to choose the technology that is most
appropriate for their biological question.
460

One advantage of Cas9 approaches in this system is the ability to screen phenotypes in
both the F0 generation (as mosaics) or in the F1 generation (3 months later). Although
the F0 phenotypes we observe are indeed subtle, they are still quantifiable. This
enables rapid screening of candidate genes for melanocyte-specific functions.
Promising hits can then be more specifically validated in the F1 generation, which is
particularly advantageous for efficiently generating biallelic knockouts, especially for
genes with recessive phenotypes like *albino*. On a per gene basis, this saves well over
3 months of work compared to germline recessive knockouts and eliminates the issue of
inbreeding, promoting healthier genetic lines and facilitating the potential for outcrossing
with any available fish line. Because we demonstrate that the Cas9 is completely
restricted to the melanocyte lineage, with no detectable off-target expression, this easily
allows us to discern the melanocyte-specific effect of pleiotropic genes very efficiently.
473
One interesting observation of our melanoma studies is evidence that loss of melanoma
genetic dependencies like *SOX10* (as suggested by the human DepMap data) can still
lead to a small number of tumors, albeit with seemingly different biological behaviors. In
the human samples we analyzed, loss of *SOX10* leads to acquisition of a *SOX9*[hi] state
instead[23]. Similarly, in our "escaper" fish, we do see an increase in SOX9 expression in
2 out of 3 fish, which would be consistent with the human data. *SOX9* is increasingly
recognized to be a potent mediator of tumor cell invasiveness, so it is possible this
could explain the somewhat more mesenchymal/invasive appearance of these
tumors[64,65]. This phenomenon clearly needs further study in future work, where the
effect could be better quantified in a larger cohort and the relevant mechanism
identified. But were this SOX10 to SOX9 switch hold up, one implication of this finding is
that clinical targeting of transcription factors like *SOX10*[66], long considered an ideal
therapeutic approach, could lead to unexpected outcomes in which tumors might
proliferate less but become more invasive or resistant to therapy. Whether this
phenomenon of *SOX10/SOX9* switching is a general phenomenon, or specific to that
class of TFs, remains to be determined. However, a similar phenomenon has been
observed for *MITF*, another melanocytic TF, suggesting this idea of TF switching needs
to be more thoroughly explored in the future[58,67,68].
492
493 Finally, while our studies clearly focused on the melanocyte lineage, highly similar
494 approaches could be taken for virtually any cellular lineage in which master regulators
495 or markers are known. Given the large number of already identified promoter/enhancer
496 fragments in the zebrafish, this would be an important extension of our system to study
497 gene variants in the context of development and other diseases[69].
498
Our approach has several limitations that may need to be addressed in future studies.
Due to the constitutive nature of the Cas9, we have little control over the timing of its
expression. Because it is knocked into the endogenous *mitfa* locus, the expression of
Cas9 will vary with the endogenous regulation of that gene. This may lead to somewhat
paradoxical and unexpected results. For example, germline deletion of *sox10* results in
the colourless phenotype, which is an animal completely devoid of melanocytes[15]. In
contrast, when we knock out *sox10* specifically in the mitfa+ cells with our *mitfa*[Cas9]
system, we see a much more subtle phenotype, and not all melanocytes are lost. We

speculate this could be due to timing: the germline allele will already have loss of sox10
function very early in the development of the melanocyte lineage (i.e. at the neural crest
stage), whereas our knockin approach will inactivate sox10 much later in development
after melanocyte specification has already occurred. This could reflect different
downstream targets of *sox10* in these two scenarios. In future iterations of this general
approach, swapping the Cas9 for an inducible version would allow for better timed
knockout of genes[70]. Because *mitfa* itself is expressed in melanoblasts, melanophores
and xanthophores during different stages of development, choosing a promoter (e.g.
PMEL) that is more tightly restricted to melanocytes may be advantageous. Another
limitation is that our fish will result in a single functional copy of *mitfa*. While this does
not result in any obvious phenotype (i.e. the melanocytes are normal) and this gene is
not currently known to exhibit haploinsufficiency, it is possible that under certain
circumstances the loss of one copy of *mitfa* could have unintended phenotypes. For
example, in the tumor setting, having only one copy of *mitfa* could theoretically reduce
tumor incidence since it can act as an oncogene in some situations[71]. This possibility of
haploinsufficiency will need to be further explored in the future. Finally, although our
system was relatively efficient, we still did not see 100% knockout. This may reflect that
it is relatively easy to bypass the CRISPR lesion in an exon with an in-frame mutation.
The use of multiple sgRNAs, or sgRNAs that target promoter regions, rather than
exons, could further augment efficiency of knockout. Alternatively, instead of targeting
DNA, we could consider using RNA targeting CRISPR enzymes. Recent work has
shown that RNA degradation induced by Cas13 works in zebrafish and may allow for
more nuanced knockdown rather than knockout of a given candidate gene[72,73]. This
may especially have advantages in the context of melanoma, where genes are
commonly dysregulated rather than completely knocked out. We envision it would be
relatively straightforward to knock Cas13 into the locus of interest using an analogous
approach to the one we have taken here.
534

Acknowledgements:
We thank members of the White lab for valuable discussions and feedback on this
project. We also thank the MSKCC flow cytometry core for assistance with cell sorting,
Integrated Genomics Operation Core, funded by an NCI Cancer Center Support Grant
(P30 CA08748) for sequencing assistance, and Molecular Cytology Core for their help
with imaging. We thank the MSKCC aquatics core facility for their support of this work.
Schematic figures were all made using Biorender.com with a paid Biorender license.
542

Funding:
RMW was funded through the NIH/NCI Cancer Center Support Grant P30 CA008748,
the Melanoma Research Alliance, The Debra and Leon Black Family Foundation, NIH
Research Program Grants R01CA229215 and R01CA238317, NIH Director's New
Innovator Award DP2CA186572, The Pershing Square Sohn Foundation, The Mark
Foundation for Cancer Research, The American Cancer Society, The Alan and Sandra
Gerry Metastasis Research Initiative at the Memorial Sloan Kettering Cancer Center,
The Harry J. Lloyd Foundation, Consano and the Starr Cancer Consortium. This work
was also supported by NIH ORIP R24OD020166 (MM). YM was supported by a Medical
Scientist Training Program grant from the NIH under award number T32GM007739 to

the Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program and Kirschstein-National Research Service Award (NRSA) predoctoral fellowship under award number F30CA265124. MVH was funded by a K99/R00 Pathway to Independence Award from the National Cancer Institute (1K99CA266931). SP was funded by a Molecular Imaging in Cancer Biology Training Grant at Memorial Sloan Kettering Cancer Center from the National Cancer institute (T32CA254875) and the Ruth L. Kirschstein National Research Service Award for Individual Predoctoral Fellows from the National Cancer Institute (F31CA271518).

Author Contributions:
                Conceptualization: SP, RMW
                Investigation: SP, YM, MVH, JBS, NMC, ZM, JX
                Visualization: SP, MVH, JBS
                Supervision: TL, MM, RMW
                Writing—original draft: SP, RMW
                Writing—review & editing: SP, MVH, NMC, RMW

Declaration of Interests:
MM has competing interests with Recombinetics Inc., LifEngine Technologies, and LifEngine Animal Health Laboratories, Inc. RMW is a paid consultant to N-of-One Therapeutics, a subsidiary of Qiagen and is on the scientific advisory board of Consano but receives no income for this. RMW receives royalty payments for the use of the *casper* zebrafish line from Carolina Biologicals. SP, YM, MVH, JBS, NMC, ZM, JX, and TL declare no competing interests.

Methods

Materials Availability: All data are available in the main text or the supplementary materials, or upon request. Plasmids, fish lines and other materials generated from this study are available upon request.

Zebrafish husbandry and ethics statement
All zebrafish experiments adhered to institutional animal protocols and were conducted in compliance with approved procedures. Fish stocks were maintained at a temperature of 28.5 °C, under 14:10 light:dark cycles, with pH set at 7.4, and salinity-controlled conditions. The zebrafish were fed a standard diet comprising brine shrimp followed by Zeigler pellets. Approval for the animal protocols outlined in this manuscript was obtained from the Memorial Sloan Kettering Cancer Center (MSKCC) Institutional Animal Care and Use Committee (IACUC), under protocol number 12-05-008. Anesthesia was conducted using Tricaine (4 g/l, Syndel, Ferndale, WA, USA) from a stock of 4 g/l and diluted to 0.16 mg/ml. Adult zebrafish of both sexes were equally employed in all experiments. Embryos, collected through natural mating, were incubated in E3 buffer (5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl2, 0.33 mM MgSO4) at 28.5 °C. The wild-type strain used was Tropical 5D Zebrafish (T5D)[27].

GeneWeld plasmid construction

A combination of restriction enzyme digestion and HiFi cloning was used to construct the GeneWeld pPRISM-nCas9n, γcry1:BFP vector for targeted integration to isolate an nls-Cas9-nls knockin. A 765 bp fragment containing the Porcine teschovirus-1 polyprotein 2A peptide sequence was amplified from the pPRISM-2A-Cre, γcry1:BFP vector (Addgene #117789) with primers Ori-F and 2A-R[24,25]. The nCas9n cDNA was amplified from the expression vector pT3TS-nCas9n (Addgene #46757) with primers Cas9-F and Cas9-R[74]. A KpnI/SpeI fragment containing the pPRISM-2A-Cre, gcry1:BFP vector backbone was assembled with the 2A and nCas9n PCR amplicons using the NEBuilder HiFi DNA Assembly Cloning Kit (NEB # E5520S) following the manufacturer's instructions.

Homology arms were designed as previously described[24,25]. Briefly, 48bp homology arms complementary to the *mitfa* target site were designed using GTagHD (http://www.genesculpt.org/gtaghd/) for the pPRISM GeneWeld plasmid series. The two pairs of complementary oligos can be found in Table S1. 5' and 3' arms were cloned sequentially into the pPRISM-nCas9n, γcry1:BFP vector. The 3' homology arm oligos were annealed and cloned into the vector using the BspQI restriction site as previously described. Due to the multiple BfuAI sites within Cas9, instead of using BfuAI restriction cloning, we alternatively used in-fusion cloning (Takara Bio) to insert a gBlock (IDT) containing the 5' homology arm into the pPRISM-nCas9n, gcry1:BFP vector. Primers and gblock sequence used for cloning are listed in Table S1. The 5' and 3' homology arms were sequenced with primers 5'homologycheckF and R3'_pgtag_seq, respectively.

Injection of GeneWeld reagents

Injections of GeneWeld reagents were performed on one-cell stage embryos. We used the Alt-R CRISPR-Cas9 system (IDT) for the initial knock-in of nCas9n, γcry1:BFP to the *mitfa* locus. In brief, 100 µM tracrRNA (IDT 1075928) was mixed at a 1:1 ratio with either *mitfa* crRNA or universal crRNA then incubated at 95 °C for 5 min. Recombinant Cas9 (IDT 1081059) was incubated with both sgRNAs for 10 min at 37 °C to form the RNP complex. The GeneWeld pPRISM-5'mitfaHom-nCas9n, gcry1:BFP-3'mitfaHom plasmid was then added to the injection mix along with phenol red (Sigma-Aldrich P0290). The final concentrations injected into the embryos were: 75 pg/nl Cas9, 12.5 pg/nl *mitfa* sgRNA, 12.5 pg/nl Universal sgRNA, and 5 pg/nl pPRISM plasmid.

Genotyping

Tail clips were taken from 2 month postfertilization (mpf) zebrafish and DNA was extracted using the DNeasy Blood and tissue kit (Qiagen). DNA was PCR amplified with Q5 high fidelity DNA polymerase (NEB). The forward primer MitfaPCR-F binds to the *mitfa* genomic region upstream of the insertion site and the reverse primer 5'homology_amplify-R binds the p2A region of the insertion cassette. Expected PCR amplicon size is 269bp. Amplicons were sequenced using sanger sequencing (Azenta Life Sciences).

RNA-FISH HCR

We adapted the HCR protocol from Ibarra-Garcia-Padilla et al.[75]. Briefly, embryos were
collected and treated with 0.0045% 1-phenyl 2-thiourea (PTU) (Sigma-Aldrich) after 24
hpf to block pigmentation. Embryos were harvested at 72 hpf and fixed in 4% PFA
(Santa Cruz) for 24 hrs at 4 °C. Embryos were then washed with PBS and
dehydrated/permeabilized with a series of MeOH (Millipore) washes. Embryos were
rehydrated with a series of MeOH/PBST washes, permeabilized with acetone (Fisher
Scientific) and incubated with Proteinase K (Millipore) for 30 min. This was followed by
further fixation with 4% PFA. Embryos were pre-hybridized in probe hybridization buffer
(PHB) before incubating them with HCR probes in PHB at concentration of 20nM. Probe
sequences for *mitfa* and Cas9 can be found in Table S1. Embryos were washed with
probe wash buffer before being pre-amplified with probe amplification buffer (PAB).
Following pre-amplification, amplifier hairpins were thawed, snap-cooled, and mixed
with PAB to a concentration of 36 nM. Embryos were incubated in the hairpin solution
followed by washes with 5×SSCT and 1× PBST. Embryos were stained with 2 µg/mL
DAPI in PBST and mounted on glass slides. Imaging was performed using a Zeiss
LSM880 confocal microscope. Melanocytes were assessed for co-expression of *mitfa*
and Cas9 within a 212-micron long section of the embryo, located directly caudal to the
yolk and typically encompassing 3–5 distinct midline melanocytes per embryo. Overlap
in expression was assessed using Fiji.
664
Guide RNA plasmids
The following plasmids were constructed using the Gateway Tol2kit:
zU6A:gRNA-NT;zU6B:gRNA-*ptena*;zU6C:gRNA-*tp53*/394,
zU6A:gRNA-NT;zU6B:gRNA-*ptenb*;zU6C:gRNA-*tp53*/394,
zU6A:gRNA-*sox10*;zU6B:gRNA-*ptena*;zU6C:gRNA-*tp53*/394,
zU6A:gRNA-*sox10*;zU6B:gRNA-*ptenb*;zU6C:gRNA-*tp53*/394.
671
In-fusion cloning (Takara Bio) was used to insert *mitfa:*GFP or *mitfa:*TdTomato into a
zU6:gRNA/394 plasmid previously developed in the lab (See Table S1 for primers
used). gRNAs were cloned into zU6:gRNA;*mitfa:*GFP/394 or
zU6:gRNA;*mitfa:*TdTomato/394 using BsmBI sites. Other plasmids used in this study
that were previously developed in the lab include zU6:gRNA-*ptena*/394, zU6:gRNA-
ptenb/394, zU6:gRNA-*tp53*/394, *mitfa*-BRAF[V600E];cmlc2:eGFP. All guide RNAs used in
this study besides *tuba1a/c* have previously been validated in zebrafish[24,32,56,76].
679
Validation of *tuba1a/c* sgRNAs
The Alt-R CRISPR-Cas9 system (IDT) was used to achieve non-cell type specific
knockout of *tuba1a/c.* Guide RNAs were designed using ChopChop[77]. In brief, 100 µM
tracrRNA (IDT 1075928) was mixed at a 1:1 ratio with crRNA then incubated at 95 °C
for 5 min. Recombinant Cas9 (IDT 1081059) was incubated with the sgRNA for 10 min
at 37 °C to form the RNP complex. Phenol red was added prior to injection into 1 cell-
stage wild-type T5D embryos. To validate on-target cutting, 4 embryos were pooled and
DNA was extracted using Quick Extract buffer (Fisher Scientific NC0302740). DNA was
PCR amplified with Q5 high fidelity DNA polymerase (NEB) and forward and reverse
primers (Table S1). Exosap IT for PCR Product Clean Up was added to PCR product

prior to sanger sequencing. Forward and reverse primers used for PCR amplification
were also used for sequencing (Table S1).
692
Transgenic lines
To generate MG-U6:sgRNA stable lines, one-cell-stage embryos from crossing WT and
mitfa:Cas9 fish were collected and injected with the 30 pg indicated plasmid and 20 pg
tol2 mRNA. Fish with GFP+ melanocytes and BFP+ eyes were selected and outcrossed
with WT fish to produce the F1 generation. F1 fish were again sorted and outcrossed to
generate F2 generation zebrafish. All fish were imaged on a Zeiss AxioZoom V16 with
Zen 2.1 software.
700
Flow cytometry of melanocytes from adult zebrafish
Zebrafish were euthanized using ice-cold water and dissected using a clean scalpel and
forceps. The epidermal and dermal layers of the skin as well as the fins were separated
from the rest of the tissues and diced into 1-3mm pieces. Cells were dissociated with
Liberase TL (Millipore Sigma 05401020001) and filtered for single cell suspensions as
previously described[32]. Samples were then FACS sorted (BD FACSAria) for GFP+ and
GFP- cells. WT fish were used as a GFP negative control. Genomic DNA was isolated
from sorted cells using the DNeasy Blood and Tissue Kit (Qiagen).
709
710   **Tumor Dissection**
711   Fish with tumors were euthanized with ice-cold water and immediately dissected with a
712   scalpel to isolate tumor tissue. Genomic DNA was purified from the tumor samples
713   using the Dneasy Blood and tissue kit (Qiagen).
714
CRISPR sequencing
Primer pairs were designed to produce amplicons 200 to 280 bp in length with the
mutation site within 100 bp from the beginning or end of amplicon (see Table S1 for
primer sequences). Genomic DNA isolated from methods detailed above was PCR
amplified with Q5 high fidelity DNA polymerase (NEB) and run on an agarose gel to
visualize PCR products. The samples were then purified using the NucleoSpin Gel and
PCR cleanup kit (Takara Bio). Deep sequencing was conducted using the CRISPR-seq
platform. Sequencing data was analyzed with CRISPResso2[78]. and indel charts were
generated with CRISPRVariants R package[79].
724
Drug treatments
Adult zebrafish were treated for 24 hr with 750nM neocuproine (Sigma-Aldrich)
dissolved in fish water with a final concentration of 0.0075% DMSO as previously
described[39]. Cells from the middle melanocyte stripe were counted within a rectangular
region of 4mm x 1.5mm. Epinephrine hydrochloride (Sigma-Aldrich) was used at 1
mg/ml in fish water. Fish were treated for 10 minutes prior to imaging.
731
Zebrafish tumor-free survival curves
To generate a transgenic melanoma model, one-cell-stage embryos from WT X
mitfa:Cas9 fish crosses were injected with *mitfa*-BRAF^{V600E};cmlc2:eGFP, 20 pg Tol2
mRNA, and the indicated gRNA plasmids for each condition. The total amount of

plasmid per embryo did not exceed 30 pg. Embryos were screened for GFP positive
hearts and sorted for BFP+ or BFP- eyes. Fish were screened for tumors using a
microscope every 2 weeks starting at 4 weeks post fertilization. Tumors were only
called if an elevated lesion was observed. Graphpad prism 9 was used to generate and
analyze Kaplan-Meier survival curves. Statistical differences were determined using log-
rank Mantel–Cox test. All screening and imaging was conducted on a Zeiss AxioZoom
V16 using Zen 2.1 software.
743
Histology
Zebrafish were sacrificed using ice-cold water and placed in 4% paraformaldehyde
(Santa Cruz 30525-89-4) in PBS for 72 h at 4°C on a rocker. Fish were then transferred
to 70% EtOH for 24 h at 4°C on a rocker. Fish were sent to Histowiz, Inc (Brooklyn, NY,
USA), where they were paraffin embedded, sectioned, stained and imaged. All the
stainings were performed at Histowiz, Inc Brooklyn, using the Leica Bond RX automated
stainer (Leica Microsystems) using a Standard Operating Procedure and fully
automated workflow. Samples were processed, embedded in paraffin, and sectioned at
4μm. The slides were dewaxed using xylene and alcohol based dewaxing solutions.
Epitope retrieval was performed by heat-induced epitope retrieval (HIER) of the
formalin-fixed, paraffin-embedded tissue using Citrate based pH 6 solution (Leica
Microsystems, AR9961) for 20 mins at 95 C. The tissues were first incubated with
peroxide block buffer (Leica Microsystems), followed by incubation with the following
antibodies for 30 min: BRAFV600E antibody (ab228461, 1:100), phospho-AKT antibody
(CST4060, 1:50), SOX9 antibody (AB_185230, 1:1000), and SOX10 antibody
(GTX128374, 1:500). This was followed by incubation with DAB secondary reagents:
polymer, DAB refine and hematoxylin (Bond Polymer Refine Detection Kit, Leica
Microsystems) according to the manufacturer's protocol. The slides were dried,
coverslipped (TissueTek-Prisma Coverslipper) and visualized using a Leica Aperio AT2
slide scanner (Leica Microsystems) at 40X.
764
765
Quantification of Sox10 intensity
FFPE slides were deparaffinized by consecutive incubations in xylene and 100-50%
ethanol. For antigen retrieval, slides were placed in 10 mM sodium citrate pH 6.2 in a
pressure cooker and heated to 95°C for 20 minutes. Slides were then cooled to room
temperature and blocked in 5% donkey serum, 1% BSA, and 0.4% Triton-X100 in PBS
for 1 hour at room temperature. The primary antibody targeting Sox10 (GeneTex,
GTX128374) was added to the blocking buffer at 1:200 concentration and incubated at
4°C overnight. Slides were then washed in PBS before incubation with the secondary
antibody (donkey anti-rabbit Alexa Fluor Plus 488, Thermo Fisher Scientific #AC32790;
1:250) and Hoescht (Thermo Fisher Scientific, #62249; 1:1000) in blocking buffer for 2
hours at room temperature. Slides were mounted in Vectashield Plus (Vector Labs, H-
1900). Slides were imaged on a LSM 880 confocal microscope using a 40X oil
immersion objective. 3 images were captured per sample. Quantification of Sox10
intensity per cell was performed using CellProfiler[80] and MATLAB r2023b (Mathworks).
Cells were segmented in CellProfiler to obtain a nuclear mask from the Hoescht signal
and using this mask the mean intensity per nucleus was quantified.

782
Re-analysis of human melanoma scRNA-seq data
Human melanoma scRNA-seq data from Wouters et al. was accessed from GEO
(GSE134432)[23]. All analysis was done in R (version 4.3.1) using the Seurat package
(version 5.0.2)[81]. Counts matrices for each sample and cell line were imported into R
using the function read.table and converted into Seurat objects with the function
CreateSeuratObject before merging into one combined Seurat object using the function
merge. Counts were normalized using the Seurat function NormalizeData with default
parameters. UMAP was calculated using the Seurat function RunUMAP with 10
dimensions. SOX10 expression was plotted using the functions FeaturePlot and
VlnPlot.

Statistical analysis
Statistical significance was analyzed using t test, Log-rank (Mantel-Cox) test, Wilcoxon
Rank Sum Test or as indicated in the figure legends. GraphPad Prism 9 software was
used for data processing and statistical analysis. *P < 0.05; **P < 0.005; ***P < 0.001;
****P < 0.0001. Data are presented as means ± SD unless otherwise indicated.

References

1   Mayer, C. S. & Huser, V. Learning important common data elements from shared
     study data: The All of Us program analysis. *PLoS One* **18**, e0283601 (2023).
     https://doi.org/10.1371/journal.pone.0283601
2   Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes
     of a wide range of complex diseases of middle and old age. *PLoS Med* **12**,
     e1001779 (2015). https://doi.org/10.1371/journal.pmed.1001779
3   Tsherniak, A. *et al.* Defining a Cancer Dependency Map. *Cell* **170**, 564-576 e516
     (2017). https://doi.org/10.1016/j.cell.2017.06.010
4   Amsterdam, A. *et al.* Identification of 315 genes essential for early zebrafish
     development. *Proc Natl Acad Sci U S A* **101**, 12792-12797 (2004).
     https://doi.org/10.1073/pnas.0403929101
5   Yin, L. *et al.* Multiplex Conditional Mutagenesis Using Transgenic Expression of
     Cas9 and sgRNAs. *Genetics* **200**, 431-441 (2015).
     https://doi.org/10.1534/genetics.115.176917
6   Ablain, J., Durand, E. M., Yang, S., Zhou, Y. & Zon, L. I. A CRISPR/Cas9 vector
     system for tissue-specific gene disruption in zebrafish. *Dev Cell* **32**, 756-764
     (2015). https://doi.org/10.1016/j.devcel.2015.01.032
7   Zhou, W. *et al.* Neutrophil-specific knockout demonstrates a role for mitochondria
     in regulating neutrophil motility in zebrafish. *Dis Model Mech* **11** (2018).
     https://doi.org/10.1242/dmm.033027
8   Ni, T. T. *et al.* Conditional control of gene function by an invertible gene trap in
     zebrafish. *Proc Natl Acad Sci U S A* **109**, 15389-15394 (2012).
     https://doi.org/10.1073/pnas.1206131109
9   Grajevskaja, V., Camerota, D., Bellipanni, G., Balciuniene, J. & Balciunas, D.
     Analysis of a conditional gene trap reveals that tbx5a is required for heart

regeneration in zebrafish. *PLoS One* **13**, e0197293 (2018).
https://doi.org/10.1371/journal.pone.0197293
10 Burg, L. *et al.* Conditional mutagenesis by oligonucleotide-mediated integration of
loxP sites in zebrafish. *PLoS Genet* **14**, e1007754 (2018).
https://doi.org/10.1371/journal.pgen.1007754
11 Kalvaityte, M. & Balciunas, D. Conditional mutagenesis strategies in zebrafish.
Trends Genet **38**, 856-868 (2022). https://doi.org/10.1016/j.tig.2022.04.007
12 Singh Angom, R., Wang, Y., Wang, E., Dutta, S. K. & Mukhopadhyay, D.
Conditional, Tissue-Specific CRISPR/Cas9 Vector System in Zebrafish Reveals
the Role of Nrp1b in Heart Regeneration. *Arterioscler Thromb Vasc Biol* **43**,
1921-1934 (2023). https://doi.org/10.1161/ATVBAHA.123.319189
13 Shin, M. *et al.* Generation and application of endogenously floxed alleles for cell-
specific knockout in zebrafish. *Dev Cell* **58**, 2614-2626 e2617 (2023).
https://doi.org/10.1016/j.devcel.2023.07.022
14 Ablain, J., Liu, S., Moriceau, G., Lo, R. S. & Zon, L. I. SPRED1 deletion confers
resistance to MAPK inhibition in melanoma. *J Exp Med* **218** (2021).
https://doi.org/10.1084/jem.20201097
15 Dutton, K. A. *et al.* Zebrafish colourless encodes sox10 and specifies non-
ectomesenchymal neural crest fates. *Development* **128**, 4113-4125 (2001).
https://doi.org/10.1242/dev.128.21.4113
16 Parichy, D. M. *et al.* Mutational analysis of endothelin receptor b1 (rose) during
neural crest and pigment pattern development in the zebrafish Danio rerio. *Dev
Biol* **227**, 294-306 (2000). https://doi.org/10.1006/dbio.2000.9899
17 Eom, D. S., Patterson, L. B., Bostic, R. R. & Parichy, D. M. Immunoglobulin
superfamily receptor Junctional adhesion molecule 3 (Jam3) requirement for
melanophore survival and patterning during formation of zebrafish stripes. *Dev
Biol* **476**, 314-327 (2021). https://doi.org/10.1016/j.ydbio.2021.04.007
18 Parichy, D. M. & Turner, J. M. Zebrafish puma mutant decouples pigment pattern
and somatic metamorphosis. *Dev Biol* **256**, 242-257 (2003).
https://doi.org/10.1016/s0012-1606(03)00015-0
19 Nguyen, P. D. *et al.* Haematopoietic stem cell induction by somite-derived
endothelial cells controlled by meox1. *Nature* **512**, 314-318 (2014).
https://doi.org/10.1038/nature13678
20 Irion, U., Singh, A. P. & Nusslein-Volhard, C. The Developmental Genetics of
Vertebrate Color Pattern Formation: Lessons from Zebrafish. *Curr Top Dev Biol*
117, 141-169 (2016). https://doi.org/10.1016/bs.ctdb.2015.12.012
21 Parichy, D. M., Rawls, J. F., Pratt, S. J., Whitfield, T. T. & Johnson, S. L.
Zebrafish sparse corresponds to an orthologue of c-kit and is required for the
morphogenesis of a subpopulation of melanocytes, but is not essential for
hematopoiesis or primordial germ cell development. *Development* **126**, 3425-
3436 (1999). https://doi.org/10.1242/dev.126.15.3425
22 Capparelli, C. *et al.* Targeting SOX10-deficient cells to reduce the dormant-
invasive phenotype state in melanoma. *Nat Commun* **13**, 1381 (2022).
https://doi.org/10.1038/s41467-022-28801-y

23   Wouters, J. *et al.* Robust gene expression programs underlie recurrent cell states
and phenotype switching in melanoma. *Nat Cell Biol* **22**, 986-998 (2020).
https://doi.org/10.1038/s41556-020-0547-3

24   Welker, J. M. *et al.* GeneWeld: Efficient Targeted Integration Directed by Short
Homology in Zebrafish. *Bio Protoc* **11**, e4100 (2021).
https://doi.org/10.21769/BioProtoc.4100

25   Wierson, W. A. *et al.* Efficient targeted integration directed by short homology in
zebrafish and mammalian cells. *Elife* **9** (2020).
https://doi.org/10.7554/eLife.53968

26   White, R. M. *et al.* Transparent adult zebrafish as a tool for in vivo transplantation
analysis. *Cell Stem Cell* **2**, 183-189 (2008).
https://doi.org/10.1016/j.stem.2007.11.002

27   Balik-Meisner, M., Truong, L., Scholl, E. H., Tanguay, R. L. & Reif, D. M.
Population genetic diversity in zebrafish lines. *Mamm Genome* **29**, 90-100
(2018). https://doi.org/10.1007/s00335-018-9735-x

28   Lister, J. A., Robertson, C. P., Lepage, T., Johnson, S. L. & Raible, D. W. nacre
encodes a zebrafish microphthalmia-related protein that regulates neural-crest-
derived pigment cell fate. *Development* **126**, 3757-3767 (1999).
https://doi.org/10.1242/dev.126.17.3757

29   Fernandez, L. P. *et al.* SLC45A2: a novel malignant melanoma-associated gene.
Hum Mutat **29**, 1161-1167 (2008). https://doi.org/10.1002/humu.20804

30   Branicki, W., Brudnik, U., Draus-Barini, J., Kupiec, T. & Wojas-Pelc, A.
Association of the SLC45A2 gene with physiological human hair colour variation.
J Hum Genet **53**, 966-971 (2008). https://doi.org/10.1007/s10038-008-0338-3

31   Streisinger, G., Singer, F., Walker, C., Knauber, D. & Dower, N. Segregation
analyses and gene-centromere distances in zebrafish. *Genetics* **112**, 311-319
(1986). https://doi.org/10.1093/genetics/112.2.311

32   Weiss, J. M. *et al.* Anatomic position determines oncogenic specificity in
melanoma. *Nature* **604**, 354-361 (2022). https://doi.org/10.1038/s41586-022-
04584-6

33   Hou, L., Arnheiter, H. & Pavan, W. J. Interspecies difference in the regulation of
melanocyte development by SOX10 and MITF. *Proc Natl Acad Sci U S A* **103**,
9081-9085 (2006). https://doi.org/10.1073/pnas.0603114103

34   Kelsh, R. N. *et al.* Zebrafish pigmentation mutations and the processes of neural
crest development. *Development* **123**, 369-389 (1996).
https://doi.org/10.1242/dev.123.1.369

35   Carney, T. J. *et al.* A direct role for Sox10 in specification of neural crest-derived
sensory neurons. *Development* **133**, 4619-4630 (2006).
https://doi.org/10.1242/dev.02668

36   Cunningham, R. L. *et al.* Functional in vivo characterization of sox10 enhancers
in neural crest and melanoma development. *Commun Biol* **4**, 695 (2021).
https://doi.org/10.1038/s42003-021-02211-0

37   Owen, J. P., Kelsh, R. N. & Yates, C. A. A quantitative modelling approach to
zebrafish pigment pattern formation. *Elife* **9** (2020).
https://doi.org/10.7554/eLife.52998

38 Nagao, Y. *et al.* Distinct interactions of Sox5 and Sox10 in fate specification of
pigment cells in medaka and zebrafish. *PLoS Genet* **14**, e1007260 (2018).
https://doi.org/10.1371/journal.pgen.1007260
39 O'Reilly-Pol, T. & Johnson, S. L. Neocuproine ablates melanocytes in adult
zebrafish. *Zebrafish* **5**, 257-264 (2008). https://doi.org/10.1089/zeb.2008.0540
40 Cushion, T. D., Leca, I. & Keays, D. A. MAPping tubulin mutations. *Front Cell*
Dev Biol **11**, 1136699 (2023). https://doi.org/10.3389/fcell.2023.1136699
41 Rogers, S. L., Tint, I. S., Fanapour, P. C. & Gelfand, V. I. Regulated bidirectional
motility of melanophore pigment granules along microtubules in vitro. *Proc Natl*
Acad Sci U S A **94**, 3720-3725 (1997). https://doi.org/10.1073/pnas.94.8.3720
42 Larson, T. A., Gordon, T. N., Lau, H. E. & Parichy, D. M. Defective adult
oligodendrocyte and Schwann cell development, pigment pattern, and
craniofacial morphology in puma mutant zebrafish having an alpha tubulin
mutation. *Dev Biol* **346**, 296-309 (2010).
https://doi.org/10.1016/j.ydbio.2010.07.035
43 Farnsworth, D. R., Saunders, L. M. & Miller, A. C. A single-cell transcriptome
atlas for zebrafish development. *Dev Biol* **459**, 100-108 (2020).
https://doi.org/10.1016/j.ydbio.2019.11.008
44 Neuhauss, S. C. *et al.* Genetic disorders of vision revealed by a behavioral
screen of 400 essential loci in zebrafish. *J Neurosci* **19**, 8603-8615 (1999).
https://doi.org/10.1523/JNEUROSCI.19-19-08603.1999
45 Myers, K. A., Bello-Espinosa, L. E., Kherani, A., Wei, X. C. & Innes, A. M.
TUBA1A Mutation Associated With Eye Abnormalities in Addition to Brain
Malformation. *Pediatr Neurol* **53**, 442-444 (2015).
https://doi.org/10.1016/j.pediatrneurol.2015.07.004
46 Veldman, M. B., Bemben, M. A. & Goldman, D. Tuba1a gene expression is
regulated by KLF6/7 and is necessary for CNS development and regeneration in
zebrafish. *Mol Cell Neurosci* **43**, 370-383 (2010).
https://doi.org/10.1016/j.mcn.2010.01.004
47 Roh, M. R. *et al.* Promoter Methylation of PTEN Is a Significant Prognostic Factor
in Melanoma Survival. *J Invest Dermatol* **136**, 1002-1011 (2016).
https://doi.org/10.1016/j.jid.2016.01.024
48 Palmieri, G. *et al.* Multiple Molecular Pathways in Melanomagenesis:
Characterization of Therapeutic Targets. *Front Oncol* **5**, 183 (2015).
https://doi.org/10.3389/fonc.2015.00183
49 Frantz, W. T. & Ceol, C. J. From Tank to Treatment: Modeling Melanoma in
Zebrafish. *Cells* **9** (2020). https://doi.org/10.3390/cells9051289
50 Ignatius, M. S. *et al.* tp53 deficiency causes a wide tumor spectrum and
increases embryonal rhabdomyosarcoma metastasis in zebrafish. *Elife* **7** (2018).
https://doi.org/10.7554/eLife.37202
51 Patton, E. E. *et al.* BRAF mutations are sufficient to promote nevi formation and
cooperate with p53 in the genesis of melanoma. *Curr Biol* **15**, 249-254 (2005).
https://doi.org/10.1016/j.cub.2005.01.031
52 He, S. *et al.* Synergistic melanoma cell death mediated by inhibition of both
MCL1 and BCL2 in high-risk tumors driven by NF1/PTEN loss. *Oncogene* **40**,
5718-5729 (2021). https://doi.org/10.1038/s41388-021-01926-y

Montal, E., Suresh, S., Ma, Y., Tagore, M. M. & White, R. M. Cancer Modeling by Transgene Electroporation in Adult Zebrafish (TEAZ). *Methods Mol Biol* **2707**, 83-97 (2024). https://doi.org/10.1007/978-1-0716-3401-1_5

Croushore, J. A. *et al.* Ptena and ptenb genes play distinct roles in zebrafish embryogenesis. *Dev Dyn* **234**, 911-921 (2005). https://doi.org/10.1002/dvdy.20576

Faucherre, A., Taylor, G. S., Overvoorde, J., Dixon, J. E. & Hertog, J. Zebrafish pten genes have overlapping and non-redundant functions in tumorigenesis and embryonic development. *Oncogene* **27**, 1079-1086 (2008). https://doi.org/10.1038/sj.onc.1210730

Kaufman, C. K. *et al.* A zebrafish melanoma model reveals emergence of neural crest identity during melanoma initiation. *Science* **351**, aad2197 (2016). https://doi.org/10.1126/science.aad2197

Shakhova, O. *et al.* Antagonistic cross-regulation between Sox9 and Sox10 controls an anti-tumorigenic program in melanoma. *PLoS Genet* **11**, e1004877 (2015). https://doi.org/10.1371/journal.pgen.1004877

Hoek, K. S. *et al.* In vivo switching of human melanoma cells between proliferative and invasive states. *Cancer Res* **68**, 650-656 (2008). https://doi.org/10.1158/0008-5472.CAN-07-2491

Solnica-Krezel, L., Schier, A. F. & Driever, W. Efficient recovery of ENU-induced mutations from the zebrafish germline. *Genetics* **136**, 1401-1420 (1994). https://doi.org/10.1093/genetics/136.4.1401

Russell, W. L. *et al.* Specific-locus test shows ethylnitrosourea to be the most potent mutagen in the mouse. *Proc Natl Acad Sci U S A* **76**, 5818-5819 (1979). https://doi.org/10.1073/pnas.76.11.5818

Cerami, E. *et al.* The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* **2**, 401-404 (2012). https://doi.org/10.1158/2159-8290.CD-12-0095

Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415-421 (2013). https://doi.org/10.1038/nature12477

Hoek, K. S. & Goding, C. R. Cancer stem cells versus phenotype-switching in melanoma. *Pigment Cell Melanoma Res* **23**, 746-759 (2010). https://doi.org/10.1111/j.1755-148X.2010.00757.x

Cheng, P. F. *et al.* Methylation-dependent SOX9 expression mediates invasion in human melanoma cells and is a negative prognostic factor in advanced melanoma. *Genome Biol* **16**, 42 (2015). https://doi.org/10.1186/s13059-015-0594-4

Yang, X. *et al.* SOX9 is a dose-dependent metastatic fate determinant in melanoma. *J Exp Clin Cancer Res* **38**, 17 (2019). https://doi.org/10.1186/s13046-018-0998-6

Takahashi, M. *et al.* DrugMap: A quantitative pan-cancer analysis of cysteine ligandability. *Cell* **187**, 2536-2556 e2530 (2024). https://doi.org/10.1016/j.cell.2024.03.027

Hossain, S. M. & Eccles, M. R. Phenotype Switching and the Melanoma Microenvironment; Impact on Immunotherapy and Drug Resistance. *Int J Mol Sci* **24** (2023). https://doi.org/10.3390/ijms24021601

68 Travnickova, J. *et al.* Zebrafish MITF-Low Melanoma Subtype Models Reveal
Transcriptional Subclusters and MITF-Independent Residual Disease. *Cancer*
Res **79**, 5769-5784 (2019). https://doi.org/10.1158/0008-5472.CAN-19-0037
69 Weiss, J. M. *et al.* Shifting the focus of zebrafish toward a model of the tumor
microenvironment. *Elife* **11** (2022). https://doi.org/10.7554/eLife.69703
70 Sun, N. *et al.* Development of drug-inducible CRISPR-Cas9 systems for large-
scale functional screening. *BMC Genomics* **20**, 225 (2019).
https://doi.org/10.1186/s12864-019-5601-9
71 Garraway, L. A. *et al.* Integrative genomic analyses identify MITF as a lineage
survival oncogene amplified in malignant melanoma. *Nature* **436**, 117-122
(2005). https://doi.org/10.1038/nature03664
72 Huang, Y. *et al.* CRISPR-dCas13-tracing reveals transcriptional memory and
limited mRNA export in developing zebrafish embryos. *Genome Biol* **24**, 15
(2023). https://doi.org/10.1186/s13059-023-02848-6
73 Kushawah, G. *et al.* CRISPR-Cas13d Induces Efficient mRNA Knockdown in
Animal Embryos. *Dev Cell* **54**, 805-817 e807 (2020).
https://doi.org/10.1016/j.devcel.2020.07.013
74 Jao, L. E., Wente, S. R. & Chen, W. Efficient multiplex biallelic zebrafish genome
editing using a CRISPR nuclease system. *Proc Natl Acad Sci U S A* **110**, 13904-
13909 (2013). https://doi.org/10.1073/pnas.1308335110
75 Ibarra-Garcia-Padilla, R., Howard, A. G. A. t., Singleton, E. W. & Uribe, R. A. A
protocol for whole-mount immuno-coupled hybridization chain reaction
(WICHCR) in zebrafish embryos and larvae. *STAR Protoc* **2**, 100709 (2021).
https://doi.org/10.1016/j.xpro.2021.100709
76 Suresh, S. *et al.* Identifying the Transcriptional Drivers of Metastasis Embedded
within Localized Melanoma. *Cancer Discov* **13**, 194-215 (2023).
https://doi.org/10.1158/2159-8290.CD-22-0427
77 Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond
genome editing. *Nucleic Acids Res* **47**, W171-W174 (2019).
https://doi.org/10.1093/nar/gkz365
78 Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing
sequence analysis. *Nat Biotechnol* **37**, 224-226 (2019).
https://doi.org/10.1038/s41587-019-0032-3
79 Lindsay, H. *et al.* CrispRVariants charts the mutation spectrum of genome
engineering experiments. *Nat Biotechnol* **34**, 701-702 (2016).
https://doi.org/10.1038/nbt.3628
80 Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability.
BMC Bioinformatics **22**, 433 (2021). https://doi.org/10.1186/s12859-021-04344-9
81 Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-
cell analysis. *Nat Biotechnol* **42**, 293-304 (2024). https://doi.org/10.1038/s41587-
023-01767-y
1049

1050
Figure 1. Targeted integration of Cas9 to the *mitfa* locus with GeneWeld knockin system.
(A) Schematic depicting integration site and GeneWeld knock-in cassette. 5' and 3' homology
arms are designed to target exon 2 of the zebrafish *mitfa* gene. The knock-in cassette includes
p2A followed by Cas9 and BFP driven by the eye-specific promoter gamma-crystallin. **(B)**
Pipeline to produce F1 mitfa^{Cas9} zebrafish. The GeneWeld knock-in vector and gRNAs targeting
the *mitfa* genomic insertion site or specific sites on the knock-in vector (UgRNA) are co-injected
into one-cell stage wild-type zebrafish embryos. Embryos are screened for BFP+ eyes marking
mosaic integration. These mosaic founder fish are then raised to adulthood and crossed with
wild-type fish. The resulting embryos are screened for BFP+ eyes and sequenced to confirm
precise integration. **(C)** Fluorescent in situ hybridization chain reaction (FISH) on 3dpf wild-type
and *mitfa*^{Cas9} embryos treated with 1-phenyl 2-thiourea (PTU). Arrows on the whole embryo
image (WT 3dpf embryo not treated with PTU) indicate the embryonic melanocyte stripe regions
where *mitfa* and Cas9 expression is expected. The presence of *mitfa* and Cas9 RNA was
assessed by confocal microscopy at 40x magnification. Maximum intensity projections are
shown. n=17 WT embryos and n=16 mitfa^{Cas9} embryos were screened. Scale bars, 100µm.

1066
Figure 2. Melanocyte lineage-specific knockout of *albino* using *mitfa*[Cas9] fish.
(A) Pipeline to generate F0 and F1 U6:gRNA; *mitfa*:GFP (MG-gRNA) zebrafish. **(B)** Adult
mitfa[Cas9] MG-NT and MG-*albino* F0 fish. **(C)** Proportion of MG-NT (n=19) and MG-*albino* (n=25)
F0 fish with loss of pigmentation phenotype. **(D)** Adult mitfa[Cas9] MG-NT and MG-*albino* F1 fish.
(E) Pigmented area/ mm^2 calculated for n=5 fish/ genotype within a defined rectangular region of
interest (ROI) encompassing the top and middle melanocyte stripes. Two-sided student's *t*-test
was used to assess statistical significance, **** p < 0.0001; error bars, SD. **(F)** 3dpf mitfa[Cas9]
MG-NT and MG-*albino* F2 fish. Representative embryos are shown for each genotype. **(G)**
Mean grey value of head melanocytes calculated for n=10 embryos/genotype within a defined
hexagonal ROI indicated as a red outline in 2F. Two-sided student's *t*-test was used to assess
statistical significance, **** p < 0.0001. **(H)** Schematic for CRISPR sequencing protocol. **(I)**
CRISPR-seq results are shown for WT and n=3 independent F1 MG-*albino* fish. Only GFP
negative cells were isolated from WT fish. Results are shown as a fraction of sequences with
indels calculated using CRISPRESSO.

Figure 3. Melanocyte lineage-specific knockout of *sox10* using *mitfa*^Cas9 fish.
(A) Adult mitfa^Cas9 MG-NT and MG-*sox10* F0 fish. **(B)** Proportion of MG-NT (n=19) and MG-
sox10 (n=16) F0 fish with disrupted stripes phenotype. **(C)** Adult mitfa^Cas9 MG-NT and MG-*sox10*
F1 fish. **(D)** The average width of melanocyte stripe 1D and xanthophore interstripe X0 was
calculated for each fish by averaging 5 stripe/interstripe width measurements. N=5 fish per
genotype. Two-sided student's *t*-test was used to assess statistical significance, *** p < 0.001;

error bars, SD. **(E)** F1 MG-NT (n=5) and MG-*sox10* (n=5) adult fish were treated with epinephrine and melanocytes were counted within a defined rectangular region of interest 3.46mm x 2.54mm encompassing the top and middle melanocyte stripes. Two-sided student's *t*-test was used to assess statistical significance, * $p < 0.05$; error bars, SD. **(F)** Schematic of neocuproine (neo) experimental setup. Adult mitfa^{Cas9} MG-NT (n= 5) and MG-*sox10* (n=5) F1 fish were treated with neocuproine for 24 hours to ablate melanocytes then imaged at day 7,15, and 70 to measure regeneration of melanocytes compared to day 0. **(G)** Quantification of melanocyte regeneration. Fish were treated with epinephrine prior to imaging to enable counting of melanocytes. Two-sided student's *t*-test was used to assess statistical significance, ** $p < 0.01$; error bars, SD. **(H)** Representative images are shown for MG-NT and MG-*sox10* fish pre- and post-neocuproine treatment.

1107
Figure 4. Non-autonomous function of *tuba1a/tuba1c* on melanocytes.
(A) 4dpf zebrafish embryos injected with either NT or *tuba1a/c* Alt-R CRISPR Cas9 gRNAs. **(B)**
Percentage of NT or *tuba1a/c* Alt-R zebrafish embryos with dispersed melanocyte phenotypes.
N=2 independent experiments. **(C)** Survival percentage is shown for NT and *tuba1a/c* Alt-R
embryos. Embryos were counted at 24hpf and again at 14dpf to determine survival. **(D)** 4dpf
mitfa[Cas9] MG-*tuba1a/c* F2 embryos (BFP+ eyes) compared to sibling controls with no mitfa[Cas9]
(BFP- eyes). Representative embryos are shown for each genotype. **(E)** Pigmented area/ mm^2
calculated for n=19 embryos/genotype from two independent MG-*tuba1a/c* F2 clutches. Two-
sided student's *t*-test was used to assess statistical significance, ns: no significance. **(F)** 4dpf
Alt-R injected zebrafish embryos imaged before and after epinephrine (epi) treatment. **(G)**
Pigmented area/ mm^2 calculated for n=20 embryos/genotype from two independent clutches.
Two-sided student's *t*-test was used to assess statistical significance, **** p < 0.0001.

1120
1121
Figure 5. Generation of a zebrafish melanoma model in *mitfa*^{Cas9} fish.

(A) Adult mitfa^{Cas9} MG-NT; *mitfa*:TdTomato;U6:NT (MTdt-NT) and MG-*ptena*;
mitfa:TdTomato;U6:*ptenb* (MTdT-*ptenb*) F0 fish. **(B)** Proportion of MG-NT; MTdT-NT (n=14) and
MG-*ptena*; MTdT-*ptenb* (n=14) F0 fish with disrupted stripes phenotype. **(C)** Schematic of
zebrafish tumorigenesis assay. Indicated plasmids are injected into one-cell stage embryos from
crosses between mitfa^{Cas9} and WT fish. The mitfa:BRAF^{V600E} plasmid includes cardiac-specific
cmlc2:GFP. Embryos were sorted for GFP+ hearts and BFP+ eyes and fish were screened
every 2 weeks for tumors. **(D)** Tumor free survival curve. N=2 independent experiments. Tumors
were tracked over the course of 30 weeks. Log-rank (Mantel-Cox) test was used to assess
statistical significance, *** p < 0.001; **** p < 0.0001; ns: no significance. **(E)** Histology was
performed on one fish from each indicated injection group. Dotted lines indicate site of
sectioning. Red chromogen was used for all IHC staining. Scale bars, 50µm. **(F)** CRISPR-seq

1134   results are shown for normal skin dissected from WT fish and tumors dissected from injection
1135   conditions 2, 3, and 4. Results are shown as a fraction of sequences with indels calculated
1136   using CRISPRESSO.

1152
Figure 6. Melanoma specific knockout of *sox10* reduces tumor burden and induces
phenotypic switching.
(**A**) Schematic of zebrafish tumorigenesis assay. Indicated plasmids are injected into one-cell
stage embryos from crosses between mitfa[Cas9] and WT fish. Embryos were sorted for GFP+
hearts and BFP+ eyes and fish were screened every 2 weeks for tumors. (**B**) Tumor free
survival curve. N=3 independent experiments. Tumors were tracked over the course of 50
weeks. Log-rank (Mantel-Cox) test was used to assess statistical significance, **** p < 0.0001.
(**C**) Histology is shown for one fish from each injection group. Dotted lines indicate site of
sectioning. Red chromogen was used for all IHC staining. Scale bars, 50µm. (**D**) Color
thresholding was used on IHC images to calculate the percentage of nuclei that stained positive
for *sox9*. NT-1 n=1222 cells, NT-2 n=1620 cells, NT-3 n=2185 cells, sox10-1 n=970 cells,
sox10-2 n=594, sox10-3 n=1046. Cells from n=3 images were analyzed for each tumor. (**E**)
Schematic of scRNA-seq experimental setup from Wouters et al. Patient derived cell lines were
treated with siRNAs targeting SOX10 or NTC and scRNA-seq was conducted at 72hrs. (**F**)
UMAP of scRNA-seq dataset for MM057 cell line from Wouters et al. Cells from siSOX10 and
siNT conditions are labeled. (**G**) Normalized expression of Sox10 per cell in UMAP space. (**H**)
Violin plots of normalized expression of Sox10 per cell. Median is shown as dashed red line.
Wilcoxon Rank Sum Test was used to assess statistical significance, **** p < 0.0001. (**I**)
Normalized expression of Sox9 per cell in UMAP space. (**J**) Violin plots of normalized

expression of Sox9 per cell. Wilcoxon Rank Sum Test was used to assess statistical
significance, **** p < 0.0001.

A

B

C

Figure S1. Validation of Cas9 knock-in.
(A) Integration efficiency of *mitfa*-targeted P2A-Cas9, gamma-cry:BFP vector. n= 4
independent experiments are shown with 50 embryos screened for each experiment.
BFP+ eyes were detected in a total of 37/200 embryos. Error bars, SD. **(B)** Sanger
sequencing results from four F1 fish with BFP+ eyes. Primers were designed with the
forward primer located in the genomic *mitfa* locus and reverse primer within the Cas9
insertion cassette so that amplification can only occur if there is integration in the correct
orientation. Snapgene was used to align sequences **(C)** Images of 3 day old and 3
month old clutchmates from a *mitfa*Cas9 incross. **(D)** Quantification of the percentage of
mitfa positive cells that have Cas9 expression in each embryo. The presence of *mitfa*
and Cas9 RNA was assessed by confocal microscopy from fluorescent in situ
hybridization chain reaction (FISH) on 3dpf wild-type and *mitfa*Cas9 embryos. Out of 28
mitfa+ cells assessed across n=7 embryos, 24 had detectable Cas9 RNA.

A F2 MG-gRNA 1mpf

B F2 MG-gRNA 1mpf

C F2 MG-gRNA 3mpf

D F2 MG-gRNA 3mpf

E

ENSDARG00000002593.8 *(albino)*

1205
Figure S2. *albino* knockout fish.
(A) Quantification of pigmentation for 1mpf mitfa^{Cas9} MG-NT and MG-*albino* F2 fish.
Pigmented area/ mm^2 calculated for n=12 fish/ genotype within a defined rectangular
region of interest (ROI) encompassing the top and middle melanocyte stripes. Two-
sided student's *t*-test was used to assess statistical significance, **** p < 0.0001; error
bars, SD. **(B)** Color and GFP images of 1mpf mitfa^{Cas9} MG-NT and MG-*albino* F2 fish.

(C) Pigmented area/ mm^2 calculated for n=12 fish/ genotype within a defined rectangular
region of interest (ROI) encompassing the top and middle melanocyte stripes. Two-
sided student's *t*-test was used to assess statistical significance, **** $p < 0.0001$; error
bars, SD. **(D)** Color images of 3mpf mitfa^{Cas9} MG-NT and MG-*albino* F2 fish. **(E**) Indel
chart for the *albino* locus produced using CRISPR Variants. **T insertion is observed
across all samples, likely resulting from a PCR or sequencing artifact.
1218
1219
1220

Figure S3

1222
1223
Figure S3. Knockout of *tuba1a*
(A) 6dpf zebrafish embryos injected with either NT or *tuba1a/c* sg1 Alt-R CRISPR Cas9
gRNAs. **(B)** Validation of Alt-R CRISPR Cas9 *tuba1a/c* sg1 targeting with sanger
sequencing. Synthego Ice software estimated 88% indels in the *tuba1a* locus and 97%
indels in the *tuba1c* locus. **(C)** Validation of Alt-R CRISPR Cas9 *tuba1a*-specific sgRNA
targeting with sanger sequencing. Synthego Ice software estimated 97% indels in the
tuba1a locus. **(D)** Validation of Alt-R CRISPR Cas9 *tuba1c*-specific sgRNA targeting
with sanger sequencing. Synthego Ice software estimated 56% indels in the *tuba1c*
locus.**(E)** Validation of Alt-R CRISPR Cas9 *tuba1a/c* sg2 targeting with sanger
sequencing. Synthego Ice software estimated 88% indels in the *tuba1a* locus and 86%
indels in the *tuba1c* locus. **(F)** Percentage of NT or *tuba1a/c* sg2 Alt-R zebrafish
embryos with dispersed melanocyte phenotypes. N=2 independent experiments. **(G)**
Adult mitfa[Cas9] MG-*tuba1a/c* F0 fish. Image is representative of n=7 F0 fish. **(H)** Adult
mitfa[Cas9] MG-*tuba1a/c* F1 fish. Image is representative of n=14 F1 fish. **(I)** CRISPR-seq
results are shown for WT and n=2 F1 MG-*tuba1a/c* fish sorted for GFP+ cells. Results
are shown as a fraction of sequences with indels calculated using CRISPRESSO.
1240
1241

1242
1243 **Figure S4. CRISPR-seq for tumor suppressor genes.**
1244 (**A**) Indel chart for the *ptena* locus produced using CRISPRVariants. (**B**) Indel chart for
1245 the *ptenb* locus produced using CRISPRVariants. (**C**) Indel chart for the *p53* locus
1246 produced using CRISPRVariants.

Figure S5

1247
Figure S5. *In vivo* and *in vitro* targeting of Sox10 leads to upregulation of Sox9.
(**A**) Histology is shown for two additional fish from each injection group. Dotted lines
indicate site of sectioning. Red chromogen was used for all IHC staining. Scale bars,
50μm. (**B**) *Sox10* intensity per nucleus from NT and *sox10* KO tumors. Intensity was
calculated from immunofluorescence staining of *Sox10*. Red lines indicate mean with

95% CI. Each point represents one nuclei. NT-1 n=1334 cells, NT-2 n=1606 cells, NT-3
n=1204 cells, sox10-1 n=869 cells, sox10-2 n=998, sox10-3 n=1046. Cells from n=3
images were analyzed for each tumor. **(C)** UMAP of scRNA-seq dataset for MM074 cell
line from Wouters et al. Cells from siSOX10 and siNT conditions are labeled. **(D)**
Normalized expression of SOX10 per cell in UMAP space in MM074 cell line. **(E)**
Normalized expression of SOX9 per cell in UMAP space in MM074 cell line. **(F)** Violin
plots of normalized expression of SOX10 per cell. Median is shown as dashed red line.
Wilcoxon Rank Sum Test was used to assess statistical significance, **** p < 0.0001.
(G) Violin plots of normalized expression of SOX9 per cell. Wilcoxon Rank Sum Test
was used to assess statistical significance; ns: no significance. **(H)** UMAP of scRNA-seq
dataset for MM087 cell line from Wouters et al. Cells from siSOX10 and siNT conditions
are labeled. **(I)** Normalized expression of SOX10 per cell in UMAP space in MM087 cell
line. **(J)** Normalized expression of SOX9 per cell in UMAP space in MM087 cell line. **(K)**
Violin plots of normalized expression of SOX10 per cell. Median is shown as dashed red
line. Wilcoxon Rank Sum Test was used to assess statistical significance, **** p <
0.0001. **(L)** Violin plots of normalized expression of SOX9 per cell. Wilcoxon Rank Sum
Test was used to assess statistical significance, **** p < 0.0001.
1270
1271