1 **Identifying *in vivo* genetic dependencies of melanocyte and melanoma**
2 **development**
3

4 Sarah Perlee[1,2], Yilun Ma[1,3,4], Miranda V. Hunter[1], Jacob B. Swanson[5,6], Zhitao Ming[7],
5 Julia Xia[1], Timothée Lionnet[5,6,8], Maura McGrail[7], Richard M. White[1,9, *]
6
[1]Department of Cancer Biology and Genetics, Memorial Sloan Kettering Cancer Center,
New York, NY, USA
[2]Gerstner Sloan Kettering Graduate School of Biomedical Sciences, Memorial Sloan
Kettering Cancer Center, New York, NY, USA.
[3]Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program, Memorial
Sloan Kettering Cancer Center, New York, NY, USA.
[4]Cell and Developmental Biology Program, Weill Cornell Graduate School of Medical
Sciences, New York, NY, USA.
[5]Institute for Systems Genetics, NYU Grossman School of Medicine, New York, NY,
USA.
[6]Department of Cell Biology, NYU Grossman School of Medicine, New York, NY, USA.
[7]Department of Genetics, Development and Cell Biology, Iowa State University, IA,
USA.
[8]Department of Biomedical Engineering, NYU Tandon School of Engineering, Brooklyn,
NY.
[9]Nuffield Department of Medicine, Ludwig Institute for Cancer Research, University of
Oxford, Oxford, UK.
[*] Correspondence to Richard M. White: Richard.white@ludwig.ox.ac.uk

25
Abstract

The advent of large-scale sequencing in both development and disease has identified
large numbers of candidate genes that may be linked to important phenotypes.
Validating the function of these candidates *in vivo* is challenging, due to low efficiency
and low throughput of most model systems. We have developed a rapid, scalable
system for assessing the role of candidate genes using zebrafish. We generated
transgenic zebrafish in which Cas9 was knocked-in to the endogenous *mitfa* locus, a
master transcription factor of the melanocyte lineage. We used this system to identify
both cell-autonomous and non-cell autonomous regulators of normal melanocyte
development. We then applied this to the melanoma setting to demonstrate that loss of
genes required for melanocyte survival can paradoxically promote more aggressive
phenotypes, highlighting that *in vitro* screens can mask *in vivo* phenotypes. Our high-
efficiency genetic approach offers a versatile tool for exploring developmental
processes and disease mechanisms that can readily be applied to other cell lineages.

40
Introduction:
Efficient cell type-specific knockout of genes is crucial for unraveling the intricate
molecular mechanisms underlying cell function, development, and transformation.
Zebrafish (*Danio rerio*) have emerged as a powerful genetic model organism for
studying these processes. The large number of animals that can be easily generated
make it advantageous (compared to other systems like mice) for querying large

numbers of genetic variants that may have links to human diseases. However, most
studies in this system use germline knockout, which makes it difficult to determine
whether the identified gene has a specific function in a particular lineage, or whether the
phenotype reflects a more general developmental defect. Furthermore, many genes
which might have lineage specific functions cannot be studied using germline knockout,
since embryonic lethality often accompanies global knockout of essential genes[1]. To
overcome these limitations, several groups have explored tissue-specific and
conditional knockout strategies in zebrafish[2-10]. For example, the MAZERATI system
uses tissue-specific promoter transgenes to drive Cas9 expression, which are
introduced via plasmid injection into one-cell stage embryos[3,11]. Despite these
advancements, efficiency remains of paramount importance. In normal development,
low-efficiency knockout of a gene can mask important phenotypes, as the mutant cells
can readily be outcompeted by wild-type or inframe mutations. Efficiency is even more
important in the cancer context. Mutation of a tumor suppressor (i.e., PTEN) need not
be highly efficient, since clones with a Darwinian advantage will inevitably outcompete
other clones. However, in situations where we want to study necessity of a tumor
promoting gene *in vivo* (i.e., an oncogene), cells which have mutations of a key driver
gene will easily be outcompeted by nearby clones. Such efficiency is critical as
sequencing projects such as All of Us, the UK Biobank and the Cancer DepMap
continue to identify hundreds of new candidate genes on a regular basis, many of which
have yet to be characterized *in vivo*[12-14].
68
Skin color pigmentation is amongst the most heterogeneous of genetic systems, with a
large number of loci linked to melanocyte development. Mutations in pleiomorphic
genes such as *sox10, ednrb, jam3b, tuba8l3, meox1* and *kit* yield clear defects in
melanocyte/melanophore development, but these mutants also have defects in other
tissues[15-21]. Many of these genes are also expressed in melanoma, but their specific *in*
vivo function in tumorigenesis remains poorly understood.
75
In this study, we wished to develop a highly efficient and scalable system to understand
in vivo genetic dependencies, using melanocytes and melanoma as an exemplar. By
integrating the Cas9 nuclease into the endogenous melanocyte-specific *mitfa* locus, we
achieved highly efficient and specific gene disruption across the melanocyte
developmental lineage. We used our *mitfa*[Cas9] knockin animals to inactivate the
pigmentation gene *albino* and the embryonic essential genes *sox10, tuba1a,* and
ptena/b in melanocytes without additional developmental or off-target phenotypes. We
also demonstrated that this system can be used to efficiently induce melanomas in wild-
type fish by inactivating the tumor suppressors *tp53* and *ptena/b* within melanocytes
also expressing the melanoma oncogene BRAF[V600E]. Inactivating the neural crest
transcription factor *sox10* in these tumors significantly decreased melanoma initiation
but resulted in rare invasive tumors that highly expressed the *sox9* transcription factor,
highlighting that unexpected *in vivo* phenotypes can emerge that aren't predicted from
in vitro studies[22,23]. Our system can thus be used to advance our understanding of
melanocyte biology and the genetic underpinnings of phenotypic switching in
melanoma, an approach that can readily be used for other cell types.
92

Results:

Targeted integration of Cas9 to the *mitfa* locus

To selectively disrupt genes in a melanocyte lineage-specific manner, we established a transgenic zebrafish line harboring Cas9 at the endogenous *mitfa* locus. *Mitfa* expression is detectable by 18 hours postfertilization (hpf) and is highly expressed in pigment cells including melanocytes, xanthophores and their progenitors. To achieve knock-in, we utilized the GeneWeld method, which enables efficient integration of DNA cassettes using homology mediated end joining (HMEJ)[24,25]. The knock-in cassette, containing Cas9 and BFP driven by the eye-specific γ-crystallin promoter, was targeted to exon 2 of the *mitfa* gene, which leads to disruption of the endogenous *mitfa* gene (Figure 1A). We strategically selected this genomic site for several reasons. First, this is the sole exon shared among the three *mitfa* protein-coding transcripts. Furthermore, targeting Cas9 to exon 2 offers the flexibility of replicating this knock-in strategy in *casper* zebrafish, a transparent fish widely used for melanoma studies that harbor a premature stop codon in *mitfa* exon 3[26].

The knock-in vector, along with Cas9 protein and guide RNAs targeting either the *mitfa* integration site or the sequences flanking the 5' and 3' homology arms, were injected into 1-cell stage wild-type Tropical 5D (T5D) zebrafish embryos (Figure 1B)[27]. An average of 19% of injected embryos had detectable ocular BFP expression at 3 days postfertilization (dpf), indicating successful integration (Figure S1A). Embryos were raised to adulthood then crossed with WT T5D fish. The resulting F1 embryos were screened for BFP+ eyes and tail clippings were sequenced to confirm precise integration of the knock-in cassette at the *mitfa* locus. We sequenced four F1 fish and detected on-target integration at the *mitfa* locus in all four fish (Figure S1B). In zebrafish, one copy of *mitfa* is sufficient for the normal development of neural crest-derived melanocytes, but loss of both copies of *mitfa* results in loss of the melanocyte lineage[26,28]. To determine the effect of a single copy of *mitfa*^Cas9 on melanocytes, *mitfa*^Cas9 fish were in-crossed and the resulting siblings were imaged at various time points (Figure S1C). WT and *mitfa*^Cas9 fish exhibited indistinguishable melanocyte patterning during both embryonic and adult stages. However, as expected, fish carrying two copies of Cas9 (*mitfa*^Cas9/Cas9) resembled the *nacre* mutant (*mitfa*-/-), characterized by complete loss of melanocytes, indicating the endogenous *mitfa* gene is disrupted on knock-in alleles[28]. We used zebrafish harboring one knock-in allele (*mitfa*^Cas9) for all subsequent experiments to allow us to test gene function in melanocytes.

To assess whether Cas9 expression was restricted to melanocytes, we performed fluorescence in situ hybridization (FISH) on 3dpf WT and *mitfa*^Cas9 embryos to measure *mitfa* and Cas9 mRNA expression (Figure 1C and S1D). In both WT and *mitfa*^Cas9 fish, *mitfa* was detected within the embryonic melanocyte stripe regions as expected. In *mitfa*^Cas9 fish, Cas9 mRNA was detected only within melanocyte stripe regions also expressing *mitfa*. These results provide functional validation for on-target integration and tissue-restricted expression of Cas9 to the melanocyte lineage.

Melanocyte-specific loss of the human skin color associated gene *slc45a2* results in robust loss of pigmentation

To assess the effectiveness of our stable *mitfa*[Cas9] line, we first targeted the *albino* gene
(*slc45a2*). Large scale GWAS studies have repeatedly identified SNPs in this gene as
tightly linked to human skin color variation, and germline disruption leads to a complete
loss of pigmentation in melanocytes[29-31]. This distinct phenotype enables the visual
observation of Cas9 activity in zebrafish injected with *albino* gRNA. To test our system,
we designed plasmids, denoted MG-gRNA, containing a zU6:gRNA cassette followed
by *mitfa*:GFP to enable visualization of cells expressing the *albino* gRNA. MG-*albino*
and MG-NT were injected into one-cell stage embryos from crosses between *mitfa*[Cas9]
and WT fish, which express Cas9 in all melanocytes but exhibit mosaic gRNA
expression (Figure 2A).
149
Embryos were screened for BFP+ eyes to identify those harboring *mitfa*[Cas9]. Upon
adulthood, 92% of F0 MG-*albino* fish had mosaic loss of pigmentation within the
melanocyte stripe regions (Figure 2B and C). Unpigmented regions maintained GFP
expression (Figure 2B), indicating that the breaks in melanocyte stripes arise from loss
of pigmentation in melanocytes rather than loss of the melanocyte lineage (Figure 2B).
As expected, control MG-NT clutch mates displayed normal pigmentation and GFP+
melanocytes were indistinguishable from GFP- regions on the same fish (Figure 2B and
C).
158
We next aimed to determine the efficiency of *mitfa*[Cas9] in the F1 generation, in which
every melanocyte stably expresses the *albino* gRNA and *mitfa*[Cas9]. To generate F1
lines, we outcrossed the F0 MG-gRNA fish to WT and sorted embryos for BFP+ eyes
and GFP+ melanocytes. F1 MG-*albino* fish exhibited a near complete loss of
pigmentation, demonstrating that our *mitfa*[Cas9] system is highly efficient in F1 fish
(Figure 2D and E). To study the effect of *albino* inactivation across various stages of
development, F1 MG-gRNA fish were once again outcrossed to WT, generating F2 MG-
gRNA fish expressing one copy of *mitfa*[Cas9]. Imaging revealed a measurable
pigmentation loss in F2 MG-*albino* melanocytes as early as 3 days post-fertilization
(dpf) (Figure 2F and G). F2 MG-*albino* fish maintained loss of pigmentation throughout
development, indicating the sustained activity of *mitfa*[Cas9] (Figure S2A-D).
170
To confirm on-target cutting of the *albino* gene in the melanocyte lineage (and not other
tissues), the skin from F1 MG-*albino* fish was dissected and GFP+ and GFP- cells were
isolated using FACS (Figure 2H). GFP+ cells encompass *mitfa*-expressing
melanocytes, xanthophores, and their progenitors, while GFP- cells represent other
non-melanocytic, non-*mitfa* expressing skin cells, such as keratinocytes, fibroblasts, and
immune cells. We performed CRISPR-seq on DNA from both populations to directly
measure Cas9 efficiency and specificity, quantifying the proportion of indels at the
albino locus. Although we have previously shown that FACS sorting melanocytic cells is
possible, it is important to note that we often observe high levels of contaminating
keratinocytes in FACS sorted melanocyte populations, which may lead to an
underestimation of the true allelic frequency in our CRISPR-seq[32]. Despite this, robust
inactivation of the *albino* gene was observed in GFP+ (*mitfa*-expressing) cells,
confirming somatic inactivation (Figure 2I). The majority of indels were frame-shift
inducing mutations. A likely PCR or sequencing artifact resulting in a "T" insertion was

excluded from the frameshift indel percentage as it was observed in all conditions
including WT fish with no pigmentation phenotype (Figure S2E). These results confirm
that our *mitfa*[Cas9] system allows us to inactivate genes within the melanocytic lineage *in*
vivo in a highly efficient and specific manner.

189
The neural crest-related gene *sox10* has specific function in adult melanocyte
patterning and regeneration
Having confirmed that our *mitfa*[Cas9] system allows us to rapidly, efficiently and
specifically manipulate melanocyte gene expression, we next wanted to target
embryonic essential genes. Detecting "negative" gene knockout phenotypes such as
cell death can be challenging *in vivo* due to the selective advantage that WT cells have
over mutants. Current tissue specific knockout systems lack the sensitivity to detect
negative phenotypes.

198
To assess the effectiveness of our *mitfa*[Cas9] model in detecting negative phenotypes, we
utilized it to target *sox10*, a transcription factor vital to the neural crest lineage and
indispensable for overall survival of both zebrafish and mice[15,33]. Consequently, a global
knockout approach is not feasible for studying the function of *sox10* in adult
melanocytes *in vivo*. Germline knockout of *sox10* in zebrafish leads to severe defects in
peripheral nerve development and the complete absence of melanocytes, and the
animals die around day 14[15,34,35]. Due to the complete absence of melanocytes and
their precursor cells, melanoblasts, in *sox10*[-/-] fish, it has not been possible to study the
specific role of *sox10* on these more differentiated cell types *in vivo*. One of the key
unanswered questions is how *sox10* balances its role in maintaining melanoblast
stemness with promoting terminal differentiation, particularly in the context of adult
melanocyte regeneration.

211
We leveraged our *mitfa*[Cas9] system to specifically inactivate *sox10* in the melanocyte
lineage. *mitfa*[Cas9] fish injected with a MG-*sox10* plasmid exhibited noticeable gaps in
their melanocyte stripes (Figure 3A-B). Germline deletion of a *sox10* enhancer region
similarly affects adult stripe patterning in zebrafish[36]. In contrast to the *albino* F0 fish,
the gaps in the stripes of the *sox10* F0 fish were not occupied by unpigmented GFP+
melanocytes, indicating a potential defect in the survival or differentiation of
melanocytes upon loss of *sox10*. This phenotype became even more pronounced in the
stable F1 lines, where several gaps are visible (Figure 3C). Zebrafish have four dark
stripes occupied by melanocytes, which alternate with light colored interstripes occupied
by yellow pigmented xanthophores[37]. Typically, melanocyte stripe 1D (dorsal) is wider
than the interstripe X0 directly below it, as observed in MG-NT F1 fish (Figure 3C and
D). However, the opposite is true for MG-*sox10* F1 fish, where we observed an
apparent widening of the interstripe region and narrowing of the melanocyte stripes
(Figure 3C and D). Melanocytes were counted in fish treated with epinephrine to
aggregate melanosomes, revealing a significant reduction in the number of melanocytes
in the F1 *sox10* KO fish compared to F1 NT fish (Figure 3E). Reduction in GFP+ cells
within the xanthophore interstripes (X0) suggested that *sox10* deficiency also negatively
affected xanthophore populations, in line with previous work indicating a role for *sox10*
in specifying both xanthophores and melanocytes in zebrafish embryos[38].

231
Melanocytes continually regenerate throughout life, typically from a progenitor
population of melanoblasts or melanocyte stem cells. We next assessed whether *sox10*
was required for this. We treated adult fish with neocuproine, a copper chelator
previously shown to kill mature, pigmented melanocytes[39]. We then compared
regeneration from these progenitors in MG-*sox10* and MG-NT F1 fish by counting the
percentage of melanocytes that emerged after 7, 15, and 70 days (Figure 3F). At days 7
and 15, when the melanocytes were still in the process of regenerating from
melanoblasts, there were no significant differences between the two groups. However,
by the time regeneration was completed we found a significant disparity between the
regenerative potential of MG-NT and MG-*sox10* fish. At 70 days post-neocuproine
treatment, MG-NT fish had regenerated an average of 82% of their melanocytes,
whereas only 53% of melanocytes had regenerated in MG-*sox10* fish (Figures 3G-H).
These findings demonstrate that *sox10* is required for adult melanocyte regeneration,
highlighting its requirement within melanoblast populations outside of neural crest
specification.
247
<u>Non-autonomous functions of *tuba1a* on melanocytes</u>
One of the significant challenges of conventional global knockout methods is the
inability to differentiate between cell-autonomous and non-autonomous phenotypes.
This distinction is particularly crucial in melanocyte studies due to their direct and
indirect interactions with diverse cell types. A notable example of genes with pleiotropic
effects is the tubulin gene family, whose α/β tubulin heterodimers form microtubules
essential for various cellular processes such as cell division, motility, and intracellular
transport across various cell types[40]. Microtubules are also critical for the transport of
melanosomes in melanocytic cells[41].
257
Mutation of tubulin genes such as *tuba8l3a* in zebrafish leads to highly pleiotropic
effects including defects in melanocyte patterning, CNS abnormalities, and altered
craniofacial morphology[42]. We chose to focus on the closely related α tubulin gene
tuba1a, which is highly expressed in many cell types of the developing zebrafish
including melanocytes, but whose specific function in these cells remains unexplored[43].
Non-cell type specific knockout of *tuba1a* with the Alt-R CRISPR Cas9 system results in
extensive embryo abnormalities including a curved tail phenotype, pericardial edema,
and melanocytes with more dispersed pigmentation (Figure 4A-B, S3A-B). We found
that compared to 71% survival of NT Alt-R embryos, only 10% of embryos injected with
tuba1a Alt-R survive to 14dpf (Figure 4C).
268
Similar to *sox10,* the embryonic lethality of global *tuba1a* knockout prevents the study of
knockout phenotypes in adult fish. To overcome this limitation, we implemented our
mitfa[Cas9] system to specifically knock out *tuba1a* in melanocytes. Unlike global knockout
of *tuba1a*, melanocyte specific loss of *tuba1a* resulted in viable adult fish with no
obvious abnormalities (Figure S3C-D). Surprisingly, we did not observe the dispersed
melanocyte phenotype in any *mitfa*[Cas9] MG-*tuba1a* embryos (Figure 4D, E). Based on
this observation, we hypothesized that *tuba1a* may be functioning in a cell non-
autonomous manner on melanocytes.

277
The dispersed pigmentation observed in *tuba1a* Alt-R embryos resembles that seen in
blind zebrafish, which fail to contract their melanosomes in response to light[44]. This
light-mediated camouflage involves retinal ganglion cells signaling hypothalamic
neurons to release melanin-concentrating hormone (MCH), which then binds to MCH
receptors on melanocytes, triggering melanosome aggregation via microtubules[44]. The
normal melanosome dispersion observed in *mitfa*[Cas9] MG-*tuba1a* fish suggests that
rather than a cell-intrinsic defect in melanosome aggregation, *tuba1a* may indirectly
influence melanocytes through potential defects in retinal and/or central nervous system
cells crucial for light-mediated camouflage. Interestingly, TUBA1A mutation in humans
can result in both ophthalmologic and brain abnormalities and morpholino-based
knockdown of tuba1a in zebrafish inhibits CNS development[45,46].
289
To further test this idea, we treated *tuba1a* Alt-R embryos with epinephrine to assess
their ability to contract melanosomes in response to a non-visual stimulus. *Tuba1a* Alt-R
embryos exposed to epinephrine had robust aggregation of melanosomes,
demonstrating that the loss of *tuba1a* did not impair melanosome transport (Figure 4F,
G)*. This finding further supports a non-autonomous role for *tuba1a* in melanocytes.
Utilizing the *mitfa*[Cas9] system distinguish cell-autonomous from non-autonomous gene
functions provides valuable insights into the complex dynamics of cell-cell interactions
and how genes function within cellular and organismal networks.
298
Targeting tumor suppressors with *mitfa*[Cas9] induces melanoma
We next turned our attention to melanoma. In humans, tumor suppressors such as
PTEN and *TP53* are somatically inactivated in melanoma[47,48]. In contrast, the most
commonly used zebrafish models of melanoma use a germline *tp53* mutation, which is
not typically seen in humans with the disease and precludes our ability to discern the
role of *tp53* inactivation specifically in melanocytes and melanoma[49]. While global *tp53*
loss is not lethal, it does lead to a wide range of non-melanoma tumors, which can
deteriorate fish health and confound melanoma studies[50]. Furthermore, current
zebrafish melanoma models typically involve the utilization of MiniCoopR *mitfa* rescue
cassettes into *casper* or *nacre* zebrafish, which are devoid of normal pigment cells.
Although this method is highly efficient, it does not accurately represent melanoma
initiation in the context of normal skin architecture and relies on artificial overexpression
of *mitfa*. To mimic human melanoma initiation more accurately, we used our *mitfa*[Cas9]
model to assess tumor initiation using melanoma-specific gene knockout in animals with
wild-type skin. Previous studies have shown that expression of BRAF[V600E] and loss of
tp53 in WT (non-*casper*) fish results in relatively low tumor burden, so we decided to
also target *ptena* and *ptenb*, the zebrafish orthologs of human *PTEN*[51]. *Ptena/ptenb* loss
in combination with *tp53* has been shown to accelerate melanoma formation in
zebrafish[52,53].
318
319 To first address the effect of *ptena/b* loss on normal melanocytes, we generated
320 melanocyte-specific knockouts of *ptena* and *ptenb* (Figure 5A). Like *sox10* and *tuba1a*,
321 germline knockout of *ptena/b* is embryonic lethal[54,55]. In our *ptena/ptenb* F0 melanocyte-
322 specific KO fish, survival was normal. We observed an aberrant expansion of the

melanocytes outside of the stripe regions they normally are restrained to, suggesting that inactivation of *ptena* and *ptenb* induces defects in melanocyte patterning (Figure 5A and B). However, loss of *ptena/b* alone was not sufficient to induce melanoma.

To generate tumors, we co-injected *mitfa*[Cas9] embryos with plasmids encoding *mitfa:*BRAF[V600E] and sgRNAs targeting *tp53* and *ptena/b* (Figure 5C). Larvae were sorted for BFP+ eyes, indicating mitfa-Cas9 integration, and monitored for tumors over 30 weeks. No tumors were detected in BFP- fish from any of the injection conditions (n=136). We also did not observe any tumors in BFP+ fish injected with either BRAF[V600E] alone (n=35) or ptena/b; *tp53* guides in the absence of BRAF[V600E] (n=17). In contrast, tumors developed in all BFP+ groups injected with both *mitfa:*BRAF[V600E] and gRNA plasmids. Approximately 5% of fish from the BRAF;*tp53* group developed tumors (Figure 5D). This aligns with a previous study where 6% of *tp53*-/- fish developed tumors when injected with a BRAF[V600E] construct [51]. Targeting of *ptena* and *ptenb* in *mitfa*[Cas9] fish co-injected with *mitfa*:BRAF[V600E] resulted in a higher tumor incidence of 29% (Figure 5D).

When all three tumor suppressors (*tp53, ptena, ptenb*) were targeted in combination with BRAF[V600E], 59% of fish developed tumors by 30 weeks post fertilization (wpf) (Figure 5D). Immunohistochemistry (IHC) revealed that all tumor samples expressed BRAF[V600E], while p-AKT, a hallmark of *PTEN* inactivation, was only detected in samples with *ptena/b* KO (Figure 5E). To confirm on-target cutting, DNA was extracted from tumors and CRISPR sequencing was performed on PCR amplicons for each tumor suppressor gene. A large proportion of reads contained indels at the target locus for each of the target genes (*ptena*, *ptenb*, and *tp53*) (Figure 5F and S4). Some tumors had an array of mutations, while others appeared to be composed largely of a one dominant clone (Figure S4A to C). Our system thus allows us to rapidly dissect the melanocyte-specific function of both oncogenes (BRAF[V600E]), as well as putative tumor suppressors (*tp53*, *ptena/b*), while avoiding pleiotropic effects of global tumor suppressor loss.

Targeting of lineage-specific oncogenes decreases tumor initiation but promotes progression

Our next objective was to leverage the mitfa[Cas9] fish as a tool for identifying or validating potential genetic dependencies within melanoma. Genetic dependencies can be difficult to study *in vivo* since inactivation of a required gene inevitably leads to selection for non-mutant alleles, as has been previously shown[56]. In melanoma, the DepMap has identified *SOX10* as the top *in vitro* genetic dependency necessary for tumor cell proliferation, suggesting it acts as a lineage-specific oncogene[14,57]. In prior work in both zebrafish and mice, loss of *sox10* is clearly associated with a loss of tumor initiating potential and cell proliferation[56,57]. Yet its specific role *in vivo* remains unclear, since it is required for neural crest and melanocyte specification, making it difficult to know whether its effects reflect loss of the entire lineage[57]. Given our data above showing that *sox10* is required specifically in melanocyte patterning and regeneration, we wished to investigate its function in melanoma.

To assess this, we generated plasmids containing multiple sgRNAs driven by three
distinct zU6 promoters (zU6A, zU6B, and zU6C) to simultaneously target both tumor
suppressors (*tp53, ptena/ptenb*) and tumor promoting genes (*sox10*) in the same cells,
which reduces the chance of selecting for non-mutant alleles[2]. We co-injected these
plasmids with *mitfa*:BRAF^{V600E}, sorted fish for BFP+ eyes, and tracked tumor free
survival over the course of 50 weeks (Figure 6A and B). Consistent with the DepMap
prediction, *sox10* inactivation markedly reduced tumor incidence, which likely reflects
loss of proliferation. Only 3/96 (3.1%) fish with *sox10* knockouts initiated melanomas
compared to the 24/94 fish (25.5%) from the NT condition that developed tumors
(Figure 6B). Using IHC, we validated that *sox10* KO tumors expressed lower levels of
Sox10 protein compared to NT tumors (Figure 6C and S5A). Quantification of Sox10
intensity with immunofluorescence confirmed this finding (Figure S5B).
380
Although *sox10* clearly reduced tumor initiation, we noted "escapers". These tumors
appeared morphologically very distinct compared to the *sox10* intact tumors.
Specifically, these tumors were highly invasive with a mesenchymal morphology
compared to the typical BRAF^{V600E} melanoma (Figure 6C). This observation raised the
possibility that these tumors had undergone "phenotype switching", a form of cell
plasticity in which cells reversibly move between opposite extremes of proliferation
versus invasive states[22,23,57,58]. In melanoma, the proliferative state is thought to be
characterized by high expression of *SOX10*, whereas the mesenchymal, invasive state
is characterized by high expression of *SOX9*[23]. These two highly related transcription
factors have seemingly opposite and possibly antagonistic roles in melanoma. *In vitro*,
knockout of *SOX10* in a variety of human cell lines is associated with acquisition of a
SOX9^{hi} state, which has been suggested to be linked to the invasive phenotype[23]. Re-
analysis of publicly available RNA-seq data using 3 patient-derived human melanoma
cell lines treated with siRNA targeting *SOX10* confirmed this observation, with the
SOX10 lines now becoming *SOX9*^{hi} (Figure 6E to J, S5C to L)[23]. These data raised the
hypothesis that the more invasive tumors we see in our escaper fish was due to gain of
sox9 expression. To test this, we stained our *sox10* CRISPR tumors for *sox9* and
discovered that 2 of the 3 *sox10* KO tumors markedly upregulated *sox9* relative to
control tumors expressing NT gRNA (Figure 6C and D). This data highlights that *in vitro*
genetic studies such as the DepMap, which rely on proliferation as the readout, can
mask more complex *in vivo* phenotypes.
402
Discussion:
The ability to model genetic dependencies *in vivo* is one of the most important uses of
model organisms. Forward genetic mutagenesis screens using compounds such as
ENU or EMS led to the identification of many key developmental genes in organisms
such as Drosophila, C. elegans, Saccharomyces, zebrafish, and many others[59,60].
These approaches have rapidly accelerated in the era of TALENs and CRISPR, readily
allowing for knockout of nearly any gene in a wide variety of non-model organisms.
Cre/Lox approaches have further augmented our ability to discern the cell-type specific
effects of these genes, although these are still laborious and time consuming, especially
in models such as mice[10].
413

Alongside these approaches have been remarkable advances in large scale sequencing
of human tissues. GWAS-like efforts such as the UK Biobank and the US All of Us
programs have identified thousands of germline variants (SNPs) that may be linked to
interesting phenotype variation[12,13]. A conceptually similar approach is also happening
in diseases such as cancer, in which efforts such as the TCGA, ICGC and DepMap
have continued to unveil new potential somatic variants (SNVs) linked to cancer[14,61,62].
420
Across these efforts, in both development and disease, the large number of candidate
variants makes it challenging to connect these genotypes to phenotypes. This is
especially acute in melanoma, where the large number of somatic variants generated by
UV radiation leads to a high number of background mutations that may or may not have
pathogenic function. Thus, there is a substantial need for *in vivo* models that allow for
rapid and efficient modeling of candidate genes. Our development of the *mitfa*-Cas9
knockin line provides such a method. One major advantage is that phenotypes can be
readily screened either in the F0 generation (as mosaics) or in the F1 generation (3
months later). The F1 fish in particular are advantageous, since they allow for highly
efficient biallelic knockout of the gene of interest, allowing for very quick identification of
genes that only have recessive phenotypes (i.e., *albino*). On a per gene basis, this
saves well over 3 months of work compared to germline knockouts and uses vastly
fewer animals since all F1 animals (even those for recessive alleles) are informative.
Additionally, this model eliminates the issue of inbreeding, promoting healthier genetic
lines and facilitating the potential for outcrossing with any available fish line. Because
we demonstrate that the Cas9 is completely restricted to the melanocyte lineage, with
no detectable off-target expression, this easily allows us to discern the melanocyte-
specific effect of pleiotropic genes very efficiently.
439
One interesting observation of our melanoma studies is evidence that loss of melanoma
genetic dependencies like *SOX10* (as suggested by the human DepMap data) can still
lead to tumors, albeit with very different biological behaviors. In both the fish and in the
human samples we analyzed, loss of *SOX10* leads to acquisition of a *SOX9*hi state
instead[23]. *SOX9* is increasingly recognized to be a potent mediator of tumor cell
invasiveness, which is what we observe in our fish[63,64]. One implication of this finding is
that clinical targeting of transcription factors like *SOX10*, long considered an ideal
therapeutic approach, could lead to unexpected outcomes in which tumors might
proliferate less but become more invasive or resistant to therapy. Whether this
phenomenon of *SOX10/SOX9* switching is a general phenomenon, or specific to that
class of TFs, remains to be determined. However, a similar phenomenon has been
observed for *MITF*, another melanocytic TF, suggesting this idea of TF switching needs
to be more thoroughly explored in the future[58,65,66].
453
454 Finally, while our studies clearly focused on the melanocyte lineage, highly similar
455 approaches could be taken for virtually any cellular lineage in which master regulators
456 or markers are known. Given the large number of already identified promoter/enhancer
457 fragments in the zebrafish, this would be an important extension of our system to study
458 gene variants in the context of development and other diseases[67].
459

Our approach has several limitations that may need to be addressed in future studies.
Due to the constitutive nature of the Cas9, we have little control over the timing of its
expression. Because it is knocked into the endogenous *mitfa* locus, the expression of
Cas9 will vary with the endogenous regulation of that gene. In future iterations of this
general approach, swapping the Cas9 for an inducible version would allow for better
timed knockout of genes[68]. Because *mitfa* itself is expressed in melanoblasts,
melanophores and xanthophores during different stages of development, choosing a
promoter (e.g. PMEL) that is more tightly restricted to melanocytes may be
advantageous. Finally, although our system was relatively efficient, we still did not see
100% knockout. This may reflect that it is relatively easy to bypass the CRISPR lesion
in an exon with an in-frame mutation. The use of multiple sgRNAs, or sgRNAs that
target promoter regions, rather than exons, could further augment efficiency of
knockout. Alternatively, instead of targeting DNA, we could consider using RNA
targeting CRISPR enzymes. Recent work has shown that RNA degradation induced by
Cas13 works in zebrafish and may allow for more nuanced knockdown rather than
knockout of a given candidate gene[69,70]. This may especially have advantages in the
context of melanoma, where genes are commonly dysregulated rather than completely
knocked out. We envision it would be relatively straightforward to knock Cas13 into the
locus of interest using an analogous approach to the one we have taken here.
479

Acknowledgements:
We thank members of the White lab for valuable discussions and feedback on this
project. We thank Nelly Cruz Esteves for helpful feedback on the manuscript. We also
thank the MSKCC flow cytometry core for assistance with cell sorting, Integrated
Genomics Operation Core, funded by an NCI Cancer Center Support Grant (P30
CA08748) for sequencing assistance, and Molecular Cytology Core for their help with
imaging. We thank the MSKCC aquatics core facility for their support of this work.
Schematic figures were all made using Biorender.com with a paid Biorender license.
488
Funding:
RMW was funded through the NIH/NCI Cancer Center Support Grant P30 CA008748,
the Melanoma Research Alliance, The Debra and Leon Black Family Foundation, NIH
Research Program Grants R01CA229215 and R01CA238317, NIH Director's New
Innovator Award DP2CA186572, The Pershing Square Sohn Foundation, The Mark
Foundation for Cancer Research, The American Cancer Society, The Alan and Sandra
Gerry Metastasis Research Initiative at the Memorial Sloan Kettering Cancer Center,
The Harry J. Lloyd Foundation, Consano and the Starr Cancer Consortium. This work
was also supported by NIH ORIP R24OD020166 (MM). YM was supported by a Medical
Scientist Training Program grant from the NIH under award number T32GM007739 to
the Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program and
Kirschstein-National Research Service Award (NRSA) predoctoral fellowship under
award number F30CA265124. MVH was funded by a K99/R00 Pathway to
Independence Award from the National Cancer Institute (1K99CA266931). SP was
funded by a Molecular Imaging in Cancer Biology Training Grant at Memorial Sloan
Kettering Cancer Center from the National Cancer institute (T32CA254875) and the

Ruth L. Kirschstein National Research Service Award for Individual Predoctoral Fellows
from the National Cancer Institute (F31CA271518).
507
Author Contributions:
Conceptualization: SP, RMW
Investigation: SP, YM, MVH, JBS, ZM, JX
Visualization: SP, MVH, JBS
Supervision: TL, MM, RMW
Writing—original draft: SP, RMW
Writing—review & editing: SP, MVH, RMW
515
Declaration of Interests:
MM has competing interests with Recombinetics Inc., LifEngine Technologies, and
LifEngine Animal Health Laboratories, Inc. RMW is a paid consultant to N-of-One
Therapeutics, a subsidiary of Qiagen and is on the scientific advisory board of Consano
but receives no income for this. RMW receives royalty payments for the use of the
casper zebrafish line from Carolina Biologicals. SP, YM, MVH, JBS, ZM, JX, and TL
declare no competing interests.
523
524
525 **Methods**
526
527 **Materials Availability:** All data are available in the main text or the supplementary
528 materials, or upon request. Plasmids, fish lines and other materials generated from this
529 study are available upon request.
530
Zebrafish husbandry and ethics statement
All zebrafish experiments adhered to institutional animal protocols and were conducted
in compliance with approved procedures. Fish stocks were maintained at a temperature
of 28.5 °C, under 14:10 light:dark cycles, with pH set at 7.4, and salinity-controlled
conditions. The zebrafish were fed a standard diet comprising brine shrimp followed by
Zeigler pellets. Approval for the animal protocols outlined in this manuscript was
obtained from the Memorial Sloan Kettering Cancer Center (MSKCC) Institutional
Animal Care and Use Committee (IACUC), under protocol number 12-05-008.
Anesthesia was conducted using Tricaine (4 g/l, Syndel, Ferndale, WA, USA) from a
stock of 4 g/l and diluted to 0.16 mg/ml. Adult zebrafish of both sexes were equally
employed in all experiments. Embryos, collected through natural mating, were
incubated in E3 buffer (5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl2, 0.33 mM MgSO4) at
28.5 °C. The wild-type strain used was Tropical 5D Zebrafish (T5D)[27].
544
GeneWeld plasmid construction
A combination of restriction enzyme digestion and HiFi cloning was used to construct
the GeneWeld pPRISM-nCas9n, γcry1:BFP vector for targeted integration to isolate an
nls-Cas9-nls knockin. A 765 bp fragment containing the Porcine teschovirus-1
polyprotein 2A peptide sequence was amplified from the pPRISM-2A-Cre, γcry1:BFP
vector (Addgene #117789) with primers Ori-F and 2A-R[24,25]. The nCas9n cDNA was

amplified from the expression vector pT3TS-nCas9n (Addgene #46757) with primers
Cas9-F and Cas9-R[71]. A KpnI/SpeI fragment containing the pPRISM-2A-Cre, gcry1:BFP
vector backbone was assembled with the 2A and nCas9n PCR amplicons using the
NEBuilder HiFi DNA Assembly Cloning Kit (NEB # E5520S) following the
manufacturer's instructions.
556
Homology arms were designed as previously described[24,25]. Briefly, 48bp homology
arms complementary to the *mitfa* target site were designed using GTagHD
(http://www.genesculpt.org/gtaghd/) for the pPRISM GeneWeld plasmid series. The two
pairs of complementary oligos can be found in Table S1. 5' and 3' arms were cloned
sequentially into the pPRISM-nCas9n, γcry1:BFP vector. The 3' homology arm oligos
were annealed and cloned into the vector using the BspQI restriction site as previously
described. Due to the multiple BfuAI sites within Cas9, instead of using BfuAI restriction
cloning, we alternatively used in-fusion cloning (Takara Bio) to insert a gBlock (IDT)
containing the 5' homology arm into the pPRISM-nCas9n, gcry1:BFP vector. Primers
and gblock sequence used for cloning are listed in Table S1. The 5' and 3' homology
arms were sequenced with primers 5'homologycheckF and R3'_pgtag_seq,
respectively.
569
Injection of GeneWeld reagents
Injections of GeneWeld reagents were performed on one-cell stage embryos. We used
the Alt-R CRISPR-Cas9 system (IDT) for the initial knock-in of nCas9n, γcry1:BFP to
the *mitfa* locus. In brief, 100 µM tracrRNA (IDT 1075928) was mixed at a 1:1 ratio with
either *mitfa* crRNA or universal crRNA then incubated at 95 °C for 5 min. Recombinant
Cas9 (IDT 1081059) was incubated with both sgRNAs for 10 min at 37 °C to form the
RNP complex. The GeneWeld pPRISM-5'mitfaHom-nCas9n, gcry1:BFP-3'mitfaHom
plasmid was then added to the injection mix along with phenol red (Sigma-Aldrich
P0290) . The final concentrations injected into the embryos were: 75 pg/nl Cas9, 12.5
pg/nl *mitfa* sgRNA, 12.5 pg/nl Universal sgRNA, and 5 pg/nl pPRISM plasmid.
580
Genotyping
Tail clips were taken from 2 month postfertilization (mpf) zebrafish and DNA was
extracted using the DNeasy Blood and tissue kit (Qiagen). DNA was PCR amplified with
Q5 high fidelity DNA polymerase (NEB). The forward primer MitfaPCR-F binds to the
mitfa genomic region upstream of the insertion site and the reverse primer
5'homology_amplify-R binds the p2A region of the insertion cassette. Expected PCR
amplicon size is 269bp. Amplicons were sequenced using sanger sequencing (Azenta
Life Sciences).
589
RNA-FISH HCR
We adapted the HCR protocol from Ibarra-Garcia-Padilla et al.[72]. Briefly, embryos were
collected and treated with 0.0045% 1-phenyl 2-thiourea (PTU) (Sigma-Aldrich) after 24
hpf to block pigmentation. Embryos were harvested at 72 hpf and fixed in 4% PFA
(Santa Cruz) for 24 hrs at 4 °C. Embryos were then washed with PBS and
dehydrated/permeabilized with a series of MeOH (Millipore) washes. Embryos were
rehydrated with a series of MeOH/PBST washes, permeabilized with acetone (Fisher

Scientific) and incubated with Proteinase K (Millipore) for 30 min. This was followed by
further fixation with 4% PFA. Embryos were pre-hybridized in probe hybridization buffer
(PHB) before incubating them with HCR probes in PHB at concentration of 20nM. Probe
sequences for *mitfa* and Cas9 can be found in Table S1. Embryos were washed with
probe wash buffer before being pre-amplified with probe amplification buffer (PAB).
Following pre-amplification, amplifier hairpins were thawed, snap-cooled, and mixed
with PAB to a concentration of 36 nM. Embryos were incubated in the hairpin solution
followed by washes with 5×SSCT and 1× PBST. Embryos were stained with 2 µg/mL
DAPI in PBST and mounted on glass slides. Imaging was performed using a Zeiss
LSM880 confocal microscope.
607
Guide RNA plasmids
The following plasmids were constructed using the Gateway Tol2kit:
zU6A:gRNA-NT;zU6B:gRNA-*ptena*;zU6C:gRNA-*tp53*/394,
zU6A:gRNA-NT;zU6B:gRNA-*ptenb*;zU6C:gRNA-*tp53*/394,
zU6A:gRNA-*sox10*;zU6B:gRNA-*ptena*;zU6C:gRNA-*tp53*/394,
zU6A:gRNA-*sox10*;zU6B:gRNA-*ptenb*;zU6C:gRNA-*tp53*/394.
614
In-fusion cloning (Takara Bio) was used to insert *mitfa:*GFP or *mitfa:*TdTomato into a
zU6:gRNA/394 plasmid previously developed in the lab (See Table S1 for primers
used). gRNAs were cloned into zU6:gRNA;*mitfa:*GFP/394 or
zU6:gRNA;*mitfa:*TdTomato/394 using BsmBI sites. Other plasmids used in this study
that were previously developed in the lab include zU6:gRNA-*ptena*/394, zU6:gRNA-
ptenb/394, zU6:gRNA-*tp53*/394, *mitfa*-BRAF[V600E];cmlc2:eGFP. All guide RNAs used in
this study besides *tuba1a* have previously been validated in zebrafish[24,32,56,73].
622
Validation of *tuba1a* sgRNA
The Alt-R CRISPR-Cas9 system (IDT) was used to achieve non-cell type specific
knockout of *tuba1a.* Guide RNAs were designed using ChopChop[74]. In brief, 100 µM
tracrRNA (IDT 1075928) was mixed at a 1:1 ratio with *tuba1a* crRNA then incubated at
95 °C for 5 min. Recombinant Cas9 (IDT 1081059) was incubated with the *tuba1a*
sgRNA for 10 min at 37 °C to form the RNP complex. Phenol red was added prior to
injection into 1 cell-stage wild-type T5D embryos. To validate on-target cutting, 4
embryos were pooled and DNA was extracted using Quick Extract buffer (Fisher
Scientific NC0302740). DNA was PCR amplified with Q5 high fidelity DNA polymerase
(NEB) and primers tuba1a_seq-F and tuba1a_seq-R. Exosap IT for PCR Product Clean
Up was added to PCR product prior to sanger sequencing. Primers tuba1a_seq-F and
tuba1a_seq-R were also used for sequencing.
635
Transgenic lines
To generate MG-U6:sgRNA stable lines, one-cell-stage embryos from crossing WT and
mitfa:Cas9 fish were collected and injected with the 30 pg indicated plasmid and 20 pg
tol2 mRNA. Fish with GFP+ melanocytes and BFP+ eyes were selected and outcrossed
with WT fish to produce the F1 generation. F1 fish were again sorted and outcrossed to
generate F2 generation zebrafish. All fish were imaged on a Zeiss AxioZoom V16 with
Zen 2.1 software.

643
Flow cytometry of melanocytes from adult zebrafish

Zebrafish were euthanized using ice-cold water and dissected using a clean scalpel and forceps. The epidermal and dermal layers of the skin as well as the fins were separated from the rest of the tissues and diced into 1-3mm pieces. Cells were dissociated with Liberase TL (Millipore Sigma 05401020001) and filtered for single cell suspensions as previously described[32]. Samples were then FACS sorted (BD FACSAria) for GFP+ and GFP- cells. WT fish were used as a GFP negative control. Genomic DNA was isolated from sorted cells using the DNeasy Blood and Tissue Kit (Qiagen).

Tumor Dissection

Fish with tumors were euthanized with ice-cold water and immediately dissected with a scalpel to isolate tumor tissue. Genomic DNA was purified from the tumor samples using the Dneasy Blood and tissue kit (Qiagen).

CRISPR sequencing

Primer pairs were designed to produce amplicons 200 to 280 bp in length with the mutation site within 100 bp from the beginning or end of amplicon (see Table S1 for primer sequences). Genomic DNA isolated from methods detailed above was PCR amplified with Q5 high fidelity DNA polymerase (NEB) and run on an agarose gel to visualize PCR products. The samples were then purified using the NucleoSpin Gel and PCR cleanup kit (Takara Bio). Deep sequencing was conducted using the CRISPR-seq platform. Sequencing data was analyzed with CRISPResso2[75]. and indel charts were generated with CRISPRVariants R package[76].

Drug treatments

Adult zebrafish were treated for 24 hr with 750nM neocuproine (Sigma-Aldrich) dissolved in fish water with a final concentration of 0.0075% DMSO as previously described [39]. Cells from the middle melanocyte stripe were counted within a rectangular region of 4mm x 1.5mm. Epinephrine hydrochloride (Sigma-Aldrich) was used at 1 mg/ml in fish water. Fish were treated for 10 minutes prior to imaging.

Zebrafish tumor-free survival curves

To generate a transgenic melanoma model, one-cell-stage embryos from WT X mitfa:Cas9 fish crosses were injected with *mitfa*-BRAF[V600E];cmlc2:eGFP, 20 pg Tol2 mRNA, and the indicated gRNA plasmids for each condition. The total amount of plasmid per embryo did not exceed 30 pg. Embryos were screened for GFP positive hearts and sorted for BFP+ or BFP- eyes. Fish were screened for tumors using a microscope every 2 weeks starting at 4 weeks post fertilization. Tumors were only called if an elevated lesion was observed. Graphpad prism 9 was used to generate and analyze Kaplan-Meier survival curves. Statistical differences were determined using log-rank Mantel–Cox test. All screening and imaging was conducted on a Zeiss AxioZoom V16 using Zen 2.1 software.

Histology

Zebrafish were sacrificed using ice-cold water and placed in 4% paraformaldehyde
(Santa Cruz 30525-89-4) in PBS for 72 h at 4°C on a rocker. Fish were then transferred
to 70% EtOH for 24 h at 4°C on a rocker. Fish were sent to Histowiz, Inc (Brooklyn, NY,
USA), where they were paraffin embedded, sectioned, stained and imaged. All the
stainings were performed at Histowiz, Inc Brooklyn, using the Leica Bond RX automated
stainer (Leica Microsystems) using a Standard Operating Procedure and fully
automated workflow. Samples were processed, embedded in paraffin, and sectioned at
4µm. The slides were dewaxed using xylene and alcohol based dewaxing solutions.
Epitope retrieval was performed by heat-induced epitope retrieval (HIER) of the
formalin-fixed, paraffin-embedded tissue using Citrate based pH 6 solution (Leica
Microsystems, AR9961) for 20 mins at 95 C. The tissues were first incubated with
peroxide block buffer (Leica Microsystems), followed by incubation with the following
antibodies for 30 min: BRAFV600E antibody (ab228461, 1:100), phospho-AKT antibody
(CST4060, 1:50), SOX9 antibody (AB_185230, 1:1000), and SOX10 antibody
(GTX128374, 1:500). This was followed by incubation with DAB secondary reagents:
polymer, DAB refine and hematoxylin (Bond Polymer Refine Detection Kit, Leica
Microsystems) according to the manufacturer's protocol. The slides were dried,
coverslipped (TissueTek-Prisma Coverslipper) and visualized using a Leica Aperio AT2
slide scanner (Leica Microsystems) at 40X.
707
708

Quantification of Sox10 intensity
FFPE slides were deparaffinized by consecutive incubations in xylene and 100-50%
ethanol. For antigen retrieval, slides were placed in 10 mM sodium citrate pH 6.2 in a
pressure cooker and heated to 95°C for 20 minutes. Slides were then cooled to room
temperature and blocked in 5% donkey serum, 1% BSA, and 0.4% Triton-X100 in PBS
for 1 hour at room temperature. The primary antibody targeting Sox10 (GeneTex,
GTX128374) was added to the blocking buffer at 1:200 concentration and incubated at
4°C overnight. Slides were then washed in PBS before incubation with the secondary
antibody (donkey anti-rabbit Alexa Fluor Plus 488, Thermo Fisher Scientific #AC32790;
1:250) and Hoescht (Thermo Fisher Scientific, #62249; 1:1000) in blocking buffer for 2
hours at room temperature. Slides were mounted in Vectashield Plus (Vector Labs, H-
1900). Slides were imaged on a LSM 880 confocal microscope using a 40X oil
immersion objective. 3 images were captured per sample. Quantification of Sox10
intensity per cell was performed using CellProfiler.[77] and MATLAB r2023b (Mathworks).
Cells were segmented in CellProfiler to obtain a nuclear mask from the Hoescht signal
and using this mask the mean intensity per nucleus was quantified.
725

Re-analysis of human melanoma scRNA-seq data
Human melanoma scRNA-seq data from Wouters et al. was accessed from GEO
(GSE134432)[23]. All analysis was done in R (version 4.3.1) using the Seurat package
(version 5.0.2)[78]. Counts matrices for each sample and cell line were imported into R
and converted into Seurat objects before merging into one combined Seurat object.
Counts were normalized using the Seurat function NormalizeData with default
parameters. UMAP was calculated using the Seurat function RunUMAP with 10
dimensions.

Statistical analysis

Statistical significance was analyzed using t test, Log-rank (Mantel-Cox) test, Wilcoxon Rank Sum Test or as indicated in the figure legends. GraphPad Prism 9 software was used for data processing and statistical analysis. *P < 0.05; **P < 0.005; ***P < 0.001; ****P < 0.0001. Data are presented as means ± SD unless otherwise indicated.

References

1. Amsterdam, A. *et al.* Identification of 315 genes essential for early zebrafish development. *Proc Natl Acad Sci U S A* **101**, 12792-7 (2004).
2. Yin, L. *et al.* Multiplex Conditional Mutagenesis Using Transgenic Expression of Cas9 and sgRNAs. *Genetics* **200**, 431-41 (2015).
3. Ablain, J., Durand, E.M., Yang, S., Zhou, Y. & Zon, L.I. A CRISPR/Cas9 vector system for tissue-specific gene disruption in zebrafish. *Dev Cell* **32**, 756-64 (2015).
4. Zhou, W. *et al.* Neutrophil-specific knockout demonstrates a role for mitochondria in regulating neutrophil motility in zebrafish. *Dis Model Mech* **11**(2018).
5. Ni, T.T. *et al.* Conditional control of gene function by an invertible gene trap in zebrafish. *Proc Natl Acad Sci U S A* **109**, 15389-94 (2012).
6. Grajevskaja, V., Camerota, D., Bellipanni, G., Balciuniene, J. & Balciunas, D. Analysis of a conditional gene trap reveals that tbx5a is required for heart regeneration in zebrafish. *PLoS One* **13**, e0197293 (2018).
7. Burg, L. *et al.* Conditional mutagenesis by oligonucleotide-mediated integration of loxP sites in zebrafish. *PLoS Genet* **14**, e1007754 (2018).
8. Kalvaityte, M. & Balciunas, D. Conditional mutagenesis strategies in zebrafish. *Trends Genet* **38**, 856-868 (2022).
9. Singh Angom, R., Wang, Y., Wang, E., Dutta, S.K. & Mukhopadhyay, D. Conditional, Tissue-Specific CRISPR/Cas9 Vector System in Zebrafish Reveals the Role of Nrp1b in Heart Regeneration. *Arterioscler Thromb Vasc Biol* **43**, 1921-1934 (2023).
10. Shin, M. *et al.* Generation and application of endogenously floxed alleles for cell-specific knockout in zebrafish. *Dev Cell* **58**, 2614-2626 e7 (2023).
11. Ablain, J., Liu, S., Moriceau, G., Lo, R.S. & Zon, L.I. SPRED1 deletion confers resistance to MAPK inhibition in melanoma. *J Exp Med* **218**(2021).
12. Mayer, C.S. & Huser, V. Learning important common data elements from shared study data: The All of Us program analysis. *PLoS One* **18**, e0283601 (2023).
13. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* **12**, e1001779 (2015).
14. Tsherniak, A. *et al.* Defining a Cancer Dependency Map. *Cell* **170**, 564-576 e16 (2017).
15. Dutton, K.A. *et al.* Zebrafish colourless encodes sox10 and specifies non-ectomesenchymal neural crest fates. *Development* **128**, 4113-25 (2001).
16. Parichy, D.M. *et al.* Mutational analysis of endothelin receptor b1 (rose) during neural crest and pigment pattern development in the zebrafish Danio rerio. *Dev Biol* **227**, 294-306 (2000).

780   17.   Eom, D.S., Patterson, L.B., Bostic, R.R. & Parichy, D.M. Immunoglobulin
781   superfamily receptor Junctional adhesion molecule 3 (Jam3) requirement for
782   melanophore survival and patterning during formation of zebrafish stripes. *Dev*
783   *Biol* **476**, 314-327 (2021).
784   18.   Parichy, D.M. & Turner, J.M. Zebrafish puma mutant decouples pigment pattern
785   and somatic metamorphosis. *Dev Biol* **256**, 242-57 (2003).
786   19.   Nguyen, P.D. *et al.* Haematopoietic stem cell induction by somite-derived
787   endothelial cells controlled by meox1. *Nature* **512**, 314-8 (2014).
788   20.   Irion, U., Singh, A.P. & Nusslein-Volhard, C. The Developmental Genetics of
789   Vertebrate Color Pattern Formation: Lessons from Zebrafish. *Curr Top Dev Biol*
790   **117**, 141-69 (2016).
791   21.   Parichy, D.M., Rawls, J.F., Pratt, S.J., Whitfield, T.T. & Johnson, S.L. Zebrafish
792   sparse corresponds to an orthologue of c-kit and is required for the
793   morphogenesis of a subpopulation of melanocytes, but is not essential for
794   hematopoiesis or primordial germ cell development. *Development* **126**, 3425-36
795   (1999).
796   22.   Capparelli, C. *et al.* Targeting SOX10-deficient cells to reduce the dormant-
797   invasive phenotype state in melanoma. *Nat Commun* **13**, 1381 (2022).
798   23.   Wouters, J. *et al.* Robust gene expression programs underlie recurrent cell states
799   and phenotype switching in melanoma. *Nat Cell Biol* **22**, 986-998 (2020).
800   24.   Welker, J.M. *et al.* GeneWeld: Efficient Targeted Integration Directed by Short
801   Homology in Zebrafish. *Bio Protoc* **11**, e4100 (2021).
802   25.   Wierson, W.A. *et al.* Efficient targeted integration directed by short homology in
803   zebrafish and mammalian cells. *Elife* **9**(2020).
804   26.   White, R.M. *et al.* Transparent adult zebrafish as a tool for in vivo transplantation
805   analysis. *Cell Stem Cell* **2**, 183-9 (2008).
806   27.   Balik-Meisner, M., Truong, L., Scholl, E.H., Tanguay, R.L. & Reif, D.M.
807   Population genetic diversity in zebrafish lines. *Mamm Genome* **29**, 90-100
808   (2018).
809   28.   Lister, J.A., Robertson, C.P., Lepage, T., Johnson, S.L. & Raible, D.W. nacre
810   encodes a zebrafish microphthalmia-related protein that regulates neural-crest-
811   derived pigment cell fate. *Development* **126**, 3757-67 (1999).
812   29.   Fernandez, L.P. *et al.* SLC45A2: a novel malignant melanoma-associated gene.
813   *Hum Mutat* **29**, 1161-7 (2008).
814   30.   Branicki, W., Brudnik, U., Draus-Barini, J., Kupiec, T. & Wojas-Pelc, A.
815   Association of the SLC45A2 gene with physiological human hair colour variation.
816   *J Hum Genet* **53**, 966-971 (2008).
817   31.   Streisinger, G., Singer, F., Walker, C., Knauber, D. & Dower, N. Segregation
818   analyses and gene-centromere distances in zebrafish. *Genetics* **112**, 311-9
819   (1986).
820   32.   Weiss, J.M. *et al.* Anatomic position determines oncogenic specificity in
821   melanoma. *Nature* **604**, 354-361 (2022).
822   33.   Hou, L., Arnheiter, H. & Pavan, W.J. Interspecies difference in the regulation of
823   melanocyte development by SOX10 and MITF. *Proc Natl Acad Sci U S A* **103**,
824   9081-5 (2006).

34. Kelsh, R.N. *et al.* Zebrafish pigmentation mutations and the processes of neural
crest development. *Development* **123**, 369-89 (1996).
35. Carney, T.J. *et al.* A direct role for Sox10 in specification of neural crest-derived
sensory neurons. *Development* **133**, 4619-30 (2006).
36. Cunningham, R.L. *et al.* Functional in vivo characterization of sox10 enhancers in
neural crest and melanoma development. *Commun Biol* **4**, 695 (2021).
37. Owen, J.P., Kelsh, R.N. & Yates, C.A. A quantitative modelling approach to
zebrafish pigment pattern formation. *Elife* **9**(2020).
38. Nagao, Y. *et al.* Distinct interactions of Sox5 and Sox10 in fate specification of
pigment cells in medaka and zebrafish. *PLoS Genet* **14**, e1007260 (2018).
39. O'Reilly-Pol, T. & Johnson, S.L. Neocuproine ablates melanocytes in adult
zebrafish. *Zebrafish* **5**, 257-64 (2008).
40. Cushion, T.D., Leca, I. & Keays, D.A. MAPping tubulin mutations. *Front Cell Dev*
Biol **11**, 1136699 (2023).
41. Rogers, S.L., Tint, I.S., Fanapour, P.C. & Gelfand, V.I. Regulated bidirectional
motility of melanophore pigment granules along microtubules in vitro. *Proc Natl*
Acad Sci U S A **94**, 3720-5 (1997).
42. Larson, T.A., Gordon, T.N., Lau, H.E. & Parichy, D.M. Defective adult
oligodendrocyte and Schwann cell development, pigment pattern, and
craniofacial morphology in puma mutant zebrafish having an alpha tubulin
mutation. *Dev Biol* **346**, 296-309 (2010).
43. Farnsworth, D.R., Saunders, L.M. & Miller, A.C. A single-cell transcriptome atlas
for zebrafish development. *Dev Biol* **459**, 100-108 (2020).
44. Neuhauss, S.C. *et al.* Genetic disorders of vision revealed by a behavioral screen
of 400 essential loci in zebrafish. *J Neurosci* **19**, 8603-15 (1999).
45. Myers, K.A., Bello-Espinosa, L.E., Kherani, A., Wei, X.C. & Innes, A.M. TUBA1A
Mutation Associated With Eye Abnormalities in Addition to Brain Malformation.
Pediatr Neurol **53**, 442-4 (2015).
46. Veldman, M.B., Bemben, M.A. & Goldman, D. Tuba1a gene expression is
regulated by KLF6/7 and is necessary for CNS development and regeneration in
zebrafish. *Mol Cell Neurosci* **43**, 370-83 (2010).
47. Roh, M.R. *et al.* Promoter Methylation of PTEN Is a Significant Prognostic Factor
in Melanoma Survival. *J Invest Dermatol* **136**, 1002-1011 (2016).
48. Palmieri, G. *et al.* Multiple Molecular Pathways in Melanomagenesis:
Characterization of Therapeutic Targets. *Front Oncol* **5**, 183 (2015).
49. Frantz, W.T. & Ceol, C.J. From Tank to Treatment: Modeling Melanoma in
Zebrafish. *Cells* **9**(2020).
50. Ignatius, M.S. *et al.* tp53 deficiency causes a wide tumor spectrum and increases
embryonal rhabdomyosarcoma metastasis in zebrafish. *Elife* **7**(2018).
51. Patton, E.E. *et al.* BRAF mutations are sufficient to promote nevi formation and
cooperate with p53 in the genesis of melanoma. *Curr Biol* **15**, 249-54 (2005).
52. He, S. *et al.* Synergistic melanoma cell death mediated by inhibition of both
MCL1 and BCL2 in high-risk tumors driven by NF1/PTEN loss. *Oncogene* **40**,
5718-5729 (2021).

53. Montal, E., Suresh, S., Ma, Y., Tagore, M.M. & White, R.M. Cancer Modeling by
Transgene Electroporation in Adult Zebrafish (TEAZ). *Methods Mol Biol* **2707**,
83-97 (2024).
54. Croushore, J.A. *et al.* Ptena and ptenb genes play distinct roles in zebrafish
embryogenesis. *Dev Dyn* **234**, 911-21 (2005).
55. Faucherre, A., Taylor, G.S., Overvoorde, J., Dixon, J.E. & Hertog, J. Zebrafish
pten genes have overlapping and non-redundant functions in tumorigenesis and
embryonic development. *Oncogene* **27**, 1079-86 (2008).
56. Kaufman, C.K. *et al.* A zebrafish melanoma model reveals emergence of neural
crest identity during melanoma initiation. *Science* **351**, aad2197 (2016).
57. Shakhova, O. *et al.* Antagonistic cross-regulation between Sox9 and Sox10
controls an anti-tumorigenic program in melanoma. *PLoS Genet* **11**, e1004877
(2015).
58. Hoek, K.S. *et al.* In vivo switching of human melanoma cells between proliferative
and invasive states. *Cancer Res* **68**, 650-6 (2008).
59. Solnica-Krezel, L., Schier, A.F. & Driever, W. Efficient recovery of ENU-induced
mutations from the zebrafish germline. *Genetics* **136**, 1401-20 (1994).
60. Russell, W.L. *et al.* Specific-locus test shows ethylnitrosourea to be the most
potent mutagen in the mouse. *Proc Natl Acad Sci U S A* **76**, 5818-9 (1979).
61. Cerami, E. *et al.* The cBio cancer genomics portal: an open platform for exploring
multidimensional cancer genomics data. *Cancer Discov* **2**, 401-4 (2012).
62. Alexandrov, L.B. *et al.* Signatures of mutational processes in human cancer.
Nature **500**, 415-21 (2013).
63. Cheng, P.F. *et al.* Methylation-dependent SOX9 expression mediates invasion in
human melanoma cells and is a negative prognostic factor in advanced
melanoma. *Genome Biol* **16**, 42 (2015).
64. Yang, X. *et al.* SOX9 is a dose-dependent metastatic fate determinant in
melanoma. *J Exp Clin Cancer Res* **38**, 17 (2019).
65. Hossain, S.M. & Eccles, M.R. Phenotype Switching and the Melanoma
Microenvironment; Impact on Immunotherapy and Drug Resistance. *Int J Mol Sci*
24(2023).
66. Travnickova, J. *et al.* Zebrafish MITF-Low Melanoma Subtype Models Reveal
Transcriptional Subclusters and MITF-Independent Residual Disease. *Cancer*
Res **79**, 5769-5784 (2019).
67. Weiss, J.M. *et al.* Shifting the focus of zebrafish toward a model of the tumor
microenvironment. *Elife* **11**(2022).
68. Sun, N. *et al.* Development of drug-inducible CRISPR-Cas9 systems for large-
scale functional screening. *BMC Genomics* **20**, 225 (2019).
69. Huang, Y. *et al.* CRISPR-dCas13-tracing reveals transcriptional memory and
limited mRNA export in developing zebrafish embryos. *Genome Biol* **24**, 15
(2023).
70. Kushawah, G. *et al.* CRISPR-Cas13d Induces Efficient mRNA Knockdown in
Animal Embryos. *Dev Cell* **54**, 805-817 e7 (2020).
71. Jao, L.E., Wente, S.R. & Chen, W. Efficient multiplex biallelic zebrafish genome
editing using a CRISPR nuclease system. *Proc Natl Acad Sci U S A* **110**, 13904-
9 (2013).

72. Ibarra-Garcia-Padilla, R., Howard, A.G.A.t., Singleton, E.W. & Uribe, R.A. A
protocol for whole-mount immuno-coupled hybridization chain reaction
(WICHCR) in zebrafish embryos and larvae. *STAR Protoc* **2**, 100709 (2021).
73. Suresh, S. *et al.* Identifying the Transcriptional Drivers of Metastasis Embedded
within Localized Melanoma. *Cancer Discov* **13**, 194-215 (2023).
74. Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond
genome editing. *Nucleic Acids Res* **47**, W171-W174 (2019).
75. Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing
sequence analysis. *Nat Biotechnol* **37**, 224-226 (2019).
76. Lindsay, H. *et al.* CrispRVariants charts the mutation spectrum of genome
engineering experiments. *Nat Biotechnol* **34**, 701-2 (2016).
77. Stirling, D.R. *et al.* CellProfiler 4: improvements in speed, utility and usability.
BMC Bioinformatics **22**, 433 (2021).
78. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-
cell analysis. *Nat Biotechnol* **42**, 293-304 (2024).
930

Figure 1. Targeted integration of Cas9 to the *mitfa* locus with GeneWeld knockin system.
(**A**) Schematic depicting integration site and GeneWeld knock-in cassette. 5' and 3' homology
arms are designed to target exon 2 of the zebrafish *mitfa* gene. The knock-in cassette includes
p2A followed by Cas9 and BFP driven by the eye-specific promoter gamma-crystallin. (**B**)
Pipeline to produce F1 mitfa^Cas9 zebrafish. The GeneWeld knock-in vector and gRNAs targeting
the *mitfa* genomic insertion site or specific sites on the knock-in vector (UgRNA) are co-injected
into one-cell stage wild-type zebrafish embryos. Embryos are screened for BFP+ eyes marking
mosaic integration. These mosaic founder fish are then raised to adulthood and crossed with
wild-type fish. The resulting embryos are screened for BFP+ eyes and sequenced to confirm
precise integration. (**C**) Fluorescent in situ hybridization chain reaction (FISH) on 3dpf wild-type
and *mitfa*^Cas9 embryos treated with 1-phenyl 2-thiourea (PTU). The presence of *mitfa* and Cas9
RNA was assessed by confocal microscopy at 40x magnification. Maximum intensity projections
are shown. n=17 WT embryos and n=16 mitfa^Cas9 embryos were screened. Scale bars, 100μm.

Figure 2. Melanocyte lineage-specific knockout of *albino* using *mitfa*^Cas9 fish.
(A) Pipeline to generate F0 and F1 *mitfa*:GFP;U6:gRNA (MG-gRNA) zebrafish. **(B)** Adult
mitfa^Cas9 MG-NT and MG-*albino* F0 fish. **(C)** Proportion of MG-NT (n=19) and MG-*albino* (n=25)
F0 fish with loss of pigmentation phenotype. **(D)** Adult mitfa^Cas9 MG-NT and MG-*albino* F1 fish.
(E) Pigmented area/ mm^2 calculated for n=5 fish/ genotype within a defined rectangular region of
interest (ROI) encompassing the top and middle melanocyte stripes. Two-sided student's *t*-test
was used to assess statistical significance, **** p < 0.0001; error bars, SD. **(F)** 3dpf mitfa^Cas9
MG-NT and MG-*albino* F2 fish. Representative embryos are shown for each genotype. **(G)**
Mean grey value of head melanocytes calculated for n=10 embryos/genotype within a defined
hexagonal ROI indicated as a red outline in 2F. Two-sided student's *t*-test was used to assess
statistical significance, **** p < 0.0001. **(H)** Schematic for CRISPR sequencing protocol. **(I)**
CRISPR-seq results are shown for WT and n=3 independent F1 MG-*albino* fish. Only GFP
negative cells were isolated from WT fish. Results are shown as a fraction of sequences with
indels calculated using CRISPRESSO.

Figure 3. Melanocyte lineage-specific knockout of *sox10* using *mitfa*Cas9 fish.
(A) Adult mitfaCas9 MG-NT and MG-*sox10* F0 fish. **(B)** Proportion of MG-NT (n=19) and MG-*sox10* (n=16) F0 fish with disrupted stripes phenotype. **(C)** Adult mitfaCas9 MG-NT and MG-*sox10* F1 fish. **(D)** The average width of melanocyte stripe 1D and xanthophore interstripe X0 was calculated for each fish by averaging 5 stripe/interstripe width measurements. N=5 fish per

genotype. Two-sided student's *t*-test was used to assess statistical significance, *** p < 0.001;
error bars, SD. **(E)** F1 MG-NT (n=5) and MG-*sox10* (n=5) adult fish were treated with
epinephrine and melanocytes were counted within a defined rectangular region of interest
3.46mm x 2.54mm encompassing the top and middle melanocyte stripes. Two-sided student's *t*-
test was used to assess statistical significance, * p < 0.05; error bars, SD. **(F)** Schematic of
neocuproine (neo) experimental setup. Adult mitfa^{Cas9} MG-NT (n= 5) and MG-*sox10* (n=5) F1
fish were treated with neocuproine for 24 hours to ablate melanocytes then imaged at day 7,15,
and 70 to measure regeneration of melanocytes compared to day 0. **(G)** Representative images
are shown for MG-NT and MG-*sox10* fish pre- and post-neocuproine treatment. **(H)**
Quantification of melanocyte regeneration. Fish were treated with epinephrine prior to imaging
to enable counting of melanocytes. Two-sided student's *t*-test was used to assess statistical
significance, ** p < 0.01; error bars, SD.
979
980

981
982
983
984

985
Figure 4. Non-autonomous function of *tuba1a* on melanocytes.
(A) 4dpf zebrafish embryos injected with either NT or *tuba1a* Alt-R CRISPR Cas9 gRNAs. **(B)**
Percentage of NT or *tuba1a* Alt-R zebrafish embryos with dispersed melanocyte phenotypes.
N=2 independent experiments. **(C)** Survival percentage is shown for NT and *tuba1a* Alt-R
embryos. Embryos were counted at 24hpf and again at 14dpf to determine survival. **(D)** 4dpf
mitfa^{Cas9} MG-*tuba1a* F2 embryos (BFP+ eyes) compared to sibling controls with no mitfa^{Cas9}
(BFP- eyes). Representative embryos are shown for each genotype. **(E)** Pigmented area/ mm^2
calculated for n=19 embryos/genotype from two independent MG-*tuba1a* F2 clutches. Two-
sided student's *t*-test was used to assess statistical significance, ns: no significance. **(F)** 4dpf
Alt-R injected zebrafish embryos imaged before and after epinephrine (epi) treatment. **(G)**
Pigmented area/ mm^2 calculated for n=20 embryos/genotype from two independent clutches.
Two-sided student's *t*-test was used to assess statistical significance, **** p < 0.0001.

Figure 5. Generation of a zebrafish melanoma model in *mitfa*^{Cas9} fish.
(A) Adult mitfa^{Cas9} MG-NT; *mitfa*:TdT;U6:NT (MTdt-NT) and MG-*ptena*; MTdT-*ptenb* F0 fish. **(B)**
Proportion of MG-NT; MTdT-NT (n=14) and MG-*ptena*; MTdT-*ptenb* (n=14) F0 fish with
disrupted stripes phenotype. **(C)** Schematic of zebrafish tumorigenesis assay. Indicated
plasmids are injected into one-cell stage embryos from crosses between mitfa^{Cas9} and WT fish.
The mitfa:BRAF^{V600E} plasmid includes cardiac-specific cmlc2:GFP. Embryos were sorted for
GFP+ hearts and BFP+ eyes and fish were screened every 2 weeks for tumors. **(D)** Tumor free
survival curve. N=2 independent experiments. Tumors were tracked over the course of 30
weeks. Log-rank (Mantel-Cox) test was used to assess statistical significance, *** p < 0.001; ****
p < 0.0001; ns: no significance. **(E)** Histology was performed on one fish from each indicated
injection group. Dotted lines indicate site of sectioning. Red chromogen was used for all IHC
staining. Scale bars, 50μm. **(F)** CRISPR-seq results are shown for normal skin dissected from
WT fish and tumors dissected from injection conditions 2, 3, and 4. Results are shown as a
fraction of sequences with indels calculated using CRISPRESSO.

1013
Figure 6. Melanoma specific knockout of *sox10* reduces tumor burden and induces
phenotypic switching.
(A) Schematic of zebrafish tumorigenesis assay. Indicated plasmids are injected into one-cell
stage embryos from crosses between mitfa^{Cas9} and WT fish. Embryos were sorted for GFP+
hearts and BFP+ eyes and fish were screened every 2 weeks for tumors. **(B)** Tumor free
survival curve. N=3 independent experiments. Tumors were tracked over the course of 50
weeks. Log-rank (Mantel-Cox) test was used to assess statistical significance, **** p < 0.0001.
(C) Histology is shown for one fish from each injection group. Dotted lines indicate site of
sectioning. Red chromogen was used for all IHC staining. Scale bars, 50µm. **(D)** Color
thresholding was used on IHC images to calculate the percentage of nuclei that stained positive
for *sox9*. NT-1 n=1222 cells, NT-2 n=1620 cells, NT-3 n=2185 cells, sox10-1 n=970 cells,
sox10-2 n=594, sox10-3 n=1046. Cells from n=3 images were analyzed for each tumor. **(E)**
Schematic of scRNA-seq experimental setup from Wouters et al. Patient derived cell lines were
treated with siRNAs targeting SOX10 or NTC and scRNA-seq was conducted at 72hrs. **(F)**
UMAP of scRNA-seq dataset for MM057 cell line from Wouters et al. Cells from siSOX10 and
siNT conditions are labeled. **(G)** Normalized expression of Sox10 per cell in UMAP space. **(H)**
Violin plots of normalized expression of Sox10 per cell. Median is shown as dashed red line.
Wilcoxon Rank Sum Test was used to assess statistical significance, **** p < 0.0001. **(I)**
Normalized expression of Sox9 per cell in UMAP space. **(J)** Violin plots of normalized
expression of Sox9 per cell. Wilcoxon Rank Sum Test was used to assess statistical
significance, **** p < 0.0001.