Basal ganglia output coding - entopeduncular nucleus - of contextual kinematics and
reward in the freely moving mouse

Abstract

The entopeduncular nucleus (EPN) is often termed as one of the output nuclei of the
basal ganglia owing to their highly convergent anatomy. The rodent EPN has been
implicated in reward and value coding whereas the primate analogue internal Globus
Pallidus has been found to be modulated by some movements and in some
circumstances. In this study we sought to understand how the rodent EPN might be
coding kinematic, reward, and difficulty parameters, particularly during locomotion.
Furthermore, we aimed to understand the level of movement representation: whole-body
or specific body parts. To this end, mice were trained in a freely moving two-alternative
forced choice task with two periods of displacement (return and go trajectories) and
performed electrophysiological recordings together with video-based tracking. We found
1) robust reward coding but not difficulty. 2) Spatio-temporal variables better explain EPN
activity during movement compared to kinematic variables, while both types of variables
were more robustly represented in reward-related movement. 3) Reward sensitive units
encode kinematics similarly to reward insensitive ones. 4) Population dynamics that best
account for differences between these two periods of movement can be explained by
allocentric references like distance to reward port. 5) The representation of paw and licks
is not mutually exclusive, discarding a somatotopic muscle-level representation of
movement in the EPN. Our data suggest that EPN activity represents movements and
reward in a complex way: highly multiplexed, influenced by the objective of the
displacement, where trajectories that lead to reward better represent spatial and
kinematic variables. Interestingly, there are intertwining representations of whole-body
movement kinematics with single paw and licking variables. Further, reward sensitive
units encode kinematics similarly to reward insensitive ones, challenging the notion of
distinct pathways for reward and movement processing.

Significance Statement

The entopeduncular nucleus is one of the main outputs of the basal ganglia whose activity
has been hypothesized to be inversely correlated with movement. This study examines
motor and reward coding simultaneously, finding that besides the great level of
multiplexing of these variables, spatio-temporal coding is better represented than
kinematic coding. The level of movement representation seems to be greatly influenced
by the goal of a movement, with spatially biased variables influencing the population
dynamics of this nucleus. Further, we uncover the coexistence of EPN modulation by
movement at different timescales and body parts. The simple overall activity of this output
nucleus cannot explain kinematic coding, challenging leading theories of basal ganglia
function.

Introduction

The entopeduncular nucleus (EPN) is often termed as the output of the basal ganglia. The basal ganglia are a group of subcortial nuclei with highly converging anatomy that have been associated with motor and non-motor functions. Whereas the striatum receives inputs from many diverse brain regions (Hintiryan et al., 2016; Hunnicutt et al., 2016) most of its projections end in the internal Globus Pallidus (GPi, primate analogue of EPN) or substantia nigra pars reticulata (SNr, often termed as the output nuclei) through the direct and indirect pathways. This is accompanied by a sharp reduction in the number of neurons: the striatum has 2-3 orders of magnitude more neurons. This highly convergent anatomy suggests information synthesis/convergence.

Previous studies focusing on the non-motor coding of the internal Globus Pallidus, have found that there is coding of reward (Hong & Hikosaka, 2008). Rodent studies have corroborated this finding and suggested the existence of value coding that serves to evaluate actions (Stephenson-Jones et al., 2016).

There is a particular interest in understanding the role of the basal ganglia on movement production since several diseases with motor disturbances have been linked to basal ganglia malfunction. Several studies have found a relationship between the GPi activity and limb movements of the primate. However, studies have found inconsistencies on the coding on movement depending on movement type (ramping vs. stepped, self-paced vs step tracking, (Georgopoulos et al., 1983)) or between identical movements performed under different cognitive states (cued vs memory-dependent (Turner & Anderson, 2005)). Further, some studies agree that self-paced movements are represented in a weaker manner (Mink & Thach, 1991a; Turner & Anderson, 2005). Overall, findings show that GPi units are selectively engaged in some types of movement, but it is so far unclear what cognitive contingency is most engaging.

On the other hand, current theories of the overall function of the basal ganglia, like the 'rate theory' (Albin et al., 1989)or the "dynamic activity model" (Nambu et al., 2023), posit that the firing rate of the output nuclei is inversely related to movement facilitation.

In this study we sought to examine how the information regarding kinematics, particularly during locomotion, was represented in the entopeduncular nucleus of freely moving mice. Given that previous studies have highlighted the importance of reward related activity, we further sought to understand whether reward sensitive units would be insensitive to kinematic coding. Finally, we investigated whether the EPN units were modulated by individual movements (limb or licking) as opposed to whole body movements.

Materials and methods

Animals. All procedures were approved by the Institutional Committee for the Care and Use of Laboratory Animals at the Instituto de Fisiología Celular (Protocol number FTA-121-17). Universidad Nacional Autónoma de México. and the National Norm for the Use of Animals (NOM-062-ZOO-1999). Experiments used C57BL/6J male (n=2) and female (n=4) mice (2-3 months of age at the start of training, 6-7 months during recordings). Animals were housed under a 12h light/dark cycle (lights on at 6am) with ad libitum access to food and water before the start of behavioral experiments. For behavioral

training, water was restricted so that the weight of animals was 80% of their original
weight.

Behavioral apparatus and task design. The behavioral apparatus consisted of a
triangular (side= 30cm, height=40cm) arena with a transparent acrylic floor with a bottom-
up camera view. The apparatus was encased in a light controlled and sound attenuated
chamber. One of the corners was termed the Waiting Corner and the other two were the
lick port corners. The latter had lick ports. The lick port was equipped with a double
infrared sensor to capture head entry and single licks separately. The lickometer IR
sensor was inside a 7mm wide slit, inside which the reward delivery spout was placed
(see **Fig. 7A** and **Supp. Video 2**). The lickometer slit was just wide enough to fit the
tongue and deep enough to evoke a clear tongue protrusion. Rewards consisted of 3μl
droplets delivered through a solenoid, which was calibrated daily. Auditory stimuli were
delivered through a tweeter that was placed outside the arena, inside the sound
attenuating chamber. The task was controlled by interfacing Bonsai, Python scripts and
Arduino chips.

The task involved a two-alternative forced choice. The animal was required to approach
the wait corner, wait for 0.8-1.3s after which an auditory stimulus was played. Entrance
to the waiting corner was controlled by establishing a region of interest on the video feed
through Bonsai. Auditory stimuli consisted of 0.5 frequency modulated sweeps with a start
frequency of 9.2kHz and an end frequency of 0, +/-0.2, +/-0.4, +/-0.6, +/-0.8 octaves.
Upwards sweeps indicated a water reward on the right lick port and downwards on the
left. The animal then had to walk to the appropriate port. Correct responses were
rewarded with a water drop and incorrect ones were punished with a 10-20s timeout
indicated by a dimming of the ambient light (see **Supp. Video 1**).

Training was conducted through successive approximations introducing a new rule after
the previous one was acquired. To reach steady performance, animals were trained 3-4
months, 6 days a week.

Recordings. Movable microwire bundles (16 microwires, 32 micrometers in diameter,
held inside a cannula, Innovative Neurophysiology, Durham, NC)] were stereotaxtically
implanted just above the entopeduncular nucleus (-0.8 AP, 1.7 ML, 3.9 DV). Post surgical
care included antibiotic, analgesic and antiinflammatory pharmacological treatment. After
5 days of recovery, animals were retrained for 1-2 weeks. Unitary activity was recorded
for 2-6 days at each dorsoventral electrode position and the session with the best
electrophysiological (signal to noise ratio (>2), stability across time) and behavioral
[performance, number of trials (>220)] quality was selected. Microwire electrodes were
advanced in 50 micrometer dorsoventral steps for 500 micrometers in total. After
experiment completion, animals were perfused with a 4% paraformaldehyde solution.
Brains were extracted, dehydrated with a 30% sucrose solution and sectioned in a
cryostat into 30micron thick slices. Slices were mounted and photographed using a light
microscope. Microwire tracks of the 16-microwire bundle were analyzed (**Fig. 2A-B**) and
only animals with tracks traversing the EPN were selected (6 out of 10). Finally, we

located the final position of microwire tips and inferred the dorsoventral recording position
of each of the recording sessions. Only units recorded within the EPN were included.

Electrophysiological recordings were acquired at 30KHz (Cereplex Direct, Black Rock
Microsystems, UT) high pass filtered at 750Hz to extract spiking activity. Units with signal
to noise ratio of at least 2 were sorted online through a hoop algorithm and further sorted
offline (Offline Sorter, Plexon). Relevant behavioral events and video synchronization
were stored as digital events.

Animal video position was extracted using DeepLabCut (Mathis et al., 2018). The several
kinematic variables like angular velocity, velocity and acceleration of displacement as well
as spatio-temporal the relative position of the animal or time to the different corners were
computed (**Fig. 1C**). We identified 4 moments per trial to align the spiking activity: 1)
return to the wait corner, defined as the start of the negative animal-corner velocity; 2)
arrival to wait corner, when animal-corner distance <0.5pix; 3) turn start, when angular
velocity >0.01 degrees/s; 4) evaluation, the moment of lick port infrared sensor crossing.
Spiking activity was convolved with a gaussian function (sigma= 25ms) and rescaled
through linear interpolation (Kobak et al., 2016) to the average segment duration across
all trials which allowed to construct peri-event time histograms (PETH), (**Fig. 2 ,3,5**).

For **Figs. 2F, 2G** and **2I** z-score was obtained for each unit across all conditions (8 stimuli,
correct and incorrect) and averaged according to the specified conditions. Statistical
testing performed in **Fig. 2I** was done with the Wilcoxon test.

Linear regressions. Simple linear regressions were employed with some variables, for
example, reaction time in **Fig. 1E, H, J** and individual variables in the examples shown in
Fig. 4 A-C. Significance testing was done through permutation (n=1000).

Multiple regression models were used in **Fig. 4**. Data was obtained by generating a PETH
with 0.2s sliding windows (every 50ms) and obtaining the value for the different variables
tested (**Fig. 4-1**) through interpolation. The models were fitted through 10-fold cross
validation using an L1 (LASSO) regularization term. Cross-validated R2 values were
obtained from untrained data for each fold and reported as the 10-fold mean. Given that
regressors were not perfectly independent (**Fig. 4-1**), the regularization term penalized
the magnitude of the coefficients.

Partial correlation coefficient analyses (**Fig. 4 H-J**) were obtained by fitting a reduced
model lacking the variable to be studied and using the residuals to obtain the correlation
coefficient. Statistical significance was obtained by pseudo-random permutation (n=1000)
of values. For distance to goal, the reduced model included all the kinematic variables
(**Fig. 4F**) and no spatio-temporal ones (given their high level of correlation, **Fig. 4-1**). For
angular velocity, the reduced model included all kinematic and spatio-termporal except
the variable in question; for body speed the reduced model also excluded head speed.

For **Fig. 3-2**, we computed the loss of predictive power, delta R2, for each of the four
variables tested, similar to (Musall et al., 2019). We created reduced models in which the
datapoints for the specified variable were shuffled. The difference in explained variance
between the full and the reduced model equals the contribution (delta R2) of that variable
to the model. This provides a metric for unique information (not shared by other variables)
that each given variable contributes to the model.

Population level analyses

Population level analyses were employed several times throughout the study (**Figs.**
3,5,6,7), using standard Principal Component Analysis (PCA) and demixed Principal
Component Analysis (dPCA) procedures (Kobak et al., 2016).

Matrix construction varied according to the period and conditions used. For all population
level analyses individual units recorded from all sessions and all animals were pooled to
construct pseudo-simultaneous population response of data mostly recorded separately.
For **Fig. 3** condition averages were obtained for Go and evaluation periods, 0.3s before
turn start and 2.5 after lick port arrival. The dataset consisted of n=118 units recorded for
16 conditions (8 stimuli, with correct and incorrect responses) for which PETH were
obtained as described before. We included units with at least two trials per condition. Note
that responses can be classified by the direction of turn (Side), where for each stimulus
correct and incorrect responses include turns to the right and left. Further, for left and
right turns, there is stimulus gradation (0, +/-0.4, +/-0.6,+/-0.8), corresponding to the
difficulty of the task. Thus, the 16 conditions per unit were used to construct a matrix of
NxSxDxCxT, where N is the total of units (n=118), S is side (2 conditions), D is difficulty
(4 conditions), C is correct and incorrect responses (2 conditions), and T is time.

This matrix was used to calculate instantaneous variance (**Fig. 3A**) for the different
conditions with the following formulas:

197
$$VarSide_{INS}(t) = \frac{1}{N} \sum_{i=i}^{N} \sum_{S=1}^{2} (r^i(t,s) - r^i(t))^2$$

198
$$VarCorrect_{INS}(t) = \frac{1}{N} \sum_{i=i}^{N} \sum_{c=1}^{2} (r^i(t,c) - r^i(t))^2$$

199
$$VarDifficulty_{INS}(t) = \frac{1}{N} \sum_{i=i}^{N} \sum_{d=1}^{4} (r^i(t,d) - r^i(t))^2$$

200 To compute the population cosine similarity presented in **Fig. 3H-I** we used the
201 aforementioned matrix, mean-subtracted it to remove condition-independent (temporal)
202 dynamics. Then the following formula to all the population vector pairs at different
203 timepoints:

204
$$CS = \frac{r^i_{t1} \cdot r^i_{t2}}{\|r^i_{t1}\| \|r^i_{t2}\|}$$

For PCA analysis, a covariance matrix was constructed by obtaining the pairwise
covariance across all conditions. Eigenvalues and eigenvectors were then computed and
sorted by variance. For **Fig. 5-1** and **Fig. 7K** the first three components were used to
project the population responses. For **Fig. 6I**, the fourth PC was used. Projections were
obtained by computing the dot product with the eigenvector.

To assess the dimensionality of the population activity we compared the eigenvalues
obtained from the data with those obtained with per-unit time-shuffled matrices. This
destroys the time dependent correlations and thus generates PCs due to chance. We
performed this 1000 times and considered first PCs that explained more variance than
the time-shuffled ones with a $p < 0.01$ as statistically significant.

The details and mathematical procedure for dPCA analysis have been outlined elsewhere
(Kobak et al., 2016). Briefly, the method decomposes neural activity into the different
chosen task variables to produce marginalized covariance matrices. The supervised part
of the algorithm consists of choosing the variables to be analyzed. The unsupervised part
of the algorithm uses a similar analysis to that of PCA on the marginalized covariance
matrices. Our analyses were largely based on an available Python implementation
(https://github.com/machenslab/dPCA). In **Fig. 4** we marginalized the population activity
on time (condition-independent dynamics), Side, Difficulty and Correct vs incorrect.

The dPCA analysis presented in **Fig. 5** was performed similarly to that of **Fig. 3** but using
different time periods and conditions. For each session, model trajectories for Go, left and
right, were constructed by averaging the body speed and angular velocity. Then, we
computed the correlation between the model trajectory and individual trial trajectories,
both from the Return and Go periods. We selected those with a correlation coefficient of
≥0.75. Finally, we obtained PETH for the selected, highly similar trajectories.

For each unit, we obtained four conditions: Return left and right, and Go left and right.
With these we constructed a matrix NxCxS, where N is units (n=118), C is context (2
conditions, Return vs Go) and S is side (2 conditions, left and right). Thus, the
marginalizations were performed on the basis of these two variables and time (condition-
independent dynamics), mainly to extract the contextual (Return vs Go) population
dynamics.

Spike - cyclic behavior phase analysis. We analyzed how spiking was coupled to two
cyclic behaviors, gait and licking. For gait, we computed the paw position by projecting
the position onto the longitudinal axis of the mouse. We found individual gait cycles by
finding the peaks of this signal (**Fig. 6A**). We used movement periods that had at least 4
cycles/paces. We applied a band-passed filter (3-6Hz) to the signal and extracted the
instantaneous phase through a Hilbert transform. We then computed the phase at each
spike timestamp in a 400ms window around the peak (which includes around 2 gait
cycles) and generated a spike- gait-phase histogram. This was done with the hindpaw
contralateral to recorded hemisphere since hindpaws display a higher amplitude than
forepaws. The average vector was computed and tested for directional bias through the

Rayleigh test for each of the eight strides analyzed (**Fig. 6E**). Mean phase and magnitude
were computed by averaging the individual per-unit x and y-axis components.

For licking behavior, we computed the lick probability around an individual lick and applied
a 4-10Hz band-pass filter. This signal was then used to compute the spike - lick-phase in
a 400ms window (around two licks) like that performed with gait.

Support Vector Classifiers (SVCs). To understand whether the population could store
information for several variables simultaneously we trained SVCs using the
implementation in scikit (Pedregosa et al., 2011). For gait cycle, the same matrix used for
population dimensionality reduction through PCA was used. Each time vector, containing
n=118 unit averages was labeled according to the variables analyzed: context, step
number, paw position and paw velocity. Note that continuous signals (position and
velocity) were downsampled to 5 states to allow for 10-fold cross-validation. To assess
for statistical significance, we repeated the procedure on data with shuffled labels
(n=1000) and assessed the probability of a higher accuracy of the original model. Given
that we could not decode paw velocity from the original matrix, we computed the PC4
weighted matrix by using the eigenvector derived from PCA (**Fig. 6F**).

For licking we employed the same matrix used for PCA (part of which is shown in **Fig.
7J**) and trained models for lick phase, left-right licks, lick number, and correct vs incorrect
licks. To assess significance, a similar permutation procedure was performed.

Results

Given that our main interest lies in how the EPN neuronal activity encodes kinematic and
reward related variables we set up a task and recorded activity in freely moving mice. We
first describe the population dynamics that underlie reward, followed by asking how
movement is encoded by the units, contrasting both periods of movement in the designed
task (Return and Go trajectories). Next, we focus on extracting the population dynamics
that make Return and Go trajectories different. Finally, we describe how EPN units are
modulated by individual movements by analyzing gait and licking behavior.

**Mice can perform psychophysical responses on a self-paced two-alternative forced
choice task based on frequency modulated sweeps**

The main goal of this study was to establish the relationships between kinematic variables
of mice, particularly during locomotion, and the activity of the entopeduncular nucleus.
However, given that previous studies have ascertained that reward is an important
variable coded in this nucleus, we sought to evaluate how reward might be related to
kinematic coding.

To this end we devised a task in which animals were required to displace towards different
corners of a triangular arena (side=35cm, **Fig. 1A-B**). In a trial, animals were required to
move to the Waiting Corner and wait for ~1s (0.8-1.5s). After waiting, an auditory stimulus
would play, which would indicate in which other corner a water reward would be available
(a two-alternative forced choice). If the animal arrived at the appropriate lick port, it was

termed a correct response, and a water droplet (3μl) was then released. Incorrect
responses were signaled by a 10-20s timeout during which ambient lights were dimmed.
Animals would then start another trial (by moving to the Waiting Corner) in a self-paced
manner. By performing tracking [DeepLabCut, (Mathis et al., 2018)] of video recordings
from a bottom-up view, we could measure several variables for each trial (distance to
different corners, angular velocity, **Fig. 1C**) which allowed to establish different events of
the task (**Fig. 1B**). Particularly, we could establish two distinct periods of
movement/displacement: Return trajectories (to reach the wait corner) and Go trajectories
(to reach a reward corner).

Auditory stimuli consisted of 0.5s frequency modulated sweeps; upward and downward
frequency modulation indicated a right and left choice, respectively (**Fig. 1A**, right). By
varying the rate at which the frequency was modulated we attempted to modulate the
certainty with which animals decided. This was evidenced by a psychometric curve fitted
to the response probability as a function of stimulus deltaFreq (**Fig. 1D**). Note that the
central stimulus was a pure tone that had no information of side. This behavior was
reproducible for different sessions before and after electrode implant (**Fig. 1D**, middle)
and in several animals included in this study (n=6; **Fig. 1D**, right).

We then quantified how the different stimuli influenced response time and kinematic
parameters. Note that stimulus pairs with higher deltaFreq (+/-0.8) were easier than those
with less (+/-0.4) or no deltaFreq. The +/-0.2 stimulus pair was dropped from some
sessions to increase trials per condition and will not be further analyzed. We quantified
Response time as the time from stimulus onset to the onset of turning (**Fig. 1B-C**), which
we defined as the moment where animals had committed to a motor action (either left or
right turn). We found a relationship between response time and the stimulus difficulty
(correlation coefficient r=0.89, p<0.01 permutation test; **Fig. 1E**). To ascertain whether
the evolving deltaFreq of the stimuli had an impact on performance, we calculated the
probability of a correct response as a function of response time by binning trials according
to response time (bin=75ms) and by stimulus difficulty (deltaFreq). For the central
stimulus, no amount of response time increases the probability of correct response since
the stimulus contains no information of the correct side. For stimuli pairs with varying
difficulty, there is a graded increase in the probability of a correct response (**Fig. 1F**).

Next, we sought to ask whether the difficulty of the stimuli and thus uncertainty in the
decision could modulate the kinematic response of the animals. We hypothesized that
responses to easier stimuli would result in faster movement. We calculated the
instantaneous angular velocity across the 150º of turning that animals had to perform
(**Fig. 1G**), showing indeed a subtle gradation according to the difficulty of the stimuli. We
found a significant correlation between average angular speed and the deltaFreq of
stimuli for correct trials but not so for incorrect trials (**Fig. 1H**, black and gray lines,
respectively). We further calculated the instantaneous speed across the trajectory that
animals had to complete (Go trajectory, **Fig. 1I**). Animals performed easy trials faster than
more difficult ones, but no significant correlation between stimulus difficulty and speed
was found (**Fig. 1J**).

From the performance of animals in this task we conclude that animals can perform a
psychophysical task based on frequency modulated sweeps, with easier stimuli resulting
in greater hits than more difficult stimuli. The difficulty of the stimuli had an impact on the

reaction time, and increasing the sampling time for stimuli with information of correct
response increased the probability of performing a correct choice. Furthermore, difficulty
had a moderate impact on the way animals turned, but not on the whole period of
locomotion.

EPN recordings and examples

After a consistent performance across at least 2 weeks (**Fig. 1D**) animals were implanted
with a movable microwire bundle to record the activity of EPN neurons. The microwires
were advanced in 50micrometer steps for 10 steps in total. We inferred the recording
position via postmortem histological analysis of the individual microwire tracks and
included for further analysis only the ones that were recorded inside of EPN (**Fig. 2B**).
Previous reports indicated that the two main cell types in the EPN [characterized by
parvalbumin+ and somatostatin+ markers, (Stephenson-Jones et al., 2016; Wallace et
al., 2017)] exhibit differences in the coefficient of variation and the duration of the action
potentials. To this end, we computed these characteristics for the units recorded in this
study (**Fig. 2C**). However, we could not separate the population into two distinct clusters.

Fig. 2D-E show two examples of the activity recorded aligned to the task relevant events
identified in **Fig. 1B**: return, wait, turn (0º, 75º, 150º), outcome. Note that we sought to
isolate the turn during the Go trajectory by identifying the start (0º), middle (75º), and end
(150º) of the turn; the top row shows the raster plots color coded by the stimulus for all
correct trials. Despite the performance of animals being highly stereotyped, there are
slight differences due to it being freely moving. Thus, in the three bottom rows we show
the rescaled activity presented in the raster plots for the correct trials, and additionally for
the incorrect trials and false alarm trials, respectively. The false alarm trials were trials in
which animals did not wait for the auditory stimulus to be presented and performed the
entire motor sequence identical to real trials (useful since they are gated by an internal
signal without external stimuli with identical motor output).

The unit in **Fig. 2D** shows a decreasing firing rate as time/distance from the goal of each
of the Return (reach the wait corner) and Go trajectories (reach the reward corner).
Further, this unit shows differential activity during left and right turns that follows the
direction of turning in correct, incorrect, and false alarm trials. On the other hand, the unit
in **Fig. 2E** exhibited a marked difference in the time/distance varying activity during Return
and Go trajectories (with no change and ramping activity, respectively). Further, this
second unit exhibits a stark difference in the evaluation period between rewarded and
unrewarded trials.

Heatmaps in **Fig. 2F** show responses for all recorded units (118 units from 6 animals)
averaged for all correct right (**Fig. 2F**, upper) and left (**Fig. 2F,** lower) trials, sorted by
peak activity in rightward trials. Even if neurons are organized employing only the
rightward trials, the population activity displays akin responses. **Fig. 2G** upper panel
shows averaged z-score EPN activity sorted by turn side and outcome (correct, incorrect,
and false alarms), with average kinematic measurements of speed and angular velocity
presented in **Fig. 2H.** These population averages show that there is population coding
associated to turn and reward. Sorting hit trials based on stimulus identity (**Fig. 2G**, lower)
preserves side representation; however, differences across stimuli are remarkably low.

We sought to assess how average population activity could be representing movement,
turning and reward. Thus, we obtained the z-score population average during both
periods of movement (Return and Go) and the wait period (**Fig. 2I**). As well, we obtained
the average activity during turns ipsilateral and contralateral to the recording site (right
hemisphere), and for correct and incorrect trials during the evaluation period. Out of all
conditions, reward (evaluation of correct trials) exhibits the highest population activity
($p<0.01$, Wilcoxon test); correct trials also exhibit a higher activity compared to
unrewarded trials despite similar speed (**Fig. 2H**). Ipsilateral turns exhibit a higher activity
compared to contralateral turns and the wait period. Finally, despite a tendency to lower
activity during both movement periods (**Fig. 2G-H**), the data does not show a statistically
significant difference on average activity during the wait period compared to the Return
and Go periods.

Population analysis of the EPN reveals its coding of reward, left and right but not
difficulty

As a first objective we sought to analyze coding of stimulus difficulty and reward, given
previous reports that GPi/EPN neurons code for the value of expected positive and
negative outcomes (Stephenson-Jones et al., 2016) as well as the spatial position of the
expected positive and negative outcomes (Hong & Hikosaka, 2008). We began by trying
to understand how these variables were represented in the population activity. For
population level analyses throughout the study, we pooled recorded units from all animals
to construct a pseudo-simultaneous population. We calculated the instantaneous
variance associated with these two variables, as well as to left and right trials (**Fig. 3A**).

In agreement with **Fig. 2G**, there is high variance associated with correct and incorrect
trials during the evaluation period of the task. Further, left/right variance is increased
during the turning period. In concordance with **Fig. 2G**, the variance associated with
difficulty (stimuli) is remarkably low.

Next, to understand the population dynamics associated to these task parameters we
performed demixed Principal Component Analysis (Kobak et al., 2016). This allowed us
to generate a low-dimensional space that captures variance related to specific task
parameters: condition independent temporal dynamics, correct and incorrect trials,
stimulus difficulty, and side (left and right) of turning. We hypothesized that condition
independent temporal dynamics would include kinematic relationships while
marginalizations in correct/incorrect would include reward signals; the left/right
marginalization would show angular velocity related activity, and the difficulty
marginalization (that is, the segregation related to the deltaFreq of the auditory stimuli)
would show a gradation related to the action value.

After performing dPCA and sorting the extracted components by the explained total
variance (ETV), we find that the first three are condition independent components
(temporal). Further, only five PCs are statistically significant (see Methods). The most
prominent dynamic, temporal dPC1 which explains 19.2% of total variance, separates the
outcome period from the wait and go periods akin to a step function. The right panel in

Fig. 3C shows the weights of individual units onto this dynamic. We show individual example units in **Fig.3-1** which correspond to the letter markings in the weight panels (**Figs. 3C-G**, right panels). The second temporal dPC (16.1% ETV) exhibits ramping activity during the Go moving period (**Fig. 3D**), which could be encoding time or the distance to goal. The third temporal dPC projection exhibits a triphasic dynamic during the moving period, potentially correlating with the speed of the animal's trajectory (**Fig. 3E**).

The fourth and fifth dPCs correspond to correct/incorrect and side, respectively. As expected, the correct/incorrect dPC separates correct from incorrect trials during the evaluation period (**Fig. 3F**). Further, the side dPC is maximally different during the turning period (**Fig. 3G**). Note that the weights of these axes are positive and negative centered around zero. This means that units encoding reward can code a rewarded trial as an increase in activity or as a decrease in activity and vice versa (example units are shown in **Fig. 3-1g** and **h**). Similarly, leftwards turns can be encoded as a decrease in activity or an increase of activity (**Fig. 3-1 a** and **b**).

Interestingly, the marginalization of difficulty (stimuli) did not yield a significant dPC. This, coupled with the comparatively low difficulty-associated variance (**Fig. 3A**) suggests that difficulty coding is not well represented in this dataset. Still, by comparing correct and incorrect trials (with stimulus) vs false alarm trials (without stimulus), some units show stimulus-associated activity (**Fig. 3-2AB**). We looked to further assess whether stimulus related activity could represent difficulty. We asked whether activity around stimulus and turning could be representing the presence of stimulus, the difficulty of the stimulus, what the animal interpreted from the stimulus (correctly or incorrectly turning to either right or left), or simply the angular velocity (**Fig. 3-2C**). Given their overlap, we calculated the delta R2 of each of these possibilities, which measures how much the goodness of fit improves when considering each variable versus only the others (see Methods). This analysis shows that including information about the difficulty or the interpretation are worse at explaining activity than the sole presence of stimuli (**Fig. 3-2D**). Further, the variable that best describes the activity in this period is angular velocity.

To assess how side and reward coding might correlate/covary with each other, we computed the population self-similarity across time (**Fig 3. H**). This self-similarity was computed from the demeaned responses, which eliminates condition-independent (temporal) coding. We found that the evaluation period is well separated from the Go period and the interaction between them is very low. We assessed the similarity of the population vector at the time points of greatest side and reward variance (t=0.2s and t=1.8s, respectively; see **Fig. 3A**) against the rest of the time bins (**Fig. 3I**). This shows that turning and reward coding are largely uncoupled.

The presented analyses in **Figs. 3** and **3-2** show that at the population level the EPN recorded activity in freely moving mice encodes reward and turning side, both of which are encoded positively and negatively. Minimal contribution of difficulty was detected.

Contextual motor coding across different periods

As a second objective, we analyzed the relationships between the activity of the recorded units and kinematic variables. Since one of the most prominent population dynamics describes a linear relationship between time/space and population rate (**Fig. 3D**), we sought to study spatio-temporal variables as well. By design, the task has two distinct epochs of displacement: Return trajectories (to the waiting corner) and Go trajectories (to either of the reward corners). We hypothesized that purely 'motor coding' would be invariant to these two contexts.

Fig. 4A-C shows three example units. Each unit has two raster plots corresponding to Return and Go trajectories, aligned to movement start and sorted by movement duration. We grouped and averaged their activity into six bins for visualization and presented the average firing rate and average speed. All units presented have significant correlations with at least three of the variables presented, which suggests that a multiple regression model is better suited for the analysis.

For the construction of the multiple regression models, we divided variables into kinematic (measured from the body of the animal) and spatio-temporal (measured in relation with the arena) and reward related (**Fig. 4F** and **Fig. 4-1**). Given the possibility of collinearity between measured variables (**Fig. 4-1**), we employed LASSO regression which introduces a regularization term that shrinks the coefficients and penalizes colinear variables, acting as a feature selection (see Methods). To assess how well the model explained the neural activity we calculated a tenfold cross-validated R2 for each unit, which ensures that no variance is predicted by chance.

The group of kinematic variables is worse at explaining single unit activity when compared to spatio-temporal and reward related variables (**Fig. 4D**). However, a model incorporating all three groups of variables outperforms any specific category.

An important goal of this study was to compare the encoding kinematic and spatio-temporal variables between the Return versus the Go trajectories (two periods of similar movement but different sub-goals).

By fitting models separately for the Return and Go trajectories, we found that the average cvR2 for models fitted with Go trajectory data is higher for kinematic, and spatio-temporal variables, as well as for a mixed model (p<0.001, Wilcoxon test; orange and green bars in **Fig 4E**). This implies both kinematic and spatio-temporal variables are best represented during the trajectories that animals performed towards the reward ports (Go trajectories). This is probably partly due to units that are more responsive during the Go trajectory (**Fig. 2E**).

Next, we sought to investigate which variables were most relevant for the models' success. To this end we constructed single variable models for each unit and plot the mean cvR2 for each of the measured variables (**Fig. 4F**). This was also done separately for return (orange) and go (green) trajectories, as well as for the entire duration model (gray). Amongst the kinematic variables the best variable at explaining the neural data variance was the body speed, followed by the head speed and the angular velocity (**Fig. 4F**). We found that relationships with tested variables are not generalizable between both contexts. Model performance is reduced drastically (p<0.001, Wilcoxon test) when switching the test data to that of the other context (**Fig. 4G**). This is unlikely to be due to over-fitting since models were trained through cross-validation. The implication of this

finding is that understanding the relationships between kinematic and spatiotemporal
variables in one context is not enough to predict how the population will respond in
another context.

To further examine why these models were not generalizable between contexts, we
calculated the partial correlation coefficient for both contexts for three of the variables:
distance to goal, speed, and angular velocity (**Fig. 4H-J**). We fitted the full model minus
the variable in question to the data of each of the two contexts and calculated the
correlation coefficient with the residuals (see Methods). We plotted the return trajectory
correlation coefficient (x-axis) against the go trajectory (y-axis). In a hypothetical scenario
where EPN units reflected purely motor coding, all data points would lie within the identity
line (dashed line). However, this analysis reveals that some units display a significant
partial correlation with the variable during either Return trajectories (orange dots) or the
Go trajectories (green dots). Even though there are units that display a significant
correlation in both trajectories (brown dots), these units do not solely lie within the I and
III quadrants of the cartesian plot, which means that some units have statistically
significant correlations with opposite signs during distinct trajectories (**Fig. 4H-J**, pie
charts). Indeed, for angular velocity and body speed only about a third (31%) have
significant correlations in both trajectories, but only about a fifth of all units have significant
correlations with congruent signs (**Fig. 4H-J**, pie charts). Overall, these analyses paint a
complex picture, where the subset of units that encode each of the variables studied in
one context is not the same as those that encode the same variable in the other context.
Further, the differing magnitude of the correlation explains why models trained on one
trajectory are not generalizable to the other trajectory. Some extreme cases of this
instability of coding are those units that have a sign switch between one context and the
other (black outer pie in **Fig. 4H-J**).

Given that we found that one unit might significantly encode multiple variables, we wished
to investigate how the significant partial correlations for these three variables might be
distributed amongst the recorded units. In **Fig. 4K** we plot Venn diagrams of the
percentage of units that significantly code each of these three variables during the Return
(top) and the Go (bottom) trajectories and their overlap. We find that there is a similar
landscape in both periods of movement wherein most units have significant correlations
with more than one variable. However, as examined before, the subset of units in each
of the categories is not the same from one context to the other.

No segregation of units coding for kinematic, spatio-temporal variables and reward

Next, we asked whether the recorded units could be grouped based upon what variables
they code. That is, we sought to understand whether groups of units might encode some
variables more strongly than others. Previous reports suggest that EPN units that project
to the lateral habenula encode reward as a decrease in firing rate. Thus, we wished to
ask whether reward encoding units can code kinematic and spatio-temporal variables as
well.

To this end, we first segregated units upon their reward coding properties: reward positive
(which increased activity with reward) and reward negative units (which decreased activity
with reward). We performed auROC on the 250ms after head entry comparing rewarded

trials and incorrect trails (p<0.001, permutation test). Mean activity of reward insensitive,
positive and negative units is shown in **Fig. 4L**. Next, we performed a dimensionality
reduction on the coefficients of the model that best explained both contexts (kinematic +
spatio-temporal model on pooled data) using UMAP (McInnes et al., 2018). We observe
a continuum rather than discrete clusters (**Fig. 4L**). Note that individual units are color
coded according to their responsivity to reward. We did not find a clear clustering either.
This implies that, overall, coding for kinematic and spatio-temporal variables is uniformly
distributed and the coding of these variables does not segregate the coding for reward.
Interestingly, reward sensitive units encoded kinematic and spatio-temporal parameters
similarly to reward insensitive units (**Fig. 4M**). Reward positive units are better at encoding
these variables than are reward negative units. However, neither type of reward sensitive
units had a significantly different cvR2 than that of reward insensitive units.

To ask whether action value might be influencing the way units encoded variables, we
segregated trials by their outcome into correct and incorrect. Additionally, we included
false alarm trials, which animals performed in the absence of any auditory stimulus. We
fitted the models independently for each subset of data. We found that false alarm trials
have a statistically significantly lower cvR2 only for the spatio-temporal model, not so for
the kinematic one. Thus, time/space varying signals are attenuated during False Alarm
trials (the example unit on **Fig. 2E** exhibits this behavior).

What makes Return and Go trajectories different at a population level?

To address this question, we next sought to understand what population level dynamics
are shared between Return and Go trajectories, and what underlying dynamics might be
explaining context differences. To this end we found highly similar Return and Go
trajectories from a motor perspective (see Methods) based on body speed and angular
velocity (**Fig. 5A**). Intriguingly, by projecting the population activity (n=118 units) onto the
first 3 PCs (**Fig. 5B**), Return and Go trajectories occupy different subspaces in this graph
(**Fig. 5-1A-B**). Indeed, while the kinematic similarity is high between Return and Go
trajectories (**Fig. 5C**, upper), population similarity exhibits two groups based on the
context (**Fig. 5C**, lower).

To find a more interpretable dynamic that may account for the differences between the
two contexts we employed dPCA, marginalizing by context, left-right, and temporal
components. We hypothesized that temporal components shared by both contexts might
exhibit kinematic variables and that left-right marginalization would encompass angular
velocity. After performing the dPCA procedure (see Methods), we sorted the components
by variance (**Fig. 5-1C**). Note that after marginalization upon the desired variables, the
extraction of the dynamics is unsupervised. This is to say that extracted dynamics do not
have to necessarily produce traces related to individual kinematic parameters, and they
are unlikely to do so given the multiplexing of encoded variables (**Fig. 4K**). However, we
sought to interpret the resulting traces based on measurable variables. Thus, we
compared the resulting trace with the variables measured on the same time periods by
calculating the correlation coefficient and testing significance through permutation testing
(p<0.001).

The contextual marginalization produces two significant dPCs that explain the differences
between Return and Go trajectories. Indeed, the contextual dPC1 explains 15% of the
total variance (**Fig. 5D**), and Return vs Go traces exhibit oppositely evolving temporal
dynamics. The spatial variable of distance from the waiting corner best explains these
contextual traces, while no kinematic variable correlates with these traces. Interestingly,
context dPC2 (**Fig. 5D**) exhibits a high similarity to the speed-correlated temporal dPC2
(**Fig. 5-1E**). However, context dPC2 shows oppositely evolving dynamics symmetric over
time between Return and Go trajectories. This suggests that there is a simultaneous
representation of speed and a spatially referenced velocity (**Fig. 5G-I**). Indeed, context
dPC2 correlates well with the velocity from the waiting corner.

Condition-independent (temporal) dPCs summarize population dynamics shared by both
Return and Go contexts (**Fig. 5-1**). Temporal dPC1 has a high correlation with distance
to goal (**Fig. 5-2D**) while temporal dPC2 (**Fig. 5-1E**) is well correlated to the speed. Note
that these traces do not correlate with one specific variable, which is to be expected given
the previously established high level of multiplexing (**Fig. 4H-K**). The marginalization for
side (side dPC1, **Fig. 5-1G**) produces traces that have significant correlation with angular
velocity, both for the Return and Go trajectories. Note that Go trajectories occurred under
the presence of an auditory stimulus, and not the Return trajectories; however, traces
corresponding to left and right turns (irrespective of goal) show good overlap. Thus, this
further confirms that the associated differences between left and right turning are well
explained by angular velocity. We conclude that context dependent dynamics explain
variables whose measurement depends on specific places of the arena. Thus, spatially
biased variables have an important representation by the EPN population activity.

EPN units are modulated by gait and licking

Our findings reveal a robust representation of whole-body speed (**Fig. 4**) as well as
spatially biased variables (**Fig. 4-5**). However, prior studies have indicated a somatotopic
representation in the GPi across various species [humans, non-human primates, cats
(Baker et al., 2010; Larsen & McBride, 1979; Nambu, 2011)]. Therefore, we examined the
EPN's modulation by individual movements of the paw and tongue, using the cyclic nature
of gait and licking to examine the potential presence of muscle-level activity since they
exhibit a clear flexor-extensor muscle relationship.

Given that animals were recorded from a bottom-up view, we were able to track individual
paws of the animal (**Fig. 6A**). We searched for Return and Go trajectories with at least
four individual strides (**Fig. 6A**, triangles) and used them to align unit activity. **Fig. 6B**
shows two example units simultaneously recorded from different channels. Note how Unit
1 has a very robust stride locked activity, while Unit 2 shows a smaller stride-based
modulation and instead becomes immersed in an evolving temporal dynamic. The
average paw position for the session (**Fig. 6B**, bottom) is well preserved, and that there
is a high level of correlation or anticorrelation, which justifies our using only one paw to
study. This procedure was performed for all units studied and is presented in **Fig. 6C**,
sorted by peak activity during the step 4 in Return trajectories. This sorting is only slightly
preserved in the rest of the steps, highlighting the dynamic nature of this gait modulation.

Since locomotion necessitates a coordinated contraction of flexor and extensors during
each stride, we hypothesized that if individual muscle activity was represented in the GPi,
we would find stride-cycle locked activity in symmetric opposing phases (flexors vs
extensors). We calculated the spike- contralateral hindlimb phase locking for each unit at
each step studied. In **Fig. 6D** we plot the average vectors of individual units with
significant phase locking (p<0.001, Rayleigh directionality test) for each of the eight steps
studied. Note how for each step units tend to point to one phase and not to the antiphase.
Further, there is a shift in the mean phase of the paw (**Fig. 6D-E** middle), for Return and
Go trajectories. Surprisingly, as the step number increases (which corresponds to steps
closer to the goal of the trajectory), a greater fraction of units has a significantly stride
phase-locked activity (**Fig. 6E,** upper panel). The magnitude of the average vector also
increases with step number (**Fig. 6E,** lower panel), suggesting an increase in phase-
locking. These results are not supportive of this activity being muscle related, since most
gait modulated units are biased toward a single phase of gait, with no antiphase
representation (**Fig. 6C**). Further, gait modulation increases with steps closer to the goal,
and phase precession would not occur if units were strictly related to muscle activity.

Given the widespread existence of this phase-locked activity, we hypothesized the
presence of a covariance plane that would encompass it. Thus, by performing PCA on
the population activity (**Fig. 6C,6F**) we could understand what variable might be
represented at a population level. We found that the population projection onto the first 3
PCs correlated best with spatio-temporal variables (**Fig. 6G,** upper panel) while the fourth
PC had a good correlation with paw velocity (**Fig. 6g,** lower). Indeed, there is a clear
overlap of PC4 projection with paw velocity (**Fig. 6I**). Finally, we sought to compare the
robustness of the representation of the different variables on the population activity. To
this end we trained linear support vector classifiers to discern between Return and Go
trajectories (context), step number and paw position and acceleration (**Fig. 6J**). Paw
velocity could only be decoded above chance on the PC4 weighted matrix, not on the
original data. These results support the existence of a relatively weak representation of
the paw velocity at a population level. Paw related PC4 captures only 5% (**Fig. 6F**) of the
variance of this matrix, which includes less time compared to other population analyses
(**Figs. 3, 5**), does not segregate by turning direction, and does not include a reward
period.

A final test of kinematic representation was performed by analyzing the relationship
between EPN activity and licking behavior. Individual licks could be recorded by a custom
made lickometer (**Fig. 7A**, see methods and **Supp. Video 2**). We found correlation
between licking behavior and EPN activity at two different timescales: as a whole lick bout
instance and at the single lick behavior. **Fig. 7B** and **7C** are raster plots of licks and a unit
activity recorded simultaneously, for left and right licking ports. Because of the apparent
similarity between lick and firing rate traces, we sought to assess whether units might be
encoding licking rate during the licking bout. To this end we computed the rescaled activity
to the average duration of licking bouts. **Fig. 7D,** upper panel shows the rescaled activity
of three simultaneously recorded units and the corresponding licking rate. Despite varying
dynamics exhibited by these three units, all three have a high degree of correlation with
the lick rate. **Fig.7D,** lower panel shows the correlation coefficient between the
instantaneous firing rate and the instantaneous lick rate within a lick bout for all units.
Indeed, most units (91%) exhibit a positive or negative significant correlation with lick rate

(p<0.001, permutation test). We then obtained PETHs of activity related to single licks in
the left and right lick ports (red and blue, respectively), as shown in **Fig. 7E** for the same
unit as in **Fig. 7C**. By extracting the lick phase from the lick probability (sensor crossings,
Fig. 7E, upper), we could generate a spike – lick-phase histogram and calculate the
resulting vector (**Fig. 7E**, right). When performing this for all units, it was clear that they
fire more strongly in a specific phase of the licking behavior. This is also shown in the
heatmap of perievent activity sorted by peak activity (**Fig. 7G**). Thus, we show that nearly
all units (99%) exhibit correlation with lick rate, single licks, or both (**Fig. 7H**).

What is the significance of such a robust representation of licking by nearly all recorded
units? We hypothesized that different aspects of a single lick might be represented. To
this end we obtained PETH for 16 different licks per unit: the first and last licks per bout,
as well as 5 licks spread out within the bout, right and left. We noticed that occasionally
animals would execute one lick when animals performed incorrect or false alarm trials,
which we termed an incorrect lick (right and left). This allowed us to analyze a similar
motor output (a single lick) performed under different contextual situations. An example
unit (**Fig. 7I**) shows that despite similarities between all these conditions, firing rate traces
exhibit clear differences. **Fig. 7J** shows a heat map for 7 right licks during a bout sorted
by the lick with highest activity. This highlights that while there is a modulation at the
single lick timescale, a whole-bout modulation is also present. After PCA dimensionality
reduction (**Fig. 7J,** lower), we projected the population responses on the three main
principal components (**Fig. 7K**). The population responses for the 16 conditions maintain
a circular (cyclic) response. However, they occupy distinct spaces in this state space
which suggests that the EPN population can distinguish amongst them. To assess this,
we trained support vector classifiers on population vectors for lick phase, left vs right licks,
lick number, and correct vs incorrect licks. All these variables could be accurately
decoded above chance. Together, these results show that most units in the EPN are
modulated by licking behavior at different timescales. Further, information about licking
kinematics is simultaneously represented with other contextual variables.

Discussion

This study highlights the dynamic nature of kinematic representation within the EPN. It is
particularly striking that spatio-temporal dynamics are better at explaining EPN activity
than are kinematic ones. Further, relationships between these variables and EPN activity
fluctuate across two largely similar periods of movement (Return vs Go). What is more,
units multiplex different functions like reward, whole-body movements, and individual
movements (paw and licks) with contextual variables. Interestingly, we were unable to
find a strong correlate of difficulty.

The principal goal of this study was to investigate how the activity of entopeduncular
nucleus is related to kinematic variables, reward and difficulty/value, particularly during
locomotion. Two previous findings had a particular influence over the study design: 1) the
existence of reward and value coding with a spatial bias, and 2) that kinematic coding
was unstable and varies depending on cognitive states. To this end, we designed a task
that involved decision confidence, reward, and two periods of displacement.

Prominent representation of evaluation period, reward, but no difficulty in the EPN

We found that there is a robust representation of evaluation period and reward. When analyzing the population dynamics (from pooled units from all animals to construct a pseudo-simultaneous population, which assumes homogeneity across subjects) underlying the Go period, we found that a dynamic that separates the evaluation period from the rest of the task captures the most variance (**Fig. 3C**). Further, a statistically significant reward dPC was found (**Fig. 3F**). Importantly, reward coding is also represented within this nucleus by reward positive and negative neurons. These findings support previous studies (Hong & Hikosaka, 2008; Stephenson-Jones et al., 2016).

Stephenson-Jones et al. found that a subset of EPN neurons projecting to the habenula encode value of expected outcomes in a classical conditioning task. We hypothesized that by varying the difficulty of stimuli (and thus increasing uncertainty during difficult choices) we could be able to modulate the perceived value of the chosen action. However, the explained variance by this hypothesis is very low (**Fig. 3**). The stimuli were able to impact the animals' response time and angular velocity (**Fig. 1E-H**) which suggests some level of metacognition of the uncertainty in their decision. However, speed was not well correlated with difficulty (**Fig. 1I-J**), which could imply that this metacognition diluted over time. Further, value coding was documented in neurons recorded during a head-fixed classical conditioning task (Stephenson-Jones et al., 2016), where the stimulus-outcome contingency might be more explicit or the coding of neurons less mixed (due to forced immobility). It is still possible that by modifying reward contingencies such as droplet size value coding could be evidenced.

In this study, we found that rewarding outcomes can be represented by EPN units through either an increase or a decrease in firing rate (Fig. 3F, 3-1g-h, 4L). While Stephenson-Jones et al., 2016 found that mice lateral habenula (LHb)-projecting neurons within the EPN of mice primarily encoded rewarding outcomes by a decrease in firing rate, Hong and Hikosaka, 2008 observed that in primates, LHb-projecting units could encode reward through either a decrease or an increase in firing rate. Thus, our results align more closely with the latter study, which also employed an operant conditioning task.

EPN activity can represent movement but with novel considerations

In this study we found that Go trajectories have a more robust representation of kinematic and spatio-temporal variables when compared to Return trajectories (**Fig. 4E**). Indeed, several studies have found an inconsistent relationship between the GPi activity and limb movements of the primate, where units were modulated by movements under certain cognitive conditions but not others. Some hypotheses have been that GPi coding depends on movement type (ramping vs. stepped; self-paced vs step tracking, (DeLong, 1971; Georgopoulos et al., 1983; Mink & Thach, 1991a) or between identical movements performed under different cognitive states (cued vs memory-dependent (Turner & Anderson, 2005)). Further, some studies agree that self-paced movements are more weekly represented (Mink & Thach, 1991b; Turner & Anderson, 2005). Interestingly, while most previous studies have focused on limb movements, here we find that whole body displacement kinematics are better at explaining EPN activity than individual limb

movements. Both Return and Go movements in this study are self-paced, and they are
mostly different in their relation to reward.

Representation of movements seems to be very dependent on what the movement is
performed for. Kinematic and spatiotemporal variables are not stably represented across
similar movements with different purposes (**Figs. 4-5**). Given the sequential nature of the
task, Return and Go trajectories are both required to eventually get a reward. While both
trajectories have a goal: Return → auditory stimulus, Go → water droplet, they differ in
that Go trajectories are performed to receive something rewarding. Indeed, we
hypothesize that movements that can lead to a reward are better represented by EPN
neurons. A similar conclusion was reached (Gdowski et al., 2001) in primates performing
wrist movements, where they found that wrist movements that led to reward modulated a
higher fraction of units than wrist movements that did not lead to a reward. The findings
in this study differ in that kinematic relationships were indeed found in the Return
trajectories, but they were attenuated.

An important finding of this study is that spatio-temporal variables are better at explaining
EPN activity than purely kinematic ones, for both Go and Return trajectories. Further,
these spatio-temporal variables are sensitive to action value (**Fig. 4N**, false alarm trials
have a lower R2 in the spatio-temporal model). Importantly, we did not find that kinematic
representation was affected by action value (**Fig. 4N**). Only false alarm trials had a weaker
relation with spatio-temporal variables than correct trials. It is possible that value
influences the spatio-temporal dynamics [akin to reward prediction error, (Dayan &
Balleine, 2002)]. A previous finding focusing on the decision properties of the GPi neurons
found that the main correlate that the GPi has, as compared to other brain regions, was
a temporal component which they called urgency (Thura & Cisek, 2017). This is similar to
our finding that spatio-temporal dynamics dominate EPN activity. They found that
decision was not as well represented in the GPi compared to other cortical regions. This
suggests that more than movement per se, the EPN might compute contextual
relationships present in the task.

There is a longstanding belief that reward related functions in the EPN are separate from
motor related functions, since anatomical studies have found two distinct projections to
habenula and motor thalamus (Hong & Hikosaka, 2008; Parent et al., 1999; Stephenson-
Jones et al., 2016; Wallace et al., 2017). In this study we did not identify recorded units by
their projections, however, we could evaluate their sensitivity to reward given the task
design. Surprisingly, we did not find differences in the ability of reward sensitive units to
encode kinematic and spatiotemporal variables compared to reward insensitive (**Fig. 7L-
M**). Given that we did not record from the entire EPN, it is still possible that another region
of the nucleus might exhibit more segregation.

We found that latent population dynamics that best explain differences between the
Return and Go trajectories can be best explained by spatially referenced variables (such
as distance and velocity from a specific place in the arena, **Fig. 5**). The similar but
contextually opposite dynamics probably reflect the units switching sign between contexts
that were found in **Fig. 4H-J** (outer pie, black). Interestingly, a previous study in primates
(Hong & Hikosaka, 2008) found that the direction of rewarded outcome greatly modulated
units in a one-direction rewarded task, and switched sign when the spatial location of the

rewarded target was alternated. They attributed the activity to the stimulus onset, and did
not study it in regards to saccade onset. However, the direction sensitive activity was
appropriately timed to coincide with saccade performance. It is thus possible that an
important function of EPN activity is to compute kinematic and spatio-temporal relations
between the subject and the reward location. In this study dynamics related to (spatial)
side of turning during the Go period and reward were found to be largely uncorrelated
(orthogonal to each other, **Fig. 3H-I**); however, both turns during the Go period
(performed to get to the reward port) have similar value. On the other hand, population
dynamics that explain the differences between Return and Go periods, which can be said
to have different action value, did exhibit sign switching (**Fig. 5D,G**). Thus, action value
(moving to or away from the rewarded location) could explain these opposing dynamics.
Indeed, reward prediction error models incorporate/rely on spatio-temporal variables
(Dayan & Balleine, 2002; Kim et al., 2020) .

Previous studies performed in humans, primates and cats (Baker et al., 2010; Larsen &
McBride, 1979; Nambu, 2011) have suggested the existence of a somatotopic organization
in the GPi. We sought to test the hypothesis that this organization could be representing
muscle contraction information. To this end, we analyzed two cyclic movements, gait and
licking, since they require a coordinated contraction of flexor and extensor muscles. The
gait coupling of EPN units has also been found in cats (Mullie et al., 2020). However, we
found that while units did exhibit a selective firing on a specific phase of both of these
cyclic movements, there was no anti-phase balance. This could imply that only flexor
muscles are represented, or, more likely, that the activity is not related to muscle
contraction. A further fact that is contrary to muscle level modulation of activity is that both
representations, gait an licking, largely overlap. That is, units that significantly modulate
to gait also do so for licking. Thus, we can conclude that in this mouse data, somatotopic
organization is not present, which is a similar conclusion reached by previous primate
studies (DeLong, 1971; Mitchell et al., 1987).

Finally, it was surprising to find such a robust representation of licking behavior in these
units. Practically all units correlate to whole bout licking rate and/or individual licks. This
implies that the same units that presented significant correlations with movement during
the displacement periods, could robustly represent licking behavior. Further, the
population could represent contextual parameters associated with licking (**Fig. 7K-L**).

What could be the purpose of having a representation of many types of movements and
variables by the same population? The data in this study suggests that action value might
be key to the level of movement representation by the EPN. Cued movements (Go,
correct trials) are better represented than uncued (False Alarm) ones (**Fig. 4N**), which in
turn are better represented than trajectories that cannot lead to reward (Return
trajectories, **Fig.4E**). Further, licking, the action of consuming a reward, very robustly
modulates the EPN (**Fig. 7**). Thus, it is possible that the EPN serves as online feedback
of the reward properties of an action.

How do the results in the present study fit in with models of basal ganglia function? Both
the 'firing rate model' (Albin et al., 1989) and the 'dynamic activity model' (Nambu et al.,
2023) rely on an inhibition of basal ganglia output activity to facilitate movement. When
solely considering the average population activity we did not find that movement periods
exhibited a statistically significant lower activity (**Fig. 2H**). In fact, reward (evaluation

period of correct trials) exhibited higher population activity than the wait period despite
presumably more motor output (licking) than waiting. Further, we found very complex
relationships between movement and EPN activity at different timescales, which could be
affected by aberrant oscillatory activity that is described in the 'firing pattern model'
(Guillery et al., 1997).

In this study we have focused on describing the representation of several variables in the
EPN activity. However, interpreting how this complex representation affects target nuclei
is further complicated by the fact that some subpopulations within the EPN form
multitransmitter synapses of excitatory and inhibitory transmission, particularly onto the
lateral habenula (Wallace et al., 2017). In fact, simultaneous recordings of GPi and the
ventral anterior thalamus, classically considered as the motor output, reveal only weak
correlations (Schwab et al., 2020).

A weakness of the current study is the lack of characterization of neuronal subtypes. An
area of opportunity for future research could be to perform photo-identification of neuronal
subtypes within the EPN which could contribute to the overall description of the
information representation. Further, detailed anatomical viral vector strategies could aid
to improve anatomical localization of recordings, reduce reliance on histological
examination, and solve some current controversies (Lazaridis et al., 2019).

We conclude that EPN neurons exhibit a high degree of multiplexing of diverse variables
including reward, spatio-temporal, kinematic. EPN activity can reflect single movements
like gait and licking as well as whole-body movements. We found that the correlation that
units have with these variables is unstable and varies depending on context. Finally,
spatial location of reward can influence population dynamics.

Supplemental Information

884 This article includes, 1 Statistics table, 3 Supplemental Figures and two Supplemental
885 Videos.

Acknowledgements

V.R.A.K. is a doctoral student from the Programa de Doctorado en Ciencias Biomédicas,
Universidad Nacional Autónoma de México (UNAM) and has received CONAHCyT
fellowship CVU 808903.

This work was supported by CONACyT grant 220412, Fronteras de la Ciencia CONACyT
grants 2022, 2019/154039, CF-2023-I-305, the DGAPA-PAPIIT-UNAM grants IN226517,
IN203420, IN203123 and the Moshinsky fellowship to F.T.

The authors would like to thank A Cesar Poot-Hernández and Carlos A Peralta-Alvaréz
from the Bioinformatics Unit for assistance, Gabriel Diaz-deLeon for proofreading the
manuscript, and Pavel Rueda-Orozco for helpful discussions.

Author Contributions

Bibliography

Albin, R. L., Young, A. B., & Penney, J. B. (1989). The functional anatomy of basal ganglia
disorders. *Trends in Neurosciences*, *12*(10), 366–375. https://doi.org/10.1016/0166-
2236(89)90074-X

Baker, K. B., Lee, J. Y. K., Mavinkurve, G., Russo, G. S., Walter, B., DeLong, M. R., Bakay, R.
A. E., & Vitek, J. L. (2010). Somatotopic organization in the internal segment of the
globus pallidus in Parkinson's disease. *Experimental Neurology*, *222*(2), 219–225.
https://doi.org/10.1016/j.expneurol.2009.12.030

Dayan, P., & Balleine, B. W. (2002). Reward, Motivation, and Reinforcement Learning.
Neuron, *36*(2), 285–298. https://doi.org/10.1016/S0896-6273(02)00963-7

DeLong, M. R. (1971). Activity of pallidal neurons during movement. *Journal of*
Neurophysiology, *34*(3), 414–427. https://doi.org/10.1152/jn.1971.34.3.414

Gdowski, M. J., Miller, L. E., Parrish, T., Nenonene, E. K., & Houk, J. C. (2001). Context
dependency in the globus pallidus internal segment during targeted arm movements.
Journal of Neurophysiology, *85*(2), 998–1004.
https://doi.org/10.1152/jn.2001.85.2.998

Georgopoulos, A., Delong, M. R., & Crutcher, M. D. (1983). Relations between parameters
of step-tracking movements and single cell discharge in the globus pallidus and
subthalamic nucleus of the behaving monkey. In *The Journal of Neuroscience* (Vol. 3,
Issue 8). https://doi.org/10.1523/JNEUROSCI.03-08-01586.1983

Guillery, R. W., Bergman, H., Feingold, A., Nini, A., Raz, A., Slovin, H., & Abeles, M. (1997).
Physiological aspects of information processing in the basal ganglia of normal and
parkinsonian primates. *2236*(97), 1–7. papers2://publication/uuid/1FDA7C9D-DC4C-
47A4-9FE5-0972EC09DC4B

Hintiryan, H., Foster, N. N., Bowman, I., Bay, M., Song, M. Y., Gou, L., Yamashita, S.,
Bienkowski, M. S., Zingg, B., Zhu, M., Yang, X. W., Shih, J. C., Toga, A. W., & Dong, H.
W. (2016). The mouse cortico-striatal projectome. *Nature Neuroscience*, *19*(8), 1100–
1114. https://doi.org/10.1038/nn.4332

Hong, S., & Hikosaka, O. (2008). The Globus Pallidus Sends Reward-Related Signals to the
Lateral Habenula. *Neuron*, *60*(4), 720–729.
https://doi.org/10.1016/j.neuron.2008.09.035

Hunnicutt, B. J., Jongbloets, B. C., Birdsong, W. T., Gertz, K. J., Zhong, H., & Mao, T. (2016).
A comprehensive excitatory input map of the striatum reveals novel functional
organization. *ELife*. https://doi.org/10.7554/eLife.19103.001

Kim, H. R., Malik, A. N., Mikhael, J. G., Bech, P., Tsutsui-Kimura, I., Sun, F., Zhang, Y., Li, Y.,
Watabe-Uchida, M., Gershman, S. J., & Uchida, N. (2020). A Unified Framework for
Dopamine Signals across Timescales. *Cell*, *183*(6), 1600-1616.e25.
https://doi.org/10.1016/j.cell.2020.11.013

Kobak, D., Brendel, W., Constantinidis, C., Feierstein, C. E., Kepecs, A., Mainen, Z. F., Qi,
X. L., Romo, R., Uchida, N., & Machens, C. K. (2016). Demixed principal component
analysis of neural population data. *ELife*, *5*(APRIL2016), 1–36.
https://doi.org/10.7554/eLife.10989

Larsen, K. D., & McBride, R. L. (1979). The organization of feline entopeduncular nucleus
projections: Anatomical studies. *Journal of Comparative Neurology*, *184*(2), 293–308.
https://doi.org/10.1002/cne.901840206

Lazaridis, I., Tzortzi, O., Weglage, M., Märtin, A., Xuan, Y., Parent, M., Johansson, Y., Fuzik,
J., Fürth, D., Fenno, L. E., Ramakrishnan, C., Silberberg, G., Deisseroth, K., Carlén, M.,
& Meletis, K. (2019). A hypothalamus-habenula circuit controls aversion. *Molecular*
Psychiatry, *24*(9), 1351–1368. https://doi.org/10.1038/s41380-019-0369-5

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., & Bethge, M.
(2018). DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. *Nature Neuroscience*, *21*(9), 1281–1289.
https://doi.org/10.1038/s41593-018-0209-y

McInnes, L., Healy, J., Saul, N., & Großberger, L. (2018). UMAP: Uniform Manifold
Approximation and Projection. *Journal of Open Source Software*, *3*(29), 861.
https://doi.org/10.21105/joss.00861

Mink, J. W., & Thach, W. T. (1991a). Basal ganglia motor control. I. Nonexclusive relation of
pallidal discharge to five movement modes. *Journal of Neurophysiology*, *65*(2), 273–
300. https://doi.org/10.1152/jn.1991.65.2.273

Mink, J. W., & Thach, W. T. (1991b). Basal ganglia motor control. I. Nonexclusive relation of
pallidal discharge to five movement modes. *Journal of Neurophysiology*, *65*(2), 273–
300. https://doi.org/10.1152/jn.1991.65.2.273

Mitchell, S. J., Richardson, R. T., Baker, F. H., & DeLong, M. R. (1987). The primate globus
pallidus: neuronal activity related to direction of movement. *Experimental Brain*
Research, *68*(3), 491–505. https://doi.org/10.1007/BF00249793

Mullie, Y., Arto, I., Yahiaoui, N., & Drew, T. (2020). Contribution of the entopeduncular
nucleus and the globus pallidus to the control of locomotion and visually guided gait
modifications in the cat. *Cerebral Cortex*, *30*(9), 5121–5146.
https://doi.org/10.1093/cercor/bhaa106

Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S., & Churchland, A. K. (2019). Single-trial
neural dynamics are dominated by richly varied movements. *Nature Neuroscience*,
22(10), 1677–1686. https://doi.org/10.1038/s41593-019-0502-4

Nambu, A. (2011). Somatotopic organization of the primate basal ganglia. *Frontiers in*
Neuroanatomy, *APRIL*. https://doi.org/10.3389/fnana.2011.00026

Nambu, A., Chiken, S., Sano, H., Hatanaka, N., & Obeso, J. A. (2023). Dynamic Activity
Model of Movement Disorders: The Fundamental Role of the Hyperdirect Pathway.
Movement Disorders, *38*(12), 2145–2150. https://doi.org/10.1002/mds.29646

Parent, M., Lévesque, M., & Parent, A. (1999). The pallidofugal projection system in
primates: Evidence for neurons branching ipsilaterally and contralaterally to the
thalamus and brainstem. *Journal of Chemical Neuroanatomy*, *16*(3), 153–165.
https://doi.org/10.1016/S0891-0618(99)00008-3

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D.,
Brucher, M., Perrot, M., & Duchesnay, É. (2011). Scikit-learn: Machine Learning in
Python. *Journal of Machine Learning Research*, *12*(85), 2825–2830.
http://jmlr.org/papers/v12/pedregosa11a.html

Schwab, B. C., Kase, D., Zimnik, A., Rosenbaum, R., Codianni, M. G., Rubin, J. E., & Turner,
R. S. (2020). Neural activity during a simple reaching task in macaques is counter to
gating and rebound in basal ganglia–thalamic communication. *PLoS Biology*, *18*(10).
https://doi.org/10.1371/journal.pbio.3000829

Stephenson-Jones, M., Yu, K., Ahrens, S., Tucciarone, J. M., Van Huijstee, A. N., Mejia, L.
A., Penzo, M. A., Tai, L. H., Wilbrecht, L., & Li, B. (2016). A basal ganglia circuit for
evaluating action outcomes. *Nature*, *539*(7628), 289–293.
https://doi.org/10.1038/nature19845

Thura, D., & Cisek, P. (2017). The Basal Ganglia Do Not Select Reach Targets but Control
the Urgency of Commitment. *Neuron*, *95*(5), 1160-1170.e5.
https://doi.org/10.1016/j.neuron.2017.07.039

Turner, R. S., & Anderson, M. E. (2005). Context-dependent modulation of movement-
related discharge in the primate globus pallidus. *Journal of Neuroscience*, *25*(11),
2965–2976. https://doi.org/10.1523/JNEUROSCI.4036-04.2005

Wallace, M. L., Saunders, A., Huang, K. W., Philson, A. C., Goldman, M., Macosko, E. Z.,
McCarroll, S. A., & Sabatini, B. L. (2017). Genetically Distinct Parallel Pathways in the
Entopeduncular Nucleus for Limbic and Sensorimotor Output of the Basal Ganglia.
Neuron, *94*(1), 138-152.e5. https://doi.org/10.1016/j.neuron.2017.03.017

1008

1009

1010

1011

1012

1013

Figure 1. Mice learn to perform a two-alternative forced choice task in a freely moving triangular arena.

A. Left: Schematic of triangular arena. Right: stimuli were frequency modulated sweeps.

B. Time schematic of task events. The two periods of movement are highlighted.

C. Video tracking extracted variables of a single trial. Note that events correspond to B.

D. Left: Psychophysical performance of an example session of an animal. Middle: Thin traces are 5 example days pre (orange) and post (black) electrode implant for an example animal. Thicker traces are averages of pre and post-performance. Right: Thin traces are individual animal averages (n=6) of the 32 recording sessions included in this study. Thicker trace is group average.

E. Average reaction time as a function of stimulus deltaFreq. Black line is the linear fit.

F. Probability of correct response as a function of reaction time. Traces are grouped per absolute deltaFreq (left and right responses are grouped). Probability of correct responses is computed in reaction time bins of 75ms.

G. Instantaneous angular speed (radial axis) of turning per stimulus condition. 90º represents the animals' position during waiting.

H. Mean angular speed per stimulus identity for correct (colored circles) and incorrect (gray circles) trials. Trendlines were drawn following a simple linear regression. Legend shows correlation coefficient and statistical significance.

I. Instantaneous displacement speed across the length of the trajectory for left and right stimuli.

J. Mean average speed per stimulus for correct (colored circles) and incorrect (gray circles) trials. Trendlines were drawn following a simple linear regression. Legend shows correlation coefficient and statistical significance.

Figure 2. Electrophysiological recordings were obtained from the EPN.

A. Microwire bundles were implanted onto the entopeduncular nucleus. Right pane shows a schematic of electrode placement (n=6).

B. Left: Photomicrograph of electrode cannula track. Right: Close up of individual microwire electrode tracks. Red arrow heads point to microwire tips.

C. Peak to valley duration of plotted against inter-spike interval coefficient of variation for recorded spikes. Insets: average waveform for four example units.

D. Example recorded unit aligned to events identified in Figure 1 B: return, wait, turn (0º,75º,150º), evaluation, averaged according to stimulus presented. First row is rescaled average for correct trials. Second row is average of incorrect trials. Third row is false alarm trials: trials where animals did not

wait long enough for the stimulus to appear but that performed the entire movement sequence either to the right or left lick ports.

E. Same as D for another example unit.

F. Population response for correct left and right responses.

G. Upper: Average z-score segregated by left and right, for correct, incorrect, and false alarm trials. Lower: Average z-score per stimulus for correct trials.

H. Average speed (upper) and angular velocity (lower) for trials segregated as in the upper pane of G.

I. Mean population activity per epoch segregated by movement, turning, and Reward. Wilcoxon signed-rank test used.

Figure 3. Stimulus, difficulty, reward and temporal population dynamics during Go trajectories.

A. Instantaneous variance associated with trials segregated by left-right, correct/incorrect, and stimuli identity.

B. Principal Components and demixed Principal Components sorted by explained variance. Note that the difficulty dPCA does not appear within the 10 first components. Statistically significant PCs (n=5) were signaled by a bigger dot.

C-G. First five dPCs sorted by variance are shown: Temporal (condition-independent), reward, and side. Note that for traces in C-E there is a near perfect overlap of all conditions (n=16). Weights for each individual dPCs are shown to the right. Lettered circles on top are individual units' weights shown on Figure 3-1. G.

H. Population self-similarity across time. Cosine similarity was calculated for population vectors on the de-meaned (temporal dynamics removed) population activity (see Methods).

I. Population vector similarity across time for two time points: t=0.2s and t=1.8, which correspond to the moments of peak side and reward variance, respectively.

Figure 3-1. Example units. a-h.
Example units are shown corresponding to lettered circles on weight plots in Figure 3.

Figure 3-2. Stimulus presence, irrespective of identity, and angular velocity best explain EPN activity around turning epoch.

A. Example unit aligned to stimulus (left) and turn (right), segregated by left and right, correct, incorrect and false alarm trials. First row is raster plots. Middle row is average firing rate. Lower row is average angular velocity. Note that left column does not contain false alarm trials since there is no stimulus to align to.

B. Similar to A for another unit.

C. Models fit to data. Four hypotheses were tested. First row is the presence of stimulus (without identity). Second row is difficulty (absolute value of deltaFreq). Third row is stimulus interpretation (incorrectly performed trials show opposite interpretation to the correctly performed counterparts). Fourth row shows angular velocity.

D. deltaR2 calculated for each of the models (see Methods and main text). Statistical differences are shown with red lines, p<0.05, Wilcoxon signed-rank test.

Figure 4. Kinematic and spatio-temporal coding during Return and Go trajectories.

A-C show three example units. Raster plots show activity for return (left) and Go (right) trajectories sorted by trajectory duration (black circle). They were then grouped into six traces, and average firing rate is presented below. Also, the average speed of the animals locomotion is presented below. Simple linear regression for four different variables presented on the right panels, for Return (orange) and Go (green) trajectories separately.

D. Average tenfold cross-validated R2 for multiple regression models fit on data from the entire trial duration (return, wait, go, and evaluation periods) for kinematic, spatio-temporal and reward related variables, as well as full model fit on all these variables (n=118 units). Red line represents a statistically significant difference (p<0.01, Wilcoxon signed-rank test).

E. Average tenfold cross-validated R2 for models for multiple regression models fit separately on Return (orange) and Go (green) trajectory data. Red line represents a statistically significant difference (p<0.01, Wilcoxon signed-rank test).

F. Mean cvR2 for single variable models. Shading corresponds to kinematic (gray) and spatio-temporal (pink) variables. Reward related variables are unshaded. Models were fit with Return (orange), Go (green) and entire trial duration (gray) data.

G. Trained models for Return and Go periods in E were tested using Return and Go data. Individual models for kinematic, spatio-temporal and a mixed kinematic and spatio-temporal model were tested. Note all models tested on data not used for training were significantly lower than when using training data (p<0.01, Wilcoxon signed-rank test).

H. Upper: Partial correlation coefficient (r) was calculated for the distance to goal variable by fitting residuals of model on kinematic variables (excluding spatio-temporal variables). This was done separately for Return (x-axis) and Go (y-axis) data. Dashed line corresponds to the identity line (x=y). Lower: Pie chart summarizing data above. Brown: Units with statistically significant correlations with the distance to goal variable in Go and Return periods. Green: units with statistically significant correlation only during the Go period. Orange: units with statistically significant correlation only during the Return period. Outer pie summarizes the sign of the correlations where gray and salmon represent negative and positive correlations respectively, in both Return and Go periods. Black represents units that switch signs in Return and Go periods.

I. Upper: Partial correlation coefficient of residuals of a model with all spatio-temporal and kinematic variables except angular velocity, separately for Return and Go trajectories. Lower: similar to H.

J. Upper: Partial correlation coefficient of residuals of a model with all spatio-temporal and kinematic variables except body speed, separately for Return and Go trajectories. Lower: similar to H.

K. Venn diagrams of percentage of total units with a significant partial correlation coefficient for the variables distance to goal, angular velocity and body speed for Return and Go trajectories.

L. Left: Mean zScore activity of units segregated according to activity during reward. Top panel shows mean of units unresponsive to reward. Middle and lower panel shows average activity of units with statistically significant difference between rewarded and unrewarded trials in a 250ms window after head entry into the lick port. Middle panel corresponds to reward positive units that showed an increase in firing rate with rewarded trials (auROC>0.5, p<0.001 permutation test) while the lower panel corresponds to reward negative units that showed a decrease in activity with rewarded trials (auROC<0.5, p<0.001 permutation test).

Right: Dimensionality reduction performed on the regression coefficients on a model fitted Return and Go data using kinematic and spatiotemporal variables. Red dots represent reward positive units

(rewarded trials with higher firing rate that unrewarded ones) and blue ones represent reward negative units (rewarded trials with lower firing rate).

M. Mean cross-validated R2 of models fit on kinematic, spatio-temporal and both types of variables, segregated by reward coding of units (segregated as described in L). Red lines indicate statistically significant difference (Mann-Whitney U test, p<0.01).

N. Mean cross-validated R2 of kinematic, spatio-temporal, and mixed models. Data was segregated according to trial type: correct, incorrect, and false alarm.

Figure 4-1. Variables used to fit models.

A. Single trial variables used to fit models in Figure 4.

B. Correlation coefficient calculated between pairs of variables.

Figure 5. Return and Go trajectories population dynamics.

A. Highly similar trajectories were found for Return and Go periods based on body speed and angular velocity (see Methods). Average body speed and angular velocity are plotted for the four conditions found.

B. Population trajectories corresponding to the four conditions shows in A projected onto three PCs.

C. Cosine similarity of the four conditions based on kinematic parameters (body speed and angular velocity) and population activity.

D. Context dPC with most explained variance. Histogram of weights is presented on the right.

E. Correlation coefficient between the extracted population dynamics in D and several variables calculated for the same periods.

F. Distance from Waiting corner calculated for the four conditions.

G. Context dPC with second most explained variance. Histogram of weights is presented on the right.

H. Correlation coefficient between the extracted population dynamics in G and several variables calculated for the same periods.

I. Velocity from Waiting corner calculated for the four conditions.

Figure 5-1. demixed Principal components for Return and Go trajectories, continued

Projections of population activity onto the first three PCs with most explained variance for the four conditions in 5A, and separately for right trajectories in A and left trajectories in B.

C. Principal Components and demixed Principal Components sorted by explained variance. Statistical significance of PCs (n=6) is indicated by bigger dots.

D-G. temporal (condition-independent) dPCs and Side in D. are presented with their weight histogram and the correlation of these components with individual kinematic and spatio-temporal variables.

Figure 6. Gait cycle modulation of units.

A. Longitudinal position of four paws of a mouse during a return, wait and go periods. Inset shows body position in time. Colored triangles show identified events for unit alignment.

B. Activity of two simultaneously recorded example units in the eight strides studied. Lower row shows average paw position corresponding to the trials shown above.

C. Population activity during the four strides.

D. Spike gait-phase average vector for all significantly modulated units during eight strides.

E. Upper: Fraction of modulated units (with a significant directionality Rayleigh test $p<0.001$). Middle: Mean phase of population vector. Lower: Mean population vector magnitude.

F. Variance explained by PCA decomposition of data shown in C. Inset shows first four PCs' weight distribution.

G. Upper: mean absolute correlation coefficient for kinematic and spatio-temporal variables for the first four PCs.

H. Population projection of data in C onto PC1, PC2 and PC4 for the four Return strides (in orange tones) and the Go strides (in green tones).

I. Left: PC4 population projection for the eight strides studied plotted on top of contralateral hindpaw velocity. Right: correlation coefficient with individual kinematic and spatio-temporal variables. Filled circles represent a statistically significant correlation (permutation testing).

J. Population Support Vector Classifiers were trained for different variables (x-axis). Accuracy was assessed for the classifier trained on the original labeled population vectors (in blue) and for vectors trained on shuffled labels (in orange). *=$p<0.001$, permutation testing. Note that paw velocity could not be decoded from the original matrix but could be decoded from the PC4 weighted matrix.

Figure 7. Licking behavior modulation of units.

A. Infrared sensor based lickometer.

B. Licks of aligned to port entry for left (red) and right (blue) correct trials.

C. Example unit activity aligned to the same port entries as in B.

D. Upper: Three simultaneously recorded example units' (blue, orange and green) activity rescaled to average lick bout duration. Lick rate is shown in gray. Lower: Histogram of Pearson correlation coefficient between instantaneous firing rate and instantaneous lick rate for all units. Black bars represent significantly correlated units (p<0.001, permutation test).

E. Example unit shown in C aligned to single left (red) and right (blue) licks. Upper inset shows the probability occurrence as well as the lick cycle phase derived from the lick probability. Right inset shows the spike lick-phase polar histogram for same unit. Black line represents resulting vector with statistically significant directionality (Rayleigh test, p<0.001).

F. Polar histogram (bin= 30º) of average vector phase direction. Polar axis is percentage of units.

G. Average single lick activity for all recorded units sorted by peak activity. Upper inset: average lick probability.

H. Venn diagram showing percentage of units of the total population (n=118) of units with statistically significant correlation with lick rate (Pearson correlation coefficient, p<0.001, permutation test) and/or with single licks (Rayleigh directionality test, p<0.001).

I. Single lick average activity segregated by first and last lick, 5 interspersed licks, and incorrect lick for left and right licks for the same unit shown in C and E.

J. Population activity for n=118 units for 7 right licks sorted by lick with most activity. Lower: fraction of variance explained by PCA decomposition PCA weight for first 3 components.

K. Projection of lick population activity for the 16 licks considered onto first 3 PCs.

L. Population vector decoders for different variables with original labels (blue) compared to population vector decoders with shuffled labels (n=1000). *p<0.001, permutation test

Table 1. Statistics table

Figure	Descriptive statistics	Type of test	p value	n
1E	Mean +/- SEM Simple linear regression	Correlation coefficient two-tailed permutation test	0.006	7 data points
1H	Mean +/- SEM Simple linear regression	Correlation coefficient two-tailed permutation test	Correct trials=0.004 Incorrect trials= 0.17	7 data points
1J	Mean +/- SEM Simple linear regression	Correlation coefficient two-tailed permutation test	Correct trials=0.30 Incorrect trials= 0.097	7 data points
2I	Mean zScore	Wilcoxon signed-rank test	Return vs Go=0.17 Return vs Wait=0.20 Return vs Contralateral=0.48 Return vs Ipsilateral=0.12 Return vs Correct=4.0×10^{-5} Return vs Incorrect=0.88 Go vs Wait=0.11 Go vs Contralateral=0.43 Go vs Ipsilateral=0.51 Go vs Correct=0.0011 Go vs Incorrect=0.36 Wait vs Contralateral=0.35 Wait vs Ipsilateral=0.0024 Wait vs Correct=1.2×10^{-5} Wait vs Incorrect=0.42 Contralateral vs Ipsilateral=0.034 Contralateral vs Correct=0.00014 Contralateral vs Incorrect=0.59 Ipsilateral vs Correct=0.0017 Ipsilateral vs Incorrect=0.13 Correct vs Incorrect=1.7×10^{-5}	118 data points per condition
3-2D	Mean delta R2 (see Methods)	Wilcoxon signed-rank test	**Model comparison** Stimulus vs Difficulty=1.8×10^{-5} Stimulus vs Interpretation=0.0021 Stimulus vs AngVel=0.012	118 data points per condition

			Difficulty vs Interpretation=0.35 Difficulty vs AngVel=3.9×10^{-9} Interpretation vs AngVel=2.8×10^{-8}	
4D	Mean	Wilcoxon signed-rank test	Kine vs spatio=5.1×10^{-10} Kine vs reward=0.016 Kine vs kine+spatio+reward=8.6×10^{-20} Spatio vs reward=0.16 Spatio vs kine+spatio+reward=6.1×10^{-20} Reward vs kine+spatio+reward=1.9×10^{-20}	118 units
4E	Mean	Wilcoxon signed-rank test	**Kinematic model** Return vs Go=5.3×10^{-8}	118 units
4E	Mean	Wilcoxon signed-rank test	**Spatio-temporal model** Return vs Go=3.2×10^{-4}	118 units
4E	Mean	Wilcoxon signed-rank test	**Kine+spatio model** Return vs Go=4.5×10^{-6}	118 units
4G	Mean +/- SEM	Wilcoxon signed-rank test	**Kinematic model trained on Return data, tested on:** Return vs Go=2.2×10^{-15} **Kinematic model trained on Go data, tested on:** Return vs Go=4.5×10^{-21}	118 units
4G	Mean +/- SEM	Wilcoxon signed-rank test	**Spatio-temporal model trained on Return data, tested on:** Return vs Go=4.0×10^{-19} **Spatio-temporal model trained on Go data, tested on:** Return vs Go=5.4×10^{-21}	118 units
4G	Mean +/- SEM	Wilcoxon signed-rank test	**Kine+spatio model trained on Return data, tested on:** Return vs Go=2.4×10^{-19} **Kine+spatio model trained on Go data, tested on:** Return vs Go=4.2×10^{-21}	118 units
4M	Mean	Mann–Whitney U test	**Kinematic model** Rew-pos vs rew-neg=0.015 Rew-pos vs no-rew=0.62 Rew-neg vs no-rew=0.051	Rew-pos= 51 units Rew-neg= 22 units

				No-rew= 45 units
4M	Mean	Mann–Whitney U test	**Spatio-temporal model** Rew-pos vs rew-neg=0.017 Rew-pos vs no-rew=0.79 Rew-neg vs no-rew=0.060	Rew-pos= 51 units Rew-neg= 22 units No-rew= 45 units
4M	Mean	Mann–Whitney U test	**Kine+spatio model** Rew-pos vs rew-neg=0.029 Rew-pos vs no-rew=0.61 Rew-neg vs no-rew=0.11	Rew-pos= 51 units Rew-neg= 22 untis No-rew= 45 units
4N	Mean	Wilcoxon signed-rank test	**Kinematic model** Correct vs incorrect=0.62 Correct vs false alarm= 0.07 Incorrect vs false alarm= 0.19	118 units
4N	Mean	Wilcoxon signed-rank test	**Spatio-temporal model** Correct vs incorrect=0.74 Correct vs false alarm=0.02 Incorrect vs false alarm=0.08	118 units
4N	Mean	Wilcoxon signed-rank test	**Kine + spatio model** Correct vs incorrect=0.72 Correct vs false alarm=0.06 Incorrect vs false alarm=0.12	118 units
5E	Pearson correlation coefficient	Pearson correlation coefficient two-tailed permutation test	**Context dPC1 (c-dPC1)** c-dPC1 vs Head speed=0.54 c-dPC1 vs Body speed=0.96 c-dPC1 vs Head acceleration=0.18 c-dPC1 vs Body acceleration=0.27 c-dPC1 vs angVel=0.88 c-dPC1 vs dist from X=0.0 c-dPC1 vs dist from Y=0.0 c-dPC1 vs dist Waiting=0.0 c-dPC1 vs vel from Waiting corner=0.01 c-dPC1 vs dist Goal=0.62 c-dPC1 vs time=0.46	4 time series (1.2s, 100Hz) per dPC vs 4 time series (1.2s, 100Hz) per variable (480 points vs 480 points)

5H	Pearson correlation coefficient	Pearson correlation coefficient two-tailed permutation test	**Context dPC2 (c-dPC2)** c-dPC2 vs Head speed=0.53 c-dPC2 vs Body speed=0.57 c-dPC2 vs Head acceleration=0.35 c-dPC2 vs Body acceleration=0.79 c-dPC2 vs angVel=0.77 c-dPC2 vs dist from X=0.0 c-dPC2 vs dist from Y=0.0 c-dPC2 vs dist Waiting=0.0 c-dPC2 vs vel from Waiting corner=0.0 c-dPC2 vs dist Goal=0.57 c-dPC2 vs time=0.85	4 time series (1.2s, 100Hz) per dPC vs 4 time series (1.2s, 100Hz) per variable (480 points vs 480 points)
5-1D	Pearson correlation coefficient	Pearson correlation coefficient two-tailed permutation test	**Temporal dPC1 (t-dPC1)** t-dPC1 vs Head speed=0.0 t-dPC1 vs Body speed=0.37 t-dPC1 vs Head acceleration=0.0 t-dPC1 vs Body acceleration=0.0 t-dPC1 vs angVel=0.93 t-dPC1 vs dist from X=0.77 t-dPC1 vs dist from Y=0.4 t-dPC1 vs dist Waiting=0.86 t-dPC1 vs vel from Waiting corner=0.32 t-dPC1 vs dist Goal=0.0 t-dPC1 vs time=0.0	4 time series (1.2s, 100Hz) per dPC vs 4 time series (1.2s, 100Hz) per variable (480 points vs 480 points)
5-1E	Pearson correlation coefficient	Pearson correlation coefficient two-tailed permutation test	**Temporal dPC2 (t-dPC2)** t-dPC2 vs Head speed=0.0 t-dPC2 vs Body speed=0.0 t-dPC2 vs Head acceleration=0.0 t-dPC2 vs Body acceleration=0.05 t-dPC2 vs angVel=0.82 t-dPC2 vs dist from X=0.01 t-dPC2 vs dist from Y=0.05 t-dPC2 vs dist Waiting=0.48 t-dPC2 vs vel from Waiting corner=0.76 t-dPC2 vs dist Goal=0.0 t-dPC2 vs time=0.0	4 time series (1.2s, 100Hz) per dPC vs 4 time series (1.2s, 100Hz) per variable (480 points vs 480 points)
5-1F	Pearson correlation coefficient	Pearson correlation coefficient	**Temporal dPC3 (t-dPC3)** t-dPC3 vs Head speed=0.32 t-dPC3 vs Body speed=0.77	4 time series (1.2s,

		two-tailed permutation test	t-dPC3 vs Head acceleration=0.94 t-dPC3 vs Body acceleration=0.0 t-dPC3 vs angVel=0.67 t-dPC3 vs dist from X=0.7 t-dPC3 vs dist from Y=0.28 t-dPC3 vs dist Waiting=0.65 t-dPC3 vs vel from Waiting corner=0.94 t-dPC3 vs dist Goal=0.0 t-dPC3 vs time=0.0	100Hz) per dPC vs 4 time series (1.2s, 100Hz) per variable (480 points vs 480 points)
5-1G	Pearson correlation coefficient	Pearson correlation coefficient two-tailed permutation test	**Side dPC1 (s-dPC1)** s-dPC1 vs Head speed=0.39 s-dPC1 vs Body speed=0.53 s-dPC1 vs Head acceleration=0.32 s-dPC1 vs Body acceleration=0.18 s-dPC1 vs angVel=0.0 s-dPC1 vs dist from X=0.0 s-dPC1 vs dist from Y=0.0 s-dPC1 vs dist Waiting=0.92 s-dPC1 vs vel from Waiting corner=0.64 s-dPC1 vs dist Goal=0.93 s-dPC1 vs time=0.42	4 time series (1.2s, 100Hz) per dPC vs 4 time series (1.2s, 100Hz) per variable (480 points vs 480 points)
6G, lower	Pearson correlation coefficient	Pearson correlation coefficient two-tailed permutation test	Paw position vs PC1=0.77 Paw position vs PC2=0.81 Paw position vs PC3=0.11 Paw position vs PC4=0.30 Paw velocity vs PC1=0.71 Paw velocity vs PC2=0.52 Paw velocity vs PC3=0.83 Paw velocity vs PC4=0.0	200 points vs 200 points
6 I, right	Pearson correlation coefficient	Pearson correlation coefficient two-tailed permutation test	PC4 vs velHead=0.0 PC4 vs velBody=0.0 PC4 vs accHead=0.58 PC4 vs accBody=0.93 PC4 vs angVel=0.07 PC4 vs paw1=0.11 PC4 vs paw2=0.11 PC4 vs paw3=1.0 PC4 vs paw4=0.3 PC4 vs paw1vel=0.0 PC4 vs paw2vel=0.0	200 points vs 200 points

			PC4 vs paw3vel=0.0 PC4 vs paw4vel=0.0 PC4 vs distX=0.03 PC4 vs distY=0.01 PC4 vs distE=0.05 PC4 vs distGoal=0.0 PC4 vs timeGoal=0.0	
6J, left	Mean +/- SEM	Permutation test	**Classifier accuracy, original matrix** Context original labels vs shuffled=0.0 Step # original labels vs shuffled=0.0 Paw position original labels vs shuffled=0.478 Paw velocity original labels vs shuffled=0.431	1 vs 1000
6J, right	Mean +/- SEM	Permutation test	**Classifier accuracy, PC4-weighted matrix** Context original labels vs shuffled=0.0 Step # original labels vs shuffled=0.0 Paw position original labels vs shuffled=0.972 Paw velocity original labels vs shuffled=0.0	1 vs 1000
7L	Mean +/- SEM	Permutation test	**Classifier accuracy** Phase original labels vs shuffled=0.0 Left-right original labels vs shuffled=0.0 Lick # original labels vs shuffled=0.0 Correct vs incorrect original labels vs shuffled=0.0	1 vs 1000