1

Attentional modulation of secondary somatosensory and visual thalamus of mice

Gordon H Petty[1,2] and Randy M Bruno[1,2,*]

[1] Department of Neuroscience, Columbia University, New York, NY 10027 USA

[2] Department of Physiology, Anatomy, & Genetics, University of Oxford, Oxford OX1 3PT, United Kingdom

* Correspondence: randy.bruno@dpag.ox.ac.uk (R.M.B.)

ABSTRACT

Each sensory modality has its own primary and secondary thalamic nuclei. While the primary thalamic nuclei are well understood to relay sensory information from the periphery to the cortex, the role of secondary sensory nuclei is elusive. One hypothesis has been that secondary nuclei may support feature-based attention. If this is true, one would also expect the activity in different nuclei to reflect the degree to which modalities are or are not behaviorally relevant in a task. We trained head-fixed mice to attend to one sensory modality while ignoring a second modality, namely to attend to touch and ignore vision, or vice versa. Arrays were used to record simultaneously from secondary somatosensory thalamus (POm) and secondary visual thalamus (LP). In mice trained to respond to tactile stimuli and ignore visual stimuli, POm was robustly activated by touch and largely unresponsive to visual stimuli. A different pattern was observed when mice were trained to respond to visual stimuli and ignore touch, with POm now more robustly activated during visual trials. This POm activity was not explained by differences in movements (i.e., whisking, licking, pupil dilation) resulting from the two tasks. Post hoc histological reconstruction of array tracks through POm revealed that subregions varied in their degree of plasticity. LP exhibited similar phenomena. We conclude that behavioral training reshapes activity in secondary thalamic nuclei. Secondary nuclei may respond to behaviorally relevant, reward-predicting stimuli regardless of stimulus modality.

INTRODUCTION

Each of the primary sensory cortices are reciprocally connected with corresponding sensory thalamic nuclei. These nuclei are broadly classified as primary or secondary (also known as high-order)[1–3]. The primary nuclei respond robustly to sensory stimulation and transmit sensory information to the cortex[4–7]. By contrast, the secondary nuclei receive sparse input from the sensory periphery and are connected with many cortical and subcortical regions. They have nonspecific or complex receptive fields, and their role in sensation and cognition is not well understood.

The rodent primary somatosensory cortex (S1) is connected to the posterior medial nucleus (POm), the secondary somatosensory thalamic nucleus[8,9]. Despite reciprocal connections with the whisker-sensitive region of S1 (barrel cortex), POm poorly encodes whisker touch and movement[10–13]. However, POm contributes to whisker-dependent behaviors[14,15], facilitates cortical plasticity[16–18], and encodes non-sensory task-related behaviors[19]. Despite these recent findings, POm activity in awake, behaving animals is relatively poorly understood.

There is evidence that the secondary nuclei also play a critical role in attention. The lateral posterior nucleus (LP) is the secondary visual thalamic nucleus in rodents, homologous to the lateral pulvinar in primates. In humans, damage to lateral pulvinar causes attentional deficits[20–22]. Similarly, pulvinar silencing in nonhuman primates induces transient impairment in attention tasks[23,24]. Secondary visual thalamus has not yet been similarly studied in rodents.

Despite belonging to ostensibly distinct sensory systems, there are several similarities in the activity and neuroanatomy between LP and POm: Both regions target primary and high-order sensory cortex in the same layer-specific way[7,19,25,26], have broad sensory receptive fields[11,26–28], and are strongly dependent on cortical activity rather than afferent sensory input[10,29–31]. These similarities suggest that the secondary nuclei could play similar roles in sensory processing and attention.

Here, we investigated how sensory-, arousal-, and movement-evoked activity in LP and POm are shaped by associative learning. We conditioned head-fixed mice to attend to a stimulus of one sensory modality (visual or tactile) and ignore a stimulus of a second modality. We then recorded simultaneously from POm and LP. In mice trained to attend to a tactile stimulus and ignore a visual stimulus, tactile responses dominated throughout POm. In contrast, POm showed widespread visual responses in mice trained to respond to a visual stimulus and ignore touch. Unexpectedly LP exhibited responses similar to POm's. Thus, behavioral training reshapes activity in secondary thalamus, inducing responses to behaviorally relevant stimuli regardless of sensory modality.

RESULTS

Mice can attend to one sensory modality while ignoring a second modality

We developed a conditioning task in which mice were trained to associate a stimulus of one sensory modality with a reward while ignoring a stimulus of a different modality. Water-restricted mice were head fixed and presented with a visual stimulus (a drifting grating on a monitor) and a tactile stimulus (an innocuous air puff to the distal ends of the whiskers) (Figure 1a). Mice were separated into two cohorts: in the "tactile" cohort a water reward was given at the offset of the air puff (Figures 1b,c, top; n=11 mice); in the "visual" cohort the water reward was given at the offset of the drifting grating (bottom; n=12 mice).

Initially we trained mice on a shaping version of the task which followed a traditional trial-based structure (Figure 1b). A stimulus of one modality or the other was randomly presented on each trial. Trial-based structures are undesirable for our purposes because the unrewarded stimulus is informative that no reward will occur and may garner the animal's attention. Therefore, after reaching criterion (see Methods), mice were moved to the full version of the task. In the full task the stimuli were presented at the same mean rates as in the shaping task but were completely uncorrelated with one another (Figure 1c). After a stimulus, the time until the next stimulus of the same type was drawn from an exponential distribution plus a linear offset (Figure 1d, top). The *hazard rate* of the stimulus (the probability of the stimulus occurring at various times since the previous stimulus) was thus flat for the majority of possible inter-stimulus intervals (ISIs) (Figure 1d, bottom). Visual and tactile stimulus intervals were drawn independently, and the stimuli could overlap. As a result, given a stimulus, a mouse could not predict the timing of the next stimulus of either type. In pilot experiments, naive mice trained on the full version of the task learned at a much slower rate than mice initially trained on the shaping version of the task and then advanced to the full task. We therefore trained all mice first on the shaping version of the task for at least three days and until they reached criterion before moving them to the full version (see Methods).

We examined anticipatory licking to assess how mice learned the stimulus-reward association (Figure 1e-j). For each stimulus presentation we computed a lick index (the difference in the number of licks during stimulus presentation and two seconds prior to stimulus onset, divided by the sum of all licks in both periods) [32]. A positive lick index indicates that a mouse responded to a stimulus by licking, a negative index indicates that it withheld licking, and an index of zero indicates that it ignored the stimulus.

An example visually conditioned mouse is presented in Figure 1e and f. During the shaping task, this mouse initially ignored both stimuli: the mouse did not lick in response to either the air puff or drifting grating, though it did lick to consume the water reward (Figure 1e, left). After several shaping sessions, the mouse learned that the drifting grating was predictive of a reward, as evidenced by an increase in licking after the onset of the drifting grating but before reward delivery (Figure 1e middle). However, this mouse also learned that the air puff predicted a *lack* of reward, as evidenced by withholding licking upon the onset of the air puff. The mouse thus displayed a positive visual lick index and a negative tactile lick index, suggesting that it attended to both the tactile and visual stimuli (Figure 1f, middle arrow).

In the full version of the task, the two stimuli are completely decorrelated, and the air puff is no longer predictive of the presence or absence of reward. After several sessions of conditioning on the full version of the task, the mouse no longer altered its licking in response to the air puff (Figure 1e, right; Figure 1f,

rightmost arrow). This indicates that the mouse learned to ignore the uninformative, distracting stimulus.
All visually conditioned mice exhibited a similar learning trajectory (Figure 1i, left; 1j, left). These mice
initially learned that the air puff indicated the absence of reward, sometimes before they responded to
the visual stimulus at all. They only learned to ignore the air puff after several sessions of conditioning on
the full version of the task. By contrast, tactilely conditioned mice rapidly learned the association between
air puff and reward, displaying a positive lick index as early as session 1 (Figure 1g, left; 1h). Tactilely
conditioned mice always ignored the drifting grating, never displaying a visual lick index significantly
different from zero (Figure 1i, right; Figure 1j right). Such a discrepancy suggests that mice found this
particular tactile stimulus more salient than the drifting grating. All mice were first trained on the shaping
version of the task for at least three sessions and until they learned the association between the
conditioned stimulus and reward (Methods). Mice were switched to the full version of the task and trained
for at least four more days and until they no longer responded to the unrewarded stimulus. All tactilely
conditioned mice reached criterion within six to nine sessions, and all visually conditioned mice learned
within nine to thirteen sessions.

Though the visual version of the task took longer to learn, there was no difference in how the two groups
responded to the conditioned stimulus in fully trained mice: tactilely conditioned mice licked in response
to the air puff with the same latency that visually conditioned mice responded to the drifting grating
(Figure S1c, tactile conditioning median latency of 1.1 s, visual conditioning latency of 1.0 s, p=0.98, signed-
rank test). In both groups we observed a broad range of response times across mice, with some mice
responding as early as 500 ms after stimulus onset and others responding as late as 1.6 s after onset (Figure
S1a, b). In summary, mice can learn to disregard uninformative stimuli, even when those stimuli are in and
of themselves highly salient.

POm responses to tactile and visual stimuli depend on conditioning paradigm
To evaluate the effect of reward conditioning on activity in secondary sensory thalamus, we performed
electrophysiological recordings in mice fully trained on the conditioning task. We inserted a 64-channel
multielectrode array that spanned both LP and POm. Probes were coated in DiI, and histology images were
aligned (see Methods) to identify the location of each putative cell (Figure 2a-c). Spike clusters were
manually curated based on waveform shape and spike autocorrelation (Methods, Figure S2a-d). We
identified 347 POm cells and 64 LP cells from 11 tactilely conditioned mice, and 257 POm cells and 67 LP
cells from 12 visually conditioned mice. Our recordings covered most of the volume of POm but were
clustered primarily in the anterior and medial portions of LP (Figure 2d-f). Cells that were within 50 μm of
a region border were excluded from analysis. We observed no difference in waveform shape between POm
and LP, though POm cells tended to have a higher amplitude than LP cells (Figure S2e-i).

During recording, mice were presented with an equal number of air puffs and drifting gratings (80-120 of
each stimulus per mouse, median 100). We first investigated how conditioning affects POm activity. We
observed a heterogeneous population of cells with varied patterns of activity. For example, in tactilely
conditioned mice, we observed certain cells with a sharp increase in firing rate at air puff onset (Figure 3a,
left) and other cells that were primarily active after the stimulus offset when the mouse was consuming
the water reward (Figure 3a, right). A large portion of cells responded most strongly to the stimulus onset,
having a peak firing rate within 250ms of the start of the air puff (70 cells, 20.2%). However, we also

observed cells with peak firing rates spanning the entire duration of the air puff and up to four seconds
after air puff offset (Figure 3c). Of cells from tactilely conditioned mice, 195 (56.2%) significantly modified
their firing rate within one second of the air puff onset (Figure 3d, bottom). Of these cells, only 23 cells
(6.6%) had significant changes in firing rate following the onset of the drifting grating (Figure 3d, top). No
cells responded to the drifting grating alone.

POm cells from visually conditioned mice exhibited a strikingly different pattern of activity (Figures 3b, e,
f). Surprisingly, many POm cells responded to the drifting grating alone or to both the drifting grating and
the air puff (Figure 3b, left). Others had no response to either stimulus but were more active during reward
consumption (Figure 3b, right). Compared to tactilely conditioned mice, a much larger portion of cells
responded to the onset of the drifting grating and a smaller portion responded to the air puff (Figure 3f,
16 air puff responding cells, 6.2%; 12 grating responding cells, 4.7%; 85 cells responding to both stimuli,
33.1%). At the population level, visually conditioned POm cells had a significantly higher firing rate than
tactilely conditioned POm cells during the first second of the drifting grating stimulus period (Figure 3g,
h). Furthermore, POm visual responses often preceded a mouse's lick response (Figure S1). In summary,
POm displayed dramatically different responses to visual and tactile stimuli depending on the task type.

In both conditioning groups, POm activity increased when mice were consuming water. We examined firing
rates during the "offset" period, defined as the two-second period after stimulus offset and during the
reward delivery (or absence of reward for the unconditioned stimulus). In both groups, cells had a greater
firing rate during the offset period of the rewarded stimulus than the unrewarded stimulus (Figure 3h,
Wilcoxon signed-rank test, $p<10^{-20}$ for both groups).

Conditioning alters POm response latencies to a tactile stimulus
Though POm poorly encodes whisker touch compared to other somatosensory brain regions, whisker
stimulation does induce low-latency (<100ms) responses in certain POm cells, even in anaesthetized
animals[10]. We observed POm cells that responded to air puff onset in both visually and tactilely
conditioned mice (Figure 3c, e, g). Such responses could be "purely sensory" (*i.e.* driven by ascending
brainstem inputs) or could result from consequent changes in movement, arousal, and behavioral state.

To investigate how conditioning might affect early tactile stimulus responses, we analyzed the mean firing
rates of each POm cell immediately after the air puff at a finer timescale (Figure 4). We limited our analysis
to the first 500ms after stimulus onset, as it precedes the majority of licking activity across mice (Figure
S1). At the population level, POm cells in both conditioning groups had a peak of activity 40ms after air
puff onset (Figure 4a). In tactilely conditioned mice POm activity remained elevated throughout the
duration of the two-second air puff, while in visually conditioned mice POm activity attenuates over air
puff presentation (Figure 4a, 3g).

As these activity peaks were similar in both timing and magnitude, we considered that there exists a subset
of POm cells in both conditioning groups with low-latency stimulus responses driven directly by whisker
stimulation. For each cell, we computed a stimulus response latency, defined as the first time after the
stimulus when the cell's firing rate differed statistically from baseline (Methods). Interestingly, we
observed a similar proportion of cells with air puff response latencies below 100ms in both conditioning
groups (Figure 4b). However, in tactilely conditioned mice there were nearly twice as many cells

responding between 100 and 500ms after air puff onset (32.0% of tactilely conditioned cells and 18.3% of
visually conditioned cells) As a result, the mean air puff latency in tactilely conditioned POm cells is later
than that visually conditioned cells (Figure 4c, p=0.003, Wilcoxon rank-sum test). Thus, conditioning alters
POm activity by changing the proportion of cells that respond later than 100 ms after air puff onset, while
the proportion of cells that respond earlier remain constant.

POm correlates with arousal and movement regardless of conditioning

In freely-whisking mice, POm activity correlates with whisking, pupil radius, and behavioral state[11–13]. To
assess whether this correlation persists in our conditioning task and whether the conditioning type is a
factor, we acquired video of the whiskers and eye contralateral to the recorded thalamus (Figure 5a). We
measured pupil radius as a metric of arousal and quantified whisking activity by *amplitude*, the difference
between the most protracted and retracted position of the whiskers on each whisk cycle (Methods).
Whisking and pupil responses to stimuli were contingent on task type (Figure 5b). Tactilely conditioned
mice displayed a robust increase in pupil radius and whisking amplitude at the onset of the air puff (left).
This movement and arousal change persisted throughout the offset period. Tactilely conditioned mice did
not whisk in response to the drifting grating; however, the grating did induce a pupil constriction as
expected from autonomic response to luminance change. We observed a similar pupil constriction at
stimulus onset in visually conditioned mice, but this was followed by pupil dilation (right). These mice also
displayed increased pupil radius and whisking in response to the unrewarded tactile stimulus, again
suggesting that the whisker air puff is more salient than the drifting grating even when a mouse has been
trained to ignore it.

Consistent with previous results, we found that POm activity was correlated with both pupil and whisking
regardless of conditioning type (Figure 5c). Interestingly, POm neurons were slightly more correlated with
licking after tactile conditioning than after visual conditioning. We have previously reported that POm
correlates with pupil radius most strongly at a time lag of about 1 second[13]. We found that cells were more
strongly correlated with pupil radius at this time lag in visual conditioning than in tactile conditioning. This
difference in correlation might be due to the difference in pupil dynamics in response to the drifting grating
across conditioning types. However, there was no effect of conditioning on POm-pupil correlation at a time
lag of zero.

Much of the stimulus-aligned POm activity observed could conceivably be due to movement and arousal
responses rather than afferent sensory signals. To disentangle these, we fit the firing rate of each cell with
a linear model using pupil radius, whisking amplitude, and lick rate as predictors on a trial-by-trial basis.
We then analyzed the model residuals as a "movement-corrected" stimulus-evoked firing rate (Figure 5d,
e). Compared to the raw firing rates (Figure 3g, h), we found that regressing out movement had little effect
on activity during the stimulus period but resulted in reduced activity in the offset period. Thus, the
apparent tactile and visual responses in POm after behavioral conditioning cannot be explained by
movement or arousal.

We next sought to determine if this state-dependent modulation was a universal trait of all POm cells, or
if only certain cells were coupled to movement and arousal while others responded to stimuli. We again
fit a linear model of each cell's firing rate, but this time included the timing of the drifting grating, air puff,

and water reward as predictors in addition to the three movement and arousal variables. Cells with licking,
pupil radius, and/or whisking coefficients that were significantly different from 0 were classified as
"movement tuned", and cells with either of the two stimuli as significant predictors were classified as
"stimulus tuned." The vast majority of POm cells were tuned to both movement and stimuli: 312/347
(89%) of cells in tactilely conditioned mice and 239/257 (93%) of cells in visually conditioned mice. A subset
of cells were tuned only to movement (33/347 (9.5%) tactile, 16/257 (6.2%) visual), and fewer still were
tuned only to the air puff or drifting grating (2/347 (0.6%) tactile, 2/257 (0.8%) visual). Movement and
arousal relative activity is thus predominant throughout POm, regardless of whether individual cells also
possess stimulus-specific responses.

Modality selectivity varies with anatomical location in POm
We considered the possibility that the differences in stimulus-aligned activity were caused by distinct
regions within POm having distinct stimulus responses. We computed a stimulus selectivity index for each
cell that significantly responded to the air puff and/or the drifting grating based on its activity within the
first second following stimulus presentation (colored points in Figure 6d and 6f). A selectivity index of +1
indicates that a cell responds only to the air puff, an index of -1 indicates that it responds only to the
drifting grating, and an index of 0 indicates that it responds to both stimuli with equal magnitude.
Unresponsive cells (gray points in 3d and 3f) were excluded from this analysis.

All responsive cells in tactilely conditioned mice were tuned to touch (Figure 6a), and we found no pattern
between a cell's anatomical location and its selectivity index (Figure 6b, left). In visually conditioned mice,
POm neurons exhibited a wide variety of selectivity (Figure 6a). In these animals, touch selective cells
clustered in the lateral and dorsal region of POm (Figure 6b, right), a subregion receiving both top-down
inputs from the barrel cortex and ascending input from the brainstem[33,34]. Linear regression of selectivity
index against a cell's conditioning type and anatomical position along each axis (Figure 6c, Methods)
revealed that only medial-lateral position was significantly correlated with selectivity index. Thus, there
appears to be a core location in the lateral most portion of POm that is always whisker sensitive, while
conditioning dictates the selectivity of the rest of POm.

Conditioning also reshapes LP activity
Conceivably, engaging in a visual or tactile task might differentially set the baseline activity of LP and POm
accordingly. We first compared spontaneous firing rates, defined as the mean rate over periods that
contained no licks, stimuli, or rewards for at least two seconds and that lasted at least 6 seconds.
Conditioning type did not statistically change mean firing rates in POm or LP, though POm cells had a higher
baseline firing rate on average (Figure 7a).

Like POm, LP exhibited a heterogeneous set of stimulus responses that was highly dependent on
conditioning (Figure 7b-d). In tactilely conditioned mice, the majority of LP cells responded to both the air
puff and the drifting grating (25 cells, 39%) or the air puff alone (23 cells, 36%), with only 4 cells (6.3%)
responding solely to the drifting grating. At the population level, their firing rates during the air puff
stimulus period were significantly higher than the drifting grating stimulus period (Figure 7e, left). In
visually conditioned mice (Figure 7d) we observed far fewer cells responding only to the air puff (6 cells,
9%), but a similar proportion of cells that responded to the drifting grating or to both stimuli (6 grating

responding cells, 11.9%; 30 cells responding to both stimuli, 44.7%). Similar to what we observed in POm,
the magnitudes of air puff and grating responses were not significantly different (Figure 7e, right; Figure
7g). Indeed, the magnitude of the LP air puff response was significantly larger in tactilely conditioned mice
than in visually conditioned mice (Figure 7g). Thus, tactile conditioning potentiates tactile responses in LP.

We also examined how conditioning affected low-latency stimulus responses in LP (Figure S3). At the
population level, we observed a peak in LP activity approximately 80 ms after visual stimulus onset (Figure
S3a). Interestingly, this early, putatively sensory-evoked activity was apparent only in tactilely conditioned
LP. LP also exhibited low-latency responses to the air puff onset in both conditioning groups, similar to air
puff-evoked activity in POm. Further mirroring POm, conditioning altered LP stimulus response latency.
Visually conditioned LP had far more cells with a drifting grating response latency between 100 and 500
ms compared to tactilely conditioned LP (Figure S3b). As a result, the mean drifting grating response
latency was significantly higher in visually conditioned LP cells compared to tactilely conditioned cells
(Figure S3c; p=0.001, Wilcoxon rank-sum test). There was no effect of conditioning on response latencies
to the air puff.

As LP correlates with movement and arousal to the same degree POm does[13], we again computed a
movement-corrected firing rate by regressing out pupil radius, whisking amplitude, and licking (Figure 7f).
This regression quashed the difference in activity between conditioning types in the offset period only.
Interestingly, we still observed an air puff response in visually conditioned LP even after this regression
(Figure 7f, right), consistent with a strong salience of facial stimuli. We did not observe a relationship
between anatomical location and visual-tactile selectivity in LP (Figure S3); however, our recordings were
concentrated on the anterior-medial portion of LP and thus may not be characteristic of the entire nucleus.
These results demonstrate that, like POm, LP activity is reshaped according to the modality of the attended
stimulus.

DISCUSSION
Here, we investigated whether the secondary sensory thalamic nuclei differentially encode relevant and
distracting stimuli. Our novel head-fixed conditioning paradigm demonstrates that mice can attend to a
visual or tactile stimulus while ignoring a stimulus of the other sensory modality. The second modality
neither enhanced nor suppressed action. Learning dramatically altered activity in POm: after visual
conditioning, a large portion of POm cells responded to the onset of the visual stimulus. POm correlated
with arousal and movement regardless of conditioning type, but movement correlations could not explain
POm stimulus responses. Learning also altered LP activity: LP responses to tactile stimuli were greater after
tactile conditioning, while visual responses were weaker. The predominant response in both POm and LP
was to the conditioned stimulus. Thus, the secondary sensory thalamic nuclei encode the behavioral
relevance of a stimulus, regardless of its modality.

Learning is shaped by interactions of stimulus salience and behavioral relevance
When trained on a traditional, trial-based structure, visually conditioned mice learned that the drifting
grating predicted reward and that the air puff was negatively correlated with reward. Both stimuli modified
action. In our novel design, however, the unrewarded stimulus was decoupled from reward entirely,

making it a true distractor. When trained on this version of the task, visually conditioned mice learned to
completely ignore the air puff. By contrast, tactilely conditioned mice always ignored the unrewarded
drifting grating regardless of task structure. These mice also learned the stimulus-reward association after
fewer sessions. The air puff was thus innately more salient than the drifting grating. This difference in
saliency is in spite of our attempts to minimize the strength of the air puff and to match the drifting grating
in terms of contrast, size, spatial frequency and temporal frequency to typically observed mouse visual
cortex receptive fields. It is thought that mice rely more strongly on their whiskers than vision for many
behaviors, and that all but the subtlest of vibrissa stimuli are salient[35]. However, the difference in saliency
is more likely related to the engagement of personal space: even humans find facial tactile stimuli
inherently more salient than arbitrary distant visual stimuli. Regardless of the mechanism of saliency, mice
can be conditioned to completely ignore a salient stimulus in favor of a more behaviorally relevant one.

POm responds to behaviorally relevant stimuli regardless of modality

POm is only weakly activated by single whisker deflections, typically requiring large deflections of multiple
whiskers to elicit a response[10,11,27]. These sensory responses likely arise from ascending input from
somatosensory brainstem and/or descending input from primary somatosensory cortex[4,36,37]. Consistent
with those studies, we observed POm cells that responded to the onset of the air puff in both conditioning
groups. Tactilely conditioned mice additionally exhibited POm responses to not only to the onset of the
air puff but also its entire seconds-long duration. Sustained POm responses to air puff were weaker and
less common in visually conditioned mice.

Remarkably, visual conditioning sensitized POm to visual stimuli. These visual responses largely resembled
the air puff responses in tactilely conditioned mice, with individual cells' activities tiling the entire duration
of the stimulus. Visual responses in POm are unexpected. Though it receives excitatory input from a wide
array of cortical regions[31,38,39], POm is not known to receive direct input from the retina or visual cortex.
Rather than a sensory response *per se*, visual-evoked activity likely reflects encoding of behavioral state or
attention. Consistent with that idea, many visually responsive cells also responded to the air puff. It could
be that POm cells respond to any increase in attention, be it due to the unconditioned but salient,
"attention grabbing" air puff or the subtler but now behaviorally relevant drifting grating.

There is growing evidence that POm is a heterogeneous region composed of multiple subnuclei defined
by distinct thalamocortical outputs and corticothalamic inputs[19,33,34,40,41]. We found that the effects of
conditioning on POm stimulus responses varied by anatomical location. While visual conditioning induced
visual responses throughout POm, cells in the lateral dorsal region of POm remained more sensitive to the
air puff than the drifting grating. Layer 5a cells in the S1 barrel field preferentially project to this region of
POm[33], and POm cells receiving S1 corticothalamic input also receive convergent projections from the
whisker-sensitive brainstem nucleus SpVi[33]. It is likely that the observed activity in lateral dorsal POm is
driven by true whisker responses in SpVi and S1.

Movement does not explain POm plasticity

A recent study demonstrated that, in a tactile Go-NoGo paradigm, POm activity correlates with a mouse's
decision to lick or withhold licking[19] and similar results have been demonstrated in a detection task using
the forepaw[14]. Decision, response, and reward consumption coincide with movement – including whisking
and licking. Responses in POm might instead be movement signals, though we have previously
demonstrated that POm-whisking correlations cannot be fully explained by sensory reafference nor motor

efference copy from S1, M1, or the superior colliculus[13]. We found here that POm activity correlated with
whisking amplitude, pupil diameter and licking regardless of conditioning.

In both conditioning groups, POm activity was elevated during reward presentation and consumption,
coincident with licking. POm receives somatotopic projections from all of S1, not just the barrel cortex[10,42].
As such, stimulation of the mouth and tongue - as occurs when the mouse licks - could contribute (via S1)
to late phases of POm activity. Nevertheless, tactilely conditioned mice had a greater correlation between
licking and POm activity. This suggests an increased sensitivity to all tactile stimulation as mice attend to
the air puff, compared to mice that are ignoring the air puff and attending to the drifting grating.

Furthermore, visually conditioned mice had a greater correlation between POm and pupil radius, but only
at long time lags (>500 ms). Conceivably, as mice attend to visual stimuli, they could be more sensitive to
all changes in retinal input. However, at time lags close to 0 ms there is no difference in POm-pupil
correlation between tactile and visual conditioning groups, and we observe very few visually responsive
cells in tactilely conditioned mice. Therefore, this activity may instead represent an increased correlation
between POm and arousal. The drifting grating is inherently less salient than the air puff; consequently,
the visual conditioning task may require greater attentional demands, which in turn could further engage
the secondary thalamic nuclei.

As licking, whisking, and pupil dilation are all significantly correlated with POm, we examined whether this
relationship could explain stimulus-evoked activity. Even after we factored out the possible contributions
of these movement variables using regression, POm responses to the onset of conditioned stimuli were
stronger than that of the unconditioned stimuli. In both tactilely and visually conditioned mice, movement
could not explain the increased firing rate at air puff onset. These low-latency responses across
conditioning groups are likely due in part to "true" sensory responses driven by S1 and SpVi. In tactilely
conditioned mice, movement regression attenuated, but did not eliminate, activity in the later portion of
the air puff. Activity in the rewarded offset period, however, was suppressed almost entirely, suggesting
that post-stimulus activity is due to non-sensory signals when mice lick the water port.

LP activity is similarly shaped by conditioning
Like POm, LP displayed varied stimulus-evoked activity that was heavily dependent on conditioning. LP
responded to the air puff robustly and with low latency, despite lacking direct somatosensory inputs.
Though LP activity correlates with whisking and air puffs induce whisking, movement could not explain
such responses in either conditioning group. The air puff response was larger in tactilely conditioned mice,
indicating that, like POm, LP is more responsive to the occurrence of any behaviorally relevant stimuli
regardless of modality.

Unlike POm, LP responded to the drifting grating in both conditioning groups. LP receives input from the
retina, the superior colliculus, and several visual cortical areas[26,29,30,43–45]. In tactilely conditioned mice, 45%
of LP cells responded to the onset of the drifting grating, consistent with previous studies of LP responses
to simple visual stimuli[28]. In visual conditioning animals, a greater portion of cells were visually responsive
(56%), suggesting the existence of a population of LP cells that are not sensitive to simple visual stimuli
but are active when a mouse is aroused and attentive. This is further reflected in the increased latency of
drifting grating responses after visual conditioning.

It is notable that a majority of LP cells had significant responses to the air puff in both conditioning groups.
This is despite the fact that LP has no known inputs from the somatosensory brainstem or the barrel cortex.

However, we have previously seen that LP activity correlates with whisking just as strongly as POm does,
and at a time lags less than 50ms[13], a fact which we attribute to LP tracking global arousal. Further, here
we observed that the air puff is inherently more salient to naive mice than the drifting grating. It is possible
that even after visual conditioning, the air puff remains salient and arousing but the mice have learned
not to respond to it. These responses could be the result of neuromodulators, such as norepinephrine or
acetylcholine, acting either directly on LP or on its inputs[46–51].

LP has several subregions with distinct cytoarchitecture, anatomical inputs, and retinotopic
maps[25,29,30,52,53]. Our recordings were clustered in the anterior and medial portion of LP, which receives
input primarily from frontal cortex and only sparse input from V1 and superior colliculus. Other regions of
LP may receive greater input from visual cortex, and thus be more responsive to visual stimuli and less
affected by conditioning, similar to the dorsal-ventral region of POm we identified. Further
characterization of the various LP subregions in behavioral contexts could reveal how specific cortical and
subcortical inputs contribute to attention.

Possible mechanisms of non-sensory responses
What might be driving these non-sensory responses in the secondary nuclei? One likely mechanism is
neuromodulation. Cortical levels of acetylcholine and norepinephrine track arousal[50]. Both acetylcholine
and norepinephrine can act directly on thalamic nuclei like VPM[47,48] and may do so on POm and LP as well.
Acetylcholine can also enhance POm tactile responses by suppressing GABAergic activity in the zona
incerta[49]. LP receives inhibitory input from the zona incerta[54], so LP visual responses are likely modulated
by cholinergic activity in the same way. Either direct modulation or indirect disinhibition could underlie
enhanced air puff responses in tactilely conditioned POm and enhanced drifting grating responses in
visually conditioned LP. State-based disinhibition of POm could also contribute to "visual" responses after
visual conditioning by potentiating non-sensory inputs to POm.

Another potential culprit is direct corticothalamic input. S1 is thought to be the primary driver of POm[10].
Though S1 activity alone does not drive arousal-related activity in POm, direct S1-to-POm projections
contribute to performance of tactile discrimination tasks[15]. Glutamatergic S1 inputs could underlie
enhanced air puff responses in POm in behavioral contexts; however, S1 activity cannot explain the same
effect in LP. Instead, visual responses in POm and tactile responses in LP might be driven by high-order
cortex. In nonhuman primates, the lateral pulvinar has been shown to facilitate communication between
the visual region V4 and higher order cortical regions. These include the lateral intraparietal area[55] and the
temporo-occipital area[56], regions that are engaged in visual attention tasks. Indeed, the secondary nuclei
have been considered possible hubs for cortico-cortical communication[24,57,58] and are known to be
innervated by posterior parietal cortex. Elevated activity in POm and LP could reflect increased cortico-
cortical communication between high-order regions, primary sensory regions, and motor regions based
on attentional needs.

The role of high-order thalamus in attention
The secondary visual thalamic nucleus in primates – the lateral pulvinar – has been known to play a role
in guided visual attention. In humans, pulvinar damage causes a variety of attentional deficits, impairing
the ability to selectively attend to a visual stimulus in the presence of a distractor[21,22,22]. In nonhuman
primates, chemical silencing of the pulvinar induces spatial neglect[23,24]. Further, pulvinar responses to a
visual stimulus are strongest when a monkey directs attention towards that stimulus[59]. Similar results have
not yet been reported in rodent LP, nor have they been directly extended to the somatosensory thalamus.

A consistent result of attention studies in humans and nonhuman primates is the enhancement of cortical
and thalamic sensory responses to an attended visual stimuli.[24,60,61] Here, we show not just enhancement
of sensory responses to stimuli within a single modality, but also across modalities. It is worth investigating
further how secondary thalamus and high-order sensory cortex encode attention to stimuli outside of
their respective modalities. Our surprising conclusion that the nuclei are equivalently activated by
behaviorally relevant stimuli is nevertheless compatible with these previous studies.

The role of secondary thalamus in learning and cortical plasticity

Experience-dependent synaptic plasticity is a substrate of learning and memory. In anesthetized mice,
POm excitation can facilitate long-term plasticity (LTP) in layer 2/3 pyramidal cells in S1, while POm
silencing blocks those same plasticity effects[17]. Similarly, whisker-dependent associative learning
potentiates POm-to-S1 synapses[16]. In visual cortex, excitation of layer 2/3 neurons paired with stimulation
of layer 4 cells induces LTP in those layer 2/3 cells[62]. LP thus likely contributes to plasticity in visual cortex.
Elevated firing rates in POm and LP could contribute to learning the behavioral relevance of novel stimuli.
That is, they may open a window of plasticity during periods of high arousal and task engagement. As such,
we expect that disruption of thalamocortical output from secondary nuclei to sensory cortex will disrupt
or even prevent associative learning. Based on our findings, disruption of POm activity could impair even
visual learning, and disruption of LP activity tactile learning, particularly if there were a strong multimodal
component in the task. Such future experiments would further illuminate how the secondary nuclei
participate in sensory processing and could pave the way for further studies of thalamocortical and
corticothalamocortical circuits.

Conclusion

Our study demonstrates that a secondary sensory thalamic nucleus is dramatically modulated by
behaviorally relevant events both in its modality as well as other modalities. This broad dependence on
task engagement across secondary nuclei suggests a potential role in gating cortical plasticity and overall
attention, particularly with regards to multimodal associative learning and transthalamic communication
between cortical areas.

METHODS

Animals

Twenty-three wild type C57/BL6 (4 female, 19 male) were used in this study. Mice were either purchased from Jackson Laboratories or bred in our own colony, and were between 12 and 24 weeks old. All experiments were conducted under the supervision and approval of the Columbia University Institutional Animal Care and Use Committee.

Surgery

Mice were administered the global analgesics carprofen (5 mg/kg) and buprenorphine (0.05-0.1 mg/kg) subcutaneously, and anesthetized with isoflurane. The scalp was injected with the local analgesic bupivacaine (5 mg/kg). Mice were then placed in a stereotax, and the scalp and fascia were removed. A custom-designed stainless steel headplate was affixed to the skull with Metabond®. Carprofen was administered subcutaneously 24 hours post-surgery. Mice were allowed to recover for seven days, during which additional carprofen was administered as deemed necessary. After recovery, mice were deprived of water, habituated to the behavioral apparatus, and conditioned (*Behavior Apparatus* and *Conditioning*, below). After behavioral training, mice were again treated with carprofen and buprenorphine, anesthetized with isoflurane, and placed in a stereotax. A ~200 μm-wide opening was cut on the left side of the skull, centered at 1.8 mm posterior to bregma and 1.5 mm lateral of the midline. This opening was sealed with Kwik-Cast™. A ground pin was inserted over the frontal cortex of the right hemisphere. Mice were allowed to recover for 24 hours, after which recordings took place. At the end of experiments, mice were deeply anesthetized with isoflurane, then perfused transcardially with phosphate buffer followed by 4% paraformaldehyde.

Behavior Apparatus

The behavior apparatus was contained in a black box with a light-blocking door. It was constructed from metal posts on an aluminum breadboard (Thorlabs). Mice were held in a 3D-printed hutch and head-fixed with a machined steel head plate holder. Two-second-long air puffs (0.5-1 bar) were delivered through a nozzle (cut p1000 pipet tip, approximately 3.5mm diameter aperture) and controlled by a solenoid. Water rewards (8-16μL) were similarly delivered by a solenoid. Licking was recorded with an infrared proximity detector. The visual stimulus was presented on a monitor (5 inch, 800x480 resolution, Eyoyo) positioned 11 cm from right side of the mouse hutch occupying 52°x 33° of visual space. The stimulus consisted of vertical drifting bars (spatial frequency 0.05°/cycle, temporal frequency 1.5Hz) with maximum contrast (maximum luminance 247 cd/m^2, minimum luminance 3.2 cd/m^2) and lasted two seconds. When the stimulus was off the monitor displayed a gray background set to the same mean luminance as the stimulus (125 cd/m^2). A photodiode was used to detect the onset of the visual stimulus to synchronize it with the rest of the apparatus.

An Arduino UNO microcontroller drove the two solenoids. A digital signal from the Arduino was used to trigger the visual stimulus, which was driven with a custom Python script (PsychoPy®). The solenoid TTL signals, the lick detector analog signal, and the photodiode output were all recorded with OpenEphys hardware and software (see "Electrophysiology" below). When capturing video, the Arduino also drove a blinking infrared LED to synchronize video data.

Habituation and Conditioning

Throughout habituation and conditioning, mice were deprived of water in their home cages and received
all daily water intake while placed in the behavior apparatus. We closely monitored their weight and
health, and *ad libitum* water was provided if weight decreased below 80% of baseline.

Prior to conditioning, mice were habituated to handling and head fixation. They were placed in the
behavior apparatus over several sessions of increasing duration, spanning 10 to 30 minutes, during which
they were given their water ration. Mice were habituated for at least three such sessions over the course
of two to three days. During habituation, mice were observed and evaluated for signs of discomfort or
distress (i.e. squeaking and trying to escape). Mice were considered habituated if they completed a 30-
minute session without displaying any such behaviors, and all mice reached habituation after three days.

Conditioning began 24 hours after mice reached habituation. In the first phase of conditioning ("shaping"),
mice were separated into two cohorts: a "tactile" cohort and a "visual" cohort. Mice were presented with
tactile stimuli (a two-second-long air puff delivered to the right whisker field) and visual stimuli (a two-
second playback of a vertical drifting grating on a monitor, positioned on the right side of the face).
Throughout conditioning, mice were monitored via webcam to ensure that the air puff only contacted the
distal tips of the whiskers and did not disturb the facial fur nor cause the mouse to blink, flinch, or
otherwise react – ensuring the stimulus was not aversive. The stimulus types were randomly ordered. In
the visual conditioning cohort, the visual stimulus was paired with a water reward (8-16µL) delivered at
time of stimulus offset. In the tactile conditioning cohort, the reward was instead paired with the offset of
the air puff. Regardless of the type of conditioning, stimulus type was a balanced 50:50. During the shaping
version of the task, stimuli were delivered with an inter-stimulus interval (ISI) of 8-12 seconds (uniform
distribution). This distribution changed in the full version of the task (see below). Mice were trained for
one session per day, with each session consisting of an equal number of visual stimuli and air puffs.
Sessions ranged from 20-60 minutes and about 40-120 of each stimulus.

Throughout training and during electrophysiology recordings, we recorded analog signals of mouth and
tongue movements with an infrared proximity detector located below the water reward spout. Individual
licks were detected by manually inspecting this signal and setting a threshold using custom MATLAB code.
Throughout conditioning, we evaluated performance by measuring the anticipatory licking behavior of a
mouse to a rewarded stimulus or unrewarded stimulus. For each stimulus presentation, we counted the
number of times a mouse licked in the two seconds prior to the stimulus onset and the number of licks
during the two-second stimulus and calculated a lick index (LI) for that stimulus[32].

522
$$LI = \frac{licks_{stimulus} - licks_{baseline}}{licks_{stimulus} + licks_{baseline}}$$

A positive lick index indicates that a mouse responds to a stimulus by licking more and a negative index
indicates that a mouse responds by withholding licking. Licks that occurred within 50ms of the stimulus
onset were excluded when computing the lick index.

Mice were trained on the shaping version of the task until the mean lick index of the CS+ for a given session
was significantly greater than 0 (sign rank test), with a minimum of three days of conditioning. After
reaching this threshold, mice were switched to the "full" version of the task. In the full task, the stimuli
and reward were identical, but stimuli were presented at uncorrelated and less predictable intervals. For
a given stimulus, the time until the next stimulus of the same type was drawn randomly from an

exponential distribution with a mean of 10 seconds plus an offset of 8-12 seconds (draw from a uniform
distribution), for a mean ISI of 20 seconds. The maximum ISI was capped at 55 seconds, and ISIs were
binned by seconds. Both the air puff and the visual stimulus ISIs were drawn from the same distribution
but were done so independently. Thus, the stimuli could overlap, the timing of the unconditioned stimulus
was uncorrelated with the CS+ or the reward, and each stimulus had a flat hazard-rate for the majority of
possible ISIs. Mice were then trained on the full version of the task until the mean LI per session for the
unconditioned stimulus was not significantly different from 0 (signed-rank test) for a minimum of four
days.

Electrophysiology

We performed simultaneous recordings from LP and POm from fully trained mice as the animal performed
the full version of the task. Using a motorized micromanipulator (Scientifica PatchStar), we inserted a 64-
channel electrode array (Cambridge Neurotech, models H3 and H9) into the craniotomy. Arrays were
inserted vertically to a depth of 3.1-3.5 mm from the cortical surface (mean 3.4 mm). Prior to recording,
the tip of the array was dipped into a DiI solution to label recording sites. Array recordings were acquired
using an OpenEphys amplifier with two digital headstages (Intan) at 30 kHz and with a bandwidth of 2.5
Hz to 7.6 kHz. The same acquisition system was used to record stimulus timing, licking, and a syncing signal
for the video.

We used KiloSort3 to detect spikes and assign them to putative single units (Pachitariu et al., 2016) and
Phy2 (https://github.com/cortex-lab/phy; Rossant, 2021) to manually inspect each unit. We merged units
that appeared to originate from the same cell, based on waveform shape, auto- and cross-correlations,
and firing rate over time. We excluded cells that lacked a refractory period (i.e. a dip in autocorrelation
within 3ms). Specifically, units with more than 10% of spikes having inter-spike intervals shorter than
3msec were considered multi-unit and excluded. Units were assigned to the array channel on which its
mean waveform amplitude was largest.

Histology

We sectioned paraformaldehyde-fixed brains into 100-µm sections using a vibratome. To visualize the
border between POm and VPM, we stained sections for endogenous biotin using fluorescently conjugated
streptavidin (Alexa Fluor 647, ThermoFischer). We imaged the sections on a slide scanner (Nikon AZ100
Multizoom). We then used SHARP-Track to align images to the Allen reference atlas and to identify the
position of the electrode array by tracing the DiI track (Shamash et al 2018, https://github.com/cortex-
lab/allenCCF). After histological alignment (see below) we used the depth of each putative cell along the
array to assign it to a brain region. When assigning cells to brain regions, we excluded cells that were within
50 µm of a region boundary.

Videography

During electrophysiology recordings, two PS3 Eye cameras were used to capture video of the mouse's eye
and whiskers. The whisker camera was positioned below the mouse's head and recorded at 125 fps. The

eye camera was positioned on the right side of the face and recorded at 60 fps. The OpenEphys Bonsai
program was used to record video.

Pupil size was measured from video offline using custom MATLAB code. We defined a region-of-interest
around the eye and applied a threshold to each video frame to create a binary image of the pupil. We then
used the function "regionprops" to fit an ellipse to the pupil and estimated pupil radius as the geometric
mean of the semi-major and semi-minor axes of that ellipse. The trace of pupil radius over time was
smoothed over 5 frames (83 ms). Large discontinuities in the pupil radius vector (such as those caused by
a blink) were excluded based on a median filter (spanning 3 seconds) and linearly interpolated.

Whiskers were automatically tracked from videos using the software package *Trace* (Clack et al., 2012).
Custom MATLAB software was used to compute the mean angle of all whiskers over each frame. The mean
angle was bandpass filtered from 4 to 30 Hz and passed through a Hilbert transform to calculate phase.
We defined the upper and lower envelopes of the unfiltered median whisking angle as the points in the
whisk cycle where phase equaled 0 (most protracted) or ±π (most retracted), respectively. Whisking
amplitude was defined as the difference between these two envelopes.

A blinking LED paired with a TTL pulse was used to synchronize the video frames to the data collected via
the OpenEphys acquisition system. After processing the videos, this signal was used to align relevant video
data (pupil radius and whisking amplitude, e.g.) to physiology and behavioral data. Video vectors were
linearly interpolated from their respective frame rates to 1 kHz. Stimulus and licking signals were
downsampled to 1 kHz, and spike times were rounded to the nearest millisecond.

Stimulus Responses

We examined each cell's response to the two task stimuli by comparing its firing rate during a "stimulus
period" to its baseline firing rate. We first excluded periods in the recording with overlapping stimuli,
defined as any stimulus occurring within 6 seconds of a stimulus of a different type. We then counted the
number of spikes that occurred within 1 second prior to the onset of each stimulus (baseline period) and
within one second of the stimulus onset (stimulus period). We classified a cell's response as significant by
first performing an ANOVA between the baseline firing rates, the rates during the air puff stimulus period,
and the rates during the drifting grating stimulus period over each stimulus presentation. This was followed
by a Holm-Bonferroni multiple comparisons correction, which was applied to cells within each
experimental group (*i.e.*, p values from tactile conditioning POm cells were corrected separately from p
values of visual conditioning LP cells). We then classified cells as responding to the visual stimulus, the air
puff, or both by performing paired Wilcoxon ranked-sum tests between stimulus period firing rates and
the corresponding baseline firing rates. No multiple-comparisons correction was applied to the ranked-
sum values.

For significantly responding cells, we computed a stimulus selectivity index (SI). The selectivity index
measures the difference between the magnitudes of the air puff response and the drifting grating
responses compared to the baseline firing rate.

607
$$SI = \frac{|FR_A - FR_B| - |FR_V - FR_B|}{|FR_A - FR_B| + |FR_V - FR_B|}$$

Where FR$_B$ is the mean baseline firing rate, FR$_A$ is the mean firing rate during the air puff stimulus period
(a one-second period after stimulus onset) and FR$_V$ is the mean firing rate during the visual stimulus period.
A positive selectivity index indicates that the cell has a larger response to an air puff, in terms of absolute
change in firing rate, than it does to the visual stimulus, and vice-versa for a negative index.

To determine if anatomical location had a significant effect on selectivity index, we fit a linear model
between the selectivity index of all POm cells and each cell's position in the dorsal-ventral (DV), medial-
lateral (ML), and anterior-posterior (AP) position, along with the conditioning type (CT) of that cell. We
also considered interactions between experiment type and position, but not between the different
positional dimensions. The model thus had the following design:

$$SI \sim 1, DV, ML, AP, CT, CT*DV, CT*ML, CT*AP$$

To measure sensory response latency, we calculated the firing rate of each cell over 10ms bins. We
measured the mean firing rate of each cell aligned to either the air puff or the visual stimulus. We then
computed the mean and standard deviation of the cell's firing rate one second prior to the stimulus
onset (baseline firing rate). We defined a sensory response as the first instance where a cell's activity
increased or decreased to cross a 99% confidence interval (2.576 standard deviations) of the baseline
firing rate for at least two consecutive time bins, or the first time a cell's activity crossed a 99.99%
confidence interval (3.891 standard deviations) for a single time bin, whichever occurred first. Baseline
firing rates were defined as the average rate over two seconds prior to stimulus onset.

Regression of Movement and Arousal Variables
When fitting linear models, we first binned firing rates, lick rates, whisking, and pupil radius into 250ms
time bins. This bin size is long enough to capture a range of firing rates (i.e., most cells had multiple spikes
per bin) while preserving the timing of licking and whisking. We tested models with bin sizes of 50, 100,
and 500 ms and found qualitatively similar results. For each cell, we fit a linear model of firing rate to
whisking amplitude, pupil radius, and lick rate using least-squares regression. We extracted the model
residuals for each cell, which correspond to a baseline-subtracted and movement-corrected firing rate.

ACKNOWLEDGMENTS
The authors thank Dahee Jang, Katy Willard, Candice Lee, Ziyue Liu, and Armin Lak for comments on the
manuscript; Chris Rodgers and Sam Benezra for advice on creating the behavioral apparatus; and David
Park for help training mice in pilot experiments. Support was provided by a Wellcome Trust Discovery
Award (225210/Z/22/Z), an Academy of Medical Sciences Professorship, NIH/NINDS R01 NS069679 and
R01 NS094659, and NIH/NEI T32 EY013933.

Figure 1: Mice can be conditioned to attend to one sensory modality while ignoring a second modality.

a, Behavioral setup. Mice were head-fixed and presented with a visual stimulus (a drifting grating on a screen) or a tactile stimulus (an air puff to the distal portion of the whiskers). **b**, Schematic for the shaping version of the task, depicting the timing of the air puff (red), the drifting grating (dark blue), and the water reward (light blue). Mice were presented with either stimulus with an inter-stimulus-interval of 8-12 seconds. In the tactile cohort, the air puff was paired with a water reward; in the visual cohort the drifting grating was paired with the water reward. **c**, Schematic for the full version of the task. Stimuli were allowed to overlap so that only the conditioned stimulus predicted the timing of the water reward. **d**, Timing of stimuli in the full task. Inter-stimulus intervals were drawn from an exponential distribution with a linear offset (top). As a result, the hazard rate was largely flat for most stimulus presentations (bottom). **e**, Example behavior from a mouse trained on the visual task. Top row: raster plot of licks aligned to the onset of the grating (blue region). Red markers indicate the first lick to occur after reward was available. Middle row: raster plot of licks aligned to the onset of the air puff (red region). Bottom row: Mean lick rate aligned to the onset of the drifting grating (blue) or the air puff (red). Gray region indicates the stimulation period. Data are from the first session of the shaping task (left column), the first day of the full task (middle column), and the final day of conditioning (right column). **f**, Learning curve of the mouse shown in E, showing the lick index (mean +/- SEM) in response to the drifting grating (blue) or the air puff (red) on each training day. Vertical dotted line indicates the day the mouse was switched from the shaping version to the full version of conditioning. Triangles: sessions shown in e. **g, h**, Similar to e and f but for an example mouse trained on the tactile task. **i,** Learning curves of all visually conditioned mice (left) and tactilely conditioned mice (right). Thin lines: mean lick index of a single mouse. Thick lines: mean lick index across all mice. **j**, Mean and SEM lick index for all mice on the first day of conditioning ("shaping day 1"), the first day of the full task ("full day 1"), the day before neural recordings ("penultimate"), and the day of recording ("recording"). There was no significant difference in lick index to the conditioned stimulus between the penultimate day and the day of recording in either conditioning group (visual conditioning p=0.18, n=12 mince, Wilcoxon signed-rank test; tactile conditioning p=0.08, n=11 mice, Wilcoxon signed-rank test).

Figure 2: Array recordings were made simultaneously from POm and LP.

a, Example coronal section. Red: DiI marking recording location. Gray: endogenous biotin stained with Alexa fluor 647 streptavidin. **b,** Zoom of the outlined region in a. The borders of LP, POm, and VPM are outlined. A 64-channel silicon probe was used to record from LP and POm. **c**, Mean waveform of an example POm cell across all probe channels. The channel with the largest mean waveform was identified and used to estimate the position of the cell (red circle in B). **d, e, f**, Estimated location of putative cells (gray circles) located in POm (red) and LP (blue) aligned to the Allen reference atlas. Cell locations are jittered along the medial-lateral and anterior-posterior axes for visualization purposes. **d**, Coronal projection over the range of cell locations in the anterior-posterior axis. **e**, Sagittal projection over range of cell locations in the medial-lateral axis. **f**, Transverse projection over range of cell locations in the dorsal-ventral axis.

Figure 3: POm responses to tactile and visual stimuli depend on conditioning paradigm.

a, Example cells from a tactilely conditioned mouse (same mouse as in figure 1g, session 7). Top row: spike
raster aligned to the onset of the drifting grating (blue). Middle row: spike raster aligned to the air puff
(red region). Bottom row: Mean firing rate over each trial aligned to the drifting grating (blue line) and the
air puff (red line). Gray region indicates the timing of both stimuli. **b,** Sample cells from a visually
conditioned mouse (same mouse as in figure 1e, session 12). **c**, Firing rate of all POm cells from tactilely
conditioned mice aligned to either the air puff (left) or the drifting grating (right). Firing rates are binned
at 250ms and cells are sorted by the timing of the peak firing rate when aligned to the air puff. n=347 cells,
11 mice, range=11-57 cells/mouse, median=33 cells/mouse. **d**, Scatter plot of POm firing rates 1-second
pre-stimulus (x-axis) and 1 second during stimulus onset (y-axis) from tactilely conditioned mice. Colored
markers: cells with significantly different firing rates pre- and post-stimulus (1-way ANOVA with Holm-
Bonferroni correction and post-hoc Wilcoxon signed-rank test). Gray markers: cells without significantly
different firing rates. **e**, Firing rates of all POm cells from visually conditioned mice aligned to either the air
puff (left) or the drifting grating (right). Cells are sorted by the timing of the peak firing rate aligned to the
grating. (n=257 cells, 11 mice, range=10-42 cells/mouse, median=19 cells/mouse). **f**, Scatter plot of POm
firing rates pre- and post-stimulus onset for visually conditioned mice, as in e. **g**, Event-triggered average
firing rates of all POm cells (mean +/-SEM) aligned to either the air puff (red) or the drifting grating (blue)
in tactilely conditioned mice (left) and visually conditioned mice (right). Gray region indicates the timing
of both stimuli. Rates binned at 250ms. **h**, Box plots of the change in firing rate from baseline for each
POm cell during the first second of stimulus onset ("stimulus period") and during the first two seconds
post-stimulus ("offset period"). In tactilely conditioned mice, the change from baseline was greater in the
air puff stimulus period than the visual stimulus period (two-way ANOVA with post-hoc signed-rank test;
conditioning type F=1.98, p=0.16; stimulus type F= 153, $p<10^{-33}$; interaction F=115, $p<10^{-26}$, signed-rank
$p<10^{-45}$). There was no difference in the change from baseline between stimulus types in visually
conditioned mice (p=0.42, signed-rank test). In visually conditioned mice, the drifting grating stimulus
response was significantly greater than that of tactilely conditioned mice (rank-sum test, $p<10^{-42}$). In both
conditioning types, firing rates in the offset period were significantly different between stimulus types
(two-way ANOVA, conditioning F=0.01, p=0.90; stimulus F=16, $p<10^{-5}$; interaction F=312, $p<10^{-62}$; signed-
rank $p<10^{-20}$ for both visually and tactilely conditioned mice).

Figure 4: Conditioning alters tactile stimulus response latencies in POm.

a, Event-triggered average firing rates of all POm cells (mean +/-SEM) aligned to either the air puff (red)
or the drifting grating (blue) in tactilely conditioned mice (left) and visually conditioned mice (right), as in
3g. Dashed line indicates stimulus onset. Firing rates are binned at 10ms. Note the truncated Y axis. **b,**
Histogram of the stimulus response latency, as a portion of total recorded POm cells for each
conditioning group. Stimulus response latency was defined as the first time a cell's firing rate crossed a
confidence interval threshold compared to baseline (Methods). Only cells with response latencies less
than 500ms are represented. Red: stimulus response latency relative to the air puff onset. Blue: response
latency relative to the drifting grating onset. **d,** Box plot of the air puff response latencies of POm cells
less than 500ms. Response latencies in tactilely conditioned mice were later than those in visually
conditioned mice (p=0.003, Wilcoxon rank-sum test, n=226 cells from tactilely conditioned mice, 101
cells from visually conditioned mice).

Figure 5: POm correlates with arousal and movement regardless of conditioning.

a, Pupil and whisker tracking. Top left: Example frame from an eye video with the edge of the pupil highlighted in green. Top Right: A sample trace of pupil radius over time (gray: raw; green: smoothed). Bottom Left: Example frame from a whisker video with whiskers highlighted in green. Bottom Right: Example trace of whisker angle over time (gray) and the most protracted and retracted whisker positions over each whisk cycle (green), which is used to compute whisking amplitude. **b**, Whisking amplitude and pupil radius aligned to stimuli in tactilely conditioned mice (n=11 mice, left column) and visually conditioned mice (n=11 mice, right column). Top row: Pupil radius of each mouse aligned to either the air puff (blue) or the drifting grating (red) (Z-scored, mean+/-SEM). Bottom row: Whisking amplitude of each mouse aligned to the two stimuli (mean+/-SEM). Gray regions indicate stimulus timing. **c**, Cross-correlations between POm activity and pupil radius (top), whisking amplitude (middle), and lick rate (bottom). Light gray: Cells from tactilely conditioned mice (mean +/-SEM, n=347 cells, 11 mice). Dark gray: Cells from visually conditioned mice (n=257, 11 mice). Negative lags indicate that POm activity precedes the movement variable and positive lag indicates that POm activity follows the movement variable. **d**, Movement-corrected firing rates. The firing rate of each cell was fit to a linear model with whisking, licking, pupil radius, and baseline rate as predictors. Model residuals of all cells (mean+/-SEM) are aligned to the drifting grating and the air puff, as in B. **e**, Box plot of the model residuals of each POm cell during the first second of stimulus onset ("stimulus period"), and during the first two seconds post-stimulus ("offset period"), the same time windows as in 3g and 3h. Adjusted firing rates in tactilely conditioned mice were significantly higher during the air puff stimulus period than the drifting grating period. (Two-way ANOVA with post-hoc Wilcoxon signed-rank test. Conditioning type F=4.23, p=0.04; stimulus type F= 68.2, p<0.001; interaction F=38.5, p<0.001; tactile conditioning Wilcoxon p<0.001; visual conditioning Wilcoxon p=0.09). In visually conditioned mice, adjusted firing rates were higher during the drifting grating offset period than the air puff offset period. (conditioning F=0.003, p=0.95; stimulus F=3.9, p=0.048; interaction F=9.45, p=0.002; tactile conditioning Wilcoxon p=0.94, visual conditioning p<0.001).

Figure 6: POm responses to tactile and visual stimuli vary based on anatomical location.

a, Sensory selectivity index for significantly responding POm cells (colored cells in 3d and 3f). Cells with a positive selectivity index primarily respond to the air puff and cells with a negative index primarily respond to the drifting grating. Cells from tactilely conditioned mice had a significantly higher selectivity index (Wilcoxon rank sum test, p<0.001, n=200 tactile conditioning cells, 120 visual conditioning cells). **b**, Position of all POm cells separated by conditioning type. Cell positions are projected on the coronal plane (top row) and sagittal plane (bottom row), as in figure 2d and 2e. Significantly responding cells are colored by their selectivity index. Gray Xs indicate unresponsive cells. Positions in the medial-lateral and anterior-posterior axes are jittered for visualization only. **c**, selectivity index of significantly responding cells as a function of their position in POm along the anterior-posterior (left), dorsal-ventral (middle), and medial-lateral axes (right). Light gray: tactile conditioning. Dark gray: visual conditioning. A linear model of selectivity index with experiment type and anatomical position as predictors revealed a significant effect of medial-lateral position and an interaction between position and experiment type. In visually conditioned mice, cells in the lateral portion of POm were more selective to the air puff (model p<10^{-60}, adjusted R^2=0.61, conditioning p<10^{-5}, medial-lateral p<10^{-4}, ML-conditioning interaction p<10^{-4}, all other terms p>0.1)

Figure 7: Conditioning also reshapes LP activity.

a, Baseline firing rates of all POm and LP cells separated by conditioning type. POm cells had higher firing
rates than LP cells, and firing rates did not significantly differ by conditioning type (two-way ANOVA, region
F=8.0, p=0.0049; conditioning type F=0.44, p=0.51; interaction F=1.44, p=0.23). **b**, Example LP cells from a
tactilely conditioned mouse (top) and a visually conditioned mouse (bottom). Mean firing rates over each
trial are aligned to the drifting grating (blue line) and the air puff (red line). Gray region indicates the timing
of both stimuli. **c**, Firing rate of all LP cells from tactilely conditioned mice aligned to either the air puff
(left) or the drifting grating (right). Cells are sorted by the timing of the peak firing rate when aligned to
the air puff. n=64 cells, 7 mice, range=4-16 cells/mouse, median=7 cells/mouse. **d**, Firing rate of all LP cells
from visually conditioned mice aligned to either the air puff (left) or the drifting grating (right). Cells are
sorted by the timing of the peak firing rate when aligned to the drifting grating. n=67 cells, 8 mice, range=1-
19 cells/mouse, median=6 cells/mouse. **e**, Event-triggered average firing rates of all LP cells (mean +/SEM)
aligned to either the air puff (red) or the drifting grating (blue) in tactilely conditioned mice (left) and
visually conditioned mice (right). Gray region indicates the timing of both stimuli. **f**, Movement-corrected
firing rates. The firing rate of each cell was fit to a linear model with whisking, licking, pupil radius, and
baseline firing rate as predictors. Model residuals of all cells (mean+/-SEM) are aligned to the drifting
grating and the air puff, as in E. **g**, Box plots of the change in firing rate from baseline for each LP cell during
the first second of stimulus onset ("stimulus" period) and during the first two seconds post-stimulus
("offset" period). In tactilely conditioned mice, the change from baseline was greater in the air puff
stimulus period than the visual stimulus period (two-way ANOVA with post-hoc signed-rank test;
conditioning type F=1.268, p=0.13; stimulus type F= 15.7, p<=0.001; interaction F=13.25, p=0.003, signed-
rank p<10^{-4}). There was no difference in the change from baseline between stimulus types in visually
conditioned mice (p=0.91, signed-rank test). The change from baseline in during the air puff stimulus
period was also significantly greater in tactilely conditioned LP cells than in visually conditioned cells
(p=0.002, rank-sum test). In both conditioning types, firing rates in the offset period were significantly
different between stimulus types (two-way ANOVA, conditioning F=2.1, p=0.15; stimulus F=0.15, p=0.69;
interaction F=19.5, p<10^{-4}; signed-rank p<10^{-4} in tactile conditioning, p=0.007 in visual conditioning).

Figure S1: Latency of anticipatory licking does not depend on the modality of the conditioned stimulus.

a, Latency to the first lick that occurred after the presentation of the air puff in tactilely conditioned mice
on the day of recording. Box plots depict the distribution of lick latencies of individual mice for all
presentations of the air puff. Only licks that occurred prior to the delivery of the water reward (t≤2s)
were counted. **b,** Latency to the first lick after the presentation of the drifting grating in visually
conditioned mice on the day of recording. **c,** Box plots of the mean lick latency for each mouse as a
function of experiment type. Visually and tactilely conditioned mice did not have significantly different
lick latencies (tactile conditioning median latency=1.1s, n=11 mice; visual conditioning median
latency=1.0s, n=12 mice; p=0.98, Wilcoxon rank sum test).

Figure S2: Manual curation of POm and LP cells

a, b, Example POm cells. *Left*, Mean spike waveform (black) with 200 randomly sampled individual
waveforms (gray). *Right*, Autocorrelogram, binned at 0.5ms. Gray lines indicate a time lag of +/-1ms. **c, d**,
Example LP cells, as in a and b. **e**, Mean waveforms of 50 randomly selected POm cells (red) and 50
randomly selected LP cells (blue). **f**, Histogram of waveform half-widths of each recorded POm cell (red,
n=604) and LP cell (blue, n=131). **g**, Box chart of waveform half-width. Half-widths were not significantly
different between LP and POm cells (POm: 0.40+/-0.07ms, LP: 0.41+/-0.08ms (mean+/-std), t-test

p=0.052). **h**, Histogram of waveform amplitudes from each recorded POm and LP cell. **i**, Box chart of
waveform amplitudes. POm cells had significantly higher amplitudes than LP cells (POm: 0.91+/-0.47mV,
LP: 0.67+/-0.33mV (mean+/-std), t-test p<0.001).

Figure S3: Conditioning alters stimulus response latencies in LP
a, Event-triggered average firing rates of all LP cells (mean +/-SEM) aligned to either the air puff (red) or
the drifting grating (blue) in tactilely conditioned mice (left) and visually conditioned mice (right), as in
7e. Dashed line indicates stimulus onset. Firing rates are binned at 10ms. **b,** Histogram of the stimulus
response latency, as a portion of total recorded LP cells for each conditioning group. Stimulus response
latency was defined as the first time a cell's firing rate crossed a confidence interval threshold compared
to baseline (Methods). Only cells with response latencies less than 500ms are represented. Red:
response latency relative to the air puff onset. Blue: response latency relative to the drifting grating
onset. **c,** Box plot of LP response latencies to the air puff (red) and the drifting grating (blue) in tactilely
conditioned mice (unshaded) and visually conditioned (shaded) mice. Drifting grating response latencies
were significantly different between conditioning groups (Drifting grating p=0.001, Wilcoxon rank-sum
test, n=32 tactilely conditioned cells, 37, visually conditioned cells. Air puff p=0.29, n=52 tactilely
conditioned cells, 46 visually conditioned cells).

Figure S4: LP responses to tactile and visual stimuli as a function of anatomical location
a, Stimulus selectivity index for significantly responding LP cells. Cells with a positive selectivity index
primarily respond to the air puff and cells with a negative index primarily respond to the drifting grating.
Cells from tactilely conditioned mice had a significantly higher selectivity index (Wilcoxon rank sum test,
p<0.001, n=50 tactile conditioning cells, 41 visual conditioning cells). **b,** Position of all LP cells separated
by conditioning type. Cell positions are projected on the coronal plane (top row) and sagittal plane
(bottom row), as in figure 2d and 2e. Significantly responding cells are colored by their selectivity index.
Gray Xs indicate unresponsive cells. Positions in the medial-lateral and anterior-posterior axes are
jittered for visualization only.

REFERENCES
1. Guillery, R.W., and Sherman, S.M. (2002). Thalamic Relay Functions and Their Role in Corticocortical
Communication: Generalizations from the Visual System. Neuron *33*, 163–175.
https://doi.org/10.1016/S0896-6273(01)00582-7.

2. Herkenham, M. (1980). Laminar organization of thalamic projections to the rat neocortex. Science
207, 532–535. https://doi.org/10.1126/science.7352263.

3. Phillips, J.W., Schulmann, A., Hara, E., Winnubst, J., Liu, C., Valakh, V., Wang, L., Shields, B.C., Korff,
W., Chandrashekar, J., et al. (2019). A repeated molecular architecture across thalamic pathways.
Nature Neuroscience *22*, 1925–1935. https://doi.org/10.1038/s41593-019-0483-3.

4. Chiaia, N.L., Rhoades, R.W., Bennett-Clarke, C.A., Fish, S.E., and Killackey, H.P. (1991). Thalamic
processing of vibrissal information in the rat. I. Afferent input to the medial ventral posterior and
posterior nuclei. Journal of Comparative Neurology *314*, 201–216.
https://doi.org/10.1002/cne.903140202.

5. Constantinople, C.M., and Bruno, R.M. (2013). Deep cortical layers are activated directly by
thalamus. Science *340*, 1591–1594. https://doi.org/10.1126/science.1236425.

6. Sherman, S.M., and Guillery, R.W. (2002). The role of the thalamus in the flow of information to the
cortex. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences *357*,
1695–1708. https://doi.org/10.1098/rstb.2002.1161.

7. Wimmer, V.C., Bruno, R.M., de Kock, C.P.J., Kuner, T., and Sakmann, B. (2010). Dimensions of a
Projection Column and Architecture of VPM and POm Axons in Rat Vibrissal Cortex. Cerebral Cortex
20, 2265–2276. https://doi.org/10.1093/cercor/bhq068.

8. Deschênes, M., Takatoh, J., Kurnikova, A., Moore, J.D., Demers, M., Elbaz, M., Furuta, T., Wang, F.,
and Kleinfeld, D. (2016). Inhibition, Not Excitation, Drives Rhythmic Whisking. Neuron *90*, 374–387.
https://doi.org/10.1016/j.neuron.2016.03.007.

9. Petersen, C.C.H. (2007). The functional organization of the barrel cortex. Neuron *56*, 339–355.
https://doi.org/10.1016/j.neuron.2007.09.017.

10. Diamond, M.E., Armstrong-James, M., Ebner, F.F., Armstrong-James, M., and Ebner, F.F. (1992).
Somatic sensory responses in the rostral sector of the posterior group (POm) and in the ventral
posterior medial nucleus (VPM) of the rat thalamus. Journal of Comparative Neurology *318*, 462–
476. https://doi.org/10.1002/cne.903180410.

11. Moore, J.D., Mercer Lindsay, N., Deschênes, M., and Kleinfeld, D. (2015). Vibrissa Self-Motion and
Touch Are Reliably Encoded along the Same Somatosensory Pathway from Brainstem through
Thalamus. PLoS biology *13*, e1002253. https://doi.org/10.1371/journal.pbio.1002253.

12. Urbain, N., Salin, P.A., Libourel, P.A., Comte, J.C., Gentet, L.J., and Petersen, C.C.H. (2015). Whisking-
Related Changes in Neuronal Firing and Membrane Potential Dynamics in the Somatosensory
Thalamus of Awake Mice. Cell Reports *13*, 647–656. https://doi.org/10.1016/j.celrep.2015.09.029.

13. Petty, G.H., Kinnischtzke, A.K., Hong, Y.K., and Bruno, R.M. (2021). Effects of arousal and movement
on secondary somatosensory and visual thalamus. eLife *10*, e67611.
https://doi.org/10.7554/eLife.67611.

14. La Terra, D., Bjerre, A.-S., Rosier, M., Masuda, R., Ryan, T.J., and Palmer, L.M. (2022). The role of
higher-order thalamus during learning and correct performance in goal-directed behavior. eLife *11*,
1–21. https://doi.org/10.7554/elife.77177.

15. Qi, J., Ye, C., Naskar, S., Inácio, A.R., and Lee, S. (2022). Posteromedial thalamic nucleus activity
significantly contributes to perceptual discrimination. PLOS Biology *20*, e3001896.
https://doi.org/10.1371/journal.pbio.3001896.

16. Audette, N.J., Bernhard, S.M., Ray, A., Stewart, L.T., and Barth, A.L. (2019). Rapid Plasticity of Higher-
Order Thalamocortical Inputs during Sensory Learning. Neuron *103*, 277-291.e4.
https://doi.org/10.1016/J.NEURON.2019.04.037.

17. Gambino, F., Pagès, S., Kehayas, V., Baptista, D., Tatti, R., Carleton, A., and Holtmaat, A. (2014).
Sensory-evoked LTP driven by dendritic plateau potentials in vivo. Nature *515*, 116–119.
https://doi.org/10.1038/nature13664.

18. Williams, L.E., and Holtmaat, A. (2019). Higher-Order Thalamocortical Inputs Gate Synaptic Long-
Term Potentiation via Disinhibition. Neuron *101*, 91-102.e4.
https://doi.org/10.1016/j.neuron.2018.10.049.

19. El-Boustani, S., Sermet, B.S., Foustoukos, G., Oram, T.B., Yizhar, O., and Petersen, C.C.H. (2020).
Anatomically and functionally distinct thalamocortical inputs to primary and secondary mouse
whisker somatosensory cortices. Nature Communications *11*, 1–12. https://doi.org/10.1038/s41467-
020-17087-7.

20. Danziger, S., Ward, R., Owen, V., and Rafal, R. (2004). Contributions of the human pulvinar to linking
vision and action. Cognitive, Affective, & Behavioral Neuroscience *4*, 89–99.
https://doi.org/10.3758/CABN.4.1.89.

21. Snow, J.C., Allen, H.A., Rafal, R.D., and Humphreys, G.W. (2009). Impaired attentional selection
following lesions to human pulvinar: evidence for homology between human and monkey.
Proceedings of the National Academy of Sciences of the United States of America *106*, 4054–4059.
https://doi.org/10.1073/pnas.0810086106.

22. Ward, R., Danziger, S., Owen, V., and Rafal, R. (2002). Deficits in spatial coding and feature binding
following damage to spatiotopic maps in the human pulvinar. Nature Neuroscience.
https://doi.org/10.1038/nn794.

23. Wilke, M., Turchi, J., Smith, K., Mishkin, M., and Leopold, D.A. (2010). Pulvinar inactivation disrupts
selection of movement plans. The Journal of neuroscience : the official journal of the Society for
Neuroscience *30*, 8650–8659. https://doi.org/10.1523/JNEUROSCI.0953-10.2010.

24. Zhou, H., Schafer, R.J., and Desimone, R. (2016). Pulvinar-Cortex Interactions in Vision and Attention.
Neuron *89*, 209–220. https://doi.org/10.1016/j.neuron.2015.11.034.

25. Nakamura, H., Hioki, H., Furuta, T., and Kaneko, T. (2015). Different cortical projections from three
subdivisions of the rat lateral posterior thalamic nucleus: A single-neuron tracing study with viral
vectors. European Journal of Neuroscience *41*, 1294–1310. https://doi.org/10.1111/ejn.12882.

26. Roth, M.M., Dahmen, J.C., Muir, D.R., Imhof, F., Martini, F.J., and Hofer, S.B. (2016). Thalamic nuclei
convey diverse contextual information to layer 1 of visual cortex. Nature Neuroscience *19*, 299–307.
https://doi.org/10.1038/nn.4197.

27. Ahissar, E., Sosnik, R., and Haidarliu, S. (2000). Transformation from temporal to rate coding in a
somatosensory thalamocortical pathway. Nature *406*, 302–306. https://doi.org/10.1038/35018568.

28. Allen, A.E., Procyk, C.A., Howarth, M., Walmsley, L., and Brown, T.M. (2016). Visual input to the
mouse lateral posterior and posterior thalamic nuclei: photoreceptive origins and retinotopic order.
The Journal of Physiology *594*, 1911–1929. https://doi.org/10.1113/JP271707.

29. Bennett, C., Gale, S.D., Garrett, M.E., Newton, M.L., Callaway, E.M., Murphy, G.J., and Olsen, S.R.
(2019). Higher-Order Thalamic Circuits Channel Parallel Streams of Visual Information in Mice.
Neuron *102*, 477-492.e5. https://doi.org/10.1016/j.neuron.2019.02.010.

30. Leow, Y.N., Zhou, B., Sullivan, H.A., Barlowe, A.R., Wickersham, I.R., and Sur, M. (2022). Brain-wide
mapping of inputs to the mouse lateral posterior (LP/Pulvinar) thalamus–anterior cingulate cortex
network. J Comp Neurol *530*, 1992–2013. https://doi.org/10.1002/cne.25317.

31. Liao, C.-C., Chen, R.-F., Lai, W.-S., Lin, R.C.S., and Yen, C.-T. (2010). Distribution of large terminal inputs
from the primary and secondary somatosensory cortices to the dorsal thalamus in the rodent.
Journal of Comparative Neurology *518*, 2592–2611. https://doi.org/10.1002/cne.22354.

32. Jurjut, O., Georgieva, P., Busse, L., and Katzner, S. (2017). Learning enhances sensory processing in
mouse V1 before improving behavior. Journal of Neuroscience *37*, 6460–6474.
https://doi.org/10.1523/JNEUROSCI.3485-16.2017.

33. Sumser, A., Mease, R.A., Sakmann, B., and Groh, A. (2017). Organization and somatotopy of
corticothalamic projections from L5B in mouse barrel cortex. Proceedings of the National Academy
of Sciences *114*, 8853–8858. https://doi.org/10.1073/PNAS.1704302114.

34. Groh, A., Bokor, H., Mease, R.A., Plattner, V.M., Hangya, B., Stroh, A., Deschenes, M., and Acsády, L.
(2014). Convergence of Cortical and Sensory Driver Inputs on Single Thalamocortical Cells. Cerebral
Cortex *24*, 3167–3179. https://doi.org/10.1093/cercor/bht173.

35. Warren, R.A., Zhang, Q., Hoffman, J.R., Li, E.Y., Hong, Y.K., Bruno, R.M., and Sawtell, N.B. (2021). A
rapid whisker-based decision underlying skilled locomotion in mice. eLife *10*, e63596.
https://doi.org/10.7554/eLife.63596.

36. Deschênes, M., Veinante, P., and Zhang, Z.-W. (1998). The organization of corticothalamic
projections: reciprocity versus parity. Brain Research Reviews *28*, 286–308.
https://doi.org/10.1016/S0165-0173(98)00017-4.

37. Veinante, P., Jacquin, M.F., and Deschênes, M. (2000). Thalamic projections from the whisker-
sensitive regions of the spinal trigeminal complex in the rat. Journal of Comparative Neurology *420*,
233–243. https://doi.org/10.1002/(SICI)1096-9861(20000501)420:2<233::AID-CNE6>3.0.CO;2-T.

38. Aldes, L.D. (1988). Thalamic connectivity of rat somatic motor cortex. Brain Research Bulletin *20*,
333–348. https://doi.org/10.1016/0361-9230(88)90063-9.

39. Gharaei, S., Honnuraiah, S., Arabzadeh, E., and Stuart, G.J. (2020). Superior colliculus modulates
cortical coding of somatosensory information. Nature communications *11*, 1693.
https://doi.org/10.1038/s41467-020-15443-1.

40. Mease, R.A., Metz, M., and Groh, A. (2016). Cortical Sensory Responses Are Enhanced by the Higher-
Order Thalamus. Cell Reports *14*, 208–215. https://doi.org/10.1016/j.celrep.2015.12.026.

41. Viaene, A.N., Petrof, I., and Sherman, S.M. (2011). Properties of the thalamic projection from the
posterior medial nucleus to primary and secondary somatosensory cortices in the mouse.
Proceedings of the National Academy of Sciences *108*, 18156–18161.
https://doi.org/10.1073/pnas.1114828108.

42. Fabri, M., and Burton, H. (1991). Topography of connections between primary somatosensory cortex
and posterior complex in rat: a multiple fluorescent tracer study. Brain Research *538*, 351–357.
https://doi.org/10.1016/0006-8993(91)90455-5.

43. Born, G., Schneider-Soupiadis, F.A., Erisken, S., Vaiceliunaite, A., Lao, C.L., Mobarhan, M.H., Spacek,
M.A., Einevoll, G.T., and Busse, L. (2021). Corticothalamic feedback sculpts visual spatial integration
in mouse thalamus. Nat Neurosci *24*, 1711–1720. https://doi.org/10.1038/s41593-021-00943-0.

44. Juavinett, A.L., Kim, E.J., Collins, H.C., and Callaway, E.M. (2020). A systematic topographical
relationship between mouse lateral posterior thalamic neurons and their visual cortical projection
targets. Journal of Comparative Neurology *528*, 99–111. https://doi.org/10.1002/cne.24737.

45. Scholl, L.R., Foik, A.T., and Lyon, D.C. (2021). Projections between visual cortex and pulvinar in the
rat. J Comp Neurol *529*, 129–140. https://doi.org/10.1002/cne.24937.

46. Eggermann, E., Kremer, Y., Crochet, S., and Petersen, C.C.H. (2014). Cholinergic Signals in Mouse
Barrel Cortex during Active Whisker Sensing. Cell Reports *9*, 1654–1660.
https://doi.org/10.1016/j.celrep.2014.11.005.

47. Aguilar, J.R., and Castro-Alamancos, M.A. (2005). Spatiotemporal Gating of Sensory Inputs in
Thalamus during Quiescent and Activated States. https://doi.org/10.1523/JNEUROSCI.3229-05.2005.

48. Hirata, A., and Castro-Alamancos, M.A. (2010). Neocortex Network Activation and Deactivation
States Controlled by the Thalamus. J Neurophysiol *103*, 1147–1157.
https://doi.org/10.1152/jn.00955.2009.

49. Masri, R., Trageser, J.C., Bezdudnaya, T., Li, Y., and Keller, A. (2006). Cholinergic regulation of the
posterior medial thalamic nucleus. Journal of Neurophysiology *96*, 2265–2273.
https://doi.org/10.1152/jn.00476.2006.

50. Reimer, J., McGinley, M.J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D.A., and Tolias, A.S. (2016).
Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. Nat Commun
7, 13289. https://doi.org/10.1038/ncomms13289.

51. Varela, C. (2014). Thalamic neuromodulation and its implications for executive networks. Frontiers in
Neural Circuits *8*. https://doi.org/10.3389/fncir.2014.00069.

52. Takahashi, T. (1985). The organization of the lateral thalamus of the hooded rat. J Comp Neurol *231*,
281–309. https://doi.org/10.1002/cne.902310302.

53. Zhou, N.A., Maire, P.S., Masterson, S.P., and Bickford, M.E. (2017). The mouse pulvinar nucleus:
Organization of the tectorecipient zones. Vis Neurosci *34*, E011.
https://doi.org/10.1017/S0952523817000050.

54. Barthó, P., Freund, T.F., and Acsády, L. (2002). Selective GABAergic innervation of thalamic nuclei
from zona incerta. European Journal of Neuroscience *16*, 999–1014. https://doi.org/10.1046/j.1460-
9568.2002.02157.x.

55. Eradath, M.K., Pinsk, M.A., and Kastner, S. (2021). A causal role for the pulvinar in coordinating task-
independent cortico–cortical interactions. Journal of Comparative Neurology *529*, 3772–3784.
https://doi.org/10.1002/cne.25193.

56. Saalmann, Y.B., Pinsk, M.A., Wang, L., Li, X., and Kastner, S. (2012). The pulvinar regulates
information transmission between cortical areas based on attention demands. Science (New York,
N.Y.) *337*, 753–756. https://doi.org/10.1126/science.1223082.

57. Lohse, M., Cooper, M., Sader, E., Langfelder, A., Kahn, M., Baxter, L., Bartram, J., Phillips, J.W., Upton,
A.L., and Mann, E.O. (2018). Motor cortex can modulate somatosensory processing via cortico-
thalamo-cortical pathway. Preprint at bioRxiv, https://doi.org/10.1101/366567
https://doi.org/10.1101/366567.

58. Theyel, B.B., Llano, D.A., and Sherman, S.M. (2010). The corticothalamocortical circuit drives higher-
order cortex in the mouse. Nat Neurosci *13*, 84–88. https://doi.org/10.1038/nn.2449.

59. Petersen, S.E., Robinson, D.L., and Keys, W. (1985). Pulvinar nuclei of the behaving rhesus monkey:
visual responses and their modulation. Journal of neurophysiology *54*, 867–886.
https://doi.org/10.1152/jn.1985.54.4.867.

60. Gregoriou, G.G., Gotts, S.J., Zhou, H., and Desimone, R. (2009). High-Frequency, long-range coupling
between prefrontal and visual cortex during attention. Science *324*, 1207–1210.
https://doi.org/10.1126/science.1171402.

61. Moran, J., and Desimone, R. (1985). Selective attention gates visual processing in the extrastriate
cortex. Science (New York, N.Y.) *229*, 782–784. https://doi.org/10.1126/science.4023713.

62. Kirkwood, A., and Bear, M.F. (1994). Hebbian synapses in visual cortex. J. Neurosci. *14*, 1634–1645.
https://doi.org/10.1523/JNEUROSCI.14-03-01634.1994.

1012

Figure 1: Mice can be conditioned to attend to one sensory modality while ignoring a second modality.

Figure 2: Array recordings were made simultaneously from POm and LP

Figure 3: POm responses to tactile and visual stimuli depend on conditioning paradigm

Figure 4: Conditioning alters tactile stimulus response latencies in POm

a

Population Response

b

c

Figure 5: POm correlates with arousal and movement regardless of conditioning.

Figure 6: POm responses to tactile and visual stimuli vary based on anatomical location.

Figure 7: Conditioning also reshapes LP activity

Figure S1: Latency of anticipatory licking does not depend on the modality of the conditioned stimulus

Figure S2: Manual curation of POm and LP cells.

Figure S3: Conditioning alters stimulus response latencies in LP

Figure S4: LP responses to tactile and visual stimuli as a function of anatomical location