TAK1-mediated phosphorylation of PLCE1 represses PIP2 hydrolysis to impede esophageal squamous cancer metastasis

Qianqian Ju[1], Wenjing Sheng[1], Meichen Zhang[1], Jing Chen[1], Liucheng Wu[2], Xiaoyu Liu[1], Wentao Fang[3,*], Hui Shi[3,*], Cheng Sun[1,*]

[1]Key Laboratory of Neuroregeneration of Jiangsu and Ministry of Education; NMPA Key Laboratory for Research and Evaluation of Tissue Engineering Technology Products; School of Medicine, Nantong University, Nantong, 226001, China.

[2]Laboratory Animal Center, Nantong University, Nantong, 226001, China.

[3]Department of Thoracic Surgery, Shanghai Chest Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, 200030, China.

* Correspondence:

Cheng Sun, Key Laboratory of Neuroregeneration of Jiangsu and Ministry of Education; NMPA Key Laboratory for Research and Evaluation of Tissue Engineering Technology Products; School of Medicine, Nantong University, China. E-mail: suncheng1975@ntu.edu.cn.

Wentao Fang, Department of Thoracic Surgery, Shanghai Chest Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China. E-mail: vwtfang@hotmail.com.

Hui Shi, Department of Thoracic Surgery, Shanghai Chest Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China. E-mail: shihui1@shchest.org.

Abstract

TAK1, a serine/threonine protein kinase, has been identified as a key regulator in a wide

variety of cellular processes. However, its function and involved mechanism in cancer

metastasis are still not well understood. Here, we found that knockdown of TAK1 promoted

esophageal squamous cancer cell (ESCC) migration and invasion, whereas overexpression of

TAK1 resulted in an opposite outcome. Moreover, these *in vitro* findings could be recapitulated

in vivo in a xenograft metastasis mouse model. Mechanistically, co-immunoprecipitation

combined with mass spectrometry demonstrated that TAK1 interacted with phospholipase C

epsilon 1 (PLCE1), and phosphorylated PLCE1 at serine 1060 (S1060). Functional studies

revealed that phosphorylation at S1060 in PLCE1 resulted in decreased enzyme activity,

leading to a repression on PIP2 hydrolysis. As a result, the degradation products of PIP2

including DAG and inositol IP3 were reduced, which thereby suppressed signal transduction in

the axis of PKC/GSK-3β/β-Catenin. Consequently, cancer metastasis related genes were

impeded by TAK1. Overall, our data indicate that TAK1 plays a negative role in ESCC

metastasis, which depends on TAK1 induced phosphorylation of PLCE1 at S1060.

Introduction

Esophageal cancer (EC) is the seventh most common malignancy worldwide with more than 600,000 new cases diagnosed annually (Sung, Ferlay et al., 2021). Clinical survey indicate that EC causes more than 500,000 deaths in 2020 globally (Sung et al., 2021), which is mainly due to its high aggressiveness and poor survival rate. The primary histological subtypes of EC include esophageal squamous cell carcinoma and esophageal adenocarcinoma (He, Xu et al., 2021, Lagergren, Smyth et al., 2017). Esophageal squamous cell carcinoma is a major subtype in China that account for more than 90% of EC (He et al., 2021). Due to the lack of specific biomarkers, the early diagnosis rate of ESCC is very low. Currently, esophageal squamous cell carcinoma diagnosis mainly relies on gastroscopy. By this way, most patients are found to be in locally advanced or metastatic stages when been diagnosed. The 5-year survival rate is approximately 15-25% (Pennathur, Gibson et al., 2013). Surgery has been increasingly used to treat esophageal squamous cell carcinoma at early stages. Neoadjuvant chemotherapy or chemoradiotherapy has become the standard treatment for locally advanced esophageal squamous cell carcinoma (He et al., 2021, Lagergren et al., 2017). In recent decades, therapeutic strategies have made great progress towards esophageal squamous cell carcinoma, and yet recurrence and metastasis are still significant challenges (Mao, Zeng et al., 2021). Therefore, it is urgent to deeply understand the mechanisms involved in cell metastasis, and thus try to explore new druggable targets for treating esophageal squamous cell carcinoma.

Transforming growth factor β-activated kinase 1 (TAK1), a member of the mitogen-activated protein kinase (MAPK) kinase kinase (MAP3K) family, is encoded by the gene *Map3k7*. As a serine/threonine protein kinase, TAK1 has been shown to play an integral role in inflammatory signal transduction across multiple pathways (Sakurai, 2012). By this mean, TAK1 was found to be involved in a wide variety of cellular processes including cell survival, cell migratory and invasive capacity, inflammation, immunity regulation, and tumorigenesis (Cho, Shim et al., 2021, Mukhopadhyay & Lee, 2020). Although TAK1 has been widely studied in cancer progression, its precise role remains controversial. For example, by acting as a tumor suppressor, TAK1 has been shown to negatively associated with tumor progression in several

human cancers including prostate cancer (Huang, Tang et al., 2021), hepatocellular carcinoma

(HCC) (Tan, Zhao et al., 2020, Wang, Zhang et al., 2021), cervical cancer (Guan, Lu et al.,

2017) and certain blood cancers (Guo, Zhang et al., 2019). On the contrary, TAK1 has been

shown to promote tumor progression across a range of human cancers, including colon cancer,

ovarian, lung cancer and breast cancer (Augeri, Langenfeld et al., 2016, Cai, Shi et al., 2014,

Xu, Niu et al., 2022). The above findings indicate that TAK1 may play a dual role in tumor

initiation, progression, and metastasis. In our previous study, TAK1 has been shown to

phosphorylate RASSF9 at serine 284 to inhibit cell proliferation by targeting the

RAS/MEK/ERK axis in ESCC (Shi, Ju et al., 2021). Up to date, there is no report regarding the

precise role of TAK1 on ESCC metastasis.

Phospholipase C epsilon 1 (PLCE1) is encoded by the *Plce1* gene on chromosome 10q23 in

human, and belongs to the phospholipase C (PLC) family (Fukami, Inanobe et al., 2010,

Kadamur & Ross, 2013). Like other PLC family members, PLCE1 is composed of the PLC

catalytic domain, PH domain, EF domain and C2 domain. In addition, PLCE1 also has unique

regions, two C-terminal Ras association (RA) domains and a N-terminal CDC25-homology

domain (Kadamur & Ross, 2013). Once activated, PLCE1 is essential for intracellular signaling

by catalyzing the hydrolysis of membrane phospholipids such as phosphatidylinositol

4,5-bisphosphate (PIP2), to produce two important secondary messengers, inositol

1,4,5-trisphosphate (IP3) and diacylglycerol (DAG), which further trigger IP3-dependent

calcium ions (Ca^{2+}) release from the endoplasmic reticulum and PKC activation (Fukami et al.,

2010, Kadamur & Ross, 2013). Accumulated evidence has shown that PLCE1 promotes

cancer cell growth, migration, and metastasis in multiple human cancers, such as

hepatocellular carcinoma, non-small cell lung cancer, head and neck cancer, bladder cancer,

gastric cancer, and prostate cancer (Abnet, Freedman et al., 2010, Fan, Fan et al., 2019, Liao,

Han et al., 2017, Ma, Wang et al., 2011, Ou, Guo et al., 2010, Wang, Liao et al., 2020, Yue,

Zhao et al., 2019). Till now, however, whether and how PLCE1 affects cancer metastasis in

ESCC are largely unknown.

In this study, we examined the potential role of TAK1 in cell metastasis in ESCC. We found

that TAK1 negatively regulates cell migration and invasion in ESCC, in which PLCE1 is a

downstream target of TAK1. In detail, TAK1 phosphorylates PLCE1 at serine 1060 (S1060) to

inhibit its enzyme activity, leading to decreased IP3 and DAG, both are the products of PLCE1

catalyzed reactions. As a result, IP3/DAG triggered signal transduction in the axis of

PKC/GSK-3β/β-Catenin was blunted. All these outcomes induced by TAK1 caused a

repression on cell migration and invasion in ESCC.

Results

TAK1 negatively regulates ESCC migration and invasion

In our previous study, it has been shown that TAK1 expression was reduced in esophageal

squamous tumor tissues, and TAK1 inhibits ESCC proliferation (Shi et al., 2021). To examine

whether TAK1 affects the epithelial-mesenchymal transition (EMT) process in ESCC, we

increased TAK1 expression in ECA-109 cells by transfecting a plasmid expressing *Map3k7*

(TAK1 gene name) and confirmed the overexpression of TAK1 (**Fig. 1A**). Due to the promising

roles of epidermal growth factor (EGF) on the EMT (Lu, Ghosh et al., 2003), it was used to

trigger the EMT in ESCC. As shown in **Fig. 1B**, EGF treatment induced spindle-shaped cell

morphology in ECA-109 cells, which was markedly prevented by TAK1. These data imply that

TAK1 is a negative regulator of the EMT in ESCC. To address this prediction, we performed

transwell assay and found that TAK1 repressed cell migration and invasion in ECA-109 cells

(**Fig. 1C;** *SI Appendix,* Fig. S1A). In addition, wound healing assay further confirmed the

negative impact of TAK1 on cancer cell migration (**Fig. 1D;** *SI Appendix,* Fig. S1B). Next, we

examined whether TAK1 affects the EMT related gene expression. Our data showed that

TAK1 increased E-cadherin and ZO-1, two epithelial molecules, while mesenchymal

molecules such as N-cadherin, Vimentin, and ZEB1 were reduced by TAK1 (**Fig. 1E;** *SI*

Appendix, Fig. S1C). The qRT-PCR data further confirmed these changes (*SI Appendix,* Fig.

S1D).

Next, we performed loss-of-function studies to examine the effects of TAK1 on cell migration

and invasion in ESCC. The knockdown efficiency of *Map3k7* siRNA on TAK1 expression was

reported previously (Shi et al., 2021). Since the EMT process involves dynamic and spatial

regulation of the cytoskeleton (Fife, McCarroll et al., 2014, Li & Wang, 2020), we therefore

analyzed the expression of F-Actin by immunofluorescence. As shown in **Fig. 1F**, TAK1

knockdown induced F-Actin expression. Meanwhile, TAK1 knockdown promoted

spindle-shaped mesenchymal morphology in ECA-109 cells (**Fig. 1G**). In accordance to these

changes, the cell invasion and migration were stimulated as evidenced by the data from

wound healing and transwell assays (**Fig. 1H;** *SI Appendix,* Fig. S2A-B). The epithelial

markers including E-cadherin and Claudin-1 were decreased by TAK1 knockdown; while the

mesenchymal markers such as N-cadherin, Vimentin, Snail, and Slug were increased (**Fig. 1I**;

SI Appendix, Fig. S2C). At mRNA level, *Cdh1* and *Cldn1* were inhibited by TAK1 knockdown,

while *Cdh2* and *Snail1* were promoted (*SI Appendix,* Fig. S2D). To verify TAK1 knockdown on

cell migration and invasion, we also downregulated TAK1 expression by using lentivirus

carrying *Map3k7* shRNA (LV-*Map3k7* shRNA) (*SI Appendix,* Fig. S3A, B). In LV-*Map3k7*

shRNA transduced cells, cell migration and invasion were obviously increased (*SI Appendix,*

Fig. S3C-E). The epithelial markers including E-cadherin, ZO-1, and Claudin-1 were repressed

by TAK1 knockdown; while the mesenchymal markers such as N-cadherin, Slug, and ZEB-1

were activated (*SI Appendix,* Fig. S3F, G). The qRT-PCR data showed similar changes in

these EMT genes (*SI Appendix,* Fig. S3H). Additionally, TAK1 knockdown was accomplished

by CRISPR-Cas9 using *Map3k7* guide RNA (gRNA) (*SI Appendix,* Fig. S4A). Again, we

observed cell migration and invasion were enhanced by *Map3k7* gRNA (*SI Appendix,* Fig.

S4B-D). The expressions in E-cadherin, ZO-1, and Claudin-1 were inhibited by *Map3k7* gRNA,

while the expressions in N-cadherin, Vimentin, Snail, and Slug were activated (*SI Appendix,*

Fig. S4E, F). Similar changes in these EMT related gene expression were also observed in

qRT-PCR analysis (*SI Appendix,* Fig. S4G). Furthermore, we inhibited TAK1 activity by Oxo,

NG25, or Takinib and observed that all these treatments induced morphological changes in

ECA-109 cells from round shape to spindle-like shape (*SI Appendix,* Fig. S5A). Cell migration

and invasion were also stimulated by TAK1 inhibition (*SI Appendix,* Fig. S5B-E). Taken

together, these findings indicate that TAK1 is a negative regulator of cell migration and

invasion in ESCC.

TAK1 phosphorylates PLCE1 at serine 1060

The above data clearly showed that TAK1 negatively regulates cell migration and invasion in

ESCC. To reveal the involved mechanism, we performed co-immunoprecipitation combined

with mass spectrometry to identify the potential downstream target of TAK1. As previously

reported, 24 proteins were found to be phosphorylated in the immunocomplex (Shi et al.,

2021). Of them, phospholipase C epsilon 1 (PLCE1) caused our attention due to its essential

roles in cell growth, migration, and metastasis in various human cancers (Abnet et al., 2010,

Chen, Wang et al., 2019, Chen, Xin et al., 2020, Gu, Zheng et al., 2018, Kadamur & Ross,

2013, WangZhou et al., 2010). According to the mass spectrometry data, the serine residue at

1060 (S1060) in PLCE1 was phosphorylated (**Fig. 2A**). Currently, there is no antibody against

phosphorylated PLCE1 at S1060 (p-PLCE1 S1060), we therefore generated an antibody for

detecting p-PLCE1 S1060. To verify the antibody specificity, S1060 in PLCE1 was mutated

into alanine (S1060A). As shown in **Fig. 2B**, wild type (WT) PLCE1 could be phosphorylated

by TAK1 at S1060, while PCLE1 S1060A was resistant to TAK1 induced phosphorylation at

S1060. Moreover, p-PLCE1 S1060 induced by TAK1 could be weakened by

(5Z)-7-Oxozeaenol (Oxo), a potent inhibitor of TAK1 (**Fig. 2C**). Of note, phosphorylated TAK1

(p-TAK1) was markedly decreased in the presence of Oxo, indicating TAK1 was inhibited (**Fig.**

2C). To further confirm these findings, other TAK1 inhibitors such as Takinib and NG25 were

employed, and similar changes in p-PLCE1 and p-TAK1 were observed (**Fig. 2D, E**).

Collectively, these results indicate that PLCE1 is a downstream target of TAK1, and PLCE1

S1060 was phosphorylated by TAK1. To further confirm this notion, we analyzed TAK1 and

p-PCLE1 expression in clinical samples. As shown **Fig. 2F**, TAK1 expression was reduced in

tumor tissues as compared to their respective adjacent normal tissues. In accordance to these

changes in TAK1, p-PLCE1 was also decreased in tumor tissues (**Fig. 2G**). The

Pearson correlation tests showing TAK1 expression was positively correlated with p-PLCE1

(**Fig. 2H**). These changes in TAK1 and p-PLCE1 were recapitulated by western blot data (**Fig.**

2I).

TAK1 phosphorylates PLCE1 to inhibit cell migration and invasion in ESCC

It has been well documented that PLCE1 plays a key role in cancer progression (Abnet et al.,

2010, Chen et al., 2019, Chen et al., 2020, Gu et al., 2018, Wang et al., 2010). Therefore, we

predicted that the negative impact of TAK1 on cell migration and invasion in ESCC may rely on

PLCE1. To verify this prediction, we first examined whether PLCE1 affects cell migration and

invasion. In ECA-109 cells, PLCE1 overexpression significantly increased cell migration and

invasion (**Fig. 3A-C**; *SI Appendix*, Fig. S6A, B). In accordance to these changes, the epithelial

marker E-cadherin was repressed by PLCE1, while the mesenchymal markers including

Vimentin, Snail, Slug, and ZEB-1 were activated (**Fig. 3D**; *SI Appendix,* Fig. S6C). The

qRT-PCR data also revealed that *Cdh1* was downregulated by PLCE1, while *Vim*, *Snail1*, and

Snail2 were upregulated (*SI Appendix,* Fig. S6D). To further confirm these findings, we

performed PLCE1 knockdown by using siRNA. As shown in **Fig. 3E**, *Plce1* expression was

markedly reduced by all three tested siRNAs. Similar changes were also observed in the

western blot data (**Fig. 3F**). Of these siRNAs, siRNA-2# exhibited the best knockdown

efficiency and it was chosen in the following experiments. On the contrary, as compared to

PLCE1 overexpression, PLCE1 knockdown impeded cell migration and invasion (**Fig. 3G, H**;

SI Appendix, Fig. S7A, B). As for genes in the EMT process, their expressions exhibited

contrary patterns with comparison of PLCE1 overexpression (**Fig. 3I**; *SI Appendix,* Fig. S7C,

D). Moreover, similar to PLCE1 overexpression, activation of PLCE1 by m-3M3FBS

potentiated cell migration and invasion (*SI Appendix,* Fig. S8A-D). Inhibition of PLCE1 by

U-73122 recapitulated the phenotypes induced by PLCE1 knockdown (*SI Appendix,* Fig.

S9A-D).

Due to PLCE1 is a lipid hydrolase, we next asked whether TAK1-induced phosphorylation in

PLCE1 affects its enzyme activity. To this aim, we transfected ECA-109 cells with a plasmid

bearing Myc-PLCE1 or TAK1, and PLCE1 was captured by pulldown using anti-Myc beads,

which was then subjected to enzyme activity assay. As shown in **Fig. 4A**, the PLCE1 activity

was reduced in the presence of TAK1; however, the PLCE1 S1060A activity was not affected

(**Fig. 4B**), indicating TAK1 induced phosphorylation at S1060 repressed PLCE1 activity. As a

lipid hydrolase, PLCE1 catalyzes PIP2 hydrolysis to produce IP3 and DAG, both are the

second messengers in diverse cellular events (Kadamur & Ross, 2013). Therefore, we

examined these two products to verify the inhibitory effect of TAK1 on PLCE1. Our data

showed that the productions of IP3 and DAG were increased by PLCE1, which could be

largely counteracted by TAK1 (**Fig. 4C-E**). As a messenger, IP3 can induce endoplasmic

reticulum to release Ca^{2+} into the cytoplasm (Kadamur & Ross, 2013). Hence, we analyzed the

cytoplasmic Ca^{2+} by Fluo-4 AM staining. As expected, the signals for the cytoplasmic Ca^{2+}

were obviously increased by PLCE1 in ECA-109 cells, and this trend was weakened by TAK1

(**Fig. 4F, G**). Meanwhile, we also directly detected the cytoplasmic Ca^{2+} with a

fluorospectrophotometer and similar changes were observed (**Fig. 4H**). Flow cytometry

analysis also evidenced the increase in the cytoplasmic Ca^{2+} induced by PLCE1 was

prevented by TAK1 (**Fig. 4I**). All these evidences indicate that TAK1 inhibits PLCE1 activity,

which is likely due to TAK1 induced phosphorylation in PLCE1 at S1060. To further verify this

notion, a series of functional experiments were performed. For instance, we observed that

TAK1 could reverse PLCE1-induced cell migration and invasion; however, TAK1 failed to affect

PLCE1 S1060A-induced cell migration and invasion (*SI Appendix,* Fig. S10A-D). Overall,

these data clearly indicate that PLCE1 has a positive impact on cell migration and invasion in

ESCC. By inducing phosphorylation at S1060, TAK1 inhibits PLCE1 enzyme activity and

thereby counteracts PLCE1-induced cell migration and invasion in ESCC. At this regard,

PLCE1 is a downstream target of TAK1 for transducing its inhibitory effects on cell migration

and invasion in ESCC.

TAK1 inhibits PLCE1-induced signal transduction in the PKC/GSK-3β/β-Catenin axis

PLCE1 hydrolyzes PIP2 to produce two important second messengers DAG and IP3 that

trigger Ca^{2+} release from the endoplasmic reticulum to the cytoplasm to activate PKC (Harden,

Hicks et al., 2009, Kadamur & Ross, 2013), suggesting PKC activation is responsible for

PLCE1-induced cell migration and invasion in ESCC. To address this prediction, we treated

cells with 2-APB, an IP3 receptor (IP3R) inhibitor, and found that 2-APB treatment almost

completely reversed PLCE1-induced the intracellular Ca^{2+} (*SI Appendix,* Fig. S11A-E).

Accordingly, PLCE1-induced cell migration and invasion were also replenished by 2-APB in

ECA-109 cells (*SI Appendix,* Fig. S12A-D). The changes in the EMT genes induced by PLCE1

were largely reversed by 2-APB (*SI Appendix,* Fig. S12E). As well as in ECA-109 cells, 2-APB

also induced similar phenotypes in KYSE-150 and TE-1 cells (*SI Appendix,* Fig. S13A-E; Fig.

S14A-E). Meanwhile, BAPTA-AM, an intracellular Ca^{2+} chelator, was found to be able to

dampen PLCE1-induced cell migration and invasion in ECA-109 cells (*SI Appendix,* Fig.

S15A-D). PKC is a promising downstream target of intracellular Ca^{2+} (Kadamur & Ross, 2013).

We therefore asked whether PLCE1-induced cell growth behaviors are depended on PKC

activation. To this end, Midostaurin was used to inhibit PKC and our data showed that

Midostaurin almost completely abolished cell migration and invasion induced by PLCE1 (*SI*

Appendix, Fig. S16A-D).

It has been shown that PKC positively regulates the expression and stability of β-Catenin in

a GSK-3β-dependent manner (Duong, Yu et al., 2017, Gwak, Cho et al., 2006, Liu, Shi et al.,

2018, Ryu & Han, 2015, Tejeda-Munoz, Gonzalez-Aguilar et al., 2015). Of note, β-Catenin is

considered as a positive regulator in the EMT process (Valenta, Hausmann et al., 2012).

Therefore, we predicted that PLCE1 stimulates cell migration and invasion via the axis of

PKC/GSK-3β/β-Catenin. To test this prediction, ECA-109 cells were treated with IP3 receptor

(IP3R) inhibitor (2-APB), Ca^{2+} chelator (BAPTA-AM), or PKC inhibitor (Midostaurin), all these

treatments successfully inhibited PKC activity as evidenced by reduced phosphorylated PKC

(p-PKC) (**Fig. 5A-C**; *SI Appendix,* S17A-C). As a result, phosphorylated GSK-3β (p-GSK3β)

was decreased by 2-APB, BAPTA-AM, and Midostaurin (**Fig. 5A-C**; *SI Appendix,* S17A-C),

indicating GSK-3β kinase activity was upregulated. Accordingly, phosphorylated β-Catenin

(p-β-Catenin) was increased, leading to degradation of β-Catenin (**Fig. 5A-C**; *SI Appendix,*

S17A-C). As a downstream target of β-Catenin, MMP2 expression was inhibited by all three

tested chemicals (**Fig. 5A-C**; *SI Appendix,* S17A-C). The immunofluorescence data also

showed that PLCE1-induced β-Catenin expression could be counteracted by 2-APB,

BAPTA-AM, and Midostaurin (**Fig. 5D**). These results clearly indicate that PLCE1-induced cell

migration and invasion in ESCC are likely via the axis of PKC/GSK-3β/β-Catenin. Furthermore,

we observed that PLCE1-induced signal transduction in the axis of PKC/GSK-3β/β-Catenin

could be inhibited by TAK1 (**Fig. 5E**; *SI Appendix,* S17D). The expression of β-Catenin

induced by PLCE1 was largely reversed by TAK1 (**Fig. 6F**). However, inactive TAK1 (TAK1

K63W) had no such effects (**Fig. 5G**; *SI Appendix,* S17E). Moreover, TAK1 failed to affect

PLCE1 S1060A induced signal cascade in the axis of PKC/GSK-3β/β-Catenin (**Fig. 5H**; *SI*

Appendix, S17F). These evidences further indicate that TAK1 phosphorylates PLCE1 at

S1060 to inhibit its activity and its downstream signal transduction in the axis of

PKC/GSK-3β/β-Catenin.

TAK1 inhibition promotes ESCC metastasis *in vivo*

To examine TAK1-regulated ESCC metastasis *in vivo*, a xenograft model with nude mice was employed. Mice was injected with ECA109 cells via the tail vein, and then mice were daily treated with Takinib (50 mg/kg) or corn oil (Vehicle) for consecutive 15 days. Eight weeks later, mice were sacrificed for analyzing cancer cell metastasis. We found 4 of 6 mice in the Takinib group developed lung metastasis, while only 1 of 6 mice in the Vehicle group had lung metastasis. Moreover, the number of metastatic nodules in lung tissues was higher in Takinib treated mice (**Fig. 6A-C**). In addition, the effects of TAK1 inhibition on the axis of PKC/GSK-3β/β-Catenin was examined. As shown in **Fig. 6D**, TAK1 inhibition by Takinib activated PKC, leading to increased p-GSK-3β. As a result, p-β-Catenin was reduced and MMP2 expression was upregulated (**Fig. 6D, E**). Of note, p-TAK1 was decreased by Takinib, indicating TAK1 activity was successfully inhibited (**Fig. 6D, E**). In accordance to this change, p-PLCE1 was decreased in Takinib-treated mice (**Fig. 6D, E**). Hence, these data indicate that TAK1 represses ESCC metastasis *in vivo*, and this benefit is likely due to TAK1-mediated phosphorylation in PLCE1 at S1060.

PLCE1 facilitates ESCC metastasis *in vivo*

To further verify the function of PLCE1 in ESCC metastasis, we generated a stable cell line with PLCE1 knockdown by using lentivirus bearing PLCE1 shRNA (LV-shPLCE1) (*SI Appendix,* Fig. S18A-B). These cells with low PLCE1 expression were injected into nude mice via the tail vein to produce a mouse xenograft tumor model. Eight weeks later, mice were sacrificed, the nodule number and incidence rate were reduced in mice injected with LV-shPLCE1 cells (**Fig. 7A-C**). In addition, western blot analysis showed that PLCE1 silencing inhibited signal transduction in the PKC/GSK-3β/β-Catenin axis, as evidenced by reduced p-PKC and p-GSK-3β, and increased p-β-Catenin (**Fig. 7D, E**). In accordance to these changes, the expression of MMP2 was decreased (**Fig. 7D, E**). These results further confirmed the notion that PLCE1 plays a key role in cancer cell metastasis in ESCC.

Discussion

TAK1 is a serine/threonine kinase and a major member of the MAPK family (Zhu, Lama et al.,

2021). In response to various cytokines, pathogens, lipopolysaccharides, hypoxia, and DNA

damage, the E3 ligase TRAF2 or TRAF6 activates TAK1 through Lys63-linked

polyubiquitylation, and then TAK1 undergoes auto-phosphorylation at Ser192 and Thr184/187

to achieve full activation (Skaug, Jiang et al., 2009, Sorrentino, Thakur et al., 2008).

Consequently, activated TAK1, in turn, phosphorylates downstream substrates to spark the

NF-kB and MAPKs (i.e. ERK, p38 MAPK, and JNK) signaling pathways, thereby participates in

cellular inflammation, immune response, fibrosis, cell death, and cancer cell invasion and

metastasis (Yang, Chen et al., 2022, Zhou, Tao et al., 2021). TAK1 has been shown to

decrease in high-grade human prostate cancer; and TAK1 deficiency promotes prostate

tumorigenesis by increasing androgen receptor (AR) protein levels and activity or activating

p38 MAPK pathway (Huang et al., 2021, Wu, Shi et al., 2012). Similarly, hepatocyte-specific

TAK1 ablation drives RIPK1 kinase-dependent inflammation to promote liver fibrosis and

hepatocellular carcinoma (Su, Gao et al., 2023, Tan et al., 2020, Xia, Ji et al., 2021).

Furthermore, TAK1 represses the transcription of human telomerase; and activate tumor

suppressor protein LKB1, all these evidences indicate that TAK1 is a tumor suppressor

(Adhikari, Xu et al., 2007, Fujiki, Miura et al., 2007, Xie, Zhang et al., 2006). Consistent with

these findings, in our previous study, we found that TAK1 expression was reduced in

esophageal squamous tumors compared to adjacent normal tissues, and TAK1 negatively

correlates with esophageal squamous tumor patient survival (Shi et al., 2021). In this study, we

extended our research on TAK1 and observed that TAK1 inhibits cell migration and invasion in

ESCC. In contrast to these findings, TAK1 has been shown to play a positive role in tumor cell

proliferation, migration, invasion, colony formation, and metastasis, especially in breast cancer,

pancreatic cancer, and non-small cell lung cancer (Kim, Kim et al., 2023, Santoro, Zanotto et

al., 2020, Tripathi, Shin et al., 2019). Therefore, TAK1 may play pleiotropic roles in different

types of cancer cells.

As a kinase, TAK1 is considered as a central regulator of tumor cell proliferation, migration

and invasion and has been demonstrated to play tumor suppression or activation role via the

NF-κB and MAPK activation (Mukhopadhyay & Lee, 2020, Wang et al., 2021, Zhang, Cheng et

al., 2023). However, the precise molecular and cellular mechanisms by which TAK1 regulates ESCC metastasis remain unclear. As previously reported, by co-immunoprecipitation coupled with mass spectrometry (MS/MS), we identified RASSF9 as a downstream target of TAK1 for transducing its repression on cell proliferation in ESCC (Shi et al., 2021). By phosphorylating RASSF9 at S284, TAK1 negatively regulates cell proliferation in ESCC (Shi et al., 2021). Meanwhile, we also found that PLCE1 was presented in the immunocomplex, which was phosphorylated at S1060 in the presence of TAK1 in ECA-109 cells. As a member of the human phosphoinositide-specific PLC family, PLCE1 can be activated by a variety of intracellular and extracellular signal molecules including hormones, cytokines, neurotransmitters, growth factors, etc. (Kadamur & Ross, 2013). Numerous studies have shown that PLCE1 plays a key role in cancer development and progression through various pathways (Abnet et al., 2010, Fan et al., 2019, Ghosh, Nataraj et al., 2021, He, Wang et al., 2016, Yue et al., 2019). Moreover, PLCE1 has been characterized as a susceptibility gene in ESCC (Abnet et al., 2010, Wang et al., 2010). All these evidences suggest that PLCE1 is a potential downstream target for transducing the negative impact of TAK1 on cell migration and invasion in ESCC.

Once activated, PLCE1 catalyzes the hydrolysis of PIP2 on the cell membrane to produce two secondary messengers IP3 and DAG; IP3 induces Ca^{2+} release from the ER into the cytoplasm via IP3R; both Ca^{2+} and DAG are potent activators of PKC. By this way, PLCE1 regulates cell growth, proliferation, and differentiation and thereby plays a key role in tumor growth and development (Bunney & Katan, 2010, Kadamur & Ross, 2013, Land & Rubin, 2017, Smrcka, Brown et al., 2012). Consistent with these notions, in this study, we observed that PLCE1-overexpressing cells exhibit higher production of intracellular IP3, DAG, and intracellular Ca^{2+}. As a consequence, we found that PLCE1 overexpression accelerates cell migration and invasion, while PLCE1 knockdown results in an opposite phenotype. Moreover, in the presence of TAK1, PLCE1-induced cell migration and invasion were largely counteracted. By considering TAK1 is a protein kinase, therefore, we proposed that TAK1 phosphorylates PLCE1 at S1060 to inhibit its enzyme activity and thus impedes cell migration and invasion in ESCC. Indeed, PLCE1 activity was blunted in the presence of TAK1, whereas

mutated PLCE1 (PLCE1 S1060A) was not affected. It has been shown that PKC positively

regulates the expression and stability of β-Catenin in a GSK-3β-dependent manner (Duong et

al., 2017, Gwak et al., 2006, Ryu & Han, 2015). Therefore, we investigated the impact of

PLCE1 on the PKC/GSK-3β/β-Catenin axis. As expected, our data revealed that PLCE1

promotes cell migration and invasion in ESCC by activating the PKC/GSK-3β/β-Catenin

pathway, results in a series of phosphorylation modification in PKC and GSK-3β. As a result,

phosphorylated β-Catenin decreased, leading to higher stability of β-Catenin, which eventually

promotes the EMT process by affecting epithelial and mesenchymal gene expression.

In summary, our findings indicate TAK1 is a negative regulator in cell migration and invasion

in ESCC. Mechanistically, TAK1 phosphorylates PLCE1 at S1060 to inhibit its phospholipase

activity, leading to reductions in IP3 and DAG. Consequently, PKC activity was blunted, which

results in decreased phosphorylated GSK-3β and increased phosphorylated β-Catenin. As a

consequence, the stability of β-Catenin was decreased and its transcriptional activity was

blunted. By this mean, the epithelial marker gene expression was upregulated by TAK1, while

the mesenchymal marker gene expression was downregulated. All these outcomes eventually

cause a breakdown in cell migration and invasion in ESCC. Hence, our data revealed a new

facet of TAK1 in the EMT process in ESCC by inhibiting PLCE1 activity and its downstream

signal transduction in the axis of PKC/GSK-3β/β-Catenin. Moreover, TAK1, together with its

downstream target PLCE1, are potential druggable targets for developing agents for dealing

with ESCC.

Methods

Cell culture

Human ESCC cell lines ECA109, KYSE-150 and TE-1 were purchased from the Shanghai

Institute of Biochemistry and Cell Biology (Shanghai, China). HEK-293 cells were purchased

from American Type Culture Collection (ATCC, Manassas, VA, USA). All cells were cultured in

Dulbecco's modified Eagle's (DMEM) medium (Hyclone, UT, USA) containing 10% fetal bovine

serum (FBS, Gibco, Carlsbad, CA), 100 U/ml penicillin and 100 μg/ml streptomycin (Life

Technologies, Carlsbad, CA, USA) at 37°C in a humidified atmosphere of 5% CO_2.

Human esophageal squamous tumor specimens

Human esophageal squamous tumor specimens were obtained from the Affiliated Hospital of Nantong University. Human sample collection for research was conducted in accordance with the recognized ethical guideline of Declaration of Helsinki and approved by the Ethics Committee of Affiliated Hospital of Nantong University.

Generation of antibodies against phospho-PLCE1 (S1060)

Rabbit polyclonal antibody that recognizes phospho-PLCE1 (S1060) was raised against c-WSARNPS(p)PGTSAK peptide at Absin (Shanghai, China). Briefly, the phosphorylated polypeptide c-WSARNPS(p)PGTSAK was synthesized as the target peptide. Then two rabbits were immunized with the target peptide antigen. Meanwhile, non-phosphorylated polypeptide c-WSARNPSPGTSAK was synthesized as a control. The rabbits were sacrificed and blood was collected for further purification. The prepared antibody was verified and stored at -20°C until use.

CRISPR-Cas9 mediated TAK1 deletion

Map3k7 was deleted by the method of CRISPR-Cas9 with the procedures describe previously (Shi et al., 2021). Briefly, we designed an integrated vector based on OriP/EBNA1 (epiCRISPR), which bears gRNA, Cas9 and puromycin resistance genes. ECA-109 cells were transfected with the plasmids using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. The cells were cultured in a medium containing 2.5 µg/ml puromycin for 1 week. The surviving stable cells were used for further experiments. The gRNA sequence is listed in Table S1.

Protein extraction and western blot analysis

Western blot assay was performed as previously described (Tan, Krueger et al., 2022). Briefly, Total protein were extracted from cells and tissues using protein lysis buffer supplemented with protease and phosphatase inhibitors (Roche Applied Science, Penzberg, Germany). Protein concentrations were assayed by the Pierce™ BCA Protein Assay Kit (Thermo Scientific,

Waltham, MA, USA). Subsequently, equal amounts of protein were separated with 6% or 10%

SDS-PAGE, and then protein were transferred onto 0.45 μm PVDF membranes. After blocked

with 5% BSA in TBST for 1 h, the membranes were incubated with primary antibodies at 4°C

overnight. Thereafter, the membranes were incubated with HRP-linked secondary antibodies

for 1 h at room temperature. Protein signals were visualized using enhanced

chemiluminescence (ECL) reagent (Thermo Fisher Scientific, Waltham, MA, USA) and

quantitatively analyzed using ImageJ (NIH). Actin were used as a loading control.

Measurement of intracellular-free Ca^{2+}

Cells were transfected with plasmid expressing PLCE1 and/or TAK1 for 6 h and then treated

with BAPTA-AM (10 μM), 2-APB (10 μM), or Midostaurin (100 nM) for additional 18 h,

respectively. Measurement of intracellular Ca^{2+} in treated cells was performed using a Fluo-4

AM Assay Kit (Beyotime, Jiangsu, China). In detail, cells were washed with PBS, and

incubated with 1 μM Fluo-4 AM in PBS at 37°C for 30 min. Cell were then washed with PBS for

3 times and further incubated for 20 min to ensure Fluo-4 AM completely transformed into

Fluo-4 in cells. Finally, Fluo-4 fluorescent intensity was detected with a confocal laser scanning

fluorescence microscope (Olympus BX51, Tokyo, Japan), fluorescence microplate (BioTek

Instruments, Inc.), or flow cytometry (FACSCalibur, BD Bioscience, Franklin Lakes, NJ, USA)

at excitation of 488 nm and emission of 520 nm to determine the change of intracellular Ca^{2+}

concentration.

PLCE1 pulldown and phospholipase activity assay

Protein pulldown was performed using a previously reported method (Shi et al., 2021). Briefly,

HEK293 cells were transfected with a plasmid expressing PLCE1-Myc or TAK1. 24 h

post-transfection, cells were harvested for preparing total cell lysates, which was then

subjected to protein pulldown assay using the Myc-Tag beads (Sepharose® Bead Conjugate;

#55464; CST, Beverley, MA, USA). After 4 h-incubation on a rotator, unbound proteins were

removed using the ice-cold wash buffer. Finally, phospholipase activity bound on the beads

was detected with a PLC Activity Assay Kit (Jining Shiye, Shanghai, China) with the provided

protocol. In rationale, PLC catalyzes the hydrolysis of O-(4-nitrophenyl) choline (NPPC) to

produce p-nitrophenol (PNP), which has an absorption maximum at 405 nm. By detecting the

increase rate of PNP at 405 nm, the magnitude of PLC enzyme activity can be calculated.

IP3 and DAG assays

IP3 and DAG levels in cells were assayed by a Human Inositol 1,4,5-triphosphate

enzyme-linked Immunosorbent Assay (ELISA) Kit and a Human Diacylglycerol commercial

ELISA Kit (Mlbio, Shanghai, China) according to the manufacturer's instructions. The

absorbance was measured at 450 nm with a microplate reader.

Co-immunoprecipitation and MS/MS spectrometry

Co-immunoprecipitation and MS/MS assay were performed as previously described (Shi et al.,

2021). Briefly, ECA-109 cells were transfected with a plasmid expressing *Map3k7*. 36 h

post-transfection, cells were harvested for co-immunoprecipitation by using an antibody

against TAK1. The resulting immunocomplex was subjected to LC-MS/MS spectrometry

analysis (Shanghai Applied Protein Technology Co., Ltd.). MS/MS spectra were searched

using MASCOT engine (Matrix Science, London, UK; version 2.4) against the UniProKB

human (161584 total entries, downloaded 20180105).

Mouse xenograft metastasis model

For *in vivo* metastasis assay, 4-week-old male BALB/c nude mice (18-20 g) were purchased

from Shanghai Slake Laboratory Animal Co. Ltd. (Shanghai, China) and randomly divided into

2 groups (n = 6 - 10 per group). Mice were housed in air-filtered laminar flow cages (5

mice/cage) with 12 h light cycle and adequate food and water. To examine the role of PLCE1

on metastasis, each mouse was intravenously (i.v.) injected with 1×10^6 ECA109 cells diluted

in 100 μl PBS, which were transduced with LV-shPLCE1 or negative control virus. Additionally,

to examine TAK1 inhibition on metastasis, each mouse was intravenously (i.v.) injected with 1

$\times 10^6$ ECA109 cells. Mice were then intraperitoneally injected with Takinib (50 mg/kg/day) or

vehicle (corn oil) for 15 days. Mouse body weight was measured once a week. Eight weeks

later, mice were sacrificed, and lungs and livers were collected for further analysis. All animal

experiments were carried out in accordance with the institutional ethical guidelines of animal

care, and were approved by the Animal Experimentation Ethics Committee of the Nantong

University (Approval ID: SYXK (SU) 2017-0046).

Statistical analysis

Statistical significance of differences between groups were tested by using analysis of

variance (ANOVA) or unpaired Student's *t*-test. Data were presented as mean ± SD from at

least three independent experiments. All statistical analyses were performed using GraphPad

Prism version 8.0 (GraphPad Software, La Jolla, CA, USA). * $p < 0.05$ was accepted as

statistical significance.

Data availability

The dataset regarding MS-based protein phosphorylation modifications can be found in

Figshare https://doi.org/10.6084/m9.figshare.25271140.

References

Abnet CC, Freedman ND, Hu N, Wang Z, Yu K, Shu XO, Yuan JM, Zheng W, Dawsey SM, Dong LM, Lee
MP, Ding T, Qiao YL, Gao YT, Koh WP, Xiang YB, Tang ZZ, Fan JH, Wang C, Wheeler W et al. (2010) A
shared susceptibility locus in PLCE1 at 10q23 for gastric adenocarcinoma and esophageal squamous
cell carcinoma. *Nat Genet* 42: 764-7

Adhikari A, Xu M, Chen ZJ (2007) Ubiquitin-mediated activation of TAK1 and IKK. *Oncogene* 26:
3214-26

Augeri DJ, Langenfeld E, Castle M, Gilleran JA, Langenfeld J (2016) Inhibition of BMP and of TGFbeta
receptors downregulates expression of XIAP and TAK1 leading to lung cancer cell death. *Mol Cancer* 15:
27

Bunney TD, Katan M (2010) Phosphoinositide signalling in cancer: beyond PI3K and PTEN. *Nat Rev*
Cancer 10: 342-52

Cai PC, Shi L, Liu VW, Tang HW, Liu IJ, Leung TH, Chan KK, Yam JW, Yao KM, Ngan HY, Chan DW (2014)
Elevated TAK1 augments tumor growth and metastatic capacities of ovarian cancer cells through
activation of NF-kappaB signaling. *Oncotarget* 5: 7549-62

Chen Y, Wang D, Peng H, Chen X, Han X, Yu J, Wang W, Liang L, Liu Z, Zheng Y, Hu J, Yang L, Li J, Zhou H,
Cui X, Li F (2019) Epigenetically upregulated oncoprotein PLCE1 drives esophageal carcinoma
angiogenesis and proliferation via activating the PI-PLCepsilon-NF-kappaB signaling pathway and
VEGF-C/ Bcl-2 expression. *Mol Cancer* 18: 1

Chen Y, Xin H, Peng H, Shi Q, Li M, Yu J, Tian Y, Han X, Chen X, Zheng Y, Li J, Yang Z, Yang L, Hu J, Huang
X, Liu Z, Huang X, Zhou H, Cui X, Li F (2020) Hypomethylation-Linked Activation of PLCE1 Impedes
Autophagy and Promotes Tumorigenesis through MDM2-Mediated Ubiquitination and Destabilization
of p53. *Cancer Res* 80: 2175-2189

Cho SH, Shim HJ, Park MR, Choi JN, Akanda MR, Hwang JE, Bae WK, Lee KH, Sun EG, Chung IJ (2021)
Lgals3bp suppresses colon inflammation and tumorigenesis through the downregulation of
TAK1-NF-kappaB signaling. *Cell Death Discov* 7: 65

Duong M, Yu X, Teng B, Schroder P, Haller H, Eschenburg S, Schiffer M (2017) Protein kinase C ϵ
stabilizes beta-catenin and regulates its subcellular localization in podocytes. *J Biol Chem* 292:
12100-12110

Fan J, Fan Y, Wang X, Niu L, Duan L, Yang J, Li L, Gao Y, Wu X, Luo C (2019) PLCepsilon regulates
prostate cancer mitochondrial oxidative metabolism and migration via upregulation of Twist1. *J Exp*
Clin Cancer Res 38: 337

Fife CM, McCarroll JA, Kavallaris M (2014) Movers and shakers: cell cytoskeleton in cancer metastasis.
Br J Pharmacol 171: 5507-23

Fujiki T, Miura T, Maura M, Shiraishi H, Nishimura S, Imada Y, Uehara N, Tashiro K, Shirahata S, Katakura
Y (2007) TAK1 represses transcription of the human telomerase reverse transcriptase gene. *Oncogene*
26: 5258-66

Fukami K, Inanobe S, Kanemaru K, Nakamura Y (2010) Phospholipase C is a key enzyme regulating
intracellular calcium and modulating the phosphoinositide balance. *Prog Lipid Res* 49: 429-37

Ghosh S, Nataraj NB, Noronha A, Patkar S, Sekar A, Mukherjee S, Winograd-Katz S, Kramarski L, Verma
A, Lindzen M, Garcia DD, Green J, Eisenberg G, Gil-Henn H, Basu A, Lender Y, Weiss S, Oren M, Lotem
M, Geiger B et al. (2021) PD-L1 recruits phospholipase C and enhances tumorigenicity of lung tumors
harboring mutant forms of EGFR. *Cell Rep* 35: 109181

Gu D, Zheng R, Xin J, Li S, Chu H, Gong W, Qiang F, Zhang Z, Wang M, Du M, Chen J (2018) Evaluation of
GWAS-Identified Genetic Variants for Gastric Cancer Survival. *EBioMedicine* 33: 82-87

Guan S, Lu J, Zhao Y, Woodfield SE, Zhang H, Xu X, Yu Y, Zhao J, Bieerkehazhi S, Liang H, Yang J, Zhang F,
Sun S (2017) TAK1 inhibitor 5Z-7-oxozeaenol sensitizes cervical cancer to doxorubicin-induced
apoptosis. *Oncotarget* 8: 33666-33675

Guo D, Zhang A, Huang J, Suo M, Zhong Y, Liang Y (2019) Suppression of HSP70 inhibits the
development of acute lymphoblastic leukemia via TAK1/Egr-1. *Biomed Pharmacother* 119: 109399

Gwak J, Cho M, Gong SJ, Won J, Kim DE, Kim EY, Lee SS, Kim M, Kim TK, Shin JG, Oh S (2006)
Protein-kinase-C-mediated beta-catenin phosphorylation negatively regulates the Wnt/beta-catenin
pathway. *J Cell Sci* 119: 4702-9

Harden TK, Hicks SN, Sondek J (2009) Phospholipase C isozymes as effectors of Ras superfamily
GTPases. *J Lipid Res* 50 Suppl: S243-8

He S, Xu J, Liu X, Zhen Y (2021) Advances and challenges in the treatment of esophageal cancer. *Acta*
Pharm Sin B 11: 3379-3392

He Y, Wang C, Wang Z, Zhou Z (2016) Genetic variant PLCE1 rs2274223 and gastric cancer: more to be
explored? *Gut* 65: 359-60

Huang Z, Tang B, Yang Y, Yang Z, Shi L, Bai Y, Yan B, Karnes RJ, Zhang J, Jimenez R, Wang L, Wei Q, Yang J,
Xu W, Jia Z, Huang H (2021) MAP3K7-IKK Inflammatory Signaling Modulates AR Protein Degradation
and Prostate Cancer Progression. *Cancer Res* 81: 4471-4484

Kadamur G, Ross EM (2013) Mammalian phospholipase C. *Annu Rev Physiol* 75: 127-54

Kim MJ, Kim JY, Shin JH, Kang Y, Lee JS, Son J, Jeong SK, Kim D, Kim DH, Chun E, Lee KY (2023) FFAR2
antagonizes TLR2- and TLR3-induced lung cancer progression via the inhibition of AMPK-TAK1
signaling axis for the activation of NF-kappaB. *Cell Biosci* 13: 102

Lagergren J, Smyth E, Cunningham D, Lagergren P (2017) Oesophageal cancer. *Lancet* 390: 2383-2396

Land M, Rubin CS (2017) A Calcium- and Diacylglycerol-Stimulated Protein Kinase C (PKC),
Caenorhabditis elegans PKC-2, Links Thermal Signals to Learned Behavior by Acting in Sensory
Neurons and Intestinal Cells. *Mol Cell Biol* 37

Li X, Wang J (2020) Mechanical tumor microenvironment and transduction: cytoskeleton mediates
cancer cell invasion and metastasis. *Int J Biol Sci* 16: 2014-2028

Liao X, Han C, Qin W, Liu X, Yu L, Zhu G, Yu T, Lu S, Su H, Liu Z, Chen Z, Yang C, Huang K, Liu Z, Liang Y,
Huang J, Dong J, Li L, Qin X, Ye X et al. (2017) PLCE1 polymorphisms and expression combined with
serum AFP level predicts survival of HBV-related hepatocellular carcinoma patients after hepatectomy.
Oncotarget 8: 29202-29219

Liu T, Shi Y, Chan MTV, Peng G, Zhang Q, Sun X, Zhu Z, Xie Y, Sham KWY, Li J, Liu X, Ho IHT, Gin T, Lu Z,
Wu WKK, Cheng CHK (2018) Developmental protein kinase C hyper-activation results in microcephaly
and behavioral abnormalities in zebrafish. *Transl Psychiatry* 8: 232

Lu Z, Ghosh S, Wang Z, Hunter T (2003) Downregulation of caveolin-1 function by EGF leads to the loss
of E-cadherin, increased transcriptional activity of beta-catenin, and enhanced tumor cell invasion.
Cancer Cell 4: 499-515

Ma H, Wang LE, Liu Z, Sturgis EM, Wei Q (2011) Association between novel PLCE1 variants identified in
published esophageal cancer genome-wide association studies and risk of squamous cell carcinoma of
the head and neck. *BMC Cancer* 11: 258

Mao C, Zeng X, Zhang C, Yang Y, Xiao X, Luan S, Zhang Y, Yuan Y (2021) Mechanisms of Pharmaceutical
Therapy and Drug Resistance in Esophageal Cancer. *Front Cell Dev Biol* 9: 612451

Mukhopadhyay H, Lee NY (2020) Multifaceted roles of TAK1 signaling in cancer. *Oncogene* 39:
1402-1413

Ou L, Guo Y, Luo C, Wu X, Zhao Y, Cai X (2010) RNA interference suppressing PLCE1 gene expression
decreases invasive power of human bladder cancer T24 cell line. *Cancer Genet Cytogenet* 200: 110-9

Pennathur A, Gibson MK, Jobe BA, Luketich JD (2013) Oesophageal carcinoma. *Lancet* 381: 400-12

Ryu JM, Han HJ (2015) Autotaxin-LPA axis regulates hMSC migration by adherent junction disruption
and cytoskeletal rearrangement via LPAR1/3-dependent PKC/GSK3beta/beta-catenin and PKC/Rho
GTPase pathways. *Stem Cells* 33: 819-32

Sakurai H (2012) Targeting of TAK1 in inflammatory disorders and cancer. *Trends Pharmacol Sci* 33:
522-30

Santoro R, Zanotto M, Simionato F, Zecchetto C, Merz V, Cavallini C, Piro G, Sabbadini F, Boschi F,
Scarpa A, Melisi D (2020) Modulating TAK1 Expression Inhibits YAP and TAZ Oncogenic Functions in
Pancreatic Cancer. *Mol Cancer Ther* 19: 247-257

Shi H, Ju Q, Mao Y, Wang Y, Ding J, Liu X, Tang X, Sun C (2021) TAK1 Phosphorylates RASSF9 and
Inhibits Esophageal Squamous Tumor Cell Proliferation by Targeting the RAS/MEK/ERK Axis. *Adv Sci*
(Weinh) 8: 2001575

Skaug B, Jiang X, Chen ZJ (2009) The role of ubiquitin in NF-kappaB regulatory pathways. *Annu Rev*
Biochem 78: 769-96

Smrcka AV, Brown JH, Holz GG (2012) Role of phospholipase Cepsilon in physiological
phosphoinositide signaling networks. *Cell Signal* 24: 1333-43

Sorrentino A, Thakur N, Grimsby S, Marcusson A, von Bulow V, Schuster N, Zhang S, Heldin CH,
Landstrom M (2008) The type I TGF-beta receptor engages TRAF6 to activate TAK1 in a receptor
kinase-independent manner. *Nat Cell Biol* 10: 1199-207

Su W, Gao W, Zhang R, Wang Q, Li L, Bu Q, Xu Z, Liu Z, Wang M, Zhu Y, Wu G, Zhou H, Wang X, Lu L

(2023) TAK1 deficiency promotes liver injury and tumorigenesis via ferroptosis and macrophage
cGAS-STING signalling. *JHEP Rep* 5: 100695

Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F (2021) Global Cancer
Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185
Countries. *CA Cancer J Clin* 71: 209-249

Tan R, Krueger RK, Gramelspacher MJ, Zhou X, Xiao Y, Ke A, Hou Z, Zhang Y (2022) Cas11 enables
genome engineering in human cells with compact CRISPR-Cas3 systems. *Mol Cell* 82: 852-867 e5

Tan S, Zhao J, Sun Z, Cao S, Niu K, Zhong Y, Wang H, Shi L, Pan H, Hu J, Qian L, Liu N, Yuan J (2020)
Hepatocyte-specific TAK1 deficiency drives RIPK1 kinase-dependent inflammation to promote liver
fibrosis and hepatocellular carcinoma. *Proc Natl Acad Sci U S A* 117: 14231-14242

Tejeda-Munoz N, Gonzalez-Aguilar H, Santoyo-Ramos P, Castaneda-Patlan MC, Robles-Flores M (2015)
Glycogen Synthase Kinase 3beta Is Positively Regulated by Protein Kinase Czeta-Mediated
Phosphorylation Induced by Wnt Agonists. *Mol Cell Biol* 36: 731-41

Tripathi V, Shin JH, Stuelten CH, Zhang YE (2019) TGF-beta-induced alternative splicing of TAK1
promotes EMT and drug resistance. *Oncogene* 38: 3185-3200

Valenta T, Hausmann G, Basler K (2012) The many faces and functions of beta-catenin. *EMBO J* 31:
2714-36

Wang L, Zhang X, Lin ZB, Yang PJ, Xu H, Duan JL, Ruan B, Song P, Liu JJ, Yue ZS, Fang ZQ, Hu H, Liu Z,
Huang XL, Yang L, Tian S, Tao KS, Han H, Dou KF (2021) Tripartite motif 16 ameliorates nonalcoholic
steatohepatitis by promoting the degradation of phospho-TAK1. *Cell Metab* 33: 1372-1388 e7

Wang LD, Zhou FY, Li XM, Sun LD, Song X, Jin Y, Li JM, Kong GQ, Qi H, Cui J, Zhang LQ, Yang JZ, Li JL, Li
XC, Ren JL, Liu ZC, Gao WJ, Yuan L, Wei W, Zhang YR et al. (2010) Genome-wide association study of
esophageal squamous cell carcinoma in Chinese subjects identifies susceptibility loci at PLCE1 and
C20orf54. *Nat Genet* 42: 759-63

Wang XK, Liao XW, Yang CK, Liu ZQ, Han QF, Zhou X, Zhang LB, Deng T, Gong YZ, Huang JL, Huang R,
Han CY, Yu TD, Su H, Ye XP, Peng T, Zhu GZ (2020) Oncogene PLCE1 may be a diagnostic biomarker and
prognostic biomarker by influencing cell cycle, proliferation, migration, and invasion ability in
hepatocellular carcinoma cell lines. *J Cell Physiol* 235: 7003-7017

Wu M, Shi L, Cimic A, Romero L, Sui G, Lees CJ, Cline JM, Seals DF, Sirintrapun JS, McCoy TP, Liu W, Kim
JW, Hawkins GA, Peehl DM, Xu J, Cramer SD (2012) Suppression of Tak1 promotes prostate
tumorigenesis. *Cancer Res* 72: 2833-43

Xia S, Ji L, Tao L, Pan Y, Lin Z, Wan Z, Pan H, Zhao J, Cai L, Xu J, Cai X (2021) TAK1 Is a Novel Target in
Hepatocellular Carcinoma and Contributes to Sorafenib Resistance. *Cell Mol Gastroenterol Hepatol* 12:
1121-1143

Xie M, Zhang D, Dyck JR, Li Y, Zhang H, Morishima M, Mann DL, Taffet GE, Baldini A, Khoury DS,
Schneider MD (2006) A pivotal role for endogenous TGF-beta-activated kinase-1 in the
LKB1/AMP-activated protein kinase energy-sensor pathway. *Proc Natl Acad Sci U S A* 103: 17378-83

Xu G, Niu L, Wang Y, Yang G, Zhu X, Yao Y, Zhao G, Wang S, Li H (2022) HDAC6-dependent deacetylation
of TAK1 enhances sIL-6R release to promote macrophage M2 polarization in colon cancer. *Cell Death*
Dis 13: 888

Yang L, Chen H, Hu Q, Liu L, Yuan Y, Zhang C, Tang J, Shen X (2022) Eupalinolide B attenuates
lipopolysaccharide-induced acute lung injury through inhibition of NF-kappaB and MAPKs signaling by
targeting TAK1 protein. *Int Immunopharmacol* 111: 109148

Yue QY, Zhao W, Tan Y, Deng XL, Zhang YH (2019) PLCE1 inhibits apoptosis of non-small cell lung cancer

via promoting PTEN methylation. *Eur Rev Med Pharmacol Sci* 23: 6211-6216

Zhang X, Cheng L, Gao C, Chen J, Liao S, Zheng Y, Xu L, He J, Wang D, Fang Z, Zhang J, Yan M, Luan Y,

Chen S, Chen L, Xia X, Deng C, Chen G, Li W, Liu Z et al. (2023) Androgen Signaling Contributes to Sex

Differences in Cancer by Inhibiting NF-kappaB Activation in T Cells and Suppressing Antitumor

Immunity. *Cancer Res* 83: 906-921

Zhou Y, Tao T, Liu G, Gao X, Gao Y, Zhuang Z, Lu Y, Wang H, Li W, Wu L, Zhang D, Hang C (2021) TRAF3

mediates neuronal apoptosis in early brain injury following subarachnoid hemorrhage via targeting

TAK1-dependent MAPKs and NF-kappaB pathways. *Cell Death Dis* 12: 10

Zhu L, Lama S, Tu L, Dusting GJ, Wang JH, Liu GS (2021) TAK1 signaling is a potential therapeutic target

for pathological angiogenesis. *Angiogenesis* 24: 453-470

Acknowledgements

We would like to thank Friedhelm Hildebrandt from University of Michigan Drive for providing the plasmid expressing PLCE1. This work was supported by the National Natural Science Foundation of China (No. 32271193); the Natural Science Foundation of Shanghai (No. 21ZR1449800); the Scientific Project from Shanghai Municipal Health Commission (202240015); the Postgraduate Research & Practice Innovation Program of Jiangsu Province (KYCX23_3411); and the Priority Academic Program Development (PAPD) of Jiangsu Higher Education Institutions.

Author contributions

QJ performed all the experiments with assistance from WS, MZ, and JC. XL designed oligonucleotide sequences for qRT-PCR, shRNA, siRNA, and mutated PLCE1. LW provided advice throughout the development of the project. CS, TW, and HS developed the study concept and experiment design, supervised the study, and wrote the manuscript with input from all the authors.

Disclosure and competing interest statement

The authors declare that they have no competing interests.

Figure legends

Figure 1. TAK1 negatively regultes ESCC migration and invasion

(A) Increased expression of TAK1 in ECA-109 cells transfected with a plasmid expressing

Map3k7.

(B) Increased expression of TAK1 inhibits the morphological changes to form spindle-shaped

mesenchymal cells induced by EGF (100 ng/ml) in ECA-109. Scale bar = 100 μm.

(C) Increased expression of TAK1 inhibits cell migration and invasion in ECA-109 cells. Cell

migration and invasion were analyzed by transwell assay. Scale bar = 500 μm. n = 4

biologically independent replicates.

(D) Wound healing assay showing cell migration was attenuated by TAK1. n = 5 biologically

independent replicates.

(E) TAK1 decreased mesenchymal marker gene expression, while increased the expression of

epithelial markers. ECA-109 cells were transfected with the plasmid carrying *Map3k7*. 24 h

post-transfection, protein samples were prepared and subjected to western blot. Actin was

used as a loading control.

(F) Knockdown of TAK1 increased the expression of F-Actin. ECA-109 cells were transfected

with *Map3k7* siRNA. Seventy-two h post-transfection, cells were subjected to

immunofluorescence analysis using an anti-F-Actin antibody (red). Hoechst was used to stain

the nucleus (blue). Scale bar = 100 μm.

(G) TAK1 knockdown induces spindle-shaped mesenchymal cell morphology in ECA-109 cells.

Scale bar = 100 μm.

(H) Reduced expression of TAK1 promotes cell migration and invasion. ECA-109 cells were

transfected with *Map3k7* siRNA. 72 h post-transfection, cell migration and invasion were

analyzed by transwell assay. n = 5 biologically independent replicates.

(I) Knockdown of TAK1 increases mesenchymal protein marker expression, and decreases

epithelial protein marker expression.

Data are presented as mean ± SD. Statistical significance was tested by unpaired Student's

t-test. *$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$.

Figure 2. TAK1 phosphorylates PLCE1 at serine 1060

(A) Tandem mass spectrometry showing serine 1060 (S1060) in PLCE1 was phosphorylated

by TAK1. ECA-109 cells were transfected with a plasmid expressing *Map3k7*. Twenty-four

h-post transfection, cells were harvested and subjected to co-immunoprecipitation. The

resulting immunocomplex was analyzed by liquid chromatography coupled with tandem mass

spectrometry (LC-MS/MS).

(B) TAK1 fails to phosphorylate PLCE1^{S1060A}. ECA-109 cells were co-transfected with the

plasmids carrying wildtype (WT) *Plce1*, mutated *Plce1* (PLCE1^{S1060A}), or *Map3k7* as indicated.

Twenty-four h-post transfection, cells were collected for western blot analysis.

(C-E) Inhibition of TAK1 reduces PLCE1 phosphorylation at S1060. ECA-109 cells were

co-transfected with the plasmids expressing *Plce1* or *Map3k7*. 6 h post-transfection, TAK1

inhibitor (5Z)-7-Oxozeaenol (Oxo; 10 μM) (C), or 10 μM Takinib (D), or 10 μM NG25 (E) was

added in culture medium, and cells were cultured for additional 18 h. Cells were then

subjected to western blot analysis. Actin was used as a loading control.

(F-G) Immunohistochemical analysis of TAK1 (F) and p-PLCE1 (G) expression in normal and

esophageal squamous tumor tissues. *n* = 4 biologically independent replicates. Scale bar = 20

μm.

(H) Correlation between p-PLCE1 and TAK1 based on immunohistochemical data as shown in

(F-G). 10 views for each sample were randomly chosen for Pearson correlation test.

(I) TAK1 and p-PLCE1 protein levels in clinical samples. Protein levels were analyzed by

western blot, and Actin was used as a loading control. *n* = 4 biologically independent replicates.

N: normal tissue; T: tumor tissue.

Figure 3. PLCE1 positively regulates ESCC migration and invasion

(A) Increased expression of PLCE1 in ECA-109 cells transfected with a plasmid expressing

Plce1.

(B-C) Increased expression of PLCE1 enhances cell migration and invasion. ECA-109 cells

were transfected with the plasmid carrying *Plce1*. Twenty-four h-post transfection, cells were

subjected to transwell (B) or wound healing (C) assay. *n* = 5 biologically independent

replicates.

(D) Increased expression of PLCE1 in ECA-109 cells induces mesenchymal protein marker

expression, while reduces epithelial protein marker expression.

(E-F) Knockdown of PLCE1. ECA-109 cells were transfected with siRNAs targeting *Plce1*.

Seventy-two h-post transfection, cells were harvested for analyzing PLCE1 expression by

qRT-PCR (E) and western blot (F).

(G-H) Reduced expression of PLCE1 inhibits cell migration and invasion. ECA-109 cells were

transfected with *Plce1* siRNA-2. Forty-eight h-post transfection, cells were subjected to

transwell (G) or wound healing (H) assay. *n* = 3-5 biologically independent replicates.

(I) Knockdown of PLCE1 promotes epithelial protein marker expression, while represses

mesenchymal protein marker expression. ECA-109 cells were transfected with *Plce1* siRNA-2.

Seventy-two h-post transfection, cells were harvested and subjected to western blot analysis.

Protein level was detected by western blot, and Actin was used as a loading control. Gene

expression was analyzed by qRT-PCR and *Gapdh* was used as a house-keeping gene. Data

are presented as mean ± SD. Statistical significance was tested by unpaired Student's *t*-test.

*$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$.*

Figure 4. TAK1 inhibits PLCE1 enzyme activity

(A-B) Effects of TAK1 on PLCE1 (A) and PLCE1 S1060A (B) enzyme activity. ECA-109 cells

were co-transfected with the plasmids expressing *Plce1-Myc* or *Map3k7*. Twenty-four h

post-transfection, cells were subjected to pull down assay by using the beads with anti-Myc

antibody. PLCE1 enzyme activity was assayed by Phospholipases C (PLC) Activity Assay Kit.

n = 3 biologically independent replicates.

(C-E) TAK1 abolishes PLCE1-induced IP3 and DAG in ECA-109 (C), KYSE-150 (D), and TE-1

cells (E). Cells were transfected with the plasmids bearing *Plce1* or *Map3k7* as indicated.

Twenty-four h post-transfection, cells were harvested for measuring IP3 and DAG. *n* = 3

biologically independent replicates.

(F) TAK1 attenuates PLCE1-induced intracellular Ca^{2+} ($[Ca^{2+}]$). ECA-109 cells were

transfected with the plasmids bearing *Plce1* or *Map3k7* as indicated. $[Ca^{2+}]$ was labeled with

Fluo-4 AM, which was then detected by a fluorescent microscope. Scale bar = 200 μm.

(G) Quantified fluorescence intensity of $[Ca^{2+}]$ in ECA-109 cells. *n* = 3 biologically independent

replicates.

(H) Fluorescence intensity of Fluo-4 in ECA-109, KYSE-150, and TE-1 cells was examined

with a fluorospectrophotometer. n = 3 biologically independent replicates.

(I) Flow cytometry analysis of $[Ca^{2+}]$. Cell treatments were described in (F).

Data are presented as mean ± SD. Statistical significance was tested by unpaired Student's

t-test (A-B) or two-tailed one-way ANOVA test (C-E, G-I). *$p < 0.05$, **$p < 0.01$, and ***$p <$

$0.001.$

Figure 5. TAK1 inhibits PLCE1-induced signal transduction in the axis of

PKC/GSK-3β/β-Catenin

(A) IP3R blocking inhibits PLCE1 induced signal transduction in the axis of

PKC/GSK-3β/β-Catenin. ECA-109 cells were transfected with the plasmid expressing *Plce1*

for 6 h and then treated with 2-APB (10 µM) for additional 18 h.

(B) $[Ca^{2+}]$ blocking represses signal transduction in the axis of PKC/GSK-3β/β-Catenin

induced by PLCE1. ECA-109 cells were transfected with the plasmid expressing *Plce1* for 6 h

and then treated with BAPTA-AM (10 µM) for additional 18 h.

(C) PKC inhibition blocks PLCE1 stimulated signal transduction in the axis of

PKC/GSK-3β/β-Catenin. ECA-109 cells were transfected with the plasmid expressing *Plce1*. 6

h post-transfection, cells were treated with 100 nM of Midostaurin for additional 18 h.

(D) PKC inhibition represses PLCE1-induced nuclear translocation of β-Catenin in ECA-109

cells. Cells were transfected with the plasmid expressing PLCE1. Six h post-transfection,

2-APB (10 µM), BAPTA-AM (10 µM), or Midostaurin (100 nM) was added in culture medium,

and cells were cultured for additional 18 h. Scale bar = 100 µm. Immunofluorescence was

used to examine subcellular distribution of β-Catenin.

(E) TAK1 counteracts PLCE1-induced signal transduction in the axis of

PKC/GSK-3β/β-Catenin. ECA-109 cells were transfected with the plasmids expressing *Plce1*

or *Map3k7* as indicated for 24 h.

(F) TAK1 reduces PLCE1-induced nuclear distribution of β-Catenin in ECA-109 cells. Cells

were transfected with the plasmids expressing *Plce1* or *Map3k7* as indicated. Scale bar = 100

µm.

(G) Dominant negative TAK1 (K63W) fails to block signal transduction in the axis of

PKC/GSK-3β/β-Catenin/MMP2 induced by PLCE1. ECA-109 cells were transfected with the

plasmids expressing *Plce1* or mutated *Map3k7* (TAK1 K63W) for 24 h.

(H) TAK1 has no effect on PLCE1 S1060A induced signal transduction in the axis of

PKC/GSK-3β/β-Catenin. ECA-109 cells were transfected with the plasmids expressing PLCE1

S1060A or TAK1 for 24 h. Protein levels were analyzed by western blot, and Actin was used as

a loading control. Representative blots were shown.

Figure 6. Inhibition of TAK1 by Takinib promotes ESCC metastasis in nude mice

Each mouse was intravenously injected with 1×10^6 ECA109 cells diluted in 100 μl PBS. Mice

were treated with Takinib at the dosage of 50 mg/kg/day for 15 days, mice in control group

were received vehicle (corn oil). Eight weeks later, mice were sacrificed, and the lungs and

livers from each group were collected and photographed.

(A) Typical images of specimens.

(B) Hematoxylin and eosin staining of metastatic nodules in lungs.

(C) The number of nodules in lungs.

(D) Takinib treatment induces signal transduction in the axis of PKC/GSK-3β/β-Catenin.

Protein levels were analyzed by western blot, and Actin was used as a loading control. *n* = 3

biologically independent replicates.

(E) Quantitative analysis of the western blot data shown in (D).

Data are presented as mean ± SD. Statistical significance was tested by unpaired Student's

t-test. *$p < 0.05$, **$p < 0.01$*, and ***$p < 0.001$.*

Figure 7. PLCE1 knockdown inhibits ESCC metastasis in nude mice

ECA-109 cells were transduced with lentivirus bearing PLCE1 shRNA (LV-shPLCE1) or NC

shRNA (LV-shPLCE1 NC). Each mouse was intravenously injected with the LV transduced

cells (1×10^6 cells/mouse). Eight weeks later, mice were sacrificed, and the lungs and livers

from each group were collected and photographed.

(A) Typical images of lung specimens.

(B) Hematoxylin and eosin staining of metastatic nodules in lungs.

(C) The number of nodules in lungs.

(D) PLCE1 knockdown represses signal transduction in the axis of PKC/GSK-3β/β-Catenin.

Protein levels were analyzed by western blot, and Actin was used as a loading control. $n = 3$

biologically independent replicates.

(E) Quantitative analysis of the western blot data shown in (D).

Data are presented as mean ± SD. Statistical significance was tested by unpaired Student's

t-test. **p < 0.01*, and ****p < 0.001.*

922 FIGURE 1

FIGURE 2

FIGURE 3

955   FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7