Transcriptomics and trans-organellar complementation reveal a limited signaling capacity

of 12-*cis*-oxo-phytodienoic acid in wounded Arabidopsis

Khansa Mekkaoui[1], Ranjit Baral[1], Fiona Smith[1], Moritz Klein[2], Ivo Feussner[2], Bettina

Hause[1,*]

[1] Department of Cell and Metabolic Biology, Leibniz Institute of Plant Biochemistry, Weinberg

3, D-06120 Halle/Saale, Germany

[2] Department of Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences and

Goettingen Center for Molecular Biosciences (GZMB), University of Goettingen, Justus-von-

Liebig-Weg 11, 37077 Göttingen, Germany

Corresponding author: Bettina Hause, Bettina.Hause@ipb-halle.de

Running title:

Signaling and compartmentalization of OPDA

Abstract

12-*cis*-Oxo-phytodienoic acid (OPDA), the main precursor of the key plant growth and defense hormone jasmonoyl-isoleucine (JA-Ile), is believed to have distinct signaling roles in plant's responses to stress. In Arabidopsis, insights into OPDA functions have been gained from studying a mutant, which is affected in the conversion of OPDA by missing OPDA REDUCTASE3 (OPR3). *opr3* mutants, however, accumulate JA-Ile through a cytosolic bypass mediated by OPR2. Therefore, wound-induced transcriptome of *opr2opr3* in comparison to wild-type and *allene oxide synthase* mutant was analyzed to unravel OPDA signaling. Results showed that OPDA lacked a distinct transcriptional signature, whereas known OPDA-response genes were wound-induced independently of OPDA. The application of OPDA to *opr2opr3* resulted in a distinct transcriptional response compared to the endogenous rise of OPDA in the same mutant, with the activation of the sulfur assimilation pathway genes occurring only with the external application of the compound. These findings suggested a compartmentalization of endogenously produced OPDA, investigated further through trans-organellar complementation. OPR3 complemented *opr2opr3* mutants in fertility and wound-induced JA-Ile production regardless of its localization. Since *in vitro* enzymatic studies revealed OPR3's activity on both OPDA and 4,5-ddh-JA, conclusions on translocation of OPDA were not unequivocal. Dissecting the conversion of either OPDA or 4,5-ddh-JA by OPR2 and OPR1 organelle variants pointed, however, to a strong OPDA compartmentalization supporting its lacking signaling function.

Introduction

Jasmonic acid (JA) and its bioactive form jasmonoyl-isoleucine (JA-Ile) function primarily as signaling molecules coordinating plant growth and reproduction and responses to biotic and abiotic stresses [1-4]. They play a central role in the defense of plants against mechanical damage and herbivory, as JA-Ile rapidly accumulates at the site of injury and acts as a signaling molecule by triggering a cascade of defense responses [5,6]. JA-Ile signaling involves a series of molecular events mediated by its perception by the F-box protein CORONATINE INSENSITIVE 1 (COI1) leading to the degradation of the JASMONATE ZIM-DOMAIN (JAZ) repressors, hence the release of the expression of JA-responsive genes [7-13].

One major intermediate in the formation of the bioactive JA-Ile, 12-*cis*-oxo-phytodienoic acid (OPDA), has been known to be a signaling molecule with independent functions from JA-Ile [14-17]. OPDA formation takes place in the chloroplast via the octadecanoid pathway, by oxygenation of α-linolenic acid by 13-lipoxygenases (LOXs) and the coupled actions of ALLENE OXIDE SYNTHASE (AOS) and ALLENE OXIDE CYCLASE (AOC) [18-20] (Fig. 1a). Metabolization of OPDA into JA occurs after its transfer to peroxisomes, where the reduction of its cyclopentenone ring by OPDA REDUCTASE3 (OPR3) and three rounds of β-oxidation take place [1,21,22]. A parallel hexadecanoid pathway gives rise to *dinor*-12-oxo-phytodienoic acid (*dn*-OPDA), which in turn is converted to JA following the same pathway as OPDA, but with two rounds of β-oxidation only [23].

The putative distinct function of OPDA in Angiosperms partially derives from its function in non-vascular plants by inducing defense- and growth inhibition-related transcriptional reprogramming in *Marchantia polymorpha* [13,24-26]. Such JA-independent functions of OPDA were also characterized in vascular plants for different vegetative processes including initiation of tendril coiling in *Bryonia dioica* [27], regulation of embryo development in *Solanum lycopersicum*

[28], and inhibition of seed germination and induction of stomatal closure in *Arabidopsis thaliana* [29-31]. Despite its perception being not identified in Arabidopsis, OPDA was shown to act as signaling molecule by transcriptionally inducing a set of genes, which is not induced by JA [32]. These OPDA-specific response genes were characterized as wound- and OPDA-responsive in a COI1-independent manner and encompassed mainly stress-responsive genes, e.g. *DRE-BINDING PROTEIN 2A* (*DREB2A)*, *FAD-LINKED OXIDOREDUCTASE* (*FAD-OXR), SALT TOLERANCE ZINC FINGER* (*ZAT10)*, *ETHYLENE RESPONIVE ELEMENT BINDING FACTOR5* (*ERF5*), *GLUTATHION TRANSFERASE6* (*GST6*), and the glutaredoxin member *GRX480*. OPDA contains an α, β-unsaturated carbonyl group which is thought to react with nucleophilic molecules by Michael addition [33]. The electrophilic nature of oxylipins could convey biological activities, as deduced from the accumulation of oxylipins adducts, including OPDA-GSH conjugates, during the hypersensitive response in pathogen-elicited tobacco leaves [34]. OPDA binding to the CYCLOPHILIN20-3 (CYP20-3) was suggested to trigger the formation of the cysteine synthase complex activating sulfur assimilation to regulate cell redox homeostasis through mediation of *TGACG-BINDING* (TGA) transcription factors [35,36]. Here, CYP20-3-dependent OPDA signaling induces high levels of GSH leading to a retrograde signaling, which coordinates expression of OPDA-induced genes [37]. These transcriptional changes in Arabidopsis induced by OPDA through CYP20-3 module and/or additional uncharacterized mechanism consolidated the hypothesis of its function not limited to being only a JA precursor [16].

The previous work characterizing OPDA signaling and JA-independent functions in Arabidopsis relied mainly on the use of the *opr3* mutant as the genetic background where OPDA and JA biosynthesis were uncoupled due to missing OPR3 [38,39]. The *opr3* mutant is male sterile like the *aos* mutant being deficient in OPDA and JA [38,40,41]. Both mutants show defects in anther dehiscence, stamen elongation and pollen development showing that these defects were due to

impairments in JA production. Persistent resistance of the *opr3* mutant against necrotrophic

pathogens initially hinted towards a defense function of OPDA, but further studies uncovered a

conditional removal of the T-DNA containing intron upon fungal infection of *opr3-1* on the one

hand [42], and an OPR3-independent pathway leading to JA/JA-Ile formation through the cytosolic

OPR2 in the *opr3-3* mutant on the other hand [43]. Low JA-Ile formation in both *opr3* mutants

rather than OPDA was shown to be responsible of its resistance, hence challenging the suggested

JA-independent OPDA signaling function [42,43]. Mainly the recently identified formation of JA-Ile

via bypassing the OPR3 (Fig. 1a) questioned previously identified functions of OPDA.

Consequently, OPDA-specific signaling exclusively analyzed with OPDA-producing but JA/JA-

Ile deficient mutants requires further studies.

This study re-addressed the question whether OPDA can function as an independent

signaling molecule in Arabidopsis besides being a JA precursor. Using the *opr2-1 opr3-3* mutant

(further named *opr2opr3*) where formation of JA/JA-Ile does not occur [43], we investigated the

effects of basal and wound-induced OPDA on gene expression in seedlings and rosette leaves of

A. thaliana. The results suggested that the endogenously formed OPDA is unlikely to be

responsible for mediating gene expression, while exogenous supply of OPDA resulted in a

distinct signaling. The activation of the sulfur assimilation pathway exclusively by exogenously

supplied OPDA hints towards its predominant detoxification within the cell but indicates a

compartmentalization of the endogenously formed OPDA. Therefore, we addressed the question

of the occurrence of OPDA within the cell and whether it could be perceived. A trans-organellar

complementation approach showed that OPDA is likely to be confined to the cell compartments

where it is synthesized (plastids), transported through (cytosol) and metabolized (peroxisomes).

Instead, 4,5-ddh-JA appears to be a mobile compound within the plant cell supporting the

assumption that OPDA has no signaling function in wounded Arabidopsis plants.

Results

A JA positive feedback loop during development is responsible for the tissue capability to produce OPDA.

The *opr2opr3* double mutant was characterized as JA-deficient, with JA formation through the peroxisomal OPR3 and alternatively through the cytosolic OPR2 being both interrupted [43]. The JA deficiency in the *opr2opr3* and *aos* mutants was confirmed by measuring JA content under both control and wounding conditions (Supplementary Fig. 1a). In the mutants, JA levels remained at the detection limit, with approximately 0.002 nmol/g of fresh weight JA detected in the wounded *opr2opr3*. This contrasts with the reported 0.2 nmol/g of fresh weight JA accumulation in the wounded *opr3* mutant plants[39], thereby confirming the JA deficiency of *opr2opr3* [43]. JA deficiency is associated with a decrease in the accumulation of JA-biosynthesis enzymes, among them specifically the AOC, due to the disruption of the JA positive feedback loop during development [44]. Consequently, lower background and wound-induced levels of OPDA were observed in the *opr2opr3* mutant compared to the wild-type (Supplementary Figure 1b), which indeed correlated with diminished AOC protein content in *opr2opr3* and *aos* seedlings in comparison to wild-type (Col-0) seedlings (Supplementary Fig. 1c, d).

Due to the partial OPDA deficiency of *opr2opr3* seedlings, it was crucial to normalize the levels with those of the wild type prior to investigating OPDA signaling within this mutant. To achieve this, we mimicked the missing JA positive feedback loop in the *opr2opr3* and *aos* mutants by supplying seedlings with 1 µM JA methyl ester (MeJA) during development to increase the AOC protein levels (Supplementary Fig. 2a). In contrast to water (mock-)pretreated seedlings, seedlings pretreated with MeJA during development exhibited increased AOC protein levels in wild-type and both mutants (Fig. 1b). Notably, this increase was more pronounced in the

mutants resulting in AOC protein levels in *opr2op3* and *aos* seedlings matching those of wild

type seedlings (Fig. 1c). Transcript accumulation of the JA biosynthesis enzymes *LOX2*, *AOS*,

AOC2 and *OPR3* determined by RT-qPCR corroborated the function of the JA positive feedback

loop: MeJA pretreatment during development resulted in either a compensation or an

enhancement of their basal transcript levels in the mutants compared to wild type (Fig. 1d). They

all are, however, highly induced upon wounding in wild-type seedlings, but their induction is

significantly lower in the mutants (Fig. 1d). This dampened induction demonstrates that the

MeJA pretreatment did not activate a wound-induced JA-Ile signaling in the JA-deficient

mutants. Most importantly, however, the restoration of AOC protein content in the *opr2op3*

enhanced its basal and wound-induced OPDA and *dn*-OPDA levels to the levels of wild-type,

whereas the *aos* mutant remained OPDA deficient (Fig. 1e). Despite the accumulation of low

residual levels of JA in *opr2opr3* and *aos* seedlings upon pretreatment, wound-induced

accumulation of JA and JA-Ile was only observed in wild-type seedlings and correlated with the

transcriptional response of the JA-responsive genes (Fig. 1d). This validates the JA deficiency of

both mutant lines within this experimental design and the exclusive OPDA enhancing role of the

developmental JA-feedback loop for the *opr2opr3* mutant (Fig. 1e).

A comparative transcriptomics approach does not show an OPDA-specific response to

wounding.

OPDA signaling in Arabidopsis was previously inferred from its capacity to mediate

transcriptional changes by inducing a specific set of OPDA-responsive genes in the *opr3-1* single

mutant [32,45]. Another putative OPDA-signaling pathway was based on the detected binding of

OPDA to CYP20-3, which in turn transcriptionally induces TGA transcription factors [35]. To

further clarify OPDA-related signaling function, a comparative transcriptomics approach was set

between wild-type and *opr2opr3* seedlings. To uncouple putative OPDA signaling from common

wound-induced signaling processes, *aos* seedlings were included as a negative control, since both

JA and OPDA production are abolished in this mutant (Supplementary Fig. 2b). mRNA

sequencing of MeJA-pretreated seedlings from Col-0, *opr2opr3* and *aos* was performed using

unwounded seedlings (control) and seedlings harvested 1 h after wounding. The overall wound-

induced response showed a significant transcriptional change in all three genotypes

(Supplementary Fig. 3). Here, a strong similarity between the transcriptomes of *opr2opr3* and *aos*

seedlings was evident in the expression heatmap, with a clear overlap observed in the principal

component analysis (PCA), irrespective of the condition (Supplementary Fig. 3). The wounding

transcriptional response showed most difference in the number of differentially expressed genes

(DEGs) between Col-0 and the JA-deficient mutants, with the DEGs being 10.84% and 7.48%

greater in Col-0 than *opr2opr3* and *aos*, respectively (Fig. 2a). A common wound response was

observed in all the genotypes which was enriched in genes involved in stress, oxygen, and abiotic

stimulus responses and was therefore rather wound-inducible but completely independent from

the JA pathway (Fig. 2a, b, Supplementary Dataset 1). Interestingly, this group contained known

stress-responsive genes with several genes being previously reported as OPDA-responsive, such

as *DREB2A* and *FAD-OXR,* which were also induced in the *aos* mutant by wounding despite its

OPDA deficiency[32] (Fig. 2c, d, Supplementary Tab. 4, Supplementary Dataset 1). Additionally,

other major OPDA-marker genes like *ZAT10, ERF5, TCH4* and *GST6* also showed a wound-

responsive induction in *aos* seedlings indicating that they are more likely to be associated with an

OPDA-independent wound-induced stress response rather than with an OPDA-mediated response

(Fig. 2d). To exclude a JA/JA-Ile dependence of their induction, transcript levels of these genes

were monitored in rosette leaves of two *coi1*-mutant lines, *coi1-16* and *coi1-30* (Supplementary

Fig. 4a, b). The data show that all tested genes are inducible by wounding to nearly the same

level as in wild type, irrespective of the lower levels of OPDA in the *coi1*-mutants reaching in

coi1-16 only 25 % of that of wounded wild type leaves, whereas there was no increase in

wounded leaves of *coi1-30* (Supplementary Fig. 4c). Hence, their transcriptional induction by

wounding did not correlate with OPDA levels despite its COI1-independence indicating that they

are rather governed by other signaling processes in the wound response. It is interesting to note

that also JA and JA-Ile levels appeared to be significantly lower in wounded *coi1* plants in

comparison to wild type pointing again to the missing positive feedback in JA signaling and

biosynthesis during plant development [44].

In wild-type seedlings, the signaling mediated by JA/JA-Ile was evident as it distinctly

shaped the transcriptome, sustaining statistically differential basal levels of 71 JA-regulated

genes when directly comparing the wild-type transcriptome to those of the mutants at control

condition (Supplementary Fig. 5a, b). Wounding of wild-type seedlings resulted in a more drastic

transcriptional change highlighted by 432 DEGs specifically induced in Col-0 and enriched in

genes related to JA, wounding, and immune responses (Supplementary Fig. 5a, c, Supplementary

Dataset 2). These transcriptional responses, notably of genes encoding JAZ transcription factors

and other JA-responsive genes, such as *CHLOROPHYLLASE 1 (CLH1)* and *N-*

ACETYLTRANSFERASE ACTIVITY 1 (NATA1), were dampened in the *opr2opr3* and *aos*

mutants confirming the JA dependence of their induction upon wounding (Supplementary Fig.

5d, Supplementary Dataset 2). These data suggest that next to a common wound response,

JA/JA-Ile are main mediators of wound-induced transcriptional changes in wild-type seedlings.

Wound-induced and exogenously applied OPDA resulted in different alterations of the

transcriptome in *opr2opr3* seedlings.

To identify OPDA-regulated genes, transcriptional differences between *opr2opr3* and *aos* were

dissected and indicated a restricted number of DEGs at control and wounding conditions (Fig.

3a). Except for the mutated genes, which showed diminished expression in the respective mutant

backgrounds (*AOS*, *OPR2*, and *OPR3*), only nine specific genes exhibited differential expression

in *opr2opr3* seedlings compared to *aos* seedlings, regardless of the experimental condition (Fig.

3b, c). These genes encode five RNA species and four proteins, among them one cyclic

nucleotide-gated ion channel (*CNGC11*), one cytochrome P450 (*CYP81D11*), and two

hypothetical proteins. This is a very narrow or almost no difference between the transcriptomic

profiles of the mutants, despite the presence of significantly high OPDA levels in the *opr2opr3*

mutant compared to its absence in *aos* (Fig. 1e). These results make an OPDA-mediated gene

expression in wounded seedlings of the *opr2opr3* mutant unlikely.

To investigate the impact of developmental stage on our findings, the wounding experiment was

done using mature four-week-old rosettes (Supplementary Fig. 2c). Pretreatment with 1 µM

MeJA enhanced OPDA levels in wounded leaves of *opr2opr3* compared to the mock pretreated

ones (Supplementary Fig. 6a). Regardless of whether the MeJA pretreatment was administrated,

wounding led to a high number of DEGs in the wild-type and a lower, but similar number of

DEGs in the *opr2opr3* and *aos* mutants (Supplementary Fig. 6b, c). Most importantly, consistent

with the findings for seedlings, the transcriptomes of the *aos* and *opr2opr3* mutants were highly

similar despite significant OPDA accumulation in the latter and differed only with the same

DEGs found in the transcriptomes of seedlings (Supplementary Fig. 6d-f).

Previous work demonstrated a signaling role of OPDA in Arabidopsis by its exogenous supply to

wild-type plants or to the *opr3* mutant [32,35,45]. To determine whether the origin of OPDA caused

the different outcomes, RNAseq data resulting from wounding of *opr2opr3* seedlings (=

endogenous rise of OPDA) were compared to transcriptomic data generated from *opr2opr3*

seedlings, which were treated with 25 µM OPDA for 30 min (= exogenous supply of OPDA).

OPDA treatment resulted in transcriptional changes that were highly different from those induced

by wounding (Supplementary Fig. 7a, Supplementary Dataset 3). The clustering of DEGs

induced by OPDA treatment in comparison to DEGs induced by wounding showed a significant

induction of a sub-cluster of genes belonging to sulfate reduction and assimilation pathway, the

phosphorelay signal transduction pathway as well as the hormone-mediated signaling pathway

(Supplementary Fig. 7b, Fig. 4a, Supplementary Dataset 3). These genes were not induced by the

endogenous rise of OPDA and included genes encoding sulfate reductases, such as SULFATE-

DEFICIENCY INDUCED (SDI1), APS REDUCTASE3 (APR3), and RESPONSE TO LOW

SULFUR3 (LSU31), and ORA59, an APETALA2/ETHYLENE RESPONSE FACTOR

(AP2/ERF) domain transcription factor which is involved in JA and ethylene signaling and in

defense [46] (Fig. 4b). This is in line with the previously reported induction of sulfur metabolism by

OPDA binding to the CYP20-3 module [35]. Here, the activation of the sulfur metabolism pathway

was a specific effect of exogenously supplied OPDA that did not correlate with an endogenous

effect of the compound. Formerly described OPDA-induced genes were induced by both OPDA

treatment and wounding of the *opr2opr3* mutant, however, their wound induction in *aos*

indicated their involvement in general stress-response pathways (Fig. 2, Fig. 4d). In addition,

OPDA application to *opr2opr3* seedlings resulted in a slight, but significant induction of JA

signaling genes (Supplementary Fig. 7a, b, Supplementary Dataset 3). To get a deeper insight

into this JA-mediated response occurring in *opr2opr3* seedlings upon OPDA treatment,

expression levels of JA responsive genes were determined in comparison to wounded seedlings

from wild-type and both mutants (Supplementary Fig. 7c). OPDA-treated *opr2opr3* and wounded

wild-type seedlings shared the up-regulation of several JA marker genes, such as those encoding

JAZ transcription factors and JA biosynthesis enzymes (Supplementary Fig. 7c). Validation of

transcript accumulation of *JAZ2*, *JAZ7*, *JAZ13* and *CHL1* by RT-qPCR confirmed their exclusive

induction in *opr2opr3* seedlings upon OPDA application but not upon endogenous OPDA

accumulation by wounding (Supplemental Fig. 7d). This hints towards the conversion of the

exogenously applied but not the endogenously formed OPDA to JA/JA-Ile despite the loss of

OPR3 and OPR2 in the *opr2opr3* background. OPDA and JA levels in *opr2opr3* seedlings

determined after wounding and OPDA treatment showed unequivocally that *opr2opr3* seedlings

accumulated slowly higher levels of JA-Ile following OPDA application in comparison to

wounding (Supplemental Fig. 7e). Despite these levels being lower than the JA-Ile levels in

wounded wild-type seedlings, they correlated with the activated JA signaling in *opr2opr3*

seedlings after OPDA application. Interestingly, the OPDA-fed *opr2opr3* seedlings accumulated

only slightly higher OPDA levels than the wounded seedlings, whereas *dn*-OPDA accumulated

significantly only following endogenous OPDA formation after wounding (Supplemental Fig.

7e). This suggests that the exogenously applied OPDA is either detoxified by conjugation to GSH

[47] or rapidly converted to 4,5-ddh-JA as shown by Ueda et al. (accompanying manuscript). In

contrast, *dn*-OPDA accumulating after wounding in *opr2opr3* seedlings might be produced from

OPDA entering the β-oxidation cycle [43] or from the parallel hexadecanoid pathway [23]. These

results raise the question whether the different patterns of gene expression detected here, are due

to a compartmentalization of OPDA synthesized following wounding, thereby not inducing the

responses that the exogenously supplied OPDA does.

Test for subcellular distribution of the endogenously produced OPDA using a trans-

organellar complementation of the *opr2opr3* mutant.

The data obtained from wounded seedlings of *opr2opr3* and described above suggest that

endogenously produced OPDA does not exhibit signaling capacity. This leads to the question,

whether OPDA is restricted to the cell compartments of its synthesis, transfer and metabolization,

namely plastids, cytosol and peroxisomes, respectively. To test its putative translocation to other

cell compartments, a trans-organellar complementation approach according to [48] using the

opr2opr3 mutant was performed. OPR3 was targeted to different organelles in the *opr2opr3*

mutant, and the fertility of these plants was checked. A rescue of the JA-deficiency phenotype of

the *opr2opr3* mutant would indicate whether its substrate, OPDA, is able to translocate to the

tested organelles (Supplementary Fig. 8).

OPR3 is a peroxisome-localized protein and is imported into the matrix of peroxisomes

through the peroxisomal targeting signal type 1 (PTS1) consisting of a tripeptide at the extreme

carboxy-tail of the protein [49]. OPR3 contains a C-terminal SRL, which is a variation of the

prototypic PTS1 signal SKL. Removal of this signal resulted in cytosolic localization of a fusion

of OPR3ΔSRL with YFP when transiently expressed in *Nicotiana benthamiana* leaves, indicating

that import into peroxisomes was abolished (Fig. 5a). Targeting of OPR3 to the nucleus, cytosol,

plastids, mitochondria, and endoplasmic reticulum (ER) was done by fusing OPR3ΔSRL with the

respective targeting signals or target peptides (Supplementary Tab. 5). Validation of the

subcellular localization of the different organelle variants of OPR3 with co-localization studies in

N. benthamiana protoplasts was performed using established organelle markers [50] and indicated

correct subcellular localizations of OPR3 compared to the markers independently of the position

of the YFP fluorescent tag (Supplementary Fig. 9). Stable transgenic *opr2opr3* lines

complemented with the organelle variants of OPR3ΔSRL fused to YFP were generated and

resulted in clear subcellular localizations of OPR3-YFP to the nucleus, cytosol, chloroplast

stroma, mitochondria, and ER (Fig. 5b). As tested by RT-qPCR, all transgenic lines exhibited a

similar level of transgene expression (Supplementary Fig. 10).

The main phenotypic feature of JA-deficient and JA-insensitive Arabidopsis mutants is the male sterility, which is also characteristic for *opr2opr3* plants and is visible by defects in filament elongation, pollen release and seed set [41]. Plants of *opr2opr3* transformed with an empty vector (EV) showed short stamens with non-dehiscing anthers (Fig. 6a, b), similar to the reported male sterility phenotype of the *opr3* mutant [38]. As a positive control, the complementation of *opr2opr3* with the peroxisome targeted OPR3 showed full rescue of the flower phenotype (Fig. 6c). Interestingly, all other tested organelle-variants of OPR3 also rescued the flower sterility and showed open flowers like the wild-type with elongated stamens and anthers releasing pollen (Fig. 6d-h). This rescue of sterility translated also in the ability of the lines to form siliques containing seeds. The only exception were plants complemented with the ER-targeted OPR3 having partial rescue of flower fertility only. These plants formed fewer and smaller siliques compared to the wild type (Fig. 6f). This suggests that OPR3 localized to the ER may have restricted access to its substrate, potentially limiting JA/JA-Ile biosynthesis compared to other subcellular compartments.

Reconstitution of the JA pathway in vegetative tissue of the *opr2opr3* complementation lines was analyzed by determination of wound-induced levels of OPDA, JA and JA-Ile in two-week-old seedlings. Hormone levels of *opr2opr3* complemented with OPR3 targeted to either peroxisomes, cytosol or mitochondria were compared to those of wild-type and *opr2opr3* transformed with an empty vector. Data from two independent transformed lines each revealed that complementation of *opr2opr3* with OPR3 restored the JA/JA-Ile levels to wild-type levels (Fig. 6i). These results show the reconstitution of JA biosynthesis by OPR3 localized to all the tested compartments.

Trans-organellar complementation with OPR2 and OPR1 to determine whether OPDA or 4, 5-ddh-JA is the translocating substrate.

JA biosynthesis in *opr2opr3* complemented with OPR3 located in different organelles would occur only if its substrate, OPDA, is able to translocate to these compartments. However, as an alternative pathway, OPDA could enter directly the β-oxidation as reported recently [43]. This would result in formation of 4,5-ddh-JA, which might be metabolized by OPR3 irrespective of the organellar location. Conversion of 4,5-ddh-JA to JA by OPR3 was tested by *in vitro* enzymatic assay and showed that OPR3 reduces 4,5-ddh-JA *in vitro*, however to a lesser extent than OPDA (Supplementary Fig. 11). Based on substrate consumption after 10 min, only half of the given 4,5-ddh-JA was reduced while OPDA was completely converted to 3-oxo-2-(20(Z)-pentenyl)-cyclopentane-1-octanoic (OPC-8). The control reactions with inactivated enzyme showed no reduction activity. The results clearly show that OPR3 is able to reduce both cyclopentenones, 4,5-ddh-JA and OPDA, *in vitro*. In addition, it has been shown by recombinant enzymatic assays that OPR2 but not OPR1 is able to reduce 4,5-ddh-JA [43]. Conversely, OPR1 is able to reduce OPDA although with very low efficiency but OPR2 is not [21]. To determine whether OPDA or 4,5-ddh-JA is the substrate able to translocate to the tested cell compartments and specifically to cell compartments unrelated to the location of JA biosynthesis, peroxisomal, cytosolic and mitochondrial variants of OPR1 and OPR2, both fused to YFP, were used to complement the *opr2opr3* mutant. Both OPRs localized to the cytosol, whereas addition of a C-terminal SKL and an N-terminal mitochondrial target sequence led to localization in peroxisomes and mitochondria, respectively (Fig. 7a). All three variants expressing OPR2 rescued the flower sterility and showed open flowers like the wild-type with elongated stamens and anthers releasing pollen (Fig. 7b) as well as formation of seed-containing, fully developed siliques (Fig. 7c). In contrast, expression of OPR1 located to peroxisomes was the only OPR1 variant leading to a

rescue of the flower fertility, whereas targeting OPR1 to the cytosol led to partial rescue of flower

fertility only and targeting OPR1 into the mitochondria did not restore the fertility (Fig. 7b, c).

These data were supported by the fact that only seedlings harboring the OPR2 variants and

peroxisomal-located OPR1 were able to accumulate JA and JA-Ile upon wounding (Fig. 7d),

although all lines accumulated similar levels of OPDA (Supplementary Figure 12). The JA/JA-Ile

levels were lower than those of wounded wild-type seedlings, but the significantly increased

levels in comparison to *opr2opr3* transformed with the empty vector showed the capability of

these plants to produce JA/JA-Ile. Most importantly, however, expression of OPR1 located to

mitochondria did not restore fertility nor JA/JA-Ile levels. Since OPR1 does not convert 4,5-ddh-

JA to JA, this result points to the fact that 4,5-ddh-JA but not OPDA might be the compound to

be converted by OPR2 or OPR3 in different organelles.

Discussion

JA plays a major role in growth and defense against pathogens, herbivory attacks and mechanical

wounding. Its signaling through the bioactive JA-Ile is well established in Angiosperms, while in

non-vascular plants, such as *M. polymorpha*, OPDA, *dn-iso*-OPDA and Δ^4-*dn-iso*-OPDA are

bioactive[13,24-26,51]. Given that *A. thaliana* plants produce OPDA and *dn*-OPDA, possible function

of OPDA as a signaling molecule independently from JA was proposed. OPDA was shown to

induce a distinct transcriptional change from JA when exogenously applied to Arabidopsis plants

[32,45]. Additionally, OPDA´s involvement in processes like seed germination and stomatal closure

was demonstrated [29-31]. The mechanism by which OPDA exerts these functions in vascular plants

is, however, still elusive [17]. A major genetic background used for studying OPDA functions was

the *opr3* mutant [41]. Recently, an OPR3-independent pathway of JA formation through the

cytosolic isoenzyme OPR2 has been identified [43]. Hence, uncoupling of JA production from

OPDA was not achieved in the *opr3* mutant, and the characterized functions of OPDA cannot be

attributed to OPDA with certainty. Here, we addressed the question of whether OPDA functions

independently from JA as a signaling molecule in Arabidopsis by utilizing the *opr2opr3* mutant,

which does not produce JA/JA-Ile upon wounding [43].

The use of the *opr2opr3* mutant is limited by its lower production of OPDA in

comparison to the wild type due to the disrupted JA positive feedback loop as indicated by

reduced protein levels of the biosynthesis enzymes, such as AOC [1]. The enhancement of the

AOC levels and subsequent OPDA production in the JA-deficient mutants was achieved by

restituting the JA feedback loop through a repetitive supply of MeJA to seedlings during

development (Fig. 1). This result validated the role of the JA positive feedback loop in the

regulation of the basal levels of the biosynthetic enzymes and setting the mutant ability to

produce OPDA to similar levels as wild-type plants do [44].

The wounding response of wild-type seedlings involves JA-dependent and independent

processes.

To inspect the separate roles of OPDA and JA upon wounding, a comparative analysis of the

transcriptomes of seedlings of wild type, *opr2opr3* and *aos* was carried out. The transcriptomic

change induced by wounding in wild-type seedlings involved an up-regulation of genes

previously identified as JA-dependent or associated with immune responses, validating the

specific signaling role of JA-Ile in the wound response of seedlings as it has been described for

mature leaves of Arabidopsis and other plant species [5,52-54]. Unexpectedly, seedlings of the *aos*

mutant being deficient in synthesis of JA and OPDA, also showed a major transcriptional change

upon wounding that was common to the *opr2opr3* mutant and the wild type (Fig. 2). The

transcriptional response was significantly enriched in general stress response pathways, with an

induction of oxidative stress- and transcription-related genes. This implies that JA-independent

signaling processes occur in response to mechanical wounding leading to transcriptional

reprogramming. Among such processes, water stress might contribute to the regulation of wound-

responsive genes in a JA-independent manner [55]. Ethylene and abscisic acid (ABA) also

contribute to the wound response by regulating photosynthesis and drought responsive genes,

respectively [56]. An ABA-dependent and JA-independent transcriptional regulation of wax

biosynthesis to seal the wounded sites of Arabidopsis leaves has been recently characterized, but

appears to be controlled post-translationally by JA [57]. As plants defective in JA/JA-Ile

biosynthesis or perception are severely diminished in their defense response [41,58-60], the JA-

independent wound response has rather limited contribution to plant defense and might

predominantly mitigate the adverse effects of wounding itself by promoting wound healing and

tissue regeneration [61].

An OPDA-specific transcriptional response is missing in wounded *opr2opr3* seedlings, but

specific pathways are activated by the exogenous application of OPDA.

The general stress response that was both JA and OPDA-independent, comprised genes

that were previously identified as OPDA-specific response genes, such as *DREB2A*, *FAD-OXR*,

ZAT10, ERF5, TCH4, and *GST6* [32]. These genes did not show differential expression when the

transcriptomes of *opr2opr3* and *aos* were compared indicating that their induction by wounding

is OPDA-independent (Fig. 2). Moreover, these genes were induced by wounding also in leaves

of two *coi1*-mutant lines excluding any induction by JA/JA-Ile possibly caused by an insufficient

JA-deficiency of the mutants. Several abiotic stresses were reported to up-regulate the same

genes [62-65], further indicating their independence from OPDA. Moreover, direct comparison of

transcriptomes from wounded *opr2opr3* and *aos* seedlings with standard cut-off parameters did

not yield a significant list of putative OPDA-regulated genes further confirming the absence of an

OPDA signaling (Fig. 3). This could be due to three facts: (i) A putative receptor for OPDA

might be missing in *A. thaliana*, (ii) OPDA is converted to 4,5-ddh-JA, which does not

accumulate to levels having signaling capacity [43], and (iii) OPDA remains largely sequestered in

the plastids by its biosynthetic enzymes [66]. Regarding perception of OPDA, evolutionary studies

of the JA-Ile co-receptor COI1 showed that the Arabidopsis protein does not perceive OPDA or

dn-OPDA, dismissing possible signaling of OPDA through COI1 in this plant [13]. Moreover, it has

been shown that *dn-iso*-OPDA rather than *dn*-OPDA is the bioactive molecule in mosses, and

that this compound does not occur in *A. thaliana* [26]. The transcriptomic data obtained from

opr2opr3, *aos*, *coi1-16* and *coi1-30* therefore point to the fact that OPDA or its homolog *dn-*

OPDA are not perceived by COI1 or any other receptor in Arabidopsis.

The transcriptional signature of OPDA was predominantly observed in experiments

relying on its exogenous application [32,35,45]. Therefore, we compared the transcriptional changes

following application of OPDA to *opr2opr3* seedlings with those by wound-induced,

endogenously produced OPDA. This comparison revealed an up-regulation of sulfur assimilation

and GSH production by OPDA being attributed as specific signaling functions of OPDA [35]. Here,

the transcriptional activation of sulfur metabolism pathway was exclusive to the exogenous

OPDA supply, contradicting an intrinsic signaling function of the endogenous OPDA (Fig. 4). As

an electrophilic species, OPDA was shown to disturb redox homeostasis [67] and to inhibit

photosynthesis by affecting the photosystem II fluorescence [33,68]. In parallel, the *in planta*

conjugation of OPDA to GSH, which is an important antioxidant substance, and its degradation

in the vacuole were also demonstrated [47]. This implies that OPDA may not directly regulate the

sulfur assimilation pathway but instead triggers this response through its potential detoxification

when exogenously applied to plants, suggesting that the observed transcriptional response to

exogenously applied OPDA could be a side effect rather than a direct effect of OPDA itself. The

main route for the conversion of applied OPDA seems to be, however, its direct conversion to

4,5-ddh-JA, which then reaches levels up to 4000 pmol per g FW (Ueda et al., accompanying

manuscript). Moreover, Ueda et al. showed convincingly that OPDA applied to Arabidopsis

seedlings does not serve as genuine bioactive form responsible for the expression of OPDA-

marker genes. As shown by Ueda et al., 4,5-ddh-JA is inducing the expression of OPDA-specific

response genes in a dose-dependent manner, possibly due to its electrophilic properties. Upon

wounding of *opr2opr3*, however, 4,5-ddh-JA accumulates up to 500 pmol per g FW [43], a level

that does not seem to be sufficient to induce a specific gene expression (Ueda et al.,

accompanying manuscript). Here, the general stress response might be the main factor for

transcriptional changes including the previously identified OPDA-marker genes as depicted from

the wound response of *aos* mutant plants. These genes, being responsive to several abiotic

stresses [62-64], including wounding in a JA-independent manner (as shown in this manuscript),

further suggest their unspecific response to OPDA. This rather indicates that applying

electrophilic compounds, such as 4,5-ddh-JA or OPDA, causes oxidative stress, resulting in the

induction of general stress-responsive genes. Next to jasmonates, there are other compounds

known that mediate at least a part of the wound response, such as reactive oxygen species (ROS)

[69] and γ-aminobutyric acid (GABA) [70]. These JA-independent responses might be necessary for

wound-healing, which seems to be a process independent on the JA-dependent defense response

[71].

Endogenously formed OPDA is sequestered to the organelles of JA biosynthesis

To address the question whether wound-induced OPDA remains largely sequestered in the
plastids or whether it is converted to 4,5-ddh-JA similarly to the applied OPDA (Ueda et al.,
accompanying manuscript), a trans-organellar complementation approach was performed by
directing OPR3 into various organelles (Figs. 5, 6, Supplementary Fig. 9). Occurrence of OPR3
in any of the tested organelles resulted in a complementation of the phenotype and JA/JA-Ile
production upon wounding in *opr2opr3* mutants. This led to the assumption that OPDA might
travel throughout the cell. Nevertheless, the possibility still exists that OPDA is translocated
rapidly via the cytosol to the peroxisomes, where it is converted to 4,5-ddh-JA by β-oxidation as
shown previously for *opr3* and *opr2opr3* [43]. Given that OPR3 is able to convert 4,5-ddh-JA to JA
and OPDA to OPC-8 *in vitro* (Supplementary Fig. 10), the complementation approach with
OPR3 could not discriminate between the presence of either OPDA or 4,5-ddh-JA outside of the
plastids or the peroxisomes. To address this, organelle variants of the isoenzymes OPR2 and
OPR1 were used to complement the *opr2opr3* mutant (Fig. 6). OPR2 and OPR1 differ in their
enzymatic activity against OPDA and 4,5-ddh-JA: OPR2 is able to convert both compounds, but
OPR1 has been shown to be inactive on 4,5-ddh-JA [43]. Since OPDA and 4,5-ddh-JA should occur
in peroxisomes and in the cytosol, reconstitution of JA biosynthesis was obtained using either
OPR1 or OPR2 targeted to peroxisomes or the cytosol. Targeting OPR1 or OPR2 to
mitochondria, however, discriminated whether OPDA or 4,5-ddh-JA is the substrate able to
translocate to other organelles (Fig. 8). Here, complementation of *opr2opr3* with a
mitochondrial-located OPR1 did not rescue fertility and JA/JA-Ile production (Fig. 7), whereas
OPR2 in mitochondria did. Additionally, OPR1 fully complemented the mutant only within the
peroxisome, and its cytosolic variant resulted in minimal JA-Ile formation and partial rescue of
fertility. These results suggest a limited occurrence of endogenously formed OPDA in the cytosol
and its preferred compartmentalization in plastids and peroxisomes, whereas 4,5-ddh-JA might

be the substrate able to translocate between cell organelles. The possible translocation of 4,5-ddh-

JA correlates with its potential role in signaling or its effect as an electrophilic compound, as

demonstrated by Ueda et al. (accompanying manuscript).

In conclusion, despite its perception and signaling in non-vascular plants, OPDA itself does not

mediate transcriptional changes in Arabidopsis. The induction of a specific transcriptional change

occurred exclusively upon exogenous supply but not upon endogenous rise of OPDA, although in

both cases OPDA is (partially) converted to 4,5-ddh-JA ([43] and Ueda et al., accompanying

manuscript). Here, the endogenous levels of 4,5-ddh-JA – being low upon wounding, but high

upon exogenously applied OPDA – might be determining, whether it acts as an electrophilic

(toxic) compound inducing expression of stress-related genes. In turn, the data obtained after

wounding suggest that the endogenous OPDA is compartmentalized and thereby its level in the

cytosol highly regulated. The discrepancies between the effects of endogenously produced and

exogenously supplied OPDA suggest an absence of signaling per se and/or a tightly regulated

compartmentalization to prevent its effects as an electrophilic species. Both scenarios imply that

OPDA does not inherently function as a signaling compound in Arabidopsis, unless such

functionality manifests under specific, yet uncharacterized conditions. It is plausible that OPDA-

oxidation by JASMONTE-INDUCED DIOXYGENASE1 (JID1) [72], or conjugation to amino

acids [73,74] or GSH [37] contribute to the regulation of the cytosolic OPDA level.

Material and methods

Plant material, growth conditions, and treatment

Arabidopsis thaliana wild-type (ecotype Col-0) and the mutant lines *opr2opr3* [43], *aos* (*dde2-2* [40],

coi1-16 [75] and *coi1-30* [76] were genotyped using the primers listed (Supplementary Tab. 6). After

surface sterilization with 4% bleach and three-day stratification at 4°C, seedlings were grown for

ten days in liquid Murashige and Skoog (MS) medium (pH 5.7, 1% sucrose), whereas adult

plants were grown for four weeks in individual pots containing steam-sterilized clay, coir fiber,

and vermiculite. Plant growth took place in Phytocabinets (Percival Scientific, www.percival-

scientific.com/) at 120 µE m^{-2} s^{-1} under short day conditions (10/14 hours light/dark cycle) at

21/19 °C and 65% relative humidity. For wounding, seedlings were squeezed with forceps with

serrated teeth. Phytohormone treatments were done by adding either 25 µM OPDA ($\geq$ 95%,

Cayman Chemical, www.caymanchem.com) or 1 µM MeJA to the medium.

RNA isolation, RNA-seq and quantitative RT-PCR analysis

Total RNA was isolated from homogenized frozen material using the RNeasy Plant Mini Kit

(Qiagen) and treated with the DNA-free™ DNA Removal Kit (Invitrogen, #AM1906). First

strand cDNA synthesis from 1 µg DNA-free RNA was carried out using the RevertAid H Minus

reverse transcriptase with oligo(dT)18 primers (ThermoFisher Scientific™,

www.thermofisher.com). Quantitative PCR was carried-out in Hard-Shell® 96-Well PCR Plates

(Bio-Rad Laboratories, www.bio-rad.com, #HSP9601) supplied with 10 µL reaction mix of 1.5

ng/µL cDNA, 1x EvaGreen QPCR Mix II (Bio&Sell, www.bio-sell.de, #BS76.580.0200) and 0.2

µM of forward and reverse primers (Supplementary Tab. 6). The reactions were run on a CFX

Connect Real-Time PCR Detection System (Bio-Rad Laboratories) with denaturation (95°C for

15 min), amplification (40 cycles of 95°C for 15 s and 60°C for 30 s) and melt curve analysis

(95°C for 10 s, 65°C heating up to 95°C with a heating rate of 0.05°C s^{-1}). Gene expression was

normalized to the housekeeping gene *PROTEIN PHOSPHATASE 2A SUBUNIT A3*

(AT1G13320) [77,78] using the $2^{-\Delta CT}$ method [79] and included biological triplicates.

Transcriptome analysis

RNA quality and integrity were assessed on Agilent 2100 Bioanalyzer system (Agilent Technologies, www.agilent.com) using the RNA 6000 Nano Kit for standard RNA sensitivity (Agilent, #5067-1511). Three biological replicates were submitted to Novogene (www.novogene.com) for mRNA paired-end short-read sequencing (150 bp length) on an Illumina NovaSeq 6000 Sequencing System and bioinformatics analysis according to their pipeline. Reads were mapped to the *A. thaliana* TAIR10 reference genome. Gene expression levels were determined using the FPKM (Fragments per kilobase per million) method. Gene expression heatmaps, principal component analysis (PCA), gene clustering, and Gene Ontology (GO) enrichment analysis were generated using the iDEP.96 (http://bioinformatics.sdstate.edu/idep96/), iDEP 1.1 (http://bioinformatics.sdstate.edu/idep11/) and ShinyGO 0.80 (http://bioinformatics.sdstate.edu/go/) software tools [80,81].

Protein extraction and immunoblotting

Proteins were extracted by incubation of 50 mg frozen material in 200 µl of extraction buffer (25 mM Tris-Cl pH 6.8, 1% SDS, 1% [v/v] β-mercaptoethanol) at 95°C for ten minutes. Extracts were treated with 1% (v/v) Halt™ Protease and Phosphatase Inhibitor Cocktail (ThermoFisher Scientific™, #78440) and quantified at 595 nm on the SPARK® multimode microplate reader (TECAN) using the Pierce™ Bradford Plus Protein Assay Kit (ThermoFisher Scientific™, #23236). Ten micrograms of total protein were incubated at 96°C for 10 min in Laemmli Buffer (1:1, [82] to dissolve AOC trimers [83]. Proteins were separated on SDS–PAGE (4% and 12% acrylamide for stacking and resolving gels, respectively), transferred to a PVDF membrane and detected using Ponceau S staining or No-Stain™ Protein Labeling Reagent (ThermoFisher Scientific™, #A44449). Membrane blocking in 5% (w/v) BSA in TBST (20 mM Tris–Cl pH 7.8, 150 mM NaCl, 0.05% [v/v] Tween) was followed by immuno-staining using anti-AtAOC

primary antibody (1:5000, [44] and a goat anti-rabbit IgG secondary antibody conjugated with

alkaline phosphatase (1:4000, Chemicon®, Sigma-Aldrich, www.sigmaaldrich.com, #AP307P).

Detection by chemiluminescence with Immun-Star AP Substrate (ThermoFisher Scientific,

#1705018) was visualized using a Fusion FX Imaging system (Vilber, www.vilber.com). AOC

protein bands were quantified relative to total protein loadings using densitometry analysis with

ImageJ (https://imagej.nih.gov/ij/index.html).

Cloning of the OPRs and plant transformation

The OPR3 (1176 bp), OPR2 (1125 bp), and OPR1 (1194 bp) coding sequences (CDS) without

stop codons were PCR-amplified from Col-0 cDNA using the Q5® High-Fidelity DNA

Polymerase (Bio Labs, www.biolabs.io, #M0491) and the listed primer pairs (Supplementary

Tab. 7). For OPR3ΔSRL, the last nine nucleotides were removed, and silent mutations C205G

and G1109C were introduced to eliminate *BsaI* and *BpiI* sites, respectively. The CDS were

cloned into the Golden Gate level 0 vectors pAGM1287 and pAGM1299 [84]. OPR fusions to YFP

were generated by in-frame assembly of the level 0 with YFP at the C-terminus connected by a

Gly-Ser linker. The final cloning cassettes containing the CaMV *35S* promoter and *tOcs*

terminator in addition to the Oleosin-RFP plant selection marker were assembled in the

pAGM55171 vector.

Plant stable transformation was carried-out by floral dip with *Agrobacterium tumefaciens* strain

GV3101 according to [85]. To circumvent the *opr2opr3* mutant sterility, MeJA was applied to

flower buds from 48 hours post-dipping until the first silique formation. Transformant selection

relied on the seed coat RFP marker and single-insertion lines were identified by segregation

analysis determining the ratio of fluorescent to non-fluorescent seeds.

Subcellular targeting of the OPRs

Organelle variants of OPR3, OPR2, and OPR1 were generated by adding organelle targeting

signals to the N- and/or C-terminal regions of the OPRs and/or YFP sequences (Supplementary

Tab. 5). For nuclear targeting, OPR3ΔSRL was fused to the nuclear localization signal from

Simian Virus 40 [86,87], while its targeting to the cytosol relied on a C-terminal nuclear export

signal from the HIV1 REV [88]. ER-localized OPR3ΔSRL was generated by flanking the sequence

between an N-terminal signal peptide from the AtCAPE3 [89] and a C-terminal HDEL retention

signal [90]. Import to the chloroplast stroma and mitochondria, relied on an N-terminal target

peptide from the AtRECA1 [91] and the Rieske protein [92], respectively.

Transient expression in *N. benthamiana* leaves and protoplasts.

A. tumefaciens strain *GV3101*, harboring OPR constructs, was cultured for 48 hours in LB

medium with the appropriate antibiotics. The bacteria were syringe-infiltrated into *N.*

benthamiana leaves according to [93]. Leaf disc imaging was performed 48-72 hours post-

infiltration.

Protoplasts were isolated from leaves of 4-week-old *N. benthamiana* plants, as described [94].

Isolated protoplasts were co-transfected with level 1 plasmids of OPR3 constructs and organelle

markers [50] using a mixture of 200 µl of protoplast suspension and 10 µg of DNA. Protoplasts

were incubated at room temperature overnight after PEG-transformation and used for imaging.

Confocal microscopy

Confocal microscopy utilized LSM880 and LSM900 laser scanning microscopes (Zeiss

Germany, www.zeiss.de). Fluorophores were excited with 514 nm (YFP) and 561 nm (mCherry)

laser lines and detected at 510–560 nm and 570-650 nm, respectively. Image acquisition and

processing were performed using ZEN software (version 3.4, 2021, Zeiss).

Phytohormone measurements

Measurements of OPDA, *dn*-OPDA, JA, and JA-Ile were performed using a standardized Ultra-

performance liquid chromatography–tandem Mass Spectrometry (UPLC– MS/MS)-based method

[95]. 50 mg of powdered frozen tissue were extracted with 500 µl 100% LC-MS methanol

supplemented with 50 ng of $[^2H_5]$OPDA, $[^2H_6]$JA, and $[^2H_2]$JA-Ile each as internal standards.

After solid-phase extraction on HR-XC (Chromabond, Macherey-Nagel, www.mn-net.com), 10

µl of the eluate were analyzed via UPLC–MS/MS, and analyte content was determined relative to

the internal standard peak heights. *dn*-OPDA content was determined using $[^2H_5]$OPDA as the

internal standard.

Determination of enzymatic activity of OPR3

OPR3 CDS inserted in pET28a(+) with *EcoR*I and *Not*I cloning sites was ordered from BioCat

GmbH (www.biocat.com) and transformed into *Escherichia coli* BL21 Star (DE3) cells

(ThermoFisher Scientific) by heat shock. Protein expression was performed in auto-induction

medium at 16°C for three days [96]. Harvested cells were resolved in buffer HisA (50 mM Tris/HCl

pH 7.8, 100 mM NaCl, 10% (v/v) glycerol, 2 mM $MgCl_2$ and 5 mM imidazole) supplemented

with lysozyme, DNAse I and 0.2 mM PMSF. After a 30-minute incubation on ice, cells were

sonicated and cleared lysate was applied to 1 mL HisTrap FastFlow column (Cytiva,

www.cytivalifesciences.com). Protein was eluted with buffer HisB (50 mM Tris/HCl pH 7.8,

100 mM NaCl, 10% (v/v) glycerol, 2 mM $MgCl_2$ and 500 mM imidazole), which was changed to

storage buffer (50 mM Tris/HCl pH 7.8, 100 mM NaCl, 10% (v/v) glycerol, 2 mM $MgCl_2$) by

dialysis (MWCO 12-14 kDa) at 4°C overnight. Following SDS-PAGE, protein fractions were

used for *in vitro* assays at 0.1 mg/mL together with 0.05 mM substrate, 1 mM NADPH and

10 mM NaCl in 20 mM Tris/HCl pH 7.8. Reactions were incubated at 30°C for 10 minutes and

stopped by one volume acetonitrile. Enzyme deactivated by one volume acetonitrile was used as

control. Assays were analyzed via UHPLC-high resolution (HR)-MS according to [97]. Data was

analyzed using the MassHunter Qualitative Analysis 10.0 software (Agilent Technologies).

Figures and statistical analysis

635   Micrographs and photographs were processed through PHOTOSHOP 12.0.4 (Adobe Systems,

636   http://www.adobe.com). The statistical analyses applied to the different datasets were performed

637   using GraphPad Prism (www.graphpad.com).

Data availability

The datasets generated and/or analyzed during this study are available from the corresponding author on request. Transcriptomic raw data were deposited in Sequence Read Archive (SRA) of NCBI (PRJNA1088739).

Funding

K.M. and B.H. were supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) grant No 400681449/GRK2498. M.K. was supported by the DFG grant No 273134146/GRK 2172. I.F. acknowledges funding from the DFG (GRK 2172-PRoTECT, INST 186/1434-1, and ZUK 45/2010).

Author contributions

Conceptualization, B.H and K.M.; Methodology, K.M, R.B, and F.S.; Investigation, K.M., R.B., F.S., and M.K.; Analyzing Data, K.M., I.F. and B.H.; Writing – Original Draft, K.M.; Writing – Review & Editing, B.H., K.M., and I.F.; Funding Acquisition, B.H. and I.F.;

Acknowledgments

We thank Hagen Stellmach (IPB Halle, Germany) for help in jasmonate quantification, Andreas Schaller (University of Hohenheim, Germany) and Mats Hamberg (Karolinska Institute, Sweden) for providing seeds of *opr2opr3* and 4,5-ddh-JA, respectively. We thank Sylvestre Marillonnet (IPB Halle) for providing the Golden Gate modules and vectors used in this study. Khabat

659   Vahabi (IPB Halle) is acknowledged for assistance in RNAseq preparation and data analysis.

660   Claus Wasternack (IPB Halle) and Alain Tissier (IPB Halle) are acknowledged for critical

661   reading of the manuscript. We thank Prof. Minoru Ueda (Tokhoku University, Japan) for sharing

662   his data and helpful discussions on our manuscript.

664   Declaration of interests

665   The authors declare no competing interests.

References

1   Wasternack, C. & Hause, B. Jasmonates: biosynthesis, perception, signal transduction and

action in plant stress response, growth and development. An update to the 2007 review in

Annals of Botany. *Ann. Bot.* **111**, 1021-1058 (2013). https://doi.org:10.1093/aob/mct067

2   Wang, J., Wu, D., Wang, Y. & Xie, D. Jasmonate action in plant defense against insects.

J. Exper. Bot. **70**, 3391-3400 (2019). https://doi.org:10.1093/jxb/erz174

3   Li, C. *et al.* Jasmonate signaling pathway modulates plant defense, growth, and their

trade-offs. *Int. J. Mol. Sci.* **23**, 3945 (2022).

4   Zhang, G. *et al.* Roles of the wound hormone jasmonate in plant regeneration. *J. Exper.*

Bot. **74**, 1198-1206 (2023). https://doi.org:10.1093/jxb/erab508

5   Koo, A. J. K. & Howe, G. A. The wound hormone jasmonate. *Phytochemistry* **70**, 1571-

1580 (2009).

6   Wasternack, C. New light on local and systemic wound signaling. *Trends Plant Sci.* **24**,

102-105 (2019). https://doi.org:https://doi.org/10.1016/j.tplants.2018.11.009

7   Xie, D.-X., Feys, B., James, S., Nieto-Rostro, M. & Turner, J. COI1: An Arabidopsis

gene required for jasmonate-regulated defense and fertility. *Science* **280**, 1091-1094

(1998).

8   Xu, L. *et al.* The SCF-coi1 ubiquitin-ligase complexes are required for jasmonate

response in Arabidopsis. *Plant Cell* **14**, 1919-1935 (2002).

9   Chini, A. *et al.* The JAZ family of repressors is the missing link in jasmonate signalling.

Nature **448**, 666-671 (2007).

10   Thines, B. *et al.* JAZ repressor proteins are targets of the SCF[COI1] complex during

jasmonate signalling. *Nature* **448**, 661-665 (2007).

11   Sheard, L. B. *et al.* Jasmonate perception by inositol-phosphate-potentiated COI1-JAZ co-
690   receptor. *Nature* **468**, 400-405 (2010).

691   12   Yan, J. *et al.* Dynamic perception of jasmonates by the F-box protein COI1. *Mol. Plant*
692   **11**, 1237-1247 (2018). https://doi.org:10.1016/j.molp.2018.07.007

693   13   Monte, I. *et al.* JAZ is essential for ligand specificity of the COI1/JAZ co-receptor. *Proc.*
694   *Natl.   Acad.   Sci.   USA   **119**,   e2212155119   (2022).*
695   https://doi.org:doi:10.1073/pnas.2212155119

696   14   Dave, A. & Graham, I. A. Oxylipin signalling: a distinct role for the jasmonic acid
697   precursor *cis*-(+)-12-oxo-phytodienoic acid (*cis*-OPDA). *Front. Plant Sci.* **3:42** (2012).
698   https://doi.org:10.3389/fpls.2012.00042

699   15   Wasternack, C. & Strnad, M. Jasmonate signaling in plant stress responses and
700   development – active and inactive compounds. *New Biotechnol.* **33**, 604-613 (2016).
701   https://doi.org:http://dx.doi.org/10.1016/j.nbt.2015.11.001

702   16   Liu, W. & Park, S.-W. 12-oxo-Phytodienoic acid: A fuse and/or switch of plant growth
703   and   defense   responses?   *Front.   Plant   Sci.*   **12**   (2021).
704   https://doi.org:10.3389/fpls.2021.724079

705   17   Jimenez-Aleman, G. H., Thirumalaikumar, V. P., Jander, G., Fernie, A. R. & Skirycz, A.
706   OPDA, more than just a jasmonate precursor. *Phytochemistry* **204**, 113432 (2022).
707   https://doi.org:https://doi.org/10.1016/j.phytochem.2022.113432

708   18   Feussner, I. & Wasternack, C. The lipoxygenase pathway. *Annu. Rev. Plant Biol.* **53**, 275-
709   297 (2002).

710   19   Andreou, A. & Feussner, I. Lipoxygenases - Structure and reaction mechanism.
711   *Phytochemistry* **70**, 1504-1510 (2009).

20 Bannenberg, G., Martínez, M., Hamberg, M. & Castresana, C. Diversity of the enzymatic

activity in the lipoxygenase gene family of *Arabidopsis thaliana*. *Lipids* **44**, 85-95 (2009).

https://doi.org:10.1007/s11745-008-3245-7

21 Schaller, F., Biesgen, C., Müssig, C., Altmann, T. & Weiler, E. W. 12-Oxophytodienoate

reductase 3 (OPR3) is the isoenzyme involved in jasmonate biosynthesis. *Planta* **210**,

979-984 (2000).

22 Schaller, A. & Stintzi, A. Enzymes in jasmonate biosynthesis - Structure, function,

regulation. *Phytochemistry* **70**, 1532-1538 (2009).

23 Weber, H., Vick, B. & Farmer, E. Dinor-oxo-phytodienoic acid: A new hexadecanoid

signal in the jasmonate family. *Proc. Natl. Acad. Sci. USA* **94**, 10473-10478 (1997).

24 Monte, I. *et al.* Ligand-receptor co-evolution shaped the jasmonate pathway in land

plants. *Nature Chem. Biol.* **14**, 480-488 (2018). https://doi.org:10.1038/s41589-018-0033-

4

25 Monte, I. *et al.* An ancient COI1-independent function for reactive electrophilic oxylipins

in   thermotolerance.   *Curr.   Biol.   **30**,   962-971.e963   (2020).

https://doi.org:10.1016/j.cub.2020.01.023

26 Chini, A., Monte, I., Zamarreño, A. M., García-Mina, J. M. & Solano, R. Evolution of the

jasmonate ligands and their biosynthetic pathways. *New Phytol.* **238**, 2236-2246 (2023).

https://doi.org:https://doi.org/10.1111/nph.18891

27 Stelmach, B. A. *et al.* Quantitation of the octadecanoid 12-oxo-phytodienoic acid, a

signalling compound in plant mechanotransduction. *Phytochemistry* **47**, 539-546 (1998).

28 Goetz, S. *et al.* Role of cis-12-oxo-phytodienoic acid in tomato embryo development.

Plant Physiol. **158**, 1715-1727 (2012). https://doi.org:10.1104/pp.111.192658

29   Dave, A. *et al.* 12-Oxo-phytodienoic acid accumulation during seed development repesses seed germination in Arabidopsis. *Plant Cell* **23**, 583-599 (2011). https://doi.org:10.1105/tpc.110.081489

30   Savchenko, T. & Dehesh, K. Drought stress modulates oxylipin signature by eliciting 12-OPDA as a potent regulator of stomatal aperture. *Plant Signal. Behavior* **9**, e28304 (2014).

31   Chang, Y. *et al.* Light-induced stomatal opening in Arabidopsis is negatively regulated by chloroplast-originated OPDA signaling. *Curr. Biol.* **33**, 1071-1081.e1075 (2023). https://doi.org:https://doi.org/10.1016/j.cub.2023.02.012

32   Taki, N. *et al.* 12-oxo-phytodienoic acid triggers expression of a distinct set of genes and plays a role in wound-induced gene expression in Arabidopsis. *Plant Physiol.* **139**, 1268-1283 (2005).

33   Alméras, E. *et al.* Reactive electrophile species activate defense gene expression in Arabidopsis. *Plant J.* **34**, 205-216 (2003).

34   Davoine, C. *et al.* Adducts of oxylipin electrophiles to glutathione reflect a 13 specificity of the downstream lipoxygenase pathway in the tobacco hypersensitive response. *Plant Physiol.* **140**, 1484-1493 (2006).

35   Park, S.-W. *et al.* Cyclophilin 20-3 relays a 12-oxo-phytodienoic acid signal during stress responsive regulation of cellular redox homeostasis. *Proc. Natl. Acad. Sci. USA* **110**, 9559-9564 (2013). https://doi.org:10.1073/pnas.1218872110

36   Müller, S. M. *et al.* The redox-sensitive module of cyclophilin 20-3, 2-cysteine peroxiredoxin and cysteine synthase integrates sulfur metabolism and oxylipin signaling in the high light acclimation response. *Plant J.* **91**, 995-1014 (2017). https://doi.org:10.1111/tpj.13622

37   Adhikari, A. & Park, S.-W. Reduced GSH acts as a metabolic cue of OPDA signaling in

coregulating photosynthesis and defense activation under stress. *Plants* **12**, 3745 (2023).

38   Stintzi, A. & Browse, J. The Arabidopsis male-sterile mutant, *opr3*, lacks the 12-

oxophytodienoic acid reductase required for jasmonate synthesis. *Proc. Natl. Acad. Sci.*

USA **97**, 10625-10630 (2000).

39   Stintzi, A., Weber, H., Reymond, P., Browse, J. & Farmer, E. Plant defense in the absence

of jasmonic acid: The role of cyclopentanones. *Proc. Natl. Acad. Sci. USA* **98**, 12837-

12842 (2001).

40   von Malek, B., van der Graaff, E., Schneitz, K. & Keller, B. The Arabidopsis male-sterile

mutant *dde2-2* is defective in the *ALLENE OXIDE SYNTHASE* gene encoding one of the

key enzymes of the jasmonic acid biosynthesis pathway. *Planta* **216**, 187-192 (2002).

41   Browse, J. The power of mutants for investigating jasmonate biosynthesis and signaling.

Phytochemistry **70**, 1539-1546 (2009).

42   Chehab, E. W. *et al.* Intronic T-DNA insertion renders Arabidopsis *opr3* a conditional

jasmonic   acid-producing   mutant.   *Plant   Physiology*   **156**,   770-778   (2011).

https://doi.org:10.1104/pp.111.174169

43   Chini, A. *et al.* An OPR3-independent pathway uses 4,5-didehydrojasmonate for

jasmonate   synthesis.   *Nature   Chem.   Biol.*   **14**,   171   (2018).

https://doi.org:10.1038/nchembio.2540

44   Stenzel, I. *et al.* Jasmonate biosynthesis and the allene oxide cyclase family of

Arabidopsis thaliana. *Plant Mol. Biol.* **51**, 895-911 (2003).

45   Ribot, C., Zimmerli, C., Farmer, E. E., Reymond, P. & Poirier, Y. Induction of the

Arabidopsis *PHO1;H10* gene by 12-oxo-phytodienoic acid but not jasmonic acid via a

CORONATINE INSENSITIVE1-dependent pathway. *Plant Physiol.* **147**, 696-706 (2008).

46   Pre, M. *et al.* The AP2/ERF domain transcription factor ORA59 integrates jasmonic acid

784   and ethylene signals in plant defense. *Plant Physiol.* **147**, 1347-1357 (2008).

785   47   Ohkama-Ohtsu, N. *et al.* 12-Oxo-phytodienoic acid-glutathione conjugate is transported

786   into the vacuole in Arabidopsis. *Plant Cell Physiol.* **52**, 205-209 (2011).

787   48   Mehrshahi, P. *et al.* Transorganellar complementation redefines the biochemical

788   continuity of endoplasmic reticulum and chloroplasts. *Proc. Natl. Acad. Sci. USA* **110**,

789   12126-12131 (2013). https://doi.org:10.1073/pnas.1306331110

790   49   Gould, S. J., Keller, G. A., Hosken, N., Wilkinson, J. & Subramani, S. A conserved

791   tripeptide sorts proteins to peroxisomes. *J. Cell Biol.* **108**, 1657-1664 (1989).

792   https://doi.org:10.1083/jcb.108.5.1657

793   50   Stellmach, H., Hose, R., Räde, A., Marillonnet, S. & Hause, B. A new set of Golden-

794   Gate-based organelle marker plasmids for colocalization studies in plants. *Plants* **11**, 2620

795   (2022).

796   51   Kaji, T. *et al.* Δ4-dn-iso-OPDA, a bioactive plant hormone of *Marchantia polymorpha*.

797   *iScience* **27**, 110191 (2024). https://doi.org:https://doi.org/10.1016/j.isci.2024.110191

798   52   Chung, H. S. *et al.* Regulation and function of Arabidopsis *JASMONATE ZIM*-domain

799   genes in response to wounding and herbivory. *Plant Physiol.* **146**, 952-964 (2008).

800   53   Howe, G. A. Jasmonates as signals in the wound response. *J. Plant Growth Regul.* **23**,

801   223-237 (2004).

802   54   Wasternack, C. in *Plant Hormone Signaling*   (eds P Hedden & SG Thomas) 185-228

803   (Blackwell Publishing, 2006).

804   55   Reymond, P., Weber, H., Damond, M. & Farmer, E. E. Differential gene expression in

805   response to mechanical wounding and insect feeding in Arabidopsis. *Plant Cell* **12**, 707-

806   719 (2000).

56 Delessert, C., Wilson, I., Straeten, D., Dennis, E. & Dolferus, R. Spatial and temporal

analysis of the local response to wounding. *Plant Mol. Biol.* **55**, 165-181 (2004).

57 Lewandowska, M. *et al.* Wound-induced triacylglycerol biosynthesis is jasmonoy-l-

isoleucin and abscisic acid independent. *Plant Biol.* **25**, 509-517 (2023).

https://doi.org:https://doi.org/10.1111/plb.13513

58 Goodspeed, D., Chehab, E. W., Min-Venditti, A., Braam, J. & Covington, M. F.

Arabidopsis synchronizes jasmonate-mediated defense with insect circadian behavior.

Proc. Natl. Acad. Sci. USA **109**, 4674-4677 (2012).

https://doi.org:10.1073/pnas.1116368109

59 Guo, Q. *et al.* JAZ repressors of metabolic defense promote growth and reproductive

fitness in *Arabidopsis. Proc. Natl. Acad. Sci. USA* **115**, E10768-E10777 (2018).

https://doi.org:10.1073/pnas.1811828115

60 Verhage, A. *et al.* Rewiring of the jasmonate signaling pathway in Arabidopsis during

insect herbivory. *Front. Plant Sci.* **2** (2011). https://doi.org:10.3389/fpls.2011.00047

61 Zhang, A. *et al.* Cell-wall damage activates DOF transcription factors to promote wound

healing and tissue regeneration in *Arabidopsis thaliana. Current Biology* **32**, 1883-

1894.e1887 (2022). https://doi.org:10.1016/j.cub.2022.02.069

62 Davletova, S., Schlauch, K., Coutu, J. & Mittler, R. The Zinc-Finger Protein Zat12 plays

a central role in reactive oxygen and abiotic stress signaling in Arabidopsis *Plant*

Physiol. **139**, 847-856 (2005). https://doi.org:10.1104/pp.105.068254

63 Mittler, R. *et al.* Gain- and loss-of-function mutations in Zat10 enhance the tolerance of

plants to abiotic stress. *FEBS Lett.* **580**, 6537-6542 (2006).

https://doi.org:https://doi.org/10.1016/j.febslet.2006.11.002

64 Mizoi, J. *et al.* Heat-induced inhibition of phosphorylation of the stress-protective

transcription factor DREB2A promotes thermotolerance of *Arabidopsis thaliana*. *J. Biol.*

Chem. **294**, 902-917 (2019). https://doi.org:https://doi.org/10.1074/jbc.RA118.002662

65 Sakuma, Y. *et al.* Dual function of an *Arabidopsis* transcription factor DREB2A in water-

stress-responsive and heat-stress-responsive gene expression. *Proc. Natl. Acad. Sci. USA*

103, 18822-18827 (2006). https://doi.org:doi:10.1073/pnas.0605639103

66 Pollmann, S. *et al.* Substrate channeling in oxylipin biosynthesis through a protein

complex in the plastid envelope of *Arabidopsis thaliana*. *J. Exper. Bot.* **70**, 1483-1495

(2019). https://doi.org:10.1093/jxb/erz015

67 Mueller, M. J. & Berger, S. Reactive electrophilic oxylipins: Pattern recognition and

signalling. *Phytochemistry* **70**, 1511-1521 (2009).

68 Mueller, S. *et al.* General detoxification and stress responses are mediated by oxidized

lipids through TGA transcription factors in *Arabidopsis*. *Plant Cell* **20**, 768-785 (2008).

69 Li, X. *et al.* Wounding induces a peroxisomal H_2O_2 decrease via glycolate oxidase-

catalase switch dependent on glutamate receptor-like channel-supported Ca^{2+} signaling in

plants. *The Plant Journal* **116**, 1325-1341 (2023).

https://doi.org:https://doi.org/10.1111/tpj.16427

70 Scholz, S. S., Reichelt, M., Mekonnen, D. W., Ludewig, F. & Mithöfer, A. Insect

herbivory-elicited GABA accumulation in plants is a wound-induced, direct, systemic,

and jasmonate-independent defense response. *Frontiers in Plant Science* **6**, 1128 (2015).

https://doi.org:10.3389/fpls.2015.01128

71 Hoermayer, L. & Friml, J. Targeted cell ablation-based insights into wound healing and

restorative patterning. *Current Opinion in Plant Biology* **52**, 124-130 (2019).

https://doi.org:https://doi.org/10.1016/j.pbi.2019.08.006

72   Yi, R. *et al.* Dioxygenase JID1 mediates the modification of OPDA to regulate jasmonate

homeostasis. *Cell Discovery* **9**, 39 (2023). https://doi.org:10.1038/s41421-023-00530-6

73   Floková, K. *et al.* A previously undescribed jasmonate compound in flowering

Arabidopsis thaliana – The identification of *cis*-(+)-OPDA-Ile. *Phytochemistry* **122**, 230-

237 (2016). https://doi.org:http://dx.doi.org/10.1016/j.phytochem.2015.11.012

74   Mik, V. *et al.* Synthetic and analytical routes to the L-amino acid conjugates of *cis*-OPDA

and their identification and quantification in plants. *Phytochemistry* **215**, 113855 (2023).

https://doi.org:https://doi.org/10.1016/j.phytochem.2023.113855

75   Ellis, C. & Turner, J. A conditionally fertile *coi1* allele indicates cross-talk between plant

hormone signalling pathways in *Arabidopsis thaliana* seeds and young seedlings. *Planta*

215, 549-556 (2002).

76   Yang, D.-L. *et al.* Plant hormone jasmonate prioritizes defense over growth by interfering

with gibberellin signaling cascade. *Proceedings of the National Academy of Sciences* **109**,

E1192–E1200 (2012). https://doi.org:10.1073/pnas.1201616109

77   Czechowski, T., Stitt, M., Altmann, T., Udvardi, M. K. & Scheible, W.-R. Genome-wide

identification and testing of superior reference genes for transcript normalization in

Arabidopsis. *Plant Physiol.* **139**, 5-17 (2005). https://doi.org:10.1104/pp.105.063743

78   Wang, H. *et al.* Reference genes for normalizing transcription in diploid and tetraploid

Arabidopsis. *Sci. Rep.* **4**, 6781 (2014). https://doi.org:10.1038/srep06781

79   Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C(T)

method. *Nature Protocols* **3**, 1101-1108 (2008).

80   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals

and   plants.   *Bioinformatics*   **36**,   2628-2629   (2019).

https://doi.org:10.1093/bioinformatics/btz931

81   Ge, S. X., Son, E. W. & Yao, R. iDEP: an integrated web application for differential

expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics* **19**, 534 (2018).

https://doi.org:10.1186/s12859-018-2486-6

82   Laemmli, U. K. Cleavage of structural proteins during the assembly of the head of

bacteriophage T4. *Nature* **227**, 680-685 (1970).

83   Otto, M., Naumann, C., Brandt, W., Wasternack, C. & Hause, B. Activity regulation by

heteromerization of Arabidopsis allene oxide cyclase family members. *Plants* **5**, 3 (2016).

84   Engler, C. *et al.* A Golden Gate modular cloning toolbox for plants. *ACS Synthetic Biol.*

3, 839-843 (2014). https://doi.org:10.1021/sb4001504

85   Zhang, X., Henriques, R., Lin, S.-S., Niu, Q.-W. & Chua, N.-H. *Agrobacterium*-mediated

transformation of *Arabidopsis thaliana* using the floral dip method. *Nature Protocols* **1**,

641-646 (2006). https://doi.org:10.1038/nprot.2006.97

86   Lu, J. *et al.* Types of nuclear localization signals and mechanisms of protein import into

the nucleus. *Cell Comm. Signal.* **19**, 60 (2021). https://doi.org:10.1186/s12964-021-

00741-y

87   Raikhel, N. Nuclear targeting in plants 1. *Plant Physiol.* **100**, 1627-1632 (1992).

https://doi.org:10.1104/pp.100.4.1627

88   Kosugi, S., Hasebe, M., Tomita, M. & Yanagawa, H. Nuclear export signal consensus

sequences defined using a localization-based yeast selection system. *Traffic* **9**, 2053-2062

(2008). https://doi.org:https://doi.org/10.1111/j.1600-0854.2008.00825.x

89   Kunze, M. & Berger, J. The similarity between N-terminal targeting signals for protein

import into different organelles and its evolutionary relevance. *Front. Physiol.* **6** (2015).

https://doi.org:10.3389/fphys.2015.00259

90   De Meyer, T. & Depicker, A. Trafficking of endoplasmic reticulum-retained recombinant

proteins is unpredictable in *Arabidopsis thaliana*. *Front. Plant Sci.* **5** (2014).

https://doi.org:10.3389/fpls.2014.00473

91   Eseverri, Á. *et al.* Transit peptides from photosynthesis-related proteins mediate import of

a marker protein into different plastid types and within different species. *Front. Plant Sci.*

11 (2020). https://doi.org:10.3389/fpls.2020.560701

92   von Heijne, G., Steppuhn, J. & Herrmann, R. G. Domain structure of mitochondrial and

chloroplast targeting peptides. *Europ. J. Biochem.* **180**, 535-545 (1989).

https://doi.org:https://doi.org/10.1111/j.1432-1033.1989.tb14679.x

93   Norkunas, K., Harding, R., Dale, J. & Dugdale, B. Improving agroinfiltration-based

transient gene expression in *Nicotiana benthamiana*. *Plant Methods* **14**, 71 (2018).

https://doi.org:10.1186/s13007-018-0343-2

94   Janik, K., Stellmach, H., Mittelberger, C. & Hause, B. in *Phytoplasmas: Methods and*

Protocols (eds Rita Musetti & Laura Pagliari) 321-331 (Springer New York, 2019).

95   Balcke, G. *et al.* An UPLC-MS/MS method for highly sensitive high-throughput analysis

of phytohormones in plant tissues. *Plant Methods* **8**, 47 (2012).

96   Studier, F. W. Protein production by auto-induction in high density shaking cultures.

Protein Expr. Purif. **41**, 207-234 (2005). https://doi.org:10.1016/j.pep.2005.01.016

97   Feussner, K., Abreu, I. N., Klein, M. & Feussner, I. in *Methods Enzymol.* Vol. 680 (ed

Joseph Jez) 325-350 (Academic Press, 2023).

Figures

Figure 1:

Fig. 1: MeJA pretreatment restitutes AOC protein levels in the mutants and OPDA content in the *opr2opr3* mutant.

(a) Synthesis of jasmonic acid (JA)/JA-Ile from α-linolenic acid (α-LeA). The biosynthesis starts with the release of α-LeA from chloroplast membrane galactolipids and its conversion to (13S)-hydroperoxyoctadecatrienoic acid (13-HPOT) by 13-LIPOXYGENASE (LOX). *cis*-(+)-12-oxophytodienoic acid (OPDA) is formed in the chloroplast (green background) from 13-HPOT through sequential actions of ALLENE OXIDE SYNTHASE (AOS) and ALLENE OXIDE CYCLASE (AOC). In the peroxisomes (yellow background) reduction of cyclopentenone ring of OPDA is catalyzed by a peroxisomal OPDA REDUCTASE (OPR3) resulting in the formation of 3-oxo-2-(20(Z)-pentenyl)-cyclopentane-1-octanoic (OPC-8:0). Shortening of the octanoic acid side chains of OPC-8:0 by three rounds of β-oxidation yields JA. The cytosol-released JA is conjugated to Isoleucine (Ile) by the enzyme JASMONATE RESISTANT 1 (JAR1) resulting in the major bulk formation of the biologically active JA-Ile. OPDA can directly undergo three rounds of β-oxidation yielding 4,5-didehydro-JA (4,5-ddh-JA) in a parallel OPR3-bypass pathway. The formed 4,5-ddh-JA is released to the cytosol and then reduced to JA by OPR2 resulting in JA-Ile formation. Enzymes shaded in green, yellow and blue are located in the plastids, peroxisomes and cytosol, respectively.

(b) AOC protein content in seedlings pretreated with 1 µM MeJA or water (Mock) visualized by immunoblot, equal loading is depicted by staining with No-Stain™ Protein Labeling Reagent.

(c) AOC band intensity quantified relative to total protein loading.

(d) RT-qPCR of JA-responsive genes *LOX2*, *AOS*, *AOC2* and *OPR3* showing transcript accumulation at control condition (-wounding) and at 1 h after wounding. Seedlings were pretreated with MeJA or water (Mock) during development.

(e) Levels of *cis*-OPDA, *dn*-OPDA, JA and JA-Ile in seedlings of wild-type (Col-0), *opr2opr3*,

and *aos* at control condition (-wounding) and 1 h after wounding. For detailed values see

Supplementary Tab. 2.

Seedlings were pretreated with MeJA or water (Mock) during development (see Supplementary

Fig. 2a for experimental set-up). Bars represent means of three biological replicates with 120

seedlings each (single dots; ±SEM). Statistically significant differences among genotypes within

each condition were calculated using One-Way ANOVA followed by Tukey-HSD and are

indicated by asterisks with *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

Figure 2:

Fig. 2: The JA/OPDA-independent wound response of seedlings. Ten-day old seedlings from Col-0, *opr2opr3* and *aos* were pretreated with MeJA during development and were either unwounded (control) or harvested 1 h post wounding with forceps (wounding).

(a) Number of differentially regulated genes (DEGs) after wounding compared to non-wounded controls in seedlings of Col-0, *opr2opr3* and *aos*, all pretreated with 1 µM MeJA during development (see Fig. S2 for experimental set-up). DEGs were identified using FDR and FC cutoffs of 0.05 and 2, respectively.

(b) Bar plot showing Gene Ontology enrichment analysis summarizing the significant biological processes (FDR=0.05) enriched commonly in all genotypes in response to wounding, with bars indicating gene fold enrichment and color scale indicating FDR values.

(c) Heatmap illustrating 50 out of 211 DEGs enriched in the GO term 'response to stress' from **(b)**. Average FPKM values of three independent biological replicates were transformed into row normalized Z-score.

(d) RT-qPCR validation of DEGs previously described as OPDA-responsive genes. Transcript accumulation of *DREB2A*, *FAD-OXR*, *ZAT10, ERF5, TCH4,* and *GST6* in Col-0, *opr2opr3* and *aos* at control and wounding conditions. Transcript levels were normalized to those of *PP2A3*. Bars represent means of three biological replicates (single dots; ±SEM). Statistically significant differences among genotypes within each condition were calculated using Two-Way ANOVA followed by Tukey-HSD and are indicated by different letters.

Figure 3:

Fig. 3: OPDA-specific transcriptional change deduced from transcriptomic comparison between seedlings of *opr2opr3* and *aos* at control and wounding conditions. Ten-day old seedlings from *opr2opr3* and *aos* were pretreated with MeJA during development and were either unwounded (control, C) or harvested 1 h post wounding with forceps (wounding, W).

(a) Venn diagram representing genes showing differential expression when *opr2opr3* is compared to *aos* at control and wounding conditions (FDR cutoff = 0.05 and FC cutoff = 2).

(b) Heatmap illustrating the expression levels of the eleven wounding-related DEGs from (A). The average FPKM values of three independent biological replicates were transformed into row-normalized Z-score.

(c) Transcript accumulation of *AOS*, *OPR2,* and *OPR3* selected from **(b)** and determined by RT-qPCR. Transcript levels were normalized to those of *PP2A3*. Bars represent means of three

994 biological replicates (single dots; ±SEM). Asterisks denote statistically significant differences

995 among genotypes as determined by Student's t-test within each condition (** $p<0.01$, ***

996 $p<0.001$, **** $p<0.0001$).

Figure 4:

1000

Fig. 4: Application of OPDA results in the induction of sulfur metabolism.

(a) Heatmap of DEGs showing enrichment in genes involved in sulfur assimilation and common

hormone signaling in seedlings of *opr2opr3* following application of 25 µM OPDA for 30 min.

DEGs were plotted and compared to those from wounded seedlings of Col-0, *opr2opr3* and *aos*

using average FPKM values transformed into row-normalized Z-score of three biological

replicates.

(b) RT-qPCR validation of the up-regulation of the sulfur metabolism genes *SDI1*, *APR3* and

LSU3 as well as the regulator of JA- and ethylene-responsive gene expression *ORA59* in

opr2opr3 seedlings treated with OPDA in comparison to wounded *opr2opr3* seedlings. OPDA-

treatment and wounding of seedlings were compared to seedlings which were non-wounded

(control) and water-treated (water), respectively.

(c) RT-qPCR validation of the up-regulation of the OPDA-responsive genes *DREB2A*, *FAD-*

OXR, *ZAT10* and *ERF5* in *opr2opr3* seedlings treated with OPDA in comparison to wounded

opr2opr3 seedlings. OPDA-treated and wounded *opr2opr3* seedlings were compared to their

respective controls, non-wounded (control) and water-treated seedlings, respectively.

Bars **(b-c)** represent means of three biological replicates with 120 seedlings each (single dots;

±SEM). Statistically significant differences among treatments were calculated using Two-Way

ANOVA followed by Tukey-HSD and are indicated by different letters.

1019

1020 **Figure 5:**

1021

1022

Fig. 5: The removal of the SRL peroxisomal signal results in a cytosolic localization of

OPR3 in *N. benthamiana* leaves and targeting of OPR3ΔSRL to the nucleus, cytosol, plastid

stroma and mitochondria in stable transformed *opr2opr3* lines.

(a-b) OPR3 (a) and OPR3ΔSRL (b) were C-terminally fused to YFP and transiently expressed in

N. benthamiana leaves under control of the CaMV *35S* promoter. Imaging was performed at 48-

72 h post infiltration.

(c-g) Stable *opr2opr3* complementation lines expressing OPR3ΔSRL targeted to the nucleus **(c)**,

the cytosol **(d)**, ER **(e)**, chloroplast stroma **(f)** and mitochondria **(g)**, all under control of the

CaMV *35S* promoter. All pictures are taken from homozygous T2 single insertion lines.

The yellow signal corresponds to YFP, whereas the magenta corresponds to chlorophyll

autofluorescence. Scale bars represent 20 µm. NLS, nuclear targeting signal; NES, nuclear export

signal; SIP, signal peptide; cTP, chloroplast targeting peptide; mTP, mitochondria targeting

peptide.

1036

Figure 6:

Fig. 6: Restitution of the fertility and JA/JA-Ile levels of *opr2opr3* plants by OPR3-targeted to different organelles.

(a-h) Open flower buds showing stamen filament elongation and pollen release by anthers (scale bars represent 0.5 mm) and plant inflorescences showing silique formation at days 50-60 post germination with asterisks marking fully developed siliques. Col-0 and *opr2opr3* transformed with an empty vector **(a, b)** show fertility and sterility, respectively. *opr2opr3* T2 lines complemented with peroxisome- **(c)**, nucleus- **(d)**, cytosol- **(e)**, ER- **(f)**, plastid stroma- **(g)** and mitochondria- **(h**) targeted OPR3 fused to YFP show the rescue of the *opr2opr3* sterility phenotype. Red asterisk show exemplary siliques containing seeds.

(i) OPDA, JA and JA-Ile accumulation at 1 h after wounding of seedlings of wild-type (WT) or *opr2opr3* transformed with either empty vector (EV) or *35S::OPR3* targeted to peroxisomes (Px), cytosol (Cyt) or mitochondria (Mt). Numbers refer to independent transgenic lines, which were selected for single insertion and homozygosity. Bars represent means of four biological replicates with 120 seedlings each (single dots; ±SEM). Statistically significant differences between *opr2opr3* transformed with empty vector and transformed with the OPR3 constructs were calculated using One-Way ANOVA followed by Tukey HSD ($p<0.05$) and are denoted by different letters.

Figure 7:

1059

1060

Fig. 7: Restitution of the fertility and JA/JA-Ile levels of *opr2opr3* plants by OPR2, but not

by OPR1, both targeted to different organelles.

Stable *opr2opr3* complementation lines expressing OPR2 (left) or OPR1 (right) targeted to the

cytosol (without any signal peptide), to the peroxisomes (OPR-SKL) or to mitochondria (mTP-

OPR), all under control of the CaMV *35S* promoter. All transgenic lines showed a similar level of

transgene expression (Supplementary Fig. 10).

(a) Micrographs showing the correct localization of the respective OPR-YFP-fusion. The yellow

signal corresponds to OPR-YFP, whereas the magenta signal corresponds to chlorophyll

autofluorescence. Scale bars represent 50 µm.

(b, c) Open flower buds to show stamen filament elongation and pollen release by anthers **(b)** and

plant inflorescences showing silique formation at days 50-60 post germination **(c)**. Note that

transformation of *opr2opr3* with all variants of OPR2 and with OPR1 targeted to peroxisomes

resulted in a rescue of fertility visible by elongated filaments and fully developed siliques.

Targeting of OPR1 to cytosol led to partial rescue shown by few fully developed siliques only

(red asterisks), whereas targeting of OPR1 to mitochondria did not rescue fertility. Scale bars in

(b) represent 0.5 mm.

(d) JA and JA-Ile accumulation at 1 h after wounding of seedlings of *opr2opr3* transformed with

either *35S::OPR2* or *35S::OPR1*, either targeted to cytosol (Cyt), peroxisomes (Px), or

mitochondria (Mt). Numbers refer to independent transgenic lines, which were selected for single

insertion and homozygosity. The mean levels of JA/JA-Ile of *opr2opr3* transformed with the

empty vector are taken from Fig. 6 and are represented by red dashed lines for comparison. Bars

represent means of four biological replicates with 120 seedlings each (single dots; ±SEM).

Statistically significant differences between *opr2opr3* transformed with the different *OPR2* and

OPR1 constructs and *opr2opr3* transformed with the empty vector were calculated using One-

Way ANOVA with Fisher LSD post-hoc test and are indicated by asterisks with $*p<0.05$,

$**p<0.01$, $***p<0.001$, and $****p<0.0001$; ns = non-significant.

1087

Figure 8:

Fig. 8: Schematic picture on the approaches leading to insights about the putative distribution of OPDA and 4,5-ddh-JA within cells of a wounded Arabidopsis seedling.

Putative subcellular distribution of OPDA and its conversion to JA deduced from the trans-organellar complementation approach.

(a) In wild-type plants, OPDA is converted to JA in the peroxisome through the OPR3 canonical route, the contribution of OPR2 located in the cytosol appears to be neglectable but comes into place if OPR3 is not functional (e.g., *opr3* single mutant, see [42]).

(b) The interruption of both routes in the *opr2opr3* mutant results in conversion of OPDA to 4,5-ddh-JA which is not metabolized to JA/JA-Ile.

(c) Targeting of OPR3 to any organelle in *opr2opr3* mutant plants (depicted for mitochondrial localization) restitutes JA biosynthesis in this mutant. If this is a result of either an ability of OPDA or of its β-oxidation product, 4,5-ddh-JA, to translocate to other cell compartments cannot be decided by this approach, since OPR3 is active to convert OPDA and 4,5-ddh-JA (red

question marks). Trans-organellar complementation by using OPR2 or OPR1 targeted into

mitochondria, however, helped to clarify this point.

(d) Mitochondrial targeting of OPR2 with its specificity to 4,5-ddh-JA complemented JA

biosynthesis favoring rather an ability of 4,5-ddh-JA to be mobile, although the conversion of

OPDA can still not be excluded due to some activity of OPR2 towards OPDA (light red question

mark).

(e) As final proof, OPR1 was targeted to mitochondria, because OPR1 is specific for OPDA and

does not convert 4,5-ddh-JA. Expression of mtOPR1 in *opr2opr3* did not restitute JA/JA-Ile

production. Even the very limited JA formation by cytosolic OPR1 suggests restricted occurrence

of OPDA in this compartment (not shown in this scheme, data see Figure 7). The lack of JA

formation by mitochondrial OPR1 suggests that OPDA may not have the ability to access this

cellular compartment and might largely compartmentalized in the organelles involved in JA

biosynthesis, the plastids and peroxisomes, and occurs in limited amounts in the cytosol.

1117

Supplementary information

Fig. S1: Seedlings of *opr2opr3* mutant show diminished AOC proteins content and produce lower OPDA levels compared to wild-type (Col-0).

Fig. S2: The experimental setup for transcriptome analyses.

Fig. S3: Hierarchical clustering and principal component analysis (PCA) illustrating the wound-induced transcriptional changes in Col-0, *opr2opr3* and *aos*.

Fig. S4: Wound-induced gene expression and hormone levels in *coi1* mutant leaves.

Fig. S5: The JA-dependent wound response occurring in wild-type seedlings.

Fig. S6: Transcriptome comparisons of wounded rosettes show limited signaling of OPDA.

Fig. S7: Application of OPDA results in the induction of JA-Ile signaling.

Fig. S8: The concept of the trans-organellar complementation of the *opr2opr3* mutant.

Fig. S9: OPR3 targeted to the ER, cytosol, nucleus, and chloroplasts colocalizes with the corresponding organelle markers in *N. benthamiana* protoplasts.

Fig. S10: Expression levels of the transgene in *opr2op3* mutants transformed with diverse OPR constructs.

Fig. S11: OPR3 reduces OPDA and 4,5-ddh-JA *in vitro*.

Fig. S12: OPDA levels at 1 h after wounding of seedlings of wild-type (WT) or *opr2opr3* transformed with either empty vector (EV), *35S::OPR2* or *35S::OPR1* targeted to peroxisomes (Px), cytosol (Cyt) or mitochondria (Mt).

Tab. S1: OPDA and JA levels shown in Fig. S1c-d.

Tab. S2: OPDA and JA levels shown in Fig. 1e.

Tab. S3: OPDA and JA levels shown in Fig. S5a.

Tab. S4: Genes selected from Taki et al. (2005) showing a similar wound-induced induction or

repression in Col-0, *opr2opr3* and *aos*.

Tab. S5: Peptide sequences of the targeting signals used for subcellular targeting.

Tab. S6: Primers sequences used for mutants genotyping and RT-qPCR.

Tab. S7: Primer sequences used for Golden Gate cloning of OPR3, OPR2 and OPR1 and their

subcellular targeting.

1148

1149 **Additional Supplementary Files**: Datasets 1-3