# Unraveling the Role of 12-*cis*-Oxo-Phytodienoic Acid in the Wound Response of *Arabidopsis thaliana*: Insights from Transcriptomic and Complementation Studies

**Khansa Mekkaoui[1], Ranjit Baral[1], Fiona Smith[1], Moritz Klein[2], Ivo Feussner[2], Bettina Hause[1,\*]**

[1] Department of Cell and Metabolic Biology, Leibniz Institute of Plant Biochemistry, Weinberg 3, D-06120 Halle/Saale, Germany

[2] Department of Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences and Goettingen Center for Molecular Biosciences (GZMB), University of Goettingen, Justus-von-Liebig-Weg 11, 37077 Göttingen, Germany

Corresponding author: Bettina Hause, Bettina.Hause@ipb-halle.de

**Running title:**

Signaling and compartmentalization of OPDA

## Abstract

In addition to jasmonoyl-isoleucine (JA-Ile), a well-established signaling molecule for plant growth and defense, its precursor, *cis*-12-oxo-phytodienoic acid (OPDA), is thought to possess independent signaling functions. Its perception in vascular plants is still uncharacterized. Several OPDA functions in Arabidopsis were inferred from a mutant that is affected in the function of the OPDA REDUCTASE3 (OPR3), catalyzing the conversion of OPDA within peroxisomes. Recently, *opr3* plants were found to accumulate JA-Ile via a cytosolic OPR2-mediated bypass. Given the uncoupling of OPDA and JA biosynthesis in the JA-deficient mutant *opr2opr3*, potential OPDA signaling was investigated by a transcriptome approach comparing wild type, *opr2opr3* and the JA- and OPDA-deficient mutant *allene oxide synthase*. Dissecting the wound response of seedlings revealed that OPDA lacked a transcriptional signature, and that previously characterized OPDA-response genes were wound-induced independently of OPDA. Exogenous application of OPDA to *opr2opr3* seedlings led to JA-Ile formation and signaling even in absence of OPR2 and OPR3 and resulted in activation of sulfur assimilation. These divergent responses to endogenously synthesized and applied OPDA suggest a compartmentalization of endogenous OPDA which was investigated by a trans-organellar complementation approach. OPR3 complemented the *opr2opr3* mutant in terms of fertility and wound-induced JA-Ile production irrespective of its subcellular localization. *In vitro* enzymatic activity of OPR3, however, showed conversion of OPDA and 4,5-didehydro-JA (4,5-ddh-JA), therefore not allowing to conclude which compound is translocated. Dissecting the conversion of either OPDA or 4,5-ddh-JA by OPR2 and OPR1 organelle variants pointed to a strong OPDA compartmentalization supporting its lacking signaling capacity.

**Introduction**

Jasmonic acid (JA) and its bioactive form jasmonoyl-isoleucine (JA-Ile) function primarily as signaling molecules coordinating plant growth and reproduction and responses to biotic and abiotic stresses (Li et al., 2022; Wang et al., 2019; Wasternack and Hause, 2013; Zhang et al., 2023). They play a central role in the defense of plants to mechanical damage and herbivory, as JA-Ile rapidly accumulates at the site of injury and acts as a signaling molecule by triggering a cascade of defense responses (Koo and Howe, 2009; Wasternack, 2019). JA-Ile signaling involves a series of molecular events mediated by its perception by the F-box protein CORONATINE INSENSITIVE 1 (COI1) leading to the degradation of the JASMONATE ZIM-DOMAIN (JAZ) repressors, hence the release of the expression of JA-responsive genes (Chini et al., 2007; Monte et al., 2022; Sheard et al., 2010; Thines et al., 2007; Xie et al., 1998; Xu et al., 2002; Yan et al., 2018).

One major intermediate in the formation of the bioactive JA-Ile, *cis*-12-oxo-phytodienoic acid (OPDA), has been known to be also a signaling molecule with independent functions from JA-Ile (Dave and Graham, 2012; Jimenez-Aleman et al., 2022; Liu and Park, 2021; Wasternack and Strnad, 2016). OPDA formation takes place in the chloroplast via the octadecanoid pathway, by oxygenation of α-linolenic acid by 13-lipoxygenases (LOXs) and the coupled actions of ALLENE OXIDE SYNTHASE (AOS) and ALLENE OXIDE CYCLASE (AOC) (Andreou and Feussner, 2009; Bannenberg et al., 2009; Feussner and Wasternack, 2002) (Figure 1A). Metabolization of OPDA into JA occurs after its transfer to peroxisomes, where the reduction of its cyclopentenone ring by OPDA REDUCTASE3 (OPR3) and three rounds of β-oxidation take place (Schaller and Stintzi, 2009; Schaller et al., 2000; Wasternack and Hause, 2013). A parallel hexadecanoid pathway gives rise to *dinor*-12-oxo-phytodienoic acid (*dn*-OPDA), which in turn is converted to JA following the same pathway as OPDA, but with two rounds of β-oxidation only (Weber et al., 1997).

The putative distinct function of OPDA in Angiosperms partially derives from its function in lower plants by inducing defense- and growth inhibition-related transcriptional reprogramming in *Marchantia polymorpha* (Chini et al., 2023; Monte *et al.*, 2022; Monte et al., 2018; Monte et al., 2020). Such JA-independent functions of OPDA were also characterized in higher plants for different vegetative processes including initiation of tendril coiling in *Bryonia dioica* (Stelmach et al., 1998), regulation of embryo development in *Solanum lycopersicum* (Goetz et al., 2012), and inhibition of seed germination and induction of stomatal closure in *Arabidopsis thaliana* (Chang et al., 2023; Dave et al., 2011; Savchenko and Dehesh, 2014). Despite its perception being not

identified in Arabidopsis, OPDA was shown to act as signaling molecule by transcriptionally inducing a set of genes, which is not induced by JA (Taki et al., 2005). These OPDA-specific response genes were characterized as wound- and OPDA-responsive in a COI1-independent manner and encompassed mainly stress-responsive genes, e.g. *DRE-BINDING PROTEIN 2A* (*DREB2A*), *FAD-LINKED OXIDOREDUCTASE* (*FAD-OXR), SALT TOLERANCE ZINC FINGER* (*ZAT10)* and the glutaredoxin member *GRX480*. OPDA contains an α, β-unsaturated carbonyl group which is thought to react with nucleophilic molecules by Michael addition (Alméras et al., 2003). The electrophilic nature of oxylipins could convey biological activities, as deduced from the accumulation of oxylipins adducts, including OPDA-GSH conjugates, during the hypersensitive response in pathogen-elicited tobacco leaves (Davoine *et al.*, 2006). OPDA binding to the CYCLOPHILIN20-3 (CYP20-3) was suggested to trigger the formation of the cysteine synthase complex activating sulfur assimilation to regulate cell redox homeostasis through mediation of *TGACG-BINDING* (TGA) transcription factors (Müller et al., 2017; Park et al., 2013). Here, CYP20-3-dependent OPDA signaling induces high levels of GSH leading to a retrograde signaling, which coordinates expression of OPDA-induced genes (Adhikari and Park, 2023). These transcriptional changes in Arabidopsis induced by OPDA through CYP20-3 module and/or additional uncharacterized mechanism consolidated the hypothesis of its function not limited to being only a JA precursor (Liu and Park, 2021).

The previous work characterizing OPDA signaling and JA-independent functions in Arabidopsis relied mainly on the use of the *opr3* mutant as the genetic background where OPDA and JA biosynthesis were uncoupled due to missing OPR3 (Stintzi and Browse, 2000; Stintzi et al., 2001). The *opr3* mutant is male sterile like the *aos* mutant being deficient in OPDA and JA (Browse, 2009; Stintzi and Browse, 2000; von Malek et al., 2002). Both mutants show defects in anther dehiscence, stamen elongation and pollen development showing that these defects were due to impairments in JA production. Persistent resistance of the *opr3* mutant against necrotrophic pathogens initially hinted towards a defense function of OPDA, but further studies uncovered an OPR3-independent pathway leading to JA/JA-Ile formation through the cytosolic OPR2 in the *opr3* mutant (Chini et al., 2018). Low JA-Ile formation in the *opr3* mutant rather than OPDA was shown to be responsible of its resistance, hence challenging the suggested JA-independent OPDA signaling function (Chini et al., 2018). Thus, the recently identified "leakiness" of the *opr3* mutant and formation of JA-Ile via bypassing the OPR3 (Figure 1A) questioned previously identified

functions of OPDA. Consequently, OPDA-specific signaling exclusively analyzed with the *opr3* mutants requires further studies.

This study re-addressed the question whether OPDA can function as an independent signaling molecule in Arabidopsis besides being a JA precursor. Using the *opr2opr3* mutant where formation of JA/JA-Ile does not occur (Chini et al., 2018), we investigated the effects of basal and wound-induced OPDA on gene expression in seedlings of *A. thaliana*. The results suggested that the endogenously formed OPDA is unlikely to be responsible to mediate gene expression, while exogenous supply of OPDA resulted in a distinct signaling. Exogenously supplied OPDA is predominantly detoxified and partially converted to JA/JA-Ile, hinting towards a compartmentalization of the endogenously formed OPDA. Therefore, we addressed the question of the occurrence of OPDA within the cell and whether it could be perceived. A trans-organellar complementation approach showed that OPDA is likely to be confined to the cell compartments where it is synthesized and metabolized. Instead, 4,5-ddh-JA appears to be the mobile compound within the plant cell supporting the assumption that OPDA has no signaling function in wounded seedlings.

**Results**

**A JA positive feedback loop during development is responsible for the tissue capability to produce OPDA.**

The *opr2opr3* double mutant was characterized as JA-deficient, with JA formation through the peroxisomal OPR3 and alternatively through the cytosolic OPR2 being both interrupted (Chini et al., 2018). JA deficiency is associated with a decrease in the accumulation of JA-biosynthesis enzymes, among them specifically the AOC, due to the disruption of the JA positive feedback loop during development (Stenzel et al., 2003). Indeed, *opr2opr3* and the *aos* accumulated statistically lower levels of the AOC proteins in their seedlings in comparison to wild-type (Col-0) seedlings (Supplemental Figure 1A, B). These diminished AOC protein levels correlated with lower background and wound-induced levels of OPDA in the *opr2opr3* mutant compared to the wild type, whereas the JA levels remained at the detection limit (around 0.002 nmol/g of fresh weight) in the mutants (Supplemental Figure 1C, D), which contrast to *opr3* mutant plants accumulating JA levels after wounding of around 0.2 nmol/g of fresh weight (Stintzi et al., 2001). This confirmed the stronger JA deficiency of *opr2opr3* compared to the *opr3* mutant.

Due to the partial OPDA deficiency of *opr2opr3* seedlings, it was crucial to normalize the

levels with those of the wild type prior to investigating OPDA signaling within this mutant. To

achieve this, we mimicked the missing JA positive feedback loop in the *opr2opr3* and *aos* mutants

by supplying seedlings with 1 µM JA methyl ester (MeJA) during development to increase the

AOC protein levels (Supplemental Figure 2A). In contrast to water (mock-)pretreated seedlings,

seedlings pretreated with MeJA during development exhibited increased AOC protein levels in

wild type and both mutants (Figure 1B). Notably, this increase was more pronounced in the mutants

resulting in AOC protein levels in *opr2op3* and *aos* seedlings matching those of wild type seedlings

(Figure 1C). Transcript accumulation of the JA biosynthesis enzymes *LOX2*, *AOS*, *AOC2* and

*OPR3* determined by RT-qPCR corroborated the function of the JA positive feedback loop: MeJA

pretreatment during development resulted in a compensation of their basal transcript levels in the

mutants compared to wild type (Figure 1D). They all are, however, highly induced upon wounding

in wild-type seedlings, but their induction is significantly lower in the mutants (Figure 1D). This

shows that the MeJA pretreatment did not lead to a wound-induced JA-Ile signaling in the JA-

deficient mutants. Most importantly, however, the restoration of AOC protein content in the

*opr2op3* enhanced its basal and wound-induced OPDA and *dn*-OPDA levels to the levels of wild

type but did not affect the OPDA deficiency of the *aos* mutant (Figure 1E). Despite the

accumulation of low residual levels of JA in *opr2opr3* and *aos* seedlings upon pretreatment, wound

induction of JA and JA-Ile was only observed in the wild-type seedlings. This validates the JA

deficiency of both mutant lines and the exclusive OPDA enhancing role of the developmental JA-

feedback loop for the *opr2opr3* mutant (Figure 1E).

**A comparative transcriptomics approach shows a JA-specific but not an OPDA-specific**

**response to wounding.**

OPDA signaling in Arabidopsis was previously inferred from its capacity to mediate transcriptional

changes by inducing a specific set of OPDA-responsive genes in the "leaky" *opr3* single mutant

(Ribot et al., 2008; Taki et al., 2005). Another putative OPDA-signaling pathway was based on the

detected binding of OPDA to CYP20-3, which in turn transcriptionally induces TGA transcription

factors (Park et al., 2013). To further clarify this signaling function, a comparative transcriptomics

approach was set between wild-type and *opr2opr3* seedlings. To uncouple putative OPDA

signaling from wound-induced signaling processes, *aos* seedlings were included as a negative

control, since both JA and OPDA production are abolished in this mutant (Supplemental Figure

2B). mRNA sequencing of MeJA-pretreated seedlings from Col-0, *opr2opr3* and *aos* was performed using unwounded seedlings (control) and seedlings harvested 1 h after wounding. The overall wound-induced response showed a significant transcriptional change in all three genotypes (Supplemental Figure 3). Here, a strong similarity between the transcriptomes of *opr2opr3* and *aos* seedlings was evident in the expression heatmap and principal component analysis (PCA) irrespective of the condition (Supplemental Figure 3). Indeed, the highest number of differentially expressed genes (DEGs) was common to all three genotypes and were enriched in genes involved in stress, oxygen, and abiotic stimulus responses (Figure 2A, B). This general stress response was therefore rather wound-inducible but independent from the JA pathway. Interestingly, this group contained known stress-responsive genes with several genes being previously reported as OPDA-responsive, such as *DREB2A*, *FAD-OXR, ZAT10* and *GRX480* (Taki et al., 2005) (Figure 2C, D). Their wound-responsive induction in *aos* seedlings indicated that they are more likely to be associated with a JA-independent wound-induced stress response than with an OPDA-mediated response.

In wild-type seedlings, the signaling mediated by JA/JA-Ile was obvious as it shaped differentially the transcriptome already by sustaining statistically differential basal levels of 58 JA-responsive genes in comparison to the mutants (Supplemental Figure 4A, B). Wounding of wild-type seedlings resulted in a transcriptional change enriched in genes related to JA and immune responses (Supplemental Figure 4C). These transcriptional responses, notably of genes encoding JAZ transcription factors and other JA-responsive genes, such as *CHLOROPHYLLASE 1* (*CLH1*) and *N-ACETYLTRANSFERASE ACTIVITY 1* (*NATA1*), were dampened in the *opr2opr3* and *aos* mutants confirming the JA dependence of their induction upon wounding (Supplemental Figure 4D). These data suggest that next to a common wound response, JA/JA-Ile are main mediators of wound-induced transcriptional changes in wild-type seedlings.

**Wound-induced and exogenously applied OPDA resulted in different alterations of the transcriptome in *opr2opr3* seedlings.**

To identify OPDA-regulated genes, transcriptional differences between *opr2opr3* and *aos* were dissected and indicated a restricted number of DEGs at control and wounding conditions (Figure 3A). Except for the mutated genes, which showed diminished expression in the respective mutant backgrounds (*AOS*, *OPR2*, and *OPR3*), only nine specific genes exhibited differential expression in *opr2opr3* seedlings compared to *aos* seedlings, regardless of the experimental condition (Figure

3B, C). These genes encode five RNA species and four proteins, among them one cyclic nucleotide-gated ion channel (*CNGC11*), one cytochrome P450 (*CYP81D11*), and two hypothetical proteins. This is a very narrow or almost no difference between transcriptomic profiles of the mutants, although there were significantly enhanced OPDA levels in the *opr2opr3* mutant compared to *aos* (Figure 1E). These results make an OPDA-mediated gene expression in wounded seedlings of the *opr2opr3* mutant unlikely.

Previous work demonstrated a signaling role of OPDA in Arabidopsis by its exogenous supply to wild-type plants or to the *opr3* mutant (Park et al., 2013; Ribot et al., 2008; Taki et al., 2005). To determine whether the origin of OPDA caused the different outcomes, RNAseq data resulting from wounding of *opr2opr3* seedlings (= endogenous rise of OPDA) were compared to transcriptomic data generated from *opr2opr3* seedlings, which were treated with 25 μM OPDA for 30 min (= exogenous supply of OPDA). OPDA treatment resulted in transcriptional changes that were highly different from those induced by wounding (Supplemental Figure 5). The clustering of DEGs induced by OPDA treatment in comparison to DEGs induced by wounding showed an exclusive induction of JA signaling and sulfate reduction by OPDA application (Supplemental Figure 5). To get a deeper insight into this JA-mediated response occurring in *opr2opr3* seedlings upon OPDA treatment, expression levels of JA responsive genes were determined in comparison to wounded seedlings from wild type and both mutants (Figure 4). OPDA-treated *opr2opr3* and wounded wild-type seedlings shared the up-regulation of several JA marker genes, such as those encoding JAZ transcription factors and JA biosynthesis enzymes (Figure 4A). Validation of transcript accumulation of *JAZ2*, *JAZ7*, *JAZ13* and *CHL1* by RT-qPCR confirmed their exclusive induction in *opr2opr3* seedlings upon OPDA application but not upon endogenous OPDA accumulation by wounding (Figure 4B). This hints towards the conversion of the exogenously applied but not the endogenously formed OPDA to JA/JA-Ile despite the loss of OPR3 and OPR2 in the *opr2opr3* background. OPDA and JA levels in *opr2opr3* seedlings determined after wounding and OPDA treatment showed unequivocally that *opr2opr3* seedlings accumulated significantly higher levels of JA-Ile following OPDA application in comparison to wounding (Figure 4C). Despite these levels being lower than the wild-type JA-Ile levels, they correlated with the activated JA signaling in *opr2opr3* seedlings after OPDA application. Interestingly, the OPDA-fed *opr2opr3* seedlings accumulated only slightly higher OPDA levels than the wounded seedlings, whereas *dn*-OPDA accumulated significantly only following endogenous OPDA formation after wounding (Figure 4C). This suggests that the exogenously applied OPDA did not engage in β-

oxidation to form *dn*-OPDA in the *opr2opr3* mutant unlike the endogenously formed OPDA, which is able to enter the β-oxidation cycle leading to *dn*-OPDA accumulation (Chini et al., 2018). Nevertheless, JA-Ile formation after OPDA application without accumulation of *dn*-OPDA hints towards its conversion to 3-oxo-2-(20(Z)-pentenyl)-cyclopentane-1-octanoic (OPC-8:0) (Figure 1A) in the cytosol and its further metabolization into JA through β-oxidation in the peroxisome. Here, the third OPR isoenzyme, OPR1, which is cytosolic and has some activity towards OPDA (Chini et al., 2018), might contribute to the conversion of OPDA.

OPDA application to the *opr2opr3* mutant showed a significant induction of a second sub-cluster of genes belonging to sulfate reduction and assimilation pathway as well as the phosphorelay signal transduction pathway (Figure 4A, Supplemental Figure 5). These genes were not induced by the endogenous rise of OPDA and included genes encoding sulfate reductases, such as APS REDUCTASE1 (APR1), APR2 and APR3, ATP-SULFURYLASE1 (APS1), and SULFATE-DEFICIENCY INDUCED (SDI1) (Figure 4D). This is in line with the previously reported induction of sulfur metabolism by OPDA binding to the CYP20-3 module (Park et al., 2013). These results raise the question whether this is due to a mis-localization of the exogenously provided compound, whereas OPDA synthesized following wounding is compartmentalized and therefore does not induce the responses that the exogenously supplied OPDA does.

**Test for subcellular distribution of the endogenously produced OPDA using a trans-organellar complementation of the *opr2opr3* mutant.**

The data obtained from wounded seedlings of *opr2opr3* and described above suggest that endogenously produced OPDA does not exhibit signaling capacity. This leads to the question, whether OPDA is restricted to the cell compartments of its synthesis, transfer and metabolization, namely plastids, cytosol and peroxisomes, respectively. To test its putative translocation to other cell compartments, a trans-organellar complementation approach according to (Mehrshahi et al., 2013) using the *opr2opr3* mutant was performed. OPR3 was targeted to different organelles in the *opr2opr3* mutant, and the fertility of these plants was checked. A rescue of the JA-deficiency phenotype of the *opr2opr3* mutant would indicate whether its substrate, OPDA, is able to translocate to the tested organelles (Supplemental Figure 6).

OPR3 is a peroxisome-localized protein and is imported into the matrix of peroxisomes through the peroxisomal targeting signal type 1 (PTS1) consisting of a tripeptide at the extreme carboxy-tail of the protein (Gould et al., 1989). OPR3 contains a C-terminal SRL which is a

variation of the prototypic PTS1 signal SKL. The removal of the SRL signal resulted in a cytosolic localization of a fusion of OPR3ΔSRL with YFP when transiently expressed in *Nicotiana benthamiana* leaves indicating that import into peroxisomes was abolished (Supplemental Figure 7A, B). Targeting of OPR3 to the nucleus, cytosol, plastids, mitochondria, and endoplasmic reticulum (ER) was done by fusing OPR3ΔSRL with the respective targeting signals or target peptides (Supplemental Table 1). Validation of the subcellular localization of the different organelle variants of OPR3 with co-localization studies in *N. benthamiana* protoplasts was performed using established organelle markers (Stellmach et al., 2022) and indicated correct subcellular localizations of OPR3 compared to the markers independently of the position of the YFP fluorescent tag (Supplemental Figure 8). Stable transgenic *opr2opr3* lines complemented with the organelle variants of OPR3ΔSRL fused to YFP were generated and resulted in clear subcellular localizations of OPR3-YFP to the nucleus, cytosol, chloroplast stroma, mitochondria, and ER (Supplemental Figure 7C-G).

The main phenotypic feature of JA-deficient and JA-insensitive Arabidopsis mutants is the male sterility, which is also characteristic for *opr2opr3* plants and is visible by defects in filament elongation, pollen release and seed set (Browse, 2009). Plants of *opr2opr3* transformed with an empty vector (EV) showed short stamens with non-dehiscing anthers (Figure 5A, B), similar to the reported male sterility phenotype of the *opr3* mutant (Stintzi and Browse, 2000). As a positive control, the complementation of *opr2opr3* with the peroxisome targeted OPR3 showed full rescue of the flower phenotype (Figure 5C). Interestingly, all other tested organelle-variants of OPR3 also rescued the flower sterility and showed open flowers like the wild type with elongated stamens and anthers releasing pollen (Figure 5D-H). This rescue of sterility translated also in the ability of the lines to form siliques containing seeds. The only exception were plants complemented with the ER-targeted OPR3 having partial rescue of flower fertility only. These plants formed fewer and smaller siliques compared to the wild type (Figure 5F). It is tempting to speculate that OPR3 localized to the ER might have access only to limited amounts of its substrate, leading to partial JA/JA-Ile biosynthesis only.

Reconstitution of the JA pathway in vegetative tissue of the *opr2opr3* complementation lines was analyzed by determination of wound-induced levels of OPDA, JA and JA-Ile in two-week-old seedlings. Hormone levels of *opr2opr3* complemented with OPR3 targeted to either peroxisomes, cytosol or mitochondria were compared to those of wild type and *opr2opr3* transformed with an empty vector. Data from two independent transformed lines each revealed that

complementation of *opr2opr3* with OPR3 restored the JA/JA-Ile levels to wild-type levels (Figure 5I). These results show the reconstitution of JA biosynthesis by OPR3 localized to all the tested compartments.

**Trans-organellar complementation with OPR2 and OPR1 to determine whether OPDA or 4, 5-ddh-JA is the translocating substrate.**

JA biosynthesis in *opr2opr3* complemented with OPR3 located in different organelles would occur only if its substrate, OPDA, is able to translocate to these compartments. However, as an alternative pathway, OPDA could enter directly the β-oxidation as reported recently (Chini et al., 2018). This would result in formation of 4,5-ddh-JA, which might be metabolized by OPR3 irrespective of the organellar location. Conversion of 4,5-ddh-JA to JA by OPR3 was tested by *in vitro* enzymatic assay and showed that OPR3 reduces 4,5-ddh-JA *in vitro*, however to a lesser extent than OPDA (Supplemental Figure 9). Based on substrate consumption after 10 min, only half of the given 4,5-ddh-JA was reduced while OPDA was completely converted to OPC-8. The control reactions with inactivated enzyme showed no reduction activity. The results clearly show that OPR3 is able to reduce both cyclopentenones, 4,5-ddh-JA and OPDA, *in vitro*. In addition, it has been shown by recombinant enzymatic assays that OPR2 but not OPR1 is able to reduce 4,5-ddh-JA (Chini et al., 2018). Conversely, OPR1 is able to reduce OPDA although with very low efficiency but OPR2 is not (Schaller et al., 2000). To determine whether OPDA or 4,5-ddh-JA is the substrate able to translocate to the tested cell compartments and specifically to cell compartments unrelated to the location of JA biosynthesis, peroxisomal, cytosolic and mitochondrial variants of OPR1 and OPR2, both fused to YFP, were used to complement the *opr2opr3* mutant. Both OPRs localized to the cytosol, whereas addition of a C-terminal SKL and an N-terminal mitochondrial target sequence led to localization in peroxisomes and mitochondria, respectively (Figure 6A). All three variants expressing OPR2 rescued the flower sterility and showed open flowers like the wild type with elongated stamens and anthers releasing pollen (Figure 6B) as well as formation of seed-containing, fully developed siliques (Figure 6C). In contrast, expression of OPR1 located to peroxisomes was the only OPR1 variant leading to a rescue of the flower sterility, whereas targeting OPR1 to the cytosol led to partial rescue of flower fertility only and targeting OPR1 into the mitochondria did not restore the fertility (Figure 6B, C). These data were supported by the fact that only seedlings harboring the OPR2 variants and peroxisomal-located OPR1 were able to accumulate JA and JA-Ile upon wounding (Figure 6D), although all lines accumulated similar levels of OPDA

(Supplemental Figure 10). The JA/JA-Ile levels were lower than those of wounded wild-type seedlings, but the significantly increased levels in comparison to *opr2opr3* showed the capability of these plants to produce JA/JA-Ile. Most importantly, however, expression of OPR1 located to mitochondria did not restore fertility nor JA/JA-Ile levels. Since OPR1 does not convert 4,5-ddh-JA to JA, this result points to the fact that 4,5-ddh-JA but not OPDA might be the compound to be converted by OPR2 or OPR3 in different organelles.

**Discussion**

JA plays a major role in growth and defense against pathogens, herbivory attacks and mechanical wounding. Its signaling through the bioactive JA-Ile is well established in Angiosperms, while in lower plants, such as *M. polymorpha*, OPDA and its β-oxidation product *dn*-OPDA are bioactive (Chini et al., 2023; Monte et al., 2022; Monte et al., 2018; Monte et al., 2020). Given that *A. thaliana* plants produce both compounds, possible function of OPDA as a signaling molecule independently from JA was proposed. OPDA was shown to induce a distinct transcriptional change from JA when exogenously applied to Arabidopsis plants (Ribot et al., 2008; Taki et al., 2005). Additionally, OPDA´s involvement in processes like seed germination and stomatal closure was demonstrated (Chang et al., 2023; Dave et al., 2011; Savchenko and Dehesh, 2014). The mechanism by which OPDA exerts these functions in vascular plants is, however, still elusive (Jimenez-Aleman et al., 2022). A major genetic background used for studying OPDA functions was the *opr3* mutant (Browse, 2009). Recently, an OPR3-independent pathway of JA formation through the cytosolic isoenzyme OPR2 has been identified (Chini et al., 2018). Hence, uncoupling of JA production from OPDA was not achieved in this mutant, and the characterized functions of OPDA cannot be attributed to OPDA with certainty. Here, we addressed the question of whether OPDA functions independently from JA as a signaling molecule in Arabidopsis, by utilizing the *opr2opr3* mutant, which is more JA-deficient than *opr3* (Chini et al., 2018).

The use of the *opr2opr3* mutant is limited by its lower production of OPDA in comparison to the wild type due to the disrupted JA positive feedback loop as indicated by reduced protein levels of the biosynthesis enzymes, such as AOC (Wasternack and Hause, 2013). The enhancement of the AOC levels and subsequent OPDA production in the JA-deficient mutants was achieved by restituting the JA feedback loop through a repetitive supply of MeJA to seedlings during development (Figure 1). This result validated the role of the JA positive feedback loop in the

364 regulation of the basal levels of the biosynthetic enzymes and setting the tissue ability to produce

365 OPDA and putatively JA (Stenzel et al., 2003).

**The wounding response of wild-type seedlings involves JA-dependent and independent processes.**

To inspect the separate roles of OPDA and JA upon wounding, a comparative analysis of the

transcriptomes of seedlings of wild type, *opr2opr3* and *aos* was carried out. The transcriptomic

change induced by wounding in wild-type seedlings involved an up-regulation of genes previously

identified as JA-dependent or associated with immune responses, validating the specific signaling

role of JA-Ile in the wound response of seedlings as it has been described for mature leaves of

Arabidopsis and other plant species (Chung et al., 2008; Howe, 2004; Koo and Howe, 2009;

Wasternack, 2006). Unexpectedly, seedlings of the *aos* mutant being deficient in synthesis of JA

and OPDA, also showed a major transcriptional change upon wounding that was common to the

*opr2opr3* mutant and the wild type (Figure 2). The transcriptional response was significantly

enriched in general stress response pathways, with an induction of oxidative stress- and

transcription-related genes. This implies that JA-independent signaling processes occur in response

to mechanical wounding leading to transcriptional reprogramming. Among such processes, water

stress might contribute to the regulation of wound-responsive genes in a JA-independent manner

(Reymond et al., 2000). Ethylene and abscisic acid (ABA) also contribute to the wound response

by regulating photosynthesis and drought responsive genes, respectively (Delessert et al., 2004).

An ABA-dependent and JA-independent transcriptional regulation of wax biosynthesis to seal the

wounded sites of Arabidopsis leaves has been recently characterized, but appears to be controlled

post-translationally by JA (Lewandowska et al., 2023). As plants defective in JA/JA-Ile

biosynthesis or perception are severely diminished in their defense response (Browse, 2009;

Goodspeed et al., 2012; Guo et al., 2018; Verhage et al., 2011), the JA-independent wound response

has rather limited contribution to plant defense and might only mitigate the adverse effects of

wounding itself.

**An OPDA-specific transcriptional response is missing in wounded *opr2opr3* seedlings, but specific pathways are activated by the exogenous application of OPDA.**

394 The general stress response that was both JA and OPDA-independent, comprised genes that were

395 previously identified as OPDA-specific response genes, such as *FAD-OXR*, *DREB2A*, *ZAT10* and

*GRX480* (Taki et al., 2005). These genes did not show differential expression when the transcriptomes of *opr2opr3* and *aos* were compared. Given that they were equally induced in seedlings of *opr2op3* and *aos* mutants, their induction by wounding is OPDA-independent (Figure 2). Several abiotic stresses were reported to up-regulate the same genes (Davletova et al., 2005; Mittler et al., 2006; Mizoi et al., 2019; Sakuma et al., 2006), further indicating their independence from OPDA. Moreover, direct comparison of transcriptomes from wounded *opr2opr3* and *aos* seedlings with standard cut-off parameters did not yield a significant list of putative OPDA-regulated genes further confirming the absence of an OPDA signaling (Figure 3). This could be due to three facts: (i) A putative receptor for OPDA might be missing in *A. thaliana*, (ii) OPDA remains sequestered in the plastids by its biosynthetic enzymes (Pollmann et al., 2019), and (iii) OPDA is converted to 4,5-ddh-JA, which does not have signaling capacity (Chini et al., 2018). Regarding perception of OPDA, evolutionary studies of the JA-Ile co-receptor COI1 showed that the Arabidopsis protein does not perceive OPDA or *dn*-OPDA, dismissing possible signaling of OPDA through COI1 in this plant (Monte et al., 2022). Moreover, it has been shown that *dn-iso*-OPDA rather than *dn*-OPDA is the bioactive molecule in mosses, and that this compound does not occur in *A. thaliana* (Chini et al., 2023). The transcriptomic data obtained from *opr2opr3* and *aos* therefore point to the fact that OPDA or its homolog *dn*-OPDA are not perceived by COI1 or any other receptor in Arabidopsis.

The transcriptional signature of OPDA was predominantly observed in experiments relying on its exogenous application (Park et al., 2013; Ribot et al., 2008; Taki et al., 2005). Therefore, we compared the transcriptional changes following application of OPDA to *opr2opr3* seedlings with those by wound-induced, endogenously produced OPDA. This comparison revealed an up-regulation of JA-Ile signaling components only by the exogenous supply with OPDA and correlated with a significant accumulation of JA-Ile (Figure 4). Beside not uncoupling JA-Ile biosynthesis from OPDA, this approach indicates that JA-Ile formation in the *opr2opr3* background could occur through OPR1 due to the reported low activity of OPR1 towards *cis*-OPDA (Schaller et al., 2000). In this case, exogenously supplied OPDA might flood the cytosol and therefore is available to the cytosolic OPR1, whereas the endogenously produced OPDA might either stay in the plastid stroma (Pollmann et al., 2019) or is rapidly converted to 4,5-ddh-JA (Chini et al., 2018) (see below).

The induction of sulfur assimilation and GSH production by OPDA were attributed as specific signaling functions of OPDA (Park et al., 2013). Here, the transcriptional activation of

sulfur metabolism pathway was exclusive to the exogenous OPDA supply, contradicting an intrinsic signaling function of the endogenous OPDA (Figure 4). As an electrophilic species, OPDA was shown to disturb redox homeostasis (Mueller and Berger, 2009) and to inhibit photosynthesis by affecting the photosystem II fluorescence (Alméras et al., 2003; Mueller et al., 2008). In parallel, the *in planta* conjugation of OPDA to GSH, which is an important antioxidant substance, and its degradation in the vacuole were also demonstrated (Ohkama-Ohtsu et al., 2011). Taken together, this implies that OPDA may not directly regulate the sulfur assimilation pathway but instead triggers this response through its potential detoxification when exogenously applied to plants.

**Endogenously formed OPDA is sequestered to the organelles of JA biosynthesis**

To address the question whether OPDA remains sequestered in the plastids or whether it is converted to 4,5-ddh-JA, a trans-organellar complementation approach was performed by directing OPR3 into various organelles (Figure 5, Supplemental Figs. 7, 8). Occurrence of OPR3 in any of the tested organelles resulted in a complementation of the phenotype and JA/JA-Ile production upon wounding in *opr2opr3* mutants. This led to the assumption that OPDA might travel throughout the cell. Nevertheless, the possibility still exists that OPDA is translocated directly to the peroxisomes, where it is converted to 4,5-ddh-JA by β-oxidation as shown previously for *opr3* and *opr2opr3* (Chini et al., 2018). Given that OPR3 is able to convert 4,5-ddh-JA to JA and OPDA to OPC-8 *in vitro* (Supplemental Figure 9), the complementation approach with OPR3 could not discriminate between the presence of either OPDA or 4,5-ddh-JA outside of the plastids or the peroxisomes. To address this, organelle variants of the isoenzymes OPR2 and OPR1 were used to complement the *opr2opr3* mutant (Figure 6). OPR2 and OPR1 differ in their enzymatic activity against OPDA and 4,5-ddh-JA: OPR2 is able to convert both compounds, but OPR1 has been shown to be inactive on 4,5-ddh-JA (Chini et al., 2018). Since OPDA and 4,5-ddh-JA should occur in peroxisomes and possibly in the cytosol, reconstitution of JA biosynthesis was obtained using either OPR1 or OPR2 targeted to peroxisomes or the cytosol. Targeting OPR1 or OPR2 to mitochondria, however, discriminated whether OPDA or 4,5-ddh-JA is the substrate able to translocate to other organelles (Figure 7). Here, complementation of *opr2opr3* with a mitochondrial-located OPR1 did not rescue fertility and JA/JA-Ile production (Figure 6), whereas OPR2 in mitochondria did. Additionally, OPR1 fully complemented the mutant only within the

peroxisome, and its cytosolic variant resulted in minimal JA-Ile formation and partial rescue of

fertility. These results suggest a restricted occurrence of OPDA in the cytosol and its preferred

compartmentalization in plastids and peroxisomes, whereas 4,5-ddh-JA might be the substrate able

to translocate between cell organelles.

In conclusion, despite its perception and signaling in lower plants, OPDA did not mediate

transcriptional changes in Arabidopsis. The induction of a specific transcriptional change occurred

exclusively upon exogenous supply but not upon endogenous rise of OPDA. The formation of JA-

Ile only by exogenous OPDA indicated that such an approach can introduce bias and suggested

that the endogenous OPDA is compartmentalized. The use of the trans-organellar complementation

approach to study the subcellular distribution of OPDA showed that OPR3 has specificity to other

JA precursors beside OPDA. When other OPR isoenzymes with higher substrate specificities were

used, the levels of restitution of JA-Ile biosynthesis in the mutant supported OPDA being limited

to the known organelles in the canonical pathway, while 4,5-ddh-JA was likely able to translocate

within the cell.

**Material and methods**

**Plant material, growth conditions, and treatment**

*Arabidopsis thaliana* wild type (ecotype Col-0) and the mutant lines *opr2opr3* (Chini et al., 2018)

and *aos* (*dde2-2*, von Malek et al., 2002) were genotyped using the primers listed (Supplemental

Table 2). After surface sterilization with 4% bleach and three-day stratification at 4°C, seedlings

were grown for ten days in liquid Murashige and Skoog (MS) medium (pH 5.7, 1% sucrose),

whereas adult plants were grown for four weeks in individual pots containing steam-sterilized clay,

coir fiber, and vermiculite. Plant growth took place in Phytocabinets (Percival Scientific,

www.percival-scientific.com/) at 120 $\mu$E m$^{-2}$ s$^{-1}$ under short day conditions (10/14 hours light/dark

cycle) at 21/19 °C and 65% relative humidity. For wounding, seedlings were squeezed with forceps

with serrated teeth. Phytohormone treatments were done by adding either 25 $\mu$M OPDA ($\geq$ 95%,

Cayman Chemical, www.caymanchem.com) or 1 $\mu$M MeJA to the medium.

**RNA isolation, RNA-seq and quantitative RT-PCR analysis**

Total RNA was isolated from homogenized frozen material using the RNeasy Plant Mini Kit (Qiagen) and treated with the DNA-free™ DNA Removal Kit (Invitrogen, #AM1906). First strand cDNA synthesis from 1 μg DNA-free RNA was carried out using the RevertAid H Minus reverse transcriptase with oligo(dT)18 primers (ThermoFisher Scientific™, www.thermofisher.com). Quantitative PCR was carried-out in Hard-Shell® 96-Well PCR Plates (Bio-Rad Laboratories, www.bio-rad.com, #HSP9601) supplied with 10 μL reaction mix of 1.5 ng/μL cDNA, 1x EvaGreen QPCR Mix II (Bio&Sell, www.bio-sell.de, #BS76.580.0200) and 0.2 μM of forward and reverse primers (Supplemental Table 2). The reactions were run on a CFX Connect Real-Time PCR Detection System (Bio-Rad Laboratories) with denaturation (95°C for 15 min), amplification (40 cycles of 95°C for 15 s and 60°C for 30 s) and melt curve analysis (95°C for 10 s, 65°C heating up to 95°C with a heating rate of 0.05°C s$^{-1}$). Gene expression was normalized to the housekeeping gene *PROTEIN PHOSPHATASE 2A SUBUNIT A3* (AT1G13320) (Czechowski et al., 2005; Wang et al., 2014) using the $2^{-\Delta CT}$ method (Schmittgen and Livak, 2008) and included biological triplicates.

**Transcriptome analysis**

RNA quality and integrity were assessed on Agilent 2100 Bioanalyzer system (Agilent Technologies, www.agilent.com) using the RNA 6000 Nano Kit for standard RNA sensitivity (Agilent, #5067-1511). Three biological replicates were submitted to Novogene (www.novogene.com) for mRNA paired-end short-read sequencing (150 bp length) on an Illumina NovaSeq 6000 Sequencing System and bioinformatics analysis according to their pipeline. Reads were mapped to the *A. thaliana* TAIR10 reference genome. Gene expression levels were determined using the FPKM (Fragments per kilobase per million) method. Gene expression heatmaps, principal component analysis (PCA), gene clustering, and Gene Ontology (GO) enrichment analysis, were generated using the iDEP .96 and iDEP 1.1 software tools (Ge et al., 2018).

**Protein extraction and immunoblotting**

Proteins were extracted by incubation of 50 mg frozen material in 200 μl of extraction buffer (25 mM Tris-Cl pH 6.8, 1% SDS, 1% [v/v] β-mercaptoethanol) at 95°C for ten minutes. Extracts were treated with 1% (v/v) Halt™ Protease and Phosphatase Inhibitor Cocktail (ThermoFisher Scientific™, #78440) and quantified at 595 nm on the SPARK® multimode microplate reader (TECAN) using the Pierce™ Bradford Plus Protein Assay Kit (ThermoFisher Scientific™,

#23236). Ten micrograms of total protein were incubated at 96°C for 10 min in Laemmli Buffer (1:1, (Laemmli, 1970) to dissolve AOC trimers (Otto et al., 2016). Proteins were separated on SDS–PAGE (4% and 12% acrylamide for stacking and resolving gels, respectively), transferred to a PVDF membrane and detected using Ponceau S staining or No-Stain™ Protein Labeling Reagent (ThermoFisher Scientific™, #A44449). Membrane blocking in 5% (w/v) BSA in TBST (20 mM Tris–Cl pH 7.8, 150 mM NaCl, 0.05% [v/v] Tween) was followed by immuno-staining using anti-AtAOC primary antibody (1:5000, Stenzel et al., 2003) and a goat anti-rabbit IgG secondary antibody conjugated with alkaline phosphatase (1:4000, Chemicon®, Sigma-Aldrich, www.sigmaaldrich.com, #AP307P). Detection by chemiluminescence with Immun-Star AP Substrate (ThermoFisher Scientific, #1705018) was visualized using a Fusion FX Imaging system (Vilber, www.vilber.com). AOC protein bands were quantified relative to total protein loadings using densitometry analysis with ImageJ (https://imagej.nih.gov/ij/index.html).

**Cloning of the OPRs and plant transformation**

The OPR3 (1176 bp), OPR2 (1125 bp), and OPR1 (1194 bp) coding sequences (CDS) without stop codons were PCR-amplified from Col-0 cDNA using the Q5® High-Fidelity DNA Polymerase (Bio Labs, www.biolabs.io, #M0491) and the listed primer pairs (Supplemental Table 3). For OPR3ΔSRL, the last nine nucleotides were removed, and silent mutations C205G and G1109C were introduced to eliminate *BsaI* and *BpiI* sites, respectively. The CDS were cloned into the Golden Gate level 0 vectors pAGM1287 and pAGM1299 (Engler et al., 2014). OPR fusions to YFP were generated by in-frame assembly of the level 0 with YFP at the C-terminus connected by a Gly-Ser linker. The final cloning cassettes containing the CaMV *35S* promoter and *tOcs* terminator in addition to the Oleosin-RFP plant selection marker were assembled in the pAGM55171 vector.

Plant stable transformation was carried-out by floral dip with *Agrobacterium tumefaciens* strain *GV3101* according to (Zhang et al., 2006). To circumvent the *opr2opr3* mutant sterility, MeJA was applied to flower buds from 48 hours post-dipping until the first silique formation. Transformant selection relied on the seed coat RFP marker and single-insertion lines were identified by segregation analysis determining the ratio of fluorescent to non-fluorescent seeds.

**Subcellular targeting of the OPRs**

Organelle variants of OPR3, OPR2, and OPR1 were generated by adding organelle targeting signals to the N- and/or C-terminal regions of the OPRs and/or YFP sequences (Supplemental Table 1). For nuclear targeting, OPR3ΔSRL was fused to the nuclear localization signal from Simian Virus 40 (Lu et al., 2021; Raikhel, 1992), while its targeting to the cytosol relied on a C-terminal nuclear export signal from the HIV1 REV (Kosugi et al., 2008). ER-localized OPR3ΔSRL was generated by flanking the sequence between an N-terminal signal peptide from the AtCAPE3 (Kunze and Berger, 2015) and a C-terminal HDEL retention signal (De Meyer and Depicker, 2014). Import to the chloroplast stroma and mitochondria, relied on an N-terminal target peptide from the AtRECA1 (Eseverri et al., 2020) and the Rieske protein (von Heijne et al., 1989), respectively.

**Transient expression in *N. benthamiana* leaves and protoplasts.**

*A. tumefaciens* strain *GV3101*, harboring OPR constructs, was cultured for 48 hours in LB medium with the appropriate antibiotics. The bacteria were syringe-infiltrated into *N. benthamiana* leaves according to (Norkunas et al., 2018). Leaf disc imaging was performed 48-72 hours post-infiltration.

Protoplasts were isolated from leaves of 4-week-old *N. benthamiana* plants, as described (Janik et al., 2019). Isolated protoplasts were co-transfected with level 1 plasmids of OPR3 constructs and organelle markers (Stellmach et al., 2022) using a mixture of 200 µl of protoplast suspension and 10 µg of DNA. Protoplasts were incubated at room temperature overnight after PEG-transformation and used for imaging.

**Confocal microscopy**

Confocal microscopy utilized LSM880 and LSM900 laser scanning microscopes (Zeiss Germany, www.zeiss.de). Fluorophores were excited with 514 nm (YFP) and 561 nm (mCherry) laser lines and detected at 510–560 nm and 570-650 nm, respectively. Image acquisition and processing were performed using ZEN software (version 3.4, 2021, Zeiss).

**Phytohormone measurements**

Measurements of OPDA, *dn*-OPDA, JA, and JA-Ile were performed using a standardized Ultra-performance liquid chromatography–tandem Mass Spectrometry (UPLC– MS/MS)-based method (Balcke et al., 2012). 50 mg of powdered frozen tissue were extracted with 500 µl 100% LC-MS methanol supplemented with 50 ng of $[^2H_5]$OPDA, $[^2H_6]$JA, and $[^2H_2]$JA-Ile each as internal

standards. After solid-phase extraction on HR-XC (Chromabond, Macherey-Nagel, www.mn-net.com), 10 µl of the eluate were analyzed via UPLC–MS/MS, and analyte content was determined relative to the internal standard peak heights. *dn*-OPDA content was determined using [$^2$H$_5$]OPDA as the internal standard.

**Determination of enzymatic activity of OPR3**

OPR3 CDS inserted in pET28a(+) with *EcoR*I and *Not*I cloning sites was ordered from BioCat GmbH (www.biocat.com) and transformed into *Escherichia coli* BL21 Star (DE3) cells (ThermoFisher Scientific) by heat shock. Protein expression was performed in auto-induction medium at 16°C for three days (Studier, 2005). Harvested cells were resolved in buffer HisA (50 mM Tris/HCl pH 7.8, 100 mM NaCl, 10% (v/v) glycerol, 2 mM MgCl$_2$ and 5 mM imidazole) supplemented with lysozyme, DNAse I and 0.2 mM PMSF. After a 30-minute incubation on ice, cells were sonicated and cleared lysate was applied to 1 mL HisTrap FastFlow column (Cytiva, www.cytivalifesciences.com). Protein was eluted with buffer HisB (50 mM Tris/HCl pH 7.8, 100 mM NaCl, 10% (v/v) glycerol, 2 mM MgCl$_2$ and 500 mM imidazole), which was changed to storage buffer (50 mM Tris/HCl pH 7.8, 100 mM NaCl, 10% (v/v) glycerol, 2 mM MgCl$_2$) by dialysis (MWCO 12-14 kDa) at 4°C overnight. Following SDS-PAGE, protein fractions were used for *in vitro* assays at 0.1 mg/mL together with 0.05 mM substrate, 1 mM NADPH and 10 mM NaCl in 20 mM Tris/HCl pH 7.8. Reactions were incubated at 30°C for 10 minutes and stopped by one volume acetonitrile. Enzyme deactivated by one volume acetonitrile was used as control. Assays were analyzed via UHPLC-high resolution (HR)-MS according to Feussner et al. (2023). Data was analyzed using the MassHunter Qualitative Analysis 10.0 software (Agilent Technologies).

**Figures and statistical analysis**

Micrographs and photographs were processed through PHOTOSHOP 12.0.4 (Adobe Systems, http://www.adobe.com). The statistical analyses applied to the different datasets were performed using GraphPad Prism (www.graphpad.com).

**Data availability**

The datasets generated and/or analyzed during this study are available from the corresponding author on request. Transcriptomic raw data were deposited in Sequence Read Archive (SRA) of NCBI (PRJNA1088739).

**Funding**

K.M. and B.H. were supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) grant No 400681449/GRK2498. M.K. was supported by the DFG grant No 273134146/GRK 2172. I.F. acknowledges funding from the DFG (GRK 2172-PRoTECT, INST 186/1434-1, and ZUK 45/2010).

**Author contributions**

Conceptualization, B.H and K.M.; Methodology, K.M, R.B, and F.S.; Investigation, K.M., R.B., F.S., and M.K.; Analyzing Data, I.F. and B.H.; Writing – Original Draft, K.M.; Writing – Review & Editing, B.H. and I.F.; Funding Acquisition, B.H. and I.F.;

**Acknowledgments**

We thank Hagen Stellmach (IPB Halle, Germany) for help in jasmonate quantification, Andreas Schaller (University of Hohenheim, Germany) and Mats Hamberg (Karolinska Institute, Sweden) for providing seeds of *opr2opr3* and 4,5-ddh-JA, respectively. We thank Sylvestre Marillonnet (IPB Halle) for providing the Golden Gate modules and vectors used in this study. Khabat Vahabi (IPB Halle) is acknowledged for assistance in RNAseq preparation and data analysis Claus Wasternack (IPB Halle) and Alain Tissier (IPB Halle) are acknowledged for critical reading of the manuscript.

**Declaration of interests**

632     The authors declare no competing interests.

## References

**Adhikari, A., and Park, S.-W.** (2023). Reduced GSH acts as a metabolic cue of OPDA signaling in coregulating photosynthesis and defense activation under stress. Plants **12**:3745.

**Alméras, E., Stolz, S., Vollenweider, S., Reymond, P., Mène-Saffrané, L., and Farmer, E.** (2003). Reactive electrophile species activate defense gene expression in Arabidopsis. Plant J. **34**:205-216.

**Andreou, A., and Feussner, I.** (2009). Lipoxygenases - Structure and reaction mechanism. Phytochemistry **70**:1504-1510.

**Balcke, G., Handrick, V., Bergau, N., Fichtner, M., Henning, A., Stellmach, H., Tissier, A., Hause, B., and Frolov, A.** (2012). An UPLC-MS/MS method for highly sensitive high-throughput analysis of phytohormones in plant tissues. Plant Methods **8**:47.

**Bannenberg, G., Martínez, M., Hamberg, M., and Castresana, C.** (2009). Diversity of the enzymatic activity in the lipoxygenase gene family of *Arabidopsis thaliana*. Lipids **44**:85-95. 10.1007/s11745-008-3245-7.

**Browse, J.** (2009). The power of mutants for investigating jasmonate biosynthesis and signaling. Phytochemistry **70**:1539-1546.

**Chang, Y., Shi, M., Sun, Y., Cheng, H., Ou, X., Zhao, Y., Zhang, X., Day, B., Miao, C., and Jiang, K.** (2023). Light-induced stomatal opening in Arabidopsis is negatively regulated by chloroplast-originated OPDA signaling. Curr. Biol. **33**:1071-1081.e1075. https://doi.org/10.1016/j.cub.2023.02.012.

**Chini, A., Monte, I., Zamarreño, A.M., García-Mina, J.M., and Solano, R.** (2023). Evolution of the jasmonate ligands and their biosynthetic pathways. New Phytol. **238**:2236-2246. https://doi.org/10.1111/nph.18891.

**Chini, A., Fonseca, S., Fernandez, G., Adie, B., Chico, J.M., Lorenzo, O., Garcia-Casado, G., Lopez-Vidriero, I., Lozano, F.M., Ponce, M.R., et al.** (2007). The JAZ family of repressors is the missing link in jasmonate signalling. Nature **448**:666-671.

**Chini, A., Monte, I., Zamarreño, A.M., Hamberg, M., Lassueur, S., Reymond, P., Weiss, S., Stintzi, A., Schaller, A., Porzel, A., et al.** (2018). An OPR3-independent pathway uses 4,5-didehydrojasmonate for jasmonate synthesis. Nature Chem. Biol. **14**:171. 10.1038/nchembio.2540.

**Chung, H.S., Koo, A.J.K., Gao, X., Jayanty, S., Thines, B., Jones, A.D., and Howe, G.A.**
(2008). Regulation and function of Arabidopsis *JASMONATE ZIM*-domain genes in
response to wounding and herbivory. Plant Physiol. **146**:952-964.

**Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K., and Scheible, W.-R.** (2005). Genome-
wide identification and testing of superior reference genes for transcript normalization in
Arabidopsis. Plant Physiol. **139**:5-17. 10.1104/pp.105.063743.

**Dave, A., and Graham, I.A.** (2012). Oxylipin signalling: a distinct role for the jasmonic acid
precursor *cis*-(+)-12-oxo-phytodienoic acid (*cis*-OPDA). Front. Plant Sci.
**3:42**10.3389/fpls.2012.00042.

**Dave, A., Hernández, M.L., He, Z., Andriotis, V.M.E., Vaistij, F.E., Larson, T.R., and**
**Graham, I.A.** (2011). 12-Oxo-phytodienoic acid accumulation during seed development
represses seed germination in Arabidopsis. Plant Cell **23**:583-599. 10.1105/tpc.110.081489.

**Davletova, S., Schlauch, K., Coutu, J., and Mittler, R.** (2005). The Zinc-Finger Protein Zat12
plays a central role in reactive oxygen and abiotic stress signaling in Arabidopsis  Plant
Physiol. **139**:847-856. 10.1104/pp.105.068254.

**Davoine, C., Falletti, O., Douki, T., Iacazio, G., Ennar, N., Montillet, J.-L., and**
**Triantaphylides, C.** (2006). Adducts of oxylipin electrophiles to glutathione reflect a 13
specificity of the downstream lipoxygenase pathway in the tobacco hypersensitive
response. Plant Physiol. **140**:1484-1493.

**De Meyer, T., and Depicker, A.** (2014). Trafficking of endoplasmic reticulum-retained
recombinant proteins is unpredictable in *Arabidopsis thaliana*. Front. Plant Sci.
**5**10.3389/fpls.2014.00473.

**Delessert, C., Wilson, I., Straeten, D., Dennis, E., and Dolferus, R.** (2004). Spatial and temporal
analysis of the local response to wounding. Plant Mol. Biol. **55**:165-181.

**Engler, C., Youles, M., Gruetzner, R., Ehnert, T.-M., Werner, S., Jones, J.D.G., Patron, N.J.,**
**and Marillonnet, S.** (2014). A Golden Gate modular cloning toolbox for plants. ACS
Synthetic Biol. **3**:839-843. 10.1021/sb4001504.

**Eseverri, Á., Baysal, C., Medina, V., Capell, T., Christou, P., Rubio, L.M., and Caro, E.**
(2020). Transit peptides from photosynthesis-related proteins mediate import of a marker
protein into different plastid types and within different species. Front. Plant Sci.
**11**10.3389/fpls.2020.560701.

**Feussner, I., and Wasternack, C.** (2002). The lipoxygenase pathway. Annu. Rev. Plant Biol.
**53**:275-297.

**Feussner, K., Abreu, I.N., Klein, M., and Feussner, I.** (2023). Chapter twelve - metabolite
fingerprinting: A powerful metabolomics approach for marker identification and functional
gene annotation. In Methods Enzymol., J. Jez, ed. (Academic Press: pp. 325-350.
https://doi.org/10.1016/bs.mie.2022.08.015.

**Ge, S.X., Son, E.W., and Yao, R.** (2018). iDEP: an integrated web application for differential
expression and pathway analysis of RNA-Seq data. BMC Bioinformatics **19**:534.
10.1186/s12859-018-2486-6.

**Goetz, S., Hellwege, A., Stenzel, I., Kutter, C., Hauptmann, V., Forner, S., McCaig, B., Hause,**
**G., Miersch, O., Wasternack, C., et al.** (2012). Role of cis-12-oxo-phytodienoic acid in
tomato embryo development. Plant Physiol. **158**:1715-1727. 10.1104/pp.111.192658.

**Goodspeed, D., Chehab, E.W., Min-Venditti, A., Braam, J., and Covington, M.F.** (2012).
Arabidopsis synchronizes jasmonate-mediated defense with insect circadian behavior.
Proc. Natl. Acad. Sci. USA **109**:4674-4677. 10.1073/pnas.1116368109.

**Gould, S.J., Keller, G.A., Hosken, N., Wilkinson, J., and Subramani, S.** (1989). A conserved
tripeptide  sorts  proteins  to  peroxisomes.  J.  Cell  Biol.  **108**:1657-1664.
10.1083/jcb.108.5.1657.

**Guo, Q., Yoshida, Y., Major, I.T., Wang, K., Sugimoto, K., Kapali, G., Havko, N.E., Benning,**
**C., and Howe, G.A.** (2018). JAZ repressors of metabolic defense promote growth and
reproductive fitness in *Arabidopsis*. Proc. Natl. Acad. Sci. USA **115**:E10768-E10777.
10.1073/pnas.1811828115.

**Howe, G.A.** (2004). Jasmonates as signals in the wound response. J. Plant Growth Regul. **23**:223-
237.

**Janik, K., Stellmach, H., Mittelberger, C., and Hause, B.** (2019). Characterization of
phytoplasmal effector protein interaction with proteinaceous plant host targets using
bimolecular fluorescence complementation (BiFC). In Phytoplasmas: Methods and
Protocols, R. Musetti and L. Pagliari, eds. (Springer New York: New York, NY), pp. 321-
331. 10.1007/978-1-4939-8837-2_24

**Jimenez-Aleman, G.H., Thirumalaikumar, V.P., Jander, G., Fernie, A.R., and Skirycz, A.**
(2022). OPDA, more than just a jasmonate precursor. Phytochemistry **204**:113432.
https://doi.org/10.1016/j.phytochem.2022.113432.

Koo, A.J.K., and Howe, G.A. (2009). The wound hormone jasmonate. Phytochemistry **70**:1571-1580.

Kosugi, S., Hasebe, M., Tomita, M., and Yanagawa, H. (2008). Nuclear export signal consensus sequences defined using a localization-based yeast selection system. Traffic **9**:2053-2062. https://doi.org/10.1111/j.1600-0854.2008.00825.x.

Kunze, M., and Berger, J. (2015). The similarity between N-terminal targeting signals for protein import into different organelles and its evolutionary relevance. Front. Physiol. **6**10.3389/fphys.2015.00259.

Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature **227**:680-685.

Lewandowska, M., Zienkiewicz, A., Feussner, K., König, S., Kunst, L., and Feussner, I. (2023). Wound-induced triacylglycerol biosynthesis is jasmonoy-l-isoleucin and abscisic acid independent. Plant Biol. **25**:509-517. https://doi.org/10.1111/plb.13513.

Li, C., Xu, M., Cai, X., Han, Z., Si, J., and Chen, D. (2022). Jasmonate signaling pathway modulates plant defense, growth, and their trade-offs. Int. J. Mol. Sci. **23**:3945.

Liu, W., and Park, S.-W. (2021). 12-oxo-Phytodienoic acid: A fuse and/or switch of plant growth and defense responses? Front. Plant Sci. **12**10.3389/fpls.2021.724079.

Lu, J., Wu, T., Zhang, B., Liu, S., Song, W., Qiao, J., and Ruan, H. (2021). Types of nuclear localization signals and mechanisms of protein import into the nucleus. Cell Comm. Signal. **19**:60. 10.1186/s12964-021-00741-y.

Mehrshahi, P., Stefano, G., Andaloro, J.M., Brandizzi, F., Froehlich, J.E., and DellaPenna, D. (2013). Transorganellar complementation redefines the biochemical continuity of endoplasmic reticulum and chloroplasts. Proc. Natl. Acad. Sci. USA **110**:12126-12131. 10.1073/pnas.1306331110.

Mittler, R., Kim, Y., Song, L., Coutu, J., Coutu, A., Ciftci-Yilmaz, S., Lee, H., Stevenson, B., and Zhu, J.-K. (2006). Gain- and loss-of-function mutations in Zat10 enhance the tolerance of plants to abiotic stress. FEBS Lett. **580**:6537-6542. https://doi.org/10.1016/j.febslet.2006.11.002.

Mizoi, J., Kanazawa, N., Kidokoro, S., Takahashi, F., Qin, F., Morimoto, K., Shinozaki, K., and Yamaguchi-Shinozaki, K. (2019). Heat-induced inhibition of phosphorylation of the stress-protective transcription factor DREB2A promotes thermotolerance of *Arabidopsis thaliana*. J. Biol. Chem. **294**:902-917. https://doi.org/10.1074/jbc.RA118.002662.

**Monte, I., Caballero, J., Zamarreño, A.M., Fernández-Barbero, G., García-Mina, J.M., and**
**Solano, R.** (2022). JAZ is essential for ligand specificity of the COI1/JAZ co-receptor.
Proc. Natl. Acad. Sci. USA **119**:e2212155119. doi:10.1073/pnas.2212155119.

**Monte, I., Kneeshaw, S., Franco-Zorrilla, J.M., Chini, A., Zamarreño, A.M., García-Mina,**
**J.M., and Solano, R.** (2020). An ancient COI1-independent function for reactive
electrophilic oxylipins in thermotolerance. Curr. Biol. **30**:962-971.e963.
10.1016/j.cub.2020.01.023.

**Monte, I., Ishida, S., Zamarreño, A.M., Hamberg, M., Franco-Zorrilla, J.M., García-Casado,**
**G., Gouhier-Darimont, C., Reymond, P., Takahashi, K., García-Mina, J.M., et al.**
(2018). Ligand-receptor co-evolution shaped the jasmonate pathway in land plants. Nature
Chem. Biol. **14**:480-488. 10.1038/s41589-018-0033-4.

**Mueller, M.J., and Berger, S.** (2009). Reactive electrophilic oxylipins: Pattern recognition and
signalling. Phytochemistry **70**:1511-1521.

**Mueller, S., Hilbert, B., Dueckershoff, K., Roitsch, T., Krischke, M., Mueller, M.J., and**
**Berger, S.** (2008). General detoxification and stress responses are mediated by oxidized
lipids through TGA transcription factors in *Arabidopsis*. Plant Cell **20**:768-785.

**Müller, S.M., Wang, S., Telman, W., Liebthal, M., Schnitzer, H., Viehhauser, A., Sticht, C.,**
**Delatorre, C., Wirtz, M., Hell, R., et al.** (2017). The redox-sensitive module of cyclophilin
20-3, 2-cysteine peroxiredoxin and cysteine synthase integrates sulfur metabolism and
oxylipin signaling in the high light acclimation response. Plant J. **91**:995-1014.
10.1111/tpj.13622.

**Norkunas, K., Harding, R., Dale, J., and Dugdale, B.** (2018). Improving agroinfiltration-based
transient gene expression in *Nicotiana benthamiana*. Plant Methods **14**:71.
10.1186/s13007-018-0343-2.

**Ohkama-Ohtsu, N., Sasaki-Sekimoto, Y., Oikawa, A., Jikumaru, Y., Shinoda, S., Inoue, E.,**
**Kamide, Y., Yokoyama, T., Hirai, M.Y., Shirasu, K., et al.** (2011). 12-Oxo-phytodienoic
acid-glutathione conjugate is transported into the vacuole in Arabidopsis. Plant Cell
Physiol. **52**:205-209.

**Otto, M., Naumann, C., Brandt, W., Wasternack, C., and Hause, B.** (2016). Activity regulation
by heteromerization of Arabidopsis allene oxide cyclase family members. Plants **5**:3.

**Park, S.-W., Li, W., Viehhauser, A., He, B., Kim, S., Nilsson, A.K., Andersson, M.X., Kittle,**
**J.D., Ambavaram, M.M.R., Luan, S., et al.** (2013). Cyclophilin 20-3 relays a 12-oxo-

phytodienoic acid signal during stress responsive regulation of cellular redox homeostasis.

Proc. Natl. Acad. Sci. USA **110**:9559-9564. 10.1073/pnas.1218872110.

**Pollmann, S., Springer, A., Rustgi, S., von Wettstein, D., Kang, C., Reinbothe, C., and**

**Reinbothe, S.** (2019). Substrate channeling in oxylipin biosynthesis through a protein

complex in the plastid envelope of *Arabidopsis thaliana*. J. Exper. Bot. **70**:1483-1495.

10.1093/jxb/erz015.

**Raikhel, N.** (1992). Nuclear targeting in plants 1. Plant Physiol. **100**:1627-1632.

10.1104/pp.100.4.1627.

**Reymond, P., Weber, H., Damond, M., and Farmer, E.** (2000). Differential gene expression in

response to mechanical wounding and insect feeding in Arabidopsis. Plant Cell **12**:707-719.

**Ribot, C., Zimmerli, C., Farmer, E.E., Reymond, P., and Poirier, Y.** (2008). Induction of the

Arabidopsis *PHO1;H10* gene by 12-oxo-phytodienoic acid but not jasmonic acid via a

*CORONATINE INSENSITIVE1*-dependent pathway. Plant Physiol. **147**:696-706.

**Sakuma, Y., Maruyama, K., Qin, F., Osakabe, Y., Shinozaki, K., and Yamaguchi-Shinozaki,**

**K.** (2006). Dual function of an *Arabidopsis* transcription factor DREB2A in water-stress-

responsive and heat-stress-responsive gene expression. Proc. Natl. Acad. Sci. USA

**103**:18822-18827. doi:10.1073/pnas.0605639103.

**Savchenko, T., and Dehesh, K.** (2014). Drought stress modulates oxylipin signature by eliciting

12-OPDA as a potent regulator of stomatal aperture. Plant Signal. Behavior **9**:e28304.

**Schaller, A., and Stintzi, A.** (2009). Enzymes in jasmonate biosynthesis - Structure, function,

regulation. Phytochemistry **70**:1532-1538.

**Schaller, F., Biesgen, C., Müssig, C., Altmann, T., and Weiler, E.W.** (2000). 12-

Oxophytodienoate reductase 3 (OPR3) is the isoenzyme involved in jasmonate

biosynthesis. Planta **210**:979-984.

**Schmittgen, T.D., and Livak, K.J.** (2008). Analyzing real-time PCR data by the comparative

C(T) method. Nature Protocols **3**:1101-1108.

**Sheard, L.B., Tan, X., Mao, H., Withers, J., Ben-Nissan, G., Hinds, T.R., Kobayashi, Y., Hsu,**

**F.-F., Sharon, M., Browse, J., et al.** (2010). Jasmonate perception by inositol-phosphate-

potentiated COI1-JAZ co-receptor. Nature **468**:400-405.

**Stellmach, H., Hose, R., Räde, A., Marillonnet, S., and Hause, B.** (2022). A new set of Golden-

Gate-based organelle marker plasmids for colocalization studies in plants. Plants **11**:2620.

**Stelmach, B.A., Müller, A., Hennig, P., Laudert, D., Andert, L., and Weiler, E.W.** (1998).
Quantitation of the octadecanoid 12-oxo-phytodienoic acid, a signalling compound in plant
mechanotransduction. Phytochemistry **47**:539-546.

**Stenzel, I., Hause, B., Miersch, O., Kurz, T., Maucher, H., Weichert, H., Ziegler, J., Feussner,**
**I., and Wasternack, C.** (2003). Jasmonate biosynthesis and the allene oxide cyclase family
of *Arabidopsis thaliana*. Plant Mol. Biol. **51**:895-911.

**Stintzi, A., and Browse, J.** (2000). The Arabidopsis male-sterile mutant, *opr3*, lacks the 12-
oxophytodienoic acid reductase required for jasmonate synthesis. Proc. Natl. Acad. Sci.
USA **97**:10625-10630.

**Stintzi, A., Weber, H., Reymond, P., Browse, J., and Farmer, E.** (2001). Plant defense in the
absence of jasmonic acid: The role of cyclopentanones. Proc. Natl. Acad. Sci. USA
**98**:12837-12842.

**Studier, F.W.** (2005). Protein production by auto-induction in high density shaking cultures.
Protein Expr. Purif. **41**:207-234. 10.1016/j.pep.2005.01.016.

**Taki, N., Sasaki-Sekimoto, Y., Obayashi, T., Kikuta, A., Kobayashi, K., Ainai, T., Yagi, K.,**
**Sakurai, N., Suzuki, H., Masuda, T., et al.** (2005). 12-oxo-phytodienoic acid triggers
expression of a distinct set of genes and plays a role in wound-induced gene expression in
Arabidopsis. Plant Physiol. **139**:1268-1283.

**Thines, B., Katsir, L., Melotto, M., Niu, Y., Mandaokar, A., Liu, G., Nomura, K., He, S.Y.,**
**Howe, G.A., and Browse, J.** (2007). JAZ repressor proteins are targets of the SCF[COI1]
complex during jasmonate signalling. Nature **448**:661-665.

**Verhage, A., Vlaardingerbroek, I., Raaijmakers, C., Van Dam, N., Dicke, M., Van Wees,**
**S.C.M., and Pieterse, C.M.J.** (2011). Rewiring of the jasmonate signaling pathway in
Arabidopsis during insect herbivory. Front. Plant Sci. **2**10.3389/fpls.2011.00047.

**von Heijne, G., Steppuhn, J., and Herrmann, R.G.** (1989). Domain structure of mitochondrial
and chloroplast targeting peptides. Europ. J. Biochem. **180**:535-545.
https://doi.org/10.1111/j.1432-1033.1989.tb14679.x.

**von Malek, B., van der Graaff, E., Schneitz, K., and Keller, B.** (2002). The Arabidopsis male-
sterile mutant *dde2-2* is defective in the *ALLENE OXIDE SYNTHASE* gene encoding one
of the key enzymes of the jasmonic acid biosynthesis pathway. Planta **216**:187-192.

**Wang, H., Wang, J., Jiang, J., Chen, S., Guan, Z., Liao, Y., and Chen, F.** (2014). Reference
genes for normalizing transcription in diploid and tetraploid Arabidopsis. Sci. Rep. **4**:6781.
10.1038/srep06781.

**Wang, J., Wu, D., Wang, Y., and Xie, D.** (2019). Jasmonate action in plant defense against
insects. J. Exper. Bot. **70**:3391-3400. 10.1093/jxb/erz174.

**Wasternack, C.** (2006). Oxylipins: biosynthesis, signal transduction and action. In Plant Hormone
Signaling, P. Hedden and S. Thomas, eds. (Blackwell Publishing: Harpenden), pp. 185-228.

**Wasternack, C.** (2019). New light on local and systemic wound signaling. Trends Plant Sci.
**24**:102-105. https://doi.org/10.1016/j.tplants.2018.11.009.

**Wasternack, C., and Hause, B.** (2013). Jasmonates: biosynthesis, perception, signal transduction
and action in plant stress response, growth and development. An update to the 2007 review
in Annals of Botany. Ann. Bot. **111**:1021-1058. 10.1093/aob/mct067.

**Wasternack, C., and Strnad, M.** (2016). Jasmonate signaling in plant stress responses and
development – active and inactive compounds. New Biotechnol. **33**:604-613.
http://dx.doi.org/10.1016/j.nbt.2015.11.001.

**Weber, H., Vick, B., and Farmer, E.** (1997). Dinor-oxo-phytodienoic acid: A new hexadecanoid
signal in the jasmonate family. Proc. Natl. Acad. Sci. USA **94**:10473-10478.

**Xie, D.-X., Feys, B., James, S., Nieto-Rostro, M., and Turner, J.** (1998). COI1: An Arabidopsis
gene required for jasmonate-regulated defense and fertility. Science **280**:1091-1094.

**Xu, L., Liu, F., Lechner, E., Genschik, P., Crosby, W., Ma, H., Peng, W., Huang, D., and Xie,**
**D.** (2002). The SCF-coi1 ubiquitin-ligase complexes are required for jasmonate response
in Arabidopsis. Plant Cell **14**:1919-1935.

**Yan, J., Yao, R., Chen, L., Li, S., Gu, M., Nan, F., and Xie, D.** (2018). Dynamic perception of
jasmonates by the F-box protein COI1. Mol. Plant **11**:1237-1247.
10.1016/j.molp.2018.07.007.

**Zhang, G., Liu, W., Gu, Z., Wu, S., E, Y., Zhou, W., Lin, J., and Xu, L.** (2023). Roles of the
wound hormone jasmonate in plant regeneration. J. Exper. Bot. **74**:1198-1206.
10.1093/jxb/erab508.

**Zhang, X., Henriques, R., Lin, S.-S., Niu, Q.-W., and Chua, N.-H.** (2006). *Agrobacterium*-
mediated transformation of *Arabidopsis thaliana* using the floral dip method. Nature
Protocols **1**:641-646. 10.1038/nprot.2006.97.

**Supplemental data**

**Figure S1**: Seedlings of *opr2opr3* mutant show diminished AOC proteins content and produce lower OPDA levels compared to wild type (Col-0).

**Figure S2**: The experimental setup for transcriptome analyses.

**Figure S3**: Hierarchical clustering and principal component analysis (PCA) illustrating the wound-induced transcriptional changes in Col-0, *opr2opr3* and *aos*.

**Figure S4**: The JA-dependent wound response occurring in wild type seedlings.

**Figure S5**: Clustering of the transcriptional response of seedlings from *opr2opr3* upon treatment with 25 µM OPDA with the response of seedlings from Col-0, *opr2opr3* and *aos* upon wounding.

**Figure S6**: The concept of the trans-organellar complementation of the *opr2opr3* mutant.

**Figure S7:** The removal of the SRL peroxisomal signal results in a cytosolic localization of OPR3 in *N. benthamiana* leaves and targeting of OPR3ΔSRL to the nucleus, cytosol, plastid stroma and mitochondria in stable transformed *opr2opr3* lines.

**Figure S8**: OPR3 targeted to the ER, cytosol, nucleus, and chloroplasts colocalizes with the corresponding organelle markers in *N. benthamiana* protoplasts.

**Figure S9**: OPR3 reduces OPDA and 4,5-ddh-JA *in vitro*.

**Figure S10**: OPDA levels at 1 h after wounding of seedlings of wild type (WT) or *opr2opr3* transformed with either empty vector (EV), *35S::OPR2* or *35S::OPR1* targeted to peroxisomes (Px), cytosol (Cyt) or mitochondria (Mt).

**Table S1:** Peptide sequences of the targeting signals used for subcellular targeting.

**Table S2:** Primers sequences used for mutants genotyping and RT-qPCR.

**Table S3:** Primer sequences used for Golden Gate cloning of OPR3, OPR2 and OPR1 and their subcellular targeting.

# Figures

**Figure 1: MeJA pretreatment restitutes AOC protein levels in the mutants and OPDA content in the *opr2opr3* mutant.**

**(A)** Synthesis of jasmonic acid (JA)/JA-Ile from α-linolenic acid (α-LeA). The biosynthesis starts with the release of α-LeA from chloroplast membrane galactolipids and its conversion to (13S)-hydroperoxyoctadecatrienoic acid (13-HPOT) by 13-LIPOXYGENASE (LOX). *cis*-(+)-12-oxophytodienoic acid (*cis*-OPDA) is formed in the chloroplast (green background) from 13-HPOT through sequential actions of ALLENE OXIDE SYNTHASE (AOS) and ALLENE OXIDE CYCLASE (AOC). In the peroxisomes (yellow background) reduction of cyclopentenone ring of *cis*-OPDA is catalyzed by a peroxisomal OPDA REDUCTASE (OPR3) resulting in the formation of 3-oxo-2-(20(Z)-pentenyl)-cyclopentane-1-octanoic (OPC-8:0). Shortening of the octanoic acid side chains of OPC-8:0 by three rounds of β-oxidation yields JA. The cytosol-released JA is conjugated to Isoleucine (Ile) by the enzyme JASMONATE RESISTANT 1 (JAR1) resulting in the major bulk formation of the biologically active JA-Ile. *cis*-OPDA can directly undergo three rounds of β-oxidation yielding 4,5-didehydro-JA (4,5-ddh-JA) in a parallel OPR3-bypass pathway. The formed 4,5-ddh-JA is released to the cytosol and then reduced to JA by OPR2 and results in JA-Ile formation. Enzymes shaded in green, yellow and blue are located in the plastids, peroxisomes and cytosol, respectively.

**(B)** AOC protein content in seedlings pretreated with 1 µM MeJA or water (Mock) visualized by immunoblot, equal loading is depicted by staining with No-Stain™ Protein Labeling Reagent.

**(C)** AOC band intensity detected by immunoblot and quantified relative to total protein loading determined by Ponceau S.

**(D)** RT-qPCR of JA-responsive genes *LOX2*, *AOS*, *AOC2* and *OPR3* showing transcript accumulation at control condition (-wounding) and at 1 h after wounding with forceps. Seedlings were either pretreated with MeJA or water (Mock) during development.

**(E)** levels of *cis*-OPDA, *dn*-OPDA, JA and JA-Ile in seedlings of wild-type (Col-0), *opr2opr3*, and *aos* at control condition (-wounding) and 1 h after wounding with forceps.

Seedlings were either pretreated with MeJA or water (Mock) during development (see Supplemental Figure 2A for experimental set-up). Bars represent means of three biological replicates with 120 seedlings each (single dots) ±SEM. Statistically significant differences among genotypes within each condition were calculated using One-Way ANOVA followed by Tukey-HSD and are indicated by asterisks with $*p<0.05$, $**p<0.01$, $***p<0.001$, and $****p<0.0001$.

Abbreviations for compounds: 13-HPOT, (13S)-hydroperoxyoctadecatrienoic acid; cis-(+)-OPDA, *cis*-(+)-12- oxophytodienoic acid; dn-OPDA, dinor-12-oxo phytodienoic acid, OPC-8:0, 3-oxo-2-(2-pentenyl)- cyclopentane-1-octanoic acid; 4,5-ddh-JA, 4,5-didehydrojasmonic acid; JA, jasmonic acid; JA-Ile, jasmonoyl isoleucine.

Abbreviations for enzymes: LOX, lipoxygenase; AOS, allene oxide synthase; AOC, allene oxide cyclase; OPR3, 12-oxophytodienoate-10,11-reductase 3; OPR2, 12-oxophytodienoate-10,11-reductase 2; JAR1, jasmonoyl-isoleucine synthetase.

**Figure 2: The JA/OPDA-independent wound response of seedlings.** Ten-day old seedlings from Col-0, *opr2opr3* and *aos* were pretreated with MeJA during development and were either unwounded (control) or harvested 1 h post wounding with forceps (wounding).

**(A)** Venn diagram showing the wound response in Col-0 compared to *opr2opr3* and *aos* mutants. Differentially expressed genes (DEGs) after wounding compared to non-wounded control for each genotype showed a common response with 1460 DEGs. DEGs were identified using FDR and FC cutoffs of 0.05 and 2, respectively.

**(B)** Hierarchical cluster tree summarizing significant biological pathways enriched in the 1460 DEGs common to all genotypes from (A). Gene ontology analysis was performed based on biological process.

**(C)** Heatmap illustrating 50 out of 211 DEGs enriched in the GO term 'response to stress' from **(B)**. Average FPKM values of three independent biological replicates were transformed into row normalized Z-Score.

**(D)** RT-qPCR validation of DEGs previously described as OPDA-responsive genes. Transcript accumulation of *FAD-OXR*, *DREB2A*, *ZAT10* and *GRX480* in Col-0, *opr2opr3* and *aos* at control and wounding conditions. Transcript levels were normalized to those of *PP2A*. Bars represent means of three biological replicates ±SEM. Statistically significant differences among genotypes within each condition were calculated using One-Way ANOVA followed by Tukey-HSD and are indicated by asterisks with ****$p<0.0001$; ns = non-significant.

**Figure 3: The OPDA-specific transcriptional change deduced from transcriptomic comparison between seedlings of *opr2opr3* and aos at control and wounding conditions.** Ten-day old seedlings from *opr2opr3* and *aos* were pretreated with MeJA during development and were either unwounded (control, C) or harvested 1 h post wounding with forceps (wounding, W).

**(A)** Venn diagram representing genes showing differential expression when *opr2opr3* is compared to *aos* at control and wounding conditions (FDR cutoff = 0.05 and FC cutoff = 2).

**(B)** Heatmap illustrating the expression levels of the eleven wounding-related DEGs from (A). The average FPKM values of three independent biological replicates were transformed into row-normalized Z-Score.

**(C)** Transcript accumulation of *AOS*, *OPR2,* and *OPR3* selected from **(B)** and determined by RT-qPCR. Transcript levels were normalized to those of *PP2A*. Bars represent means of three biological replicates (single dots; ±SEM). Asterisks denote statistically significant differences among genotypes as determined by Student's t-test within each condition (** p<0.01, *** p<0.001, **** p<0.0001).

**Figure 4: Application of OPDA results in the induction of JA-Ile signaling and sulfur metabolism.**

**(A)** Heatmap of DEGs showing enrichment in JA signaling pathway genes and genes involved in sulfur assimilation in seedlings of *opr2opr3* following application of 25 μM OPDA for 30 min. DEGs were plotted and compared to those from wounded seedlings of Col-0, *opr2opr3* and *aos*

using average FPKM values transformed into row-normalized Z-score of three biological replicates.

**(B)** RT-qPCR validation of the up-regulation of JA-responsive genes *JAZ2*, *JAZ7*, *JAZ13* and *CLH1* in *opr2opr3* seedlings treated with OPDA in comparison to wounded *opr2opr3* seedlings. OPDA-treated and wounded *opr2opr3* seedlings were compared to their respective controls, non-wounded (control) and water-treated seedlings, respectively.

**(C)** *cis*-OPDA, *dn*-OPDA, JA and JA-Ile accumulation in *opr2opr3* seedlings treated with OPDA in comparison to wounded *opr2opr3* seedlings producing OPDA endogenously.

**(D)** RT-qPCR validation of the up-regulation of the sulfur metabolism genes *SDI1*, *ORA59*, *APR1* and *APS3* in *opr2opr3* seedlings treated with OPDA in comparison to wounded *opr2opr3* seedlings. OPDA-treatment and wounding of seedlings were compared to seedlings which were non-wounded (control) and water-treated (water), respectively.

Bars **(B-D)** represent means of three biological replicates with 120 seedlings each (single dots; ±SEM). Statistically significant differences among treatments were calculated using One-Way ANOVA followed by Tukey-HSD and are indicated by asterisks with $*p<0.05$, $**p<0.01$, $***p<0.001$, and $****p<0.0001$; ns = non-significant.

**Figure 5: Restitution of the fertility and JA/JA-Ile levels of *opr2opr3* plants by OPR3-targeted to different organelles.**

**(A-H)** Open flower buds showing stamen filament elongation and pollen release by anthers (scale bars represent 0.5 mm) and plant inflorescences showing silique formation at days 50-60 post germination with asterisks marking fully developed siliques. Col-0 and *opr2opr3* transformed with an empty vector **(A, B)** show fertility and sterility, respectively. *opr2opr3* T2 lines complemented with peroxisome- **(C)**, nucleus- **(D)**, cytosol- **(E)**, ER- **(F)**, plastid stroma- **(G)** and mitochondria- **(H)** targeted OPR3 fused to YFP show the rescue of the *opr2opr3* sterility phenotype.

**(I)** *cis*-OPDA, JA and JA-Ile accumulation at 1 h after wounding of seedlings of wild type (WT) or *opr2opr3* transformed with either empty vector (EV) or *35S::OPR3* targeted to peroxisomes (Px), cytosol (Cyt) or mitochondria (Mt). Numbers refer to independent transgenic lines, which were selected for single insertion and homozygosity. Bars represent means of four biological replicates with 120 seedlings each (single dots; ±SEM). Statistically significant differences between *opr2opr3* transformed with empty vector and transformed with the OPR3 constructs were calculated using One-Way ANOVA followed by Tukey HSD ($p < 0.05$) and are denoted by different letters.

**Figure 6: Restitution of the fertility and JA/JA-Ile levels of *opr2opr3* plants by OPR2, but not by OPR1, both targeted to different organelles.**

Stable *opr2opr3* complementation lines expressing OPR2 (left) or OPR1 (right) targeted to the cytosol (without any signal peptide), to the peroxisomes (OPR-SKL) or to mitochondria (mTP-OPR), all under control of the CaMV *35S* promoter.

**(A)** Micrographs showing the correct localization of the respective OPR-YFP-fusion. The yellow signal corresponds to OPR-YFP, whereas the magenta signal corresponds to chlorophyll autofluorescence. Scale bars represent 50 μm.

**(B, C)** Open flower buds to show stamen filament elongation and pollen release by anthers **(B)** and plant inflorescences showing silique formation at days 50-60 post germination **(C)**. Note that transformation of *opr2opr3* with all variants of OPR2 and with OPR1 targeted to peroxisomes resulted in a rescue of fertility visible by elongated filaments and fully developed siliques. Targeting of OPR1 to cytosol led to partial rescue shown by few fully developed siliques only (red

asterisks), whereas targeting of OPR1 to mitochondria did not rescue fertility. Scale bars in **(B)** represent 0.5 mm.

**(D)** JA and JA-Ile accumulation at 1 h after wounding of seedlings of *opr2opr3* transformed with either *35S::OPR2* or *35S::OPR1*, either targeted to cytosol (Cyt), peroxisomes (Px), or mitochondria (Mt). Numbers refer to independent transgenic lines, which were selected for single insertion and homozygosity. The mean levels of JA/JA-Ile of *opr2opr3* are taken from Figure 7 and are represented by dashed lines for comparison. Bars represent means of four biological replicates with 120 seedlings each (single dots; ±SEM). Statistically significant differences between *opr2opr3* transformed with the different *OPR2* and *OPR1* constructs and *opr2opr3* were calculated using One-Way ANOVA with Fisher LSD post-hoc test and are indicated by asterisks with $*p<0.05$, $**p<0.01$, $***p<0.001$, and $****p<0.0001$; ns = non-significant.

**Figure 7: Schematic picture on the approaches leading to insights about the putative distribution of OPDA and 4,5-ddh-JA within cells of a wounded Arabidopsis seedling.**

Putative subcellular distribution of OPDA and its conversion to JA deduced from the trans-organellar complementation approach.

**(A)** In wild-type plants, OPDA is converted to JA in the peroxisome through the OPR3 canonical route, the contribution of OPR2 located in the cytosol appears to be neglectable but comes into place if OPR3 is not functional (e.g., *opr3* single mutant, Chini et al., 2018).

**(B)** The interruption of both routes in the *opr2opr3* mutant results in conversion of OPDA to 4,5-ddh-JA which is not metabolized to JA/JA-Ile.

**(C)** Targeting of OPR3 to any organelle in *opr2opr3* mutant plants (depicted for mitochondrial localization) restitutes JA biosynthesis in this mutant. If this is a result of either an ability of OPDA or of its β-oxidation product, 4,5-ddh-JA, to translocate to other cell compartments cannot be decided by this approach, since OPR3 is active to convert OPDA and 4,5-ddh-JA (red question marks). Trans-organellar complementation by using OPR2 or OPR1 targeted into mitochondria, however, helped to clarify this point.

**(D)** Mitochondrial targeting of OPR2 with its specificity to 4,5-ddh-JA complemented JA biosynthesis favoring rather an ability of 4,5-ddh-JA to be mobile, although the conversion of OPDA can still not be excluded due to some activity of OPR2 towards OPDA (light red question mark).

**(E)** As final proof, OPR1 was targeted to mitochondria, because OPR1 is specific for OPDA and does not convert 4,5-ddh-JA. Expression of mtOPR1 in *opr2opr3* did not restitute JA/JA-Ile production. Even the very limited JA formation by cytosolic OPR1 suggests restricted occurrence of OPDA in this compartment (not shown in this scheme, data see Figure 6). The lack of JA formation by mitochondrial OPR1 and the diminished JA formation by cytosolic OPR1 suggest that OPDA may not have the ability to access these cellular compartments and might strongly compartmentalized in the organelles involved in JA biosynthesis, the plastids and peroxisomes.