Unexpectedly low recombination rates and presence of hotspots in termite genomes

Turid Everitt[1#], Tilman Rönneburg[1#], Daniel Elsner[2], Anna Olsson[1], Yuanzhen Liu[1], Tuuli

Larva[1], Judith Korb[2,3], Matthew T Webster[1,4*]

1) Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden

2) Evolutionary Biology and Ecology, University of Freiburg, Freiburg, Germany

3) Research Institute for the Environment and Livelihoods, Charles Darwin University,

Casuarina Campus, Darwin, Australia

4) Science for Life Laboratory, Uppsala University, Uppsala, Sweden

# equal contribution

*correspondence to matthew.webster@imbim.uu.se

Abstract

Meiotic recombination is a fundamental evolutionary process that facilitates adaptation and

the removal of deleterious genetic variation. Social Hymenoptera exhibit some of the

highest recombination rates among metazoans, whereas high recombination rates have not

been found among non-social species from this insect order. It is unknown whether

elevated recombination rates are a ubiquitous feature of all social insects. In many

metazoan taxa, recombination is mainly restricted to hotspots a few kilobases in length.

However, little is known about the prevalence of recombination hotspots in insect genomes.

Here we infer recombination rate and its fine-scale variation across the genomes of two

social species from the insect order Blattodea: the termites *Macrotermes bellicosus* and

Cryptotermes secundus. We used linkage-disequilibrium-based methods to infer

recombination rate. We infer that recombination rates are less than 1 cM/Mb in both

species, which is lower than the average metazoan rate. We also observed a highly punctate

distribution of recombination in both termite genomes, indicative of the presence of

recombination hotspots. We infer the presence of full-length PRDM9 genes in the genomes

of both species, which suggests recombination hotspots in termites might be determined by

PRDM9, as they are in mammals. We also find that recombination rates in genes are

correlated with inferred levels of germline DNA methylation. The finding of low

recombination rates in termites indicates that eusociality is not universally connected to

elevated recombination rate. We speculate that intense selection among haploid males

promotes high recombination rates in social Hymenoptera.

Introduction

Meiotic recombination has two fundamental roles: It is essential for correct disjunction of

chromosomes during cell division and it generates new combinations of genetic variants

that form the raw material of evolution (Hartfield and Keightley 2012). However,

recombination can also be deleterious, as it can generate structural mutations and break up

favourable combinations of alleles. Recombination varies among species, with optimal

recombination rate in a genome likely determined by the optimal trade-off of its positive

and negative evolutionary and mechanistic effects (Stapley et al. 2017).

The highest recombination rates in metazoans observed so far are found in social insects

(Wilfert et al. 2007; Stapley et al. 2017). However, until now recombination rate has only

been studied in social insects from the order Hymenoptera, to which the majority of social

insects belong. Among Hymenoptera, the highest rates are found in the genus *Apis*: the

Western honey bee *Apis mellifera* (20.8 cM/Mb) and its relatives *Apis cerana* (17.4 cM/Mb)

Apis florea (20.8 cM/Mb) and *Apis dorsata* (25.1 cM/Mb) (Beye et al. 2006; Shi et al. 2013;

Liu et al. 2015; Wallberg et al. 2015; Rueppell et al. 2016; Kawakami et al. 2019). Other

social Hymenoptera also have high rates including the bumblebee *Bombus terrestris* (8.9)

(Liu et al. 2017; Kawakami et al. 2019), the stingless bee *Frieseomelitta varia* (9.3 – 12.5

cM/Mb) (Waiker et al. 2021), the wasp *Vespula vulgaris* (9.7 cM/Mb) (Sirviö et al. 2011a)

and the ants *Pogonomyrmex rugosus* (11.1 cM/Mb) (Sirviö et al. 2011b) and *Acromymex*

echinatior (6.1) (Sirviö et al. 2006). In contrast, the solitary bee *Megachile rotundata* and the

solitary wasp *Nasonia* have relatively low rates (1.0 and 1.5 cM/Mb respectively) (Niehuis et

al. 2010; Jones et al. 2019). For comparison, the average recombination rate of 15 insect

species outside of Hymenoptera is 2.2 cM/Mb (Wilfert et al. 2007).

We can learn about the forces shaping the evolution of recombination rates in general by

understanding why high recombination rates have evolved in social Hymenoptera. Several

hypotheses have been proposed to explain this observation. One set of explanations

focusses on the effects of recombination on intra-colony genetic diversity. For example,

increasing the genetic diversity of a colony could promote task specialization among

workers (Kent et al. 2012; Kent and Zayed 2013). Similarly, elevated genetic variation in a

colony could prevent invasion by pathogens or parasites due to diversification of immune

genes (Fischer and Schmid-Hempel 2005). Recombination could also reduce variance in

relatedness between nestmates, thereby reducing potential kin conflict within colonies

(Sherman 1979; Templeton 1979; Wilfert et al. 2007). Additionally, it has been proposed

that high recombination rates could have facilitated the evolution of eusociality over a

longer evolutionary timescale, because recombination between genes that take on

functions in different castes permits them to evolve more independently (Kent and Zayed

2013). Several studies have addressed these hypotheses (Kent et al. 2012; Liu et al. 2015;

Wallberg et al. 2015; Rueppell et al. 2016; Liu et al. 2017; Jones et al. 2019; Waiker et al.

2021; Kawakami et al. 2019) but so far none is strongly supported. It is also not clear

whether eusociality *per se* selects for higher recombination rates, or whether high rates are

a specific feature of eusocial Hymenoptera.

The main characteristics of eusocial insect species are the presence of a worker caste, which

forgo reproduction in order to care for brood of other colony members, and a single (or very

few) reproducing individuals (queens, and in termites, also kings). Eusocial insects are found

mainly in Hymenoptera, among bees, wasps and ants, and also Blattodea, in which all

termites (Infraorder: Isoptera) are eusocial but exhibit varying levels of social complexity.

These insect orders likely diverged ~400 million years ago (Misof et al. 2014) and there are

substantial differences between social insects from the two orders (Korb 2008; Korb and

Thorne 2017). Hymentoptera belong to the superorder Holometabola, which is the most

diverse insect superorder and contains 11 orders including Lepidoptera (butterflies, moths),

Coleoptera (beetles), and Diptera (true flies), which all undergo complete metamorphosis.

Blattodea is a hemimetabolous order, which undergoes partial metamorphosis.

Hymenoptera have haplodiploid sex determination in which males develop from haploid,

unfertilized eggs and females derive from diploid, fertilized eggs. In Blattodea, both males

and females are diploid. Members of the worker caste in eusocial Hymenoptera are all

female, whereas eusocial Blattodea workers are of both sexes. Hymenoptera colonies are

headed by one or a small number of queens, whereas Blattodea colonies contain both a king

and a queen.

The hypotheses proposed to explain the high recombination rates observed in eusocial

Hymenoptera predict that recombination should be elevated in all social insects. However,

so far it is unknown whether social insects from other orders also have high rates. As

described above, there are fundamental differences in the sex determination system and

colony compositions of social insects from Hymenoptera and Blattodea. The robustness of

the association between eusociality and high recombination rates can therefore be

addressed by assessing mean genomic recombination rates in social Blattodea (termites).

In addition to variation between species, the rate of recombination also varies along

chromosomes. In mammals, crossover events are mainly restricted to regions that contain

specific motifs that act as binding sites for the PRDM9 (PR/SET domain 9) protein that marks

regions for double-stranded breaks, which are repaired by recombination during meiosis

(Baudat et al. 2010; Myers et al. 2010; Parvanov et al. 2010). Species with an active PRDM9

include nearly all mammals and many vertebrate taxa (Baker et al. 2017). In species without

PRDM9, recombination can be directed to hotspots defined by other features, such as CpG

islands and promoters that have open chromatin. Such hotspots have been characterised in

the genomes of diverse taxa including yeast, birds, and dogs (Axelsson et al. 2012; Lam and

Keeney 2015; Singhal et al. 2015; Berglund et al. 2014). In insects, the prevalence of

recombination hotspots is not well studied. Fine-scale recombination maps in the fruit fly

Drosophila melanogaster and honey bee *A. mellifera* indicate a paucity of recombination

hotspots (Chan et al. 2012; Smukowski Heil et al. 2015; Wallberg et al. 2015), whereas

hotspots appear to be present in the genome of the butterfly *Leptidea sinapis* (Torres et al.

2023).

Several other factors may also modulate recombination rate along chromosomes, such as

germline DNA methylation. Many vertebrate species have elevated recombination in CpG

islands, which are usually unmethylated in the germline (Axelsson et al. 2012; Singhal et al.

2015; Berglund et al. 2014). In insect genomes, methylation is not always present, but

commonly restricted to gene bodies (Arsala et al. 2022; Wang et al. 2013; Lyko et al. 2010;

Ventós-Alfonso et al. 2020; Bewick et al. 2019; Harrison et al. 2018). In the honey bee *A.*

mellifera and solitary bee *M. rotundata* recombination is reduced in genes, and particularly

in those inferred to have the highest levels of germline methylation (Wallberg et al. 2015;

Jones et al. 2019). In addition, some studies have identified correlations between caste-

biases in gene expression and recombination. In honey bees, genes with worker-biased gene

expression have been found to have elevated recombination rates (Kent et al. 2012) but this

is likely an indirect result of differences in germline methylation between sets of genes

(Wallberg et al. 2015).

Here we estimate recombination rate and its fine-scale variation across the genomes of two

distantly related termite species with contrasting social complexities (Korb and Hartfelder

2008; Korb and Thorne 2017). *Macrotermes bellicosus* (Termitidae: Macrotermitinae) is a

fungus-growing termite with a high level of social complexity, with colonies of several

millions of individuals and sterile workers. It belongs to the foraging (multiple-pieces

nesting) termites, where workers leave the nest to forage for food. The drywood termite

Cryptotermes secundus (Kalotermitidae) has a low level of social complexity. Its colonies

consist of a few hundred individuals and totipotent workers from which queens and kings

develop (Hoffmann et al. 2012). It belongs to the wood-dwelling (one-piece nesting)

termites that nest in a single piece of wood that also serves as their food source.

The study has three main aims. Firstly, we aim to test the robustness of the association

between sociality and high recombination rate. If sociality is a universal driver of high

recombination rates, we would expect both termite species to have elevated recombination

rates, and that it is particularly elevated in *M. bellicosus*, which has a higher level of social

complexity. Secondly, we aim to determine whether recombination hotspots exist in the

two termite genomes, and infer whether active full-length PRDM9 genes are present in their

genomes. Thirdly, we aim to analyse the factors that govern variation in recombination

across the genome. In particular, whether genomic variation in recombination rates is

associated with patterns of methylation and gene expression as has been reported in other

studies.

Results

Genetic variation in two termite species is typical for social insects

We utilized genome assemblies of two termite species: *M. bellicosus* and *C. secundus*. The

genome assembly of *M. bellicosus* (Qiu et al. 2023) is 1.3 Gbp in total length across 275

scaffolds with scaffold N50 of 22.4 Mbp. The genome assembly of *C. secundus* (Csec_1.0)

(Harrison et al. 2018) is divided into 55,483 scaffolds with a scaffold N50 of 1.2 Mbp and a

total length of 1.0 Gbp.

We sequenced 10 unrelated individuals each of *M. bellicosus* and *C. secundus* to a mean

depth of 34x and 31x per sample, respectively (Supplementary Table S1). We mapped reads

to the appropriate genome assemblies and called variants in each species. In *M. bellicosus*

we called 5.6 million SNPs and in *C. secundus* 15.1 million SNPs (Table 1). One individual of

C. secundus (CS_8) was excluded due to a low proportion of mapped reads, poor mapping

quality and strand bias in mapping. Estimates of variation based on θ_W (Watterson 1975) are

0.14% and 0.44% in *M. bellicosus* and *C. secundus* respectively, which are broadly

comparable to levels of variation in the honey bee *A. mellifera* (0.3 - 0.8%) (Wallberg et al.

2014).

We used these estimates of genetic variation to estimate effective population size (N_E) for

each of the sampled populations (Table 1). These rely on an estimate of mutation rate, μ, in

each species. As there is no estimate of μ for Blattodea, we considered a range of estimates

from insects compiled by (Lynch et al. 2023) that includes estimates from Diptera

(*Drosophila*) (Keightley et al. 2009, 2014), Hymenoptera (Liu et al. 2017; Yang et al. 2015)

and Lepidoptera (Keightley et al. 2015). These estimates range between a minimum of 2.7 x

10^{-10} bp^{-1}gen^{-1} in *Acyrthosiphon pisum* (Fazalova and Nevado 2020) to a maximum of 8.1 x

10^{-9} bp^{-1}gen^{-1} in *Drosophila pseudoobscura* (Krasovec 2021). The average of multiple

estimates in *Drosophila melanogaster* is 4.5 x 10^{-9} bp^{-1}gen^{-1}, which is typical of insects (Lynch

et al. 2023).

Our estimates of N_E based on θ_W are 42,898 – 1,278,982 in *M. bellicosus*, which has high

social complexity, and 136,869 – 4,080,725 in *C. secundus*, which has a lower level of social

complexity (Table 1). Estimates of N_E assuming the typical mutation rate of 4.5 x 10^{-9} bp^{-}

^{1}gen^{-1} are 76,876 and 245,280 respectively. These estimates are consistent with previous

evidence that eusocial species with large colonies and high social complexity tend to have

lower effective population sizes and the values are broadly comparable to levels of N_E found

in social and solitary Hymenoptera (Leffler et al. 2012; Romiguier et al. 2014).

Low average recombination rate in termite genomes

We next examined the decay of linkage disequilibrium (LD) based on the statistic r^2. Both

termite species show more extensive LD compared to *A. mellifera*, which has an extremely

high recombination rate (Wallberg et al. 2015) (Figure 1). As N_E is broadly similar in all these

species, these differences are expected to largely represent differences in recombination

rates, and indicate that the two termite species do not exhibit the elevated genome average

recombination rate that is observed in honey bees.

We estimated variation in the population recombination rate, ρ, in both datasets using

LDhat (Auton and McVean 2007) (Table 1). Average rates of ρ/kb are 1.7 and 4.8 in *M.*

bellicosus and *C. secundus* respectively. These estimates can be converted to cM/Mb using

estimates of N_E, which gives estimates of average recombination rate of 0.033 - 0.986

cM/Mb in *M. bellicosus* and 0.030 - 0.885 cM/Mb in *C. secundus*, which is much lower than

the average among insects (Stapley et al. 2017), and particularly lower than the elevated

recombination rates inferred in eusocial Hymenoptera. Estimates that assume a mutation

rate close to the average for insects are 0.550 and 0.494 cM/Mb in *M. bellicosus* and *C.*

secundus respectively. This represents the first estimation of recombination rate in social

insects outside of Hymenoptera and demonstrates that eusociality is not a universal driver

of high recombination rates.

Evidence for active recombination hotspots in termite genomes

Variation in recombination rate across the genome in *M. bellicosus* and *C. secundus* appears

highly variable (Figure 2). We characterised the distribution of recombination events across

the two termite genomes using cumulative distribution plots (Figure 3). We found that 50%

of the recombination occurs in 3.5% of the genome in *M.* bellicosus and 4.8% of the genome

in *C. secundus*. This indicates that the majority of recombination events are restricted to a

much smaller portion of the genome than observed in *A. mellifera*, where 50% of

recombination events occur in >30% of the genome (Wallberg et al. 2015).

We further investigated whether recombination hotspots are present in the two termite

genomes by looking at the distribution of ρ in windows of 1 kbp, 10 kbp and 100 kbp across

the genome (Figure S1). For all window sizes, a small subset of windows was observed with

ρ greatly exceeding the mean in both termite species. For example, the percentage of 1 kbp

windows with recombination rate at least four standard deviations above the genome wide

mean value is 0.88% for *M. bellicosus* and 1.20% for *C. secundus*. By contrast, *A. mellifera*

has no windows above this limit for any window size, consistent with a lack of

recombination hotspots.

We next defined hotspots as 2 kbp regions with more than 5-fold higher recombination rate

than the local background as defined as the surrounding 100 kbp and used STREME (Bailey

2021) to identify 8 to 20-mers that were enriched in hotspots compared to the rest of the

genome. This yielded significantly enriched motifs in both species (8 motifs for *M. bellicosus*,

and 12 motifs for *C. secundus* after multiple testing; Supplementary Tables S2 and S3).

However, none the motifs we identified are present in more than 5% of hotspots and exhibit

more than 2-fold difference relative to the background, suggesting they are not viable

candidate motifs for promoting recombination in hotspots.

Identification and characterization of the PRDM9 gene in termites

We searched the *M. bellicosus* and *C. secundus* genomes for evidence of complete PRDM9

genes, which could potentially govern the presence of recombination hotspots in these

genomes, as it has been shown to do in mammalian genomes (Myers et al. 2010; Baudat et

al. 2010; Parvanov et al. 2010). The *C. secundus* genome contains a PRDM9 homolog

(XP_023708049.2) with annotated KRAB, SSXRD, SET and zinc finger domains (Figure 4).

There is no complete homolog of PRDM9 in the *M. bellicosus* annotation. We therefore used

BLAST to search for sequences homologous to the conserved functional domains from *C.*

secundus. We identified a putative PRDM9 orthologue on Scaffold 42, containing all

domains in the correct order. We identified retroviral conserved domains in an intron 5' of

the zinc-finger domains in this gene in the *M. bellicosus* genome assembly, which are not

predicted to alter the transcript. We confirmed the presence of a full-length transcript as

well as presence and correct order of conserved domains by alignment of the

Zoothermopsis nevadensis PRDM9 protein sequence to the *M. bellicosus* genome using

GeneWise (Madeira et al. 2024)(Figure 4 and Supplementary Table S4).

We used a zinc-finger prediction algorithm (Persikov and Singh 2014) to predict DNA binding

motifs from the zinc-finger domain present in PRDM9 in the *M. bellicosus* and *C. secundus*

genome assemblies, for which an accurate annotation was available. The most probable

consensus motifs were TATGGAACGACAGGAACAACGACATCATCAGCCGCCTAAT and

TAATAAGTAGAATCGTTTAG for *M. bellicosus* and *C. secundus* respectively. Base probability

matrices for these motifs are shown in Supplementary Table S5. We tested whether the

presence of these motifs corresponded to elevated recombination considering 1 kbp around

the motif. However, recombination rate was not significantly elevated around these

predicted motifs for either species, either when searching based on the most likely motif, or

by searching based on the probability matrices.

Genomic correlates of recombination rate

We analysed genomic correlates of recombination rate in 10 kbp windows. There is a

significant correlation between recombination and $CpG_{O/E}$ in both species with Spearman's ρ

= 0.24, $p < 1 \times 10^{-4}$ for *M. bellicosus* and Spearman's ρ = 0.10, $p < 1 \times 10^{-4}$ for *C. secundus*

(Figure S2). A reduction of recombination rates in regions of low CpG content is consistent

with interference of germline methylation with recombination. However, as the correlation

coefficients are low, the explanatory power is weak. Correlations of similar magnitudes were

observed between recombination and GC content (Spearman's ρ =0.017 for *M. bellicosus*, p

$< 1 \times 10^{-4}$ and Spearman's ρ =0.23 for *C. secundus*, $p < 1 \times 10^{-4}$). This correlation has been

associated with an effect of recombination in driving GC through GC biased gene conversion

but the relatively low correlation coefficients indicate that this force is not strong in

termites. By contrast, the correlation between GC content and recombination rate is much

stronger in honey bee (R^2 = 0.506), which is likely caused by high recombination rates and

intense GC-biased gene conversion (Wallberg et al. 2015).

There is no significant correlation between overall repeat element density and

recombination rate in either of the two genomes. There is a weak positive but highly

significant correlation between simple repeats and recombination in both genomes

(Spearman's ρ = 0.06, p < 1 x 10^{-4} for *M. bellicosus* and Spearman's ρ = 0.07, p < 1 x 10^{-4} for

C. secundus). SINE and LINE elements are also only weakly associated with recombination

rate in both genomes (SINEs: Spearman's ρ = -0.01, p > 0.05 for *M. bellicosus* and

Spearman's ρ = 0.02, p < 1 x 10^{-4} for *C. secundus*; LINEs: Spearman's ρ = -0.01, p < 1 x 10^{-4} for

M. bellicosus and Spearman's ρ = 0.001, p > 0.05 for *C. secundus*). There is a significant

negative correlation between gene density and recombination for the *M. bellicosus* genome

(Spearman's ρ = -0.20, p < 1 x 10^{-4}), but no significant correlation for *C. secundus*. We also

tested whether hotspots differ from the background with respect to GC content. The GC

content does not differ substantially between hotspots and the rest of the genome in either

species (means of 40.1% and 40.4% for *M. bellicosus* and 40.5% in both categories for *C.*

secundus).

We found a highly significant correlation between nucleotide diversity (π) and

recombination in both species (Figure S2), with Spearman's ρ = 0.52, p < 1 x 10^{-4} for *M.*

bellicosus and Spearman's ρ = 0.32, p < 1 x 10^{-4} for *C. secundus*. This correlation is found

across a wide range of eukaryotic taxa and is generally accepted to be caused by the

interaction between recombination and linked selection (Nachman 2001; Begun and

Aquadro 1992).

Recombination and gene expression patterns correlate with CpG$_{O/E}$

Analysis of the distribution of CpG$_{O/E}$ in both termite species showed that it is bimodally

distributed, with substantially lower values in introns and exons compared to flanking

noncoding regions (Figure S3). This indicates that germline DNA methylation is biased

towards gene bodies in the two termite genomes, a pattern that is also found in honey bees

and other insect genomes with functional DNA methylation (Harrison et al. 2018; Elango et

al. 2009; Sarda et al. 2012). We found that exons have lower $CpG_{O/E}$ compared to introns,

which are both lower than flanking regions (Figure 5, top row; all comparisons p < 0.001

after Bonferroni correction) consistent with higher levels of germline methylation in exons

(Supplementary Table S6). We next analysed how recombination rate varies among genomic

features. We found that ρ/kbp is also significantly reduced in genes compared to flanking

regions, with exons showing reduced ρ/kbp compared to introns (Figure 5, bottom row, all

p-values <0.005 after Bonferroni correction). This effect was seen in both species and is

more pronounced for *C. secundus* than for *M. bellicosus.* There was a significant correlation

between $CpG_{O/E}$ and recombination rate among genes (Spearman's ρ = 0.369, p < 1 x 10^{-4} for

M. bellicosus; Spearman's ρ = 0.166, p < 1 x 10^{-4} for *C. secundus*).

We next investigated whether patterns of gene expression between castes and sexes

(hereafter, for short 'caste-biased gene expression') were correlated with $CpG_{O/E}$ and

recombination rate using gene expression data from two studies (Lin et al. 2021; Elsner et

al. 2018). It has been proposed that it is evolutionarily advantageous for genes with worker-

biased expression to be located in regions with elevated recombination (Kent et al. 2012).

For *M. bellicosus*, we defined several categories based on gene expression difference among

castes (Elsner et al. 2018). The categories we used were queen-biased, king-biased, worked-

biased, male-biased, female-biased, reproduction-biased, and differentially expressed (see

methods for details). For *C. secundus*, worker-biased and queen-biased genes were

identified (Lin et al. 2021). Comparisons of $CpG_{O/E}$ and recombination rate for the original

gene expression categories reported for *M. bellicosus* are shown in Figure S4. We find that

mean $CpG_{O/E}$ in gene bodies shows significant variation between gene expression categories

in both *M. bellicosus* and *C. secundus* (Figure 6). However, this variation is not consistent

between species. In *M. bellicosus*, queen-biased and female-biased genes have low $CpG_{O/E}$

and king-biased and male-biased genes have high $CpG_{O/E}$. In *C. secundus*, queen-biased

genes have higher than average $CpG_{O/E}$.

The differences in $CpG_{O/E}$ between caste-biased gene expression categories are mirrored by

differences in ρ/kbp (Figure 6). Gene expression categories with elevated $CpG_{O/E}$

consistently show elevated ρ/kbp, and this pattern is found in both species. Studies in the

honey bee have found that worker-biased genes have lower CpG and higher recombination

rates (Kent et al. 2012; Wallberg et al. 2015). In *M. bellicosus,* we find that worker-biased

genes have slightly elevated CpG$_{O/E}$ and ρ/kbp. However, in *C. secundus*, these values are

reduced in worker-biased genes, whereas queen-biased genes have a slightly elevated

CpG$_{O/E}$ and ρ/kbp in this species.

We investigated the relative roles of CpG$_{O/E}$ and gene expression in determining ρ in genes

using a linear model, with CpG$_{O/E,}$ ρ in flanking region and expression categories as

explanatory variables. In both *M. bellicosus* and *C. secundus*, we found that CpG$_{O/E}$ was the

strongest predictor of recombination rate variation among genes (Supplementary Tables S7

and S8) in line with the analyses above. We found that caste-biased expression had no

significant impact on the recombination rate, with the exception of king-biased genes in *M.*

bellicosus. These have higher ρ/kbp than expected based on CpG$_{O/E}$ alone. Generally, CpG$_{O/E}$

is a good predictor of ρ/kbp*,* and the variation in ρ/kbp among expression categories likely

mainly reflects differences in CpG$_{O/E}$, as accounting for it removes most significant

interactions.

Discussion

The main findings we present here are 1) that two distantly related termite species with

varying social complexity both have relatively low genomic average rates of meiotic

recombination, 2) both genomes possess recombination hotspots and likely contain full-

length copies of PRDM9 and, 3) there are reduced levels of recombination in genes and

genome regions depleted in CpG sites. Our findings contrast with the extremely elevated

recombination rates observed in eusocial Hymenoptera, which are likely to be generated by

a feature that is specific to these taxa. The finding of recombination hotspots is one of the

first in insects. This could reflect a conserved function of PRDM9 in initiating recombination

events in both vertebrates and invertebrates. Our results also support a role for germline

methylation in suppressing recombination.

Low recombination rates in termites

We estimate that recombination rates in both termite species are less than 1 cM/Mb. These

estimates of recombination rate per meiosis depend on estimates of mutation rate. This is

not known in termites so we considered a range of estimates from insects. Data from a wide

range of taxa indicate that at least one crossover per chromosome tetrad is essential for

correct chromosomal segregation during meiosis. The chromosome numbers *M. bellicosus*

and *C. secundus* are 2n = 42 and 2n = 40 respectively (Jankásek et al. 2021). Considering the

assembly lengths, this corresponds to a minumum average genomic recombination rate of

0.92 cM/Mb and 1.01 cM/Mb would be necessary for accurate meiosis to occur in the two

species respectively. These values are closest to the maximum rates of 0.99 and 0.89

cM/Mb estimated when using the mutation rate estimated from *D. pseudoobscura* which is

the highest estimate reported in insects (Lynch et al. 2023). The estimates based on higher

mutation rates are therefore more likely to be true, indicating that termite mutation rates

are likely to be among higher range for insects and that Ne in this species is relatively low

(~40,000 for *M. bellicosus* and 140,000 for *C. secundus* based on Table 1).

It should be noted that our estimates of recombination rate are sex-averaged rates. It is

possible that recombination is absent in one of the termite sexes, which would make a

lower sex-averaged recombination rate more feasible. In addition, sex-linked contigs have

not been identified in the termite genome assemblies utilized here. *M. bellicosus* and *C.*

secundus have X1X2/Y1Y2 and X/Y sex-determining systems respectively (Jankásek et al.

2021). The inclusion of sex chromosomes in the analysis leads to errors in determining the

genome-wide recombination rate per meiosis because they have lower N_E than the rest of

the genome. In addition, sex chromosomes are more prone to assembly and read mapping

errors, due to reads aligning with the incorrect copy of the chromosome. However,

considering that sex chromosomes only comprise a minor portion of the genome, this is

unlikely to substantially alter our results.

Eusociality is not a universal driver of high rates of meiotic recombination

Elevated genomic rates of recombination are characteristic of all eusocial Hymenoptera so

far investigated. Several hypotheses have been advanced to explain this phenomenon,

which are all based on an evolutionary advantage of recombination in eusocial species

(Wilfert et al. 2007). For instance, it has been suggested that elevated recombination could

increase genetic diversity in a colony. This could promote diversity in the workforce,

enabling more efficient task specialisation among workers (Kent et al. 2012). Alternatively, it

could render the colony less susceptible to invasion by parasites and pathogens (Fischer and

Schmid-Hempel 2005). Elevated recombination might also favour rapid evolution of caste-

specifically expressed genes (Kent and Zayed 2013). In addition, increased recombination

could lead to reduced variance in relatedness, which could prevent kin conflict in colonies

(Sherman 1979; Templeton 1979).

All these hypotheses predict that recombination rate should be elevated in all eusocial

insects. Here, however, in the first report of recombination rate in social insects outside of

Hymenoptera, we find overall low levels of recombination. Both termite species studied

here are eusocial, and in particular *M. bellicosus* has extremely large and complex colonies

characteristic of eusociality (Korb and Thorne 2017). The low recombination rate inferred in

the two termite genomes prompts re-evaluation of hypotheses connecting eusociality with

high recombination rates. Although both termites and social Hymenoptera share many

traits, they differ in others that reflect their different ancestries.

The chromosome numbers of both termite species (*M. bellicosus*, 2n = 42; *C. secundus*, 2n =

40) are typical for termites, although variation exists between taxa (Jankásek et al. 2021).

These values do not differ from those observed in Hymenoptera, which have a similar range

(e.g. *A. mellifera*, 2n = 32) and do not vary between solitary and eusocial taxa (Cardoso et al.

2018; Cunha et al. 2021; Ross et al. 2015). This suggests that differences in chromosome

number are unlikely to be relevant in explaining the differences in recombination rates

among these taxa.

One proposed advantage of sex and recombination is that it is associated with sexual

selection and a higher intensity of selection on males. This could be advantageous because

differential male mating success can reduce mutational load in sexual populations by

causing deleterious mutations to segregate at a lower equilibrium frequency on average

(Siller 2001; Agrawal 2001). Recombination is essential for effective purging of deleterious

alleles as it unlinks deleterious variants from nearby beneficial ones thus preventing

selection interference (Hartfield and Keightley 2012). Selection has been found to be

stronger in males than females across animal species (Janicke et al. 2016; Winkler et al.

2021). Species with a male-biased sex ratio are expected to experience particularly intense

selection on males. It is therefore possible that recombination rate could be particularly

elevated in species in which males face intense competition as this increases the

evolutionary advantage of recombination.

The genus *Apis* produces highly male-biased reproductive offspring and thousands of male drones compete to fertilise a small number of female queens (Winston 1991; Koeniger et al. 2011; Baer 2005). It is also in this genus that the most extreme recombination rates have been inferred (Beye et al. 2006; Shi et al. 2013; Liu et al. 2015; Wallberg et al. 2015; Rueppell et al. 2016; Kawakami et al. 2019). Male-biased sex ratios are also observed in other social bees and wasps. Both male and female biased operational sex ratios have been observed in ants, but are difficult to measure because workers invest more resources in female offspring and may manipulate sex ratios. In non-social Hymenoptera, females lay roughly equal numbers of haploid and diploid eggs (Trivers and Hare 1976). The sex ratio of reproductive males and females is also relatively even in most termite species (Roisin 2001). It is possible that the male-biased sex ratio observed in many eusocial Hymenoptera could favour elevated recombination rates in the species so far studied.

The effects of higher intensity of selection on males on recombination rate could be even more pronounced when males are haploid as in Hymenoptera. Haploid males can be considered analogous to the gametes of females. However, as many more genes are expressed in adult haploid males compared to gametes there are more targets of selection. Selection against deleterious alleles is particularly effective in haploid males because recessive deleterious alleles are exposed (Hedrick and Parker 1997).

Although its effects on overall recombination rates are unclear, haploid selection is believed be key to driving sex-specific differences in recombination rate (Immler 2019). It has been argued that differences in the intensity of selection between sexes in the haploid phase can drive differences in recombination rates between sexes among animal and plant species (heterochiasmy) (Burt et al. 1991; Trivers 1988; Lenormand 2003; Lenormand and Dutheil 2005). It is unclear how theory and observations regarding the prevalence and nature of heterochiasmy applies to haplodiploids, in which only females recombine. However, it appears that the intensity of selection on the haploid phase likely has the potential to drive selective pressure on recombination rates.

Further research is needed to determine the cause of variation in recombination rates among social and non-social insects. For example, simulations could be used to determine the effect of selection on haploid males on recombination rates, considering different sex ratios. In addition, it is necessary to estimate recombination rate in a wider range of social

and non-social species to determine whether the strength of selection on males is a key

factor in determining these rates.

Full length PRDM9 gene and recombination hotspots in termite genomes

Our results are also indicative of the presence of recombination hotspots in the two termite

genomes. The fine-scale distribution of recombination rate in both termite genomes is

similar to that in humans where 50% of recombination events occur in ~6% of the genome

(Myers et al. 2005). This contrasts to the honey bee and solitary bees where fine-scale maps

are available and recombination is relatively uniformly distributed in the genome (Wallberg

et al. 2015; Jones et al. 2019).

Two main mechanisms have been shown to lead to recombination hotspots. The first, which

is best understood in human and mouse, is governed by the PRDM9 protein. A zinc-finger

domain recognises a specific DNA sequence motif and then performs a histone modification

in the vicinity, which marks the sequence for a DNA double-stranded break that is repaired

by recombination during meiosis (Baudat et al. 2010; Myers et al. 2010; Parvanov et al.

2010). In species with this mechanism, the zinc finger domain is fast evolving, which results

in a rapid turnover of hotspot locations (Myers et al. 2010). Full-length PRDM9 orthologues

are present among a range of highly-diverged vertebrates including many fish, reptiles and

mammals, but many full or partial losses of the gene have also been inferred. In a study of

225 vertebrate genomes, a minimum of six partial and three complete losses were inferred

(Baker et al. 2017). However, it is unclear whether PRMD9-hotspots are common in

invertebrates.

A second mechanism is found in vertebrates that lack PRDM9, in which recombination

hotspots are localised to regions of open chromatin such as CpG islands and promoters. This

is found in dogs and in most or all birds, in which PRDM9 has been lost, and results in

hotspots with stable rather than rapidly evolving locations (Axelsson et al. 2012; Singhal et

al. 2015). A similar pattern is also observed in the yeast *Saccharomyces cerevisiae* in which

recombination preferentially occurs in promoter regions (Wu and Lichten 1994). PRDM9-

independent hotspots can be identified in CpG islands is some vertebrate genomes even

when PRDM9 is present (Joseph et al. 2024). In invertebrates such as *Drosophila* and

Caenorhabditis elegans, recombination hotspots appear to be absent (Coop and Przeworski

2007; Chan et al. 2012; Smukowski Heil et al. 2015). In insects, fine-scale maps of social and

solitary bees have not revealed recombination hotspots (Wallberg et al. 2015; Jones et al.

2019), but there is evidence for hotspots in a butterfly genome (Torres et al. 2023).

We find evidence that recombination is directed towards hotspots in both termite species.

We also identify a full-length PRDM9 gene in both species, containing all of the domains

needed to initiate recombination. However, we are unable to demonstrate that PRMD9 is

responsible for the presence of recombination hotspots in the two genomes. We used a

motif prediction algorithm to predict the zinc-finger binding motif in the *C. secundus*

genome, but we did not observe elevated recombination rate around instances of the motif

in the genome. Furthermore, we did not identify any common sequence motifs that were

enriched in recombination hotspots in either of the termite genomes. A plausible

interpretation of these observations is that the PRDM9 protein indeed initiates

recombination in hotspots in both species, but that positive selection leads to frequent

shifts in the target motifs, which results in a lack of a strong association between a specific

motif and LD-based hotspots (Myers et al. 2010). It is also possible that the sequences of the

zinc-finger motifs are not correctly represented in the genome assemblies, which could

result from problems with assembly around minisatellites. However, an unknown

mechanism could also be responsible for hotspots in termites.

Evidence that germline DNA methylation suppresses recombination

In insects with a functioning DNA methyltransferase machinery, the $CpG_{O/E}$ statistic shows

substantial variation along the genome, which mainly reflects variation in levels of germline

DNA methylation (Elango et al. 2009). In contrast to vertebrate genomes, methylation is

strongly skewed towards gene bodies in the genomes of insects (Glastad et al. 2011). The

overall genomic correlation between $CpG_{O/E}$ and recombination rate is significant but weak

in the two termite genomes. This could reflect that the majority of the genomes are not

methylated, so that much of the variation in $CpG_{O/E}$ on the genome scale is not strongly

influenced by methylation. In the two termite genomes studied here, we find lowest values

of $CpG_{O/E}$ in exons, whereas noncoding flanking regions are biased towards higher values,

indicative of lower levels of DNA methylation (Figure 5, Figure S3). All categories display a

predominantly bimodal distribution of $CpG_{O/E}$, likely reflective of the presence of

methylated and unmethylated regions.

The differences in $CpG_{O/E}$ are also reflected by differences in recombination rate (ρ/kb),

which are reduced in regions inferred to be more highly methylated. Our inference of

reduced recombination rates in exons could be influenced by lower N_E in these regions due

to the effects of linked selection, which could result in more extensive LD (Charlesworth

2009). However, the observation that differences in $CpG_{O/E}$ among genes are associated

with recombination rate indicates that linked selection is unlikely to be a major factor

driving the observed differences in recombination rate.

Using linear models, we identified $CpG_{O/E}$ as the factor with the strongest influence on

recombination rate variation among genes, consistent with an effect of DNA methylation on

recombination rate. This factor had a stronger effect than differences in gene expression

patterns across castes, a result that was also observed in the honey bee genome (Wallberg

et al. 2015). This indicates that recombination rate variation is not modulated by differences

in gene expression in these genomes, which might be expected if natural selection favours

increased recombination rates in certain genes. For example, it has been proposed that

natural selection could favour elevated recombination rates in genes with roles in worker

behaviour or in immune function due to their important roles in colony function (Wilfert et

al. 2007; Kent et al. 2012). However, the results presented here suggest that differences in

recombination rate between genes with caste-biased patterns of gene expression are mainly

due to underlying differences in methylation levels among genes.

A direct effect of germline DNA methylation in suppressing recombination could influence

variation in recombination landscapes among vertebrates and invertebrates. Vertebrate

genomes are usually highly methylated with the exception of CpG islands. In vertebrate

genomes that lack a functional PRDM9 gene, recombination events tend to localise to CpG

islands (for example in dogs and birds)(Singhal et al. 2015; Axelsson et al. 2012). The effect

of DNA methylation on recombination rate variation in invertebrates is less clear, but the

results presented here also support a role of germline methylation in suppressing

recombination.

Materials and Methods

Sample collection

Macrotermes bellicosus samples were collected from colonies in Kakpin, next to the Comoé

National Park in Côte d'Ivoire (coordinates 8°39'N 3°46'W) (Elsner et al. 2018). For both

species, each of the sequenced samples came from a different colony. *Cryptotermes*

secundus colonies were collected from dead *Ceriops tagal* mangrove trees near Palmerston-

Channel Island (Darwin Harbor, Northern Territory, Australia; 12°30'S 131°00'E). Colonies

were then kept in *Pinus radiata* wood blocks in climate rooms in Germany, providing 28 °C,

70% relative humidity, and a 12-h day/night cycle (Lin et al. 2021).

Population sequencing

The termites were cut in two, approximately between head and thorax. DNA was extracted

from both parts using the QIAGEN Blood and Tissue Kit following the standard protocol,

including treating with 4 µL RNase and 25 µL Proteinase K. The DNA extraction product was

cleaned and concentrated with ZymoResearch Genomic DNA Clean & Concentrator kit. For

most samples, the RNase treatment was repeated prior to cleaning and concentrating. One

DNA sample from each individual was chosen, based on the DNA amount and quality, to be

sequenced. Sequencing libraries were produced using the Illumina DNA PCR-free library

preparation kit according to the manufacturers protocol, using an input DNA quantity of 200

ng per sample. This protocol yields insert sizes of approximately 350 bp. We performed

Illumina short-read sequencing on the libraries using NovaSeq 6000 S4 flow cell with 2 x 150

bp paired end reads.

Read mapping and variant calling

The sequence data from the two termite species were mapped to the reference genomes

for *M. bellicosus* (Qiu et al. 2023) and *C. secundus* (Csec_1.0) (Harrison et al. 2018)

respectively, using Burrows-Wheeler alignment tool BWA version 0.7.17 (Li & Durbin 2009)

with the BWA-MEM algorithm. Sorting and indexing of the BAM-files was done with the

Samtools package version 1.17 (Li et al. 2009), followed by adding read groups and marking

duplicate reads with Picard toolkit version 1.118 (http://broadinstitute.github.io/picard/).

After mapping, we excluded all scaffolds < 1kb from the *C. secundus* genome, which in total

correspond to 2% of the genome size, as they are potentially of lower quality and not

informative for analysis of linkage disequilibrium or recombination. For *M. bellicosus* there

are no scaffolds shorter than 1 kbp, but we excluded the following five scaffolds, which

together correspond to 3.78% of the genome length, due to unexpectedly high

heterozygosity: 93, 62, 41, 35 and 32.

One of the *C. secundus* individuals did not map well to the reference genome (71% mapped

reads, unequal distribution between forward and reverse strand and an average mapping

quality of 35 while all the other samples had an average mapping quality >45). This

individual was excluded from all further analyses and hence the total number of termite

individuals analysed in this study is 19, out of which 9 were *C. secundus*.

Variant calling was done using the tools HaplotypeCaller, GenomicsDBImport and

GenotypeGVCFs from GATK version 4.3.0.0 as described in their best practices workflow

(https://gatk.broadinstitute.org/hc/en-us/articles/360035535932-Germline-short-variant-

discovery-SNPs-Indels-, last accessed 2023-07-11). The variants were filtered in multiple

steps, first removing extended regions of low mapping quality or deviating read depth as

those regions are likely to be unreliable. This was done based on statistics from the

Samtools mpileup and depth tools, respectively, removing 100 kb windows with mean

mapping quality below 70 or mean read depth more than two standard deviations from the

genome-wide mean. In the later recombination rate and LD decay calculations, values in

intervals spanning those regions were also removed. Indels were removed using VCFtools

version 0.1.16 (Danecek et al. 2011) before GATK VariantFiltration was run with the

following limits, which are all equal to or stricter than the ones recommended

(https://gatk.broadinstitute.org/hc/en-us/articles/360035890471-Hard-filtering-germline-

short-variants, last accessed 2023-07-11) and adapted to the observed distributions of the

corresponding statistics: QD<2; FS>50; MQ<40; ReadPosRankSum< -4; ReadPosRankSum>4;

SOR>3; ExcessHet>5 for both species as well as MQRankSum with limits -5 and 5 for *C.*

secundus and limits -6 and 4 for *M. bellicosus*. This was followed by additional filtering with

VCFtools to select only bi-allelic sites with a minimum quality score of 30 and maximum of

40% missing genotypes. The SNPs with mean read depth in the lowest or highest 2.5% of the

depth distribution were also filtered out, which translates to a depth below 10 or above 34

for *C. secundus* and below 5 or above 40 for *M. bellicosus.* For the recombination rate

calculations, the SNPs were also filtered for a minor allele count >= 2, as rare variants

appearing on only one chromosome have a relatively high false positive rate and are not

informative regarding LD. The average proportion of missing genotypes per individual was

4.5 x 10^{-5} for *C. secundus* and 1.2 x 10^{-4} for *M. bellicosus*. After variant calling and filtering,

the data was phased and imputed using Beagle version 5.1 (Browning and Browning 2007;

Browning et al. 2018).

Estimation of LD decay

The decay of linkage disequilibrium, r2, with increasing physical distance was estimated with

the software PopLDDecay version 3.42 (Zhang et al. 2019) over a maximum distance of 10

kbp. For this analysis, sequence data from *A. mellifera scutellata* was included for

comparison ((Wallberg et al. 2017); BioProject PRJNA357367). The *A. m. scutellata*

sequences were mapped to the Amel_HAv3.1 reference genome (Wallberg et al. 2019) and

processed and filtered in the same way as described above for the termite data (GATK

VariantFiltration with the limits QD<2, FS>25, MQ<40, MQRankSum<-4, MQRankSum>4,

ReadPosRankSum<-4, ReadPosRankSum>4, SOR>3, ExcessHet>5 and VCFtools to select bi-

allelic sites with mean read depth between 5-13, quality score >= 30, minor allele count >=2

and <=40% missing genotypes).

Estimation of population recombination rate

The recombination rate was estimated using the complete, interval and stat functions of

LDhat (https://github.com/auton1/Ldhat, downloaded 2023-04-19) (Auton and McVean

2007) with 10 million iterations, sampling every 5000th iteration, taking the first 20 samples

as burn-in.

We evaluated different block penalties (1, 5 and 20), where a lower value allows the

recombination rate to change more easily across the map. As there were only minor

differences between the results for both species, we decided to use a block penalty of 1 as

this has been used previously in similar studies (Wallberg et al. 2015; Jones et al. 2019).

The output from the LDhat stat program gives the value of the parameter ρ/kbp between

each pair of consecutive SNPs. Those values were converted to equally sized windows across

each scaffold, as well as the coordinates of individual genomic features, by taking a

weighted average of values from overlapping intervals. The *M. bellicosus* scaffolds contain a

few gaps of unknown size and the values between SNPs across such gaps were removed.

The parameter ρ estimated by LDhat is proportional to the recombination rate r and the

effective population size N_E, which in turn can be estimated based on the mutation rate μ

and the number of segregating sites, S. In order to convert ρ to r, the following equations

were used, where n is the number of haploid sequences. The number of segregating sites

was calculated before the filter on minor allele count >=2 was applied, as this filter likely

removes many true SNPs as well, which could affect the estimates.

θ_W was estimated for each sample using the equation below from (Watterson 1975):

$$\theta_W = K \bigg/ \sum_{i=1}^{n-1} \frac{1}{i}$$

where K is the number of SNPs and i is the number of haploid genomes sampled. We used

the value of θ_W to estimate N_E using:

$$N_E = \frac{\theta}{4\mu}$$

where μ is the mutation rate. We estimated N_E using a range of mutation rates estimated

from other insect species (Keightley et al. 2009, 2014, 2015; Yang et al. 2015; Liu et al. 2017;

Lynch et al. 2023). We then estimated the recombination rate using the following equation,

assuming a range of values of N_E.

$$r = \frac{\rho}{4N_e}$$

Cumulative plot of rho ρ/kbp

A cumulative plot of the proportion of rho versus the proportion of the physical distance

along the genome was constructed based on the output from LDhat for the two termite

species and previously generated LDhat output for *A. mellifera* (Wallberg et al. 2015;

PRJNA236426). The ρ/kbp estimates between markers were placed in decreasing order

before multiplication with the respective physical distances between the markers. Then

cumulative sums were calculated for the ρ-values as well as the corresponding physical

distances and divided by the total genome-wide sums.

k-mer enrichment in recombination hotspots

In order to identify k-mers associated with elevated recombination rate, we defined

recombination hotspots as 2kb segments with at least five-fold elevated recombination rate

compared to the local background, as defined by 50 kb flanking up and downstream of the

segment, resulting in 39837 regions for *M. bellicosus and* 40894 regions for *C. secundus.*

Then, we used these hotspots to search for motifs associated with elevated recombination

rate by comparing against an equivalent number of randomly chosen 2kb regions from the

rest of the genome. This was done using STREME 5.5.4 (Bailey 2021) using a minimum and

maximum size of k-mer of 8 and 20, (--minw 8 ; --maxw 20) and otherwise default options.

From all motifs with significant enrichment after adjusting for multiple testing, we then

considered only motifs present in more than 5% of all hotspots and an at least two-fold

enrichment compared to the background as credible candidates.

Identifying PRDM9 orthologues in M. bellicosus and C. secundus

A full-length copy of PRDM9 is present in the *C. secundus* genome annotation, but not in *M.*

bellicosus. Using Orthofinder (Emms and Kelly 2019), and the genome annotation files for

both species, we identified a single orthologue to the *C. secundus* PRDM9 in *M. bellicosus*.

We analysed this region using the NCBI conserved domain search (Wang et al. 2023). We

also searched the *M. bellicosus* assembly for the PR/SET domains derived from *C. secundus*

and *Zootermopsis nevadensis* using BLAST (Camacho et al. 2009) and subsequently for all

other PRDM9 domains (KRAB, SSXRD, ZF-Casette) to identify hits that contain all other

PRDM9 domains in the vicinity in the correct order. We used GeneWise (Madeira et al.

2024) with default options to perform gapped alignment of a the PRDM9 protein sequence

from *Zoothermopsis nevadensis* against the target region to predict exon-intron borders.

Predicting binding sites from zinc-finger-motifs

Using the ZF-motifs from *C.secundus,* we obtained letter probability matrices for predicted

binding motifs using a webserver (http://zf.princeton.edu/index.php ; (Persikov and Singh

2014) which were used to search for binding sites across the genome using FIMO (Grant et

al. 2011) with default options. Subsequently, the mean rho for the surrounding 1 kb bin was

used to test for a difference in mean rho between bins associated with a suspected binding

site, and a similar number of randomly chosen bins without association. For significance

testing, we repeated this draw 10.000 times to compare the associated bins against the 95%

confidence interval of the resulting distribution.

Analysis of genomic correlates of recombination

We divided the genome into 10 kb windows in which average ρ and other genomic features

were estimated. Windows shorter than 5 kb (half of the specified window size), which

appear in short scaffolds or at the ends at longer scaffolds, were excluded from the analyses

(excluding less than 1% of the sequence for each species). Correlation coefficients were

estimated with Spearmans ρ, with significance estimated by permutation tests with 10,000

iterations.

In order to determine the repeat content across the *M. bellicosus* and *C. secundus* genome

assemblies, we first generated custom repeat libraries for each species with RepeatModeler

(Flynn et al. 2020) using default options. This was then combined with Repbase 29 (Bao et

al. 2015) and DFAM 3.8 (Storer et al. 2021) and used to run RepeatMasker with default

options on each genome. We estimated the proportion of sequence in each repeat class in

10 kbp windows and used Spearman's rank correlation to correlate this with estimates of

recombination rate.

Per gene recombination rate and CpG$_{O/E}$ ratio

The observed/expected frequency of CpG sites (CpG$_{O/E}$) and mean ρ were calculated on a

per-gene basis, as well as for exons and introns, and 50kb flanking regions 10kb upstream

and downstream of the gene, akin to Wallberg et al (2015). For recombination rate, the data

from the interval function in LDhat (Auton and McVean 2007) was used, utilising a weighted

mean when the element spanned multiple markers. For visualisation, extreme CpG$_{O/E}$ values

due to low GC content were set to a maximum of 4.

We tested for significant difference in mean between ρ and CpG$_{O/E}$ in flanking regions, exons

and introns using paired t-tests corrected for multiple testing using a Bonferroni threshold.

Differential expression data

We classified genes in both species according to their caste-biased patterns of expression

from two studies (Elsner et al. 2018; Lin et al. 2021). Elsner et al. (2018) analysed four castes

of *M. bellicosus*, with two age classes of each: major and minor workers, queens and kings.

They analysed differences in gene expression among these classes using RNAseq data. We

reclassified the differential expression data by grouping the direct caste-vs-caste

comparisons as "queen-biased", "king-biased", "worker-biased", "reproduction-biased",

"male-biased" or "female-biased" if they were differentially upregulated for this caste (or

group) in comparison to other castes (or groups). Since Elser *et al.* (2018) initially mapped

the *M. bellicosus* reads against the *Macrotermes natalensis* genome, We used orthofinder

(Emms and Kelly 2019) to identify the *M. bellicosus* orthologues corresponding to the

differentially expressed genes. Only single-copy genes were considered for this analysis.

Lin et al. (2021), performed RNAseq on samples of workers and queens. Here, we used the

data from untreated queens and workers. Differentially expressed *C. secundus* genes were

classified as "worker-biased" or "queen-biased" if they were differentially expressed with a

bias to the respective caste, or unbiased if they were not, and subsequently compared

between classes. The lists of genes we identified in the caste-biased categories in both

species are provided as Supplementary Tables S9 and S10.

Correlates of recombination rate variation among genes

In order to assess the relative importance of $CpG_{O/E}$ ratio and differential expression

patterns on recombination rate among genes, an ordinary least squares model was built

using statsmodels (Seabold and Perktold 2010). $CpG_{O/E}$, differential expression categories,

flanking regions $CpG_{O/E}$ and flanking region recombination rate were used as exogenous

variables.

Data access

Custom scripts are available on GitHub at https://github.com/troe27/termite-

recombination-rate and included as Supplemental Material. The Illumina whole-genome

sequencing data generated in this study have been submitted to the NCBI BioProject

database (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number

PRJNA1021607.

Competing interest statement

The authors declare no competing interests.

Acknowledgements

We acknowledge grant 2018-03896 from The Swedish Research Council to MTW and from

the Deutsche Forschungsgemeinschaft (DFG) KO-1895/26-1: 417980976 to JK. We

acknowledge support from the National Genomics Infrastructure in Stockholm funded by

Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish

Research Council. The computations were enabled by resources in project NAISS 2023/23-

402 provided by the National Academic Infrastructure for Supercomputing in Sweden

(NAISS) at UPPMAX, funded by the Swedish Research Council through grant agreement no.

2022-06725. We thank Göran Arnqvist for helpful discussions on the evolution of sex and

recombination. The Parks and Wildlife Commission, Northern Territory, and the Department

of the Environment, Water, Heritage and the Arts, Australia, gave permission to collect

(Permit number 59044) and export (PWS2016-001559) the *C. secundus* termites. The Office

Ivoirien des Parcs et Réserves (OIPR) provided sampling and export permits for the *M.*

bellicosus samples. The study was conducted according to the Nagoya protocol.

References

Agrawal AF. 2001. Sexual selection and the maintenance of sexual reproduction. *Nature*
755   **411**: 692–695.

756   Arsala D, Wu X, Yi SV, Lynch JA. 2022. Dnmt1a is essential for gene body methylation and
757   the regulation of the zygotic genome in a wasp. *PLOS Genetics* **18**: e1010181.

758   Auton A, McVean G. 2007. Recombination rate estimation in the presence of hotspots.
759   *Genome Research* **17**: 1219–27.

760   Axelsson E, Webster MT, Ratnakumar A, Consortium L, Ponting CP, Lindblad-Toh K. 2012.
761   Death of PRDM9 coincides with stabilization of the recombination landscape in the
762   dog genome. *Genome Res* **22**: 51–63.

763   Baer B. 2005. Sexual selection in Apis bees. *Apidologie* **36**: 187–200.

764   Bailey TL. 2021. STREME: accurate and versatile sequence motif discovery. *Bioinformatics*
765   **37**: 2834–2840.

Baker Z, Schumer M, Haba Y, Bashkirova L, Holland C, Rosenthal GG, Przeworski M. 2017. Repeated losses of PRDM9-directed recombination despite the conservation of PRDM9 across vertebrates ed. B. De Massy. *eLife* **6**: e24133.

Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mobile DNA* **6**: 11.

Baudat F, Buard J, Grey C, Fledel-Alon A, Ober C, Przeworski M, Coop G, de Massy B. 2010. PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. *Science* **327**: 836–840.

Begun DJ, Aquadro CF. 1992. Levels of naturally occurring DNA polymorphism correlate with recombination rates in D. melanogaster. *Nature* **356**: 519–520.

Berglund J, Quilez J, Arndt PF, Webster MT. 2014. Germline methylation patterns determine the distribution of recombination events in the dog genome. *Genome Biol Evol* **7**: 522–530.

Bewick AJ, Sanchez Z, Mckinney EC, Moore AJ, Moore PJ, Schmitz RJ. 2019. Dnmt1 is essential for egg production and embryo viability in the large milkweed bug, Oncopeltus fasciatus. *Epigenetics & Chromatin* **12**: 6.

Beye M, Gattermeier I, Hasselmann M, Gempe T, Schioett M, Baines JF, Schlipalius D, Mougel F, Emore C, Rueppell O, et al. 2006. Exceptionally high levels of recombination across the honey bee genome. *Genome research* **16**: 1339–44.

Browning BL, Zhou Y, Browning SR. 2018. A One-Penny Imputed Genome from Next-Generation Reference Panels. *The American Journal of Human Genetics* **103**: 338–348.

Browning SR, Browning BL. 2007. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am J Hum Genet* **81**: 1084–1097.

Burt A, Bell G, Harvey PH. 1991. Sex differences in recombination. *Journal of Evolutionary Biology* **4**: 259–277.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.

Cardoso DC, Santos HG, Cristiano MP. 2018. The Ant Chromosome database – ACdb: an online resource for ant (Hymenoptera: Formicidae) chromosome researchers. *Myrmecological News* **27**: 87–91.

Chan AH, Jenkins PA, Song YS. 2012. Genome-Wide Fine-Scale Recombination Rate Variation in Drosophila melanogaster. *PLOS Genetics* **8**: e1003090.

Charlesworth B. 2009. Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. *Nat Rev Genet* **10**: 195–205.

Coop G, Przeworski M. 2007. An evolutionary view of human recombination. *Nature reviews*
Genetics **8**: 23–34.

Cunha MS, Cardoso DC, Cristiano MP, de Oliveira Campos LA, Lopes DM. 2021. The Bee
Chromosome database (Hymenoptera: Apidae). *Apidologie* **52**: 493–502.

Elango N, Hunt BG, Goodisman MA, Yi SV. 2009. DNA methylation is widespread and
associated with differential gene expression in castes of the honeybee, Apis
mellifera. *Proceedings of the National Academy of Sciences of the United States of*
America **106**: 11206–11.

Elsner D, Meusemann K, Korb J. 2018. Longevity and transposon defense, the case of
termite reproductives. *PNAS* **115**: 5504–5509.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology* **20**: 238.

Fazalova V, Nevado B. 2020. Low Spontaneous Mutation Rate and Pleistocene Radiation of
Pea Aphids. *Mol Biol Evol* **37**: 2045–2051.

Fischer O, Schmid-Hempel P. 2005. Selection by parasites may increase host recombination
frequency. *Biol Lett* **1**: 193–195.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020.
RepeatModeler2 for automated genomic discovery of transposable element families.
Proceedings of the National Academy of Sciences **117**: 9451–9457.

Glastad KM, Hunt BG, Yi SV, Goodisman M a. D. 2011. DNA methylation in insects: on the
brink of the epigenomic era. *Insect Molecular Biology* **20**: 553–565.

Grant CE, Bailey TL, Noble WS. 2011. FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**: 1017–1018.

Harrison MC, Jongepier E, Robertson HM, Arning N, Bitard-Feildel T, Chao H, Childers CP,
Dinh H, Doddapaneni H, Dugan S, et al. 2018. Hemimetabolous genomes reveal
molecular basis of termite eusociality. *Nature Ecology & Evolution* **2**: 557–566.

Hartfield M, Keightley PD. 2012. Current hypotheses for the evolution of sex and
recombination. *Integrative Zoology* **7**: 192–209.

Hedrick PW, Parker JD. 1997. Evolutionary Genetics and Genetic Variation of Haplodiploids
and X-Linked Genes. *Annual Review of Ecology and Systematics* **28**: 55–83.

Hoffmann K, Foster KR, Korb J. 2012. Nest value mediates reproductive decision making
within termite societies. *Behavioral Ecology* **23**: 1203–1208.

Immler S. 2019. Haploid Selection in "Diploid" Organisms. *Annual Review of Ecology,*
Evolution, and Systematics **50**: 219–236.

Janicke T, Häderer IK, Lajeunesse MJ, Anthes N. 2016. Darwinian sex roles confirmed across
the animal kingdom. *Science Advances* **2**: e1500983.

Jankásek M, Kotyková Varadínová Z, Šťáhlavský F. 2021. Blattodea Karyotype Database. *EJE*
118: 192–199.

Jones JC, Wallberg A, Christmas MJ, Kapheim KM, Webster MT. 2019. Extreme Differences in
Recombination Rate between the Genomes of a Solitary and a Social Bee. *Mol Biol*
Evol **36**: 2277–2291.

Joseph J, Prentout D, Laverré A, Tricou T, Duret L. 2024. High prevalence of PRDM9-
independent recombination hotspots in placental mammals. *Proceedings of the*
National Academy of Sciences **121**: e2401973121.

Kawakami T, Wallberg A, Olsson A, Wintermantel D, de Miranda JR, Allsopp M, Rundlöf M,
Webster MT. 2019. Substantial Heritable Variation in Recombination Rate on
Multiple Scales in Honeybees and Bumblebees. *Genetics* **212**: 1101–1119.

Keightley PD, Ness RW, Halligan DL, Haddrill PR. 2014. Estimation of the Spontaneous
Mutation Rate per Nucleotide Site in a Drosophila melanogaster Full-Sib Family.
Genetics **196**: 313–320.

Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW,
Jiggins CD. 2015. Estimation of the Spontaneous Mutation Rate in Heliconius
melpomene. *Mol Biol Evol* **32**: 239–243.

Keightley PD, Trivedi U, Thomson M, Oliver F, Kumar S, Blaxter ML. 2009. Analysis of the
genome sequences of three Drosophila melanogaster spontaneous mutation
accumulation lines. *Genome Res* **19**: 1195–1201.

Kent CF, Minaei S, Harpur BA, Zayed A. 2012. Recombination is associated with the
evolution of genome structure and worker behavior in honey bees. *PNAS* **109**:
18012–18017.

Kent CF, Zayed A. 2013. Evolution of recombination and genome structure in eusocial
insects. *Communicative & Integrative Biology* **6**: e22919.

Koeniger G, Koeniger N, Phiancharoen M. 2011. Comparative Reproductive Biology of
Honeybees. In *Honeybees of Asia* (eds. H.R. Hepburn and S.E. Radloff), pp. 159–206,
Springer, Berlin, Heidelberg https://doi.org/10.1007/978-3-642-16422-4_8 (Accessed
February 13, 2024).

Korb J. 2008. Termites, hemimetabolous diploid white ants? *Frontiers in Zoology* **5**: 15.

Korb J, Hartfelder K. 2008. Life history and development - a framework for understanding
developmental plasticity in lower termites. *Biological Reviews* **83**: 295–313.

Korb J, Thorne B. 2017. Sociality in Termites. In *Comparative Social Evolution* (eds. D.R.
Rubenstein and P. Abbot), pp. 124–153, Cambridge University Press, Cambridge

https://www.cambridge.org/core/books/comparative-social-evolution/sociality-in-
termites/11A80CB9FA947760889676E03899E85C (Accessed August 9, 2023).

Krasovec M. 2021. The spontaneous mutation rate of Drosophila pseudoobscura. *G3*
(Bethesda) **11**: jkab151.

Lam I, Keeney S. 2015. Nonparadoxical evolutionary stability of the recombination initiation
landscape in yeast. *Science* **350**: 932–937.

Leffler EM, Bullaughey K, Matute DR, Meyer WK, Ségurel L, Venkat A, Andolfatto P,
Przeworski M. 2012. Revisiting an Old Riddle: What Determines Genetic Diversity
Levels within Species? *PLoS Biol* **10**: e1001388.

Lenormand T. 2003. The evolution of sex dimorphism in recombination. *Genetics* **163**: 811–
22.

Lenormand T, Dutheil J. 2005. Recombination Difference between Sexes: A Role for Haploid
Selection. *PLoS Biology* **3**: e63.

Lin S, Werle J, Korb J. 2021. Transcriptomic analyses of the termite, Cryptotermes secundus,
reveal a gene network underlying a long lifespan and high fecundity. *Commun Biol* **4**:
1–12.

Liu H, Jia Y, Sun X, Tian D, Hurst LD, Yang S. 2017. Direct Determination of the Mutation Rate
in the Bumblebee Reveals Evidence for Weak Recombination-Associated Mutation
and an Approximate Rate Constancy in Insects. *Mol Biol Evol* **34**: 119–130.

Liu H, Zhang X, Huang J, Chen J-Q, Tian D, Hurst LD, Yang S. 2015. Causes and consequences
of crossing-over evidenced via a high-resolution recombinational landscape of the
honey bee. *Genome Biol* **16**: 15.

Lyko F, Foret S, Kucharski R, Wolf S, Falckenhayn C, Maleszka R. 2010. The Honey Bee
Epigenomes: Differential Methylation of Brain DNA in Queens and Workers. *PLoS Biol*
8: e1000506.

Lynch M, Ali F, Lin T, Wang Y, Ni J, Long H. 2023. The divergence of mutation rates and
spectra across the Tree of Life. *EMBO reports* **24**: e57561.

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S.
2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024.
Nucleic Acids Research gkae241.

Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T,
Beutel RG, et al. 2014. Phylogenomics resolves the timing and pattern of insect
evolution. *Science* **346**: 763–767.

Myers S, Bottolo L, Freeman C, McVean G, Donnelly P. 2005. A fine-scale map of
recombination rates and hotspots across the human genome. *Science* **310**: 321–4.

Myers S, Bowden R, Tumian A, Bontrop RE, Freeman C, Macfie TS, McVean G, Donnelly P.
2010. Drive against hotspot motifs in primates implicates the PRDM9 gene in meiotic
recombination. *Science* **327**: 876–9.

Nachman MW. 2001. Single nucleotide polymorphisms and recombination rate in humans.
Trends in genetics : TIG **17**: 481–5.

Niehuis O, Gibson JD, Rosenberg MS, Pannebakker BA, Koevoets T, Judson AK, Desjardins
CA, Kennedy K, Duggan D, Beukeboom LW, et al. 2010. Recombination and Its Impact
on the Genome of the Haplodiploid Parasitoid Wasp Nasonia. *PLOS ONE* **5**: e8597.

Parvanov ED, Petkov PM, Paigen K. 2010. Prdm9 controls activation of mammalian
recombination hotspots. *Science* **327**: 835–835.

Persikov AV, Singh M. 2014. De novo prediction of DNA-binding specificities for Cys2His2
zinc finger proteins. *Nucleic Acids Research* **42**: 97–108.

Qiu B, Elsner D, Korb J. 2023. High-quality long-read genome assemblies reveal evolutionary
patterns of transposable elements and DNA methylation in termites.
2023.10.31.564968.
https://www.biorxiv.org/content/10.1101/2023.10.31.564968v1 (Accessed January
8, 2024).

Roisin Y. 2001. Caste sex ratios, sex linkage, and reproductive strategies in termites. *Insectes*
soc **48**: 224–230.

Romiguier J, Lourenco J, Gayral P, Faivre N, Weinert LA, Ravel S, Ballenghien M, Cahais V,
Bernard A, Loire E, et al. 2014. Population genomics of eusocial insects: the costs of a
vertebrate-like effective population size. *Journal of Evolutionary Biology* **27**: 593–
603.

Ross L, Blackmon H, Lorite P, Gokhman VE, Hardy NB. 2015. Recombination, chromosome
number and eusociality in the Hymenoptera. *Journal of Evolutionary Biology* **28**:
105–116.

Rueppell O, Kuster R, Miller K, Fouks B, Rubio Correa S, Collazo J, Phaincharoen M, Tingek S,
Koeniger N. 2016. A New Metazoan Recombination Rate Record and Consistently
High Recombination Rates in the Honey Bee Genus Apis Accompanied by Frequent
Inversions but Not Translocations. *Genome Biol Evol* **8**: 3653–3660.

Sarda S, Zeng J, Hunt BG, Yi SV. 2012. The Evolution of Invertebrate Gene Body Methylation.
Mol Biol Evol **29**: 1907–1916.

Seabold S, Perktold J. 2010. Statsmodels: Econometric and Statistical Modeling with Python.
In *Proceedings of the 9th Python in Science Conference* (eds. S. Van der Walt and J.
Millman), pp. 92–96.

Sherman PW. 1979. Insect Chromosome Numbers and Eusociality. *The American Naturalist*
113: 925–935.

Shi YY, Sun LX, Huang ZY, Wu XB, Zhu YQ, Zheng HJ, Zeng ZJ. 2013. A SNP Based High-Density
Linkage Map of Apis cerana Reveals a High Recombination Rate Similar to Apis
mellifera. *PLOS ONE* **8**: e76459.

Siller S. 2001. Sexual selection and the maintenance of sex. *Nature* **411**: 689–692.

Singhal S, Leffler EM, Sannareddy K, Turner I, Venn O, Hooper DM, Strand AI, Li Q, Raney B,
Balakrishnan CN, et al. 2015. Stable recombination hotspots in birds. *Science* **350**:
928–932.

Sirviö A, Gadau J, Rueppell O, Lamatsch D, Boomsma JJ, Pamilo P, PAGE Jr. RE. 2006. High
recombination frequency creates genotypic diversity in colonies of the leaf-cutting
ant Acromyrmex echinatior. *Journal of Evolutionary Biology* **19**: 1475–1485.

Sirviö A, Johnston JS, Wenseleers T, Pamilo P. 2011a. A high recombination rate in eusocial
Hymenoptera: evidence from the common wasp Vespula vulgaris. *BMC Genetics* **12**:
95.

Sirviö A, Pamilo P, Johnson RA, Page Jr Robert E, Gadau J. 2011b. ORIGIN AND EVOLUTION
OF THE DEPENDENT LINEAGES IN THE GENETIC CASTE DETERMINATION SYSTEM OF
POGONOMYRMEX ANTS. *Evolution* **65**: 869–884.

Smukowski Heil CS, Ellison C, Dubin M, Noor MAF. 2015. Recombining without Hotspots: A
Comprehensive Evolutionary Portrait of Recombination in Two Closely Related
Species of Drosophila. *Genome Biology and Evolution* **7**: 2829–2842.

Stapley J, Feulner PGD, Johnston SE, Santure AW, Smadja CM. 2017. Variation in
recombination frequency and distribution across eukaryotes: patterns and
processes. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**:
20160455.

Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. 2021. The Dfam community resource of
transposable element families, sequence models, and genome annotations. *Mobile
DNA* **12**: 2.

Templeton AR. 1979. Chromosome Number, Quantitative Genetics and Eusociality. *The
American Naturalist* **113**: 937–941.

Torres AP i, Höök L, Näsvall K, Shipilina D, Wiklund C, Vila R, Pruisscher P, Backström N.
2023. The fine-scale recombination rate variation and associations with genomic
features in a butterfly. *Genome Res* **33**: 810–823.

Trivers R. 1988. Sex differences in rates of recombination and sexual selection. In *The
Evolution of Sex*, pp. 270–286, Sinauer Associates, Sunderland, MA.

Trivers RL, Hare H. 1976. Haploidploidy and the Evolution of the Social Insect. *Science* **191**:
249–263.

Ventós-Alfonso A, Ylla G, Montañes J-C, Belles X. 2020. DNMT1 Promotes Genome
Methylation and Early Embryo Development in Cockroaches. *iScience* **23**: 101778.

Waiker P, de Abreu FCP, Luna-Lucena D, Freitas FCP, Simões ZLP, Rueppell O. 2021.
Recombination mapping of the Brazilian stingless bee Frieseomelitta varia confirms
high recombination rates in social hymenoptera. *BMC Genomics* **22**: 673.

Wallberg A, Bunikis I, Pettersson OV, Mosbech M-B, Childers AK, Evans JD, Mikheyev AS,
Robertson HM, Robinson GE, Webster MT. 2019. A hybrid de novo genome assembly
of the honeybee, Apis mellifera, with chromosome-length scaffolds. *BMC Genomics*
20: 275.

Wallberg A, Glémin S, Webster MT. 2015. Extreme recombination frequencies shape
genome variation and evolution in the honeybee, *Apis mellifera*. *PLoS Genetics* **11**:
e1005189.

Wallberg A, Han F, Wellhagen G, Dahle B, Kawata M, Haddad N, Simões ZLP, Allsopp MH,
Kandemir I, De la Rúa P, et al. 2014. A worldwide survey of genome sequence
variation provides insight into the evolutionary history of the honeybee *Apis*
mellifera. *Nat Genet* **46**: 1081–1088.

Wallberg A, Schöning C, Webster MT, Hasselmann M. 2017. Two extended haplotype blocks
are associated with adaptation to high altitude habitats in East African honey bees.
PLoS Genetics **13**: e1006792.

Wang J, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu S, Marchler GH, Song JS,
Thanki N, Yamashita RA, et al. 2023. The conserved domain database in 2023.
Nucleic Acids Res **51**: D384–D388.

Wang X, Wheeler D, Avery A, Rago A, Choi J-H, Colbourne JK, Clark AG, Werren JH. 2013.
Function and Evolution of DNA Methylation in Nasonia vitripennis. *PLoS Genet* **9**:
e1003872.

Watterson GA. 1975. On the number of segregating sites in genetical models without
recombination. *Theoretical Population Biology* **7**: 256–276.

Wilfert L, Gadau J, Schmid-Hempel P. 2007. Variation in genomic recombination rates
among animal taxa and the case of social insects. *Heredity (Edinb)* **98**: 189–197.

Winkler L, Moiron M, Morrow EH, Janicke T. 2021. Stronger net selection on males across
animals eds. P.J. Wittkopp and L. Holman. *eLife* **10**: e68316.

Winston ML. 1991. *The Biology of the Honey Bee*. Revised edition. Harvard University Press,
Cambridge, Mass.

Wu T-C, Lichten M. 1994. Meiosis-Induced Double-Strand Break Sites Determined by Yeast
Chromatin Structure. *Science* **263**: 515–518.

Yang S, Wang L, Huang J, Zhang X, Yuan Y, Chen J-Q, Hurst LD, Tian D. 2015. Parent–progeny
sequencing indicates higher mutation rates in heterozygotes. *Nature* **523**: 463–467.

Zhang C, Dong S-S, Xu J-Y, He W-M, Yang T-L. 2019. PopLDdecay: a fast and effective tool for
linkage disequilibrium decay analysis based on variant call format files.
Bioinformatics **35**: 1786–1788.

1019

Figure Legends:

1021

Figure 1. Genome-wide decay of LD measured as r^2 versus distance (bp) between SNPs for the two termite species (*M. bellicosus*, dark red; *C. secundus*, yellow) and honey bees (*A. mellifera scutellata*, blue; data from Wallberg et al. 2017).

Figure 2. Variation in recombination rate estimated a ρ/kbp across representative genomic scaffolds in a) *M. bellicosus* and b) *C. secundus*. Estimates of ρ/kbp are calculated in 10 Kbp and 100 Kbp windows.

Figure 3. Cumulative plot of proportion of recombination events versus proportion of the genome for the two termite species *M. bellicosus* and *C. secundus* compared to the honey bee *A. mellifera* (Wallberg et al. 2015). Dashed lines show the proportion of the genome where 50 % of the recombination occurs, which for *M. bellicosus* is 3.47 %, *C. secundus* is 4.77 %, and *A. mellifera* is 32.05 %.

Figure 4. PRMD9 structure and predicted zinc-finger motifs in the *M. bellicosus* and *C. secundus* genomes. A) Predicted structure of the PRDM9 protein sequence in *C. secundus.* B) Predicted structure of the PRDM9 gene in *M. bellicosus* identified by homology searches. Exons and conserved functional domains are marked. The mapped exons from *Zoothermopsis nevadensis* are also shown. C) Structure of PRDM9 gene in *C. secundus*. Exons and conserved functional domains are marked. D) Predicted zinc finger binding motif in *M. bellicosus,* E) Predicted zinc finger binding motif in *C. secundus*.

Figure 5. Boxplots showing variation in levels of CpG$_{O/E}$ and recombination rate (ρ/kbp; rho) in genic and flanking regions in the *M. bellicosus* and *C. secundus* genomes. The differences between all categories are significant.

Figure 6. Boxplots showing variation in levels of CpG$_{O/E}$ and recombination rate (ρ/kbp) in the coding region of genes in the *M. bellicosus* and *C. secundus* genomes classified according to their patterns of gene expression (Q = queen-biased, K = king-biased, W = worker-biased, F = female-biased, R = reproductive-biased).

1047

1048 **Table 1.** Estimates of levels genetic variation, effective population size and recombination
1049 rate in two termite species.

Species	Assembly length (Mb)	No. SNPs	θ_W/bp	ρ/kb	μ bp^{-1} gen^{-1}	N_E	r (cM/Mb)
M. bellicosus	1,139	5,581,662	0.14%	1.69	2.7 x 10^{-10}	1,278,982	0.033
					4.5 x 10^{-9}	76,876	0.550
					8.1 x 10^{-9}	42,898	0.986
C. secundus	992	15,037,468	0.44%	4.85	2.7 x 10^{-10}	4,080,725	0.030
					4.5 x 10^{-9}	245,280	0.494
					8.1 x 10^{-9}	136,869	0.885

θ_W/bp - Watterson's θ per base pair

ρ/kb - population recombination parameter, ρ, per kilobase

μ bp^{-1} gen^{-1} - mutation rate per base pair per generation taken from literature. Estimates correspond
to *Acyrthosiphon pisum* (2.7 x 10^{-10}), *Drosophila melanogaster* (4.5 x 10^{-9}) and *Drosophila*
pseudoobscura (8.1 x 10^{-9})(Lynch et al. 2023).

N_E - effective population size

r (cM/Mb) - recombination rate in centimorgans per megabase

1057

1058

1059
1060   Figure 1
1061

1062
1063   Figure 2
1064

1065
1066   Figure 3
1067

1068
1069   Figure 4
1070

1071
1072   Figure 5
1073

1074

1075

1076   Figure 6